

ROYAL FITNESS CLUB

RFC PREMIUM GUIDE #17

INTERMEDIATE ANABOLIC CYCLE BLUEPRINT

16-Week Protocol with Second Compound Integration

16 Weeks · 5 Chapters · Intermediate Level · Stack Strategy

© Royal Fitness Club — For Members Only

TABLE OF CONTENTS

Chapter 1 — Intermediate Cycle Principles	3
Chapter 2 — Stack Options: Adding a Second Compound	4
Chapter 3 — 16-Week Lean Bulk Protocol	5
Chapter 4 — Estrogen & Prolactin Management	6
Chapter 5 — Advanced Nutrition for Cycle	7

DISCLAIMER

For educational and harm-reduction purposes only. Not medical advice. Consult a qualified healthcare professional before starting any protocol. Individual results vary.

CHAPTER 1 — INTERMEDIATE CYCLE PRINCIPLES

"The intermediate cycle is the most important step in your development. Go too conservative and miss the potential. Go too aggressive and invite avoidable problems."

An intermediate cycle is appropriate for users who have: (1) completed at least one testosterone-only cycle with good recovery, (2) confirmed natural testosterone recovery via blood work, and (3) maximised gains from testosterone alone. The intermediate step typically involves adding a second compound to the testosterone base.

Criterion	Requirement for Intermediate Cycle
Previous cycles	Minimum 1 completed with full PCT recovery
Testosterone recovery	Confirmed via blood work — at least 80% of baseline
Training experience	3+ years; first cycle gains consolidated
Blood work knowledge	Understand all markers in your panel
Side effect management	Know what AI dose works for you from cycle 1
New compound research	Full understanding of pharmacology, half-life, risks of new compound

CHAPTER 2 — ADDING A SECOND COMPOUND

Best Second Compound Options

Compound	Synergy with Test	Primary Benefit	Main Risk	Beginner-Intermediate Suitability
Nandrolone (Deca)	Excellent — low E2	Joint support, mass, recovery	Prolactin elevation, sexual side effects	★★★★ (best first add)
Anavar (Oxandrolone)	Excellent — oral	Lean gains, strength, low risk	Liver strain (oral), expensive	★★★★ (safest oral)
Masteron	Excellent — anti-E	Hardening, anti-oestrogenic	Only works < 12% body fat	★★★ (cutting only)
Boldenone (EQ)	Good — mild	Appetite, endurance, lean mass	Slow acting, high RBC, long ester	★★★ (patience required)
Dianabol	Good — kick-start	Fast mass and strength gains	Liver, water retention, BP	★★★ (short-term oral)

CHAPTER 3 — 16-WEEK LEAN BULK PROTOCOL

Recommended Intermediate Stack: Testosterone Enanthate + NPP (Nandrolone Phenylpropionate)

Week	Test E	NPP	AI	Cabergoline	Notes
1–4	500mg/wk	300mg/wk	Anastrozole 0.5mg E3D	None (monitor)	Assess NPP response
5–12	500mg/wk	350mg/wk	Adjust to E2 bloods	0.25mg 2×/wk if prolactin rises	Peak phase
13–16	500mg/wk	200mg/wk	Continue AI	Continue Caber	Taper NPP early (long recovery)
17–18	None	None	Stop AI	Stop Caber	Clearance (NPP = 1 week)
19–22 (PCT)	None	None	None	None	Clomid + Nolvadex full 4 weeks

■ **NPP (short ester Nandrolone) is preferred over Deca for intermediate users — it clears faster, making side effect management and PCT timing more controllable.**

CHAPTER 4 — ESTROGEN & PROLACTIN MANAGEMENT

Nandrolone elevates prolactin via 19-nor compound mechanism. This requires a different management approach compared to testosterone-only cycles. Prolactin elevation causes sexual dysfunction (the infamous "Deca dick"), potential gynecomastia, and mood changes.

Compound	Estrogenic	Progestenic	Prolactin	Management
Testosterone	Yes (aromatises)	No	No	AI if needed
Nandrolone (NPP/Deca)	Low	Yes	Yes (significant)	Cabergoline + AI
Dianabol	High	No	Mild	AI essential
Masteron	None	No	No	None required
Anavar	None	No	No	None required

■ ***Cabergoline protocol for Nandrolone use: 0.25mg twice weekly. Do not use bromocriptine as an alternative — it has significantly worse side effects.***

CHAPTER 5 — ADVANCED NUTRITION FOR CYCLE

"An intermediate cycle deserves intermediate-level nutrition diligence. Tracking macros accurately is non-negotiable at this level."

Phase	Duration	Caloric Strategy	Protein Target	Carb Strategy
Weeks 1–4 (Setup)	4 weeks	+200 kcal surplus	2.0g/kg BW	Moderate — 3–4g/kg
Weeks 5–12 (Growth)	8 weeks	+400 kcal surplus	2.2g/kg BW	High — 4–5g/kg peri-workout
Weeks 13–16 (Lean out)	4 weeks	Maintenance to –100	2.5g/kg BW	Reduce slightly, maintain timing
PCT Phase	4–6 weeks	Maintenance to +100	2.3g/kg BW	Moderate — 3g/kg for hormone support

- Caloric surplus for Nandrolone + Test stack: 400–500 kcal above TDEE optimal
- Higher protein required: Nandrolone significantly increases nitrogen retention — supply the substrate
- Carbohydrate timing: 50% of daily carbs around training window (pre + post)
- Alcohol: zero during cycle. Nandrolone is processed differently with alcohol present