

12 WEEKS

FEMALE PHYSIOLOGY

TRAINING + DIET

SUPPLEMENT STACK

WOMEN'S COMPLETE BODY TRANSFORMATION GUIDE

12-Week Training · Nutrition · Hormone Awareness · Supplements · Anavar Protocol

Science-designed for female physiology — lean, strong, and confident

ELITE FITNESS GUIDE SERIES

Research-Based · Professional Edition · 2024

■ TABLE OF CONTENTS

1. Female Physiology & Body Composition Science
2. The 12-Week Transformation Framework
3. Weeks 1–4: Foundation Phase
4. Weeks 5–8: Intensification Phase
5. Weeks 9–12: Peak Phase
6. Nutrition for Women's Body Recomposition
7. Cardio & Fat Burning Protocols for Women
8. Training With Your Menstrual Cycle
9. Evidence-Based Supplement Stack for Women
10. Anavar Protocol for Women (Advanced)
11. Primobolan for Women (Advanced)
12. Progress Tracking & Long-Term Strategies

■ 1. FEMALE PHYSIOLOGY & BODY COMPOSITION SCIENCE

Women's body composition responds to diet and training differently than men's due to distinct hormonal profiles — primarily higher estrogen and lower testosterone. Understanding these differences enables a training and nutrition approach that **works with** female physiology, not against it.

Factor	Women	Men	Strategic Implication
Testosterone	20–60 ng/dL (15–30× lower)	300–1000 ng/dL	Slower muscle growth; longer timeframe needed — be patient
Estrogen	Cyclical 30–400 pg/mL	15–60 pg/mL (stable)	Promotes fat storage in hips/thighs; estrogen-dominant fat patterns
Fat oxidation	Higher at rest and during exercise	More glucose-dependent	Women are natural fat-burners; LISS cardio highly effective
Muscle fiber distribution	Higher type I (endurance) ratio	Higher type II (fast-twitch)	Higher reps (12–20) often produce better female hypertrophy
Strength:weight ratio	Generally lower	Generally higher	But women gain strength at similar RATE as men — neural adaptation
Cortisol sensitivity	Higher cortisol response to stress	Lower	Sleep and stress management are CRITICAL for women's physique goals

■ 2. 12-WEEK TRANSFORMATION FRAMEWORK

Phase	Weeks	Caloric Target	Training Focus	Cardio	Expected Result
Foundation	1–4	Mild deficit (~250 kcal)	Learn movements; build neuromuscular connection	3× 30min LISS/week	5–8 lbs fat loss; strength foundation built
Intensification	5–8	Moderate deficit (~400 kcal)	Increase volume; progressive overload; add intensity techniques	4× 35min/week + 1× HIIT	Additional 5–7 lbs fat; visible muscle shape
Peak	9–12	Aggressive deficit (~500 kcal)	Max volume; minimal rest; focus on weak points	5× 40min LISS + 2× HIIT	Final 3–5 lbs fat; conditioning peak; lean physique

■ 3–5. PHASE TRAINING PROGRAMS

Weeks 1–4 — Foundation Program (3 days/week):

Day	Muscle Groups	Key Exercises	Sets × Reps
Day A	Full Body	Goblet Squat · Romanian Deadlift · DB Bench · Seated Row · Shoulder Press · Plank	3×12–15 each
Day B	Full Body	Bulgarian Split Squat · Hip Thrust · Incline DB Press · Pull-Ups/Assisted · Lateral Raises · Dead Bug	3×12–15 each
Day C	Full Body	Sumo Deadlift · Step-Ups · Cable Flies · Face Pull · Arnold Press · Russian Twist	3×15–20 each

Weeks 5–8 — Intensification (4 days/week, Upper/Lower):

Day	Focus	Exercises (4 sets each)	Intensity Technique
Mon/Thu	Upper (Strength+Volume)	Bench Press · OHP · Barbell Row · Chin-Ups · Cable Flies · Lateral Raises	Drop sets on last set of isolation
Tue/Fri	Lower (Glutes/Legs priority)	Hip Thrust · RDL · Hack Squat · Leg Curl · Walking Lunge · Calf Raise	Pause reps on hip thrust and RDL

Weeks 9–12 — Peak (5 days/week, PPL-style): Full PPL split — same exercises as intensification phase with added volume (5 sets), shorter rest (60–75s), supersets for isolation work, and additional glute/shoulder specialization work on accessory days.

■ 6–7. NUTRITION & CARDIO FOR WOMEN

Macro	Target	Why	Top Foods
Protein	0.8–1.2 g/lb body weight	Muscle preservation + satiety — women often under-eat protein	Paneer, Greek yogurt, eggs, chicken, fish, tofu, legumes
Carbohydrates	100–180 g/day	Training performance; thyroid function; serotonin production (mood)	Oats, brown rice, sweet potato, fruits, vegetables
Fats	50–70 g/day	Estrogen production; skin health; fat-soluble vitamins	Avocado, nuts, olive oil, fatty fish, ghee

Macro	Target	Why	Top Foods
Calories	BMR × 1.4 - 300–500	Moderate deficit for sustainable results without hormonal disruption	Quality whole foods; avoid ultra-processed foods

Women's Cardio Priority Order:

- **Priority 1 — LISS Walking (daily):** 8,000–12,000 steps/day. Zero muscle stress; high fat burning; cortisol-neutral
- **Priority 2 — Moderate Cardio (3–4×/week):** 30–40 min cycling, swimming, elliptical at 65–70% max HR
- **Priority 3 — HIIT (max 2×/week):** 20 min sessions maximum; avoid during luteal phase; counterproductive if overdone
- **Fasted cardio:** Effective (20% more fat oxidized fasted) — but not mandatory; adherence > perfect protocol

■ 8. TRAINING WITH YOUR MENSTRUAL CYCLE

Phase	Days	Best Training	Adjust Diet	Key Notes
Menstrual	1–5	Light movement; yoga; walking	Normal calories; prioritize iron-rich foods	Energy often low; don't skip training entirely — movement helps cramps
Follicular	6–13	Heavy strength training; HIIT; PRs	Normal to slight surplus carbs	Rising estrogen = peak strength and pain tolerance; best time for max effort
Ovulatory	14	Absolute maximum performance day	Normal	Peak power output; set new personal records; joints slightly looser (injury awareness)
Luteal	15–28	Moderate intensity; more rest; reduce HIIT	Increase calories by 100–200 kcal; complex carbs	Progesterone dominates; fatigue, cravings increase; body temperature elevated

■ 9. EVIDENCE-BASED SUPPLEMENT STACK FOR WOMEN

Supplement	Dose	Evidence for Women
Iron (if deficient)	18–27 mg/day (with 500mg Vitamin C)	Menstrual iron loss; deficiency causes fatigue, poor recovery, reduced VO2 max
Magnesium Glycinate	300–400 mg nightly	Reduces PMS severity 40% in trials; improves sleep quality; muscle recovery
Omega-3 (EPA+DHA)	2–3 g/day	Reduces menstrual pain (prostaglandin inhibition); improves body composition; anti-inflammatory
Vitamin D3 + K2	3000 IU D3 + 100 mcg K2	Bone density; immune; testosterone production (even in women)
Ashwagandha (KSM-66)	300 mg twice daily	Reduces cortisol 20–30%; improves thyroid T4→T3 conversion; reduces stress eating
Creatine Monohydrate	3–5 g/day	Often overlooked for women — multiple studies show 5–15% strength gain; improves brain health; no fat gain
Collagen Peptides	10–15 g/day	Joint integrity; skin elasticity; connective tissue — important for high-volume training

■ 10–11. ANAVAR & PRIMOBOLAN FOR WOMEN (ADVANCED)

■■ **IMPORTANT:** These compounds are controlled substances in many jurisdictions. This information is provided for educational purposes only. Consult a physician. Women are significantly more sensitive to anabolic compounds — virilization (voice deepening, clitoral enlargement, body hair) can be PERMANENT.

Anavar (Oxandrolone) — Most Female-Friendly Anabolic:

Parameter	Conservative	Moderate	Aggressive
Dose	2.5–5 mg/day	5–10 mg/day	10–15 mg/day
Duration	6 weeks	6–8 weeks	8 weeks MAX
Virilization Risk	Very low	Low	Moderate
Expected Results	Lean maintenance + strength	5–8 lbs lean mass; hardness	10+ lbs lean mass; significant virilization risk
PCT	None required	None typically	None — but monitor hormones

Primobolan (Methenolone Acetate) for Women:

- Dose: 25–75 mg/week injectable OR 25–50 mg/day oral acetate form
- Duration: 8–12 weeks — excellent lean mass quality with minimal virilization
- Often considered the 'queen of women's compounds' — very low androgenic activity

- Stop immediately at first sign of voice changes, clitoral sensitivity changes, or excess body hair growth