

Reliable approximation in electronic structure methods:
from real space partitioning to quantum Monte Carlo approaches
Amanda Duménil Seminar at Oak Ridge National Laboratory

When is DFT not enough?

- DFT citation/publications demonstrate its applicability and often reliable performance
 - In strongly correlated regimes, or when functional and system physics do not agree
-

Reduce cost through various system reductions

Sampling Configurations Quantum Monte Carlo: applications and developments

Hilbert Space

selected CI as approximate trial wave functions

Real Space

Determining molecular fragments with unsupervised machine learning (UML)

QMC application: periodic absorption¹

H on graphene

Method	Binding energy (meV)
DMC	-691 \pm 19
PW91	-810 to -830, -870
PBE	-790, -840, -980

- functional dependent binding energy
- a need for benchmark values

¹A. Duménil et al, 'The Binding of Atomic Hydrogen on Graphene from Density Functional Theory and Diffusion Monte Carlo Calculations', *The Journal of Chemical Physics*, 156.14 (2022), 144702DOI:[10.1063/5.0085982](https://doi.org/10.1063/5.0085982).

Diffusion Monte Carlo

- Recast the Schrodinger equation in imaginary time

$$\frac{\partial |\Psi\rangle}{\partial \tau} = -\hat{H} |\Psi\rangle$$

- A formal solution to this:

$$|\psi(\tau_1 + \delta\tau)\rangle = e^{-\hat{H}\delta\tau} |\psi(\tau_1)\rangle$$

Anything non-orthogonal to the ground state is going to decay out exponentially

$$\lim_{\tau \rightarrow \infty} |\psi(\tau)\rangle = c_0 e^{-\epsilon_0 \tau} |\phi_0\rangle$$

$$-\frac{\partial \psi(\mathbf{R}, \tau)}{\partial \tau} =$$

$$\left[\sum_{i=1}^N -\frac{1}{2} \nabla_i^2 \psi(\mathbf{R}, \tau) \right] \text{ diffusion term}$$

+

$$(V(\mathbf{R}) - E_T) \psi(\mathbf{R}, \tau) \text{ branching term}$$

Importance sampling

- use of trial wavefunction for efficient sampling.

Fixed-node approximation

- Antisymmetry of fermions causes sampling problems
- Solution: fix nodes of trial wavefunction
- necessitates accurate nodal surface of trial wavefunction

QMC application: periodic absorption

H on graphene

Method	Binding energy (meV)
PBE	-820
PBE	-871
PBE0	-851 (-800)
HSE	-794 (-743)
DMC	-691 \pm 19
PW91	-810 to -830, -870
PBE	-790, -840, -980

- QMCPACK binding energies provide benchmark.
- Hybrid functionals are very close.
- Even with close binding energies, density seems to disagree in bonding region.

Reduce cost through various system reductions

Sampling Configurations Quantum Monte Carlo: applications and developments

Hilbert Space

selected CI as approximate trial wave functions

Real Space

Determining molecular fragments with unsupervised machine learning (UML)

QMC application: non-valence correlation bound anion²

NVCB rely on accurate description of correlation to bind

$$E_{corr} = E_{true} - E_{HF}$$

model system: H₂O₄

Radial integration of Hartree-Fock orbital density, $R = 4$ Angstroms

Selected-CI

$$\Psi_n = \sum c_I^{(n)} |D_I\rangle$$

$$e_\alpha = \frac{\langle \Psi^{(n)} | \hat{H} | \alpha \rangle^2}{E^{(n)} - \langle \alpha | \hat{H} | \alpha \rangle}.$$

$\{|\alpha\rangle\}$

single & double excitations

$\{|\alpha\rangle\}_*^{(n)}$

Ψ_n

Ψ_{n+1}

H₂O₄ Results

	wave function	basis set	EBE (meV)
D/HF		aug-cc-pVDZ+7s7p	183 ± 10
SD/HF		aug-cc-pVDZ	176 ± 12
SD/HF		cc-pVDZ	-528 ± 25
SD/B3LYP		aug-cc-pVDZ+7s7p	212 ± 11
SD/HF(N)//SD/NO SDCI(A)		aug-cc-pVDZ+7s7p	205 ± 10
SD/HF(N)//MD/NO SDCI(A)		aug-cc-pVDZ+7s7p	202 ± 12

²S. Upadhyay et al, ‘The Role of High-Order Electron Correlation Effects in a Model System for Non-Valence Correlation-Bound Anions’, *The Journal of Chemical Physics*, 153.22 (2020), 224118DOI:[10.1063/5.0030942](https://doi.org/10.1063/5.0030942).

wave function	basis set	EBE (meV)
MD/CIPSI NO	aug-cc-pVDZ+3s1p	190 ± 9

- Summary of selected-CI success/challenges and overall performance of other methods.

$R = 7$

- HF binds electron -> correlation not essential
- still orbital shape has noticeable changes
- DFT can recover those changes.

$R = 4$

- NCVB regime
- HF does not bind
- CIPSI shape changes slight, impacts energy.
- SDCI shape seems wrong

Follow-up Questions Ongoing projects at Sandia National Laboratories

- Can various selected CI approaches or later optimizations improve the compactness and quality of the trial wave function?
- Is there a balance in how well we are capturing dynamic vs static correlation in these systems? Especially in cases of energy differences.

↪ Aluminum systematic study: exploring various multideterminant generation and optimization schemes

- Selected CI captures static correlation, but should we take a second look at how dynamic correlation is captured?

↪ Jastrow form study

Jastrow Factor forms

$$\Psi(\{r_i\}\{r_I\}) = \exp(\mathcal{J})D(\{r_i\})$$

standard: parameterize separate 1,2, and 2-body terms. optimize parameters with VMC

A possible alternative form:

- Using framework of SNAP atomic potential
- represent the particle densities as bispectrum components.^{3 4}
- A perspective of particle neighborhoods which require fewer parameters for VMC optimization

Projection of the ρ particle density on the surface of the 4-D sphere:

$$E_{SNAP}^i(\mathbf{B}^i) = \beta_0^{\alpha_i} + \sum_{k=1}^K \beta_k^{\alpha_i} B_k^i = \beta_0^{\alpha_i} + \beta^{\alpha_i} \cdot \mathbf{B}^i$$

$$B_{j_1, j_2, j} = \sum_{m'_1, m_1 = -j_1}^{j_1} \sum_{m'_2, m_2 = -j_2}^{j_2} \sum_{m', m = -j}^j (c_{m' m}^j)^* C_{j_1 m_1 j_2 m_2}^{j m}$$

$$\times C_{j_1 m'_1 j_2 m'_2}^{j m'} c_{m'_1 m_1}^{j_1} c_{m'_2 m_2}^{j_2}$$

TO DO:

initial python sandbox for testing unit tests ensure single and multiple species work accessing LAMMPS API object interface with QMCPack

Reduce cost through various system reductions

Sampling Configurations Quantum Monte Carlo: applications and developments

Hilbert Space

selected CI as approximate trial wave functions

Real Space


Determining molecular fragments with unsupervised machine learning (UML)

³A. P. Bartók, R. Kondor, and G. Csányi, ‘On Representing Chemical Environments’, *Phys. Rev. B*, 87 (2013), 184115 DOI:[10.1103/PhysRevB.87.184115](https://doi.org/10.1103/PhysRevB.87.184115).

⁴A. P. Thompson et al, ‘Spectral Neighbor Analysis Method for Automated Generation of Quantum-Accurate Interatomic Potentials’, *Journal of Computational Physics*, 285 (2015), 316–30 DOI:<https://doi.org/10.1016/j.jcp.2014.12.018>.


Fragmenting with unsupervised machine learning

Problem





Fragmenting with unsupervised machine learning

Problem




Approach



Performance

Systems:

- water clusters
- methylthiophenes
- small polymer systems



Performance:

Summary:

- UML methods are able to resolve fragments
- spectral clustering performs well and is insensitive to descriptor

Acknowledgements

- Dr. Kenneth D. Jordan
- Dr. Daniel S. Lambrecht
- Dr. Luke Shulenburger, Dr. Raymond Clay

Computational Resources

- Center for Research computing, University of Pittsburgh
- Argonne Leadership Computing Facility
- Sandia National Laboratories

Funding:

- ACS: DNI-63213
- NSF: 1807638
- Sandia National Laboratories is a multimission laboratory managed and operated by National Technology & Engineering Solutions of Sandia, LLC, a wholly owned subsidiary of Honeywell International Inc., for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA0003525.

Conclusions

In regimes of strong electron correlation, mean field methods may be problematic and require need alternatives

Presented on three approaches:

- QMC: application to H_2O_4 and H on graphene \leftrightarrow systems where DFT varies.
- SCI methods in H_2O_4 as trial wave function
- UML MF: offer an approach to determining molecular fragmentation automatically, with very little information

References

- Albert P. Bartók, Risi Kondor, & Gábor Csányi, ‘On Representing Chemical Environments’, *Phys. Rev. B*, 87 (2013), 184115DOI:[10.1103/PhysRevB.87.184115](https://doi.org/10.1103/PhysRevB.87.184115)
- Amanda Dumí, Shiv Upadhyay, Leonardo Bernasconi, Hyeondeok Shin, Anouar Benali, & Kenneth D. Jordan, ‘The Binding of Atomic Hydrogen on Graphene from Density Functional Theory and Diffusion Monte Carlo Calculations’, *The Journal of Chemical Physics*, 156.14 (2022), 144702DOI:[10.1063/5.0085982](https://doi.org/10.1063/5.0085982)
- A. P. Thompson, L. P. Swiler, C. R. Trott, S. M. Foiles, & G. J. Tucker, ‘Spectral Neighbor Analysis Method for Automated Generation of Quantum-Accurate Interatomic Potentials’, *Journal of Computational Physics*, 285 (2015), 316–30DOI:<https://doi.org/10.1016/j.jcp.2014.12.018>
- Shiv Upadhyay, Amanda Dumí, James Shee, & Kenneth D. Jordan, ‘The Role of High-Order Electron Correlation Effects in a Model System for Non-Valence Correlation-Bound Anions’, *The Journal of Chemical Physics*, 153.22 (2020), 224118DOI:[10.1063/5.0030942](https://doi.org/10.1063/5.0030942)