

CIRCULATING COPY

LOAN COPY ONLY

UNIVERSITY OF CALIFORNIA, SAN DIEGO

Ecological and biogeochemical aspects of microbial degradation of phenolic
materials in the California coastal marine environment

A dissertation submitted in partial satisfaction of the requirements for the
Doctor of Philosophy in Marine Biology

by

Thomas J. Boyd

Committee in charge:

Professor A.F. Carlucci, Co-chair
Professor William F. Fenical, Co-chair
Professor Jeffery L. Bada
Professor Douglas H. Bartlett
Professor Willie C. Brown

1993

ABSTRACT OF THE DISSERTATION

Ecological and biogeochemical aspects of microbial degradation of phenolic materials in the California coastal marine environment

by

Thomas J. Boyd
Doctor of Philosophy in Marine Biology
University of California, San Diego, 1993
Professor A.F. Carlucci, Co-chair
Professor William F. Fenical, Co-chair

Phenolic materials are found in numerous environments. They are particularly well represented in coastal environments due to anthropogenic pollution, *in situ* production by marine organisms, and as a component of riverine humic materials flowing into estuaries. They were thus chosen as model compounds for the study of microbially mediated organic carbon dynamics in coastal systems. To determine the rates and utilization dynamics of phenolic materials by coastal marine bacteria, a sequential approach was used in which marine bacteria were first assayed to determine their ability to utilize phenolics, various sites in the California coastal zone were then analyzed qualitatively and quantitatively for phenolic materials, a method was developed to determine microbial utilization rates of phenolic materials using a high specific activity radiolabeled tracer and finally, utilization rates of *p*-cresol were determined at ambient concentrations.

Phenol, cresol isomers (*o*-, *m*-, and *p*-), catechols, and methoxyphenols were the predominant phenolic materials identified in several representative coastal environments off California. Concentrations ranged from about 2.5 to 370 ng·l⁻¹. Biodegradation rates of *p*-cresol were measured in five coastal areas using high specific activity ³H-labeled *p*-cresol added at no more than 10% of the ambient *p*-cresol concentration. Rates of microbial utilization were high, up to 35.5 ng·l⁻¹·hr⁻¹ in San Francisco Bay. Turnover times were calculated using the measured ambient concentration of *p*-cresol and ranged from 1.72 hours (San Francisco Bay) to 37 hours (at Spanish Landing in San Diego Bay). Utilization kinetics indicated microbial degradation of more complex, humic type material containing phenolic moieties.

From this work, it is concluded that biodegradation of phenolic compounds plays a major role in the biogeochemistry of organic materials in coastal marine environments. Rates determined for the breakdown of the phenolic component of humic materials in estuarine environments indicate that microbially mediated turnover times are considerably faster than for other means suggested, such as photodegradation or sedimentation.

ACKNOWLEDGMENTS

Research was funded in part by a grant from the National Sea Grant College Program, National Oceanic and Atmospheric Administration, U.S. Department of Commerce, under grant number NA89AA-D-SG138, project number R/CZ-98 through the California Sea Grant College, and in part by the California State Resources Agency. The views expressed herein are those of the author and do not necessarily reflect the views of NOAA or any of its sub-agencies. The U.S. Government is authorized to reproduce and distribute for governmental purposes. Funds and logistical support were also provided by the Office of Naval Research under grant N00014-87-K-0148 and by the Marine Life Research Group at Scripps Institution of Oceanography. Special thanks are given to the County Sanitation District of Los Angeles County and the crew of the vessel *RV Ocean Sentinel*, the Los Angeles Sanitation District and the crew of the vessel *RV La Mer*, and John D. Faulkner and Russel Vernon-Clark for review and comments on this manuscript.

