

UNIVERSITY OF CALIFORNIA, SAN DIEGO

Acid soluble peptides from the hemocytes of the ascidians: Characterization and  
antimicrobial activity.

A dissertation submitted in partial satisfaction of the requirements for the degree

Doctor of Philosophy

in

Marine Biology

by

John Andrew Tincu

2003

Committee in charge:

Victor D. Vacquier, Chair  
Kathy Barbeau  
William Fenical  
Robert Lehrer  
William Loomis  
Steven Taylor

## ABSTRACT OF THE DISSERTATION

Acid soluble peptides from the hemocytes of the ascidians: Characterization and antimicrobial activity.

by

John Andrew Tincu

Doctor of Philosophy in Marine Biology

University of California, San Diego, 2003

Victor D. Vacquier, Chair

The goal of this research is to investigate the biochemistry of solitary ascidians, with particular focus on the hemocytes (blood cells). *Styela plicata* and *Ciona intestinalis* have a high local abundance and their hemocytes are easily isolated, making them an appropriate model for histochemical studies. Because ascidian hemocytes mediate host defense reactions and accumulate specific metals, they are a potential source of novel antimicrobial and metal binding agents. Acid soluble peptides from ascidian hemocytes were investigated to determine their antimicrobial properties. Chapter II of this thesis presents a comprehensive review of marine invertebrate antimicrobial peptides. Chapter III describes the characterization of plicatamide, a modified octapeptide from hemocytes of the ascidian *S. plicata*. Chapter IV investigates the molecular origins of plicatamide and Chapter V describes

attempts to generate a polyclonal antibody to plicatamide. Chapter VI presents, in detail, the antimicrobial properties of plicatamide. Chapter VII describes two novel, partially characterized, antimicrobial peptides from the ascidian *C. intestinalis*. Chapter VIII introduces a novel, potential metal binding agent from the acid extracts of *S. plicata* hemocytes. Appendix A reviews natural peptide antibiotics from tunicates and appendix B presents the unusual gas-phase intramolecular rearrangement of tunichrome *Sp-1* presented in Chapter VIII.

Trainee Support  
Lehrer / Taylor; R/MP-93  
California Sea Grant