

The Development of the PKX Myxosporean, the Causative Agent of
Proliferative Kidney Disease, in Rainbow Trout Salmo gairdneri
Richardson

By

For the ONLY

MICHAEL LAWRENCE KENT
B.S. (Humboldt State University) 1977
M.S. (San Diego State University) 1981

DISSERTATION

Submitted in partial satisfaction of the requirements for the degree of

DOCTOR OF PHILOSOPHY

in

Comparative Pathology

in the

GRADUATE DIVISION

of the

UNIVERSITY OF CALIFORNIA

DAVIS

Approved:

Ronald P. Hedrick
James J. Olander
J. H. T. Herlihy

Committee in Charge

1985

The Development of the PKX Myxosporean, the Causative Agent of Proliferative Kidney Disease, in Rainbow Trout Salmo gairdneri Richardson.

Abstract

The endogenous development of PKX, the causative agent of proliferative kidney disease, is described in rainbow trout, Salmo gairdneri, by light and electron microscopy. Parasites multiplied by endogeny, binary fission and possibly plasmotomy in the renal interstitium, and elicited an interstitial nephritis. As the disease progressed parasites were found in the lumens of the tubules, and these represented daughter cells released from PKX parasites that had migrated to the tubules. Internal sporoblasts formed within these daughter cells (enveloping cells). The internal sporoblasts contained up to six daughter cells and some sporoblasts organized into spores with two spherical polar capsules. Although the spores persisted for several months after the interstitial PKX and associated inflammation had subsided, they did not complete their development as indicated by the fact that they remained within the enveloping cell and did not form valves.

The relationship of this spore to the vegetative stages was studied further by a transmission experiment. The myxosporean forms were only observed in the kidney tubules, while typical PKX also occurred in the blood and spleen. Parasite-free fish injected with the blood and spleen of infected fish developed both forms of the parasite, which further substantiates that PKX is the prespore form of the intraluminal myxosporean.

Because only incomplete spores were observed, the more precise taxonomic status of PKX was not determined. However, intraluminal stages of PKX show similarities to the genera Sphaerospora, Mitraspora and Parvicapsula. Furthermore, interstitial stages of PKX resemble the early stages of Sphaerospora.

The incomplete spore development and severe inflammatory response to PKX indicate that salmonids may be abnormal hosts. Fish were immunosuppressed with cortisol implants, and they exhibited greater densities of interstitial and intraluminal PKX but less interstitial hypercellularity. However, spores did not develop further than those observed in natural epizootics. This study supports the hypothesis that PKX is an early form of a myxosporean and that salmonids are abnormal hosts. A non-salmonid fish may be the primary host for PKX in which sporulation is completed.

Acknowledgements

I thank those who made this study possible. I am especially grateful to Dr. R.P. Hedrick for providing continual assistance and guidance throughout the study. Appreciation is also expressed to Drs. H. J. Olander and J. H. Theis for valuable suggestions.

I am also indebted to the following: Dr. J. Lom and C. Smith for valuable advice, R. Munn and P. Lee for electron microscopy, Dr. G. H. Stabenfeldt and T. Madley for cortisol analysis, Dr. C. B. Schreck for advice on cortisol implantation, G. Hoskins and H. Ramsey for providing specimens, D. Evins, R. Gunter, D. Manzer, J. Modin and J. Hyde for providing samples and cooperation with the study, and the California Seagrass College Program for providing financial support.

I also thank Drs. G. L. Hoffman and A. C. Olson, Jr. for inspiring my studies of fish parasites. I am especially indebted to my wife Cindy, who has provided continual love and support throughout my academic years.

This work is a result of research sponsored in part by NOAA, National Sea Grant College Program, Dept. of Commerce, under grant numbers NA80AA-D-00120 and NA85AA-D-SG140, project number R/F-100, through the California Sea Grant College Program, and in part by the California State Resources Agency. The U.S. Government is authorized to reproduce and distribute for governmental purposes.