

DESCRIPTION OF THE LARVAL
STAGES OF FIVE NORTHERN
CALIFORNIA SPECIES OF ROCKFISHES
(FAMILY SCORPAENIDAE) FROM REARING
STUDIES

CIRCULATING COPY
Sea Grant Depository

A Thesis presented to the Faculty
of
California State University, Stanislaus
and
Moss Landing Marine Laboratories

In Partial Fulfillment
Of the Requirements for the Degree
Master of Science in Marine Science

by
Guillermo Moreno
October, 1990

ABSTRACT

There are about 72 species of *Sebastes* (Family Scorpaenidae) along the eastern Pacific coast of North America, some of which are heavily exploited by both commercial and sport fisheries. Larval identifications are needed for management-related studies of these fishes. However, due to the large number of species and recent evolutionary divergence in this genus, the identification of early life stages has progressed slowly. In this study, four species of rockfish (*Sebastes mystinus*, *S. carnatus*, *S. atrovirens*, and *S. rastrelliger*) were reared and described, and larvae of another (*S. melanops*) reared elsewhere were described. The larvae were fed a mix of rotifers, brine shrimp (*Artemia*) nauplii, and daily-caught plankton. Of the descriptions of eastern Pacific *Sebastes* larvae to date, only nine have been based on reared larvae, due to the difficulty in rearing them past the yolk absorption stage.

Two general pigmentation patterns were discerned: (1) a short row of ventral midline melanophores on the trunk, and no or very little postero-dorsal pigmentation (*S. mystinus* and *S. melanops*); and (2) complete ventral midline pigmentation on the trunk, and anterior and postero-dorsal melanophores (*S. carnatus*, *S. atrovirens*, and *S. rastrelliger*). With the exception of very early stages of *S. carnatus* and *S. atrovirens*, these five species can be distinguished from each other based

on pigmentation characteristics. The morphometric proportions did not demonstrate major differences among species.

Although difficult, identification of the larval stages of *Sebastes* spp. is possible for most of the species described to date. Culture and descriptive techniques need to be applied to other species of *Sebastes* to help reduce the number of factors that confuse the taxonomy of this complex group.

ACKNOWLEDGEMENTS

First, I would like to thank my parents María Elena and Ricardo Moreno for their moral support and encouragement through these years.

I owe many thanks to my committee members, Drs. G. M. Cailliet, V. Loeb, and P. Roe for their support and interest in my work, and for critically reviewing this manuscript. The final draft of this thesis would not have been possible without their comments and ideas.

Many people helped in this project, however, I cannot mention all of them. Suffice it to say that I am indebted to all of them. Many thanks are due to L. McMasters who drew the larvae and helped many times with drafting and photographic work.

I would like to thank C. Miller for providing the *S. melanops* larvae. Food rearing techniques were improved with the help of R. Orhun at Hubbs/Sea World Research Institute.

This research would not have been possible without the help and collaboration of the Monterey Bay Aquarium. I am grateful to the Husbandry Department at the MBA for its continuous interest and logistical support.

Many thanks are due to the faculty, staff, and students of Moss Landing Marine Laboratories. Their help and encouragement have been extraordinary.

This work is the result of research sponsored in part by NOAA,

National Sea Grant College Program, Department of Commerce, under grant number NA85AA-D-56140, project number R/F-115, through the California Sea Grant College Program, and in part by the California State Resources Agency. The U.S. Government is authorized to reproduce and distribute for governmental purposes. This grant was awarded to Drs. G. M. Cailliet, and V. Loeb. Additional support was given by the David and Lucile Packard Foundation.