

LOAN COPY ONLY

UNIVERSITY OF CALIFORNIA

SANTA CRUZ

DOMOIC ACID ANALYSES OF
ZOOPLANKTON, FISH AND RHODOPHYTA IN MONTEREY BAY.

A thesis submitted in partial satisfaction
of the requirements for the degree of

MASTER OF SCIENCE

in


MARINE SCIENCES

by

Graeme John Haywood

May 1995

The thesis of Graeme John Haywood is
approved:

Three handwritten signatures are written on three horizontal lines. The first signature is 'Mary S. ...', the second is 'Bennard', and the third is a stylized signature.

Dean of Graduate Studies and Research

ACKNOWLEDGMENTS

Mary Silver has been a fabulous supervisor, who ensured that my research project was interesting and challenging, introduced me to the world of biological oceanography and plankton biology and provided me with unique experiences to examine mesopelagic zooplankton from a remotely operated vehicle.

Much of the research for this thesis was completed in the HPLC laboratory of R Tjeerdema at UC, Santa Cruz. Ron gave his time and resources generously toward this project, and I always felt that I had his full support throughout my degree program.

Bess Ward, my other adviser has always treated me as a friend which I appreciate. Thankyou for reading my manuscript

Despite having his own workload and projects to complete, Mark Cattey taught me how to prepare samples for analysis, operate the chemstation and HPLC and then assist in the interpretation of data. Since no previous work has been carried out on domoic acid in zooplankton, considerable trial and error was necessary. I could not have completed this project without Mark's patience and assistance.

I would like to acknowledge Francisco Chavez of MBARI for permitting me to use zooplankton collected on his cruise for this research project, and Kurt Buck for sharing his extensive knowledge with me on several occasions. The staff at MBARI, including the crews of the Point Lobos, and Point Sur created opportunities which were highlights of my time in California.

Most importantly, I thank John Ashen, Susan Coale, David Garrison, Linda Goff, Marcia Gowing, Peter Miller, Mike Murrell, Rafael Olivieri, Gordon Smith, Debbie Steinberg, Joe Zuccarello and Peter Walz, for their support and assistance through this project.

Jenny Spiegel, my life partner and Herb Matthew who inspired me to 'don't ever give up' have been my long term support team over the several years which I really appreciate.

This research was funded by a grant from the National Sea Grant College Program, National Oceanic and Atmospheric Administration (NOAA), US Department of Commerce, under grant number NA36RG0537, project number R/CZ-124, through the California Sea Grant College. The views expressed herein are those of the authors and do not necessarily reflect the views of NOAA or any of its sub-agencies. Also, the US Government is authorized to reproduce and distribute for governmental purposes.

This research project was also supported by grants from the UCSC Institute of Marine Sciences, and the Ether and Earl Myers Trust. Funding was also received from the UC Toxic Substances Research and Teaching Program, Coastal Toxicology Component.

CHAPTER 1.
DOMOIC ACID IN ZOOPLANKTON AND FISH;
MONTEREY BAY CALIFORNIA

ABSTRACT

In late summer 1991, seabird fatalities drew attention, for the first time, to the presence of domoic acid in Monterey Bay. The purpose of the present study was to discover whether the toxin has been present in earlier years, using samples of plankton collected at UCSC from as early as 1979. Examination of the samples revealed that this was not a new phenomenon: domoic acid is detectable in zooplankton obtained during at least nine of the preceding thirteen years. Furthermore, domoic acid permeates the food web, entering all tested taxa of zooplankton and small fish. At each trophic level, there is an increase in the amount of domoic acid found in each animal. On a concentration basis, however, domoic acid is not bioaccumulated.

CHAPTER 2.**THE RESPONSE OF COPEPODS TO DIETARY DOMOIC ACID.****ABSTRACT.**

Domoic acid is a naturally occurring neurotoxin produced by some *Pseudo-nitzschia* species. Copepods given a diet of this diatom fed heavily and toxin body burdens reached a steady state with bioaccumulation occurring during the first 10 min. of exposure. When switched to an alternate non toxic species, domoic acid was rapidly depurated, suggesting that it was only present in the alimentary canal. No domoic acid related mortality was observed in the copepod and the results from this study indicate that copepods provide only a short-term vector during plankton blooms for domoic acid in marine food chains.

CHAPTER 3

A survey of macroalgae for the presence of domoic acid. Santa Cruz, California.

ABSTRACT.

Domoic acid is a powerful neurotoxin that has been found in a few Rhodophytes and some species of *Pseudo-nitzschia*. The main toxin producing source of domoic acid in Monterey identified to date is *Pseudonitzschia australis*. In seeking another possible source of the toxin, a survey of the more common red macroalgae in the Santa Cruz area was made. Domoic acid was not detected in the specimens tested.

National Sea Grant Depository
Pell Library Building - GSO
University of Rhode Island
Narragansett, RI 02882-1197USA
