

LOAN COPY ONLY

Fisheries Models of Pelagic Species

By

JOSÉ ANTÓNIO MARTINS GOULART

B.S. (Santa Clara University) 1969

M.S. (Santa Clara University) 1971

CIRCULATING COPY
Sea Grant Depository

DISSERTATION

Submitted in partial satisfaction of the requirements for the degree of

DOCTOR OF PHILOSOPHY

in

Applied Mathematics

in the

GRADUATE DIVISION

of the

UNIVERSITY OF CALIFORNIA

DAVIS

Approved:

Mane Mangel
Chris W. Sargent
John H. Hellawell

Committee in charge

1989

Fisheries Models of Pelagic Species

ABSTRACT

An optimal vessel allocation strategy aimed at maximizing the season's catch is developed for a K -vessel fleet operating on a region consisting of N adjacent cells over a M -period fishing season. The direction of pelagic fish movement and the "a priori" probability about the location of fish are assumed to be known from historical data; the model formulation relies on Bayesian updating and dynamic programming. Two examples of the model's performance are given for $K = 2$ and $K = 3$, with $N = 3$ and $M = 2$.

A new method for the determination of abundance of a pelagic stock is derived. The model refers to a multi-cell fishing region and applies search theory to the description of the detection process, since search effort represents a considerable fraction of the total fishing effort. The concept of random search for patches of schools and non-random search for schools within a patch is employed. An operational characterization of the search parameter, the pattern of fish movement and tagging data (number of detections and corresponding search times) provides the necessary fishery information. The abundance estimate is calculated through maximum likelihood estimation (MLE). Computer simulations for the special case of a two-cell

fishing area and a two-week fishing season confirm the method's good predictive ability.

The knowledge of the mechanisms which govern the movement of pelagic fish (e.g. tuna) is important for the study of fisheries interactions and the definition of fishing policy, both at the national and international levels. The basic model assumes that the spatiotemporal distribution of tuna can be described by a two-dimensional Brownian motion process with drift and is adjusted to account for: (a) instant catching and perfect catchability; (b) instant catching and imperfect catchability; (c) catch interval, stock depletion and operational catchability; and (d) catch interval, stock depletion, operational catchability and natural mortality. Bayesian and maximum likelihood estimation techniques are applied to determine the drift and diffusion parameters associated with tagged tuna movements. Using simulated tagging data sets, the method's performance is illustrated for different sampling conditions and parameter configurations.

occasion to participate in stimulating discussions on the various topics of the research project.

I express particular appreciation to my father and mother for their constant attentions and loving care. They were always present in times of need and no act can ever repay so much giving.

A final word is addressed to my friends, some of whom have been uniquely generous with their time and resources. To all, I am grateful for the companionship and understanding of life's predicaments.

This work is a result of research sponsored in part by NOAA, National Sea Grant College Program, Department of Commerce, under grant number NA85AA-D-SG140, project number R/F-109, through the California Sea Grant College Program. The U.S. Government is authorized to reproduce and distribute for governmental purposes.

Additional support was provided by the Secretaria Regional da Agricultura e Pescas (Regional Government of the Azores), in compliance with the Azores Tuna Stock Assessment Research Program, and the University of the Azores, through a scholarship grant that covered the nominal expenses of the doctoral program. The data on the South Pacific tuna were furnished by Dr. Ray Hilborn while he was associated with the South Pacific Commission, Noumea, New Caledonia.