

LEAH COA-1

SEA GRANT COLLEGE PROGRAM

**CIRCULATING COPY**  
**Sea Grant Depository**

UNIVERSITY OF CALIFORNIA, SAN DIEGO  
SAN DIEGO STATE UNIVERSITY

Chitobiase and Chitinase from *Vibrio harveyi*

A dissertation submitted in partial satisfaction of the  
requirements for the degree Doctor of Philosophy in  
Biology

by

Rafael W. Soto-Gil

Committee in charge:

University of California, San Diego

Professor Donald R. Helinski  
Professor Douglas W. Smith

San Diego State University

Professor Judith W. Zyskind, Chairperson  
Professor A. Stephen Dahms  
Professor Sanford I. Bernstein

1988

California Sea Grant College Program

Zyskind; R/F-93  
NA80AA-D-00120 & NA85AA-D-SG140

# ABSTRACT OF THE DISERTATION

Chitobiase and Chitinase from *Vibrio harveyi*

by

Rafael W. Soto-Gil

Doctor of Philosophy in Biology

University of California, San Diego, 1988

Professor Judith W. Zyskind, Chairperson

The *Vibrio harveyi* N,N'-diacetylchitobiase (chitobiase) and chitinase genes were cloned in *E. coli*. Chitobiase activity was found to be strongly induced by chitobiose in *V. harveyi*, whereas, when cloned in *E. coli*, chitobiase gene (*chb*) expression was constitutive. Chitobiase was localized to the outer membrane in *E. coli* clones harboring the *chb* gene, and it was exported with concomitant removal of a signal peptide. Maturation of chitobiase may follow a pathway similar to that of the major outer membrane lipoprotein (Lpp) of *E. coli*. A region six amino acids in length surrounding the Lpp processing site was found to be identical to a corresponding region of chitobiase, and processing of chitobiase is inhibited by the signal peptidase II specific inhibitor, globomycin. A protein purification scheme was developed where chitobiase was obtained in up to 30% yield by detergent removal from the surface of *E. coli* cells containing the *chb* gene, followed by a HPLC purification step using ion-exchange chromatography. The purified protein had a specific activity of 104 U/mg and an

apparent  $M_r$  of 92,000 deduced by SDS-PAGE. Transcriptional analysis of the *chb* gene promoter revealed three mRNA start sites in *E. coli* harboring the *chb* gene, two of these are also used by *V. harveyi*. An amino acid sequence comparison is made between chitobiase and the human hexosaminidase A. Two chitinase genes from *V. harveyi* were cloned, and the nucleotide sequence of one of these genes (*chiA*) is described. Chitinase expressed from a truncated (missing ~580 nucleotides at the 3' end) version of the *chiA* gene was excreted from *E. coli* cells. A comparison is made between the deduced amino acid sequences of two *S. marcescens* chitinases and *V. harveyi* chitinase.