

**LOAN COPY ONLY**

**BETWEEN-YEAR DIFFERENCES IN ABUNDANCE AND DISTRIBUTION OF  
LARVAL FISHES AND ASSOCIATED ENVIRONMENTAL CONDITIONS  
OFF DAVENPORT, CALIFORNIA, 1991-1993**

**A Thesis Presented to the Faculty  
of  
California State University, Stanislaus  
through  
Moss Landing Marine Laboratories**

**In Partial Fulfillment  
of the Requirements for the Degree  
of Master of Science in Marine Science**

**By  
Brendan J. Daly  
December 1997**

CERTIFICATION OF APPROVAL

BETWEEN-YEAR DIFFERENCES IN ABUNDANCE AND DISTRIBUTION OF  
LARVAL FISHES AND ASSOCIATED ENVIRONMENTAL CONDITIONS  
OFF DAVENPORT, CALIFORNIA, 1991-1993

By  
Brendan J. Daly

Gregor M. Cailliet  
Dr. Gregor M. Cailliet  
Professor of Ichthyology

12/3/97  
Date

Valerie J. Loeb  
Dr. Valerie J. Loeb  
Adjunct Professor of Biological Oceanography

3 Dec '97  
Date

Mary M. Yoklavich  
Mary M. Yoklavich  
Research Fishery Biologist

3 Dec 97  
Date

Pamela Roe  
Dr. Pamela Roe  
Professor of Zoology

Dec. 3, 1997  
Date

## ACKNOWLEDGMENTS

This study was funded in part by a grant from the National Sea Grant College Program, National Oceanographic and Atmospheric Administration, U.S. Department of Commerce, under grant number NA89AA-D-SG138, project number R/F-142 to V.J. Loeb, M.M. Yoklavich, and G.M. Cailliet through the California Sea Grant College, and the California State Resources Agency. The views expressed herein are those of the author and do not necessarily reflect the views of NOAA or any of its sub-agencies. Additional funding was provided by the Dr. Earl H. Myers and Ethel M. Myers Oceanographic and Marine Biology Trust.

I sincerely thank the members of my graduate advisory committee, Dr. Gregor M. Cailliet, Dr. Valerie J. Loeb, Dr. Pamela Roe, and Mary M. Yoklavich, for their advice, patience, and support throughout this project.

## ABSTRACT

Ichthyoplankton was sampled from December 1991 through April 1993 at five stations along an onshore-offshore transect near Davenport, California. In total, 66,301 larval fishes representing 63 taxa were collected. The five most abundant taxa (*Merluccius productus*, *Engraulis mordax*, *Sebastes* spp., *Stenobranchius leucopsarus* and *Genyonemus lineatus*) comprised > 94% of all larvae. *Engraulis mordax* abundance during January-April 1992 was three times higher than in comparable samples from 1993; in contrast, *M. productus* abundance was more than 50 times lower during January-April 1992 than in comparable samples from 1993. Additionally, mean January-April abundance of many larval fish taxa peaked further inshore in 1992 than 1993. These differences were attributed to oceanographic conditions associated with the 1992-93 El Niño. Cluster analysis of data from January-April samples revealed ecologically significant clusters in 1992 and 1993. Finally, a survey was completed comparing ichthyoplankton abundance and taxonomic composition in Davenport samples with those inside Monterey Bay.

