

Neurosecretory Staining in the Cerebral and Pleural-Pedal
Ganglia of *Haliotis discus hannai* and *Trochus niloticus*,
and its Relationship to Reproduction.

By

LOAN COPY ONLY

KIRK OTTO HAHN
B.S. (University of California, Irvine) 1975

DISSERTATION

Submitted in partial satisfaction of the requirements for the degree of

DOCTOR OF PHILOSOPHY

in

ENDOCRINOLOGY

in the


GRADUATE DIVISION

of the

UNIVERSITY OF CALIFORNIA

DAVIS

Approved:



Committee in Charge

1990

REPRODUCIBLE COPY
See Grant Depository

CIRCULATING COPY
See Grant Depository

ABSTRACT

Neurosecretory staining in the brain ganglia of two economically important prosobranch gastropods, Haliotis discus hannai and Trochus niloticus, was studied during their respective reproductive cycles, and the variation in stain affinity of the neurosecretory cells and amount of neurosecretory material contained in the neurosecretory cells were measured.

Neurosecretory activity was found in Cell Types A and B in the cerebral ganglia of Haliotis discus hannai, and the amount of neurosecretory material in the cells and stain intensity varied with the reproductive cycle. Cell Types C and D were not neurosecretory. The stain intensity of Cell Type A showed a correlation with vitellogenesis in the ovary. The stain intensity of Cell Type A in males showed no correlation with a physiological function in the testis. Cell Type B in both sexes did not show a correlation with either gametogenesis, vitellogenesis or spawning. The pleural-pedal ganglion contained 8 different cell types. The quantity of different cell types in the pleural-pedal ganglion was unusual for a prosobranch gastropod. Cell Types, #1- and #7-cells, showed neurosecretory staining and variation in the stain intensity during the reproductive cycle. Cell Type #1 was the most abundant cell in the pleural-pedal ganglion and the variation of neurosecretory material clearly correlated with gametogenesis.

Neurosecretion in Cell Type #7 showed a strong correlation with the induction of spawning. A possible neurohemal organ (pleural organ) was found in close proximity to the #7-cells in the statocyst tissue. The pleural organ is the first ectodermal neurohemal organ reported in a prosobranch gastropod. A nervous connection (stato-pleural nerve) with a possible sensory function was found between the statocyst and the pleural-pedal ganglion. The stato-cerebral nerve probably does not transmit the nervous impulses from the statocyst as has been previously reported.

The reproduction of Trochus niloticus was very unusual and was not similar to any previously described prosobranch. The population as a whole was asynchronous, nevertheless, the population contained several sub-groups which had synchronized reproductive cycles. Spawning occurred primarily at the new moon lunar phase (the day of the new moon and the 3 following days). The reproductive cycle was approximately 8 lunar months. T. niloticus was a seasonal spawner with complete spawning and each individual had only one co-hort of gametes developing at a time. Gametogenesis occurred during the first 1 to 2 lunar months after spawning, and then there was a relatively linear increase in the size of the oocytes during vitellogenesis. A second layer formed around the oocyte at the end of gonad development.

Two major cell types were found in the cerebral ganglion. The AF stain intensity of Cell Type 1 remained relatively constant at a high level for most of the reproductive cycle;

however, there were decreases in the intensity during two periods; first to third lunar month (period of rapid oocyte growth) and eighth to ninth lunar month (period of rapid expansion of secondary layer). The stain intensity of Cell Type 1 in the cerebral ganglion of males showed no variation. The stain intensity of Cell Type 2 was very low in most individuals and the stain intensity in the left cerebral ganglion was consistently reduced compared to the right cerebral ganglion. Eleven cell types were found in the pleural-pedal ganglion. The larger cells were along the edge of the ganglion with the medulla relatively empty. Most cells of all cell types were located along the cerebro-pedal and cerebro-pleural connectives, and along the curve of the pleural-pedal ganglion surrounding the statocysts. The cells near the statocyst showed neurosecretory staining with AF and Herlant's. P-cells contained intensely staining neurosecretory material, and the amount of neurosecretory granules and the intensity of staining varied during the experiment. The oscillation of the stain intensity in P-cells did not show any correlation with the reproductive cycle. The structure of the statocysts was typical of gastropods. The morphology of the statocyst was almost identical to the statocyst in abalone. A stato-pleural nerve was found connecting the sensory cells of the statocyst to the pleural-pedal ganglion.

A preliminary study on the production of yolk in the red abalone, Haliotis rufescens, is reported. The molecular

weight of the major yolk protein is approximately 2×10^6 daltons. The yolk protein in the red abalone is insoluble in dilute buffer solutions and must be isolated in distilled water. Experiments were conducted to measure the concentration of the major yolk protein in the blood during the reproductive cycle, however, the results were inconclusive.

ACKNOWLEDGMENTS

The path to a doctoral degree is long and circuitous, and along the way many people contribute to the final product. My parents, Joy and Otto, supported me through all my years as a student and encouraged me to learn. They were always there for me and helped me make those difficult decisions along the way.

I especially want to thank my major professor, Ernest S. Chang, for his support during my graduate career and his review of this thesis. His guidance has made me a better researcher and helped me look at problems from different perspectives. I would also like to thank my thesis committee, Wally Clark and Bruce Hammock, for their help with my doctoral research and review of this thesis. The students and staff of Dr. Bruce Hammock's laboratory are especially thanked for their help in teaching me the techniques of ELISA and helping me evaluate the resulting data.

I want to thank Franz Engelmann (University of California, Los Angeles) for teaching me the basic principles of research and attention to detail. He allowed me to begin my search for knowledge of endocrinology in abalone.

The Japan Ministry of Education, Tohoku University (Sendai, Japan), and the University of California gave me a Monbusho Scholarship to spend 1 year in Japan to study abalone endocrinology and reproduction. My study of abalone reproduction in Japan formed the foundation of my doctoral

research and, more significantly, the thrust of my future research and career. Tetsuo Seki (Oyster Research Institute) and Nagahisa Uki (Tohoku Regional Fisheries Research Laboratory) spent many hours with me discussing the role of research in the progress of abalone aquaculture. Their assistance and friendship made my stay in Japan very enjoyable.

Mitsue Doi was invaluable to my studies in Japan. She helped me learn Japanese and assisted in dealing with the many day-to-day problems. She also translated many Japanese research papers into English and did an excellent job in the difficult task of translating these highly technical papers. Her friendship and help over the years has been greatly appreciated.

George Trevelyan and Nancy Coultrup assisted me in collecting data for my research on the reproduction of the tropical top shell. Their help during the busy and very exciting days on Moorea was critical to the success of the project.

I would like to thank Dr. W. Miller for giving me representative histological slides from his study on the neurosecretion of the red abalone.

Eleanor Uhlinger was invaluable in helping me find the background research papers needed to investigate the wide ranging questions posed by my doctoral research. Her work as research librarian of the Bodega Marine Laboratory greatly reduced my time spent locating research papers. She was able

to locate obscure and difficult to find papers published throughout the world.

The National Sea Grant College Program and James Sullivan, Director of the California Sea Grant College Program, are thanked for supporting my graduate research with a traineeship. The traineeship was invaluable and allowed me to devote my full effort to research.

I thank NOAA, National Sea Grant College Program, and the Department of Commerce for partial support of my research on abalone and tropical top shell (Grant NA80AA-D-00120, Project R/A-61 to Ernest S. Chang). I also received partial funding from the University of California Jastro-Shields Research Scholarship, Lerner-Gray Fund for Marine Research, Sigma-Xi and University of California Research Expeditions Program.

The members of Ernest Chang's laboratory were an integral part of helping me complete my doctoral research. I want to thank George Trevelyan, Mark Snyder, Mike Brody, Sherry Fitzsimmons, Jin-Hua Cheng, Art Hertz, Marilyn Bruce and Diane Aronstein. George Trevelyan was a good friend and colleague, and I will always remember those late nights working in the lab with him.

I thank all the professors, students, and staff at the Bodega Marine Laboratory, University of California, Davis for their help and friendship during my stay at the laboratory.