

AGE, GROWTH, AND RADIOMETRIC AGE
VALIDATION OF THE BLACKGILL ROCKFISH,
SEBASTES MELANOSTOMUS

A thesis submitted to the faculty of
San Francisco State University
In partial fulfillment of
The requirements for
The degree

Master of Science
In
Marine Science

By

Melissa Morgan Mahoney

San Francisco, California

December, 2002

ABSTRACT

As nearshore fish populations in the eastern Pacific Ocean become depleted, many commercial fishers shift their efforts toward deeper continental slope habitats to target fishes for which biological information is limited. One such fishery developed in the early 1980s for the blackgill rockfish, *Sebastes melanostomus*, a deep-dwelling (300-800 m) species that congregates over rocky pinnacles, mainly from southern California to southern Oregon. Growth zone-derived age estimates from otolith thin sections were compared to ages obtained from the radioactive disequilibria of ^{210}Pb , relative to its parent, ^{226}Ra , in otolith cores of blackgill rockfish collected off the U.S. west coast in 1985 and 1998-2000. Age estimates were validated up to at least age 41, with a strong pattern of agreement supporting longevity exceeding 90 years. Age and length data fitted to the von Bertalanffy growth function indicate *Sebastes melanostomus* is slow-growing ($k = 0.045$) and that females grow slower than males, but reach a larger asymptotic length. Estimates of age at 50% maturity, derived from previously published length-at-maturity estimates, are 17 years for males and 21 years for females. Results of this study agree with general life history traits already recognized for many *Sebastes* species, such as long life, slow growth, and late age at maturation. These traits may undermine the sustainability of blackgill rockfish populations when heavy fishing pressure, such as that which occurred in the 1980s, is applied.

ACKNOWLEDGEMENTS

We wish to thank John Butler, Don Pearson, and Cindy Taylor of the Southwest Fisheries Science Center, Mark Wilkins, Jerry Hoff, Waldo Wakefield and Bob Lauth of the Alaska Fisheries Science Center, and Bob Lea of CDFG for donating specimens for this study. Mary Yoklavich (NMFS), Di Tracey and Larry Paul (NIWA – New Zealand), Kristen Munk (ADFG), Don Pearson (NMFS), Tom Laidig (NMFS), and Steve Campana (DFO – Canada) provided valuable insight into blackgill rockfish growth patterns. Patrick McDonald of the Oregon Department of Fish and Wildlife aged otolith sections. Pete Holden at University of California, Santa Cruz and Dr. Craig Lundstrom of the University of Illinois measured radium in the refined samples using TIMS. This work was supported by the National Sea Grant College Program of the U.S. Department of Commerce's National Oceanic and Atmospheric Administration under NOAA Grant number NA06RG0142, project number R/F-182, through the California Sea Grant College Program; and in part by the California State Resources Agency. The views expressed herein do not necessarily reflect the views of any of those organizations. The U.S. government is authorized to reproduce and distribute for governmental purposes.