

HABITAT-DEPENDENT RECRUITMENT OF TWO TEMPERATE REEF
FISHES AT MULTIPLE SPATIAL SCALES

A Thesis

Presented to the

Faculty of

San Diego State University

In Partial Fulfillment

of the Requirements for the Degree

Master of Science

in

Biology

by

Kelly S. Andrews

Spring 2003

ABSTRACT

The distribution and abundance of reef fishes often has been attributed to several processes that result in a measure of recruitment success. In this study, I employ a large-scale experimental reef system to examine patterns of recruitment of two rocky reef fishes, the California sheephead (*Semicossyphus pulcher*) and the blackeye goby (*Rhinogobiops nicholsii*). The experimental system covers 3.3 km of coastline and consists of seven blocks of eight reefs, each reef 1600 m² in area. I examined three reefs per block that represented low, medium, and high percentage cover of hard substrate habitat in the fall of 2001 and 2002 for newly recruited California sheephead and blackeye goby. Among different treatments of habitat coverage, recruitment of California sheephead was significantly higher on reefs of medium coverage than for other coverages, while the blackeye goby exhibited lower recruitment on reefs of low coverage than for other coverages. Within reefs, recruitment to "edge" (ecotone) habitat was significantly lower than "inside" the reef for each species in 2001. At the smallest scale, several measures of microhabitat structure were quantified, and rugosity was important in predicting the presence of a recruit for each species at this small scale.

The densities of recruits of California sheephead corresponded to the densities of age 1+ individuals the following year, suggesting that the spatial patterns of abundance among treatments of habitat coverage may be established very early in life, and the abundance of predators does not appear to influence the patterns observed in any survey. Very low densities of recruits for each species may have led to habitat-dependent patterns of recruitment through habitat selection at settlement and density-independent mortality because of the very low densities of recruits observed here. Longer temporal studies with variable recruitment are needed to determine the importance of habitat structure relative to other processes.

ACKNOWLEDGMENTS

I thank my thesis advisor, Todd Anderson, for the years of mentoring, advising, and providing stimulating thought on ecological issues of importance. Many thanks go to Dane Bowker for his more than able diving assistance while working at SCAR, and to Dan Reed, Steve Schroeter, and the UCSB divers for information about the reefs and visibility updates. Also, I thank Colin Brauner and Joe Mahaffy for offering critical reviews of this manuscript and my fellow lab mates and other diving assistants for their efforts. Finally, I thank my wife, Megan, for dealing graciously with my obsession with this research. This publication was supported in part by the National Sea Grant College Program of the U.S. Department of Commerce's National Oceanic and Atmospheric Administration under NOAA Grant # NA06RG0142, project # R/F-188 (to T. Anderson), through the California Sea Grant College Program; and in part by the California State Resources Agency. Funding was also provided by the Department of Biology at San Diego State University through the Mabel Myers Scholarship to K. Andrews. The views expressed herein do not necessarily reflect the views of any of those organizations.