

CIRCULATING COPY

UNIVERSITY OF CALIFORNIA
Santa Barbara

Mechanism of Iron Acquisition by the Marine Bacterium *Alteromonas*
luteoviolacea

A Dissertation submitted in partial satisfaction
of the requirements for the Degree of

Doctor of Philosophy

in

Chemistry

by

Richard Thomas Reid

1994

LOAN COPY ONLY

Committee in charge:

Professor Alison Butler, Chairperson

Professor Peter Ford

Professor Thomas Hooker

Professor Stanley Parsons

ABSTRACT

Mechanisms of Iron Acquisition by the Marine Bacterium

Alteromonas luteoviolacea

by

Richard Thomas Reid

Marine bacteria exist in an environment that can be extremely iron poor. The levels of iron found in the open ocean are many orders of magnitude lower than that needed to produce normal growth under laboratory culture conditions, yet very little is known about the mechanisms by which marine bacteria may sequester iron. Terrestrial and enteric bacteria have been shown to produce a class of compound called siderophores when grown under conditions of iron limitation. The function of the siderophores is to solubilize and chelate any iron sources, and to make the metal available to the cell.

The marine bacterium *Alteromonas luteoviolacea*, was investigated in an attempt to understand what mechanisms this bacterium may have evolved to adjust to low iron levels in the oceanic environment. Two siderophores, alterobactin A and

B, were isolated from the spent cell culture supernatants of bacteria grown under conditions of iron limitation. The compounds were purified using a combination of solid phase extraction, hydrophobic interaction and gel filtration chromatography. The structures of alterobactins A and B were determined using mass spectral, amino acid and NMR analyses. Alterobactin B is comprised of N⁸-(2,3-dihydroxybenzoyl-4,8-diamino-3-hydroxyoctanoyl-seryl-glycyl-aspartyl-(β-hydroxyaspartyl)-glycyl-(β-hydroxyaspartate). Alterobactin A is the corresponding depsipeptide formed from the condensation between the serine hydroxyl group and the terminal carboxylate. The lactone ring of alterobactin A is unstable, and undergoes a base catalyzed hydrolysis to form Alterobactin B with a half-life of 11.6 hours at pH 8.0.

Alterobactin A coordinates iron(III) with a stoichiometry of 1:1 and an estimated β_{110} (the proton independent stability constant) of 10^{48} . Alterobactin B coordinates iron(III) with a 2:1 ligand:iron(III) stoichiometry. The presence of both compounds in the cell culture supernatant is regulated by exogenous iron; the synthesis of both compounds is repressed under high iron conditions. Alterobactin A and B both also function to transport iron into the cells, confirming their function as microbial siderophores.

Acknowledgements

I would like to thank all the people who have helped me through this process. My friends in the lab who kept me working and those outside of science who kept me sane. My parents and in-laws for their moral support for all these years. I would also like to acknowledge California Sea Grant trainee program for financial support of the project.

Finally and most importantly, I must thank my wife Lisa, for her unending love and support, without which I could have never completed this work.

Butler: R/MP-53