

Addressing Treatment Switching Bias with G-methods: Exploring the Impact of Model Specification

Amani Al Tawil^{*1,2}, Sean McGrath³, Robin Ristl^{†4}, and Ulrich Mansmann^{†1,2}

¹*Institute for Medical Information Processing, Biometry, and Epidemiology (IBE), Faculty of Medicine, Ludwig-Maximilians-Universität München*

²*Pettenkofer School of Public Health, Faculty of Medicine, Ludwig-Maximilians-Universität München*

³*Department of Biostatistics, Harvard T.H. Chan School of Public Health*

⁴*Center for Medical Data Science, Medical University of Vienna*

Electronic Supplementary Material 4

Covariates used in the inverse probability of censoring weights and parametric gformula methods

^{*}Correspondence: altawil@ibe.med.uni-muenchen.de

[†]Equally contributed

Tables

Table S3: Covariates used in the inverse probability of censoring weights and parametric gformula methods

	Covariates	Abbreviations
Baseline covariates	Age	AGE, AGE ² , AGE ^S
	Sex	SEX ₂
	Race	RACE ₂
	Initial diagnosis stage	IDS ₄
	Measurable intracranial Central Nervous System disease	ICS ₂
	Lung involvement at study entry	LI ₄
	ECOG score	ECOG ₂
	Strata at randomization*	ST ₄
	Smoking history	SM ₂
	Prior radiation therapy	RT ₂
Time-varying covariates	Follow-up time	FUT, FUT ² , FUT ^S
	ECOG score	ECOG ₂ , ECOG ₃
	Target-lesion size	TLS, TLS ² , TLS ^S
	Intracranial disease progression	IDP ₂
	Disease progression	DP ₂
	Time to disease progression	TDP, TDP ² , TDP ^S
	Treatment	A ₂

Numerical Variables are labeled as variablename^{*n*}, where *n* indicates the type of transformation applied:

1: linear term, 2: quadratic term, *S*: spline transformation.

Categorical Variables are labeled as variablename_{*n*}, where *n* indicates the number of categories.

*Strata at randomization include baseline brain metastases and previous chemotherapy