


# Multi-Resolution Analysis

---

Dr. Tushar Sandhan


# What does image convey

- 'A picture is worth a thousand words' – H. Ibsen (1828~1906)
  - info at various scales (resolutions)
  - vary the window size (W)
  - vary the image size itself & keeping W fixed




# What does image convey

- 'A picture is worth a thousand words' – H. Ibsen (1828~1906)
  - info at various scales (resolutions)
  - vary the window size (W)
  - vary the image size itself & keeping W fixed
- Larger objects can be analyzed @ \_\_\_\_\_ resolution



# What does image convey


- 'A picture is worth a thousand words' – H. Ibsen (1828~1906)
  - info at various scales (resolutions)
  - vary the window size (W)
  - vary the image size itself & keeping W fixed
- Larger objects can be analyzed @ \_\_\_\_\_ resolution
- Smaller ones can be analyzed @ \_\_\_\_\_ resolution



# Resolution relations

- Fourier analysis

- info abt freq content in a signal
- can we find out at what time the freq has occurred?
  - STFT (Gabor'1946)
  - if the time window is short: can assume stationarity
- can't locate sudden signal changes (edges)
- FT is ideal for stationary signals




upchirp


# Resolution relations

- Fourier analysis

- info abt freq content in a signal
- can we find out at what time the freq has occurred?
  - STFT (Gabor'1946)
  - if the time window is short: can assume stationarity
- can't locate sudden signal changes (edges)
- FT is ideal for stationary signals



upchirp





downchirp




# Resolution relations

## ■ Fourier analysis

- info abt freq content in a signal
- can we find out at what time the freq has occurred?
  - STFT (Gabor'1946)
  - if the time window is short: can assume stationarity
- can't locate sudden signal changes (edges)
- FT is ideal for stationary signals



upchirp




downchirp

# What do we need?

---

- Capture incremental info in the signal
  - low resolution
  - mid res.
  - high res.






# What do we need?


---

- Capture incremental info in the signal
  - low resolution
  - mid res.
  - high res.
- Does the shape of objects change in an image?
  - car, person, building etc.



# What do we need?

- Capture incremental info in the signal
  - low resolution
  - mid res.
  - high res.
- Does the shape of objects change in an image?
  - car, person, building etc.
- Do high freq. components always contain useful info?
  - noisy image
  - high freq is good near abrupt transitions (e.g. edgy regions)



# MRA: Multi-Resolution Analysis

---

- Find a space which can be fragmented in to subspaces
  - basis should be independent
  - basis functions: orthonormal
  - a function in  $L^2$  to control scale (freq): ( $s$  or  $n$ ) and space (time): ( $\tau$  or  $m$ )

$$\psi \in L^2 \longrightarrow \psi_{m,n}$$

# MRA: Multi-Resolution Analysis

---

- Find a space which can be fragmented in to subspaces
  - basis should be independent
  - basis functions: orthonormal
  - a function in  $L^2$  to control scale (freq): ( $s$  or  $n$ ) and space (time): ( $\tau$  or  $m$ )

$$\psi \in L^2 \longrightarrow \psi_{m,n}$$

$$(\psi_{m,n}, \psi_{k,l}) = \int_{-\infty}^{\infty} \psi_{m,n}(x) \psi_{k,l}(x) dx = \delta_{m,k} \delta_{n,l}$$

# MRA: Multi-Resolution Analysis

---

- Find a space which can be fragmented in to subspaces
  - basis should be independent
  - basis functions: orthonormal
  - a function in  $L^2$  to control scale (freq): ( $s$  or  $n$ ) and space (time): ( $\tau$  or  $m$ )

$$\psi \in L^2 \longrightarrow \psi_{m,n}$$

$$(\psi_{m,n}, \psi_{k,l}) = \int_{-\infty}^{\infty} \psi_{m,n}(x) \psi_{k,l}(x) dx = \delta_{m,k} \delta_{n,l}$$

$$\delta_{ij} = \begin{cases} 0 & i \neq j \\ 1 & i = j \end{cases} \quad m, n, i, j \text{ are integers}$$

# MRA: Multi-Resolution Analysis

---

- Find a space which can be fragmented in to subspaces
  - basis should be independent
  - basis functions: orthonormal
  - a function in  $L^2$  to control scale (freq): ( $s$  or  $n$ ) and space (time): ( $\tau$  or  $m$ )

$$\psi \in L^2 \longrightarrow \psi_{m,n}$$

$$(\psi_{m,n}, \psi_{k,l}) = \int_{-\infty}^{\infty} \psi_{m,n}(x) \psi_{k,l}(x) dx = \delta_{m,k} \delta_{n,l}$$

$$f_1(x) = x^2$$

$$f_2(x) = x^3$$

$$\delta_{ij} = \begin{cases} 0 & i \neq j \\ 1 & i = j \end{cases} \quad m, n, i, j \text{ are integers}$$

# MRA: Multi-Resolution Analysis

---

- Find a space which can be fragmented in to subspaces
  - basis should be independent
  - basis functions: orthonormal
  - a function in  $L^2$  to control scale (freq): ( $s$  or  $n$ ) and space (time): ( $\tau$  or  $m$ )

$$\psi \in L^2 \longrightarrow \psi_{m,n}$$

$$(\psi_{m,n}, \psi_{k,l}) = \int_{-\infty}^{\infty} \psi_{m,n}(x) \psi_{k,l}(x) dx = \delta_{m,k} \delta_{n,l}$$

$$\delta_{ij} = \begin{cases} 0 & i \neq j \\ 1 & i = j \end{cases} \quad m, n, i, j \text{ are integers}$$

$$f_1(x) = x^2$$

$$f_2(x) = x^3$$

$$(f_1, f_2) = \int_{-1}^1 x^2 \cdot x^3 dx$$



# MRA: represent a function

---

- Wavelet is the unified parametrized family of basis functions
  - wavelet  $\psi$  is called orthogonal, if the family of functions  $\psi_{m,n}$  are orthogonal
  - then a given signal can be uniquely expressed as sum of infinite series:

$$f(t) = \sum_{n,m=-\infty}^{\infty} c_{m,n} \psi_{m,n}(t)$$

# MRA: represent a function

---

- Wavelet is the unified parametrized family of basis functions
  - wavelet  $\psi$  is called orthogonal, if the family of functions  $\psi_{m,n}$  are orthogonal
  - then a given signal can be uniquely expressed as sum of infinite series:

$$f(t) = \sum_{n,m=-\infty}^{\infty} c_{m,n} \psi_{m,n}(t)$$

$$c_{m,n} = (f, \psi_{m,n}) \quad \dots \text{(from orthogonality)}$$

# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

1. Assimilation

- Subspaces can be embedded into each other

# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

Dyadic: resolution changes  
by a factor of 2 between spaces

1. Assimilation
  - Subspaces can be embedded into each other

# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

Dyadic: resolution changes  
by a factor of 2 between spaces

## 1. Assimilation

- Subspaces can be embedded into each other

$$V_0 \subset V_1 \subset V_2 \cdots V_m \subset V_{m+1}$$

# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

Dyadic: resolution changes  
by a factor of 2 between spaces

## 1. Assimilation

- Subspaces can be embedded into each other

$$V_0 \subset V_1 \subset V_2 \cdots V_m \subset V_{m+1}$$

## 2. Completeness (full span)

- Union of all embedded subspaces spans the entire  $L^2$

# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

Dyadic: resolution changes  
by a factor of 2 between spaces

## 1. Assimilation

- Subspaces can be embedded into each other

$$V_0 \subset V_1 \subset V_2 \cdots V_m \subset V_{m+1}$$

## 2. Completeness (full span)

- Union of all embedded subspaces spans the entire  $L^2$

$$\bigcup_{m=-\infty}^{\infty} V_m \quad \text{is dense in } L^2$$



# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

## 3. No redundancy

- Intersection of subspaces is a trivial set

# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

## 3. No redundancy

- Intersection of subspaces is a trivial set

$$\bigcap_{m=-\infty}^{\infty} V_m \text{ is trivial set } \{0\}$$

# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

## 3. No redundancy

- Intersection of subspaces is a trivial set

$$\bigcap_{m=-\infty}^{\infty} V_m \text{ is trivial set } \{0\}$$

## 4. Consistency in resolution of $f(t)$

- Superspace and subspace should maintain consistency of resolutions of input function  $f(t)$

# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

## 3. No redundancy

- Intersection of subspaces is a trivial set

$$\bigcap_{m=-\infty}^{\infty} V_m \text{ is trivial set } \{0\}$$

## 4. Consistency in resolution of $f(t)$

- Superspace and subspace should maintain consistency of resolutions of input function  $f(t)$

$$f(t) \in V_m$$

$$\text{iff } f(2t) \in V_{m+1}, \forall m \in I$$

# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

## 5. Info or energy preservation

- $\exists$  a function  $\phi \in V_0$  such that  $\{ \phi_{0,n}(x) = \phi(x - n) \}$  is orthogonal basis for  $V_0$  so that

- $\phi$  is called scaling func (or father wavelet)
- $\phi$  when contracted, then its integer translates can span any  $V_m$ :

$$V_m = \text{span}\{ \phi(2^m x - n) \}$$

# MRA: space properties

---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

## 5. Info or energy preservation

- $\exists$  a function  $\phi \in V_0$  such that  $\{\phi_{0,n}(x) = \phi(x - n)\}$  is orthogonal basis for  $V_0$  so that

$$\|f\|^2 = \int_{-\infty}^{\infty} |f(x)|^2 dx = \sum_{-\infty}^{\infty} |(f, \phi_{0,n})|^2$$

- $\phi$  is called scaling func (or father wavelet)
- $\phi$  when contracted, then its integer translates can span any  $V_m$ :

$$V_m = \text{span}\{\phi(2^m x - n)\}$$



# MRA: space properties


---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

## 5. Info or energy preservation

- $\exists$  a function  $\phi \in V_0$  such that  $\{\phi_{0,n}(x) = \phi(x - n)\}$  is orthogonal basis for  $V_0$  so that

$$\|f\|^2 = \int_{-\infty}^{\infty} |f(x)|^2 dx = \sum_{-\infty}^{\infty} |(f, \phi_{0,n})|^2$$



- $\phi$  is called scaling func (or father wavelet)
- $\phi$  when contracted, then its integer translates can span any  $V_m$ :

$$V_m = \text{span}\{\phi(2^m x - n)\}$$

# MRA: space properties


---

- sequence of embedded subspaces of  $L^2$  :  $\{V_m\}$  where  $m \in I$  (integers)

## 5. Info or energy preservation

- $\exists$  a function  $\phi \in V_0$  such that  $\{\phi_{0,n}(x) = \phi(x - n)\}$  is orthogonal basis for  $V_0$  so that

$$\|f\|^2 = \int_{-\infty}^{\infty} |f(x)|^2 dx = \sum_{-\infty}^{\infty} |(f, \phi_{0,n})|^2$$



- $\phi$  is called scaling func (or father wavelet)
- $\phi$  when contracted, then its integer translates can span any  $V_m$ :

$$V_m = \text{span}\{\phi(2^m x - n)\}$$

- $\phi$  generates MRA: if  $\{V_m\}$  is a multiresolution of  $L^2$  and if  $V_0$  is closed subspace generated by translates of a single function  $\phi$

# MRA: refinement equation

---

- we have to reflect  $\phi(x)$  in to the higher resolution or refined space  $V_1$ 
  - high resolution is with finer scale factor of  $2^{-1}$
- $\phi$  is scaling func for  $V_0$  and since  $V_0 \subset V_1$ 
  - it becomes scaling func for  $V_1$

# MRA: refinement equation

---

- we have to reflect  $\phi(x)$  in to the higher resolution or refined space  $V_1$ 
  - high resolution is with finer scale factor of  $2^{-1}$
- $\phi$  is scaling func for  $V_0$  and since  $V_0 \subset V_1$ 
  - it becomes scaling func for  $V_1$

$$\phi(x) = \sum_{-\infty}^{\infty} c_n \phi_{1,n}(x) = \sqrt{2} \sum_{-\infty}^{\infty} c_n \phi(2x - n)$$

# MRA: refinement equation

---

- we have to reflect  $\phi(x)$  in to the higher resolution or refined space  $V_1$ 
  - high resolution is with finer scale factor of  $2^{-1}$
- $\phi$  is scaling func for  $V_0$  and since  $V_0 \subset V_1$ 
  - it becomes scaling func for  $V_1$

$$\phi(x) = \sum_{-\infty}^{\infty} c_n \phi_{1,n}(x) = \sqrt{2} \sum_{-\infty}^{\infty} c_n \phi(2x - n)$$

$$c_n = (\phi, \phi_{1,n})$$

# MRA: refinement equation

---

- we have to reflect  $\phi(x)$  in to the higher resolution or refined space  $V_1$ 
  - high resolution is with finer scale factor of  $2^{-1}$
- $\phi$  is scaling func for  $V_0$  and since  $V_0 \subset V_1$ 
  - it becomes scaling func for  $V_1$

$$\phi(x) = \sum_{-\infty}^{\infty} c_n \phi_{1,n}(x) = \sqrt{2} \sum_{-\infty}^{\infty} c_n \phi(2x - n)$$

$$c_n = (\phi, \phi_{1,n}) \qquad \sum_{-\infty}^{\infty} |c_n|^2 = 1$$

# Scaling function in orthogonal complement?

---

- If  $U$  &  $W$  are subspaces then their sum is the subspace

$$U + W = \{\mathbf{u} + \mathbf{w} : \mathbf{u} \in U, \mathbf{w} \in W\}$$

- The direct sum is the sum of independent subspaces

$$U \oplus W$$



# Scaling function in orthogonal complement?

---

- If  $U$  &  $W$  are subspaces then their sum is the subspace

$$U + W = \{\mathbf{u} + \mathbf{w} : \mathbf{u} \in U, \mathbf{w} \in W\}$$

- The direct sum is the sum of independent subspaces

$$U \oplus W$$

- $V_m \subset V_{m+1}$  thus we can define orthogonal complement of  $V_m$  in  $V_{m+1}$  as

# Scaling function in orthogonal complement?

---

- If  $U$  &  $W$  are subspaces then their sum is the subspace

$$U + W = \{\mathbf{u} + \mathbf{w} : \mathbf{u} \in U, \mathbf{w} \in W\}$$

- The direct sum is the sum of independent subspaces

$$U \oplus W$$

- $V_m \subset V_{m+1}$  thus we can define orthogonal complement of  $V_m$  in  $V_{m+1}$  as

$$V_{m+1} = V_m \oplus W_m$$

# Scaling function in orthogonal complement?

---

- If  $U$  &  $W$  are subspaces then their sum is the subspace

$$U + W = \{\mathbf{u} + \mathbf{w} : \mathbf{u} \in U, \mathbf{w} \in W\}$$

- The direct sum is the sum of independent subspaces

$$U \oplus W$$

- $V_m \subset V_{m+1}$  thus we can define orthogonal complement of  $V_m$  in  $V_{m+1}$  as

$$V_{m+1} = V_m \oplus W_m \qquad V_m \perp W_m$$

# Orthogonal complementary space

---

$$\begin{aligned} V_{m+1} &= V_m \oplus W_m \\ &= (V_{m-1} \oplus W_{m-1}) \oplus W_m \\ &\vdots \\ &= V_0 \oplus W_0 \oplus W_1 \oplus \dots \oplus W_m \\ &= V_0 \oplus \left( \bigoplus_{n=0}^m W_n \right) . \end{aligned}$$

# Orthogonal complementary space

---

$$\begin{aligned} V_{m+1} &= V_m \oplus W_m \\ &= (V_{m-1} \oplus W_{m-1}) \oplus W_m \\ &\vdots \\ &= V_0 \oplus W_0 \oplus W_1 \oplus \dots \oplus W_m \\ &= V_0 \oplus \left( \bigoplus_{n=0}^m W_n \right) . \end{aligned}$$

$$m \rightarrow \infty$$

# Orthogonal complementary space

---

$$\begin{aligned} V_{m+1} &= V_m \oplus W_m \\ &= (V_{m-1} \oplus W_{m-1}) \oplus W_m \\ &\vdots \\ &= V_0 \oplus W_0 \oplus W_1 \oplus \cdots \oplus W_m \\ &= V_0 \oplus \left( \bigoplus_{n=0}^m W_n \right) . \end{aligned}$$

$$m \rightarrow \infty$$

$$V_0 \oplus \left( \bigoplus_{n=0}^{\infty} W_n \right) = L^2$$

# Orthogonal complementary space

---

$$\begin{aligned} V_{m+1} &= V_m \oplus W_m \\ &= (V_{m-1} \oplus W_{m-1}) \oplus W_m \\ &\vdots \\ &= V_0 \oplus W_0 \oplus W_1 \oplus \dots \oplus W_m \\ &= V_0 \oplus \left( \bigoplus_{n=0}^m W_n \right) . \end{aligned}$$

$$m \rightarrow \infty$$

$$V_0 \oplus \left( \bigoplus_{n=0}^{\infty} W_n \right) = L^2$$

$V_0$  is just a reference space  
(reference can be any index, i.e.  
resolution can scale-up or down)

$$\bigoplus_{n=-\infty}^{\infty} W_n = L^2$$

# MRA: mother wavelet

---

- complementary space also spans  $L^2$ 
  - $\exists$  a function  $\psi \in W_0$  such that  $\{ \psi_{0,n}(x) = \psi(x - n) \}$  is orthogonal basis for  $W_0$
  - Orthogonal basis for  $W_m$  are given as:

$$\psi_{m,n}(x) = 2^{m/2} \psi(2^m x - n)$$

- $\psi$  is called a mother wavelet



# MRA: mother wavelet

---

- complementary space also spans  $L^2$ 
  - $\exists$  a function  $\psi \in W_0$  such that  $\{ \psi_{0,n}(x) = \psi(x - n) \}$  is orthogonal basis for  $W_0$
  - Orthogonal basis for  $W_m$  are given as:

$$\psi_{m,n}(x) = 2^{m/2} \psi(2^m x - n)$$

- $\psi$  is called a mother wavelet
- $\psi$  spans orthogonal complement subset  $W_m$ , while scaling function  $\phi$  spans the subsets  $V_m$

# MRA: WT

---

- wavelet

$$\psi_{a,b}(t) = \frac{1}{\sqrt{b}}\psi\left(\frac{t-a}{b}\right)$$

- decomposition

# MRA: WT

---

- wavelet

$$\psi_{a,b}(t) = \frac{1}{\sqrt{b}} \psi\left(\frac{t-a}{b}\right)$$

- decomposition

$$W(a,b) = K \int_{-\infty}^{+\infty} \psi^*\left(\frac{x-a}{b}\right) f(x) dx$$


# Haar

---

- Haar wavelet

# Haar


- Haar wavelet



# Haar

- Haar wavelet


$$\psi(t) = \begin{cases} 1 & 0 \leq t < \frac{1}{2}, \\ -1 & \frac{1}{2} \leq t < 1, \\ 0 & \text{otherwise.} \end{cases}$$




# Haar

- Haar wavelet

$$\psi(t) = \begin{cases} 1 & 0 \leq t < \frac{1}{2}, \\ -1 & \frac{1}{2} \leq t < 1, \\ 0 & \text{otherwise.} \end{cases}$$



$$\psi_{m,n}(x) = 2^{m/2} \psi(2^m x - n)$$



# Conclusion

- MRA



# Conclusion

- MRA

□ Image analysis in multiple resolutions

□ MRA

- Spaces
- Basis functions
- Wavelets