



University of  
**Nottingham**  
UK | CHINA | MALAYSIA

Autumn 2018

# Econometric Theory I

## Computer Lab Class III

Juergen Amann

[juergen.amann@nottingham.ac.uk](mailto:juergen.amann@nottingham.ac.uk)

Wednesday 12:00 - 13:00, C42 SCGB

## Last week

Two exercises:

- **Exercise 1:** Examine the evidence of a gender pay gap.
  - Dummy variables.
  - What does it mean when we '*control for (a) variable(s)*'?
- **Exercise 2:** Examine the determinants of educational attainment.
  - Joint parameter hypothesis testing of coefficients; e.g.  
 $H_0 : \beta_2 = \beta_3$ .
  - Restricted Least Squares.

## This week

Model the birth weights of infants:

- Identify factors that may determine a child's birth weight.
- Interpret Stata output.
- One- vs. two-sided hypothesis testing.
- Over- and under-specification of an econometric model.
- Homoskedasticity vs. heteroskedasticity.

## This week

- What are the factors that determine a child's birth weight?

## This week

- What are the factors that determine a child's birth weight?
- Our data set `bwght.xls` contains  $n = 1,388$  observations on the following variables:
  - `bwghti`: birth weight in ounces
  - `faminci`: family income in USD 1,000
  - `orderi`: birth order of child
  - `cigsi`: cigarettes smoked per day by mother during pregnancy
  - `motheduci`: number of years mother spent in full-time education

## This week

- What are the factors that determine a child's birth weight?
- Our data set `bwght.xls` contains  $n = 1,388$  observations on the following variables:
  - `bwghti`: birth weight in ounces
  - `faminci`: family income in USD 1,000
  - `orderi`: birth order of child
  - `cigsi`: cigarettes smoked per day by mother during pregnancy
  - `motheduci`: number of years mother spent in full-time education
- What is the expected sign for the variables? Which variables do you expect to be significant?

## Birth weight: regression 1

- Download `bwght.xls` and import it into Stata.

## Birth weight: regression 1

- Download `bwght.xls` and import it into Stata.
- Estimate a regression of `birth weight` on a `constant`, `family income` and `birth order`.



## Birth weight: regression 1

- Download `bwght.xls` and import it into Stata.
- Estimate a regression of `birth weight` on a `constant`, `family income` and `birth order`.

```
. regress bwght faminc order
```

## Birth weight: regression 1

- Download `bwght.xls` and import it into Stata.
- Estimate a regression of `birth weight` on a `constant`, `family income` and `birth order`.

```
. regress bwght faminc order
```

Source	SS	df	MS	Number of obs	=	1,388
Model	9103.2286	2	4551.6143	F(2, 1385)	=	11.15
Residual	565508.491	1,385	408.30938	Prob > F	=	0.0000
				R-squared	=	0.0158
				Adj R-squared	=	0.0144
Total	574611.72	1,387	414.283864	Root MSE	=	20.207

	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
bwght						
faminc	.1235904	.0290392	4.26	0.000	.0666249	.1805559
order	1.439648	.6086762	2.37	0.018	.2456214	2.633675
_cons	112.7618	1.45622	77.43	0.000	109.9052	115.6185

- Test if variables are significant, interpret signs and magnitudes.

## Birth weight: regression 1

Results of regression 1:

- `_cons`: For `order = faminc = 0`, the average expected birth weight is 112.76 ounces, i.e.  $\approx 3.18\text{kg}$ .

## Birth weight: regression 1

Results of regression 1:

- `_cons`: For `order = faminc = 0`, the average expected birth weight is 112.76 ounces, i.e.  $\approx 3.18kg$ .
- `faminc`: Keeping `order` constant, a USD 1,000 increase in family income corresponds to an increase of expected birth weight of 0.12 ounces, i.e.  $\approx 3.4g$ .

## Birth weight: regression 1

### Results of regression 1:

- `_cons`: For `order = faminc = 0`, the average expected birth weight is 112.76 ounces, i.e.  $\approx 3.18kg$ .
- `faminc`: Keeping `order` constant, a USD 1,000 increase in family income corresponds to an increase of expected birth weight of 0.12 ounces, i.e.  $\approx 3.4g$ .
- `order`: For the same family income, the next-born child is expected to be approx. 1.14 ounces ( $\approx 32.4g$ ) heavier than his/her sibling born right before him/her.

## Birth weight: regression 1

### Results of regression 1:

- `_cons`: For `order = faminc = 0`, the average expected birth weight is 112.76 ounces, i.e.  $\approx 3.18kg$ .
- `faminc`: Keeping `order` constant, a USD 1,000 increase in family income corresponds to an increase of expected birth weight of 0.12 ounces, i.e.  $\approx 3.4g$ .
- `order`: For the same family income, the next-born child is expected to be approx. 1.14 ounces ( $\approx 32.4g$ ) heavier than his/her sibling born right before him/her.

## Birth weight: regression 1

### Results of regression 1:

- `_cons`: For `order = faminc = 0`, the average expected birth weight is 112.76 ounces, i.e.  $\approx 3.18kg$ .
- `faminc`: Keeping `order` constant, a USD 1,000 increase in family income corresponds to an increase of expected birth weight of 0.12 ounces, i.e.  $\approx 3.4g$ .
- `order`: For the same family income, the next-born child is expected to be approx. 1.14 ounces ( $\approx 32.4g$ ) heavier than his/her sibling born right before him/her.
- All coefficients are found to be highly significant.

## Birth weight: regression 1

- Look at the test for overall relation.



## Birth weight: regression 1

- Look at the test for overall relation.

Source	SS	df	MS	Number of obs	=	1,388
Model	9103.2286	2	4551.6143	F(2, 1385)	=	11.15
Residual	565508.491	1,385	408.30938	Prob > F	=	0.0000
				R-squared	=	0.0158
				Adj R-squared	=	0.0144
Total	574611.72	1,387	414.283864	Root MSE	=	20.207

- Remember [Slides 4, p.10]: *'One useful special case of the F-test is in testing whether all the explanatory variables taken together are significant in the regression model.'*

## Birth weight: regression 1

- Look at the test for overall relation.

Source	SS	df	MS	Number of obs	=	1,388
Model	9103.2286	2	4551.6143	F(2, 1385)	=	11.15
Residual	565508.491	1,385	408.30938	Prob > F	=	0.0000
				R-squared	=	0.0158
				Adj R-squared	=	0.0144
Total	574611.72	1,387	414.283864	Root MSE	=	20.207

- Remember [Slides 4, p.10]: *'One useful special case of the F-test is in testing whether all the explanatory variables taken together are significant in the regression model.'*

For our multivariate regression

$$\text{bwght}_i = \beta_1 + \beta_2 \text{faminc}_i + \beta_3 \text{order}_i + u_i$$

this means  $H_0 : \beta_2 = \beta_3 = 0$ .

## Birth weight: regression 1

- Look at the test for overall relation.

Source	SS	df	MS	Number of obs	=	1,388
Model	9103.2286	2	4551.6143	F(2, 1385)	=	11.15
Residual	565508.491	1,385	408.30938	Prob > F	=	0.0000
Total	574611.72	1,387	414.283864	R-squared	=	0.0158
				Adj R-squared	=	0.0144
				Root MSE	=	20.207

- Remember [Slides 4, p.10]: *'One useful special case of the F-test is in testing whether all the explanatory variables taken together are significant in the regression model.'*

For our multivariate regression

$$\text{bwght}_i = \beta_1 + \beta_2 \text{faminc}_i + \beta_3 \text{order}_i + u_i$$

this means  $H_0 : \beta_2 = \beta_3 = 0$ .

- This test statistic follows an  $F_{(q, n-k)}$  distribution under  $H_0$  where  $q = 2$  (no. of tested parameters) and  $(n - k) = 1,385$ .

## Birth weight: regression 1

- Look at the test for overall relation.

Source	SS	df	MS	Number of obs	=	1,388
				F(2, 1385)	=	11.15
Model	9103.2286	2	4551.6143	Prob > F	=	0.0000
Residual	565508.491	1,385	408.30938	R-squared	=	0.0158
				Adj R-squared	=	0.0144
Total	574611.72	1,387	414.283864	Root MSE	=	20.207

- Remember [Slides 4, p.10]: *'One useful special case of the F-test is in testing whether all the explanatory variables taken together are significant in the regression model.'*

For our multivariate regression

$$\text{bwght}_i = \beta_1 + \beta_2 \text{faminc}_i + \beta_3 \text{order}_i + u_i$$

this means  $H_0 : \beta_2 = \beta_3 = 0$ .

- This test statistic follows an  $F_{(q, n-k)}$  distribution under  $H_0$  where  $q = 2$  (no. of tested parameters) and  $(n - k) = 1,385$ .
- Compare test statistic against critical value (and/or use p-value): We reject  $H_0$  here; variables are 'jointly significant'. ▶▶ Let's do the same 'by hand'!

## Birth weight: regression 1

- Interpret the **goodness-of-fit** of the regression, i.e. the  $R^2$ .

## Birth weight: regression 1

- Interpret the **goodness-of-fit** of the regression, i.e. the  $R^2$ .

Source	SS	df	MS	Number of obs	=	1,388
Model	9103.2286	2	4551.6143	F(2, 1385)	=	11.15
Residual	565508.491	1,385	408.30938	Prob > F	=	0.0000
				R-squared	=	0.0158
				Adj R-squared	=	0.0144
Total	574611.72	1,387	414.283864	Root MSE	=	20.207

- See Slides 4, p.19-21 for more details.

## Birth weight: regression 1

- Interpret the **goodness-of-fit** of the regression, i.e. the  $R^2$ .

Source	SS	df	MS	Number of obs	=	1,388
				F(2, 1385)	=	11.15
Model	9103.2286	2	4551.6143	Prob > F	=	0.0000
Residual	565508.491	1,385	408.30938	R-squared	=	0.0158
				Adj R-squared	=	0.0144
Total	574611.72	1,387	414.283864	Root MSE	=	20.207

- See Slides 4, p.19-21 for more details.
- Main takeaway here: Our model fits the data only very poorly (very low  $R^2$ ).

## Birth weight: regression 1

- Interpret the **goodness-of-fit** of the regression, i.e. the  $R^2$ .

Source	SS	df	MS	Number of obs	=	1,388
				F(2, 1385)	=	11.15
Model	9103.2286	2	4551.6143	Prob > F	=	0.0000
Residual	565508.491	1,385	408.30938	R-squared	=	0.0158
				Adj R-squared	=	0.0144
Total	574611.72	1,387	414.283864	Root MSE	=	20.207

- See Slides 4, p.19-21 for more details.
- Main takeaway here: Our model fits the data only very poorly (very low  $R^2$ ).
- Only about 1.58% of the variation of the dependent variable about its mean is explained by **regression 1**.



## Birth weight: regression 1

- Interpret the **goodness-of-fit** of the regression, i.e. the  $R^2$ .

Source	SS	df	MS	Number of obs	=	1,388
				F(2, 1385)	=	11.15
Model	9103.2286	2	4551.6143	Prob > F	=	0.0000
Residual	565508.491	1,385	408.30938	R-squared	=	0.0158
				Adj R-squared	=	0.0144
Total	574611.72	1,387	414.283864	Root MSE	=	20.207

- See Slides 4, p.19-21 for more details.
- Main takeaway here: Our model fits the data only very poorly (very low  $R^2$ ).
- Only about 1.58% of the variation of the dependent variable about its mean is explained by **regression 1**.
- As before, we can calculate the  $R^2$  using other parts of the regression output. [▶ See how it's done!](#)

## Birth weight: regression 2

- Estimate a regression of birth weight on a constant, family income, birth order and **cigs**.

## Birth weight: regression 2

- Estimate a regression of birth weight on a constant, family income, birth order and cigs.

```
. regress bwght faminc order cigs
```

## Birth weight: regression 2

- Estimate a regression of birth weight on a constant, family income, birth order and **cigs**.

```
. regress bwght faminc order cigs
```

Source	SS	df	MS	Number of obs	=	1,388
				F(3, 1384)	=	16.63
Model	19996.5211	3	6665.50703	Prob > F	=	0.0000
Residual	554615.199	1,384	400.733525	R-squared	=	0.0348
				Adj R-squared	=	0.0327
Total	574611.72	1,387	414.283864	Root MSE	=	20.018

  

bwght	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
faminc	.0979201	.0291868	3.35	0.001	.040665	.1551752
order	1.616372	.603955	2.68	0.008	.4316058	2.801138
cigs	-.4771537	.091518	-5.21	0.000	-.6566827	-.2976247
_cons	114.2143	1.4693	77.73	0.000	111.3321	117.0966

- Observe how the regression output has changed!

## Birth weight: regression 2

- Estimate a regression of birth weight on a constant, family income, birth order and **cigs**.

```
. regress bwght faminc order cigs
```

Source	SS	df	MS	Number of obs	=	1,388
Model	19996.5211	3	6665.50703	F(3, 1384)	=	16.63
Residual	554615.199	1,384	400.733525	Prob > F	=	0.0000
Total	574611.72	1,387	414.283864	R-squared	=	0.0348
				Adj R-squared	=	0.0327
				Root MSE	=	20.018

  

bwght	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
faminc	.0979201	.0291868	3.35	0.001	.040665	.1551752
order	1.616372	.603955	2.68	0.008	.4316058	2.801138
cigs	-.4771537	.091518	-5.21	0.000	-.6566827	-.2976247
_cons	114.2143	1.4693	77.73	0.000	111.3321	117.0966

- Observe how the regression output has changed!
- Conduct a 1-sided test on the **cigs** coefficient using statistical tables!

## Birth weight: regression 2

One- vs two-sided tests?

- Two-sided:
  - Testing for the possibility of the relationship in both directions.

## Birth weight: regression 2

One- vs two-sided tests?

- Two-sided:
  - Testing for the possibility of the relationship in both directions.
  - IOW A two-sided test will test both if the coefficient is significantly bigger **and** significantly smaller than zero.

## Birth weight: regression 2

One- vs two-sided tests?

- Two-sided:
  - Testing for the possibility of the relationship in both directions.
  - IOW A two-sided test will test both if the coefficient is significantly bigger **and** significantly smaller than zero.
  - Given a fixed significance level of, say,  $\alpha = 0.05$ , a two-sided test allots half of our  $\alpha$  to testing the statistical significance in *one* and the other half of  $\alpha$  to the other direction.



## Birth weight: regression 2

### One- vs two-sided tests?

- Two-sided:
  - Testing for the possibility of the relationship in both directions.
  - IOW A two-sided test will test both if the coefficient is significantly bigger **and** significantly smaller than zero.
  - Given a fixed significance level of, say,  $\alpha = 0.05$ , a two-sided test allots half of our  $\alpha$  to testing the statistical significance in *one* and the other half of  $\alpha$  to the other direction.
  - This means `cigs` is considered significantly different from 0 if the test statistic is in the top 2.5% or bottom 2.5% of its probability distribution, resulting in a p-value less than 0.05.

## Birth weight: regression 2

One- vs two-sided tests?

- One-sided:
  - Tests either if the estimated parameter is significantly greater **or** significantly smaller than zero, but not both.

## Birth weight: regression 2

One- vs two-sided tests?

- One-sided:
  - Tests either if the estimated parameter is significantly greater **or** significantly smaller than zero, but not both.
  - IOW it tests for the possibility of the relationship in one direction and completely disregarding the other direction.

## Birth weight: regression 2

One- vs two-sided tests?

- One-sided:
  - Tests either if the estimated parameter is significantly greater **or** significantly smaller than zero, but not both.
  - IOW it tests for the possibility of the relationship in one direction and completely disregarding the other direction.
  - A one-sided test allots **all** of our  $\alpha$  to testing the statistical significance in *one* direction only.

## Birth weight: regression 2

### One- vs two-sided tests?

- One-sided:
  - Tests either if the estimated parameter is significantly greater **or** significantly smaller than zero, but not both.
  - IOW it tests for the possibility of the relationship in one direction and completely disregarding the other direction.
  - A one-sided test allots **all** of our  $\alpha$  to testing the statistical significance in *one* direction only.
  - This means that, depending on the chosen tail, *cigs* is considered significantly lower (higher) than 0 if the test statistic is in the bottom (top) 5% of its probability distribution, resulting in a p-value less than 0.05.

## Birth weight: regression 2

### One- vs two-sided tests?

- One-sided:
  - Tests either if the estimated parameter is significantly greater **or** significantly smaller than zero, but not both.
  - IOW it tests for the possibility of the relationship in one direction and completely disregarding the other direction.
  - A one-sided test allots **all** of our  $\alpha$  to testing the statistical significance in *one* direction only.
  - This means that, depending on the chosen tail, *cigs* is considered significantly lower (higher) than 0 if the test statistic is in the bottom (top) 5% of its probability distribution, resulting in a p-value less than 0.05.
  - The one-tailed test provides more power to detect an effect in one direction by not testing the effect in the other direction.

## Birth weight: regression 2

- One-sided test in our example

## Birth weight: regression 2

- One-sided test in our example

```
. regress bwght faminc order cigs
```



## Birth weight: regression 2

- One-sided test in our example

```
. regress bwght faminc order cigs
```

	bwght	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
faminc		.0979201	.0291868	3.35	0.001	.040665	.1551752
order		1.616372	.603955	2.68	0.008	.4316058	2.801138
cigs		-.4771537	.091518	-5.21	0.000	-.6566827	-.2976247
_cons		114.2143	1.4693	77.73	0.000	111.3321	117.0966

- Above  $H_0 : \beta_{cigs} = 0$ ,  $H_a : \beta_{cigs} \neq 0$  (two-sided).
- We want to test for the one-sided alternative that  $\beta_{cigs} < 0$ . Note that the effect has the correct 'predicted sign'.
- To get the p-value, simply divide  $p - Val_{cigs}/2$  because the effect is going in the 'predicted' direction. This is  $P(| - 5.21|)$ .
- We reject  $H_0$  in favour of the one-sided alternative indicating that smoking has a significantly negative impact on birth weight.

## Birth weight: regression 3

- Estimate a regression of birth weight on a constant, family income, birth order, cigs and mothereduc.

## Birth weight: regression 3

- Estimate a regression of birth weight on a constant, family income, birth order, cigs and mothereduc.

```
. regress bwght faminc order cigs mothereduc
```

## Birth weight: regression 3

- Estimate a regression of birth weight on a constant, family income, birth order, cigs and mothereduc.

```
. regress bwght faminc order cigs mothereduc
```

Source	SS	df	MS	Number of obs	=	1,388
Model	20031.4556	4	5007.8639	F(4, 1383)	=	12.49
Residual	554580.264	1,383	400.998022	Prob > F	=	0.0000
Total	574611.72	1,387	414.283864	R-squared	=	0.0349
				Adj R-squared	=	0.0321
				Root MSE	=	20.025

bwght	Coef.	Std. Err.	t	P> t	[95% Conf. Interval]	
faminc	.0937846	.0323843	2.90	0.004	.0302569	.1573122
order	1.631028	.6061915	2.69	0.007	.4418743	2.820183
cigs	-.4730827	.0925814	-5.11	0.000	-.6546978	-.2914676
mothereduc	.076161	.2580337	0.30	0.768	-.4300187	.5823408
_cons	113.317	3.376943	33.56	0.000	106.6925	119.9415

- Interpret results!

## Model specification

From the lecture [slides 4, pp. 14]:

- If an important explanatory variable is **missing** from our model, the estimated coefficients are **biased** (omitted variable bias)!

## Model specification

From the lecture [slides 4, pp. 14]:

- If an important explanatory variable is **missing** from our model, the estimated coefficients are **biased** (omitted variable bias)!
- If an irrelevant explanatory variable is included in our model, the estimated coefficients are **still unbiased** but **inefficient**, i.e. its variance is greater when the irrelevant variable is included.

## Model specification

From the lecture [slides 4, pp. 14]:

- If an important explanatory variable is **missing** from our model, the estimated coefficients are **biased** (omitted variable bias)!
- If an irrelevant explanatory variable is included in our model, the estimated coefficients are **still unbiased** but **inefficient**, i.e. its variance is greater when the irrelevant variable is included.

Which is the best specification for our problem?

## Model specification

From the lecture [slides 4, pp. 14]:

- If an important explanatory variable is **missing** from our model, the estimated coefficients are **biased** (omitted variable bias)!
- If an irrelevant explanatory variable is included in our model, the estimated coefficients are **still unbiased** but **inefficient**, i.e. its variance is greater when the irrelevant variable is included.

Which is the best specification for our problem?

- **regression 1**: low  $R^2$ , cigs missing; **probably under-specified**.



## Model specification

From the lecture [slides 4, pp. 14]:

- If an important explanatory variable is **missing** from our model, the estimated coefficients are **biased** (omitted variable bias)!
- If an irrelevant explanatory variable is included in our model, the estimated coefficients are **still unbiased** but **inefficient**, i.e. its variance is greater when the irrelevant variable is included.

Which is the best specification for our problem?

- **regression 1**: low  $R^2$ , cigs missing; **probably under-sepcified**.
- **regression 3**: mothereduc probably irrelevant; **probably over-sepcified**.

## Model specification

From the lecture [slides 4, pp. 14]:

- If an important explanatory variable is **missing** from our model, the estimated coefficients are **biased** (omitted variable bias)!
- If an irrelevant explanatory variable is included in our model, the estimated coefficients are **still unbiased** but **inefficient**, i.e. its variance is greater when the irrelevant variable is included.

Which is the best specification for our problem?

- **regression 1**: low  $R^2$ , cigs missing; **probably under-specified**.
- **regression 3**: mothereduc probably irrelevant; **probably over-specified**.
- **regression 2**: note that  $adj. R^2_{reg.3} < adj. R^2_{reg.2}$ ; **probably best fit**.

## Homo- vs. heteroskedasticity

See lecture [slides 4, pp. 37]:

## Homo- vs. heteroskedasticity

See lecture [slides 4, pp. 37]:

- **Homoskedasticity**: One of the classical assumptions of the linear regression model [Slides 3, p.3].

## Homo- vs. heteroskedasticity

See lecture [slides 4, pp. 37]:

- **Homoskedasticity:** One of the classical assumptions of the linear regression model [Slides 3, p.3].
- **Heteroskedasticity:** Errors have non-constant variances.

## Homo- vs. heteroskedasticity

See lecture [slides 4, pp. 37]:

- **Homoskedasticity:** One of the classical assumptions of the linear regression model [Slides 3, p.3].
- **Heteroskedasticity:** Errors have non-constant variances.
  - **Example:** More variation in consumption behaviour for rich compared to poor families.

## Homo- vs. heteroskedasticity

See lecture [slides 4, pp. 37]:

- **Homoskedasticity:** One of the classical assumptions of the linear regression model [Slides 3, p.3].
- **Heteroskedasticity:** Errors have non-constant variances.
  - **Example:** More variation in consumption behaviour for rich compared to poor families.
  - **Causes:** measurement error, sub-population differences, miss-specification (omitted variables) etc. (many causes possible).

## Homo- vs. heteroskedasticity

See lecture [slides 4, pp. 37]:

- **Homoskedasticity:** One of the classical assumptions of the linear regression model [Slides 3, p.3].
- **Heteroskedasticity:** Errors have non-constant variances.
  - **Example:** More variation in consumption behaviour for rich compared to poor families.
  - **Causes:** measurement error, sub-population differences, miss-specification (omitted variables) etc. (many causes possible).
  - **Consequences:** Coefficient estimates remain unbiased. However, OLS is no longer efficient (smallest variance) **and** estimated standard errors are biased.



## Homo- vs. heteroskedasticity

See lecture [slides 4, pp. 37]:

- **Homoskedasticity:** One of the classical assumptions of the linear regression model [Slides 3, p.3].
- **Heteroskedasticity:** Errors have non-constant variances.
  - **Example:** More variation in consumption behaviour for rich compared to poor families.
  - **Causes:** measurement error, sub-population differences, miss-specification (omitted variables) etc. (many causes possible).
  - **Consequences:** Coefficient estimates remain unbiased. However, OLS is no longer efficient (smallest variance) **and** estimated standard errors are biased.  
⇒ Hypothesis testing is invalid!

## Homo- vs. heteroskedasticity

- Re-run regression 2 and check for heteroskedasticity.

## Homo- vs. heteroskedasticity

- Re-run [regression 2](#) and check for heteroskedasticity.

```
. regress bwght faminc order cigs
```

## Homo- vs. heteroskedasticity

- Re-run [regression 2](#) and check for heteroskedasticity.

```
. regress bwght faminc order cigs
```

- Visual analysis first; e.g. for faminc

## Homo- vs. heteroskedasticity

- Re-run [regression 2](#) and check for heteroskedasticity.

```
. regress bwght faminc order cigs
```

- Visual analysis first; e.g. for faminc

```
. rvppplot faminc
```

## Homo- vs. heteroskedasticity

- Re-run [regression 2](#) and check for heteroskedasticity.

```
. regress bwght faminc order cigs
```

- Visual analysis first; e.g. for faminc

```
. rvppplot faminc
```

- White test for heteroskedasticity:

## Homo- vs. heteroskedasticity

- Re-run [regression 2](#) and check for heteroskedasticity.

```
. regress bwght faminc order cigs
```

- Visual analysis first; e.g. for faminc

```
. rvppplot faminc
```

- White test for heteroskedasticity:

```
. estat imtest, white
```

## Homo- vs. heteroskedasticity

- Re-run [regression 2](#) and check for heteroskedasticity.

```
. regress bwght faminc order cigs
```

- Visual analysis first; e.g. for faminc

```
. rvppplot faminc
```

- White test for heteroskedasticity:

```
. estat imtest, white
```

White's test for  $H_0$ : homoskedasticity  
against  $H_a$ : unrestricted heteroskedasticity

```
chi2(9)      =      0.63  
Prob > chi2  =      0.9999
```

- We fail to reject  $H_0$  of homoskedasticity; no statistical evidence of heteroskedasticity.
- This is in line with our visual analysis.



Thank you and see you in  
two weeks!

Juergen Amann

[juergen.amann@nottingham.ac.uk](mailto:juergen.amann@nottingham.ac.uk)

Wednesday 12:00 - 13:00, C42 SCGB

---

## Overall relation 'by hand' [◀ Go back](#)

There are two ways how we can recover the previous statistic ourselves:

## Overall relation 'by hand' [◀ Go back](#)

There are two ways how we can recover the previous statistic ourselves:

- 1 Joint parameter test (remember, we did this last week!)

```
. test faminc = order = 0  
  
( 1) faminc - order = 0  
( 2) faminc = 0  
  
F( 2, 1385) = 11.15  
Prob > F = 0.0000
```

## Overall relation 'by hand' [◀ Go back](#)

There are two ways how we can recover the previous statistic ourselves:

- 1 Joint parameter test (remember, we did this last week!)

```
. test faminc = order = 0  
  
( 1) faminc - order = 0  
( 2) faminc = 0  
  
F( 2, 1385) = 11.15  
Prob > F = 0.0000
```

- 2 Use one of the results from lecture[Slides 4, p.11], e.g.:  $F^* = \frac{R^2/(q)}{(1-R^2)/(n-k)}$

## Overall relation 'by hand' [◀ Go back](#)

There are two ways how we can recover the previous statistic ourselves:

- 1 Joint parameter test (remember, we did this last week!)

```
. test faminc = order = 0  
  
( 1) faminc - order = 0  
( 2) faminc = 0  
  
      F( 2, 1385) =    11.15  
      Prob > F =    0.0000
```

- 2 Use one of the results from lecture[Slides 4, p.11], e.g.:  $F^* = \frac{R^2/(q)}{(1-R^2)/(n-k)}$

```
. display 0.0158/(1 - 0.0158) * 1385/2  
11.117151
```

## $R^2$ 'by hand' [◀ Go back](#)

From the lecture slides [Slides 3, pp.20]:

- The  $R^2$  measure of goodness-of-fit is defined by  $R^2 = \frac{ESS}{TSS} = 1 - \frac{RSS}{TSS}$ .

## $R^2$ 'by hand' [◀ Go back](#)

From the lecture slides [Slides 3, pp.20]:

- The  $R^2$  measure of goodness-of-fit is defined by  $R^2 = \frac{ESS}{TSS} = 1 - \frac{RSS}{TSS}$ .

```
. regress bwght faminc order cigs mothereduc
```

## $R^2$ 'by hand' [◀ Go back](#)

From the lecture slides [Slides 3, pp.20]:

- The  $R^2$  measure of goodness-of-fit is defined by  $R^2 = \frac{ESS}{TSS} = 1 - \frac{RSS}{TSS}$ .

```
. regress bwght faminc order cigs mothereduc
```

Source	SS	df	MS	Number of obs	=	1,388
				F(4, 1383)	=	12.49
Model	20031.4556	4	5007.8639	Prob > F	=	0.0000
Residual	554580.264	1,383	400.998022	R-squared	=	0.0349
				Adj R-squared	=	0.0321
Total	574611.72	1,387	414.283864	Root MSE	=	20.025



## $R^2$ 'by hand' [◀ Go back](#)

From the lecture slides [Slides 3, pp.20]:

- The  $R^2$  measure of goodness-of-fit is defined by  $R^2 = \frac{ESS}{TSS} = 1 - \frac{RSS}{TSS}$ .

```
. regress bwght faminc order cigs mothereduc
```

Source	SS	df	MS	Number of obs	=	1,388
-----				F(4, 1383)	=	12.49
Model	20031.4556	4	5007.8639	Prob > F	=	0.0000
Residual	554580.264	1,383	400.998022	R-squared	=	0.0349
-----				Adj R-squared	=	0.0321
Total	574611.72	1,387	414.283864	Root MSE	=	20.025

```
. display 20031.4556 / 574611.72  
0.03486085
```

## $R^2$ 'by hand' [◀ Go back](#)

From the lecture slides [Slides 3, pp.20]:

- The  $R^2$  measure of goodness-of-fit is defined by  $R^2 = \frac{ESS}{TSS} = 1 - \frac{RSS}{TSS}$ .

```
. regress bwght faminc order cigs mothereduc
```

Source	SS	df	MS	Number of obs	=	1,388
				F(4, 1383)	=	12.49
Model	20031.4556	4	5007.8639	Prob > F	=	0.0000
Residual	554580.264	1,383	400.998022	R-squared	=	0.0349
				Adj R-squared	=	0.0321
Total	574611.72	1,387	414.283864	Root MSE	=	20.025

```
. display 20031.4556 / 574611.72  
0.03486085
```

- The adjusted  $R^2$  incorporates a penalty for including additional variables:

$$adj.R^2 = 1 - \left[ \left( \frac{T-1}{T-k} \right) (1 - R^2) \right]$$

## $R^2$ 'by hand' [◀ Go back](#)

From the lecture slides [Slides 3, pp.20]:

- The  $R^2$  measure of goodness-of-fit is defined by  $R^2 = \frac{ESS}{TSS} = 1 - \frac{RSS}{TSS}$ .

```
. regress bwght faminc order cigs mothereduc
```

Source	SS	df	MS	Number of obs	=	1,388
				F(4, 1383)	=	12.49
Model	20031.4556	4	5007.8639	Prob > F	=	0.0000
Residual	554580.264	1,383	400.998022	R-squared	=	0.0349
				Adj R-squared	=	0.0321
Total	574611.72	1,387	414.283864	Root MSE	=	20.025

```
. display 20031.4556 / 574611.72  
0.03486085
```

- The adjusted  $R^2$  incorporates a penalty for including additional variables:

$$adj.R^2 = 1 - \left[ \left( \frac{T-1}{T-k} \right) (1 - R^2) \right]$$

```
. display 1 - ((1388-1)/(1383)*(1-20031.4556/574611.72))  
0.03206942
```