ExamGOAL
PYQ PDF

Free · With answer key

JEE MAIN · MATHEMATICS · CHAPTER-WISE PYQS

Matrices and Determinants

JEE Main Mathematics · Previous Year Questions with Answer Key

415	202	2002 – 2026	54	English
QUESTIONS	PAPERS COVERED	YEARS	OUT OF SYLLABUS	LANGUAGE

INSIDE THIS PDF

MCQ (Single Correct Answer)	290 questions
Numerical	71 questions
Out of syllabus	54 questions
Answer key	at the end

Practise online

Every question with its full solution, free on **questions.examside.com**

1. **Solve**
   Print it, or work through it on screen.
2. **Check**
   The answer key is on the last pages.
3. **Understand**
   Scan the QR code beside any question for its full solution.

MCQ (Single Correct Answer) 290 questions

Q1. JEE Main 2026 (Online) 8th April Evening Shift

If the system of linear equations :

$x + y + z = 6$
$x + 2y + 5z = 10$
$2x + 3y + \lambda z = \mu$

has infinitely many solutions, then the value of $\lambda + \mu$ equals:

(A) 12 (B) 16 (C) 22 (D) 28

Open solution

Q2. JEE Main 2026 (Online) 8th April Evening Shift

Let $A = \begin{bmatrix} \alpha & 1 & 2 \\ 2 & 3 & 0 \\ 0 & 4 & 5 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 & 0 \\ 0 & -5\alpha & 0 \\ 0 & 4\alpha & -2\alpha \end{bmatrix} + \text{adj}(A)$. If $\det(B) = 66$, then $\det(\text{adj}(A))$ equals :

(A) 289 (B) 361 (C) 441 (D) 529

Open solution

Q3. JEE Main 2026 (Online) 6th April Evening Shift

The sum of all possible values of $\theta \in [0, 2\pi]$, for which the system of equations :

$x\cos 3\theta - 8y - 12z = 0$
$x\cos 2\theta + 3y + 3z = 0$
$x + y + 3z = 0$

has a non-trivial solution, is equal to :

(A) π (B) 2π (C) 3π (D) 4π

Open solution

Q4. JEE Main 2026 (Online) 6th April Evening Shift

Let $\mathrm{A} = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 9 & 3 & 1 \end{bmatrix}$ and $\mathrm{B} = [\mathrm{b}_{ij}], 1 \le i, j \le 3$. If $\mathrm{B} = \mathrm{A}^{99} - \mathrm{I}$, then the value of $\frac{\mathrm{b}_{31} - \mathrm{b}_{21}}{\mathrm{b}_{32}}$ is :

(A) 99 (B) 199 (C) 149 (D) 159

Open solution

Q5. JEE Main 2026 (Online) 5th April Evening Shift

If $f : \mathbf{N} \to \mathbf{Z}$ is defined by

$$f(n) = \begin{vmatrix} n & -1 & -5 \\ -2n^2 & 3(2k+1) & 2k+1 \\ -3n^3 & 3k(2k+1) & 3k(k+2)+1 \end{vmatrix}, k \in N,$$

and $\sum_{n=1}^{k} f(n) = 98$, then k is equal to :

(A) 3 (B) 4 (C) 5 (D) 6

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q6. JEE Main 2026 (Online) 5th April Evening Shift

Open solution

Let M be a 3×3 matrix such that $\mathrm{M}\begin{pmatrix}1\\0\\0\end{pmatrix} = \begin{pmatrix}1\\2\\3\end{pmatrix}, \mathrm{M}\begin{pmatrix}0\\1\\0\end{pmatrix} = \begin{pmatrix}0\\1\\2\end{pmatrix}$ and $\mathrm{M}\begin{pmatrix}0\\0\\1\end{pmatrix} = \begin{pmatrix}-1\\1\\1\end{pmatrix}$. If $\mathrm{M}\begin{pmatrix}x\\y\\z\end{pmatrix} = \begin{pmatrix}1\\7\\11\end{pmatrix}$, then $x+y+z$ equals :

(A) 4 (B) 5 (C) 7 (D) 11

Q7. JEE Main 2026 (Online) 5th April Morning Shift

Open solution

Let A be a 3×3 matrix such that

$\mathrm{A}^{\mathrm{T}}\begin{bmatrix}1\\0\\1\end{bmatrix} = \begin{bmatrix}5\\2\\2\end{bmatrix}, \mathrm{A}^{\mathrm{T}}\begin{bmatrix}0\\0\\1\end{bmatrix} = \begin{bmatrix}3\\1\\1\end{bmatrix}, \mathrm{A}\begin{bmatrix}1\\0\\1\end{bmatrix} = \begin{bmatrix}3\\4\\4\end{bmatrix}$ and $\mathrm{A}\begin{bmatrix}0\\0\\1\end{bmatrix} = \begin{bmatrix}1\\3\\1\end{bmatrix}$.

If $\det(A) = 1$, then $\det\left(\text{adj}\left(A^2 + A\right)\right)$ is equal to:

(A) 16 (B) 25 (C) 49 (D) 64

Q8. JEE Main 2026 (Online) 5th April Morning Shift

Open solution

Consider the system of linear equations in x, y, z :

$x + 2y + tz = 0$
$6x + y + 5tz = 0$
$3x + t^2y + f(t)z = 0$

where $f : \mathbb{R} \to \mathbb{R}$ is a differentiable function. If this system has infinitely many solutions for all $t \in \mathbb{R}$, then f

(A) is a constant function (B) is strictly increasing on $\mathbb{R}$
(C) is strictly decreasing on R (D) has two critical points

Q9. JEE Main 2026 (Online) 4th April Evening Shift

Open solution

If the system of equations :

$x + y + z = 5$
$x + 2y + 3z = 9$
$x + 3y + \lambda z = \mu$

has infinitely many solutions, then the value of $\lambda + \mu$ is :

(A) 16 (B) 18 (C) 19 (D) 21

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q10. JEE Main 2026 (Online) 4th April Evening Shift

Let $A=\begin{bmatrix}1 & 2 & 7\\ 4 & -2 & 8\\ 3 & 8 & -7\end{bmatrix}$ and $\det(A-\alpha I)=0$, where α is a real number. If the largest possible value of α is p, then the circle $(x-p)^2+(y-2p)^2=320$, intersects the co-ordinate axes at

(A) 1 point (B) 2 points (C) 3 points (D) 4 points

Open solution

Q11. JEE Main 2026 (Online) 4th April Morning Shift

Let $S=\left\{A=\begin{bmatrix}a & b\\ c & d\end{bmatrix}: a,b,c,d\in\{0,1,2,3,4\} \text{ and } A^2-4A+3I=0\right\}$ be a set of 2×2 matrices. Then the number of matrices in S , for which the sum of the diagonal elements is equal to 4 , is :

(A) 20 (B) 17 (C) 21 (D) 19

Open solution

Q12. JEE Main 2026 (Online) 4th April Morning Shift

Let $A=\begin{bmatrix}1 & 1 & 2\\ -2 & 0 & 1\\ 1 & 3 & 5\end{bmatrix}$. Then the sum of all elements of the matrix $\operatorname{adj}\left(\operatorname{adj}\left(2(\operatorname{adj} A)^{-1}\right)\right)$ is equal to:

(A) 3 (B) 4 (C) -4 (D) -3

Open solution

Q13. JEE Main 2026 (Online) 2nd April Evening Shift

If the system of equations

$x+5y+6z=4$

$2x+3y+4z=7$

$x+6y+az=b$

has infinitely many solutions, then the point (a,b) lies on the line

(A) $y-x=3$ (B) $x-y=3$ (C) $x+y=11$ (D) $x+y=12$

Open solution

Q14. JEE Main 2026 (Online) 2nd April Morning Shift

Let $\alpha,\beta\in\mathbb{R}$ be such that the system of linear equations

$$x+2y+z=5$$
$$2x+y+\alpha z=5$$
$$8x+4y+\beta z=18$$

has no solution. Then $\frac{\beta}{\alpha}$ is equal to :

(A) -4 (B) 4 (C) 8 (D) -8

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q15. JEE Main 2026 (Online) 2nd April Morning Shift

Open solution

Let $A=\begin{bmatrix}1 & 2\\ 1 & \alpha\end{bmatrix}$ and $B=\begin{bmatrix}3 & 3\\ \beta & 2\end{bmatrix}$. If $A^2-4A+I=O$ and $B^2-5B-6I=O$, then among the two statements:

(S1) : $[(B-A)(B+A)]^T=\begin{bmatrix}13 & 15\\ 7 & 10\end{bmatrix}$

and

(S2) : $\det(\text{adj}(A+B))=-5$

(A) only (S1) is correct

(B) only (S2) is correct

(C) both (S1) and (S2) are correct

(D) both (S1) and (S2) are wrong

Q16. JEE Main 2026 (Online) 28th January Morning Shift

Open solution

Let A, B and C be three 2×2 matrices with real entries such that $B=(I+A)^{-1}$ and $\mathrm{A}+\mathrm{C}=\mathrm{I}$.

If $\mathrm{BC}=\begin{bmatrix}1 & -5\\ -1 & 2\end{bmatrix}$ and $\mathrm{CB}\begin{bmatrix}x_1\\ x_2\end{bmatrix}=\begin{bmatrix}12\\ -6\end{bmatrix}$, then x_1+x_2 is

(A) 4

(B) 2

(C) 0

(D) -2

Q17. JEE Main 2026 (Online) 24th January Evening Shift

Open solution

Let $P=[p_{ij}]$ and $Q=[q_{ij}]$ be two square matrices of order 3 such that $q_{\mathrm{ij}}=2^{(\mathrm{i}+\mathrm{j}-1)}\mathrm{p}_{\mathrm{ij}}$ and $\det(\mathrm{Q})=2^{10}$. Then the value of $\det(\text{adj}(\text{adj}\,\mathrm{P}))$ is:

(A) 81

(B) 16

(C) 124

(D) 32

Q18. JEE Main 2026 (Online) 24th January Evening Shift

Open solution

Let $f(x)=\int\frac{7x^{10}+9x^8}{(1+x^2+2x^9)^2}dx, x>0, \lim\limits_{x\to 0}f(x)=0$ and $f(1)=\frac{1}{4}$.

If $\mathrm{A}=\begin{bmatrix}0 & 0 & 1\\ \frac{1}{4} & f'(1) & 1\\ \alpha^2 & 4 & 1\end{bmatrix}$ and $\mathrm{B}=\text{adj}(\text{adj}\,\mathrm{A})$ be such that $|\mathrm{B}|=81$, then α^2 is equal to

(A) 2

(B) 4

(C) 3

(D) 1

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q19. JEE Main 2026 (Online) 23rd January Evening Shift

Open solution

The system of linear equations

$x + y + z = 6$
$2x + 5y + az = 36$
$x + 2y + 3z = b$

has :

(A) unique solution for $a = 8$ and $b = 16$

(B) infinitely many solutions for $a = 8$ and $b = 14$

(C) infinitely many solutions for $a = 8$ and $b = 16$

(D) unique solution for $a = 8$ and $b = 14$

Q20. JEE Main 2026 (Online) 23rd January Morning Shift

Open solution

Among the statements :

I: If $\begin{vmatrix} 1 & \cos\alpha & \cos\beta \\ \cos\alpha & 1 & \cos\gamma \\ \cos\beta & \cos\gamma & 1 \end{vmatrix} = \begin{vmatrix} 0 & \cos\alpha & \cos\beta \\ \cos\alpha & 0 & \cos\gamma \\ \cos\beta & \cos\gamma & 0 \end{vmatrix}$, then $\cos^2\alpha + \cos^2\beta + \cos^2\gamma = \frac{3}{2}$, and

II: If $\begin{vmatrix} x^2+x & x+1 & x-2 \\ 2x^2+3x-1 & 3x & 3x-3 \\ x^2+2x+3 & 2x-1 & 2x-1 \end{vmatrix} = px + q$, then $p^2 = 196q^2$,

(A) both are true (B) both are false (C) only I is true (D) only II is true

Q21. JEE Main 2026 (Online) 22nd January Evening Shift

Open solution

Let n be the number obtained on rolling a fair die. If the probability that the system

$x - ny + z = 6$
$x + (n-2)y + (n+1)z = 8$
$(n-1)y + z = 1$

has a unique solution is $\frac{k}{6}$, then the sum of k and all possible values of n is :

(A) 22 (B) 20 (C) 24 (D) 21

Q22. JEE Main 2026 (Online) 22nd January Evening Shift

Open solution

If $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ is a solution of the system of equations $AX = B$, where $\text{adj } A = \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & 5 \\ 1 & -2 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} 4 \\ 0 \\ 2 \end{bmatrix}$, then $|x + y + z|$ is equal to :

(A) 3 (B) 2 (C) $\frac{3}{2}$ (D) 1

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q23. JEE Main 2026 (Online) 22nd January Morning Shift

If $A=\begin{bmatrix}2 & 3\\ 3 & 5\end{bmatrix}$, then the determinant of the matrix $\left(A^{2025}-3A^{2024}+A^{2023}\right)$ is

(A) 12 (B) 24 (C) 28 (D) 16

Open solution

Q24. JEE Main 2026 (Online) 21st January Evening Shift

If the system of equations

$3x+y+4z=3$

$2x+\alpha y-z=-3$

$x+2y+z=4$

has no solution, then the value of α is equal to:

(A) 13 (B) 4 (C) 19 (D) 23

Open solution

Q25. JEE Main 2026 (Online) 21st January Evening Shift

For the matrices $A=\begin{bmatrix}3 & -4\\ 1 & -1\end{bmatrix}$ and $B=\begin{bmatrix}-29 & 49\\ -13 & 18\end{bmatrix}$, if $(A^{15}+B)\begin{bmatrix}x\\ y\end{bmatrix}=\begin{bmatrix}0\\ 0\end{bmatrix}$, then among the following which one is **true**?

(A) $x=16, y=3$ (B) $x=5, y=7$ (C) $x=11, y=2$ (D) $x=18, y=11$

Open solution

Q26. JEE Main 2025 (Online) 8th April Evening Shift

Let α be a solution of $x^2+x+1=0$, and for some *a* and *b* in $\mathbb{R}$,

$\begin{bmatrix}4 & a & b\end{bmatrix}\begin{bmatrix}1 & 16 & 13\\ -1 & -1 & 2\\ -2 & -14 & -8\end{bmatrix}=\begin{bmatrix}0 & 0 & 0\end{bmatrix}$. If $\frac{4}{\alpha^4}+\frac{m}{\alpha^a}+\frac{n}{\alpha^b}=3$, then *m* + *n* is equal to _____

(A) 11 (B) 3 (C) 8 (D) 7

Open solution

Q27. JEE Main 2025 (Online) 8th April Evening Shift

Let $A=\begin{bmatrix}2 & 2+p & 2+p+q\\ 4 & 6+2p & 8+3p+2q\\ 6 & 12+3p & 20+6p+3q\end{bmatrix}$.

If $\det(\text{adj}(\text{adj}(3A)))=2^m\cdot 3^n, m,n\in\mathbb{N}$, then $m+n$ is equal to

(A) 22 (B) 20 (C) 24 (D) 26

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q28. JEE Main 2025 (Online) 7th April Evening Shift

Open solution

Let the system of equations

x + 5y - z = 1

4x + 3y - 3z = 7

24x + y + λz = μ

λ, μ ∈ ℝ, have infinitely many solutions. Then the number of the solutions of this system,

if x, y, z are integers and satisfy 7 ≤ x + y + z ≤ 77, is :

(A) 4 (B) 5 (C) 3 (D) 6

Q29. JEE Main 2025 (Online) 7th April Morning Shift

Open solution

Let A be a 3×3 matrix such that $|\text{adj}(\text{adj}(\text{adj } A))| = 81$.

If $S = \left\{ n \in \mathbb{Z} : (|\text{ adj}(\text{adj } A)|)^{\frac{(n-1)^2}{2}} = |A|^{(3n^2-5n-4)} \right\}$, then $\sum_{n \in S} \left| A^{(n^2+n)} \right|$ is equal to :

(A) 820 (B) 866 (C) 750 (D) 732

Q30. JEE Main 2025 (Online) 7th April Morning Shift

Open solution

Let the system of equations :

$2x + 3y + 5z = 9$
$7x + 3y - 2z = 8$
$12x + 3y - (4 + \lambda)z = 16 - \mu$

have infinitely many solutions. Then the radius of the circle centred at (λ, μ) and touching the line $4x = 3y$ is :

(A) $\frac{7}{5}$ (B) $\frac{21}{5}$ (C) 7 (D) $\frac{17}{5}$

Q31. JEE Main 2025 (Online) 4th April Evening Shift

Open solution

Let the matrix $A = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$ satisfy $A^n = A^{n-2} + A^2 - I$ for $n \geqslant 3$. Then the sum of all the elements of A^{50} is :

(A) 44 (B) 39 (C) 52 (D) 53

Q32. JEE Main 2025 (Online) 3rd April Morning Shift

Open solution

Let A be a matrix of order 3×3 and $|A| = 5$. If $|2 \text{ adj}(3A \text{ adj}(2A))| = 2^\alpha \cdot 3^\beta \cdot 5^\gamma, \alpha, \beta, \gamma \in N$, then $\alpha + \beta + \gamma$ is equal to

(A) 26 (B) 27 (C) 25 (D) 28

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q33. JEE Main 2025 (Online) 2nd April Evening Shift

Open solution

If the system of equations

$2x + \lambda y + 3z = 5$
$3x + 2y - z = 7$
$4x + 5y + \mu z = 9$

has infinitely many solutions, then $(\lambda^2 + \mu^2)$ is equal to :

(A) 30 (B) 26 (C) 22 (D) 18

Q34. JEE Main 2025 (Online) 2nd April Evening Shift

Open solution

Let A be a 3×3 real matrix such that $A^2(A - 2I) - 4(A - I) = O$, where I and O are the identity and null matrices, respectively. If $A^5 = \alpha A^2 + \beta A + \gamma I$, where α, β, and γ are real constants, then $\alpha + \beta + \gamma$ is equal to :

(A) 76 (B) 12 (C) 4 (D) 20

Q35. JEE Main 2025 (Online) 2nd April Morning Shift

Open solution

Let $\mathrm{A} = \begin{bmatrix} \alpha & -1 \\ 6 & \beta \end{bmatrix}, \alpha > 0$, such that $\det(\mathrm{A}) = 0$ and $\alpha + \beta = 1$. If I denotes 2×2 identity matrix, then the matrix $(I + A)^8$ is :

(A) $\begin{bmatrix} 257 & -64 \\ 514 & -127 \end{bmatrix}$ (B) $\begin{bmatrix} 766 & -255 \\ 1530 & -509 \end{bmatrix}$

(C) $\begin{bmatrix} 1025 & -511 \\ 2024 & -1024 \end{bmatrix}$ (D) $\begin{bmatrix} 4 & -1 \\ 6 & -1 \end{bmatrix}$

Q36. JEE Main 2025 (Online) 2nd April Morning Shift

Open solution

Let $a \in R$ and A be a matrix of order 3×3 such that $\det(A) = -4$ and $A + I = \begin{bmatrix} 1 & a & 1 \\ 2 & 1 & 0 \\ a & 1 & 2 \end{bmatrix}$, where I is the identity matrix of order 3×3. If $\det((a+1)\,\mathrm{adj}((a-1)A))$ is $2^{\mathrm{m}}3^{\mathrm{n}}, \mathrm{m}, \mathrm{n} \in \{0, 1, 2, \ldots, 20\}$, then $\mathrm{m} + \mathrm{n}$ is equal to :

(A) 14 (B) 17 (C) 15 (D) 16

Q37. JEE Main 2025 (Online) 2nd April Morning Shift

Open solution

If the system of linear equations

$3x + y + \beta z = 3$
$2x + \alpha y - z = -3$
$x + 2y + z = 4$

has infinitely many solutions, then the value of $22\beta - 9\alpha$ is :

(A) 31 (B) 37 (C) 43 (D) 49

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q38. JEE Main 2025 (Online) 29th January Evening Shift

Let $A = [a_{ij}]$ be a 2×2 matrix such that $a_{ij} \in \{0, 1\}$ for all i and j. Let the random variable X denote the possible values of the determinant of the matrix A. Then, the variance of X is:

(A) $\frac{5}{8}$ (B) $\frac{1}{4}$ (C) $\frac{3}{4}$ (D) $\frac{3}{8}$

Open solution

Q39. JEE Main 2025 (Online) 29th January Evening Shift

Let $\alpha, \beta\ (\alpha \neq \beta)$ be the values of m, for which the equations $x + y + z = 1$, $x + 2y + 4z = m$ and $x + 4y + 10z = m^2$ have infinitely many solutions. Then the value of $\sum_{n=1}^{10}(n^{\alpha} + n^{\beta})$ is equal to :

(A) 3410 (B) 560 (C) 3080 (D) 440

Open solution

Q40. JEE Main 2025 (Online) 29th January Evening Shift

Let $\mathrm{A} = [a_{ij}]$ be a matrix of order 3×3, with $a_{ij} = (\sqrt{2})^{i+j}$. If the sum of all the elements in the third row of A^2 is $\alpha + \beta\sqrt{2}, \alpha, \beta \in \mathbf{Z}$, then $\alpha + \beta$ is equal to :

(A) 210 (B) 280 (C) 224 (D) 168

Open solution

Q41. JEE Main 2025 (Online) 29th January Morning Shift

Let $A = [a_{ij}] = \begin{bmatrix} \log_5 128 & \log_4 5 \\ \log_5 8 & \log_4 25 \end{bmatrix}$. If A_{ij} is the cofactor of a_{ij}, $C_{ij} = \sum_{k=1}^{2} a_{ik}A_{jk}, 1 \leq i, j \leq 2$, and $C = [C_{ij}]$, then $8|C|$ is equal to :

(A) 288 (B) 262 (C) 222 (D) 242

Open solution

Q42. JEE Main 2025 (Online) 29th January Morning Shift

Let M and m respectively be the maximum and the minimum values of

$$f(x) = \begin{vmatrix} 1 + \sin^2 x & \cos^2 x & 4\sin 4x \\ \sin^2 x & 1 + \cos^2 x & 4\sin 4x \\ \sin^2 x & \cos^2 x & 1 + 4\sin 4x \end{vmatrix}, x \in R$$

Then $M^4 - m^4$ is equal to :

(A) 1280 (B) 1040 (C) 1215 (D) 1295

Open solution

Q43. JEE Main 2025 (Online) 28th January Evening Shift

Let $\mathrm{A} = \begin{bmatrix} \frac{1}{\sqrt{2}} & -2 \\ 0 & 1 \end{bmatrix}$ and $\mathrm{P} = \begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}, \theta > 0$. If $\mathrm{B} = \mathrm{PAP}^{\top}, \mathrm{C} = \mathrm{P}^{\top}\mathrm{B}^{10}\mathrm{P}$ and the sum of the diagonal elements of C is $\frac{m}{n}$, where $\gcd(m, n) = 1$, then $m + n$ is :

(A) 127 (B) 2049 (C) 258 (D) 65

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q44. JEE Main 2025 (Online) 24th January Evening Shift

For some a, b, let $f(x) = \begin{vmatrix} a + \frac{\sin x}{x} & 1 & b \\ a & 1 + \frac{\sin x}{x} & b \\ a & 1 & b + \frac{\sin x}{x} \end{vmatrix}, x \neq 0, \lim_{x \to 0} f(x) = \lambda + \mu a + \nu b$. Then $(\lambda + \mu + v)^2$ is equal to :

Open solution

(A) 25 (B) 16 (C) 9 (D) 36

Q45. JEE Main 2025 (Online) 24th January Evening Shift

If the system of equations

$x + 2y - 3z = 2$
$2x + \lambda y + 5z = 5$
$14x + 3y + \mu z = 33$

has infinitely many solutions, then $\lambda + \mu$ is equal to :

Open solution

(A) 13 (B) 10 (C) 12 (D) 11

Q46. JEE Main 2025 (Online) 24th January Morning Shift

If the system of equations
$2x - y + z = 4$
$5x + \lambda y + 3z = 12$
$100x - 47y + \mu z = 212$
has infinitely many solutions, then $\mu - 2\lambda$ is equal to

Open solution

(A) 56 (B) 59 (C) 57 (D) 55

Q47. JEE Main 2025 (Online) 23rd January Evening Shift

The system of equations
$x + y + z = 6,$
$x + 2y + 5z = 9,$
$x + 5y + \lambda z = \mu,$
has no solution if

Open solution

(A) $\lambda = 17, \mu = 18$ (B) $\lambda = 17, \mu \neq 18$ (C) $\lambda = 15, \mu \neq 17$ (D) $\lambda \neq 17, \mu \neq 18$

Q48. JEE Main 2025 (Online) 23rd January Evening Shift

Let $A = [a_{ij}]$ be a 3×3 matrix such that $A \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, A \begin{bmatrix} 4 \\ 1 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$ and $A \begin{bmatrix} 2 \\ 1 \\ 2 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$, then a_{23} equals :

Open solution

(A) 2 (B) -1 (C) 1 (D) 0

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q49. JEE Main 2025 (Online) 23rd January Morning Shift

Open solution

If the system of equations

$(\lambda-1)x+(\lambda-4)y+\lambda z=5$
$\lambda x+(\lambda-1)y+(\lambda-4)z=7$
$(\lambda+1)x+(\lambda+2)y-(\lambda+2)z=9$

has infinitely many solutions, then $\lambda^2+\lambda$ is equal to

(A) 20 (B) 10 (C) 6 (D) 12

Q50. JEE Main 2025 (Online) 23rd January Morning Shift

Open solution

If $\mathrm{A}, \mathrm{B}, \text{and } \left(\operatorname{adj}\left(\mathrm{A}^{-1}\right)+\operatorname{adj}\left(\mathrm{B}^{-1}\right)\right)$ are non-singular matrices of same order, then the inverse of $A\left(\operatorname{adj}\left(A^{-1}\right)+\operatorname{adj}\left(B^{-1}\right)\right)^{-1}B$, is equal to

(A) $\frac{AB^{-1}}{|A|}+\frac{BA^{-1}}{|B|}$

(B) $\operatorname{adj}\left(\mathrm{B}^{-1}\right)+\operatorname{adj}\left(\mathrm{A}^{-1}\right)$

(C) $\mathrm{AB}^{-1}+\mathrm{A}^{-1}\,\mathrm{B}$

(D) $\frac{1}{|AB|}(\operatorname{adj}(B)+\operatorname{adj}(A))$

Q51. JEE Main 2025 (Online) 22nd January Evening Shift

Open solution

If the system of linear equations :

$x+y+2z=6$
$2x+3y+\mathrm{az}=\mathrm{a}+1$
$-x-3y+\mathrm{bz}=2\,\mathrm{b}$

where $a, b \in \mathbf{R}$, has infinitely many solutions, then $7a+3b$ is equal to :

(A) 12 (B) 9 (C) 22 (D) 16

Q52. JEE Main 2025 (Online) 22nd January Evening Shift

Open solution

For a 3×3 matrix M, let trace (M) denote the sum of all the diagonal elements of M. Let A be a 3×3 matrix such that $|A|=\frac{1}{2}$ and trace $(A)=3$. If $B=\operatorname{adj}(\operatorname{adj}(2A))$, then the value of $|B|+$ trace (B) equals :

(A) 56 (B) 132 (C) 174 (D) 280

Q53. JEE Main 2024 (Online) 9th April Evening Shift

Open solution

Let $B=\begin{bmatrix}1 & 3\\ 1 & 5\end{bmatrix}$ and A be a 2×2 matrix such that $AB^{-1}=A^{-1}$. If $BCB^{-1}=A$ and $C^4+\alpha C^2+\beta I=O$, then $2\beta-\alpha$ is equal to

(A) 16 (B) 10 (C) 8 (D) 2

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q54. JEE Main 2024 (Online) 9th April Morning Shift

Let $\lambda, \mu \in \mathbf{R}$. If the system of equations

$3x + 5y + \lambda z = 3$

$7x + 11y - 9z = 2$

$97x + 155y - 189z = \mu$

has infinitely many solutions, then $\mu + 2\lambda$ is equal to :

(A) 24 (B) 25 (C) 27 (D) 22

Open solution

Q55. JEE Main 2024 (Online) 8th April Evening Shift

If $\alpha \neq \mathrm{a}, \beta \neq \mathrm{b}, \gamma \neq \mathrm{c}$ and $\begin{vmatrix} \alpha & \mathrm{b} & \mathrm{c} \\ \mathrm{a} & \beta & \mathrm{c} \\ \mathrm{a} & \mathrm{b} & \gamma \end{vmatrix} = 0$, then $\frac{\mathrm{a}}{\alpha-\mathrm{a}} + \frac{\mathrm{b}}{\beta-\mathrm{b}} + \frac{\gamma}{\gamma-\mathrm{c}}$ is equal to :

(A) 2 (B) 3 (C) 1 (D) 0

Open solution

Q56. JEE Main 2024 (Online) 8th April Evening Shift

If the system of equations $x + 4y - z = \lambda, 7x + 9y + \mu z = -3, 5x + y + 2z = -1$ has infinitely many solutions, then $(2\mu + 3\lambda)$ is equal to :

(A) -2 (B) 2 (C) 3 (D) -3

Open solution

Q57. JEE Main 2024 (Online) 8th April Morning Shift

Let $A = \begin{bmatrix} 2 & a & 0 \\ 1 & 3 & 1 \\ 0 & 5 & b \end{bmatrix}$. If $A^3 = 4A^2 - A - 21I$, where I is the identity matrix of order 3×3, then $2a + 3b$ is equal to

(A) -10 (B) -12 (C) -13 (D) -9

Open solution

Q58. JEE Main 2024 (Online) 6th April Evening Shift

If A is a square matrix of order 3 such that $\det(A) = 3$ and $\det\left(\operatorname{adj}\left(-4\operatorname{adj}\left(-3\operatorname{adj}\left(3\operatorname{adj}\left((2\,\mathrm{A})^{-1}\right)\right)\right)\right)\right) = 2^{\mathrm{m}}3^{\mathrm{n}}$, then $\mathrm{m} + 2\mathrm{n}$ is equal to :

(A) 2 (B) 4 (C) 3 (D) 6

Open solution

Q59. JEE Main 2024 (Online) 6th April Morning Shift

For $\alpha, \beta \in \mathbb{R}$ and a natural number n, let $A_r = \begin{vmatrix} r & 1 & \frac{n^2}{2} + \alpha \\ 2r & 2 & n^2 - \beta \\ 3r - 2 & 3 & \frac{n(3n-1)}{2} \end{vmatrix}$. Then $2A_{10} - A_8$ is

(A) $4\alpha + 2\beta$ (B) 0 (C) $2n$ (D) $2\alpha + 4\beta$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q60. JEE Main 2024 (Online) 5th April Evening Shift

Open solution

The values of m, n, for which the system of equations

$x + y + z = 4,$
$2x + 5y + 5z = 17,$
$x + 2y + mz = n$

has infinitely many solutions, satisfy the equation :

(A) $m^2 + n^2 - m - n = 46$ (B) $m^2 + n^2 + mn = 68$

(C) $m^2 + n^2 - mn = 39$ (D) $m^2 + n^2 + m + n = 64$

Q61. JEE Main 2024 (Online) 5th April Evening Shift

Open solution

Let $\alpha\beta \neq 0$ and $A = \begin{bmatrix} \beta & \alpha & 3 \\ \alpha & \alpha & \beta \\ -\beta & \alpha & 2\alpha \end{bmatrix}$. If $B = \begin{bmatrix} 3\alpha & -9 & 3\alpha \\ -\alpha & 7 & -2\alpha \\ -2\alpha & 5 & -2\beta \end{bmatrix}$ is the matrix of cofactors of the elements of A, then $\det(AB)$ is equal to :

(A) 64 (B) 343 (C) 125 (D) 216

Q62. JEE Main 2024 (Online) 5th April Morning Shift

Open solution

Let A and B be two square matrices of order 3 such that $|A| = 3$ and $|B| = 2$. Then $|A^T A(\text{adj}(2A))^{-1}(\text{adj}(4B))(\text{adj}(AB))^{-1}AA^T|$ is equal to :

(A) 32 (B) 81 (C) 64 (D) 108

Q63. JEE Main 2024 (Online) 5th April Morning Shift

Open solution

If the system of equations

$11x + y + \lambda z = -5$
$2x + 3y + 5z = 3$
$8x - 19y - 39z = \mu$

has infinitely many solutions, then $\lambda^4 - \mu$ is equal to :

(A) 51 (B) 45 (C) 47 (D) 49

Q64. JEE Main 2024 (Online) 4th April Evening Shift

Open solution

Let $A = \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}$ and $B = I + \text{adj}(A) + (\text{adj } A)^2 + \ldots + (\text{adj } A)^{10}$. Then, the sum of all the elements of the matrix B is:

(A) -110 (B) 22 (C) -124 (D) -88

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q65. JEE Main 2024 (Online) 4th April Morning Shift

Let $\alpha \in (0, \infty)$ and $A = \begin{bmatrix} 1 & 2 & \alpha \\ 1 & 0 & 1 \\ 0 & 1 & 2 \end{bmatrix}$. If $\det\left(\text{adj}\left(2A - A^T\right) \cdot \text{adj}\left(A - 2A^T\right)\right) = 2^8$, then $(\det(A))^2$ is equal to:

Open solution

(A) 16 (B) 36 (C) 49 (D) 1

Q66. JEE Main 2024 (Online) 4th April Morning Shift

If the system of equations

$x + (\sqrt{2}\sin\alpha)y + (\sqrt{2}\cos\alpha)z = 0$
$x + (\cos\alpha)y + (\sin\alpha)z = 0$
$x + (\sin\alpha)y - (\cos\alpha)z = 0$

has a non-trivial solution, then $\alpha \in \left(0, \frac{\pi}{2}\right)$ is equal to :

Open solution

(A) $\frac{5\pi}{24}$ (B) $\frac{11\pi}{24}$ (C) $\frac{7\pi}{24}$ (D) $\frac{3\pi}{4}$

Q67. JEE Main 2024 (Online) 1st February Evening Shift

Let the system of equations $x + 2y + 3z = 5, 2x + 3y + z = 9, 4x + 3y + \lambda z = \mu$ have infinite number of solutions. Then $\lambda + 2\mu$ is equal to :

Open solution

(A) 22 (B) 17 (C) 15 (D) 28

Q68. JEE Main 2024 (Online) 1st February Morning Shift

If $\mathrm{A} = \begin{bmatrix} \sqrt{2} & 1 \\ -1 & \sqrt{2} \end{bmatrix}, \mathrm{B} = \begin{bmatrix} 1 & 0 \\ 1 & 1 \end{bmatrix}, \mathrm{C} = \mathrm{ABA}^{\mathrm{T}}$ and $\mathrm{X} = \mathrm{A}^{\mathrm{T}}\mathrm{C}^2\,\mathrm{A}$, then $\det \mathrm{X}$ is equal to :

Open solution

(A) 243 (B) 729 (C) 27 (D) 891

Q69. JEE Main 2024 (Online) 1st February Morning Shift

If the system of equations

$2x + 3y - z = 5$

$x + \alpha y + 3z = -4$

$3x - y + \beta z = 7$

has infinitely many solutions, then $13\alpha\beta$ is equal to :

Open solution

(A) 1110 (B) 1120 (C) 1210 (D) 1220

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q70. JEE Main 2024 (Online) 31st January Evening Shift

Open solution

Let A be a 3×3 real matrix such that

$$A\begin{pmatrix}1\\0\\1\end{pmatrix} = 2\begin{pmatrix}1\\0\\1\end{pmatrix}, A\begin{pmatrix}-1\\0\\1\end{pmatrix} = 4\begin{pmatrix}-1\\0\\1\end{pmatrix}, A\begin{pmatrix}0\\1\\0\end{pmatrix} = 2\begin{pmatrix}0\\1\\0\end{pmatrix}.$$

Then, the system $(A - 3I)\begin{pmatrix}x\\y\\z\end{pmatrix} = \begin{pmatrix}1\\2\\3\end{pmatrix}$ has :

(A) exactly two solutions

(B) infinitely many solutions

(C) unique solution

(D) no solution

Q71. JEE Main 2024 (Online) 31st January Morning Shift

Open solution

If the system of linear equations

$x - 2y + z = -4$
$2x + \alpha y + 3z = 5$
$3x - y + \beta z = 3$

has infinitely many solutions, then $12\alpha + 13\beta$ is equal to

(A) 60 (B) 54 (C) 64 (D) 58

Q72. JEE Main 2024 (Online) 30th January Evening Shift

Open solution

Let $R = \begin{pmatrix}x & 0 & 0\\0 & y & 0\\0 & 0 & z\end{pmatrix}$ be a non-zero 3×3 matrix, where $x\sin\theta = y\sin\left(\theta + \frac{2\pi}{3}\right) = z\sin\left(\theta + \frac{4\pi}{3}\right) \neq 0, \theta \in (0, 2\pi)$. For a square matrix M, let trace (M) denote the sum of all the diagonal entries of M. Then, among the statements:

(I) Trace $(R) = 0$

(II) If trace $(\text{adj}(\text{adj}(R))) = 0$, then R has exactly one non-zero entry.

(A) Only (I) is true

(B) Only (II) is true

(C) Both (I) and (II) are true

(D) Neither (I) nor (II) is true

Q73. JEE Main 2024 (Online) 30th January Evening Shift

Open solution

Consider the system of linear equations $x + y + z = 5, x + 2y + \lambda^2 z = 9, x + 3y + \lambda z = \mu$, where $\lambda, \mu \in \mathbb{R}$. Then, which of the following statements is NOT correct?

(A) System is consistent if $\lambda \neq 1$ and $\mu = 13$

(B) System is inconsistent if $\lambda = 1$ and $\mu \neq 13$

(C) System has unique solution if $\lambda \neq 1$ and $\mu \neq 13$

(D) System has infinite number of solutions if $\lambda = 1$ and $\mu = 13$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q74. JEE Main 2024 (Online) 30th January Morning Shift

Consider the system of linear equations
$x+y+z=4\mu, x+2y+2\lambda z=10\mu, x+3y+4\lambda^2 z=\mu^2+15$ where $\lambda, \mu \in \mathbf{R}$. Which one of the following statements is NOT correct ?

(A) The system has unique solution if $\lambda \neq \frac{1}{2}$ and $\mu \neq 1, 15$

(B) The system has infinite number of solutions if $\lambda = \frac{1}{2}$ and $\mu = 15$

(C) The system is consistent if $\lambda \neq \frac{1}{2}$

(D) The system is inconsistent if $\lambda = \frac{1}{2}$ and $\mu \neq 1$

Open solution

Q75. JEE Main 2024 (Online) 29th January Evening Shift

Let $A=\begin{bmatrix}2 & 1 & 2\\ 6 & 2 & 11\\ 3 & 3 & 2\end{bmatrix}$ and $P=\begin{bmatrix}1 & 2 & 0\\ 5 & 0 & 2\\ 7 & 1 & 5\end{bmatrix}$. The sum of the prime factors of $\left|P^{-1}AP-2I\right|$ is equal to

(A) 66 (B) 27 (C) 23 (D) 26

Open solution

Q76. JEE Main 2024 (Online) 29th January Morning Shift

Let $A=\begin{bmatrix}1 & 0 & 0\\ 0 & \alpha & \beta\\ 0 & \beta & \alpha\end{bmatrix}$ and $|2\,\mathrm{A}|^3=2^{21}$ where $\alpha, \beta \in Z$. Then a value of α is

(A) 9 (B) 17 (C) 3 (D) 5

Open solution

Q77. JEE Main 2024 (Online) 29th January Morning Shift

Let A be a square matrix such that $\mathrm{AA}^{\mathrm{T}}=\mathrm{I}$. Then $\frac{1}{2}A\left[\left(A+A^T\right)^2+\left(A-A^T\right)^2\right]$ is equal to

(A) $\mathrm{A}^2+\mathrm{A}^{\mathrm{T}}$ (B) $\mathrm{A}^3+\mathrm{I}$

(C) $\mathrm{A}^3+\mathrm{A}^{\mathrm{T}}$ (D) $\mathrm{A}^2+\mathrm{I}$

Open solution

Q78. JEE Main 2024 (Online) 27th January Evening Shift

The values of α, for which $\begin{vmatrix}1 & \frac{3}{2} & \alpha+\frac{3}{2}\\ 1 & \frac{1}{3} & \alpha+\frac{1}{3}\\ 2\alpha+3 & 3\alpha+1 & 0\end{vmatrix}=0$, lie in the interval

(A) $(-2,1)$ (B) $\left(-\frac{3}{2},\frac{3}{2}\right)$

(C) $(-3,0)$ (D) $(0,3)$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q79. JEE Main 2024 (Online) 27th January Morning Shift

Consider the matrix $f(x) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix}$.

Open solution

Given below are two statements :

Statement I : $f(-x)$ is the inverse of the matrix $f(x)$.

Statement II : $f(x)f(y) = f(x+y)$.

In the light of the above statements, choose the correct answer from the options given below :

(A) Statement I is false but Statement II is true

(B) Both Statement I and Statement II are false

(C) Both Statement I and Statement II are true

(D) Statement I is true but Statement II is false

Q80. JEE Main 2023 (Online) 13th April Evening Shift

Open solution

If the system of equations

$2x + y - z = 5$

$2x - 5y + \lambda z = \mu$

$x + 2y - 5z = 7$

has infinitely many solutions, then $(\lambda+\mu)^2 + (\lambda-\mu)^2$ is equal to

(A) 916 (B) 912 (C) 920 (D) 904

Q81. JEE Main 2023 (Online) 13th April Morning Shift

Open solution

For the system of linear equations

$2x + 4y + 2az = b$

$x + 2y + 3z = 4$

$2x - 5y + 2z = 8$

which of the following is NOT correct?

(A) It has infinitely many solutions if $a = 3, b = 8$

(B) It has infinitely many solutions if $a = 3, b = 6$

(C) It has unique solution if $a = b = 8$

(D) It has unique solution if $a = b = 6$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q82. JEE Main 2023 (Online) 13th April Morning Shift

Let $B=\begin{bmatrix}1 & 3 & \alpha\\ 1 & 2 & 3\\ \alpha & \alpha & 4\end{bmatrix}$, $\alpha>2$ be the adjoint of a matrix A and $|A|=2$. Then $\begin{bmatrix}\alpha & -2\alpha & \alpha\end{bmatrix}B\begin{bmatrix}\alpha\\ -2\alpha\\ \alpha\end{bmatrix}$ is equal to :

Open solution

(A) 32 (B) -16 (C) 0 (D) 16

Q83. JEE Main 2023 (Online) 13th April Morning Shift

The number of symmetric matrices of order 3, with all the entries from the set $\{0,1,2,3,4,5,6,7,8,9\}$ is :

Open solution

(A) 10^9 (B) 9^{10} (C) 10^6 (D) 6^{10}

Q84. JEE Main 2023 (Online) 12th April Morning Shift

Let $A=\begin{bmatrix}1 & \frac{1}{51}\\ 0 & 1\end{bmatrix}$. If $B=\begin{bmatrix}1 & 2\\ -1 & -1\end{bmatrix}A\begin{bmatrix}-1 & -2\\ 1 & 1\end{bmatrix}$, then the sum of all the elements of the matrix $\sum_{n=1}^{50}B^n$ is equal to

Open solution

(A) 50 (B) 75 (C) 100 (D) 125

Q85. JEE Main 2023 (Online) 11th April Evening Shift

If the system of linear equations

$7x+11y+\alpha z=13$

$5x+4y+7z=\beta$

$175x+194y+57z=361$

has infinitely many solutions, then $\alpha+\beta+2$ is equal to :

Open solution

(A) 6 (B) 4 (C) 5 (D) 3

Q86. JEE Main 2023 (Online) 11th April Evening Shift

$\begin{vmatrix}x+1 & x & x\\ x & x+\lambda & x\\ x & x & x+\lambda^2\end{vmatrix}=\frac{9}{8}(103x+81)$, then λ, $\frac{\lambda}{3}$ are the roots of the equation :

Open solution

(A) $4x^2+24x-27=0$ (B) $4x^2-24x+27=0$

(C) $4x^2-24x-27=0$ (D) $4x^2+24x+27=0$

Q87. JEE Main 2023 (Online) 11th April Morning Shift

Let A be a 2×2 matrix with real entries such that $A'=\alpha A+I$, where $\alpha\in\mathbb{R}-\{-1,1\}$. If $\det\left(A^2-A\right)=4$, then the sum of all possible values of α is equal to :

Open solution

(A) 2 (B) $\frac{3}{2}$ (C) 0 (D) $\frac{5}{2}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q88. JEE Main 2023 (Online) 10th April Morning Shift

Open solution

For the system of linear equations

$2x - y + 3z = 5$

$3x + 2y - z = 7$

$4x + 5y + \alpha z = \beta,$

which of the following is **NOT** correct?

(A) The system has infinitely many solutions for $\alpha = -6$ and $\beta = 9$

(B) The system has a unique solution for $\alpha \neq -5$ and $\beta = 8$

(C) The system is inconsistent for $\alpha = -5$ and $\beta = 8$

(D) The system has infinitely many solutions for $\alpha = -5$ and $\beta = 9$

Q89. JEE Main 2023 (Online) 8th April Evening Shift

Open solution

If $A = \begin{bmatrix} 1 & 5 \\ \lambda & 10 \end{bmatrix}$, $A^{-1} = \alpha A + \beta I$ and $\alpha + \beta = -2$, then $4\alpha^2 + \beta^2 + \lambda^2$ is equal to :

(A) 12 (B) 10 (C) 19 (D) 14

Q90. JEE Main 2023 (Online) 8th April Evening Shift

Open solution

Let S be the set of all values of $\theta \in [-\pi, \pi]$ for which the system of linear equations

$x + y + \sqrt{3}z = 0$

$-x + (\tan\theta)y + \sqrt{7}z = 0$

$x + y + (\tan\theta)z = 0$

has non-trivial solution. Then $\frac{120}{\pi} \sum_{\theta \in S} \theta$ is equal to :

(A) 40 (B) 30 (C) 10 (D) 20

Q91. JEE Main 2023 (Online) 8th April Morning Shift

Open solution

Let $A = \begin{bmatrix} 2 & 1 & 0 \\ 1 & 2 & -1 \\ 0 & -1 & 2 \end{bmatrix}$. If $|\text{adj}(\text{adj}(\text{adj}\, 2A))| = (16)^n$, then n is equal to :

(A) 9 (B) 8 (C) 10 (D) 12

Q92. JEE Main 2023 (Online) 8th April Morning Shift

Open solution

Let $P = \begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$, $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ and $Q = PAP^T$. If $P^T Q^{2007} P = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, then $2a + b - 3c - 4d$ equal to :

(A) 2004 (B) 2006 (C) 2007 (D) 2005

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q93. JEE Main 2023 (Online) 6th April Evening Shift

Let P be a square matrix such that $P^2 = I - P$. For $\alpha, \beta, \gamma, \delta \in \mathbb{N}$, if $P^\alpha + P^\beta = \gamma I - 29P$ and $P^\alpha - P^\beta = \delta I - 13P$, then $\alpha + \beta + \gamma - \delta$ is equal to :

(A) 18 (B) 22 (C) 24 (D) 40

Open solution

Q94. JEE Main 2023 (Online) 6th April Evening Shift

For the system of equations

$x + y + z = 6$

$x + 2y + \alpha z = 10$

$x + 3y + 5z = \beta$, which one of the following is **NOT** true?

(A) System has a unique solution for $\alpha = 3, \beta \neq 14$.

(B) System has infinitely many solutions for $\alpha = 3, \beta = 14$.

(C) System has no solution for $\alpha = 3, \beta = 24$.

(D) System has a unique solution for $\alpha = -3, \beta = 14$.

Open solution

Q95. JEE Main 2023 (Online) 6th April Morning Shift

If the system of equations

$x + y + az = b$

$2x + 5y + 2z = 6$

$x + 2y + 3z = 3$

has infinitely many solutions, then $2a + 3b$ is equal to :

(A) 28 (B) 25 (C) 20 (D) 23

Open solution

Q96. JEE Main 2023 (Online) 6th April Morning Shift

Let $\mathrm{A} = [\mathrm{a_{ij}}]_{2\times 2}$, where $\mathrm{a_{ij}} \neq 0$ for all i, j and $\mathrm{A}^2 = \mathrm{I}$. Let a be the sum of all diagonal elements of A and $\mathrm{b} = |\mathrm{A}|$. Then $3a^2 + 4b^2$ is equal to :

(A) 4 (B) 3 (C) 14 (D) 7

Open solution

Q97. JEE Main 2023 (Online) 1st February Evening Shift

For the system of linear equations $\alpha x + y + z = 1, x + \alpha y + z = 1, x + y + \alpha z = \beta$, which one of the following statements is **NOT** correct?

(A) It has infinitely many solutions if $\alpha = 1$ and $\beta = 1$

(B) It has infinitely many solutions if $\alpha = 2$ and $\beta = -1$

(C) $x + y + z = \frac{3}{4}$ if $\alpha = 2$ and $\beta = 1$

(D) It has no solution if $\alpha = -2$ and $\beta = 1$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q98. JEE Main 2023 (Online) 1st February Evening Shift

If $A = \frac{1}{2}\begin{bmatrix} 1 & \sqrt{3} \\ -\sqrt{3} & 1 \end{bmatrix}$, then :

(A) $A^{30} - A^{25} = 2I$ (B) $A^{30} + A^{25} - A = I$ (C) $A^{30} = A^{25}$ (D) $A^{30} + A^{25} + A = I$

Open solution

Q99. JEE Main 2023 (Online) 1st February Morning Shift

Let S denote the set of all real values of λ such that the system of equations

$\lambda x + y + z = 1$

$x + \lambda y + z = 1$

$x + y + \lambda z = 1$

is inconsistent, then $\sum_{\lambda \in S} \left(|\lambda|^2 + |\lambda|\right)$ is equal to

(A) 12 (B) 2 (C) 4 (D) 6

Open solution

Q100. JEE Main 2023 (Online) 1st February Morning Shift

Let $f(x) = \begin{vmatrix} 1+\sin^2 x & \cos^2 x & \sin 2x \\ \sin^2 x & 1+\cos^2 x & \sin 2x \\ \sin^2 x & \cos^2 x & 1+\sin 2x \end{vmatrix}$, $x \in \left[\frac{\pi}{6}, \frac{\pi}{3}\right]$. If α and β respectively are the maximum and the minimum values of f, then

(A) $\alpha^2 - \beta^2 = 4\sqrt{3}$ (B) $\beta^2 - 2\sqrt{\alpha} = \frac{19}{4}$

(C) $\beta^2 + 2\sqrt{\alpha} = \frac{19}{4}$ (D) $\alpha^2 + \beta^2 = \frac{9}{2}$

Open solution

Q101. JEE Main 2023 (Online) 31st January Morning Shift

For the system of linear equations

$x + y + z = 6$

$\alpha x + \beta y + 7z = 3$

$x + 2y + 3z = 14$

which of the following is **NOT** true ?

(A) If $\alpha = \beta = 7$, then the system has no solution

(B) For every point $(\alpha, \beta) \neq (7, 7)$ on the line $x - 2y + 7 = 0$, the system has infinitely many solutions

(C) There is a unique point (α, β) on the line $x + 2y + 18 = 0$ for which the system has infinitely many solutions

(D) If $\alpha = \beta$ and $\alpha \neq 7$, then the system has a unique solution

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q102. JEE Main 2023 (Online) 31st January Morning Shift

Let $A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 4 & -1 \\ 0 & 12 & -3 \end{pmatrix}$. Then the sum of the diagonal elements of the matrix $(A + I)^{11}$ is equal to :

Open solution

(A) 4094 (B) 2050 (C) 6144 (D) 4097

Q103. JEE Main 2023 (Online) 30th January Evening Shift

For $\alpha, \beta \in \mathbb{R}$, suppose the system of linear equations

$$x - y + z = 5$$
$$2x + 2y + \alpha z = 8$$
$$3x - y + 4z = \beta$$

has infinitely many solutions. Then α and β are the roots of :

Open solution

(A) $x^2 + 18x + 56 = 0$ (B) $x^2 - 10x + 16 = 0$ (C) $x^2 + 14x + 24 = 0$ (D) $x^2 - 18x + 56 = 0$

Q104. JEE Main 2023 (Online) 30th January Morning Shift

Let the system of linear equations

$x + y + kz = 2$

$2x + 3y - z = 1$

$3x + 4y + 2z = k$

have infinitely many solutions. Then the system

$(k + 1)x + (2k - 1)y = 7$

$(2k + 1)x + (k + 5)y = 10$

has :

Open solution

(A) unique solution satisfying $x - y = 1$ (B) infinitely many solutions

(C) no solution (D) unique solution satisfying $x + y = 1$

Q105. JEE Main 2023 (Online) 29th January Morning Shift

Let α and β be real numbers. Consider a 3 $\times$ 3 matrix A such that $A^2 = 3A + \alpha I$. If $A^4 = 21A + \beta I$, then

Open solution

(A) $\alpha = 1$ (B) $\alpha = 4$ (C) $\beta = 8$ (D) $\beta = -8$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q106. JEE Main 2023 (Online) 29th January Morning Shift

Open solution

Consider the following system of equations

$\alpha x + 2y + z = 1$

$2\alpha x + 3y + z = 1$

$3x + \alpha y + 2z = \beta$

for some $\alpha, \beta \in \mathbb{R}$. Then which of the following is NOT correct.

(A) It has a solution for all $\alpha \neq -1$ and $\beta = 2$

(B) It has no solution if $\alpha = -1$ and $\beta \neq 2$

(C) It has no solution for $\alpha = -1$ and for all $\beta \in \mathbb{R}$

(D) It has no solution for $\alpha = 3$ and for all $\beta \neq 2$

Q107. JEE Main 2023 (Online) 25th January Evening Shift

Open solution

Let A, B, C be 3 × 3 matrices such that A is symmetric and B and C are skew-symmetric. Consider the statements

(S1) $A^{13} B^{26} - B^{26} A^{13}$ is symmetric

(S2) $A^{26} C^{13} - C^{13} A^{26}$ is symmetric

Then,

(A) Only S2 is true

(B) Only S1 is true

(C) Both S1 and S2 are false

(D) Both S1 and S2 are true

Q108. JEE Main 2023 (Online) 25th January Evening Shift

Open solution

Let $A = \begin{bmatrix} \frac{1}{\sqrt{10}} & \frac{3}{\sqrt{10}} \\ \frac{-3}{\sqrt{10}} & \frac{1}{\sqrt{10}} \end{bmatrix}$ and $B = \begin{bmatrix} 1 & -i \\ 0 & 1 \end{bmatrix}$, where $i = \sqrt{-1}$. If $M = A^T B A$, then the inverse of the matrix $AM^{2023}A^T$ is

(A) $\begin{bmatrix} 1 & -2023i \\ 0 & 1 \end{bmatrix}$

(B) $\begin{bmatrix} 1 & 0 \\ 2023i & 1 \end{bmatrix}$

(C) $\begin{bmatrix} 1 & 2023i \\ 0 & 1 \end{bmatrix}$

(D) $\begin{bmatrix} 1 & 0 \\ -2023i & 1 \end{bmatrix}$

Q109. JEE Main 2023 (Online) 25th January Morning Shift

Open solution

Let S_1 and S_2 be respectively the sets of all $a \in \mathbb{R} - \{0\}$ for which the system of linear equations

$ax + 2ay - 3az = 1$

$(2a+1)x + (2a+3)y + (a+1)z = 2$

$(3a+5)x + (a+5)y + (a+2)z = 3$

has unique solution and infinitely many solutions. Then

(A) $n(S_1) = 2$ and S_2 is an infinite set

(B) $S_1 = \Phi$ and $S_2 = \mathbb{R} - \{0\}$

(C) $S_1 = \mathbb{R} - \{0\}$ and $S_2 = \Phi$

(D) S_1 is an infinite set and n(S_2) = 2

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q110. JEE Main 2023 (Online) 24th January Evening Shift

Open solution

If the system of equations

$x + 2y + 3z = 3$

$4x + 3y - 4z = 4$

$8x + 4y - \lambda z = 9 + \mu$

has infinitely many solutions, then the ordered pair (λ, μ) is equal to :

(A) $\left(\frac{72}{5}, \frac{21}{5}\right)$

(B) $\left(-\frac{72}{5}, -\frac{21}{5}\right)$

(C) $\left(-\frac{72}{5}, \frac{21}{5}\right)$

(D) $\left(\frac{72}{5}, -\frac{21}{5}\right)$

Q111. JEE Main 2023 (Online) 24th January Morning Shift

Open solution

If A and B are two non-zero n $\times$ n matrices such that $A^2 + B = A^2B$, then :

(A) $A^2B = I$

(B) $A^2 = I$ or $B = I$

(C) $A^2B = BA^2$

(D) $AB = I$

Q112. JEE Main 2022 (Online) 30th July Evening Shift

Open solution

If the system of linear equations

$\lambda x + y - z = -1$

$x - y - 3z = 2$

$-x + y + z = \mu$

has infinitely many solutions, then the equation of the line passing through the points $(\lambda + 2\mu,\ 2\lambda + \mu)$ and $(1,\ \lambda\mu)$ is

(A) $4x - 6y = -5$

(B) $2x - 2y = 5$

(C) $2x - 2y = -1$

(D) $4x + 6y = 5$

Q113. JEE Main 2022 (Online) 29th July Evening Shift

Open solution

Which of the following matrices can NOT be obtained from the matrix $\begin{bmatrix} -1 & 2 \\ 1 & -1 \end{bmatrix}$ by a single elementary row operation ?

(A) $\begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix}$

(B) $\begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}$

(C) $\begin{bmatrix} -1 & 2 \\ -2 & 7 \end{bmatrix}$

(D) $\begin{bmatrix} -1 & 2 \\ -1 & 3 \end{bmatrix}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q114. JEE Main 2022 (Online) 29th July Evening Shift

If the system of equations

$x + y + z = 6$
$2x + 5y + \alpha z = \beta$
$x + 2y + 3z = 14$

has infinitely many solutions, then $\alpha + \beta$ is equal to

(A) 8 (B) 36 (C) 44 (D) 48

Open solution

Q115. JEE Main 2022 (Online) 29th July Morning Shift

Let A and B be two 3×3 non-zero real matrices such that AB is a zero matrix. Then

(A) the system of linear equations $AX = 0$ has a unique solution

(B) the system of linear equations $AX = 0$ has infinitely many solutions

(C) B is an invertible matrix

(D) $\mathrm{adj}(\mathrm{A})$ is an invertible matrix

Open solution

Q116. JEE Main 2022 (Online) 28th July Evening Shift

Let A and B be any two 3×3 symmetric and skew symmetric matrices respectively. Then which of the following is NOT true?

(A) $\mathrm{A}^4 - \mathrm{B}^4$ is a symmetric matrix

(B) $\mathrm{AB} - \mathrm{BA}$ is a symmetric matrix

(C) $\mathrm{B}^5 - \mathrm{A}^5$ is a skew-symmetric matrix

(D) $\mathrm{AB} + \mathrm{BA}$ is a skew-symmetric matrix

Open solution

Q117. JEE Main 2022 (Online) 27th July Morning Shift

Let $A = \begin{pmatrix} 1 & 2 \\ -2 & -5 \end{pmatrix}$. Let $\alpha, \beta \in \mathbb{R}$ be such that $\alpha A^2 + \beta A = 2I$. Then $\alpha + \beta$ is equal to

(A) -10 (B) -6 (C) 6 (D) 10

Open solution

Q118. JEE Main 2022 (Online) 26th July Evening Shift

Let $A = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $B = \begin{bmatrix} 9^2 & -10^2 & 11^2 \\ 12^2 & 13^2 & -14^2 \\ -15^2 & 16^2 & 17^2 \end{bmatrix}$, then the value of $A'BA$ is:

(A) 1224 (B) 1042 (C) 540 (D) 539

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q119. JEE Main 2022 (Online) 26th July Morning Shift

Open solution

If the system of linear equations.

$8x + y + 4z = -2$

$x + y + z = 0$

$\lambda x - 3y = \mu$

has infinitely many solutions, then the distance of the point $\left(\lambda, \mu, -\frac{1}{2}\right)$ from the plane $8x + y + 4z + 2 = 0$ is :

(A) $3\sqrt{5}$ (B) 4 (C) $\frac{26}{9}$ (D) $\frac{10}{3}$

Q120. JEE Main 2022 (Online) 25th July Evening Shift

Open solution

The number of real values of λ, such that the system of linear equations

$2x - 3y + 5z = 9$

$x + 3y - z = -18$

$3x - y + (\lambda^2 - |\lambda|)z = 16$

has no solutions, is

(A) 0 (B) 1 (C) 2 (D) 4

Q121. JEE Main 2022 (Online) 25th July Morning Shift

Open solution

The number of $\theta \in (0, 4\pi)$ for which the system of linear equations

$3(\sin 3\theta)x - y + z = 2$

$3(\cos 2\theta)x + 4y + 3z = 3$

$6x + 7y + 7z = 9$

has no solution, is :

(A) 6 (B) 7 (C) 8 (D) 9

Q122. JEE Main 2022 (Online) 30th June Morning Shift

Open solution

Let $A = \begin{bmatrix} 1 & -2 & \alpha \\ \alpha & 2 & -1 \end{bmatrix}$ and $B = \begin{bmatrix} 2 & \alpha \\ -1 & 2 \\ 4 & -5 \end{bmatrix}$, $\alpha \in C$. Then the absolute value of the sum of all values of α for which det(AB) = 0 is :

(A) 3 (B) 4 (C) 2 (D) 5

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q123. JEE Main 2022 (Online) 29th June Morning Shift

Open solution

If the system of linear equations

$2x + y - z = 7$

$x - 3y + 2z = 1$

$x + 4y + \delta z = k$, where $\delta, k \in R$ has infinitely many solutions, then $\delta + k$ is equal to:

(A) -3 (B) 3 (C) 6 (D) 9

Q124. JEE Main 2022 (Online) 29th June Morning Shift

Open solution

Let $A = [a_{ij}]$ be a square matrix of order 3 such that $a_{ij} = 2^{j-i}$, for all i, j = 1, 2, 3. Then, the matrix $A^2 + A^3 + ... + A^{10}$ is equal to :

(A) $\left(\frac{3^{10}-3}{2}\right)A$ (B) $\left(\frac{3^{10}-1}{2}\right)A$

(C) $\left(\frac{3^{10}+1}{2}\right)A$ (D) $\left(\frac{3^{10}+3}{2}\right)A$

Q125. JEE Main 2022 (Online) 28th June Morning Shift

Open solution

If the system of linear equations

$2x + 3y - z = -2$

$x + y + z = 4$

$x - y + |\lambda|z = 4\lambda - 4$

where, $\lambda \in R$, has no solution, then

(A) $\lambda = 7$ (B) $\lambda = -7$ (C) $\lambda = 8$ (D) $\lambda^2 = 1$

Q126. JEE Main 2022 (Online) 27th June Morning Shift

Open solution

Let the system of linear equations

$x + 2y + z = 2$,

$\alpha x + 3y - z = \alpha$,

$-\alpha x + y + 2z = -\alpha$

be inconsistent. Then α is equal to :

(A) $\frac{5}{2}$ (B) $-\frac{5}{2}$ (C) $\frac{7}{2}$ (D) $-\frac{7}{2}$

Q127. JEE Main 2022 (Online) 26th June Evening Shift

Open solution

If the system of equations

$\alpha x + y + z = 5$, $x + 2y + 3z = 4$, $x + 3y + 5z = \beta$

has infinitely many solutions, then the ordered pair (α, β) is equal to :

(A) $(1, -3)$ (B) $(-1, 3)$ (C) $(1, 3)$ (D) $(-1, -3)$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q128. JEE Main 2022 (Online) 26th June Morning Shift

Open solution

The ordered pair (a, b), for which the system of linear equations

$3x - 2y + z = b$

$5x - 8y + 9z = 3$

$2x + y + az = -1$

has no solution, is :

(A) $(3, \frac{1}{3})$ (B) $(-3, \frac{1}{3})$

(C) $(-3, -\frac{1}{3})$ (D) $(3, -\frac{1}{3})$

Q129. JEE Main 2022 (Online) 25th June Evening Shift

Open solution

The system of equations

$-kx + 3y - 14z = 25$

$-15x + 4y - kz = 3$

$-4x + y + 3z = 4$

is consistent for all k in the set

(A) R (B) R $-$ {$-11, 13$} (C) R $-$ {13} (D) R $-$ {$-11, 11$}

Q130. JEE Main 2022 (Online) 25th June Morning Shift

Open solution

Let A be a 3×3 real matrix such that

$A\begin{pmatrix}1\\1\\0\end{pmatrix} = \begin{pmatrix}1\\1\\0\end{pmatrix}; A\begin{pmatrix}1\\0\\1\end{pmatrix} = \begin{pmatrix}-1\\0\\1\end{pmatrix}$ and $A\begin{pmatrix}0\\0\\1\end{pmatrix} = \begin{pmatrix}1\\1\\2\end{pmatrix}$.

If $X = (x_1, x_2, x_3)^T$ and I is an identity matrix of order 3, then the system $(A - 2I)X = \begin{pmatrix}4\\1\\1\end{pmatrix}$ has :

(A) no solution (B) infinitely many solutions

(C) unique solution (D) exactly two solutions

Q131. JEE Main 2022 (Online) 25th June Morning Shift

Open solution

Let $A = \begin{bmatrix}0 & -2\\2 & 0\end{bmatrix}$. If M and N are two matrices given by $M = \sum_{k=1}^{10} A^{2k}$ and $N = \sum_{k=1}^{10} A^{2k-1}$ then MN^2 is :

(A) a non-identity symmetric matrix

(B) a skew-symmetric matrix

(C) neither symmetric nor skew-symmetric matrix

(D) an identity matrix

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q132. JEE Main 2022 (Online) 24th June Evening Shift

Open solution

Let the system of linear equations

$x + y + \alpha z = 2$

$3x + y + z = 4$

$x + 2z = 1$

have a unique solution (x^*, y^*, z^*). If (α, x^*), (y^*, α) and $(x^*, -y^*)$ are collinear points, then the sum of absolute values of all possible values of α is

(A) 4 (B) 3 (C) 2 (D) 1

Q133. JEE Main 2022 (Online) 24th June Morning Shift

Open solution

The number of values of α for which the system of equations :

$x + y + z = \alpha$

$\alpha x + 2\alpha y + 3z = -1$

$x + 3\alpha y + 5z = 4$

is inconsistent, is

(A) 0 (B) 1 (C) 2 (D) 3

Q134. JEE Main 2021 (Online) 1st September Evening Shift

Open solution

Consider the system of linear equations

$-x + y + 2z = 0$

$3x - ay + 5z = 1$

$2x - 2y - az = 7$

Let S_1 be the set of all $a \in R$ for which the system is inconsistent and S_2 be the set of all $a \in R$ for which the system has infinitely many solutions. If $n(S_1)$ and $n(S_2)$ denote the number of elements in S_1 and S_2 respectively, then

(A) $n(S_1) = 2, n(S_2) = 2$ (B) $n(S_1) = 1, n(S_2) = 0$

(C) $n(S_1) = 2, n(S_2) = 0$ (D) $n(S_1) = 0, n(S_2) = 2$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q135. JEE Main 2021 (Online) 31st August Evening Shift

If $\alpha + \beta + \gamma = 2\pi$, then the system of equations

$x + (\cos\gamma)y + (\cos\beta)z = 0$

$(\cos\gamma)x + y + (\cos\alpha)z = 0$

$(\cos\beta)x + (\cos\alpha)y + z = 0$

has :

(A) no solution

(B) infinitely many solution

(C) exactly two solutions

(D) a unique solution

Open solution

Q136. JEE Main 2021 (Online) 31st August Morning Shift

If the following system of linear equations

$2x + y + z = 5$

$x - y + z = 3$

$x + y + az = b$

has no solution, then :

(A) $a = -\frac{1}{3}, b \neq \frac{7}{3}$

(B) $a \neq \frac{1}{3}, b = \frac{7}{3}$

(C) $a \neq -\frac{1}{3}, b = \frac{7}{3}$

(D) $a = \frac{1}{3}, b \neq \frac{7}{3}$

Open solution

Q137. JEE Main 2021 (Online) 31st August Morning Shift

If $a_r = \cos\frac{2r\pi}{9} + i\sin\frac{2r\pi}{9}$, r = 1, 2, 3, ..., i = $\sqrt{-1}$, then the determinant $\begin{vmatrix} a_1 & a_2 & a_3 \\ a_4 & a_5 & a_6 \\ a_7 & a_8 & a_9 \end{vmatrix}$ is equal to :

(A) $a_2a_6 - a_4a_8$

(B) a_9

(C) $a_1a_9 - a_3a_7$

(D) a_5

Open solution

Q138. JEE Main 2021 (Online) 27th August Evening Shift

Let $A = \begin{pmatrix} [x+1] & [x+2] & [x+3] \\ [x] & [x+3] & [x+3] \\ [x] & [x+2] & [x+4] \end{pmatrix}$, where [t] denotes the greatest integer less than or equal to t. If det(A) = 192, then the set of values of x is the interval :

(A) [68, 69)

(B) [62, 63)

(C) [65, 66)

(D) [60, 61)

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q139. JEE Main 2021 (Online) 27th August Evening Shift

Open solution

Let $[\lambda]$ be the greatest integer less than or equal to λ. The set of all values of λ for which the system of linear equations

$x + y + z = 4$,

$3x + 2y + 5z = 3$,

$9x + 4y + (28 + [\lambda])z = [\lambda]$ has a solution is :

(A) R

(B) $(-\infty, -9) \cup (-9, \infty)$

(C) $[-9, -8)$

(D) $(-\infty, -9) \cup [-8, \infty)$

Q140. JEE Main 2021 (Online) 27th August Morning Shift

Open solution

If the matrix $A = \begin{pmatrix} 0 & 2 \\ K & -1 \end{pmatrix}$ satisfies $A(A^3 + 3I) = 2I$, then the value of K is :

(A) $\frac{1}{2}$

(B) $-\frac{1}{2}$

(C) -1

(D) 1

Q141. JEE Main 2021 (Online) 26th August Evening Shift

Open solution

Let $A = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 0 \end{pmatrix}$. Then $A^{2025} - A^{2020}$ is equal to :

(A) $A^6 - A$

(B) A^5

(C) $A^5 - A$

(D) A^6

Q142. JEE Main 2021 (Online) 26th August Morning Shift

Open solution

Let $\theta \in \left(0, \frac{\pi}{2}\right)$. If the system of linear equations

$$(1 + \cos^2\theta)x + \sin^2\theta y + 4\sin 3\theta z = 0$$

$$\cos^2\theta x + (1 + \sin^2\theta)y + 4\sin 3\theta z = 0$$

$$\cos^2\theta x + \sin^2\theta y + (1 + 4\sin 3\theta)z = 0$$

has a non-trivial solution, then the value of θ is :

(A) $\frac{4\pi}{9}$

(B) $\frac{7\pi}{18}$

(C) $\frac{\pi}{18}$

(D) $\frac{5\pi}{18}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q143. JEE Main 2021 (Online) 26th August Morning Shift

If $A = \begin{pmatrix} \frac{1}{\sqrt{5}} & \frac{2}{\sqrt{5}} \\ \frac{-2}{\sqrt{5}} & \frac{1}{\sqrt{5}} \end{pmatrix}$, $B = \begin{pmatrix} 1 & 0 \\ i & 1 \end{pmatrix}$, $i = \sqrt{-1}$, and Q = $A^T BA$, then the inverse of the matrix A Q^{2021} A^T is equal to :

Open solution

(A) $\begin{pmatrix} \frac{1}{\sqrt{5}} & -2021 \\ 2021 & \frac{1}{\sqrt{5}} \end{pmatrix}$

(B) $\begin{pmatrix} 1 & 0 \\ -2021i & 1 \end{pmatrix}$

(C) $\begin{pmatrix} 1 & 0 \\ 2021i & 1 \end{pmatrix}$

(D) $\begin{pmatrix} 1 & -2021i \\ 0 & 1 \end{pmatrix}$

Q144. JEE Main 2021 (Online) 27th July Morning Shift

Let $A = \begin{bmatrix} 1 & 2 \\ -1 & 4 \end{bmatrix}$. If $A^{-1} = \alpha I + \beta A$, $\alpha, \beta \in R$, I is a 2×2 identity matrix then $4(\alpha - \beta)$ is equal to :

Open solution

(A) 5 (B) $\frac{8}{3}$ (C) 2 (D) 4

Q145. JEE Main 2021 (Online) 25th July Evening Shift

The number of distinct real roots

of $\begin{vmatrix} \sin x & \cos x & \cos x \\ \cos x & \sin x & \cos x \\ \cos x & \cos x & \sin x \end{vmatrix} = 0$ in the interval $-\frac{\pi}{4} \le x \le \frac{\pi}{4}$ is :

Open solution

(A) 4 (B) 1 (C) 2 (D) 3

Q146. JEE Main 2021 (Online) 25th July Evening Shift

If $P = \begin{bmatrix} 1 & 0 \\ \frac{1}{2} & 1 \end{bmatrix}$, then P^{50} is :

Open solution

(A) $\begin{bmatrix} 1 & 0 \\ 25 & 1 \end{bmatrix}$ (B) $\begin{bmatrix} 1 & 50 \\ 0 & 1 \end{bmatrix}$

(C) $\begin{bmatrix} 1 & 25 \\ 0 & 1 \end{bmatrix}$ (D) $\begin{bmatrix} 1 & 0 \\ 50 & 1 \end{bmatrix}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q147. JEE Main 2021 (Online) 25th July Morning Shift

The values of a and b, for which the system of equations

$2x + 3y + 6z = 8$

$x + 2y + az = 5$

$3x + 5y + 9z = b$

has no solution, are :

(A) $a = 3, b \neq 13$ (B) $a \neq 3, b \neq 13$ (C) $a \neq 3, b = 3$ (D) $a = 3, b = 13$

Open solution

Q148. JEE Main 2021 (Online) 22th July Evening Shift

The values of λ and μ such that the system of equations $x + y + z = 6, 3x + 5y + 5z = 26$, $x + 2y + \lambda z = \mu$ has no solution, are :

(A) $\lambda = 3, \mu = 5$ (B) $\lambda = 3, \mu \neq 10$

(C) $\lambda \neq 2, \mu = 10$ (D) $\lambda = 2, \mu \neq 10$

Open solution

Q149. JEE Main 2021 (Online) 22th July Evening Shift

Let $A = [a_{ij}]$ be a real matrix of order 3×3, such that $a_{i1} + a_{i2} + a_{i3} = 1$, for i = 1, 2, 3. Then, the sum of all the entries of the matrix A^3 is equal to :

(A) 2 (B) 1 (C) 3 (D) 9

Open solution

Q150. JEE Main 2021 (Online) 20th July Evening Shift

The value of $k \in R$, for which the following system of linear equations

$3x - y + 4z = 3,$

$x + 2y - 3z = -2$

$6x + 5y + kz = -3,$

has infinitely many solutions, is :

(A) 3 (B) -5 (C) 5 (D) -3

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q151. JEE Main 2021 (Online) 20th July Morning Shift

Let $A = \begin{bmatrix} 2 & 3 \\ a & 0 \end{bmatrix}$, a∈R be written as P + Q where P is a symmetric matrix and Q is skew symmetric matrix. If det(Q) = 9, then the modulus of the sum of all possible values of determinant of P is equal to :

Open solution

(A) 36 (B) 24 (C) 45 (D) 18

Q152. JEE Main 2021 (Online) 18th March Evening Shift

Let the system of linear equations

$4x + \lambda y + 2z = 0$

$2x - y + z = 0$

$\mu x + 2y + 3z = 0$, $\lambda, \mu \in R$.

has a non-trivial solution. Then which of the following is true?

Open solution

(A) $\mu = 6, \lambda \in R$ (B) $\lambda = 3, \mu \in R$

(C) $\mu = -6, \lambda \in R$ (D) $\lambda = 2, \mu \in R$

Q153. JEE Main 2021 (Online) 18th March Morning Shift

The solutions of the equation $\begin{vmatrix} 1+\sin^2 x & \sin^2 x & \sin^2 x \\ \cos^2 x & 1+\cos^2 x & \cos^2 x \\ 4\sin 2x & 4\sin 2x & 1+4\sin 2x \end{vmatrix} = 0, (0 < x < \pi)$, are

Open solution

(A) $\frac{\pi}{12}, \frac{\pi}{6}$ (B) $\frac{\pi}{6}, \frac{5\pi}{6}$

(C) $\frac{5\pi}{12}, \frac{7\pi}{12}$ (D) $\frac{7\pi}{12}, \frac{11\pi}{12}$

Q154. JEE Main 2021 (Online) 18th March Morning Shift

Let α, β, γ be the real roots of the equation, $x^3 + ax^2 + bx + c = 0$, (a, b, c ∈ R and a, b ≠ 0). If the system of equations (in u, v, w) given by $\alpha u + \beta v + \gamma w = 0$, $\beta u + \gamma v + \alpha w = 0$; $\gamma u + \alpha v + \beta w = 0$ has non-trivial solution, then the value of $\frac{a^2}{b}$ is

Open solution

(A) 5 (B) 3 (C) 1 (D) 0

Q155. JEE Main 2021 (Online) 18th March Morning Shift

Let $A + 2B = \begin{bmatrix} 1 & 2 & 0 \\ 6 & -3 & 3 \\ -5 & 3 & 1 \end{bmatrix}$ and $2A - B = \begin{bmatrix} 2 & -1 & 5 \\ 2 & -1 & 6 \\ 0 & 1 & 2 \end{bmatrix}$. If Tr(A) denotes the sum of all diagonal elements of the matrix A, then Tr(A) − Tr(B) has value equal to

Open solution

(A) 1 (B) 2 (C) 0 (D) 3

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q156. JEE Main 2021 (Online) 17th March Evening Shift

If x, y, z are in arithmetic progression with common difference d, $x \neq 3d$, and the determinant of the matrix $\begin{bmatrix} 3 & 4\sqrt{2} & x \\ 4 & 5\sqrt{2} & y \\ 5 & k & z \end{bmatrix}$ is zero, then the value of k^2 is :

Open solution

(A) 72 (B) 12 (C) 36 (D) 6

Q157. JEE Main 2021 (Online) 17th March Morning Shift

The system of equations kx + y + z = 1, x + ky + z = k and x + y + zk = k^2 has no solution if k is equal to :

Open solution

(A) 0 (B) -1 (C) -2 (D) 1

Q158. JEE Main 2021 (Online) 16th March Evening Shift

The maximum value of

$$f(x) = \begin{vmatrix} \sin^2 x & 1+\cos^2 x & \cos 2x \\ 1+\sin^2 x & \cos^2 x & \cos 2x \\ \sin^2 x & \cos^2 x & \sin 2x \end{vmatrix}, x \in R \text{ is :}$$

Open solution

(A) $\sqrt{5}$ (B) $\frac{3}{4}$ (C) 5 (D) $\sqrt{7}$

Q159. JEE Main 2021 (Online) 16th March Morning Shift

Let $A = \begin{bmatrix} i & -i \\ -i & i \end{bmatrix}, i = \sqrt{-1}$. Then, the system of linear equations $A^8 \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 8 \\ 64 \end{bmatrix}$ has :

Open solution

(A) Exactly two solutions (B) Infinitely many solutions

(C) A unique solution (D) No solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q160. JEE Main 2021 (Online) 26th February Evening Shift

Open solution

Consider the following system of equations :

$x + 2y - 3z = a$

$2x + 6y - 11z = b$

$x - 2y + 7z = c,$

where a, b and c are real constants. Then the system of equations :

(A) has no solution for all a, b and c

(B) has a unique solution when 5a = 2b + c

(C) has infinite number of solutions when 5a = 2b + c

(D) has a unique solution for all a, b and c

Q161. JEE Main 2021 (Online) 26th February Morning Shift

Open solution

Let A be a symmetric matrix of order 2 with integer entries. If the sum of the diagonal elements of A^2 is 1, then the possible number of such matrices is :

(A) 6 (B) 4 (C) 1 (D) 12

Q162. JEE Main 2021 (Online) 26th February Morning Shift

Open solution

The value of $\begin{vmatrix} (a+1)(a+2) & a+2 & 1 \\ (a+2)(a+3) & a+3 & 1 \\ (a+3)(a+4) & a+4 & 1 \end{vmatrix}$ is :

(A) -2

(B) 0

(C) (a + 2)(a + 3)(a + 4)

(D) (a + 1)(a + 2)(a + 3)

Q163. JEE Main 2021 (Online) 25th February Evening Shift

Open solution

Let A be a 3×3 matrix with det(A) = 4. Let R_i denote the ith row of A. If a matrix B is obtained by performing the operation $R_2 \to 2R_2 + 5R_3$ on 2A, then det(B) is equal to :

(A) 64 (B) 16 (C) 128 (D) 80

Q164. JEE Main 2021 (Online) 25th February Evening Shift

Open solution

If for the matrix, $A = \begin{bmatrix} 1 & -\alpha \\ \alpha & \beta \end{bmatrix}$, $AA^T = I_2$, then the value of $\alpha^4 + \beta^4$ is :

(A) 3 (B) 2 (C) 1 (D) 4

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q165. JEE Main 2021 (Online) 25th February Evening Shift

Open solution

The following system of linear equations

$2x + 3y + 2z = 9$

$3x + 2y + 2z = 9$

$x - y + 4z = 8$

(A) does not have any solution

(B) has a solution (α, β, γ) satisfying $\alpha + \beta^2 + \gamma^3 = 12$

(C) has a unique solution

(D) has infinitely many solutions

Q166. JEE Main 2021 (Online) 24th February Evening Shift

Open solution

Let A and B be 3×3 real matrices such that A is symmetric matrix and B is skew-symmetric matrix. Then the system of linear equations $(A^2B^2 - B^2A^2)$ X = O, where X is a 3×1 column matrix of unknown variables and O is a 3×1 null matrix, has :

(A) no solution

(B) exactly two solutions

(C) infinitely many solutions

(D) a unique solution

Q167. JEE Main 2021 (Online) 24th February Evening Shift

Open solution

For the system of linear equations:

$$x - 2y = 1, x - y + kz = -2, ky + 4z = 6, k \in R,$$

consider the following statements :

(A) The system has unique solution if $k \neq 2, k \neq -2$.

(B) The system has unique solution if k = -2

(C) The system has unique solution if k = 2

(D) The system has no solution if k = 2

(E) The system has infinite number of solutions if k $\neq -2$.

Which of the following statements are correct?

(A) (B) and (E) only **(B)** (C) and (D) only **(C)** (A) and (E) only **(D)** (A) and (D) only

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q168. JEE Main 2021 (Online) 24th February Morning Shift

Open solution

The system of linear equations
3x - 2y - kz = 10
2x - 4y - 2z = 6
x+2y - z = 5m
is inconsistent if :

(A) $k \neq 3, m \in \mathbf{R}$
(B) $k = 3, m \neq \frac{4}{5}$
(C) $k = 3, m = \frac{4}{5}$
(D) $k \neq 3, m \neq \frac{4}{5}$

Q169. JEE Main 2020 (Online) 6th September Evening Slot

Open solution

Let $\theta = \frac{\pi}{5}$ and $A = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix}$.

If $B = A + A^4$, then det (B) :

(A) lies in (1, 2)
(B) lies in (2, 3).
(C) is zero.
(D) is one.

Q170. JEE Main 2020 (Online) 6th September Morning Slot

Open solution

The values of λ and μ for which the system of linear equations
x + y + z = 2
x + 2y + 3z = 5
x + 3y + λz = μ
has infinitely many solutions are, respectively:

(A) 6 and 8
(B) 5 and 8
(C) 5 and 7
(D) 4 and 9

Q171. JEE Main 2020 (Online) 6th September Morning Slot

Open solution

Let m and M be respectively the minimum and maximum values of

$$\begin{vmatrix} \cos^2 x & 1+\sin^2 x & \sin 2x \\ 1+\cos^2 x & \sin^2 x & \sin 2x \\ \cos^2 x & \sin^2 x & 1+\sin 2x \end{vmatrix}$$

Then the ordered pair (m, M) is equal to :

(A) (−3, −1)
(B) (−4, −1)
(C) (1, 3)
(D) (−3, 3)

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q172. JEE Main 2020 (Online) 5th September Evening Slot

Open solution

If a + x = b + y = c + z + 1, where a, b, c, x, y, z are non-zero distinct real numbers, then $\begin{vmatrix} x & a+y & x+a \\ y & b+y & y+b \\ z & c+y & z+c \end{vmatrix}$ is equal to :

(A) y(b − a) (B) y(a − b) (C) y(a − c) (D) 0

Q173. JEE Main 2020 (Online) 5th September Evening Slot

Open solution

If the system of linear equations

x + y + 3z = 0

$x + 3y + k^2z = 0$

3x + y + 3z = 0

has a non-zero solution (x, y, z) for some $k \in R$, then $x + \left(\frac{y}{z}\right)$ is equal to :

(A) 9 (B) 3 (C) -9 (D) -3

Q174. JEE Main 2020 (Online) 5th September Morning Slot

Open solution

If the minimum and the maximum values of the function $f : \left[\frac{\pi}{4}, \frac{\pi}{2}\right] \to R$, defined by $f(\theta) = \begin{vmatrix} -\sin^2\theta & -1-\sin^2\theta & 1 \\ -\cos^2\theta & -1-\cos^2\theta & 1 \\ 12 & 10 & -2 \end{vmatrix}$ are m and M respectively, then the ordered pair (m,M) is equal to :

(A) $\left(0, 2\sqrt{2}\right)$ (B) (-4, 0) (C) (-4, 4) (D) (0, 4)

Q175. JEE Main 2020 (Online) 5th September Morning Slot

Open solution

Let $\lambda \in R$. The system of linear equations

$2x_1 - 4x_2 + \lambda x_3 = 1$

$x_1 - 6x_2 + x_3 = 2$

$\lambda x_1 - 10x_2 + 4x_3 = 3$

is inconsistent for:

(A) exactly one positive value of λ (B) exactly one negative value of λ

(C) exactly two values of λ (D) every value of λ

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q176. JEE Main 2020 (Online) 4th September Evening Slot

Open solution

Suppose the vectors x_1, x_2 and x_3 are the solutions of the system of linear equations, Ax = b when the vector b on the right side is equal to b_1, b_2 and b_3 respectively. if

$$x_1 = \begin{bmatrix}1\\1\\1\end{bmatrix}, x_2 = \begin{bmatrix}0\\2\\1\end{bmatrix}, x_3 = \begin{bmatrix}0\\0\\1\end{bmatrix}$$

$$b_1 = \begin{bmatrix}1\\0\\0\end{bmatrix}, b_2 = \begin{bmatrix}0\\2\\0\end{bmatrix} \text{ and } b_3 = \begin{bmatrix}0\\0\\2\end{bmatrix},$$

then the determinant of A is equal to :

(A) $\frac{3}{2}$ (B) 4 (C) 2 (D) $\frac{1}{2}$

Q177. JEE Main 2020 (Online) 4th September Evening Slot

Open solution

If the system of equations
x+y+z=2
2x+4y−z=6
3x+2y+λz=μ
has infinitely many solutions, then

(A) $2\lambda - \mu = 5$ (B) $\lambda - 2\mu = -5$ (C) $2\lambda + \mu = 14$ (D) $\lambda + 2\mu = 14$

Q178. JEE Main 2020 (Online) 4th September Morning Slot

Open solution

If $A = \begin{bmatrix}\cos\theta & i\sin\theta\\ i\sin\theta & \cos\theta\end{bmatrix}, \left(\theta = \frac{\pi}{24}\right)$

and $A^5 = \begin{bmatrix}a & b\\ c & d\end{bmatrix}$, where $i = \sqrt{-1}$ then which one of the following is not true?

(A) $a^2 - c^2 = 1$ (B) $0 \le a^2 + b^2 \le 1$ (C) $a^2 - d^2 = 0$ (D) $a^2 - b^2 = \frac{1}{2}$

Q179. JEE Main 2020 (Online) 3rd September Morning Slot

Open solution

If $\Delta = \begin{vmatrix}x-2 & 2x-3 & 3x-4\\ 2x-3 & 3x-4 & 4x-5\\ 3x-5 & 5x-8 & 10x-17\end{vmatrix}$ =

$Ax^3 + Bx^2 + Cx + D$, then B + C is equal to :

(A) -1 (B) -3 (C) 9 (D) 1

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q180. JEE Main 2020 (Online) 2nd September Evening Slot

Open solution

Let a, b, c $\in$ R be all non-zero and satisfy $a^3 + b^3 + c^3 = 2$. If the matrix

$$A = \begin{pmatrix} a & b & c \\ b & c & a \\ c & a & b \end{pmatrix}$$

satisfies $A^TA = I$, then a value of **abc** can be :

(A) 3 (B) $\frac{1}{3}$ (C) $-\frac{1}{3}$ (D) $\frac{2}{3}$

Q181. JEE Main 2020 (Online) 2nd September Evening Slot

Open solution

Let $A = \{X = (x, y, z)^T : PX = 0$ and

$x^2 + y^2 + z^2 = 1\}$ where

$$P = \begin{bmatrix} 1 & 2 & 1 \\ -2 & 3 & -4 \\ 1 & 9 & -1 \end{bmatrix},$$

then the set A :

(A) is an empty set.
(B) contains more than two elements.
(C) contains exactly two elements.
(D) is a singleton.

Q182. JEE Main 2020 (Online) 2nd September Morning Slot

Open solution

Let S be the set of all $\lambda \in$ R for which the system of linear equations

$2x - y + 2z = 2$
$x - 2y + \lambda z = -4$
$x + \lambda y + z = 4$

has no solution. Then the set S :

(A) contains more than two elements.
(B) contains exactly two elements.
(C) is a singleton.
(D) is an empty set.

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q183. JEE Main 2020 (Online) 2nd September Morning Slot

Let A be a 2×2 real matrix with entries from {0, 1} and $|A| \neq 0$. Consider the following two statements :

(P) If $A \neq I_2$, then $|A| = -1$

(Q) If $|A| = 1$, then $tr(A) = 2$,

where I_2 denotes 2×2 identity matrix and tr(A) denotes the sum of the diagonal entries of A. Then :

(A) (P) is true and (Q) is false

(B) Both (P) and (Q) are false

(C) Both (P) and (Q) are true

(D) (P) is false and (Q) is true

Open solution

Q184. JEE Main 2020 (Online) 9th January Evening Slot

The following system of linear equations

$7x + 6y - 2z = 0$

$3x + 4y + 2z = 0$

$x - 2y - 6z = 0$, has

(A) no solution

(B) infinitely many solutions, (x, y, z) satisfying $y = 2z$

(C) infinitely many solutions, (x, y, z) satisfying $x = 2z$

(D) only the trivial solution

Open solution

Q185. JEE Main 2020 (Online) 9th January Evening Slot

Let $a - 2b + c = 1$.

If $f(x) = \begin{vmatrix} x+a & x+2 & x+1 \\ x+b & x+3 & x+2 \\ x+c & x+4 & x+3 \end{vmatrix}$, then:

(A) f(50) = 1 **(B)** f(−50) = −1 **(C)** f(50) = −501 **(D)** f(−50) = 501

Open solution

Q186. JEE Main 2020 (Online) 9th January Morning Slot

If for some α and β in R, the intersection of the following three planes

$x + 4y - 2z = 1$

$x + 7y - 5z = \beta$

$x + 5y + \alpha z = 5$

is a line in R^3, then $\alpha + \beta$ is equal to :

(A) -10 **(B)** 0 **(C)** 10 **(D)** 2

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q187. JEE Main 2020 (Online) 8th January Evening Slot

The system of linear equations
$\lambda x + 2y + 2z = 5$
$2\lambda x + 3y + 5z = 8$
$4x + \lambda y + 6z = 10$ has

(A) a unique solution when $\lambda = -8$
(B) no solution when $\lambda = 2$
(C) infinitely many solutions when $\lambda = 2$
(D) no solution when $\lambda = 8$

Open solution

Q188. JEE Main 2020 (Online) 8th January Evening Slot

If $A = \begin{pmatrix} 2 & 2 \\ 9 & 4 \end{pmatrix}$ and $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ then $10A^{-1}$ is equal to :

(A) 6I − A
(B) 4I − A
(C) A − 6I
(D) A − 4I

Open solution

Q189. JEE Main 2020 (Online) 8th January Morning Slot

For which of the following ordered pairs (μ, δ), the system of linear equations
$x + 2y + 3z = 1$
$3x + 4y + 5z = \mu$
$4x + 4y + 4z = \delta$
is inconsistent ?

(A) (1, 0)
(B) (4, 3)
(C) (4, 6)
(D) (3, 4)

Open solution

Q190. JEE Main 2020 (Online) 7th January Morning Slot

Let α be a root of the equation $x^2 + x + 1 = 0$ and the
matrix $A = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & 1 & 1 \\ 1 & \alpha & \alpha^2 \\ 1 & \alpha^2 & \alpha^4 \end{bmatrix}$

then the matrix A^{31} is equal to

(A) A^2
(B) A
(C) I_3
(D) A^3

Open solution

Q191. JEE Main 2020 (Online) 7th January Morning Slot

If the system of linear equations
$2x + 2ay + az = 0$
$2x + 3by + bz = 0$
$2x + 4cy + cz = 0,$
where a, b, c $\in$ R are non-zero distinct; has a non-zero solution, then:

(A) $\frac{1}{a}, \frac{1}{b}, \frac{1}{c}$ are in A.P.
(B) a + b + c = 0
(C) a, b, c are in G.P.
(D) a,b,c are in A.P.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q192. JEE Main 2019 (Online) 12th April Evening Slot

Open solution

A value of $\theta \in \left(0, \frac{\pi}{3}\right)$, for which $\begin{vmatrix} 1+\cos^2\theta & \sin^2\theta & 4\cos 6\theta \\ \cos^2\theta & 1+\sin^2\theta & 4\cos 6\theta \\ \cos^2\theta & \sin^2\theta & 1+4\cos 6\theta \end{vmatrix} = 0$, is :

(A) $\frac{\pi}{18}$ (B) $\frac{\pi}{9}$ (C) $\frac{7\pi}{24}$ (D) $\frac{7\pi}{36}$

Q193. JEE Main 2019 (Online) 12th April Morning Slot

Open solution

If A is a symmetric matrix and B is a skew-symmetric matrix such that A + B = $\begin{bmatrix} 2 & 3 \\ 5 & -1 \end{bmatrix}$, then AB is equal to :

(A) $\begin{bmatrix} 4 & -2 \\ 1 & -4 \end{bmatrix}$

(B) $\begin{bmatrix} -4 & -2 \\ -1 & 4 \end{bmatrix}$

(C) $\begin{bmatrix} -4 & 2 \\ 1 & 4 \end{bmatrix}$

(D) $\begin{bmatrix} 4 & -2 \\ -1 & -4 \end{bmatrix}$

Q194. JEE Main 2019 (Online) 12th April Morning Slot

Open solution

If $B = \begin{bmatrix} 5 & 2\alpha & 1 \\ 0 & 2 & 1 \\ \alpha & 3 & -1 \end{bmatrix}$ is the inverse of a 3 × 3 matrix A, then the sum of all values of α for which det(A) + 1 = 0, is :

(A) 2 (B) - 1 (C) 0 (D) 1

Q195. JEE Main 2019 (Online) 10th April Evening Slot

Open solution

Let λ be a real number for which the system of linear equations x + y + z = 6, 4x + λy − λz = λ − 2, 3x + 2y − 4z = − 5 has infinitely many solutions. Then λ is a root of the quadratic equation:

(A) λ^2 + λ - 6 = 0 (B) λ^2 - λ - 6 = 0

(C) λ^2 - 3λ - 4 = 0 (D) λ^2 + 3λ - 4 = 0

Q196. JEE Main 2019 (Online) 10th April Evening Slot

Open solution

The sum of the real roots of the equation $\begin{vmatrix} x & -6 & -1 \\ 2 & -3x & x-3 \\ -3 & 2x & x+2 \end{vmatrix} = 0$, is equal to :

(A) - 4 (B) 0 (C) 1 (D) 6

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q197. JEE Main 2019 (Online) 10th April Morning Slot

Open solution

If $\Delta_1 = \begin{vmatrix} x & \sin\theta & \cos\theta \\ -\sin\theta & -x & 1 \\ \cos\theta & 1 & x \end{vmatrix}$ and

$\Delta_2 = \begin{vmatrix} x & \sin 2\theta & \cos 2\theta \\ -\sin 2\theta & -x & 1 \\ \cos 2\theta & 1 & x \end{vmatrix}$, $x \neq 0$;

then for all $\theta \in \left(0, \frac{\pi}{2}\right)$:

(A) $\Delta_1 - \Delta_2$ = x (cos 2θ − cos 4θ)

(B) $\Delta_1 + \Delta_2$ = - 2x^3

(C) $\Delta_1 + \Delta_2$ = − 2(x^3 + x −1)

(D) $\Delta_1 - \Delta_2$ = - 2x^3

Q198. JEE Main 2019 (Online) 10th April Morning Slot

Open solution

If the system of linear equations

x + y + z = 5

x + 2y + 2z = 6

x + 3y + λz = μ, (λ, $\mu \in$ R), has infinitely many solutions, then the value of $\lambda + \mu$ is :

(A) 10 (B) 9 (C) 7 (D) 12

Q199. JEE Main 2019 (Online) 9th April Evening Slot

Open solution

If the system of equations 2x + 3y − z = 0, x + ky − 2z = 0 and 2x − y + z = 0 has a non-trival solution (x, y, z), then $\frac{x}{y} + \frac{y}{z} + \frac{z}{x} + k$ is equal to :-

(A) -4 (B) $\frac{3}{4}$ (C) $\frac{1}{2}$ (D) $-\frac{1}{4}$

Q200. JEE Main 2019 (Online) 9th April Evening Slot

Open solution

The total number of matrices

$A = \begin{pmatrix} 0 & 2y & 1 \\ 2x & y & -1 \\ 2x & -y & 1 \end{pmatrix}$

(x, y $\in$ R,x $\neq$ y) for which A^{T}A = 3I$_3$ is :-

(A) 3 (B) 4 (C) 2 (D) 6

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q201. JEE Main 2019 (Online) 9th April Morning Slot

Open solution

Let α and β be the roots of the equation $x^2 + x + 1 = 0$. Then for $y \neq 0$ in R,

$$\begin{vmatrix} y+1 & \alpha & \beta \\ \alpha & y+\beta & 1 \\ \beta & 1 & y+\alpha \end{vmatrix}$$

is equal to

(A) $y(y^2 - 1)$ (B) $y(y^2 - 3)$ (C) y^3 (D) $y^3 - 1$

Q202. JEE Main 2019 (Online) 9th April Morning Slot

Open solution

If $\begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}\begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix}\begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix}\ldots\begin{bmatrix} 1 & n-1 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 78 \\ 0 & 1 \end{bmatrix}$,

then the inverse of $\begin{bmatrix} 1 & n \\ 0 & 1 \end{bmatrix}$ is

(A) $\begin{bmatrix} 1 & 0 \\ 12 & 1 \end{bmatrix}$ (B) $\begin{bmatrix} 1 & 0 \\ 13 & 1 \end{bmatrix}$

(C) $\begin{bmatrix} 1 & -13 \\ 0 & 1 \end{bmatrix}$ (D) $\begin{bmatrix} 1 & -12 \\ 0 & 1 \end{bmatrix}$

Q203. JEE Main 2019 (Online) 8th April Evening Slot

Open solution

If the system of linear equations

$x - 2y + kz = 1$
$2x + y + z = 2$
$3x - y - kz = 3$

has a solution (x,y,z), $z \neq 0$, then (x,y) lies on the straight line whose equation is :

(A) $4x - 3y - 4 = 0$ (B) $3x - 4y - 1 = 0$ (C) $4x - 3y - 1 = 0$ (D) $3x - 4y - 4 = 0$

Q204. JEE Main 2019 (Online) 8th April Evening Slot

Open solution

Let the number 2,b,c be in an A.P. and

A = $\begin{bmatrix} 1 & 1 & 1 \\ 2 & b & c \\ 4 & b^2 & c^2 \end{bmatrix}$. If $\det(A) \in [2, 16]$, then c lies in the interval :

(A) [2, 3) (B) [4, 6] (C) $(2 + 2^{3/4}, 4)$ (D) $[3, 2 + 2^{3/4}]$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q205. JEE Main 2019 (Online) 8th April Morning Slot

Let $A = \begin{pmatrix} \cos\alpha & -\sin\alpha \\ \sin\alpha & \cos\alpha \end{pmatrix}$, $(\alpha \in R)$

such that $A^{32} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ then a value of α is

Open solution

(A) 0 (B) $\frac{\pi}{16}$ (C) $\frac{\pi}{32}$ (D) $\frac{\pi}{64}$

Q206. JEE Main 2019 (Online) 8th April Morning Slot

The greatest value of c $\in$ R for which the system of linear equations

x − cy − cz = 0

cx − y + cz = 0

cx + cy − z = 0

has a non-trivial solution, is :

Open solution

(A) -1 (B) 0 (C) 1/2 (D) 2

Q207. JEE Main 2019 (Online) 12th January Evening Slot

If A = $\begin{bmatrix} 1 & \sin\theta & 1 \\ -\sin\theta & 1 & \sin\theta \\ -1 & -\sin\theta & 1 \end{bmatrix}$;

Open solution

then for all $\theta \in \left(\frac{3\pi}{4}, \frac{5\pi}{4}\right)$, det (A) lies in the interval :

(A) $\left(\frac{3}{2}, 3\right]$ (B) $\left(0, \frac{3}{2}\right]$ (C) $\left[\frac{5}{2}, 4\right)$ (D) $\left(1, \frac{5}{2}\right]$

Q208. JEE Main 2019 (Online) 12th January Evening Slot

The set of all values of λ for which the system of linear equations

x − 2y − 2z = λx

x + 2y + z = λy

− x − y = λz

has a non-trivial solution :

Open solution

(A) is an empty set (B) contains more than two elements

(C) is a singleton (D) contains exactly two elements

Q209. JEE Main 2019 (Online) 12th January Morning Slot

An ordered pair (α, β) for which the system of linear equations

$(1 + \alpha)$ x + βy + z = 2

αx + $(1 + \beta)$y + z = 3

αx + βy + 2z = 2

has a unique solution, is :

Open solution

(A) (−3, 1) (B) (1, −3) (C) (−4, 2) (D) (2, 4)

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q210. JEE Main 2019 (Online) 12th January Morning Slot

Let $P = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 9 & 3 & 1 \end{bmatrix}$ and $Q = [q_{ij}]$ be two 3×3 matrices such that $Q - P^5 = I_3$.

Open solution

Then $\frac{q_{21}+q_{31}}{q_{32}}$ is equal to :

(A) 15 (B) 9 (C) 135 (D) 10

Q211. JEE Main 2019 (Online) 11th January Evening Slot

If $\begin{vmatrix} a-b-c & 2a & 2a \\ 2b & b-c-a & 2b \\ 2c & 2c & c-a-b \end{vmatrix}$

$= (a + b + c)(x + a + b + c)^2$, $x \neq 0$,

Open solution

then x is equal to :

(A) $-2(a + b + c)$ (B) $2(a + b + c)$ (C) abc (D) $-(a + b + c)$

Q212. JEE Main 2019 (Online) 11th January Morning Slot

If the system of linear equations

$2x + 2y + 3z = a$

$3x - y + 5z = b$

$x - 3y + 2z = c$

where a, b, c are non zero real numbers, has more than one solution, then :

Open solution

(A) $b - c - a = 0$ (B) $a + b + c = 0$ (C) $b - c + a = 0$ (D) $b + c - a = 0$

Q213. JEE Main 2019 (Online) 11th January Morning Slot

Let $A = \begin{pmatrix} 0 & 2q & r \\ p & q & -r \\ p & -q & r \end{pmatrix}$. If $AA^T = I_3$, then $|p|$ is :

Open solution

(A) $\frac{1}{\sqrt{2}}$ (B) $\frac{1}{\sqrt{5}}$ (C) $\frac{1}{\sqrt{6}}$ (D) $\frac{1}{\sqrt{3}}$

Q214. JEE Main 2019 (Online) 10th January Evening Slot

Let $A = \begin{bmatrix} 2 & b & 1 \\ b & b^2+1 & b \\ 1 & b & 2 \end{bmatrix}$ where $b > 0$.

Open solution

Then the minimum value of $\frac{\det(A)}{b}$ is -

(A) $\sqrt{3}$ (B) $-2\sqrt{3}$ (C) $-\sqrt{3}$ (D) $2\sqrt{3}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q215. JEE Main 2019 (Online) 10th January Evening Slot

The number of values of $\theta \in (0, \pi)$ for which the system of linear equations

$x + 3y + 7z = 0$

$-x + 4y + 7z = 0$

$(\sin 3\theta)x + (\cos 2\theta)y + 2z = 0.$

has a non-trivial solution, is -

(A) two (B) one (C) four (D) three

Open solution

Q216. JEE Main 2019 (Online) 10th January Morning Slot

If the system of equations

$x + y + z = 5$

$x + 2y + 3z = 9$

$x + 3y + \alpha z = \beta$

has infinitely many solutions, then $\beta - \alpha$ equals -

(A) 8 (B) 21 (C) 18 (D) 5

Open solution

Q217. JEE Main 2019 (Online) 10th January Morning Slot

Let $d \in R$, and

$$A = \begin{bmatrix} -2 & 4+d & (\sin\theta)-2 \\ 1 & (\sin\theta)+2 & d \\ 5 & (2\sin\theta)-d & (-\sin\theta)+2+2d \end{bmatrix},$$

$\theta \in [0, 2\pi]$ If the minimum value of det(A) is 8, then a value of d is -

(A) -7 (B) $2\left(\sqrt{2}+2\right)$ (C) -5 (D) $2\left(\sqrt{2}+1\right)$

Open solution

Q218. JEE Main 2019 (Online) 9th January Evening Slot

If the system of linear equations

$x - 4y + 7z = g$

$3y - 5z = h$

$-2x + 5y - 9z = k$

is consistent, then :

(A) $g + 2h + k = 0$ (B) $g + h + 2k = 0$ (C) $2g + h + k = 0$ (D) $g + h + k = 0$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q219. JEE Main 2019 (Online) 9th January Evening Slot

If $A=\begin{bmatrix} e^{t} & e^{-t}\cos t & e^{-t}\sin t \\ e^{t} & -e^{-t}\cos t - e^{-t}\sin t & -e^{-t}\sin t + e^{-t}cost \\ e^{t} & 2e^{-t}\sin t & -2e^{-t}\cos t \end{bmatrix}$

then A is :

(A) invertible for all t∈**R**.

(B) invertible only if $t=\pi$

(C) not invertible for any t∈**R**

(D) invertible only if $t=\frac{\pi}{2}$.

Open solution

Q220. JEE Main 2019 (Online) 9th January Morning Slot

If $A=\begin{bmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{bmatrix}$, then the matrix A^{-50} when $\theta=\frac{\pi}{12}$, is equal to :

(A) $\begin{bmatrix} \frac{\sqrt{3}}{2} & -\frac{1}{2} \\ \frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$

(B) $\begin{bmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{-1}{2} \end{bmatrix}$

(C) $\begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$

(D) $\begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix}$

Open solution

Q221. JEE Main 2019 (Online) 9th January Morning Slot

The system of linear equations

x + y + z = 2

2x + 3y + 2z = 5

$2x + 3y + (a^2 - 1)z = a + 1$ then

(A) has infinitely many solutions for a = 4

(B) has a unique solution for $|a| = \sqrt{3}$

(C) is inconsistent when $|a| = \sqrt{3}$

(D) is inconsistent when a = 4

Open solution

Q222. JEE Main 2018 (Online) 16th April Morning Slot

Let $A=\begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix}$ and $B = A^{20}$. Then the sum of the elements of the first column of B is :

(A) 210 (B) 211 (C) 231 (D) 251

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q223. JEE Main 2018 (Online) 16th April Morning Slot

The number of values of k for which the system of linear equations,
(k + 2)x + 10y = k
kx + (k +3)y = k -1
has **no solution,** is :

(A) 1 (B) 2 (C) 3 (D) infinitely many

Open solution

Q224. JEE Main 2018 (Offline)

If the system of linear equations

x + ky + 3z = 0
3x + ky - 2z = 0
2x + 4y - 3z = 0

has a non-zero solution (x, y, z), then $\frac{xz}{y^2}$ is equal to

(A) 30 (B) -10 (C) 10 (D) -30

Open solution

Q225. JEE Main 2018 (Offline)

If $\begin{vmatrix} x-4 & 2x & 2x \\ 2x & x-4 & 2x \\ 2x & 2x & x-4 \end{vmatrix} = (A+Bx)(x-A)^2$

then the ordered pair (A, B) is equal to :

(A) (4, 5) (B) (-4, -5) (C) (-4, 3) (D) (-4, 5)

Open solution

Q226. JEE Main 2018 (Online) 15th April Evening Slot

Suppose A is any 3$\times$ 3 non-singular matrix and (A $-$ 3I) (A $-$ 5I) = O where I = I_3 and O = O_3. If αA + βA^{-1} = 4I, then $\alpha + \beta$ is equal to :

(A) 8 (B) 7 (C) 13 (D) 12

Open solution

Q227. JEE Main 2018 (Online) 15th April Evening Slot

If the system of linear equations
x + ay + z = 3
x + 2y + 2z = 6
x + 5y + 3z = b
has no solution, then :

(A) a = $-$ 1, b = 9 (B) a = $-$ 1, b $\neq$ 9
(C) a $\neq -$ 1, b = 9 (D) a = 1, b $\neq$ 9

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q228. JEE Main 2018 (Online) 15th April Morning Slot

Open solution

Let A be a matrix such that $A.\begin{bmatrix}1 & 2\\ 0 & 3\end{bmatrix}$ is a scalar matrix and |3A| = 108.

Then A^2 equals :

(A) $\begin{bmatrix}4 & -32\\ 0 & 36\end{bmatrix}$

(B) $\begin{bmatrix}36 & 0\\ -32 & 4\end{bmatrix}$

(C) $\begin{bmatrix}4 & 0\\ -32 & 36\end{bmatrix}$

(D) $\begin{bmatrix}36 & -32\\ 0 & 4\end{bmatrix}$

Q229. JEE Main 2018 (Online) 15th April Morning Slot

Open solution

Let S be the set of all real values of k for which the systemof linear equations

x + y + z = 2

2x + y − z = 3

3x + 2y + kz = 4

has a unique solution. Then S is :

(A) an empty set (B) equal to {0} (C) equal to **R** (D) equal to **R** − {0}

Q230. JEE Main 2017 (Online) 9th April Morning Slot

Open solution

For two 3 × 3 matrices A and B, let A + B = $2B^T$ and 3A + 2B = I_3, where B^T is the transpose of B and I_3 is 3 × 3 identity matrix. Then :

(A) 5A + 10B = $2I_3$ (B) 10A + 5B = $3I_3$ (C) B + 2A = I_3 (D) 3A + 6B = $2I_3$

Q231. JEE Main 2017 (Online) 8th April Morning Slot

Open solution

The number of real values of λ for which the system of linear equations

2x + 4y − λz = 0

4x + λy + 2z = 0

λx + 2y + 2z = 0

has infinitely many solutions, is :

(A) 0 (B) 1 (C) 2 (D) 3

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q232. JEE Main 2017 (Online) 8th April Morning Slot

Let A be any 3 × 3 invertible matrix. Then which one of the following is **not** always true ?

(A) adj (A) = |A|.A^{-1}

(B) adj (adj(A)) = |A|.A

(C) adj (adj(A)) = $|A|^2$.$(adj(A))^{-1}$

(D) adj (adj(A)) = | A | .$(adj(A))^{-1}$

Open solution

Q233. JEE Main 2017 (Online) 8th April Morning Slot

If

$$S = \left\{ x \in [0, 2\pi] : \begin{vmatrix} 0 & \cos x & -\sin x \\ \sin x & 0 & \cos x \\ \cos x & \sin x & 0 \end{vmatrix} = 0 \right\},$$

then $\sum_{x \in S} \tan\left(\frac{\pi}{3} + x\right)$ is equal to :

(A) $4 + 2\sqrt{3}$ (B) $-2 + \sqrt{3}$ (C) $-2 - \sqrt{3}$ (D) $-4 - 2\sqrt{3}$

Open solution

Q234. JEE Main 2017 (Offline)

If $A = \begin{bmatrix} 2 & -3 \\ -4 & 1 \end{bmatrix}$,

then adj($3A^2$ + 12A) is equal to

(A) $\begin{bmatrix} 51 & 63 \\ 84 & 72 \end{bmatrix}$

(B) $\begin{bmatrix} 51 & 84 \\ 63 & 72 \end{bmatrix}$

(C) $\begin{bmatrix} 72 & -63 \\ -84 & 51 \end{bmatrix}$

(D) $\begin{bmatrix} 72 & -84 \\ -63 & 51 \end{bmatrix}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q235. JEE Main 2017 (Offline)

Open solution

If S is the set of distinct values of 'b' for which the following system of linear equations

$x + y + z = 1$
$x + ay + z = 1$
$ax + by + z = 0$

has no solution, then S is :

(A) an empty set

(B) an infinite set

(C) a finite set containing two or more elements

(D) a singleton

Q236. JEE Main 2016 (Online) 10th April Morning Slot

Open solution

Let A be a 3×3 matrix such that $A^2 - 5A + 7I = 0$

Statement - I :

$A^{-1} = \frac{1}{7}(5I - A)$.

Statement - II :

The polynomial $A^3 - 2A^2 - 3A + I$ can be reduced to $5(A - 4I)$.

Then :

(A) Statement-I is true, but Statement-II is false.

(B) Statement-I is false, but Statement-II is true.

(C) Both the statements are true.

(D) Both the statements are false

Q237. JEE Main 2016 (Online) 10th April Morning Slot

Open solution

If $A = \begin{bmatrix} -4 & -1 \\ 3 & 1 \end{bmatrix}$,

then the determinant of the matrix $(A^{2016} - 2A^{2015} - A^{2014})$ is :

(A) 2014 (B) -175 (C) 2016 (D) -25

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q238. JEE Main 2016 (Online) 9th April Morning Slot

If P = $\begin{bmatrix} \frac{\sqrt{3}}{2} & \frac{1}{2} \\ -\frac{1}{2} & \frac{\sqrt{3}}{2} \end{bmatrix}$, $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$

Open solution

Q = PAP^T, then P^T Q^{2015} P is :

(A) $\begin{bmatrix} 0 & 2015 \\ 0 & 0 \end{bmatrix}$

(B) $\begin{bmatrix} 2015 & 1 \\ 0 & 2015 \end{bmatrix}$

(C) $\begin{bmatrix} 2015 & 0 \\ 1 & 2015 \end{bmatrix}$

(D) $\begin{bmatrix} 1 & 2015 \\ 0 & 1 \end{bmatrix}$

Q239. JEE Main 2016 (Online) 9th April Morning Slot

The number of distinct real roots of the equation,

$\begin{vmatrix} \cos x & \sin x & \sin x \\ \sin x & \cos x & \sin x \\ \sin x & \sin x & \cos x \end{vmatrix} = 0$ in the interval $\left[-\frac{\pi}{4}, \frac{\pi}{4}\right]$ is :

Open solution

(A) 4 (B) 3 (C) 2 (D) 1

Q240. JEE Main 2016 (Offline)

If $A = \begin{bmatrix} 5a & -b \\ 3 & 2 \end{bmatrix}$ and A adj $A = A\,A^T$, then $5a + b$ is equal to :

Open solution

(A) 4 (B) 13 (C) -1 (D) 5

Q241. JEE Main 2016 (Offline)

The system of linear equations

$x + \lambda y - z = 0$

$\lambda x - y - z = 0$

$x + y - \lambda z = 0$

has a non-trivial solution for :

Open solution

(A) infinitely many values of λ.

(B) exactly one value of λ.

(C) exactly two values of λ.

(D) exactly three values of λ.

Q242. JEE Main 2015 (Online) 11th April Morning Slot

If A is a 3 $\times$ 3 matrix such that $|5.\,adjA|$ = 5, then $|A|$ is equal to :

Open solution

(A) $\pm\,\frac{1}{5}$ (B) $\pm\,5$ (C) $\pm\,1$ (D) $\pm\,\frac{1}{25}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q243. JEE Main 2015 (Online) 10th April Morning Slot

Open solution

The least value of the product xyz for which the determinant

$$\begin{vmatrix} x & 1 & 1 \\ 1 & y & 1 \\ 1 & 1 & z \end{vmatrix}$$ is non-negative, is :

(A) $-2\sqrt{2}$ (B) $-16\sqrt{2}$ (C) -8 (D) -1

Q244. JEE Main 2015 (Online) 10th April Morning Slot

Open solution

If $A = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$,

then which one of the following statements is **not** correct ?

(A) $A^4 - I = A^2 + I$

(B) $A^3 - I = A(A - I)$

(C) $A^2 + I = A(A^2 - I)$

(D) $A^3 + I = A(A^3 - I)$

Q245. JEE Main 2015 (Offline)

Open solution

If $A = \begin{bmatrix} 1 & 2 & 2 \\ 2 & 1 & -2 \\ a & 2 & b \end{bmatrix}$ is a matrix satisfying the equation

$AA^T = 9I$, where I is 3×3 identity matrix, then the ordered

pair (a, b) is equal to :

(A) $(2, 1)$ (B) $(-2, -1)$ (C) $(2, -1)$ (D) $(-2, 1)$

Q246. JEE Main 2015 (Offline)

Open solution

The set of all values of λ for which the system of linear equations:

$$\begin{aligned} 2x_1 - 2x_2 + x_3 &= \lambda x_1 \\ 2x_1 - 3x_2 + 2x_3 &= \lambda x_2 \\ -x_1 + 2x_2 &= \lambda x_3 \end{aligned}$$

has a non-trivial solution

(A) contains two elements

(B) contains more than two elements

(C) in an empty set

(D) is a singleton

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q247. JEE Main 2014 (Online) 19th April Morning Slot

Let A and B be any two 3×3 matrices. If A is symmetric and B is skewsymmetric, then the matrix AB $-$ BA is :

Open solution

(A) skewsymmetric

(B) symmetric

(C) neither symmetric nor skewsymmetric

(D) I or $-$ I, where I is an identity matrix.

Q248. JEE Main 2014 (Online) 12th April Morning Slot

Open solution

If $A = \begin{bmatrix} 1 & 2 & x \\ 3 & -1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} y \\ x \\ 1 \end{bmatrix}$ be such that $AB = \begin{bmatrix} 6 \\ 8 \end{bmatrix}$, then :

(A) y = 2x (B) y = $-$ 2x (C) y = x (D) y = $-$ x

Q249. JEE Main 2014 (Online) 12th April Morning Slot

Open solution

If $f(\theta) = \begin{vmatrix} 1 & \cos\theta & 1 \\ -\sin\theta & 1 & -\cos\theta \\ -1 & \sin\theta & 1 \end{vmatrix}$ and A and B are respectively the maximum and the minimum values of f (θ), then (A, B) is equal to :

(A) (3, $-$ 1) (B) $(4, 2-\sqrt{2})$

(C) $(2+\sqrt{2}, 2-\sqrt{2})$ (D) $(2+\sqrt{2}, -1)$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q250. JEE Main 2014 (Online) 11th April Morning Slot

Open solution

Let A be a 3×3 matrix such that

$$A\begin{bmatrix}1 & 2 & 3\\ 0 & 2 & 3\\ 0 & 1 & 1\end{bmatrix} = \begin{bmatrix}0 & 0 & 1\\ 1 & 0 & 0\\ 0 & 1 & 0\end{bmatrix}$$

Then A^{-1} is :

(A) $\begin{bmatrix}3 & 1 & 2\\ 3 & 0 & 2\\ 1 & 0 & 1\end{bmatrix}$

(B) $\begin{bmatrix}3 & 2 & 1\\ 3 & 2 & 0\\ 1 & 1 & 0\end{bmatrix}$

(C) $\begin{bmatrix}0 & 1 & 3\\ 0 & 2 & 3\\ 1 & 1 & 1\end{bmatrix}$

(D) $\begin{bmatrix}1 & 2 & 3\\ 0 & 1 & 1\\ 0 & 2 & 3\end{bmatrix}$

Q251. JEE Main 2014 (Online) 9th April Morning Slot

Open solution

If a, b, c are non - zero real numbers and if the system of equations

$(a - 1)x = y + z,$

$(b - 1)y = z + x,$

$(c - 1)z = x + y,$

has a non - trivial solution, then ab + bc + ca equals :

(A) a + b + c (B) abc (C) 1 (D) -1

Q252. JEE Main 2014 (Online) 9th April Morning Slot

Open solution

If B is a 3×3 matrix such that $B^2 = 0$,

then det. $[(I + B)^{50} - 50B]$ is equal to :

(A) 1 (B) 2 (C) 3 (D) 50

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q253. JEE Main 2014 (Offline)

Open solution

If $\alpha, \beta \neq 0$, and $f(n) = \alpha^n + \beta^n$ and

$$\begin{vmatrix} 3 & 1+f(1) & 1+f(2) \\ 1+f(1) & 1+f(2) & 1+f(3) \\ 1+f(2) & 1+f(3) & 1+f(4) \end{vmatrix}$$

$= K(1-\alpha)^2(1-\beta)^2(\alpha-\beta)^2$, then K is equal to :

(A) 1 (B) -1 (C) $\alpha\beta$ (D) $\frac{1}{\alpha\beta}$

Q254. JEE Main 2014 (Offline)

Open solution

If A is a 3×3 non-singular matrix such that $AA' = A'A$ and $B = A^{-1}A'$, then BB' equals:

(A) B^{-1} (B) $\left(B^{-1}\right)'$ (C) $I + B$ (D) I

Q255. JEE Main 2013 (Online) 25th April Morning Slot

Open solution

Let S = $\left\{ \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix} : a_{ij} \in \{0,1,2\}, a_{11} = a_{22} \right\}$

Then the number of non-singular matrices in the set S is:

(A) 27 (B) 24 (C) 10 (D) 20

Q256. JEE Main 2013 (Online) 25th April Morning Slot

Open solution

Consider the system of equations:
x + ay = 0, y + az = 0 and z + ax = 0. Then the set of all real values of 'a' for which the system has a unique solution is

(A) R - {1} (B) R - {-1} (C) {1, -1} (D) {1, 0, -1}

Q257. JEE Main 2013 (Online) 23th April Morning Slot

Open solution

Let A, other than I or -I, be a 2 $\times$ 2 real matrix such that A^2 = I, I being the unit matrix. Let Tr(A) be the sum of diagonal elements of A.
Statement 1: Tr(A) = 0
Statement 2: det(A) = -1

(A) Statement 1 is true; Statement 2 is false

(B) Statement 1 is true; Statement 2 is true; Statement 2 is **not** a correct explanation for Statement 1

(C) Statement 1 is true; Statement 2 is true; Statement 2 is a correct explanation for Statement 1

(D) Statement 1 is false; Statement 2 is true

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q258. JEE Main 2013 (Online) 23th April Morning Slot

Open solution

Statement 1: The system of linear equations
x + (sin α) y + (cos α) z = 0
x + (cos α) y + (sin α) z = 0
x - (sin α) y - (cos α) z = 0
has a non-trivial solution for only one value of α lying in the interval $\left(0, \frac{\pi}{2}\right)$

Statement 2: The equation in α

$$\begin{vmatrix} \cos\alpha & \sin\alpha & \cos\alpha \\ \sin\alpha & \cos\alpha & \sin\alpha \\ \cos\alpha & -\sin\alpha & -\cos\alpha \end{vmatrix} = 0$$

has only one solution lying in the interval $\left(0, \frac{\pi}{2}\right)$

(A) Statement 1 is true; Statement 2 is true; Statement 2 is **not** correct explanation for Statement 1

(B) Statement 1 is true; Statement 2 is true; Statement 2 is correct explanation for Statement 1

(C) Statement 1 is true; Statement 2 is false

(D) Statement 1 is false; Statement 2 is true

Q259. JEE Main 2013 (Online) 22th April Morning Slot

Open solution

If p, q, r are 3 real numbers satisfying the matrix equation, $[pqr]\begin{bmatrix} 3 & 4 & 1 \\ 3 & 2 & 3 \\ 2 & 0 & 2 \end{bmatrix} = [3 \quad 0 \quad 1]$ then 2p + q - r equals

(A) -3 (B) -1 (C) 4 (D) 2

Q260. JEE Main 2013 (Online) 22th April Morning Slot

Open solution

If the system of linear equations:
$x_1 + 2x_2 + 3x_3 = 6$
$x_1 + 3x_2 + 5x_3 = 9$
$2x_1 + 5x_2 + ax_3 = b$
is consistent and has infinite number of solutions, then:

(A) a = 8, b can be any real number

(B) b = 15, a can be any real number

(C) a $\in$ R - {8} and b $\in$ R - {15}

(D) a = 8, b = 15

Q261. JEE Main 2013 (Online) 9th April Morning Slot

Open solution

The matrix $A^2 + 4A - 5I$, where I is identity matrix and $A = \begin{bmatrix} 1 & 2 \\ 4 & -3 \end{bmatrix}$, equals

(A) $4\begin{bmatrix} 2 & 1 \\ 2 & 0 \end{bmatrix}$

(B) $4\begin{bmatrix} 0 & -1 \\ 2 & 2 \end{bmatrix}$

(C) $32\begin{bmatrix} 2 & 1 \\ 2 & 0 \end{bmatrix}$

(D) $32\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q262. JEE Main 2013 (Offline)

The number of values of k, for which the system of equations :

$$(k+1)x + 8y = 4k$$
$$kx + (k+3)y = 3k - 1$$

has no solution, is

(A) infinite (B) 1 (C) 2 (D) 3

Open solution

Q263. AIEEE 2012 (Online) 26th May

If $A = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ -3 & 2 & 1 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 7 & -2 & 1 \end{bmatrix}$ then AB equals

(A) I (B) A (C) B (D) 0

Open solution

Q264. AIEEE 2012 (Online) 26th May

Statement 1: If the system of equations $x + ky + 3z = 0,\ 3x + ky - 2z = 0,\ 2x + 3y - 4z = 0$ has a nontrivial solution, then the value of k is $\frac{31}{2}$.
Statement 2: A system of three homogeneous equations in three variables has a non trivial solution if the determinant of the coefficient matrix is zero.

(A) Statement 1 is false, Statement 2 is true.

(B) Statement 1 is true, Statement 2 is true, Statement 2 is a correct explanation for Statement 1.

(C) Statement 1 is true, Statement 2 is true, Statement 2 is not a correct explanation for Statement 1.

(D) Statement 1 is true, Statement 2 is false.

Open solution

Q265. AIEEE 2012 (Online) 19th May

If A^T denotes the transpose of the matrix $A = \begin{bmatrix} 0 & 0 & a \\ 0 & b & c \\ d & e & f \end{bmatrix}$, where a, b, c, d, e and f are integers such that $abd \neq 0$, then the number of such matrices for which $A^{-1} = A^T$ is

(A) $2(3!)$ (B) $3(2!)$ (C) 2^3 (D) 3^2

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q266. AIEEE 2012 (Online) 12th May

Let A and B be real matrices of the form $\begin{bmatrix}\alpha & 0\\ 0 & \beta\end{bmatrix}$ and $\begin{bmatrix}0 & \gamma\\ \delta & 0\end{bmatrix}$, respectively.

Statement 1: $AB - BA$ is always an invertible matrix.

Statement 2: $AB - BA$ is never an identity matrix.

(A) Statement 1 is true, Statement 2 is false.

(B) Statement 1 is false, Statement 2 is true.

(C) Statement 1 is true, Statement 2 is true; Statement 2 is a correct explanation of Statement 1.

(D) Statement 1 is true, Statement 2 is true; Statement 2 is not a correct explanation of Statement 1.

Open solution

Q267. AIEEE 2012 (Online) 7th May

If $A = \begin{pmatrix}\alpha - 1\\ 0\\ 0\end{pmatrix}$, $B = \begin{pmatrix}\alpha + 1\\ 0\\ 0\end{pmatrix}$ be two matrices, then AB^T is a non-zero matrix for $|\alpha|$ not equal to

(A) 2 (B) 0 (C) 1 (D) 3

Open solution

Q268. AIEEE 2012 (Online) 7th May

If the system of equations

$x + y + z = 6$

$x + 2y + 3z = 10$

$x + 2y + \lambda z = 0$

has a unique solution, then λ is not equal to

(A) 1 (B) 0 (C) 2 (D) 3

Open solution

Q269. AIEEE 2011

Let A and B be two symmetric matrices of order 3.

Statement - 1 : $A(BA)$ and $(AB)A$ are symmetric matrices.

Statement - 2 : AB is symmetric matrix if matrix multiplication of A with B is commutative.

(A) statement - 1 is true, statement - 2 is true; statement - 2 is **not** a correct explanation for statement - 1.

(B) statement - 1 is true, statement - 2 is false.

(C) statement - 1 is false, statement -2 is true

(D) statement -1 is true, statement - 2 is true; statement - 2 is a correct explanation for statement - 1.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q270. AIEEE 2011

The number of values of k for which the linear equations $4x + ky + 2z = 0, kx + 4y + z = 0$ and $2x + 2y + z = 0$ possess a non-zero solution is :

(A) 2 (B) 1 (C) zero (D) 3

Open solution

Q271. AIEEE 2010

Let A be a 2×2 matrix with non-zero entries and let $A^2 = I$,
where I is 2×2 identity matrix. Define
$Tr(A) =$ sum of diagonal elements of A and $|A| =$ determinant of matrix A.
Statement- 1: $Tr(A) = 0$.
Statement- 2: $|A| = 1$.

(A) statement - 1 is true, statement - 2 is true; statement - 2 is **not** a correct explanation for statement - 1.

(B) statement - 1 is true, statement - 2 is false.

(C) statement - 1 is false, statement -2 is true

(D) statement -1 is true, statement - 2 is true; statement - 2 is a correct explanation for statement - 1.

Open solution

Q272. AIEEE 2010

Consider the system of linear equations; $\begin{aligned} x_1 + 2x_2 + x_3 &= 3 \\ 2x_1 + 3x_2 + x_3 &= 3 \\ 3x_1 + 5x_2 + 2x_3 &= 1 \end{aligned}$

The system has :

(A) exactly 3 solutions (B) a unique solution

(C) no solution (D) infinite number of solutions

Open solution

Q273. AIEEE 2010

The number of 3×3 non-singular matrices, with four entries as 1 and all other entries as 0, is :

(A) 5 (B) 6 (C) at least 7 (D) less than 4

Open solution

Q274. AIEEE 2009

Let a, b, c be such that $b(a + c) \neq 0$ if $\begin{vmatrix} a & a+1 & a-1 \\ -b & b+1 & b-1 \\ c & c-1 & c+1 \end{vmatrix} + \begin{vmatrix} a+1 & b+1 & c-1 \\ a-1 & b-1 & c+1 \\ (-1)^{n+2}a & (-1)^{n+1}b & (-1)^n c \end{vmatrix} = 0$

then the value of n :

(A) any even integer (B) any odd integer (C) any integer (D) zero

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q275. AIEEE 2008

Let A be a 2×2 matrix with real entries. Let I be the 2×2 identity matrix. Denote by tr(A), the sum of diagonal entries of A. Assume that $A^2 = I$.

Statement-1 : If $A \neq I$ and $A \neq -I$, then det$(A) = -1$

Statement- 2 : If $A \neq I$ and $A \neq -I$, then tr $(A) \neq 0$.

(A) statement - 1 is false, statement -2 is true

(B) statement -1 is true, statement - 2 is true; statement - 2 is a correct explanation for statement - 1.

(C) statement - 1 is true, statement - 2 is true; statement - 2 is not a correct explanation for statement - 1.

(D) statement - 1 is true, statement - 2 is false.

Open solution

Q276. AIEEE 2008

Let a, b, c be any real numbers. Suppose that there are real numbers x, y, z not all zero such that $x = cy + bz$, $y = az + cx$, and $z = bx + ay$. Then $a^2 + b^2 + c^2 + 2abc$ is equal to :

(A) 2 (B) -1 (C) 0 (D) 1

Open solution

Q277. AIEEE 2007

If $D = \begin{vmatrix} 1 & 1 & 1 \\ 1 & 1+x & 1 \\ 1 & 1 & 1+y \end{vmatrix}$ for $x \neq 0, y \neq 0$, then D is :

(A) divisible by x but not y

(B) divisible by y but not x

(C) divisible by neither x nor y

(D) divisible by both x and y

Open solution

Q278. AIEEE 2006

If A and B are square matrices of size $n \times n$ such that $A^2 - B^2 = (A - B)(A + B)$, then which of the following will be always true?

(A) $A = B$

(B) $AB = BA$

(C) either of A or B is a zero matrix

(D) either of A or B is identity matrix

Open solution

Q279. AIEEE 2006

Let $A = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix}, a, b \in N$. Then

(A) there cannot exist any B such that $AB = BA$

(B) there exist more then one but finite number of B's such that $AB = BA$

(C) there exists exactly one B such that $AB = BA$

(D) there exist infinitely many B's such that $AB = BA$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q280. AIEEE 2005

If $A^2 - A + I = 0$, then the inverse of A is :

(A) $A + I$ (B) A (C) $A - I$ (D) $I - A$

Open solution

Q281. AIEEE 2005

The system of equations
$\alpha x + y + z = \alpha - 1$
$x + \alpha y + z = \alpha - 1$
$x + y + \alpha z = \alpha - 1$
has no solutions, if α is :

(A) -2 (B) either -2 or 1 (C) not -2 (D) 1

Open solution

Q282. AIEEE 2005

If $a^2 + b^2 + c^2 = -2$ and

$$f(x) = \begin{vmatrix} 1 + a^2x & (1 + b^2)x & (1 + c^2)x \\ (1 + a^2)x & 1 + b^2x & (1 + c^2)x \\ (1 + a^2)x & (1 + b^2)x & 1 + c^2x \end{vmatrix},$$

then $f(x)$ is a polynomial of degree :

(A) 1 (B) 0 (C) 3 (D) 2

Open solution

Q283. AIEEE 2005

If $a_1, a_2, a_3, \ldots\ldots\ldots, a_n, \ldots\ldots$ are in G.P., then the determinant $\Delta = \begin{vmatrix} \log a_n & \log a_{n+1} & \log a_{n+2} \\ \log a_{n+3} & \log a_{n+4} & \log a_{n+5} \\ \log a_{n+6} & \log a_{n+7} & \log a_{n+8} \end{vmatrix}$

is equal to :

(A) 1 (B) 0 (C) 4 (D) 2

Open solution

Q284. AIEEE 2004

Let $A = \begin{pmatrix} 0 & 0 & -1 \\ 0 & -1 & 0 \\ -1 & 0 & 0 \end{pmatrix}$. The only correct statement about the matrix A is

(A) $A^2 = I$ (B) $A = (-1)I$, where I is a unit matrix
(C) A^{-1} does not exist (D) A is a zero matrix

Open solution

Q285. AIEEE 2004

Let $A = \begin{pmatrix} 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{pmatrix}$ and $10B = \begin{pmatrix} 4 & 2 & 2 \\ -5 & 0 & \alpha \\ 1 & -2 & 3 \end{pmatrix}$. If B is the inverse of matrix A, then α is

(A) 5 (B) -1 (C) 2 (D) -2

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q286. AIEEE 2004

If $a_1, a_2, a_3, \ldots\ldots\ldots, a_n, \ldots\ldots$ are in G.P., then the value of the determinant $\begin{vmatrix} \log a_n & \log a_{n+1} & \log a_{n+2} \\ \log a_{n+3} & \log a_{n+4} & \log a_{n+5} \\ \log a_{n+6} & \log a_{n+7} & \log a_{n+8} \end{vmatrix}$, is

(A) -2 (B) 1 (C) 2 (D) 0

Open solution

Q287. AIEEE 2003

If the system of linear equations
$x + 2ay + az = 0;\ x + 3by + bz = 0;\ x + 4cy + cz = 0;$
has a non - zero solution, then $a, b, c.$

(A) satisfy $a + 2b + 3c = 0$ (B) are in A.P

(C) are in G.P (D) are in H.P.

Open solution

Q288. AIEEE 2003

If $1, \omega, \omega^2$ are the cube roots of unity, then
$\Delta = \begin{vmatrix} 1 & \omega^n & \omega^{2n} \\ \omega^n & \omega^{2n} & 1 \\ \omega^{2n} & 1 & \omega^n \end{vmatrix}$ is equal to

(A) ω^2 (B) 0 (C) 1 (D) ω

Open solution

Q289. AIEEE 2003

If $A = \begin{bmatrix} a & b \\ b & a \end{bmatrix}$ and $A^2 = \begin{bmatrix} \alpha & \beta \\ \beta & \alpha \end{bmatrix}$, then

(A) $\alpha = 2ab,\ \beta = a^2 + b^2$ (B) $\alpha = a^2 + b^2,\ \beta = ab$

(C) $\alpha = a^2 + b^2,\ \beta = 2ab$ (D) $\alpha = a^2 + b^2,\ \beta = a^2 - b^2$

Open solution

Q290. AIEEE 2002

If $a > 0$ and discriminant of $ax^2 + 2bx + c$ is $-ve$, then
$\begin{vmatrix} a & b & ax + b \\ b & c & bx + c \\ ax + b & bx + c & 0 \end{vmatrix}$ is equal to

(A) $+ve$ (B) $\left(ac - b^2\right)\left(ax^2 + 2bx + c\right)$

(C) $-ve$ (D) 0

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Numerical 71 questions

Q291. JEE Main 2026 (Online) 6th April Morning Shift

Let $A=\begin{bmatrix}-1 & 1 & -1\\ 1 & 0 & 1\\ 0 & 0 & 1\end{bmatrix}$ satisfy

$\mathrm{A}^2+\alpha(\mathrm{adj}(\mathrm{adj}(\mathrm{A})))+\beta(\mathrm{adj}(\mathrm{A})(\mathrm{adj}(\mathrm{adj}(\mathrm{A}))))=\begin{bmatrix}2 & -2 & 2\\ -2 & 0 & -1\\ 0 & 0 & -1\end{bmatrix}$ for some $\alpha,\beta\in\mathbb{R}$.

Then $(\alpha-\beta)^2$ is equal to ____

Open solution

Q292. JEE Main 2026 (Online) 2nd April Evening Shift

Consider the matrices $A=\begin{bmatrix}2 & -2\\ 4 & -2\end{bmatrix}$ and $B=\begin{bmatrix}3 & 9\\ 1 & 3\end{bmatrix}$. If matrices P and Q are such that $PA=B$ and $AQ=B$, then the absolute value of the sum of the diagonal elements of $2(P+Q)$ is _____.

Open solution

Q293. JEE Main 2026 (Online) 28th January Evening Shift

Let $A=\begin{bmatrix}3 & -4\\ 1 & -1\end{bmatrix}$ and B be two matrices such that $A^{100}=100B+I$. Then the sum of all the elements of B^{100} is _____

Open solution

Q294. JEE Main 2026 (Online) 24th January Morning Shift

The number of 3×2 matrices A , which can be formed using the elements of the set $\{-2,-1,0,1,2\}$ such that the sum of all the diagonal elements of $\mathrm{A}^{\mathrm{T}}\mathrm{A}$ is 5 , is ____

Open solution

Q295. JEE Main 2026 (Online) 23rd January Evening Shift

Let $A=\begin{bmatrix}0 & 2 & -3\\ -2 & 0 & 1\\ 3 & -1 & 0\end{bmatrix}$ and B be a matrix such that $B(I-A)=I+A$. Then the sum of the diagonal elements of $\mathrm{B}^{\mathrm{T}}\mathrm{B}$ is equal to ____

Open solution

Q296. JEE Main 2026 (Online) 23rd January Morning Shift

Let $|\mathrm{A}|=6$, where A is a 3×3 matrix. If $\left|\mathrm{adj}\left(3\,\mathrm{adj}\left(\mathrm{A}^2\cdot\mathrm{adj}(2\,\mathrm{A})\right)\right)\right|=2^{\mathrm{m}}\cdot 3^{\mathrm{n}},\mathrm{m},\mathrm{n}\in\mathbf{N}$, then $\mathrm{m}+\mathrm{n}$ is equal to

____ .

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q297. JEE Main 2026 (Online) 22nd January Morning Shift

Let A be a 3×3 matrix such that $\mathrm{A}+\mathrm{A}^{\mathrm{T}}=\mathrm{O}$. If $\mathrm{A}\begin{bmatrix}1\\-1\\0\end{bmatrix}=\begin{bmatrix}3\\3\\2\end{bmatrix}, \mathrm{A}^2\begin{bmatrix}1\\-1\\0\end{bmatrix}=\begin{bmatrix}-3\\19\\-24\end{bmatrix}$ and $\det(\operatorname{adj}(2\operatorname{adj}(\mathrm{A}+\mathrm{I})))=(2)^{\alpha}\cdot(3)^{\beta}\cdot(11)^{\gamma}, \alpha, \beta, \gamma$ are non-negative integers, then $\alpha+\beta+\gamma$ is equal to ____

Open solution

Q298. JEE Main 2026 (Online) 21st January Morning Shift

For some $\alpha, \beta \in \mathbf{R}$, let $A=\begin{bmatrix}\alpha & 2\\1 & 2\end{bmatrix}$ and $B=\begin{bmatrix}1 & 1\\1 & \beta\end{bmatrix}$ be such that $A^2-4A+2I=B^2-3B+I=O$. Then $\left(\det\left(\operatorname{adj}\left(A^3-B^3\right)\right)\right)^2$ is equal to ____ .

Open solution

Q299. JEE Main 2025 (Online) 7th April Morning Shift

The number of singular matrices of order 2 , whose elements are from the set $\{2,3,6,9\}$, is _____.

Open solution

Q300. JEE Main 2025 (Online) 4th April Morning Shift

Let $A=\begin{bmatrix}\cos\theta & 0 & -\sin\theta\\0 & 1 & 0\\\sin\theta & 0 & \cos\theta\end{bmatrix}$. If for some $\theta \in (0,\pi), A^2=A^T$, then the sum of the diagonal elements of the matrix $(\mathrm{A}+\mathrm{I})^3+(\mathrm{A}-\mathrm{I})^3-6\,\mathrm{A}$ is equal to _____ .

Open solution

Q301. JEE Main 2025 (Online) 3rd April Evening Shift

Let I be the identity matrix of order 3×3 and for the matrix $A=\begin{bmatrix}\lambda & 2 & 3\\4 & 5 & 6\\7 & -1 & 2\end{bmatrix}, |A|=-1$. Let B be the inverse of the matrix $\operatorname{adj}\left(A \operatorname{adj}\left(A^2\right)\right)$. Then $|(\lambda \mathrm{B}+\mathrm{I})|$ is equal to_____

Open solution

Q302. JEE Main 2025 (Online) 29th January Morning Shift

Let $S=\left\{m \in \mathbf{Z}: A^{m^2}+A^m=3I-A^{-6}\right\}$, where $A=\begin{bmatrix}2 & -1\\1 & 0\end{bmatrix}$. Then $n(S)$ is equal to _____.

Open solution

Q303. JEE Main 2025 (Online) 28th January Morning Shift

Let M denote the set of all real matrices of order 3×3 and let $\mathrm{S}=\{-3,-2,-1,1,2\}$. Let
$\mathrm{S}_1=\left\{\mathrm{A}=[a_{\mathrm{ij}}] \in \mathrm{M}: \mathrm{A}=\mathrm{A}^{\mathrm{T}} \text{ and } a_{\mathrm{ij}} \in \mathrm{S}, \forall \mathrm{i}, \mathrm{j}\right\}$,
$\mathrm{S}_2=\left\{\mathrm{A}=[a_{\mathrm{ij}}] \in \mathrm{M}: \mathrm{A}=-\mathrm{A}^{\mathrm{T}} \text{ and } a_{\mathrm{ij}} \in \mathrm{S}, \forall \mathrm{i}, \mathrm{j}\right\}$,
$\mathrm{S}_3=\left\{\mathrm{A}=[a_{\mathrm{ij}}] \in \mathrm{M}: a_{11}+a_{22}+a_{33}=0 \text{ and } a_{\mathrm{ij}} \in \mathrm{S}, \forall \mathrm{i}, \mathrm{j}\right\}$.
If $n\left(S_1 \cup S_2 \cup S_3\right)=125\alpha$, then α equals _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q304. JEE Main 2025 (Online) 24th January Morning Shift

Let A be a 3×3 matrix such that $\mathrm{X}^{\mathrm{T}}\mathrm{AX} = \mathrm{O}$ for all nonzero 3×1 matrices $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$. If $\mathrm{A}\begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 4 \\ -5 \end{bmatrix}, \mathrm{A}\begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 0 \\ 4 \\ -8 \end{bmatrix}$, and $\det(\mathrm{adj}(2(\mathrm{A} + \mathrm{I}))) = 2^{\alpha}3^{\beta}5^{\gamma}, \alpha, \beta, \gamma \in N$, then $\alpha^2 + \beta^2 + \gamma^2$ is

Open solution

Q305. JEE Main 2025 (Online) 22nd January Morning Shift

Let A be a square matrix of order 3 such that $\det(A) = -2$ and $\det(3\,\mathrm{adj}(-6\,\mathrm{adj}(3A))) = 2^{m+n} \cdot 3^{mn}, m > n$. Then $4m + 2n$ is equal to _____.

Open solution

Q306. JEE Main 2024 (Online) 9th April Evening Shift

Consider the matrices : $A = \begin{bmatrix} 2 & -5 \\ 3 & m \end{bmatrix}, B = \begin{bmatrix} 20 \\ m \end{bmatrix}$ and $X = \begin{bmatrix} x \\ y \end{bmatrix}$. Let the set of all m, for which the system of equations $AX = B$ has a negative solution (i.e., $x < 0$ and $y < 0$), be the interval (a, b). Then $8\int_a^b |A| dm$ is equal to _____.

Open solution

Q307. JEE Main 2024 (Online) 9th April Morning Shift

Let A be a non-singular matrix of order 3. If $\det(3\,\mathrm{adj}(2\,\mathrm{adj}((\det A)A))) = 3^{-13} \cdot 2^{-10}$ and $\det(3\,\mathrm{adj}(2\mathrm{A})) = 2^{\mathrm{m}} \cdot 3^{\mathrm{n}}$, then $|3\,\mathrm{m} + 2\mathrm{n}|$ is equal to _____.

Open solution

Q308. JEE Main 2024 (Online) 8th April Morning Shift

Let $A = \begin{bmatrix} 2 & -1 \\ 1 & 1 \end{bmatrix}$. If the sum of the diagonal elements of A^{13} is 3^n, then n is equal to _____.

Open solution

Q309. JEE Main 2024 (Online) 6th April Evening Shift

If the system of equations
$2x + 7y + \lambda z = 3$
$3x + 2y + 5z = 4$
$x + \mu y + 32z = -1$
has infinitely many solutions, then $(\lambda - \mu)$ is equal to _____ :

Open solution

Q310. JEE Main 2024 (Online) 6th April Morning Shift

Let $\alpha\beta\gamma = 45; \alpha, \beta, \gamma \in \mathbb{R}$. If $x(\alpha, 1, 2) + y(1, \beta, 2) + z(2, 3, \gamma) = (0, 0, 0)$ for some $x, y, z \in \mathbb{R}, xyz \neq 0$, then $6\alpha + 4\beta + \gamma$ is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q311. JEE Main 2024 (Online) 4th April Evening Shift

Let A be a 2×2 symmetric matrix such that $A\begin{bmatrix}1\\1\end{bmatrix} = \begin{bmatrix}3\\7\end{bmatrix}$ and the determinant of A be 1 . If $A^{-1} = \alpha A + \beta I$, where I is an identity matrix of order 2×2, then $\alpha + \beta$ equals _____.

Open solution

Q312. JEE Main 2024 (Online) 4th April Morning Shift

Let A be a square matrix of order 2 such that $|A| = 2$ and the sum of its diagonal elements is -3 . If the points (x, y) satisfying $\mathrm{A}^2 + x\,\mathrm{A} + y\mathrm{I} = \mathrm{O}$ lie on a hyperbola, whose transverse axis is parallel to the x-axis, eccentricity is e and the length of the latus rectum is l, then $\mathrm{e}^4 + l^4$ is equal to _____.

Open solution

Q313. JEE Main 2024 (Online) 4th April Morning Shift

Let A be a 3×3 matrix of non-negative real elements such that $A\begin{bmatrix}1\\1\\1\end{bmatrix} = 3\begin{bmatrix}1\\1\\1\end{bmatrix}$. Then the maximum value of $\det(\mathrm{A})$ is _____.

Open solution

Q314. JEE Main 2024 (Online) 1st February Evening Shift

Let $A = I_2 - 2MM^T$, where M is a real matrix of order 2×1 such that the relation $M^TM = I_1$ holds. If λ is a real number such that the relation $AX = \lambda X$ holds for some non-zero real matrix X of order 2×1, then the sum of squares of all possible values of λ is equal to _____.

Open solution

Q315. JEE Main 2024 (Online) 31st January Evening Shift

Let A be a 3×3 matrix and $\det(A) = 2$. If $n = \det(\underbrace{\mathrm{adj}(\mathrm{adj}(\ldots.(\mathrm{adj}\,A)))}_{2024-\text{ times}})$, then the remainder when n is divided by 9 is equal to _____.

Open solution

Q316. JEE Main 2024 (Online) 29th January Evening Shift

Let for any three distinct consecutive terms a, b, c of an A.P, the lines $ax + by + c = 0$ be concurrent at the point P and $Q(\alpha, \beta)$ be a point such that the system of equations

$x + y + z = 6$,

$2x + 5y + \alpha z = \beta$ and

$x + 2y + 3z = 4$, has infinitely many solutions. Then $(PQ)^2$ is equal to _____.

Open solution

Q317. JEE Main 2024 (Online) 27th January Evening Shift

Let A be a 2×2 real matrix and I be the identity matrix of order 2. If the roots of the equation $|\mathrm{A} - x\mathrm{I}| = 0$ be -1 and 3, then the sum of the diagonal elements of the matrix A^2 is

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q318. JEE Main 2024 (Online) 27th January Morning Shift

Let $A=\begin{bmatrix}2 & 0 & 1\\ 1 & 1 & 0\\ 1 & 0 & 1\end{bmatrix}$, $B=[B_1,B_2,B_3]$, where B_1,B_2,B_3 are column matrices, and

$$AB_1=\begin{bmatrix}1\\0\\0\end{bmatrix}, AB_2=\begin{bmatrix}2\\3\\0\end{bmatrix}, \quad AB_3=\begin{bmatrix}3\\2\\1\end{bmatrix}$$

If $\alpha=|B|$ and β is the sum of all the diagonal elements of B, then $\alpha^3+\beta^3$ is equal to _____.

Open solution

Q319. JEE Main 2023 (Online) 12th April Morning Shift

Let $D_k=\begin{vmatrix}1 & 2k & 2k-1\\ n & n^2+n+2 & n^2\\ n & n^2+n & n^2+n+2\end{vmatrix}$. If $\sum_{k=1}^{n} D_k=96$, then n is equal to _____.

Open solution

Q320. JEE Main 2023 (Online) 11th April Morning Shift

Let $A=\begin{bmatrix}0 & 1 & 2\\ a & 0 & 3\\ 1 & c & 0\end{bmatrix}$, where $a,c\in\mathbb{R}$. If $A^3=A$ and the positive value of a belongs to the interval $(n-1,n]$, where $n\in\mathbb{N}$, then n is equal to _____.

Open solution

Q321. JEE Main 2023 (Online) 10th April Evening Shift

Let S be the set of values of λ, for which the system of equations

$6\lambda x-3y+3z=4\lambda^2$,

$2x+6\lambda y+4z=1$,

$3x+2y+3\lambda z=\lambda$ has no solution. Then $12\sum_{\lambda\in S}|\lambda|$ is equal to _____.

Open solution

Q322. JEE Main 2023 (Online) 29th January Evening Shift

Let A be a symmetric matrix such that $|A|=2$ and $\begin{bmatrix}2 & 1\\ 3 & \frac{3}{2}\end{bmatrix}A=\begin{bmatrix}1 & 2\\ \alpha & \beta\end{bmatrix}$. If the sum of the diagonal elements of A is s, then $\frac{\beta s}{\alpha^2}$ is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q323. JEE Main 2023 (Online) 25th January Morning Shift

Let A_1, A_2, A_3 be the three A.P. with the same common difference d and having their first terms as $A, A+1, A+2$, respectively. Let a, b, c be the $7^{th}, 9^{th}, 17^{th}$ terms of A_1, A_2, A_3, respectively, such that $\begin{vmatrix} a & 7 & 1 \\ 2b & 17 & 1 \\ c & 17 & 1 \end{vmatrix} + 70 = 0$.

If $a = 29$, then the sum of first 20 terms of an AP whose first term is $c - a - b$ and common difference is $\frac{d}{12}$, is equal to _____.

Open solution

Q324. JEE Main 2022 (Online) 29th July Evening Shift

Let $X = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ and $A = \begin{bmatrix} -1 & 2 & 3 \\ 0 & 1 & 6 \\ 0 & 0 & -1 \end{bmatrix}$. For $k \in N$, if $X'A^kX = 33$, then k is equal to _____.

Open solution

Q325. JEE Main 2022 (Online) 29th July Morning Shift

Let p and p + 2 be prime numbers and let

$\Delta = \begin{vmatrix} p! & (p+1)! & (p+2)! \\ (p+1)! & (p+2)! & (p+3)! \\ (p+2)! & (p+3)! & (p+4)! \end{vmatrix}$

Then the sum of the maximum values of α and β, such that p^α and $(p+2)^\beta$ divide Δ, is _____.

Open solution

Q326. JEE Main 2022 (Online) 28th July Morning Shift

Let $A = \begin{bmatrix} 1 & -1 \\ 2 & \alpha \end{bmatrix}$ and $B = \begin{bmatrix} \beta & 1 \\ 1 & 0 \end{bmatrix}, \alpha, \beta \in \mathbf{R}$. Let α_1 be the value of α which satisfies $(A+B)^2 = A^2 + \begin{bmatrix} 2 & 2 \\ 2 & 2 \end{bmatrix}$ and α_2 be the value of α which satisfies $(A+B)^2 = B^2$. Then $|\alpha_1 - \alpha_2|$ is equal to _____.

Open solution

Q327. JEE Main 2022 (Online) 27th July Morning Shift

Let S be the set containing all 3×3 matrices with entries from $\{-1, 0, 1\}$. The total number of matrices $A \in S$ such that the sum of all the diagonal elements of A^TA is 6 is _____.

Open solution

Q328. JEE Main 2022 (Online) 26th July Evening Shift

The number of matrices $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, where $a, b, c, d \in \{-1, 0, 1, 2, 3, \ldots\ldots, 10\}$, such that $A = A^{-1}$, is _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q329. JEE Main 2022 (Online) 25th July Evening Shift

Let $A = \begin{bmatrix} 1 & a & a \\ 0 & 1 & b \\ 0 & 0 & 1 \end{bmatrix}, a, b \in \mathbb{R}$. If for some

$n \in \mathbb{N}, A^n = \begin{bmatrix} 1 & 48 & 2160 \\ 0 & 1 & 96 \\ 0 & 0 & 1 \end{bmatrix}$ then $n + a + b$ is equal to _____.

Open solution

Q330. JEE Main 2022 (Online) 25th July Morning Shift

Let $A = \begin{pmatrix} 2 & -1 & -1 \\ 1 & 0 & -1 \\ 1 & -1 & 0 \end{pmatrix}$ and $B = A - I$. If $\omega = \frac{\sqrt{3}i-1}{2}$, then the number of elements in the set $\{n \in \{1, 2, \ldots, 100\} : A^n + (\omega B)^n = A + B\}$ is equal to _____.

Open solution

Q331. JEE Main 2022 (Online) 29th June Evening Shift

Let $M = \begin{bmatrix} 0 & -\alpha \\ \alpha & 0 \end{bmatrix}$, where α is a non-zero real number an $N = \sum_{k=1}^{49} M^{2k}$. If $(I - M^2)N = -2I$, then the positive integral value of α is _____.

Open solution

Q332. JEE Main 2022 (Online) 28th June Evening Shift

If the system of linear equations
$2x - 3y = \gamma + 5$,
$\alpha x + 5y = \beta + 1$, where $\alpha, \beta, \gamma \in$ R has infinitely many solutions then the value of | $9\alpha + 3\beta + 5\gamma$ | is equal to _____.

Open solution

Q333. JEE Main 2022 (Online) 28th June Evening Shift

Let $A = \begin{pmatrix} 1+i & 1 \\ -i & 0 \end{pmatrix}$ where $i = \sqrt{-1}$. Then, the number of elements in the set { n ∈ {1, 2, ..., 100} : $A^n = A$ } is _____.

Open solution

Q334. JEE Main 2022 (Online) 26th June Evening Shift

Let $X = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$, $Y = \alpha I + \beta X + \gamma X^2$ and $Z = \alpha^2 I - \alpha\beta X + (\beta^2 - \alpha\gamma)X^2$, $\alpha, \beta, \gamma \in$ R. If $Y^{-1} = \begin{bmatrix} \frac{1}{5} & \frac{-2}{5} & \frac{1}{5} \\ 0 & \frac{1}{5} & \frac{-2}{5} \\ 0 & 0 & \frac{1}{5} \end{bmatrix}$, then $(\alpha - \beta + \gamma)^2$ is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q335. JEE Main 2022 (Online) 25th June Evening Shift

Let $A = \begin{pmatrix} 2 & -2 \\ 1 & -1 \end{pmatrix}$ and $B = \begin{pmatrix} -1 & 2 \\ -1 & 2 \end{pmatrix}$. Then the number of elements in the set {(n, m) : n, m $\in$ {1, 2, ..., 10} and $nA^n + mB^m = I$} is _____.

Open solution

Q336. JEE Main 2022 (Online) 24th June Evening Shift

Let $S = \left\{ \begin{pmatrix} -1 & a \\ 0 & b \end{pmatrix} ; a, b \in \{1, 2, 3, \ldots . 100\} \right\}$ and let $T_n = \{A \in S : A^{n(n+1)} = I\}$. Then the number of elements in $\bigcap_{n=1}^{100} T_n$ is _____.

Open solution

Q337. JEE Main 2021 (Online) 31st August Evening Shift

The number of elements in the set $\left\{ A = \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} : a, b, d \in \{-1, 0, 1\} \, and \, (I - A)^3 = I - A^3 \right\}$, where I is 2×2 identity matrix, is :

Open solution

Q338. JEE Main 2021 (Online) 27th August Morning Shift

If the system of linear equations

$2x + y - z = 3$

$x - y - z = \alpha$

$3x + 3y + \beta z = 3$

has infinitely many solutions, then $\alpha + \beta - \alpha\beta$ is equal to _____.

Open solution

Q339. JEE Main 2021 (Online) 27th July Evening Shift

If $A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$ and $M = A + A^2 + A^3 + ... + A^{20}$, then the sum of all the elements of the matrix M is equal to _____.

Open solution

Q340. JEE Main 2021 (Online) 27th July Morning Shift

For real numbers α and β, consider the following system of linear equations :

$x + y - z = 2$, $x + 2y + \alpha z = 1$, $2x - y + z = \beta$. If the system has infinite solutions, then $\alpha + \beta$ is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q341. JEE Main 2021 (Online) 27th July Morning Shift

Let $f(x) = \begin{vmatrix} \sin^2 x & -2+\cos^2 x & \cos 2x \\ 2+\sin^2 x & \cos^2 x & \cos 2x \\ \sin^2 x & \cos^2 x & 1+\cos 2x \end{vmatrix}, x \in [0, \pi]$. Then the maximum value of f(x) is equal to _____.

Open solution

Q342. JEE Main 2021 (Online) 25th July Morning Shift

Let $S = \left\{ n \in N \middle| \begin{pmatrix} 0 & i \\ 1 & 0 \end{pmatrix}^n \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \forall a, b, c, d \in R \right\}$, where i = $\sqrt{-1}$. Then the number of 2-digit numbers in the set S is _____.

Open solution

Q343. JEE Main 2021 (Online) 22th July Evening Shift

Let $A = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$. Then the number of 3 $\times$ 3 matrices B with entries from the set {1, 2, 3, 4, 5} and satisfying AB = BA is _____.

Open solution

Q344. JEE Main 2021 (Online) 20th July Morning Shift

Let $A = \begin{pmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{pmatrix}$ and B = $7A^{20} - 20A^7 + 2I$, where I is an identity matrix of order 3 $\times$ 3. If B = $[b_{ij}]$, then $b_{13}$is equal to _____.

Open solution

Q345. JEE Main 2021 (Online) 20th July Morning Shift

Let a, b, c, d in arithmetic progression with common difference λ. If $\begin{vmatrix} x+a-c & x+b & x+a \\ x-1 & x+c & x+b \\ x-b+d & x+d & x+c \end{vmatrix} = 2$, then value of λ^2 is equal to _____.

Open solution

Q346. JEE Main 2021 (Online) 18th March Evening Shift

Let I be an identity matrix of order 2 $\times$ 2 and P = $\begin{bmatrix} 2 & -1 \\ 5 & -3 \end{bmatrix}$. Then the value of n$\in$N for which P^n = 5I $-$ 8P is equal to _____.

Open solution

Q347. JEE Main 2021 (Online) 17th March Evening Shift

Let $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ and $B = \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix}$ such that AB = B and a + d = 2021, then the value of ad $-$ bc is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q348. JEE Main 2021 (Online) 17th March Evening Shift

If 1, $\log_{10}(4^x - 2)$ and $\log_{10}\left(4^x + \frac{18}{5}\right)$ are in arithmetic progression for a real number x, then the value of the determinant $\begin{vmatrix} 2\left(x - \frac{1}{2}\right) & x - 1 & x^2 \\ 1 & 0 & x \\ x & 1 & 0 \end{vmatrix}$ is equal to :

Open solution

Q349. JEE Main 2021 (Online) 16th March Evening Shift

Let $A = \begin{bmatrix} a_1 \\ a_2 \end{bmatrix}$ and $B = \begin{bmatrix} b_1 \\ b_2 \end{bmatrix}$ be two 2 × 1 matrices with real entries such that A = XB, where

$X = \frac{1}{\sqrt{3}} \begin{bmatrix} 1 & -1 \\ 1 & k \end{bmatrix}$, and k∈R.

If $a_1^2 + a_2^2 = \frac{2}{3}(b_1^2 + b_2^2)$ and $(k^2 + 1)\, b_2^2 \neq -2b_1b_2$, then the value of k is _____.

Open solution

Q350. JEE Main 2021 (Online) 16th March Morning Shift

The total number of 3 × 3 matrices A having entries from the set {0, 1, 2, 3} such that the sum of all the diagonal entries of AA^T is 9, is equal to _____.

Open solution

Q351. JEE Main 2021 (Online) 26th February Evening Shift

If the matrix $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 3 & 0 & -1 \end{bmatrix}$ satisfies the equation

$A^{20} + \alpha A^{19} + \beta A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ for some real numbers α and β, then $\beta - \alpha$ is equal to _____.

Open solution

Q352. JEE Main 2021 (Online) 25th February Morning Shift

If $A = \begin{bmatrix} 0 & -\tan\left(\frac{\theta}{2}\right) \\ \tan\left(\frac{\theta}{2}\right) & 0 \end{bmatrix}$ and

$(I_2 + A)(I_2 - A)^{-1} = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$, then $13(a^2 + b^2)$ is equal to

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q353. JEE Main 2021 (Online) 25th February Morning Shift

If the system of equations

kx + y + 2z = 1

3x − y − 2z = 2

−2x −2y −4z = 3

has infinitely many solutions, then k is equal to _____.

Open solution

Q354. JEE Main 2021 (Online) 24th February Morning Shift

Let P = $\begin{bmatrix} 3 & -1 & -2 \\ 2 & 0 & \alpha \\ 3 & -5 & 0 \end{bmatrix}$, where $\alpha \in$ R. Suppose Q = [q_{ij}] is a matrix satisfying PQ = kI_3 for some non-zero k $\in$ R.

If $q_{23} = -\frac{k}{8}$ and |Q| = $\frac{k^2}{2}$, then $\alpha^2 + k^2$ is equal to _____.

Open solution

Q355. JEE Main 2021 (Online) 24th February Morning Shift

Let M be any 3 × 3 matrix with entries from the set {0, 1, 2}. The maximum number of such matrices, for which the sum of diagonal elements of M^TM is seven, is _____.

Open solution

Q356. JEE Main 2020 (Online) 6th September Evening Slot

The sum of distinct values of λ for which the system of equations

$$(\lambda - 1)x + (3\lambda + 1)y + 2\lambda z = 0$$
$$(\lambda - 1)x + (4\lambda - 2)y + (\lambda + 3)z = 0$$
$$2x + (3\lambda + 1)y + 3(\lambda - 1)z = 0$$

has non-zero solutions, is _____ .

Open solution

Q357. JEE Main 2020 (Online) 4th September Morning Slot

If the system of equations
x - 2y + 3z = 9
2x + y + z = b
x - 7y + az = 24,
has infinitely many solutions, then a - b is equal to.....

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q358. JEE Main 2020 (Online) 3rd September Evening Slot

Let S be the set of all integer solutions, (x, y, z), of the system of equations

$x - 2y + 5z = 0$

$-2x + 4y + z = 0$

$-7x + 14y + 9z = 0$

such that $15 \leq x^2 + y^2 + z^2 \leq 150$. Then, the number of elements in the set S is equal to _____ .

Open solution

Q359. JEE Main 2020 (Online) 3rd September Morning Slot

Let $A = \begin{bmatrix} x & 1 \\ 1 & 0 \end{bmatrix}$, $x \in R$ and $A^4 = [a_{ij}]$.

If $a_{11} = 109$, then a_{22} is equal to _____ .

Open solution

Q360. JEE Main 2020 (Online) 8th January Morning Slot

The number of all 3 × 3 matrices A, with enteries from the set {−1, 0, 1} such that the sum of the diagonal elements of AA^T is 3, is

Open solution

Q361. JEE Main 2020 (Online) 7th January Evening Slot

If the system of linear equations,

$x + y + z = 6$

$x + 2y + 3z = 10$

$3x + 2y + \lambda z = \mu$

has more than two solutions, then $\mu - \lambda^2$ is equal to _____.

Open solution

Out of syllabus 54 questions

These questions are on topics no longer in the current syllabus. They are kept for practice, after everything else.

Q362. JEE Main 2023 (Online) 15th April Morning Shift · MCQ (Single Correct Answer)

Let the determinant of a square matrix A of order m be $m - n$, where m and n satisfy $4m + n = 22$ and $17m + 4n = 93$.

If $\det(n\,\mathrm{adj}(\mathrm{adj}(mA))) = 3^a 5^b 6^c$ then $a + b + c$ is equal to :

(A) 96 (B) 84 (C) 109 (D) 101

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q363. JEE Main 2023 (Online) 15th April Morning Shift · MCQ (Single Correct Answer)

Open solution

Let the system of linear equations

$-x+2y-9z=7$

$-x+3y+7z=9$

$-2x+y+5z=8$

$-3x+y+13z=\lambda$

has a unique solution $x=\alpha, y=\beta, z=\gamma$. Then the distance of the point

(α,β,γ) from the plane $2x-2y+z=\lambda$ is :

(A) 11 (B) 7 (C) 13 (D) 9

Q364. JEE Main 2023 (Online) 13th April Evening Shift · MCQ (Single Correct Answer)

Open solution

Let for $A=\begin{bmatrix}1 & 2 & 3\\ \alpha & 3 & 1\\ 1 & 1 & 2\end{bmatrix}$, $|A|=2$. If $|2\,\mathrm{adj}\,(2\,\mathrm{adj}\,(2\mathrm{A}))|=32^{\mathrm{n}}$, then $3n+\alpha$ is equal to

(A) 11 (B) 9 (C) 12 (D) 10

Q365. JEE Main 2023 (Online) 10th April Evening Shift · MCQ (Single Correct Answer)

Open solution

If $\mathrm{A}=\frac{1}{5!6!7!}\begin{bmatrix}5! & 6! & 7!\\ 6! & 7! & 8!\\ 7! & 8! & 9!\end{bmatrix}$, then $|\mathrm{adj}(\mathrm{adj}(2\,\mathrm{A}))|$ is equal to :

(A) 2^{12} (B) 2^{20} (C) 2^{8} (D) 2^{16}

Q366. JEE Main 2023 (Online) 10th April Morning Shift · MCQ (Single Correct Answer)

Open solution

If A is a 3×3 matrix and $|A|=2$, then $|3\,adj\,(|3A|A^2)|$ is equal to :

(A) $3^{12}\,.\,6^{10}$ (B) $3^{11}\,.\,6^{10}$ (C) $3^{12}\,.\,6^{11}$ (D) $3^{10}\,.\,6^{11}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q367. JEE Main 2023 (Online) 31st January Evening Shift · MCQ (Single Correct Answer)

If a point $\mathrm{P}(\alpha, \beta, \gamma)$ satisfying

$$(\alpha \quad \beta \quad \gamma)\begin{pmatrix}2 & 10 & 8\\ 9 & 3 & 8\\ 8 & 4 & 8\end{pmatrix} = (0 \quad 0 \quad 0)$$

lies on the plane $2x + 4y + 3z = 5$, then $6\alpha + 9\beta + 7\gamma$ is equal to :

(A) $\frac{11}{5}$ (B) 11 (C) -1 (D) $\frac{5}{4}$

Open solution

Q368. JEE Main 2023 (Online) 30th January Evening Shift · MCQ (Single Correct Answer)

If P is a 3×3 real matrix such that $P^T = aP + (a-1)I$, where $a > 1$, then :

(A) $|AdjP| = 1$ (B) $|AdjP| > 1$

(C) $|AdjP| = \frac{1}{2}$ (D) P is a singular matrix

Open solution

Q369. JEE Main 2023 (Online) 30th January Morning Shift · MCQ (Single Correct Answer)

Let $A = \begin{pmatrix}\mathrm{m} & \mathrm{n}\\ \mathrm{p} & \mathrm{q}\end{pmatrix}, \mathrm{d} = |\mathrm{A}| \neq 0$ and $|\mathrm{A} - \mathrm{d}(\mathrm{AdjA})| = 0$. Then

(A) $1 + \mathrm{d}^2 = \mathrm{m}^2 + \mathrm{q}^2$ (B) $1 + d^2 = (m+q)^2$

(C) $(1+d)^2 = m^2 + q^2$ (D) $(1+d)^2 = (m+q)^2$

Open solution

Q370. JEE Main 2023 (Online) 29th January Evening Shift · MCQ (Single Correct Answer)

The set of all values of $\mathrm{t} \in \mathbb{R}$, for which the matrix

$$\begin{bmatrix}e^t & e^{-t}(\sin t - 2\cos t) & e^{-t}(-2\sin t - \cos t)\\ e^t & e^{-t}(2\sin t + \cos t) & e^{-t}(\sin t - 2\cos t)\\ e^t & e^{-t}\cos t & e^{-t}\sin t\end{bmatrix}$$ is invertible, is :

(A) $\{k\pi, k \in \mathbb{Z}\}$ (B) $\mathbb{R}$

(C) $\left\{(2k+1)\frac{\pi}{2}, k \in \mathbb{Z}\right\}$ (D) $\left\{k\pi + \frac{\pi}{4}, k \in \mathbb{Z}\right\}$

Open solution

Q371. JEE Main 2023 (Online) 25th January Morning Shift · MCQ (Single Correct Answer)

Let $x, y, z > 1$ and $A = \begin{bmatrix}1 & \log_x y & \log_x z\\ \log_y x & 2 & \log_y z\\ \log_z x & \log_z y & 3\end{bmatrix}$. Then $|\mathrm{adj}\ (\mathrm{adj}\ \mathrm{A}^2)|$ is equal to

(A) 6^4 (B) 2^8 (C) 4^8 (D) 2^4

Open solution

Q372. JEE Main 2023 (Online) 24th January Evening Shift · MCQ (Single Correct Answer)

Let A be a 3×3 matrix such that $|\mathrm{adj}(\mathrm{adj}(\mathrm{adj}\ \mathrm{A}))| = 12^4$. Then $|\mathrm{A}^{-1}\ \mathrm{adj}\ \mathrm{A}|$ is equal to

(A) 12 (B) $2\sqrt{3}$ (C) 1 (D) $\sqrt{6}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q373. JEE Main 2023 (Online) 24th January Morning Shift · MCQ (Single Correct Answer)

Let α be a root of the equation $(a-c)x^2+(b-a)x+(c-b)=0$ where a, b, c are distinct real numbers such that the matrix $\begin{bmatrix}\alpha^2 & \alpha & 1\\ 1 & 1 & 1\\ a & b & c\end{bmatrix}$ is singular. Then, the value of $\frac{(a-c)^2}{(b-a)(c-b)}+\frac{(b-a)^2}{(a-c)(c-b)}+\frac{(c-b)^2}{(a-c)(b-a)}$ is

(A) 3 (B) 6 (C) 12 (D) 9

Open solution

Q374. JEE Main 2022 (Online) 30th July Evening Shift · MCQ (Single Correct Answer)

Let A be a matrix and $\begin{bmatrix}1 & 2\\ 0 & 1\end{bmatrix}A\begin{bmatrix}2 & 0\\ -1 & 1\end{bmatrix}=\begin{bmatrix}1 & 0\\ 0 & 1\end{bmatrix}$. If $|A|$ and $|A^2|$ are the roots of the quadratic equation $ax^2+bx+3=0$, then $a+b-ab$ is equal to

(A) -34 (B) 50 (C) -390 (D) 438

Open solution

Q375. JEE Main 2022 (Online) 28th July Morning Shift · MCQ (Single Correct Answer)

Let the matrix $A=\begin{bmatrix}0 & 1 & 0\\ 0 & 0 & 1\\ 1 & 0 & 0\end{bmatrix}$ and the matrix $B_0=A^{49}+2A^{98}$. If $B_n=Adj\,(B_{n-1})$ for all $n\geq 1$, then $\det\,(B_4)$ is equal to :

(A) 3^{28} (B) 3^{30} (C) 3^{32} (D) 3^{36}

Open solution

Q376. JEE Main 2022 (Online) 27th July Evening Shift · MCQ (Single Correct Answer)

Let $A=\begin{pmatrix}4 & -2\\ \alpha & \beta\end{pmatrix}$.

If $\mathrm{A}^2+\gamma\mathrm{A}+18\mathrm{I}=\mathrm{O}$, then $\det(\mathrm{A})$ is equal to _____.

(A) -18 (B) 18 (C) -50 (D) 50

Open solution

Q377. JEE Main 2022 (Online) 26th July Morning Shift · MCQ (Single Correct Answer)

Let A be a 2×2 matrix with det (A) = -1 and det ((A + I) (Adj (A) + I)) = 4. Then the sum of the diagonal elements of A can be :

(A) -1 (B) 2 (C) 1 (D) $-\sqrt{2}$

Open solution

Q378. JEE Main 2022 (Online) 30th June Morning Shift · MCQ (Single Correct Answer)

Let A and B be two square matrices of order 2. If $det\,(A)=2$, $det\,(B)=3$ and $\det\left((\det\,(5(\det\,A)B))A^2\right)=2^a3^b5^c$ for some a, b, c, $\in$ N, then a + b + c is equal to :

(A) 10 (B) 12 (C) 13 (D) 14

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q379. JEE Main 2022 (Online) 29th June Evening Shift · MCQ (Single Correct Answer)

Let $A=\begin{pmatrix}2 & -1\\ 0 & 2\end{pmatrix}$. If $B=I-{}^5C_1(adj\,A)+{}^5C_2(adj\,A)^2-\ldots..-{}^5C_5(adj\,A)^5$, then the sum of all elements of the matrix B is

(A) -5 (B) -6 (C) -7 (D) -8

Open solution

Q380. JEE Main 2022 (Online) 28th June Morning Shift · MCQ (Single Correct Answer)

Let A be a matrix of order 3 × 3 and det (A) = 2. Then det (det (A) adj (5 adj (A^3))) is equal to _____.

(A) 512×10^6 (B) 256×10^6 (C) 1024×10^6 (D) 256×10^{11}

Open solution

Q381. JEE Main 2022 (Online) 27th June Evening Shift · MCQ (Single Correct Answer)

Let $f(x)=\begin{vmatrix}a & -1 & 0\\ ax & a & -1\\ ax^2 & ax & a\end{vmatrix}$, $a\in R$. Then the sum of the squares of all the values of a, for which $2f'(10)-f'(5)+100=0$, is

(A) 117 (B) 106 (C) 125 (D) 136

Open solution

Q382. JEE Main 2022 (Online) 27th June Evening Shift · MCQ (Single Correct Answer)

Let for some real numbers α and β, $a=\alpha-i\beta$. If the system of equations $4ix+(1+i)y=0$ and $8\left(\cos\frac{2\pi}{3}+i\sin\frac{2\pi}{3}\right)x+\bar{a}y=0$ has more than one solution, then $\frac{\alpha}{\beta}$ is equal to

(A) $-2+\sqrt{3}$ (B) $2-\sqrt{3}$ (C) $2+\sqrt{3}$ (D) $-2-\sqrt{3}$

Open solution

Q383. JEE Main 2022 (Online) 27th June Evening Shift · MCQ (Single Correct Answer)

Let A and B be two 3 × 3 matrices such that $AB=I$ and $|A|=\frac{1}{8}$. Then $|adj\,(B\,adj(2A))|$ is equal to

(A) 16 (B) 32 (C) 64 (D) 128

Open solution

Q384. JEE Main 2022 (Online) 26th June Morning Shift · MCQ (Single Correct Answer)

Let A be a 3 × 3 invertible matrix. If |adj (24A)| = |adj (3 adj (2A))|, then $|A|^2$ is equal to :

(A) 6^6 (B) 2^{12} (C) 2^6 (D) 1

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q385. JEE Main 2022 (Online) 24th June Morning Shift · MCQ (Single Correct Answer)

Let S = $\{\sqrt{n} : 1 \le n \le 50$ and n is odd$\}$.

Let a $\in$ S and $A = \begin{bmatrix} 1 & 0 & a \\ -1 & 1 & 0 \\ -a & 0 & 1 \end{bmatrix}$.

If $\sum_{a \in S} \det(adj\, A) = 100\lambda$, then λ is equal to :

(A) 218 (B) 221 (C) 663 (D) 1717

Open solution

Q386. JEE Main 2021 (Online) 27th July Evening Shift · MCQ (Single Correct Answer)

Let A and B be two 3 $\times$ 3 real matrices such that $(A^2 - B^2)$ is invertible matrix. If $A^5 = B^5$ and $A^3B^2 = A^2B^3$, then the value of the determinant of the matrix $A^3 + B^3$ is equal to :

(A) 2 (B) 4 (C) 1 (D) 0

Open solution

Q387. JEE Main 2021 (Online) 17th March Morning Shift · MCQ (Single Correct Answer)

If $A = \begin{pmatrix} 0 & \sin\alpha \\ \sin\alpha & 0 \end{pmatrix}$ and $\det\left(A^2 - \frac{1}{2}I\right) = 0$, then a possible value of α is :

(A) $\frac{\pi}{4}$ (B) $\frac{\pi}{6}$ (C) $\frac{\pi}{2}$ (D) $\frac{\pi}{3}$

Open solution

Q388. JEE Main 2020 (Online) 3rd September Evening Slot · MCQ (Single Correct Answer)

Let A be a 3 $\times$ 3 matrix such that

adj A = $\begin{bmatrix} 2 & -1 & 1 \\ -1 & 0 & 2 \\ 1 & -2 & -1 \end{bmatrix}$ and B = adj(adj A).

If |A| = λ and $|(B^{-1})^T| = \mu$, then the ordered pair, ($|\lambda|$, μ) is equal to :

(A) (3, 81) (B) $\left(9, \frac{1}{9}\right)$

(C) $\left(3, \frac{1}{81}\right)$ (D) $\left(9, \frac{1}{81}\right)$

Open solution

Q389. JEE Main 2020 (Online) 9th January Morning Slot · MCQ (Single Correct Answer)

If the matrices A = $\begin{bmatrix} 1 & 1 & 2 \\ 1 & 3 & 4 \\ 1 & -1 & 3 \end{bmatrix}$,

B = adjA and C = 3A, then $\frac{|adjB|}{|C|}$ is equal to :

(A) 8 (B) 2 (C) 72 (D) 16

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q390. JEE Main 2020 (Online) 7th January Evening Slot · MCQ (Single Correct Answer)

Let A = $[a_{ij}]$ and B = $[b_{ij}]$ be two 3 × 3 real matrices such that $b_{ij} = (3)^{(i+j-2)}a_{ji}$, where i, j = 1, 2, 3. If the determinant of B is 81, then the determinant of A is:

(A) 3 (B) $\frac{1}{3}$ (C) $\frac{1}{9}$ (D) $\frac{1}{81}$

Open solution

Q391. JEE Main 2019 (Online) 12th April Evening Slot · MCQ (Single Correct Answer)

If [x] denotes the greatest integer $\leq$ x, then the system of linear equations $[\sin\theta]x + [-\cos\theta]y = 0$, $[\cot\theta]x + y = 0$

(A) has a unique solution if $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$ and have infinitely many solutions if $\theta \in \left(\pi, \frac{7\pi}{6}\right)$

(B) have infinitely many solutions if $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right)$ and has a unique solution if $\theta \in \left(\pi, \frac{7\pi}{6}\right)$

(C) have infinitely many solutions if $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right) \cup \left(\pi, \frac{7\pi}{6}\right)$

(D) has a unique solution if $\theta \in \left(\frac{\pi}{2}, \frac{2\pi}{3}\right) \cup \left(\pi, \frac{7\pi}{6}\right)$

Open solution

Q392. JEE Main 2019 (Online) 11th January Evening Slot · MCQ (Single Correct Answer)

Let A and B be two invertible matrices of order 3 $\times$ 3. If $\det(ABA^T) = 8$ and $\det(AB^{-1}) = 8$, then $\det(BA^{-1}B^T)$ is equal to :

(A) $\frac{1}{4}$ (B) 16 (C) $\frac{1}{16}$ (D) 1

Open solution

Q393. JEE Main 2017 (Online) 9th April Morning Slot · MCQ (Single Correct Answer)

If x = a, y = b, z = c is a solution of the system of linear equations

x + 8y + 7z = 0

9x + 2y + 3z = 0

x + y + z = 0

such that the point (a, b, c) lies on the plane x + 2y + z = 6, then 2a + b + c equals :

(A) -1 (B) 0 (C) 1 (D) 2

Open solution

Q394. JEE Main 2015 (Online) 11th April Morning Slot · MCQ (Single Correct Answer)

If $\begin{vmatrix} x^2+x & x+1 & x-2 \\ 2x^2+3x-1 & 3x & 3x-3 \\ x^2+2x+3 & 2x-1 & 2x-1 \end{vmatrix}$ = ax − 12,

then 'a' is equal to :

(A) 12 (B) 24 (C) -12 (D) -24

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q395. JEE Main 2014 (Online) 19th April Morning Slot · MCQ (Single Correct Answer)

$$\Delta_r = \begin{vmatrix} r & 2r-1 & 3r-2 \\ \frac{n}{2} & n-1 & a \\ \frac{1}{2}n(n-1) & (n-1)^2 & \frac{1}{2}(n-1)(3n+4) \end{vmatrix},$$

then the value of $\sum_{r=1}^{n-1} \Delta_r$:

(A) depends only on a

(B) depends only on n

(C) depends both on a and n

(D) is independent of both a and n.

Open solution

Q396. JEE Main 2014 (Online) 12th April Morning Slot · MCQ (Single Correct Answer)

$$\begin{vmatrix} a^2 & b^2 & c^2 \\ (a+\lambda)^2 & (b+\lambda)^2 & (c+\lambda)^2 \\ (a-\lambda)^2 & (b-\lambda)^2 & (c-\lambda)^2 \end{vmatrix} = k\lambda \begin{vmatrix} a^2 & b^2 & c^2 \\ a & b & c \\ 1 & 1 & 1 \end{vmatrix}, \lambda \neq 0,$$

then k is equal to :

(A) $4\lambda abc$

(B) $-4\lambda abc$

(C) $4\lambda^2$

(D) $-4\lambda^2$

Open solution

Q397. JEE Main 2014 (Online) 11th April Morning Slot · MCQ (Single Correct Answer)

Let for i = 1, 2, 3, $p_i(x)$ be a polynomial of degree 2 in x, $p_i'(x)$ and $p_i''(x)$ be the first and second order derivatives of $p_i(x)$ respectively. Let,

$$A(x) = \begin{bmatrix} p_1(x) & p_1'(x) & p_1''(x) \\ p_2(x) & p_2'(x) & p_2''(x) \\ p_3(x) & p_3'(x) & p_3''(x) \end{bmatrix}$$

and $B(x) = [A(x)]^T A(x)$. Then determinant of B(x) :

(A) is a polynomial of degree 6 in x.

(B) is a polynomial of degree 3 in x.

(C) is a polynomial of degree 2 in x.

(D) does not depend on x.

Open solution

Q398. JEE Main 2013 (Online) 9th April Morning Slot · MCQ (Single Correct Answer)

If a, b, c are sides of a scalene triangle, then the value of $\begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix}$ is

(A) non - negative

(B) negative

(C) positive

(D) non-positive

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q399. JEE Main 2013 (Offline) · MCQ (Single Correct Answer)

If $P=\begin{bmatrix}1 & \alpha & 3\\ 1 & 3 & 3\\ 2 & 4 & 4\end{bmatrix}$ is the adjoint of a 3×3 matrix A and $|A|=4$, then α is equal to :

Open solution

(A) 4 (B) 11 (C) 5 (D) 0

Q400. AIEEE 2012 (Online) 19th May · MCQ (Single Correct Answer)

If a,b,c are non zero complex numbers satisfying $a^2+b^2+c^2=0$ and $\begin{vmatrix} b^2+c^2 & ab & ac\\ ab & c^2+a^2 & bc\\ ac & bc & a^2+b^2\end{vmatrix}=ka^2b^2c^2$, then k is equal to

Open solution

(A) 1 (B) 3 (C) 4 (D) 2

Q401. AIEEE 2012 (Online) 12th May · MCQ (Single Correct Answer)

If $\begin{vmatrix} -2a & a+b & a+c\\ b+a & -2b & b+c\\ c+a & b+c & -2c\end{vmatrix}=\alpha(a+b)(b+c)(c+a)\neq 0$ then α is equal to

Open solution

(A) $a+b+c$ (B) abc (C) 4 (D) 1

Q402. AIEEE 2012 · MCQ (Single Correct Answer)

Let P and Q be 3×3 matrices $P\neq Q$. If $P^3=Q^3$ and $P^2Q=Q^2P$ then determinant of $\left(P^2+Q^2\right)$ is equal to :

Open solution

(A) -2 (B) 1 (C) 0 (D) -1

Q403. AIEEE 2012 · MCQ (Single Correct Answer)

Let $A=\begin{pmatrix}1 & 0 & 0\\ 2 & 1 & 0\\ 3 & 2 & 1\end{pmatrix}$. If u_1 and u_2 are column matrices such that $Au_1=\begin{pmatrix}1\\0\\0\end{pmatrix}$ and $Au_2=\begin{pmatrix}0\\1\\0\end{pmatrix}$, then u_1+u_2 is equal to :

Open solution

(A) $\begin{pmatrix}-1\\1\\0\end{pmatrix}$ (B) $\begin{pmatrix}-1\\1\\-1\end{pmatrix}$

(C) $\begin{pmatrix}-1\\-1\\0\end{pmatrix}$ (D) $\begin{pmatrix}1\\-1\\-1\end{pmatrix}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q404. AIEEE 2009 · MCQ (Single Correct Answer)

Open solution

Let A be a 2×2 matrix

Statement - 1 : $adj\,(adj\,A) = A$

Statement - 2 : $|adj\,A| = |A|$

(A) statement - 1 is true, statement - 2 is true; statement - 2 is not a correct explanation for statement - 1.

(B) statement - 1 is true, statement - 2 is false.

(C) statement - 1 is false, statement -2 is true

(D) statement -1 is true, statement - 2 is true; statement - 2 is a correct explanation for statement - 1.

Q405. AIEEE 2008 · MCQ (Single Correct Answer)

Open solution

Let A be a square matrix all of whose entries are integers.
Then which one of the following is true?

(A) If det $A = \pm 1$, then A^{-1} exists but all its entries are not necessarily integers

(B) If det $A \neq \pm 1$, then A^{-1} exists and all its entries are non integers

(C) If det $A = \pm 1$, then A^{-1} exists but all its entries are integers

(D) If det $A = \pm 1$, then A^{-1} need not exists

Q406. AIEEE 2007 · MCQ (Single Correct Answer)

Open solution

Let $A = \begin{vmatrix} 5 & 5\alpha & \alpha \\ 0 & \alpha & 5\alpha \\ 0 & 0 & 5 \end{vmatrix}$. If $|A^2| = 25$, then $|\alpha|$ equals

(A) $1/5$ (B) 5 (C) 5^2 (D) 1

Q407. JEE Main 2023 (Online) 31st January Evening Shift · Numerical

Open solution

Let A be a $n \times n$ matrix such that $|\mathrm{A}| = 2$. If the determinant of the matrix $\mathrm{Adj}\left(2 \cdot \mathrm{Adj}\left(2\,\mathrm{A}^{-1}\right)\right)$ is 2^{84}, then n is equal to :

Q408. JEE Main 2022 (Online) 27th July Evening Shift · Numerical

Open solution

Consider a matrix $A = \begin{bmatrix} \alpha & \beta & \gamma \\ \alpha^2 & \beta^2 & \gamma^2 \\ \beta+\gamma & \gamma+\alpha & \alpha+\beta \end{bmatrix}$, where α, β, γ are three distinct natural numbers.

If $\frac{\det(\mathrm{adj}(\mathrm{adj}(\mathrm{adj}(\mathrm{adj}\,A))))}{(\alpha-\beta)^{16}(\beta-\gamma)^{16}(\gamma-\alpha)^{16}} = 2^{32} \times 3^{16}$, then the number of such 3 - tuples (α, β, γ) is _____.

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q409. JEE Main 2022 (Online) 27th June Morning Shift · Numerical

The positive value of the determinant of the matrix A, whose Adj(Adj(A)) = $\begin{pmatrix} 14 & 28 & -14 \\ -14 & 14 & 28 \\ 28 & -14 & 14 \end{pmatrix}$, is _____.

Open solution

Q410. JEE Main 2021 (Online) 26th August Evening Shift · Numerical

Let A be a 3 × 3 real matrix. If det(2Adj(2 Adj(Adj(2A)))) = 2^{41}, then the value of det(A^2) is equal to _____.

Open solution

Q411. JEE Main 2021 (Online) 25th July Morning Shift · Numerical

Let $M = \left\{A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} : a, b, c, d \in \{\pm 3, \pm 2, \pm 1, 0\}\right\}$. Define f : M → Z, as f(A) = det(A), for all A∈M, where z is set of all integers. Then the number of A∈M such that f(A) = 15 is equal to _____.

Open solution

Q412. JEE Main 2021 (Online) 20th July Evening Shift · Numerical

Let $A = \{a_{ij}\}$ be a 3 × 3 matrix,

where $a_{ij} = \begin{cases} (-1)^{j-i} & if \quad i < j, \\ 2 & if \quad i = j, \\ (-1)^{i+j} & if \quad i > j \end{cases}$

then $\det(3Adj(2A^{-1}))$ is equal to _____.

Open solution

Q413. JEE Main 2021 (Online) 17th March Morning Shift · Numerical

If $A = \begin{bmatrix} 2 & 3 \\ 0 & -1 \end{bmatrix}$, then the value of det($A^4$) + det($A^{10} - (\text{Adj}(2A))^{10}$) is equal to _____.

Open solution

Q414. JEE Main 2021 (Online) 16th March Morning Shift · Numerical

Let $P = \begin{bmatrix} -30 & 20 & 56 \\ 90 & 140 & 112 \\ 120 & 60 & 14 \end{bmatrix}$ and

$A = \begin{bmatrix} 2 & 7 & \omega^2 \\ -1 & -\omega & 1 \\ 0 & -\omega & -\omega + 1 \end{bmatrix}$ where

$\omega = \frac{-1+i\sqrt{3}}{2}$, and I_3 be the identity matrix of order 3. If the determinant of the matrix $(P^{-1}AP - I_3)^2$ is $\alpha\omega^2$, then the value of α is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q415. JEE Main 2021 (Online) 25th February Morning Shift · **Numerical**

Let $A = \begin{bmatrix} x & y & z \\ y & z & x \\ z & x & y \end{bmatrix}$, where x, y and z are real numbers such that x + y + z > 0 and xyz = 2. If $A^2 = I_3$, then the value of $x^3 + y^3 + z^3$ is _____.

Open solution

ExamGOAL LEARNING APP
Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Answer Key

MCQ (Single Correct Answer)

1. **C**	2. **C**	3. **D**	4. **C**	5. **A**
6. **B**	7. **D**	8. **B**	9. **B**	10. **C**
11. **D**	12. **D**	13. **B**	14. **B**	15. **B**
16. **C**	17. **B**	18. **B**	19. **B**	20. **B**
21. **A**	22. **B**	23. **D**	24. **C**	25. **C**
26. **A**	27. **C**	28. **C**	29. **D**	30. **A**
31. **D**	32. **B**	33. **B**	34. **B**	35. **B**
36. **D**	37. **A**	38. **D**	39. **D**	40. **C**
41. **D**	42. **A**	43. **D**	44. **B**	45. **C**
46. **C**	47. **B**	48. **B**	49. **D**	50. **D**
51. **D**	52. **D**	53. **B**	54. **B**	55. **D**
56. **D**	57. **C**	58. **B**	59. **A**	60. **C**
61. **D**	62. **C**	63. **C**	64. **D**	65. **A**
66. **A**	67. **B**	68. **B**	69. **B**	70. **C**
71. **D**	72. **D**	73. **C**	74. **D**	75. **D**
76. **D**	77. **C**	78. **C**	79. **C**	80. **A**
81. **B**	82. **B**	83. **C**	84. **C**	85. **B**
86. **B**	87. **D**	88. **A**	89. **D**	90. **D**
91. **C**	92. **D**	93. **C**	94. **A**	95. **D**
96. **A**	97. **B**	98. **B**	99. **D**	100. **B**
101. **B**	102. **D**	103. **D**	104. **D**	105. **D**
106. **C**	107. **A**	108. **C**	109. **C**	110. **D**
111. **C**	112. **C**	113. **C**	114. **C**	115. **B**
116. **C**	117. **D**	118. **D**	119. **D**	120. **C**
121. **B**	122. **A**	123. **B**	124. **A**	125. **B**
126. **D**	127. **C**	128. **C**	129. **D**	130. **B**
131. **A**	132. **C**	133. **B**	134. **C**	135. **B**
136. **D**	137. **C**	138. **B**	139. **A**	140. **A**
141. **A**	142. **B**	143. **B**	144. **D**	145. **B**
146. **A**	147. **A**	148. **D**	149. **C**	150. **B**

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

151. **A**	152. **A**	153. **D**	154. **B**	155. **B**
156. **A**	157. **C**	158. **A**	159. **D**	160. **C**
161. **B**	162. **A**	163. **A**	164. **C**	165. **C**
166. **C**	167. **D**	168. **B**	169. **A**	170. **B**
171. **A**	172. **B**	173. **D**	174. **B**	175. **B**
176. **C**	177. **C**	178. **D**	179. **B**	180. **B**
181. **C**	182. **B**	183. **D**	184. **C**	185. **A**
186. **C**	187. **B**	188. **C**	189. **B**	190. **D**
191. **A**	192. **B**	193. **D**	194. **D**	195. **B**
196. **B**	197. **B**	198. **A**	199. **C**	200. **B**
201. **C**	202. **C**	203. **A**	204. **B**	205. **D**
206. **C**	207. **A**	208. **C**	209. **D**	210. **D**
211. **A**	212. **A**	213. **A**	214. **D**	215. **A**
216. **A**	217. **C**	218. **C**	219. **A**	220. **C**
221. **C**	222. **C**	223. **A**	224. **C**	225. **D**
226. **A**	227. **B**	228. **D**	229. **D**	230. **B**
231. **B**	232. **D**	233. **C**	234. **A**	235. **D**
236. **C**	237. **D**	238. **D**	239. **C**	240. **D**
241. **D**	242. **A**	243. *Scan QR*	244. **C**	245. **B**
246. **A**	247. **B**	248. **A**	249. **C**	250. **A**
251. **B**	252. **A**	253. **A**	254. **D**	255. **D**
256. **B**	257. **B**	258. **A**	259. **A**	260. **D**
261. **A**	262. **B**	263. **A**	264. **A**	265. **C**
266. **B**	267. **C**	268. **D**	269. **A**	270. **A**
271. **B**	272. **C**	273. **C**	274. **B**	275. **D**
276. **D**	277. **D**	278. **B**	279. **D**	280. **D**
281. **A**	282. **D**	283. **B**	284. **A**	285. **A**
286. **D**	287. **D**	288. **B**	289. **C**	290. **C**

Numerical

291. **4**	292. **34**	293. **0**	294. **312**	295. **3**
296. **62**	297. **18**	298. **225**	299. **36**	300. **6**
301. **38 or -38**	302. **2**	303. **1613**	304. **44**	305. **34**
306. **450**	307. **14**	308. **7**	309. **38**	310. **55**

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

311. **5**	312. **25**	313. **27**	314. **2**	315. **7**
316. **113**	317. **10**	318. **28**	319. **6**	320. **2**
321. **24**	322. **5**	323. **495**	324. **10**	325. **4**
326. **2**	327. **5376**	328. **50**	329. **24**	330. **17**
331. **1**	332. **58**	333. **25**	334. **100**	335. **1**
336. **100**	337. **8**	338. **5**	339. **2020**	340. **5**
341. **6**	342. **11**	343. **3125**	344. **910**	345. **1**
346. **6**	347. **2020**	348. **2**	349. **1**	350. **766**
351. **4**	352. **13**	353. **21**	354. **17**	355. **540**
356. **3**	357. **5**	358. **8**	359. **10**	360. **672**
361. **13**				

Out of syllabus

362. **A**	363. **B**	364. **A**	365. **D**	366. **B**
367. **B**	368. **A**	369. **D**	370. **B**	371. **B**
372. **B**	373. **A**	374. **D**	375. **C**	376. **B**
377. **B**	378. **B**	379. **C**	380. **A**	381. **C**
382. **B**	383. **C**	384. **C**	385. **B**	386. **D**
387. **A**	388. **C**	389. **A**	390. **C**	391. **B**
392. **C**	393. **C**	394. **B**	395. **D**	396. **C**
397. **D**	398. **B**	399. **B**	400. **C**	401. **C**
402. **C**	403. **D**	404. **D**	405. **C**	406. **A**
407. **5**	408. **42**	409. **14**	410. **4**	411. **16**
412. **108**	413. **16**	414. **36**	415. **7**	

For numerical answers given as a range, any value within the range is correct. Full solutions: scan the QR code next to each question, or visit questions.examside.com.

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

ExamGOAL
PYQ PDF

Your Exam. Our Responsibility.

From first chapter to exam day.

Practise with the questions your exam has already asked, then test yourself under real exam conditions.

PAST YEAR QUESTIONS

Every past paper, chapter-wise.

Previous year questions sorted the way you study, each with a full solution.

- Chapter-wise and year-wise
- Weightage and topic tags on every chapter
- Your progress on each one, at a glance

MOCK TESTS

Mock tests with a real exam feel.

Practise under the same pressure as the real paper, on your phone or on the big screen.

- Timed sections, just like exam day
- Mark questions for review and come back

ANALYSIS

Know your score. Know your rank.

A full report after every test, so you know exactly where you stand.

- Your percentile among everyone who took it
- Predicted real-exam percentile and All India Rank
- Subject breakdown and time analysis

PRACTICE

Solve. Check. Understand.

Learn from every question, not just the ones you get right.

- Instant answers as you practise
- Step-by-step explanations
- See how everyone else answered

NCERT Exemplar
With full solutions

Exam syllabus
Subject and chapter-wise

Bookmarks
Organised in groups

Your notes
On any question

ExamGOAL
LEARNING APP

ExamGOAL: Exam Preparation App

One app for every exam. Free trial for new users, on Android and the web.

GET IT ON Google Play

Scan to get the app

www.examgoal.com
Practice and mock tests

questions.examside.com
Free solutions

pdf.questions.examgoal.com
Free PYQ PDFs