ExamGOAL
PYQ PDF

Free · With answer key

JEE MAIN · MATHEMATICS · CHAPTER-WISE PYQS

Inverse Trigonometric Functions

JEE Main Mathematics · Previous Year Questions with Answer Key

106	100	2002 – 2026	English
QUESTIONS	PAPERS COVERED	YEARS	LANGUAGE

INSIDE THIS PDF

MCQ (Single Correct Answer)	89 questions
Numerical	16 questions
MCQ (More than One Correct Answer)	1 question
Answer key	at the end

Practise online

Every question with its full solution, free on **questions.examside.com**

1. **Solve**
   Print it, or work through it on screen.
2. **Check**
   The answer key is on the last pages.
3. **Understand**
   Scan the QR code beside any question for its full solution.

MCQ (Single Correct Answer) 89 questions

Q1. JEE Main 2026 (Online) 8th April Evening Shift

Let $\alpha = 3\sin^{-1}\left(\frac{6}{11}\right)$ and $\beta = 3\cos^{-1}\left(\frac{4}{9}\right)$, where inverse trigonometric functions take only the principal values.

Given below are two statements :

Statement I : $\cos(\alpha+\beta) > 0$.

Statement II : $\cos(\alpha) < 0$.

In the light of the above statements, choose the correct answer from the options given below :

(A) Both Statement I and Statement II are true

(B) Both Statement I and Statement II are false

(C) Statement I is true but Statement II is false

(D) Statement I is false but Statement II is true

Open solution

Q2. JEE Main 2026 (Online) 6th April Evening Shift

If $\sin\left(\tan^{-1}(x\sqrt{2})\right) = \cot\left(\sin^{-1}\sqrt{1-x^2}\right), x \in (0,1)$, then the value of x is:

(A) $\frac{1}{2}$ (B) $\frac{1}{3}$ (C) $\frac{2}{3}$ (D) $\frac{5}{8}$

Open solution

Q3. JEE Main 2026 (Online) 6th April Morning Shift

Let $0 < \alpha < 1, \beta = \frac{1}{3\alpha}$ and $\tan^{-1}(1-\alpha) + \tan^{-1}(1-\beta) = \frac{\pi}{4}$. Then $6(\alpha+\beta)$ is equal to:

(A) 6 (B) 7 (C) 8 (D) 9

Open solution

Q4. JEE Main 2026 (Online) 28th January Evening Shift

Considering the principal values of inverse trigonometric functions, the value of the expression

$\tan\left(2\sin^{-1}\left(\frac{2}{\sqrt{13}}\right) - 2\cos^{-1}\left(\frac{3}{\sqrt{10}}\right)\right)$

is equal to :

(A) $\frac{33}{56}$ (B) $-\frac{33}{56}$ (C) $-\frac{16}{63}$ (D) $\frac{16}{63}$

Open solution

Q5. JEE Main 2026 (Online) 24th January Evening Shift

If the domain of the function $f(x) = \sin^{-1}\left(\frac{1}{x^2-2x-2}\right)$, is $(-\infty,\alpha] \cup [\beta,\gamma] \cup [\delta,\infty)$, then $\alpha+\beta+\gamma+\delta$ is equal to

(A) 4 (B) 2 (C) 5 (D) 3

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q6. JEE Main 2026 (Online) 22nd January Morning Shift

The number of solutions of $\tan^{-1} 4x + \tan^{-1} 6x = \frac{\pi}{6}$, where $-\frac{1}{2\sqrt{6}} < x < \frac{1}{2\sqrt{6}}$, is equal to :

(A) 2 (B) 0 (C) 3 (D) 1

Open solution

Q7. JEE Main 2026 (Online) 21st January Morning Shift

If the domain of the function $f(x) = \cos^{-1}\left(\frac{2x-5}{11-3x}\right) + \sin^{-1}\left(2x^2 - 3x + 1\right)$ is the interval $[\alpha, \beta]$, then $\alpha + 2\beta$ is equal to :

(A) 5 (B) 2 (C) 3 (D) 1

Open solution

Q8. JEE Main 2025 (Online) 8th April Evening Shift

The value of $\cot^{-1}\left(\frac{\sqrt{1+\tan^2(2)}-1}{\tan(2)}\right) - \cot^{-1}\left(\frac{\sqrt{1+\tan^2\left(\frac{1}{2}\right)}+1}{\tan\left(\frac{1}{2}\right)}\right)$ is equal to

(A) $\pi - \frac{3}{2}$ (B) $\pi + \frac{5}{2}$ (C) $\pi - \frac{5}{4}$ (D) $\pi + \frac{3}{2}$

Open solution

Q9. JEE Main 2025 (Online) 4th April Evening Shift

The sum of the infinite series $\cot^{-1}\left(\frac{7}{4}\right) + \cot^{-1}\left(\frac{19}{4}\right) + \cot^{-1}\left(\frac{39}{4}\right) + \cot^{-1}\left(\frac{67}{4}\right) + \ldots$ is :

(A) $\frac{\pi}{2} + \cot^{-1}\left(\frac{1}{2}\right)$ (B) $\frac{\pi}{2} - \cot^{-1}\left(\frac{1}{2}\right)$

(C) $\frac{\pi}{2} - \tan^{-1}\left(\frac{1}{2}\right)$ (D) $\frac{\pi}{2} + \tan^{-1}\left(\frac{1}{2}\right)$

Open solution

Q10. JEE Main 2025 (Online) 4th April Morning Shift

Considering the principal values of the inverse trigonometric functions, $\sin^{-1}\left(\frac{\sqrt{3}}{2}x + \frac{1}{2}\sqrt{1-x^2}\right), -\frac{1}{2} < x < \frac{1}{\sqrt{2}}$, is equal to

(A) $\frac{-5\pi}{6} - \sin^{-1} x$ (B) $\frac{5\pi}{6} - \sin^{-1} x$ (C) $\frac{\pi}{6} + \sin^{-1} x$ (D) $\frac{\pi}{4} + \sin^{-1} x$

Open solution

Q11. JEE Main 2025 (Online) 28th January Evening Shift

Let [x] denote the greatest integer less than or equal to x. Then the domain of $f(x) = \sec^{-1}(2[x] + 1)$ is:

(A) $(-\infty, \infty)$ (B) $(-\infty, \infty) - \{0\}$ (C) $(-\infty, -1] \cup [0, \infty)$ (D) $(-\infty, -1] \cup [1, \infty)$

Open solution

Q12. JEE Main 2025 (Online) 28th January Morning Shift

$\cos\left(\sin^{-1}\frac{3}{5} + \sin^{-1}\frac{5}{13} + \sin^{-1}\frac{33}{65}\right)$ is equal to:

(A) $\frac{33}{65}$ (B) 1 (C) $\frac{32}{65}$ (D) 0

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q13. JEE Main 2025 (Online) 24th January Evening Shift

If $\alpha > \beta > \gamma > 0$, then the expression $\cot^{-1}\left\{\beta + \frac{(1+\beta^2)}{(\alpha-\beta)}\right\} + \cot^{-1}\left\{\gamma + \frac{(1+\gamma^2)}{(\beta-\gamma)}\right\} + \cot^{-1}\left\{\alpha + \frac{(1+\alpha^2)}{(\gamma-\alpha)}\right\}$ is equal to :

(A) 3π (B) $\frac{\pi}{2} - (\alpha+\beta+\gamma)$

(C) π (D) 0

Open solution

Q14. JEE Main 2025 (Online) 23rd January Morning Shift

If $\frac{\pi}{2} \le x \le \frac{3\pi}{4}$, then $\cos^{-1}\left(\frac{12}{13}\cos x + \frac{5}{13}\sin x\right)$ is equal to

(A) $x + \tan^{-1}\frac{5}{12}$ (B) $x - \tan^{-1}\frac{4}{3}$ (C) $x + \tan^{-1}\frac{4}{5}$ (D) $x - \tan^{-1}\frac{5}{12}$

Open solution

Q15. JEE Main 2025 (Online) 22nd January Morning Shift

Using the principal values of the inverse trigonometric functions, the sum of the maximum and the minimum values of $16\left(\left(\sec^{-1}x\right)^2 + \left(\operatorname{cosec}^{-1}x\right)^2\right)$ is :

(A) $24\pi^2$ (B) $18\pi^2$ (C) $22\pi^2$ (D) $31\pi^2$

Open solution

Q16. JEE Main 2024 (Online) 4th April Evening Shift

Given that the inverse trigonometric function assumes principal values only. Let x, y be any two real numbers in $[-1, 1]$ such that $\cos^{-1}x - \sin^{-1}y = \alpha$, $\frac{-\pi}{2} \le \alpha \le \pi$. Then, the minimum value of $x^2 + y^2 + 2xy\sin\alpha$ is

(A) 0 (B) -1 (C) $\frac{1}{2}$ (D) $\frac{-1}{2}$

Open solution

Q17. JEE Main 2024 (Online) 4th April Morning Shift

If the domain of the function $\sin^{-1}\left(\frac{3x-22}{2x-19}\right) + \log_e\left(\frac{3x^2-8x+5}{x^2-3x-10}\right)$ is $(\alpha, \beta]$, then $3\alpha + 10\beta$ is equal to:

(A) 95 (B) 100 (C) 97 (D) 98

Open solution

Q18. JEE Main 2024 (Online) 31st January Evening Shift

If $a = \sin^{-1}(\sin(5))$ and $b = \cos^{-1}(\cos(5))$, then $a^2 + b^2$ is equal to

(A) 25 (B) $4\pi^2 + 25$ (C) $8\pi^2 - 40\pi + 50$ (D) $4\pi^2 - 20\pi + 50$

Open solution

Q19. JEE Main 2024 (Online) 31st January Morning Shift

For $\alpha, \beta, \gamma \neq 0$, if $\sin^{-1}\alpha + \sin^{-1}\beta + \sin^{-1}\gamma = \pi$ and $(\alpha+\beta+\gamma)(\alpha-\gamma+\beta) = 3\alpha\beta$, then γ equals

(A) $\sqrt{3}$ (B) $\frac{\sqrt{3}}{2}$ (C) $\frac{1}{\sqrt{2}}$ (D) $\frac{\sqrt{3}-1}{2\sqrt{2}}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q20. JEE Main 2024 (Online) 29th January Evening Shift

Let $x = \frac{m}{n}$ (m, n are co-prime natural numbers) be a solution of the equation $\cos\left(2\sin^{-1}x\right) = \frac{1}{9}$ and let $\alpha, \beta(\alpha > \beta)$ be the roots of the equation $mx^2 - nx - m + n = 0$. Then the point (α, β) lies on the line

Open solution

(A) $3x - 2y = -2$ (B) $3x + 2y = 2$ (C) $5x + 8y = 9$ (D) $5x - 8y = -9$

Q21. JEE Main 2024 (Online) 27th January Evening Shift

Considering only the principal values of inverse trigonometric functions, the number of positive real values of x satisfying $\tan^{-1}(x) + \tan^{-1}(2x) = \frac{\pi}{4}$ is :

Open solution

(A) more than 2 (B) 2 (C) 0 (D) 1

Q22. JEE Main 2023 (Online) 15th April Morning Shift

If the domain of the function

$f(x) = \log_e\left(4x^2 + 11x + 6\right) + \sin^{-1}(4x + 3) + \cos^{-1}\left(\frac{10x+6}{3}\right)$ is $(\alpha, \beta]$, then

$36|\alpha + \beta|$ is equal to :

Open solution

(A) 72 (B) 54 (C) 45 (D) 63

Q23. JEE Main 2023 (Online) 1st February Evening Shift

Let $S = \left\{x \in R : 0 < x < 1 \text{ and } 2\tan^{-1}\left(\frac{1-x}{1+x}\right) = \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)\right\}$.

If $\mathrm{n(S)}$ denotes the number of elements in S then :

Open solution

(A) $\mathrm{n(S)} = 0$

(B) $\mathrm{n(S)} = 2$ and only one element in S is less than $\frac{1}{2}$.

(C) $\mathrm{n(S)} = 1$ and the element in S is more than $\frac{1}{2}$.

(D) $\mathrm{n(S)} = 1$ and the element in S is less than $\frac{1}{2}$.

Q24. JEE Main 2023 (Online) 1st February Morning Shift

Let S be the set of all solutions of the equation $\cos^{-1}(2x) - 2\cos^{-1}\left(\sqrt{1-x^2}\right) = \pi, x \in \left[-\frac{1}{2}, \frac{1}{2}\right]$.

Then $\sum_{x \in S} 2\sin^{-1}\left(x^2 - 1\right)$ is equal to :

Open solution

(A) $\pi - 2\sin^{-1}\left(\frac{\sqrt{3}}{4}\right)$ (B) $\pi - \sin^{-1}\left(\frac{\sqrt{3}}{4}\right)$

(C) $\frac{-2\pi}{3}$ (D) 0

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q25. JEE Main 2023 (Online) 31st January Evening Shift

Let (a, b) $\subset (0, 2\pi)$ be the largest interval for which $\sin^{-1}(\sin\theta) - \cos^{-1}(\sin\theta) > 0, \theta \in (0, 2\pi)$, holds.

If $\alpha x^2 + \beta x + \sin^{-1}\left(x^2 - 6x + 10\right) + \cos^{-1}\left(x^2 - 6x + 10\right) = 0$ and $\alpha - \beta = b - a$, then α is equal to :

(A) $\frac{\pi}{16}$ (B) $\frac{\pi}{48}$ (C) $\frac{\pi}{8}$ (D) $\frac{\pi}{12}$

Open solution

Q26. JEE Main 2023 (Online) 31st January Morning Shift

If $\sin^{-1}\frac{\alpha}{17} + \cos^{-1}\frac{4}{5} - \tan^{-1}\frac{77}{36} = 0, 0 < \alpha < 13$, then $\sin^{-1}(\sin\alpha) + \cos^{-1}(\cos\alpha)$ is equal to :

(A) 16 (B) π (C) $16 - 5\pi$ (D) 0

Open solution

Q27. JEE Main 2023 (Online) 24th January Morning Shift

$\tan^{-1}\left(\frac{1+\sqrt{3}}{3+\sqrt{3}}\right) + \sec^{-1}\left(\sqrt{\frac{8+4\sqrt{3}}{6+3\sqrt{3}}}\right)$ is equal to :

(A) $\frac{\pi}{2}$ (B) $\frac{\pi}{3}$ (C) $\frac{\pi}{6}$ (D) $\frac{\pi}{4}$

Open solution

Q28. JEE Main 2022 (Online) 29th July Evening Shift

The domain of the function $f(x) = \sin^{-1}\left(\frac{x^2-3x+2}{x^2+2x+7}\right)$ is :

(A) $[1, \infty)$ (B) $[-1, 2]$ (C) $[-1, \infty)$ (D) $(-\infty, 2]$

Open solution

Q29. JEE Main 2022 (Online) 28th July Evening Shift

The sum of the absolute maximum and absolute minimum values of the function $f(x) = \tan^{-1}(\sin x - \cos x)$ in the interval $[0, \pi]$ is :

(A) 0 (B) $\tan^{-1}\left(\frac{1}{\sqrt{2}}\right) - \frac{\pi}{4}$

(C) $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right) - \frac{\pi}{4}$ (D) $\frac{-\pi}{12}$

Open solution

Q30. JEE Main 2022 (Online) 28th July Morning Shift

Considering only the principal values of the inverse trigonometric functions, the domain of the function $f(x) = \cos^{-1}\left(\frac{x^2-4x+2}{x^2+3}\right)$ is :

(A) $\left(-\infty, \frac{1}{4}\right]$ (B) $\left[-\frac{1}{4}, \infty\right)$

(C) $(-1/3, \infty)$ (D) $\left(-\infty, \frac{1}{3}\right]$

Open solution

Q31. JEE Main 2022 (Online) 28th July Morning Shift

Considering the principal values of the inverse trigonometric functions, the sum of all the solutions of the equation $\cos^{-1}(x) - 2\sin^{-1}(x) = \cos^{-1}(2x)$ is equal to :

(A) 0 (B) 1 (C) $\frac{1}{2}$ (D) $-\frac{1}{2}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q32. JEE Main 2022 (Online) 27th July Evening Shift

The domain of the function $f(x)=\sin^{-1}\left[2x^2-3\right]+\log_2\left(\log_{\frac{1}{2}}\left(x^2-5x+5\right)\right)$, where [t] is the greatest integer function, is :

(A) $\left(-\sqrt{\frac{5}{2}}, \frac{5-\sqrt{5}}{2}\right)$

(B) $\left(\frac{5-\sqrt{5}}{2}, \frac{5+\sqrt{5}}{2}\right)$

(C) $\left(1, \frac{5-\sqrt{5}}{2}\right)$

(D) $\left[1, \frac{5+\sqrt{5}}{2}\right)$

Open solution

Q33. JEE Main 2022 (Online) 26th July Evening Shift

If $0<x<\frac{1}{\sqrt{2}}$ and $\frac{\sin^{-1}x}{\alpha}=\frac{\cos^{-1}x}{\beta}$, then the value of $\sin\left(\frac{2\pi\alpha}{\alpha+\beta}\right)$ is :

(A) $4\sqrt{(1-x^2)}\left(1-2x^2\right)$

(B) $4x\sqrt{(1-x^2)}\left(1-2x^2\right)$

(C) $2x\sqrt{(1-x^2)}\left(1-4x^2\right)$

(D) $4\sqrt{(1-x^2)}\left(1-4x^2\right)$

Open solution

Q34. JEE Main 2022 (Online) 26th July Morning Shift

$\tan\left(2\tan^{-1}\frac{1}{5}+\sec^{-1}\frac{\sqrt{5}}{2}+2\tan^{-1}\frac{1}{8}\right)$ is equal to :

(A) 1

(B) 2

(C) $\frac{1}{4}$

(D) $\frac{5}{4}$

Open solution

Q35. JEE Main 2022 (Online) 30th June Morning Shift

Let m and M respectively be the minimum and the maximum values of $f(x)=\sin^{-1}2x+\sin 2x+\cos^{-1}2x+\cos 2x,\ x\in\left[0,\frac{\pi}{8}\right]$. Then m + M is equal to :

(A) $1+\sqrt{2}+\pi$

(B) $\left(1+\sqrt{2}\right)\pi$

(C) $\pi+\sqrt{2}$

(D) $1+\pi$

Open solution

Q36. JEE Main 2022 (Online) 30th June Morning Shift

Let $\alpha=\tan\left(\frac{5\pi}{16}\sin\left(2\cos^{-1}\left(\frac{1}{\sqrt{5}}\right)\right)\right)$ and $\beta=\cos\left(\sin^{-1}\left(\frac{4}{5}\right)+\sec^{-1}\left(\frac{5}{3}\right)\right)$ where the inverse trigonometric functions take principal values. Then, the equation whose roots are α and β is :

(A) $15x^2-8x-7=0$

(B) $5x^2-12x+7=0$

(C) $25x^2-18x-7=0$

(D) $25x^2-32x+7=0$

Open solution

Q37. JEE Main 2022 (Online) 29th June Morning Shift

The domain of the function $\cos^{-1}\left(\frac{2\sin^{-1}\left(\frac{1}{4x^2-1}\right)}{\pi}\right)$ is :

(A) $R-\left\{-\frac{1}{2},\frac{1}{2}\right\}$

(B) $(-\infty,-1]\cup[1,\infty)\cup\{0\}$

(C) $\left(-\infty,\frac{-1}{2}\right)\cup\left(\frac{1}{2},\infty\right)\cup\{0\}$

(D) $\left(-\infty,\frac{-1}{\sqrt{2}}\right]\cup\left[\frac{1}{\sqrt{2}},\infty\right)\cup\{0\}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q38. JEE Main 2022 (Online) 27th June Evening Shift

The value of $\cot\left(\sum_{n=1}^{50}\tan^{-1}\left(\frac{1}{1+n+n^2}\right)\right)$ is :

(A) $\frac{26}{25}$ (B) $\frac{25}{26}$ (C) $\frac{50}{51}$ (D) $\frac{52}{51}$

Open solution

Q39. JEE Main 2022 (Online) 27th June Morning Shift

$\sin^{-1}\left(\sin\frac{2\pi}{3}\right)+\cos^{-1}\left(\cos\frac{7\pi}{6}\right)+\tan^{-1}\left(\tan\frac{3\pi}{4}\right)$ is equal to :

(A) $\frac{11\pi}{12}$ (B) $\frac{17\pi}{12}$

(C) $\frac{31\pi}{12}$ (D) $-\frac{3\pi}{4}$

Open solution

Q40. JEE Main 2022 (Online) 26th June Evening Shift

If the inverse trigonometric functions take principal values then

$\cos^{-1}\left(\frac{3}{10}\cos\left(\tan^{-1}\left(\frac{4}{3}\right)\right)+\frac{2}{5}\sin\left(\tan^{-1}\left(\frac{4}{3}\right)\right)\right)$ is equal to :

(A) 0 (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{3}$ (D) $\frac{\pi}{6}$

Open solution

Q41. JEE Main 2022 (Online) 25th June Evening Shift

The value of $\tan^{-1}\left(\frac{\cos\left(\frac{15\pi}{4}\right)-1}{\sin\left(\frac{\pi}{4}\right)}\right)$ is equal to :

(A) $-\frac{\pi}{4}$ (B) $-\frac{\pi}{8}$ (C) $-\frac{5\pi}{12}$ (D) $-\frac{4\pi}{9}$

Open solution

Q42. JEE Main 2022 (Online) 24th June Evening Shift

Let $x*y=x^2+y^3$ and $(x*1)*1=x*(1*1)$.

Then a value of $2\sin^{-1}\left(\frac{x^4+x^2-2}{x^4+x^2+2}\right)$ is :

(A) $\frac{\pi}{4}$ (B) $\frac{\pi}{3}$ (C) $\frac{\pi}{2}$ (D) $\frac{\pi}{6}$

Open solution

Q43. JEE Main 2022 (Online) 24th June Morning Shift

The set of all values of k for which

$(\tan^{-1}x)^3+(\cot^{-1}x)^3=k\pi^3,\ x\in R$, is the interval :

(A) $\left[\frac{1}{32},\frac{7}{8}\right)$ (B) $\left(\frac{1}{24},\frac{13}{16}\right)$

(C) $\left[\frac{1}{48},\frac{13}{16}\right]$ (D) $\left[\frac{1}{32},\frac{9}{8}\right)$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q44. JEE Main 2022 (Online) 24th June Morning Shift

The domain of the function

$f(x) = \frac{\cos^{-1}\left(\frac{x^2-5x+6}{x^2-9}\right)}{\log_e(x^2-3x+2)}$ is :

(A) $(-\infty, 1) \cup (2, \infty)$

(B) $(2, \infty)$

(C) $\left[-\frac{1}{2}, 1\right) \cup (2, \infty)$

(D) $\left[-\frac{1}{2}, 1\right) \cup (2, \infty) - \left\{3, \frac{3+\sqrt{5}}{2}, \frac{3-\sqrt{5}}{2}\right\}$

Open solution

Q45. JEE Main 2021 (Online) 1st September Evening Shift

$\cos^{-1}(\cos(-5)) + \sin^{-1}(\sin(6)) - \tan^{-1}(\tan(12))$ is equal to :

(The inverse trigonometric functions take the principal values)

(A) $3\pi - 11$ (B) $4\pi - 9$ (C) $4\pi - 11$ (D) $3\pi + 1$

Open solution

Q46. JEE Main 2021 (Online) 31st August Evening Shift

The domain of the function

$f(x) = \sin^{-1}\left(\frac{3x^2+x-1}{(x-1)^2}\right) + \cos^{-1}\left(\frac{x-1}{x+1}\right)$ is :

(A) $\left[0, \frac{1}{4}\right]$

(B) $[-2, 0] \cup \left[\frac{1}{4}, \frac{1}{2}\right]$

(C) $\left[\frac{1}{4}, \frac{1}{2}\right] \cup \{0\}$

(D) $\left[0, \frac{1}{2}\right]$

Open solution

Q47. JEE Main 2021 (Online) 27th August Evening Shift

Let M and m respectively be the maximum and minimum values of the function f(x) = $\tan^{-1}$ (sin x + cos x) in $\left[0, \frac{\pi}{2}\right]$, then the value of tan(M − m) is equal to :

(A) $2 + \sqrt{3}$ (B) $2 - \sqrt{3}$ (C) $3 + 2\sqrt{2}$ (D) $3 - 2\sqrt{2}$

Open solution

Q48. JEE Main 2021 (Online) 27th August Morning Shift

If $(\sin^{-1}x)^2 - (\cos^{-1}x)^2 = a$; 0 < x < 1, a $\neq$ 0, then the value of $2x^2 - 1$ is :

(A) $\cos\left(\frac{4a}{\pi}\right)$

(B) $\sin\left(\frac{2a}{\pi}\right)$

(C) $\cos\left(\frac{2a}{\pi}\right)$

(D) $\sin\left(\frac{4a}{\pi}\right)$

Open solution

Q49. JEE Main 2021 (Online) 26th August Evening Shift

The domain of the function $\operatorname{cosec}^{-1}\left(\frac{1+x}{x}\right)$ is :

(A) $\left(-1, -\frac{1}{2}\right] \cup (0, \infty)$

(B) $\left[-\frac{1}{2}, 0\right) \cup [1, \infty)$

(C) $\left(-\frac{1}{2}, \infty\right) - \{0\}$

(D) $\left[-\frac{1}{2}, \infty\right) - \{0\}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q50. JEE Main 2021 (Online) 26th August Evening Shift

If $\sum_{r=1}^{50} \tan^{-1}\frac{1}{2r^2} = p$, then the value of tan p is :

(A) $\frac{101}{102}$ (B) $\frac{50}{51}$ (C) 100 (D) $\frac{51}{50}$

Open solution

Q51. JEE Main 2021 (Online) 22th July Evening Shift

If the domain of the function $f(x) = \frac{\cos^{-1}\sqrt{x^2-x+1}}{\sqrt{\sin^{-1}\left(\frac{2x-1}{2}\right)}}$ is the interval $(\alpha, \beta]$, then $\alpha + \beta$ is equal to :

(A) $\frac{3}{2}$ (B) 2 (C) $\frac{1}{2}$ (D) 1

Open solution

Q52. JEE Main 2021 (Online) 20th July Evening Shift

The value of $\tan\left(2\tan^{-1}\left(\frac{3}{5}\right) + \sin^{-1}\left(\frac{5}{13}\right)\right)$ is equal to :

(A) $\frac{-181}{69}$ (B) $\frac{220}{21}$ (C) $\frac{-291}{76}$ (D) $\frac{151}{63}$

Open solution

Q53. JEE Main 2021 (Online) 20th July Morning Shift

The number of real roots of the equation $\tan^{-1}\sqrt{x(x+1)} + \sin^{-1}\sqrt{x^2+x+1} = \frac{\pi}{4}$ is :

(A) 1 (B) 2 (C) 4 (D) 0

Open solution

Q54. JEE Main 2021 (Online) 17th March Evening Shift

The number of solutions of the equation

$\sin^{-1}\left[x^2 + \frac{1}{3}\right] + \cos^{-1}\left[x^2 - \frac{2}{3}\right] = x^2$, for x$\in$[$-$1, 1], and [x] denotes the greatest integer less than or equal to x, is :

(A) 0 (B) Infinite (C) 2 (D) 4

Open solution

Q55. JEE Main 2021 (Online) 17th March Morning Shift

The sum of possible values of x for

$\tan^{-1}(x+1) + \cot^{-1}\left(\frac{1}{x-1}\right) = \tan^{-1}\left(\frac{8}{31}\right)$ is :

(A) $-\frac{32}{4}$ (B) $-\frac{33}{4}$
(C) $-\frac{31}{4}$ (D) $-\frac{30}{4}$

Open solution

Q56. JEE Main 2021 (Online) 17th March Morning Shift

If $\cot^{-1}(\alpha) = \cot^{-1} 2 + \cot^{-1} 8 + \cot^{-1} 18 + \cot^{-1} 32 +$ ..... upto 100 terms, then α is :

(A) 1.02 (B) 1.03 (C) 1.01 (D) 1.00

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q57. JEE Main 2021 (Online) 16th March Evening Shift

Given that the inverse trigonometric functions take principal values only. Then, the number of real values of x which satisfy

$\sin^{-1}\left(\frac{3x}{5}\right)+\sin^{-1}\left(\frac{4x}{5}\right)=\sin^{-1}x$ is equal to :

(A) 2 (B) 0 (C) 3 (D) 1

Open solution

Q58. JEE Main 2021 (Online) 26th February Evening Shift

If 0 < a, b < 1, and $\tan^{-1}a+\tan^{-1}b=\frac{\pi}{4}$, then the value of

$(a+b)-\left(\frac{a^2+b^2}{2}\right)+\left(\frac{a^3+b^3}{3}\right)-\left(\frac{a^4+b^4}{4}\right)+\ldots\ldots$ is :

(A) $\log_e 2$ (B) e

(C) $\log_e\left(\frac{e}{2}\right)$ (D) e^2-1

Open solution

Q59. JEE Main 2021 (Online) 26th February Morning Shift

If $\frac{\sin^{-1}x}{a}=\frac{\cos^{-1}x}{b}=\frac{\tan^{-1}y}{c};\ 0<x<1,$
then the value of $\cos\left(\frac{\pi c}{a+b}\right)$ is :

(A) $\frac{1-y^2}{2y}$ (B) $\frac{1-y^2}{y\sqrt{y}}$ (C) $1-y^2$ (D) $\frac{1-y^2}{1+y^2}$

Open solution

Q60. JEE Main 2021 (Online) 25th February Evening Shift

$\text{cosec}\left[2\cot^{-1}(5)+\cos^{-1}\left(\frac{4}{5}\right)\right]$ is equal to :

(A) $\frac{75}{56}$ (B) $\frac{65}{56}$ (C) $\frac{56}{33}$ (D) $\frac{65}{33}$

Open solution

Q61. JEE Main 2021 (Online) 24th February Evening Shift

A possible value of $\tan\left(\frac{1}{4}\sin^{-1}\frac{\sqrt{63}}{8}\right)$ is :

(A) $\sqrt{7}-1$ (B) $\frac{1}{\sqrt{7}}$ (C) $2\sqrt{2}-1$ (D) $\frac{1}{2\sqrt{2}}$

Open solution

Q62. JEE Main 2020 (Online) 5th September Morning Slot

If S is the sum of the first 10 terms of the series

$\tan^{-1}\left(\frac{1}{3}\right)+\tan^{-1}\left(\frac{1}{7}\right)+\tan^{-1}\left(\frac{1}{13}\right)+\tan^{-1}\left(\frac{1}{21}\right)+\ldots.$

then tan(S) is equal to :

(A) $\frac{10}{11}$ (B) $\frac{5}{11}$ (C) $-\frac{6}{5}$ (D) $\frac{5}{6}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q63. JEE Main 2020 (Online) 3rd September Morning Slot

2π - $\left(\sin^{-1}\frac{4}{5}+\sin^{-1}\frac{5}{13}+\sin^{-1}\frac{16}{65}\right)$ is equal to :

(A) $\frac{7\pi}{4}$ (B) $\frac{5\pi}{4}$ (C) $\frac{3\pi}{2}$ (D) $\frac{\pi}{2}$

Open solution

Q64. JEE Main 2020 (Online) 2nd September Morning Slot

The domain of the function
f(x) = $\sin^{-1}\left(\frac{|x|+5}{x^2+1}\right)$ is $(-\infty, -a]\cup[a, \infty)$. Then a is equal to :

(A) $\frac{\sqrt{17}-1}{2}$ (B) $\frac{1+\sqrt{17}}{2}$ (C) $\frac{\sqrt{17}}{2}+1$ (D) $\frac{\sqrt{17}}{2}$

Open solution

Q65. JEE Main 2019 (Online) 12th April Morning Slot

The value of $\sin^{-1}\left(\frac{12}{13}\right)-\sin^{-1}\left(\frac{3}{5}\right)$ is equal to :

(A) $\pi-\sin^{-1}\left(\frac{63}{65}\right)$ (B) $\frac{\pi}{2}-\sin^{-1}\left(\frac{56}{65}\right)$
(C) $\frac{\pi}{2}-\cos^{-1}\left(\frac{9}{65}\right)$ (D) $\pi-\cos^{-1}\left(\frac{33}{65}\right)$

Open solution

Q66. JEE Main 2019 (Online) 10th April Evening Slot

If $\cos^{-1}x-\cos^{-1}\frac{y}{2}=\alpha$,where $-1\le x\le 1$, $-2\le y\le 2$, $x\le\frac{y}{2}$, then for all x, y, $4x^2-4xy\cos\alpha+y^2$ is equal to :

(A) $4\sin^2\alpha$ (B) $2\sin^2\alpha$
(C) $4\sin^2\alpha - 2x^2y^2$ (D) $4\cos^2\alpha + 2x^2y^2$

Open solution

Q67. JEE Main 2019 (Online) 8th April Morning Slot

If $\alpha=\cos^{-1}\left(\frac{3}{5}\right)$, $\beta=\tan^{-1}\left(\frac{1}{3}\right)$ where $0<\alpha,\beta<\frac{\pi}{2}$, then $\alpha-\beta$ is equal to :

(A) $\tan^{-1}\left(\frac{9}{14}\right)$ (B) $\sin^{-1}\left(\frac{9}{5\sqrt{10}}\right)$
(C) $\cos^{-1}\left(\frac{9}{5\sqrt{10}}\right)$ (D) $\tan^{-1}\left(\frac{9}{5\sqrt{10}}\right)$

Open solution

Q68. JEE Main 2019 (Online) 12th January Morning Slot

Considering only the principal values of inverse functions, the set
A = $\{x\ge 0: \tan^{-1}(2x)+\tan^{-1}(3x)=\frac{\pi}{4}\}$

(A) contains two elements (B) contains more than two elements
(C) is an empty set (D) is a singleton

Open solution

Q69. JEE Main 2019 (Online) 11th January Evening Slot

All x satisfying the inequality $(\cot^{-1}x)^2-7(\cot^{-1}x)+10>0$, lie in the interval :

(A) (cot 2, ∞) (B) $(-\infty, \cot 5)\cup(\cot 2, \infty)$
(C) (cot 5, cot 4) (D) $(-\infty, \cot 5)\cup(\cot 4, \cot 2)$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q70. JEE Main 2019 (Online) 10th January Evening Slot

The value of $\cot\left(\sum_{n=1}^{19}\cot^{-1}\left(1+\sum_{p=1}^{n}2p\right)\right)$ is :

(A) $\frac{22}{23}$ (B) $\frac{23}{22}$ (C) $\frac{21}{19}$ (D) $\frac{19}{21}$

Open solution

Q71. JEE Main 2019 (Online) 9th January Evening Slot

If x = $\sin^{-1}(\sin 10)$ and y = $\cos^{-1}(\cos 10)$, then y − x is equal to :

(A) 0 (B) 10 (C) 7π (D) π

Open solution

Q72. JEE Main 2019 (Online) 9th January Morning Slot

If $\cos^{-1}\left(\frac{2}{3x}\right)+\cos^{-1}\left(\frac{3}{4x}\right)=\frac{\pi}{2}$ (x > $\frac{3}{4}$), then x is equal to :

(A) $\frac{\sqrt{145}}{10}$ (B) $\frac{\sqrt{145}}{11}$ (C) $\frac{\sqrt{145}}{12}$ (D) $\frac{\sqrt{146}}{12}$

Open solution

Q73. JEE Main 2017 (Online) 9th April Morning Slot

A value of x satisfying the equation sin[$\cot^{-1}$ (1+ x)] = cos [$\tan^{-1}$ x], is :

(A) $-\frac{1}{2}$ (B) -1 (C) 0 (D) $\frac{1}{2}$

Open solution

Q74. JEE Main 2017 (Online) 8th April Morning Slot

The value of $\tan^{-1}\left[\frac{\sqrt{1+x^2}+\sqrt{1-x^2}}{\sqrt{1+x^2}-\sqrt{1-x^2}}\right]$, $|x|<\frac{1}{2}, x\neq 0$, is equal to :

(A) $\frac{\pi}{4}+\frac{1}{2}\cos^{-1}x^2$ (B) $\frac{\pi}{4}+\cos^{-1}x^2$

(C) $\frac{\pi}{4}-\frac{1}{2}\cos^{-1}x^2$ (D) $\frac{\pi}{4}-\cos^{-1}x^2$

Open solution

Q75. JEE Main 2015 (Online) 10th April Morning Slot

If f(x) = $2\tan^{-1}x + \sin^{-1}\left(\frac{2x}{1+x^2}\right)$, x > 1, then f (5) is equal to :

(A) $\frac{\pi}{2}$ (B) π

(C) $4\tan^{-1}(5)$ (D) $\tan^{-1}\left(\frac{65}{156}\right)$

Open solution

Q76. JEE Main 2015 (Offline)

Let $\tan^{-1}y=\tan^{-1}x+\tan^{-1}\left(\frac{2x}{1-x^2}\right)$,

where $|x|<\frac{1}{\sqrt{3}}$. Then a value of y is :

(A) $\frac{3x-x^3}{1+3x^2}$ (B) $\frac{3x+x^3}{1+3x^2}$

(C) $\frac{3x-x^3}{1-3x^2}$ (D) $\frac{3x+x^3}{1-3x^2}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q77. JEE Main 2014 (Online) 19th April Morning Slot

The principal value of $\tan^{-1}\left(\cot\frac{43\pi}{4}\right)$ is :

(A) $-\frac{3\pi}{4}$ (B) $\frac{3\pi}{4}$ (C) $-\frac{\pi}{4}$ (D) $\frac{\pi}{4}$

Open solution

Q78. JEE Main 2014 (Online) 12th April Morning Slot

Statement I : The equation $(\sin^{-1}x)^3 + (\cos^{-1}x)^3 - a\pi^3 = 0$ has a solution for all $a \geq \frac{1}{32}$.

Statement II : For any $x \in \mathbf{R}$,
$\sin^{-1}x + \cos^{-1}x = \frac{\pi}{2}$ and
$0 \leq \left(\sin^{-1}x - \frac{\pi}{4}\right)^2 \leq \frac{9\pi^2}{16}$.

(A) Both statements I and II are true.

(B) Both statements I and II are false.

(C) Statement I is true and statement II is false.

(D) Statement I is false and statement II is true.

Open solution

Q79. JEE Main 2013 (Online) 25th April Morning Slot

Let $x \in (0, 1)$. The set of all x such that $\sin^{-1}x > \cos^{-1}x$, is the interval:

(A) $\left(\frac{1}{2}, \frac{1}{\sqrt{2}}\right)$ (B) $\left(\frac{1}{\sqrt{2}}, 1\right)$ (C) $(0, 1)$ (D) $\left(0, \frac{\sqrt{3}}{2}\right)$

Open solution

Q80. JEE Main 2013 (Online) 23th April Morning Slot

$S = \tan^{-1}\left(\frac{1}{n^2+n+1}\right) + \tan^{-1}\left(\frac{1}{n^2+3n+1}\right) + \ldots + \tan^{-1}\left(\frac{1}{1+(n+19)(n+20)}\right)$, then tan S is equal to:

(A) $\frac{20}{401+20n}$ (B) $\frac{n}{n^2+20n+1}$ (C) $\frac{20}{n^2+20n+1}$ (D) $\frac{n}{401+20n}$

Open solution

Q81. JEE Main 2013 (Online) 22th April Morning Slot

The number of solutions of the equation, $\sin^{-1}x = 2\tan^{-1}x$ (in principal values) is:

(A) 1 (B) 4 (C) 2 (D) 3

Open solution

Q82. JEE Main 2013 (Online) 9th April Morning Slot

The value of x for which $\sin\left(\cot^{-1}(1+x)\right) = \cos(\tan^{-1}x)$ is

(A) $-\frac{1}{2}$ (B) 1 (C) 0 (D) $\frac{1}{2}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q83. JEE Main 2013 (Offline)

If x, y, z are in A.P. and $\tan^{-1}x, \tan^{-1}y$ and $\tan^{-1}z$ are also in A.P., then :

(A) $x = y = z$ (B) $2x = 3y = 6z$ (C) $6x = 3y = 2z$ (D) $6x = 4y = 3z$

Open solution

Q84. AIEEE 2012 (Online) 19th May

A value of $\tan^{-1}\left(\sin\left(\cos^{-1}\left(\sqrt{\frac{2}{3}}\right)\right)\right)$ is

(A) $\frac{\pi}{4}$ (B) $\frac{\pi}{2}$ (C) $\frac{\pi}{3}$ (D) $\frac{\pi}{6}$

Open solution

Q85. AIEEE 2008

The value of $\cot\left(\operatorname{cosec}^{-1}\frac{5}{3} + \tan^{-1}\frac{2}{3}\right)$ is :

(A) $\frac{6}{17}$ (B) $\frac{3}{17}$ (C) $\frac{4}{17}$ (D) $\frac{5}{17}$

Open solution

Q86. AIEEE 2007

If $\sin^{-1}\left(\frac{x}{5}\right) + \operatorname{cosec}^{-1}\left(\frac{5}{4}\right) = \frac{\pi}{2}$, then the value of x is :

(A) 4 (B) 5 (C) 1 (D) 3

Open solution

Q87. AIEEE 2005

If $\cos^{-1}x - \cos^{-1}\frac{y}{2} = \alpha$, then $4x^2 - 4xy\cos\alpha + y^2$ is equal to :

(A) $2\sin 2\alpha$ (B) 4 (C) $4\sin^2\alpha$ (D) $-4\sin^2\alpha$

Open solution

Q88. AIEEE 2003

The trigonometric equation $\sin^{-1}x = 2\sin^{-1}a$ has a solution for :

(A) $|a| \geq \frac{1}{\sqrt{2}}$ (B) $\frac{1}{2} < |a| < \frac{1}{\sqrt{2}}$

(C) all real values of a (D) $|a| \leq \frac{1}{\sqrt{2}}$

Open solution

Q89. AIEEE 2002

$\cot^{-1}\left(\sqrt{\cos\alpha}\right) - \tan^{-1}\left(\sqrt{\cos\alpha}\right) = x$, then sin x is equal to :

(A) $\tan^2\left(\frac{\alpha}{2}\right)$ (B) $\cot^2\left(\frac{\alpha}{2}\right)$

(C) $\tan\alpha$ (D) $\cot\left(\frac{\alpha}{2}\right)$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Numerical 16 questions

Q90. JEE Main 2026 (Online) 5th April Morning Shift

If $\frac{\pi}{4}+\sum_{p=1}^{11}\tan^{-1}\left(\frac{2^{p-1}}{1+2^{2p-1}}\right)=\alpha$, then $\tan\alpha$ is equal to ____ .

Open solution

Q91. JEE Main 2026 (Online) 28th January Morning Shift

If $k=\tan\left(\frac{\pi}{4}+\frac{1}{2}\cos^{-1}\left(\frac{2}{3}\right)\right)+\tan\left(\frac{1}{2}\sin^{-1}\left(\frac{2}{3}\right)\right)$, then

the number of solutions of the equation $\sin^{-1}(kx-1)=\sin^{-1}x-\cos^{-1}x$ is ____.

Open solution

Q92. JEE Main 2026 (Online) 21st January Evening Shift

Let the maximum value of $\left(\sin^{-1}x\right)^2+\left(\cos^{-1}x\right)^2$ for $x\in\left[-\frac{\sqrt{3}}{2},\frac{1}{\sqrt{2}}\right]$ be $\frac{\mathrm{m}}{\mathrm{n}}\pi^2$, where $\gcd(\mathrm{m},\mathrm{n})=1$. Then $\mathrm{m}+\mathrm{n}$ is equal to ____。

Open solution

Q93. JEE Main 2025 (Online) 2nd April Evening Shift

If $y=\cos\left(\frac{\pi}{3}+\cos^{-1}\frac{x}{2}\right)$, then $(x-y)^2+3y^2$ is equal to _____.

Open solution

Q94. JEE Main 2025 (Online) 29th January Morning Shift

Let S = $\left\{x:\cos^{-1}x=\pi+\sin^{-1}x+\sin^{-1}(2x+1)\right\}$. Then $\sum_{x\in S}(2x-1)^2$ is equal to _____.

Open solution

Q95. JEE Main 2025 (Online) 24th January Morning Shift

If for some $\alpha,\beta;\alpha\le\beta,\alpha+\beta=8$ and $\sec^2\left(\tan^{-1}\alpha\right)+\mathrm{cosec}^2\left(\cot^{-1}\beta\right)=36$, then $\alpha^2+\beta$ is _____

Open solution

Q96. JEE Main 2024 (Online) 9th April Evening Shift

Let the inverse trigonometric functions take principal values. The number of real solutions of the equation $2\sin^{-1}x+3\cos^{-1}x=\frac{2\pi}{5}$, is _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q97. JEE Main 2024 (Online) 6th April Morning Shift

For $n \in \mathrm{N}$, if $\cot^{-1} 3 + \cot^{-1} 4 + \cot^{-1} 5 + \cot^{-1} n = \frac{\pi}{4}$, then n is equal to _____.

Open solution

Q98. JEE Main 2023 (Online) 13th April Evening Shift

For $x \in (-1, 1]$, the number of solutions of the equation $\sin^{-1} x = 2\tan^{-1} x$ is equal to _____.

Open solution

Q99. JEE Main 2023 (Online) 13th April Morning Shift

If $S = \left\{x \in \mathbb{R} : \sin^{-1}\left(\frac{x+1}{\sqrt{x^2+2x+2}}\right) - \sin^{-1}\left(\frac{x}{\sqrt{x^2+1}}\right) = \frac{\pi}{4}\right\}$, then $\sum\limits_{x \in s}\left(\sin\left(\left(x^2+x+5\right)\frac{\pi}{2}\right) - \cos\left(\left(x^2+x+5\right)\pi\right)\right)$ is equal to _____.

Open solution

Q100. JEE Main 2023 (Online) 10th April Evening Shift

If the domain of the function $f(x) = \sec^{-1}\left(\frac{2x}{5x+3}\right)$ is $[\alpha, \beta) \mathrm{U} (\gamma, \delta]$, then $|3\alpha + 10(\beta + \gamma) + 21\delta|$ is equal to _____.

Open solution

Q101. JEE Main 2023 (Online) 25th January Morning Shift

If the sum of all the solutions of $\tan^{-1}\left(\frac{2x}{1-x^2}\right) + \cot^{-1}\left(\frac{1-x^2}{2x}\right) = \frac{\pi}{3}, -1 < x < 1, x \neq 0$, is $\alpha - \frac{4}{\sqrt{3}}$, then α is equal to _____.

Open solution

Q102. JEE Main 2022 (Online) 30th July Evening Shift

Let $S = \left\{\theta \in [0, 2\pi] - \left\{\frac{\pi}{2}, \frac{3\pi}{2}\right\} : \sin^{-1}(\sin\theta) + \cos^{-1}(\cos\theta) + \tan^{-1}(\tan\theta) = \frac{4\pi}{5}\right\}$.

Then $\frac{30}{\pi}\sum\limits_{\theta \in S}\theta$ is equal to _____.

Open solution

Q103. JEE Main 2022 (Online) 27th July Morning Shift

For $k \in \mathbb{R}$, let the solutions of the equation $\cos\left(\sin^{-1}\left(x\cot\left(\tan^{-1}\left(\cos\left(\sin^{-1} x\right)\right)\right)\right)\right) = k, 0 < |x| < \frac{1}{\sqrt{2}}$ be α and β, where the inverse trigonometric functions take only principal values. If the solutions of the equation $x^2 - bx - 5 = 0$ are $\frac{1}{\alpha^2} + \frac{1}{\beta^2}$ and $\frac{\alpha}{\beta}$, then $\frac{b}{k^2}$ is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q104. JEE Main 2022 (Online) 25th July Evening Shift

Let $x = \sin(2\tan^{-1}\alpha)$ and $y = \sin\left(\frac{1}{2}\tan^{-1}\frac{4}{3}\right)$. If $S = \{\alpha \in R : y^2 = 1 - x\}$, then $\sum_{\alpha\in S} 16\alpha^3$ is equal to _____.

Open solution

Q105. JEE Main 2022 (Online) 29th June Morning Shift

$50\tan\left(3\tan^{-1}\left(\frac{1}{2}\right) + 2\cos^{-1}\left(\frac{1}{\sqrt{5}}\right)\right) + 4\sqrt{2}\tan\left(\frac{1}{2}\tan^{-1}(2\sqrt{2})\right)$ is equal to _____.

Open solution

MCQ (More than One Correct Answer) 1 questions

Q106. JEE Main 2023 (Online) 30th January Evening Shift

Let $a_1 = 1, a_2, a_3, a_4, \ldots.$ be consecutive natural numbers.

Then $\tan^{-1}\left(\frac{1}{1+a_1a_2}\right) + \tan^{-1}\left(\frac{1}{1+a_2a_3}\right) + \ldots.. + \tan^{-1}\left(\frac{1}{1+a_{2021}a_{2022}}\right)$ is equal to :

(A) $\frac{\pi}{4} - \cot^{-1}(2022)$ (B) $\frac{\pi}{4} - \tan^{-1}(2022)$ (C) $\cot^{-1}(2022) - \frac{\pi}{4}$ (D) $\tan^{-1}(2022) - \frac{\pi}{4}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Answer Key

MCQ (Single Correct Answer)

1. **A**	2. **A**	3. **B**	4. **A**	5. **A**
6. **D**	7. **C**	8. **C**	9. **C**	10. **C**
11. **A**	12. **D**	13. **C**	14. **D**	15. **C**
16. **A**	17. **C**	18. **C**	19. **B**	20. **C**
21. **D**	22. **C**	23. **D**	24. **D**	25. **D**
26. **B**	27. **B**	28. **C**	29. **C**	30. **B**
31. **A**	32. **C**	33. **B**	34. **B**	35. **A**
36. **C**	37. **D**	38. **A**	39. **A**	40. **C**
41. **B**	42. **B**	43. **A**	44. **D**	45. **C**
46. **C**	47. **D**	48. **B**	49. **D**	50. **B**
51. **A**	52. **B**	53. **D**	54. **A**	55. **A**
56. **C**	57. **C**	58. **A**	59. **D**	60. **B**
61. **B**	62. **D**	63. **C**	64. **B**	65. **B**
66. **A**	67. **B**	68. **D**	69. **A**	70. **C**
71. **D**	72. **C**	73. **A**	74. **A**	75. **B**
76. **C**	77. **C**	78. **D**	79. **B**	80. **C**
81. **D**	82. **A**	83. **A**	84. **D**	85. **A**
86. **D**	87. **C**	88. **D**	89. **A**	

Numerical

90. **2048**	91. **1**	92. **65**	93. **3**	94. **5**
95. **14**	96. **0**	97. **47**	98. **2**	99. **4**
100. **24**	101. **2**	102. **68**	103. **12**	104. **130**
105. **29**				

MCQ (More than One Correct Answer)

106. **A, D**

For numerical answers given as a range, any value within the range is correct. Full solutions: scan the QR code next to each question, or visit questions.examside.com.

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

ExamGOAL
PYQ PDF

Your Exam. Our Responsibility.

From first chapter to exam day.

Practise with the questions your exam has already asked, then test yourself under real exam conditions.

PAST YEAR QUESTIONS

Every past paper, chapter-wise.

Previous year questions sorted the way you study, each with a full solution.

- Chapter-wise and year-wise
- Weightage and topic tags on every chapter
- Your progress on each one, at a glance

MOCK TESTS

Mock tests with a real exam feel.

Practise under the same pressure as the real paper, on your phone or on the big screen.

- Timed sections, just like exam day
- Mark questions for review and come back

ANALYSIS

Know your score. Know your rank.

A full report after every test, so you know exactly where you stand.

- Your percentile among everyone who took it
- Predicted real-exam percentile and All India Rank
- Subject breakdown and time analysis

PRACTICE

Solve. Check. Understand.

Learn from every question, not just the ones you get right.

- Instant answers as you practise
- Step-by-step explanations
- See how everyone else answered

NCERT Exemplar
With full solutions

Exam syllabus
Subject and chapter-wise

Bookmarks
Organised in groups

Your notes
On any question

ExamGOAL
LEARNING APP

ExamGOAL: Exam Preparation App

One app for every exam. Free trial for new users, on Android and the web.

GET IT ON Google Play

Scan to get the app

www.examgoal.com
Practice and mock tests

questions.examside.com
Free solutions

pdf.questions.examgoal.com
Free PYQ PDFs