ExamGOAL
PYQ PDF

Free · With answer key

JEE MAIN · MATHEMATICS · CHAPTER-WISE PYQS

Limits, Continuity and Differentiability

JEE Main Mathematics · Previous Year Questions with Answer Key

296	187	2002 – 2026	English
QUESTIONS	PAPERS COVERED	YEARS	LANGUAGE

INSIDE THIS PDF

MCQ (Single Correct Answer)	234 questions
Numerical	62 questions
Answer key	at the end

Practise online

Every question with its full solution, free on **questions.examside.com**

1. **Solve**
   Print it, or work through it on screen.
2. **Check**
   The answer key is on the last pages.
3. **Understand**
   Scan the QR code beside any question for its full solution.

MCQ (Single Correct Answer) 234 questions

Q1. JEE Main 2026 (Online) 8th April Evening Shift

For the function $f(x) = e^{\sin|x|} - |x|, x \in \mathbf{R}$, consider the following statements :

Statement I : f is differentiable for all $x \in \mathbf{R}$.

Statement II : f is increasing in $\left(-\pi, -\frac{\pi}{2}\right)$.

In the light of the above statements, choose the correct answer from the options given below :

(A) Both Statement I and Statement II are true

(B) Both Statement I and Statement II are false

(C) Statement I is true but Statement II is false

(D) Statement I is false but Statement II is true

Open solution

Q2. JEE Main 2026 (Online) 6th April Morning Shift

The value of $\lim\limits_{x\to 0}\left(\frac{x^2\sin^2 x}{x^2-\sin^2 x}\right)$ is :

(A) 2 (B) 3 (C) 4 (D) 6

Open solution

Q3. JEE Main 2026 (Online) 5th April Evening Shift

Let $f(x) = \lim\limits_{y\to 0}\frac{(1-\cos(xy))\tan(xy)}{y^3}$. Then the number of solutions of the equation $f(x) = \sin x, x \in \mathbf{R}$ is :

(A) 0 (B) 2 (C) 3 (D) 1

Open solution

Q4. JEE Main 2026 (Online) 5th April Evening Shift

Let $f(x)$ and $g(x)$ be twice differentiable functions satisfying $f''(x) = g''(x)$ for all $x \in \mathbf{R}, f'(1) = 2g'(1) = 4$ and $g(2) = 3f(2) = 9$. Then $f(25) - g(25)$ is equal to :

(A) 20 (B) 40 (C) -20 (D) -40

Open solution

Q5. JEE Main 2026 (Online) 5th April Morning Shift

The product of all possible values of α, for which

$\lim\limits_{x\to 0}\left(\frac{1-\cos(\alpha x)\cos((\alpha+1)x)\cos((\alpha+2)x)}{\sin^2((\alpha+1)x)}\right) = 2$, is :

(A) -2 (B) 1 (C) -1 (D) $\frac{5}{4}$

Open solution

Q6. JEE Main 2026 (Online) 2nd April Morning Shift

If $\lim\limits_{x\to 2}\frac{\sin(x^3-5x^2+ax+b)}{\left(\sqrt{x-1}-1\right)\log_e(x-1)} = m$, then $a+b+m$ is equal to :

(A) 5 (B) 6 (C) 8 (D) 10

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q7. JEE Main 2026 (Online) 28th January Evening Shift

Let $f(x)=\lim\limits_{\theta\to 0}\left(\frac{\cos\pi x-x^{\left(\frac{2}{\theta}\right)}\sin(x-1)}{1+x^{\left(\frac{2}{\theta}\right)}(x-1)}\right)$, $x\in\mathbb{R}$. Consider the following two statements :

(I) $f(x)$ is discontinuous at $x=1$.

(II) $f(x)$ is continuous at $x=-1$.

Then,

(A) Neither (I) nor (II) is True

(B) Only (II) is True

(C) Only (I) is True

(D) Both (I) and (II) are True

Open solution

Q8. JEE Main 2026 (Online) 28th January Morning Shift

The value of

$$\lim_{x\to 0}\frac{\log_e\left(\sec(ex)\cdot\sec\left(e^2x\right)\cdot\ldots\cdot\sec\left(e^{10}x\right)\right)}{e^2-e^{2\cos x}}$$

is equal to

(A) $\frac{\left(e^{10}-1\right)}{2e^2\left(e^2-1\right)}$

(B) $\frac{\left(e^{20}-1\right)}{2e^2\left(e^2-1\right)}$

(C) $\frac{\left(e^{10}-1\right)}{2\left(e^2-1\right)}$

(D) $\frac{\left(e^{20}-1\right)}{2\left(e^2-1\right)}$

Open solution

Q9. JEE Main 2026 (Online) 24th January Evening Shift

Let $y=y(x)$ be a differentiable function in the interval $(0,\infty)$ such that $y(1)=2$, and $\lim\limits_{t\to x}\left(\frac{t^2y(x)-x^2y(t)}{x-t}\right)=3$ for each $x>0$. Then $2y(2)$ is equal to :

(A) 27

(B) 18

(C) 23

(D) 12

Open solution

Q10. JEE Main 2026 (Online) 24th January Evening Shift

Let $[t]$ denote the greatest integer less than or equal to t. If the function

$$f(x)=\begin{cases}b^2\sin\left(\frac{\pi}{2}\left[\frac{\pi}{2}(\cos x+\sin x)\cos x\right]\right), & x<0\\ \frac{\sin x-\frac{1}{2}\sin 2x}{x^3} & ,x>0\\ a & ,x=0\end{cases}$$

is continuous at $x=0$, then a^2+b^2 is equal to :

(A) $\frac{1}{2}$

(B) $\frac{5}{8}$

(C) $\frac{3}{4}$

(D) $\frac{9}{16}$

Open solution

Q11. JEE Main 2026 (Online) 24th January Morning Shift

Let $\alpha,\beta\in\mathbb{R}$ be such that the function $f(x)=\begin{cases}2\alpha\left(x^2-2\right)+2\beta x & ,x<1\\ (\alpha+3)x+(\alpha-\beta) & ,x\geq 1\end{cases}$ be differentiable at all $x\in\mathbb{R}$. Then $34(\alpha+\beta)$ is equal to

(A) 48

(B) 84

(C) 36

(D) 24

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q12. JEE Main 2026 (Online) 24th January Morning Shift

If the function $f(x)=\frac{e^x(e^{\tan x-x}-1)+\log_e(\sec x+\tan x)-x}{\tan x-x}$ is continuous at $x=0$, then the value of $f(0)$ is equal to

(A) $\frac{2}{3}$ (B) $\frac{1}{2}$ (C) 2 (D) $\frac{3}{2}$

Open solution

Q13. JEE Main 2026 (Online) 23rd January Evening Shift

If $f(x)=\begin{cases}\frac{a|x|+x^2-2(\sin|x|)(\cos|x|)}{x} & ,x\neq 0\\ b & ,x=0\end{cases}$

is continuous at $x=0$, then $a+b$ is equal to :

(A) 1 (B) 2 (C) 0 (D) 4

Open solution

Q14. JEE Main 2026 (Online) 23rd January Morning Shift

Let $f(x)=\begin{cases}\frac{ax^2+2ax+3}{4x^2+4x-3} & ,x\neq -\frac{3}{2},\frac{1}{2}\\ \mathrm{b} & ,x=-\frac{3}{2},\frac{1}{2}\end{cases}$ be continuous at $x=-\frac{3}{2}$. If $f\circ f(x)=\frac{7}{5}$, then x is equal to:

(A) 1.4 (B) 2 (C) 1 (D) 0

Open solution

Q15. JEE Main 2026 (Online) 22nd January Evening Shift

If $\lim\limits_{x\to 0}\frac{\mathrm{e}^{(\mathrm{a}-1)x}+2\cos\,\mathrm{b}x+(\mathrm{c}-2)\mathrm{e}^{-x}}{x\cos x-\log_{\mathrm{e}}(1+x)}=2$, then $\mathrm{a}^2+\mathrm{b}^2+\mathrm{c}^2$ is equal to :

(A) 3 (B) 5 (C) 9 (D) 7

Open solution

Q16. JEE Main 2026 (Online) 22nd January Evening Shift

Let $[\cdot]$ denote the greatest integer function, and let $f(x)=\min\left\{\sqrt{2}x,x^2\right\}$.

Let $\mathrm{S}=\{x\in(-2,2):$ the function $\mathrm{g}(x)=|x|\left[x^2\right]$ is discontinuous at $x\}$.

Then $\sum\limits_{x\in\,\mathrm{S}}f(x)$ equals

(A) $2-\sqrt{2}$ (B) $2\sqrt{6}-3\sqrt{2}$ (C) $1-\sqrt{2}$ (D) $\sqrt{6}-2\sqrt{2}$

Open solution

Q17. JEE Main 2026 (Online) 21st January Morning Shift

Let $f:\mathbf{R}\to(0,\infty)$ be a twice differentiable function such that $f(3)=18, f'(3)=0$ and $f''(3)=4$.

Then $\lim\limits_{x\to 1}\left(\log_e\left(\frac{f(2+x)}{f(3)}\right)^{\frac{18}{(x-1)^2}}\right)$ is equal to :

(A) 9 (B) 18 (C) 1 (D) 2

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q18. JEE Main 2025 (Online) 8th April Evening Shift

Open solution

Given below are two statements:

Statement I: $\lim\limits_{x\to 0}\left(\frac{\tan^{-1}x+\log_e\sqrt{\frac{1+x}{1-x}}-2x}{x^5}\right)=\frac{2}{5}$

Statement II: $\lim\limits_{x\to 1}\left(x^{\frac{2}{1-x}}\right)=\frac{1}{e^2}$

In the light of the above statements, choose the **correct** answer from the options given below:

(A) Statement I is false but Statement II is true

(B) Both Statement I and Statement II are false

(C) Both Statement I and Statement II are true

(D) Statement I is true but Statement II is false

Q19. JEE Main 2025 (Online) 7th April Morning Shift

Open solution

$\lim\limits_{x\to 0^+}\frac{\tan\left(5(x)^{\frac{1}{3}}\right)\log_e\left(1+3x^2\right)}{\left(\tan^{-1}3\sqrt{x}\right)^2\left(e^{5(x)^{\frac{4}{3}}}-1\right)}$ is equal to

(A) $\frac{5}{3}$ (B) 1 (C) $\frac{1}{3}$ (D) $\frac{1}{15}$

Q20. JEE Main 2025 (Online) 4th April Evening Shift

Open solution

Let f be a differentiable function on $\mathbf{R}$ such that $f(2)=1, f'(2)=4$. Let $\lim\limits_{x\to 0}(f(2+x))^{3/x}=\mathrm{e}^{\alpha}$. Then the number of times the curve $y=4x^3-4x^2-4(\alpha-7)x-\alpha$ meets x-axis is :

(A) 3 (B) 1 (C) 2 (D) 0

Q21. JEE Main 2025 (Online) 4th April Morning Shift

Open solution

Let $f:\mathbb{R}\to\mathbb{R}$ be a continuous function satisfying $f(0)=1$ and $f(2x)-f(x)=x$ for all $x\in\mathbb{R}$. If $\lim\limits_{n\to\infty}\left\{f(x)-f\left(\frac{x}{2^n}\right)\right\}=G(x)$, then $\sum\limits_{r=1}^{10}G\left(r^2\right)$ is equal to

(A) 215 (B) 420 (C) 385 (D) 540

Q22. JEE Main 2025 (Online) 4th April Morning Shift

Open solution

If $\lim\limits_{x\to 1^+}\frac{(x-1)(6+\lambda\cos(x-1))+\mu\sin(1-x)}{(x-1)^3}=-1$, where $\lambda,\mu\in\mathbb{R}$, then $\lambda+\mu$ is equal to

(A) 20 (B) 19 (C) 18 (D) 17

Q23. JEE Main 2025 (Online) 3rd April Morning Shift

Open solution

Let $f(x)=\begin{cases}(1+ax)^{1/x}, & x<0\\ 1+b, & x=0\\ \frac{(x+4)^{1/2}-2}{(x+c)^{1/3}-2}, & x>0\end{cases}$ be continuous at $x=0$. Then e^abc is equal to:

(A) 64 (B) 48 (C) 36 (D) 72

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q24. JEE Main 2025 (Online) 2nd April Evening Shift

If $\lim\limits_{x\to 0}\frac{\cos(2x)+a\cos(4x)-b}{x^4}$ is finite, then $(a+b)$ is equal to:

(A) 0 (B) $\frac{3}{4}$ (C) -1 (D) $\frac{1}{2}$

Open solution

Q25. JEE Main 2025 (Online) 2nd April Morning Shift

For $\alpha,\beta,\gamma\in\mathbf{R}$, if $\lim\limits_{x\to 0}\frac{x^2\sin\alpha x+(\gamma-1)e^{x^2}}{\sin 2x-\beta x}=3$, then $\beta+\gamma-\alpha$ is equal to :

(A) -1 (B) 4 (C) 6 (D) 7

Open solution

Q26. JEE Main 2025 (Online) 29th January Evening Shift

Let the function $f(x)=\left(x^2-1\right)\left|x^2-ax+2\right|+\cos|x|$ be not differentiable at the two points $x=\alpha=2$ and $x=\beta$. Then the distance of the point (α,β) from the line $12x+5y+10=0$ is equal to :

(A) 5 (B) 2 (C) 4 (D) 3

Open solution

Q27. JEE Main 2025 (Online) 29th January Morning Shift

The value of $\lim\limits_{n\to\infty}\left(\sum\limits_{k=1}^{n}\frac{k^3+6k^2+11k+5}{(k+3)!}\right)$ is :

(A) 5/3 (B) 2 (C) 4/3 (D) 7/3

Open solution

Q28. JEE Main 2025 (Online) 24th January Evening Shift

Let $[x]$ denote the greatest integer function, and let m and n respectively be the numbers of the points, where the function $f(x)=[x]+|x-2|,-2<x<3$, is not continuous and not differentiable. Then $\mathrm{m}+\mathrm{n}$ is equal to :

(A) 6 (B) 9 (C) 8 (D) 7

Open solution

Q29. JEE Main 2025 (Online) 24th January Morning Shift

$\lim\limits_{x\to 0}\operatorname{cosec}x\left(\sqrt{2\cos^2x+3\cos x}-\sqrt{\cos^2x+\sin x+4}\right)$ is:

(A) $\frac{1}{\sqrt{15}}$ (B) $\frac{1}{2\sqrt{5}}$ (C) 0 (D) $-\frac{1}{2\sqrt{5}}$

Open solution

Q30. JEE Main 2025 (Online) 24th January Morning Shift

Let $f:\mathbb{R}-\{0\}\to\mathbb{R}$ be a function such that $f(x)-6f\left(\frac{1}{x}\right)=\frac{35}{3x}-\frac{5}{2}$. If the $\lim\limits_{x\to 0}\left(\frac{1}{\alpha x}+f(x)\right)=\beta;\alpha,\beta\in\mathbb{R}$, then $\alpha+2\beta$ is equal to

(A) 6 (B) 5 (C) 3 (D) 4

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q31. JEE Main 2025 (Online) 23rd January Evening Shift

$\lim\limits_{x\to\infty}\frac{(2x^2-3x+5)(3x-1)^{\frac{x}{2}}}{(3x^2+5x+4)\sqrt{(3x+2)^x}}$ is equal to :

(A) $\frac{2e}{3}$ (B) $\frac{2}{3\sqrt{e}}$ (C) $\frac{2e}{\sqrt{3}}$ (D) $\frac{2}{\sqrt{3e}}$

Open solution

Q32. JEE Main 2025 (Online) 23rd January Morning Shift

If the function

$$f(x)=\begin{cases}\frac{2}{x}\{\sin(k_1+1)x+\sin(k_2-1)x\}, & x<0\\ 4, & x=0\\ \frac{2}{x}\log_e\left(\frac{2+k_1x}{2+k_2x}\right), & x>0\end{cases}$$

is continuous at $x=0$, then $k_1^2+k_2^2$ is equal to :

(A) 5 (B) 10 (C) 20 (D) 8

Open solution

Q33. JEE Main 2025 (Online) 22nd January Evening Shift

If $\lim\limits_{x\to\infty}\left(\left(\frac{e}{1-e}\right)\left(\frac{1}{e}-\frac{x}{1+x}\right)\right)^x=\alpha$, then the value of $\frac{\log_e\alpha}{1+\log_e\alpha}$ equals :

(A) e^{-2} (B) e^2 (C) e (D) e^{-1}

Open solution

Q34. JEE Main 2025 (Online) 22nd January Morning Shift

If $\sum\limits_{r=1}^{n}T_r=\frac{(2n-1)(2n+1)(2n+3)(2n+5)}{64}$, then $\lim\limits_{n\to\infty}\sum\limits_{r=1}^{n}\left(\frac{1}{T_r}\right)$ is equal to :

(A) $\frac{2}{3}$ (B) $\frac{1}{3}$ (C) 1 (D) 0

Open solution

Q35. JEE Main 2024 (Online) 9th April Evening Shift

$\lim\limits_{x\to0}\frac{e-(1+2x)^{\frac{1}{2x}}}{x}$ is equal to

(A) $\frac{-2}{e}$ (B) $e-e^2$ (C) 0 (D) e

Open solution

Q36. JEE Main 2024 (Online) 8th April Evening Shift

For $a,b>0$, let $f(x)=\begin{cases}\frac{\tan((a+1)x)+b\tan x}{x}, & x<0\\ 3, & x=0\\ \frac{\sqrt{ax+b^2x^2}-\sqrt{ax}}{b\sqrt{a}x\sqrt{x}}, & x>0\end{cases}$ be a continuous function at $x=0$. Then $\frac{b}{a}$ is equal to :

(A) 4 (B) 5 (C) 8 (D) 6

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q37. JEE Main 2024 (Online) 6th April Evening Shift

$\lim\limits_{n\to\infty}\frac{(1^2-1)(n-1)+(2^2-2)(n-2)+\cdots+((n-1)^2-(n-1))\cdot 1}{(1^3+2^3+\cdots\cdots+n^3)-(1^2+2^2+\cdots\cdots+n^2)}$ is equal to :

(A) $\frac{2}{3}$ (B) $\frac{1}{2}$ (C) $\frac{3}{4}$ (D) $\frac{1}{3}$

Open solution

Q38. JEE Main 2024 (Online) 5th April Evening Shift

Let $f:[-1,2]\to\mathbf{R}$ be given by $f(x)=2x^2+x+[x^2]-[x]$, where $[t]$ denotes the greatest integer less than or equal to t. The number of points, where f is not continuous, is :

(A) 5 (B) 6 (C) 4 (D) 3

Open solution

Q39. JEE Main 2024 (Online) 5th April Morning Shift

If the function $f(x)=\frac{\sin 3x+\alpha\sin x-\beta\cos 3x}{x^3}, x\in\mathbf{R}$, is continuous at $x=0$, then $f(0)$ is equal to :

(A) 4 (B) -2 (C) -4 (D) 2

Open solution

Q40. JEE Main 2024 (Online) 4th April Evening Shift

If the function

$$f(x)=\begin{cases}\frac{72^x-9^x-8^x+1}{\sqrt{2}-\sqrt{1+\cos x}}, & x\neq 0\\ a\log_e 2\log_e 3 & ,x=0\end{cases}$$

is continuous at $x=0$, then the value of a^2 is equal to

(A) 968 (B) 1250 (C) 1152 (D) 746

Open solution

Q41. JEE Main 2024 (Online) 4th April Morning Shift

Let $f:\mathbf{R}\to\mathbf{R}$ be a function given by

$$f(x)=\begin{cases}\frac{1-\cos 2x}{x^2}, & x<0\\ \alpha, & x=0,\\ \frac{\beta\sqrt{1-\cos x}}{x}, & x>0\end{cases}$$

where $\alpha,\beta\in\mathbf{R}$. If f is continuous at $x=0$, then $\alpha^2+\beta^2$ is equal to :

(A) 48 (B) 6 (C) 3 (D) 12

Open solution

Q42. JEE Main 2024 (Online) 1st February Evening Shift

Let $f(x)=\left|2x^2+5|x|-3\right|, x\in\mathbf{R}$. If m and n denote the number of points where f is not continuous and not differentiable respectively, then $\mathrm{m}+\mathrm{n}$ is equal to :

(A) 5 (B) 3 (C) 2 (D) 0

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q43. JEE Main 2024 (Online) 1st February Evening Shift

Let $f(x)=\begin{cases}x-1, x \text{ is even,} \\ 2x, \quad x \text{ is odd,}\end{cases} x \in \mathbf{N}$.

If for some $\mathrm{a} \in \mathbf{N}, f(f(f(\mathrm{a})))=21$, then $\lim\limits_{x \to \mathrm{a}^-}\left\{\frac{|x|^3}{\mathrm{a}}-\left[\frac{x}{\mathrm{a}}\right]\right\}$, where $[t]$ denotes the greatest integer less than or equal to t, is equal to :

Open solution

(A) 169 (B) 121 (C) 225 (D) 144

Q44. JEE Main 2024 (Online) 1st February Morning Shift

Let $f : \mathbf{R} \to \mathbf{R}$ be defined as :

$$f(x)=\begin{cases}\frac{a-b\cos 2x}{x^2}; & x<0 \\ x^2+cx+2; & 0 \le x \le 1 \\ 2x+1; & x>1\end{cases}$$

If f is continuous everywhere in $\mathbf{R}$ and m is the number of points where f is NOT differentiable then $\mathrm{m}+\mathrm{a}+\mathrm{b}+\mathrm{c}$ equals :

Open solution

(A) 1 (B) 4 (C) 3 (D) 2

Q45. JEE Main 2024 (Online) 31st January Evening Shift

Let $f : \mathbb{R} \to (0, \infty)$ be a strictly increasing function such that $\lim\limits_{x \to \infty}\frac{f(7x)}{f(x)}=1$. Then, the value of $\lim\limits_{x \to \infty}\left[\frac{f(5x)}{f(x)}-1\right]$ is equal to

Open solution

(A) 0 (B) 4 (C) 1 (D) 7/5

Q46. JEE Main 2024 (Online) 31st January Evening Shift

Consider the function $f : (0, \infty) \to \mathbb{R}$ defined by $f(x)=e^{-|\log_e x|}$. If m and n be respectively the number of points at which f is not continuous and f is not differentiable, then $m+n$ is

Open solution

(A) 0 (B) 1 (C) 2 (D) 3

Q47. JEE Main 2024 (Online) 31st January Morning Shift

$\lim\limits_{x \to 0}\frac{e^{2|\sin x|}-2|\sin x|-1}{x^2}$

Open solution

(A) is equal to 1 (B) does not exist (C) is equal to -1 (D) is equal to 2

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q48. JEE Main 2024 (Online) 31st January Morning Shift

Let $g(x)$ be a linear function and $f(x)=\begin{cases} g(x) & ,x\le 0 \\ \left(\frac{1+x}{2+x}\right)^{\frac{1}{x}} & ,x>0 \end{cases}$, is continuous at $x=0$. If $f'(1)=f(-1)$, then the value $g(3)$ is

Open solution

(A) $\log_e\left(\frac{4}{9}\right)-1$

(B) $\frac{1}{3}\log_e\left(\frac{4}{9e^{1/3}}\right)$

(C) $\log_e\left(\frac{4}{9e^{1/3}}\right)$

(D) $\frac{1}{3}\log_e\left(\frac{4}{9}\right)+1$

Q49. JEE Main 2024 (Online) 27th January Evening Shift

Consider the function $f:(0,2)\to\mathbf{R}$ defined by $f(x)=\frac{x}{2}+\frac{2}{x}$ and the function $g(x)$ defined by $g(x)=\begin{cases}\min\{f(t)\}, & 0<t\le x \text{ and } 0<x\le 1 \\ \frac{3}{2}+x, & 1<x<2\end{cases}$. Then,

Open solution

(A) g is continuous but not differentiable at $x=1$

(B) g is continuous and differentiable for all $x\in(0,2)$

(C) g is not continuous for all $x\in(0,2)$

(D) g is neither continuous nor differentiable at $x=1$

Q50. JEE Main 2024 (Online) 27th January Evening Shift

If $\lim\limits_{x\to 0}\frac{3+\alpha\sin x+\beta\cos x+\log_e(1-x)}{3\tan^2 x}=\frac{1}{3}$, then $2\alpha-\beta$ is equal to :

Open solution

(A) 2 (B) 1 (C) 5 (D) 7

Q51. JEE Main 2024 (Online) 27th January Morning Shift

Consider the function.

Open solution

$$f(x)=\begin{cases}\frac{a(7x-12-x^2)}{b|x^2-7x+12|} & ,x<3 \\ 2^{\frac{\sin(x-3)}{x-[x]}} & ,x>3 \\ b & ,x=3,\end{cases}$$

where $[x]$ denotes the greatest integer less than or equal to x. If S denotes the set of all ordered pairs (a, b) such that $f(x)$ is continuous at $x=3$, then the number of elements in S is :

(A) Infinitely many (B) 4 (C) 2 (D) 1

Q52. JEE Main 2024 (Online) 27th January Morning Shift

If $a=\lim\limits_{x\to 0}\frac{\sqrt{1+\sqrt{1+x^4}}-\sqrt{2}}{x^4}$ and $b=\lim\limits_{x\to 0}\frac{\sin^2 x}{\sqrt{2}-\sqrt{1+\cos x}}$, then the value of ab^3 is :

Open solution

(A) 36 (B) 25 (C) 32 (D) 30

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q53. JEE Main 2023 (Online) 15th April Morning Shift

Let $[x]$ denote the greatest integer function and

$f(x) = \max\{1 + x + [x], 2 + x, x + 2[x]\}, 0 \le x \le 2$. Let m be the number of points in $[0, 2]$, where f is not continuous and n be the number of points in $(0, 2)$, where f is not differentiable. Then $(m + n)^2 + 2$ is equal to :

Open solution

(A) 3 (B) 6 (C) 2 (D) 11

Q54. JEE Main 2023 (Online) 13th April Evening Shift

If $\lim\limits_{x \to 0} \frac{e^{ax} - \cos(bx) - \frac{cxe^{-cx}}{2}}{1 - \cos(2x)} = 17$, then $5a^2 + b^2$ is equal to

Open solution

(A) 64 (B) 68 (C) 72 (D) 76

Q55. JEE Main 2023 (Online) 11th April Evening Shift

Let f and g be two functions defined by

$f(x) = \begin{cases} x + 1, & x < 0 \\ |x - 1|, & x \ge 0 \end{cases}$ and $g(x) = \begin{cases} x + 1, & x < 0 \\ 1, & x \ge 0 \end{cases}$

Open solution

Then $(g \circ f)(x)$ is :

(A) continuous everywhere but not differentiable at $x = 1$

(B) differentiable everywhere

(C) not continuous at $x = -1$

(D) continuous everywhere but not differentiable exactly at one point

Q56. JEE Main 2023 (Online) 11th April Morning Shift

Let $f(x) = [x^2 - x] + |-x + [x]|$, where $x \in \mathbb{R}$ and $[t]$ denotes the greatest integer less than or equal to t. Then, f is :

Open solution

(A) continuous at $x = 0$, but not continuous at $x = 1$

(B) continuous at $x = 0$ and $x = 1$

(C) continuous at $x = 1$, but not continuous at $x = 0$

(D) not continuous at $x = 0$ and $x = 1$

Q57. JEE Main 2023 (Online) 8th April Evening Shift

If $\alpha > \beta > 0$ are the roots of the equation $ax^2 + bx + 1 = 0$, and $\lim\limits_{x \to \frac{1}{\alpha}} \left(\frac{1 - \cos(x^2 + bx + a)}{2(1 - \alpha x)^2}\right)^{\frac{1}{2}} = \frac{1}{k}\left(\frac{1}{\beta} - \frac{1}{\alpha}\right)$, then k is equal to :

Open solution

(A) 2β (B) β (C) α (D) 2α

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q58. JEE Main 2023 (Online) 8th April Morning Shift

$\lim\limits_{x\to 0}\left(\left(\frac{(1-\cos^2(3x))}{\cos^3(4x)}\right)\left(\frac{\sin^3(4x)}{(\log_e(2x+1))^5}\right)\right)$ is equal to _____.

(A) 15 (B) 18 (C) 9 (D) 24

Open solution

Q59. JEE Main 2023 (Online) 6th April Evening Shift

$\lim\limits_{n\to\infty}\left\{\left(2^{\frac{1}{2}}-2^{\frac{1}{3}}\right)\left(2^{\frac{1}{2}}-2^{\frac{1}{5}}\right).....\left(2^{\frac{1}{2}}-2^{\frac{1}{2n+1}}\right)\right\}$ is equal to :

(A) $\sqrt{2}$ (B) 1 (C) $\frac{1}{\sqrt{2}}$ (D) 0

Open solution

Q60. JEE Main 2023 (Online) 6th April Morning Shift

Let $a_1, a_2, a_3, \ldots, a_n$ be n positive consecutive terms of an arithmetic progression. If $d > 0$ is its common difference, then

$\lim\limits_{n\to\infty}\sqrt{\frac{d}{n}}\left(\frac{1}{\sqrt{a_1}+\sqrt{a_2}}+\frac{1}{\sqrt{a_2}+\sqrt{a_3}}+.....+\frac{1}{\sqrt{a_{n-1}}+\sqrt{a_n}}\right)$ is

(A) $\frac{1}{\sqrt{d}}$ (B) 1 (C) 0 (D) $\sqrt{d}$

Open solution

Q61. JEE Main 2023 (Online) 31st January Evening Shift

$\lim\limits_{x\to\infty}\frac{(\sqrt{3x+1}+\sqrt{3x-1})^6+(\sqrt{3x+1}-\sqrt{3x-1})^6}{\left(x+\sqrt{x^2-1}\right)^6+\left(x-\sqrt{x^2-1}\right)^6}x^3$

(A) is equal to 9 (B) is equal to $\frac{27}{2}$

(C) does not exist (D) is equal to 27

Open solution

Q62. JEE Main 2023 (Online) 30th January Evening Shift

Let f, g and h be the real valued functions defined on $\mathbb{R}$ as

$$f(x)=\begin{cases}\frac{x}{|x|}, & x\neq 0\\ 1, & x=0\end{cases}$$

$$g(x)=\begin{cases}\frac{\sin(x+1)}{(x+1)}, & x\neq -1\\ 1, & x=-1\end{cases}$$

and $h(x)=2[x]-f(x)$, where $[x]$ is the greatest integer $\leq x$. Then the

value of $\lim\limits_{x\to 1} g(h(x-1))$ is :

(A) 1 (B) -1 (C) $\sin(1)$ (D) 0

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q63. JEE Main 2023 (Online) 29th January Morning Shift

Let $x = 2$ be a root of the equation $x^2 + px + q = 0$ and $f(x) = \begin{cases} \frac{1-\cos(x^2-4px+q^2+8q+16)}{(x-2p)^4}, & x \neq 2p \\ 0, & x = 2p \end{cases}$

Then $\lim\limits_{x \to 2p^+} [f(x)]$, where $[.]$ denotes greatest integer function, is

(A) 2 (B) 1 (C) 0 (D) -1

Open solution

Q64. JEE Main 2023 (Online) 25th January Evening Shift

If the function $f(x) = \begin{cases} (1 + |\cos x|)^{\frac{\lambda}{|\cos x|}}, & 0 < x < \frac{\pi}{2} \\ \mu, & x = \frac{\pi}{2} \\ e^{\frac{\cot 6x}{\cot 4x}}, & \frac{\pi}{2} < x < \pi \end{cases}$

is continuous at $x = \frac{\pi}{2}$, then $9\lambda + 6\log_e \mu + \mu^6 - e^{6\lambda}$ is equal to

(A) 11 (B) 10 (C) 8 (D) $2e^4 + 8$

Open solution

Q65. JEE Main 2023 (Online) 25th January Morning Shift

The value of $\lim\limits_{n \to \infty} \frac{1+2-3+4+5-6+ \ldots + (3n-2)+(3n-1)-3n}{\sqrt{2n^4+4n+3}-\sqrt{n^4+5n+4}}$ is :

(A) $\frac{3}{2\sqrt{2}}$ (B) $\frac{3}{2}(\sqrt{2}+1)$ (C) $3(\sqrt{2}+1)$ (D) $\frac{\sqrt{2}+1}{2}$

Open solution

Q66. JEE Main 2023 (Online) 24th January Evening Shift

The set of all values of a for which $\lim\limits_{x \to a}([x-5] - [2x+2]) = 0$, where $[\alpha]$ denotes the greatest integer less than or equal to α is equal to

(A) $[-7.5, -6.5]$ (B) $(-7.5, -6.5]$ (C) $[-7.5, -6.5)$ (D) $(-7.5, -6.5)$

Open solution

Q67. JEE Main 2023 (Online) 24th January Morning Shift

$\lim\limits_{t \to 0} \left(1^{\frac{1}{\sin^2 t}} + 2^{\frac{1}{\sin^2 t}} + \ldots + n^{\frac{1}{\sin^2 t}}\right)^{\sin^2 t}$ is equal to

(A) $\frac{n(n+1)}{2}$ (B) n (C) $n^2 + n$ (D) n^2

Open solution

Q68. JEE Main 2023 (Online) 24th January Morning Shift

Let $f(x) = \begin{cases} x^2 \sin\left(\frac{1}{x}\right), & x \neq 0 \\ 0, & x = 0 \end{cases}$

Then at $x = 0$

(A) f is continuous but f' is not continuous (B) f and f' both are continuous

(C) f is continuous but not differentiable (D) f' is continuous but not differentiable

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q69. JEE Main 2022 (Online) 30th July Evening Shift

Open solution

Let

$$f(x)=\begin{cases}\dfrac{\sin(ax+3x)+\sin x}{x}, & x<0\\ 2, & x=0\\ \dfrac{(x+6bx^2)^{\frac{1}{3}}-x^{\frac{1}{3}}}{3x^{4/3}}, & x>0\end{cases}$$

be continuous at $x=0$. If $(1-k)\,f\left(\dfrac{7}{18}\right)=ab$, then k is equal to

(A) -6 (B) 15 (C) -13 (D) 8

Q70. JEE Main 2022 (Online) 29th July Evening Shift

Open solution

Let the function $f(x)=\begin{cases}\frac{\log_e(1+5x)-\log_e(1+\alpha x)}{x} & ;\text{ if } x\neq 0\\ 10 & ;\text{ if } x=0\end{cases}$ be continuous at $x=0$.

Then α is equal to

(A) 10 (B) -10 (C) 5 (D) -5

Q71. JEE Main 2022 (Online) 29th July Morning Shift

Open solution

If $\lim\limits_{x\to 0}\frac{\alpha e^x+\beta e^{-x}+\gamma\sin x}{x\sin^2 x}=\frac{2}{3}$, where $\alpha,\beta,\gamma\in\mathbf{R}$, then which of the following is NOT correct?

(A) $\alpha^2+\beta^2+\gamma^2=6$ (B) $\alpha\beta+\beta\gamma+\gamma\alpha+1=0$

(C) $\alpha\beta^2+\beta\gamma^2+\gamma\alpha^2+3=0$ (D) $\alpha^2-\beta^2+\gamma^2=4$

Q72. JEE Main 2022 (Online) 29th July Morning Shift

Open solution

The number of points, where the function $f:\mathbf{R}\to\mathbf{R}$,

$f(x)=|x-1|\cos|x-2|\sin|x-1|+(x-3)\left|x^2-5x+4\right|$, is NOT differentiable, is :

(A) 1 (B) 2 (C) 3 (D) 4

Q73. JEE Main 2022 (Online) 28th July Evening Shift

Open solution

The function $f:\mathbb{R}\to\mathbb{R}$ defined by

$f(x)=\lim\limits_{n\to\infty}\frac{\cos(2\pi x)-x^{2n}\sin(x-1)}{1+x^{2n+1}-x^{2n}}$ is continuous for all x in :

(A) $R-\{-1\}$ (B) $\mathbb{R}-\{-1,1\}$ (C) $R-\{1\}$ (D) $R-\{0\}$

Q74. JEE Main 2022 (Online) 27th July Evening Shift

Open solution

If for $\mathrm{p}\neq\mathrm{q}\neq 0$, the function $f(x)=\frac{\sqrt[7]{\mathrm{p}(729+x)}-3}{\sqrt[3]{729+\mathrm{q}x}-9}$ is continuous at $x=0$, then :

(A) $7pq\,f(0)-1=0$ (B) $63q\,f(0)-\mathrm{p}^2=0$ (C) $21q\,f(0)-\mathrm{p}^2=0$ (D) $7pq\,f(0)-9=0$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q75. JEE Main 2022 (Online) 26th July Evening Shift

Let $\beta = \lim\limits_{x\to 0} \frac{\alpha x - (e^{3x}-1)}{\alpha x(e^{3x}-1)}$ for some $\alpha \in \mathbb{R}$. Then the value of $\alpha + \beta$ is :

(A) $\frac{14}{5}$ (B) $\frac{3}{2}$ (C) $\frac{5}{2}$ (D) $\frac{7}{2}$

Open solution

Q76. JEE Main 2022 (Online) 26th July Morning Shift

Let f : R $\to$ R be a continuous function such that $f(3x) - f(x) = x$. If $f(8) = 7$, then $f(14)$ is equal to :

(A) 4 (B) 10 (C) 11 (D) 16

Open solution

Q77. JEE Main 2022 (Online) 26th July Morning Shift

If the function $f(x) = \begin{cases} \frac{\log_e(1-x+x^2)+\log_e(1+x+x^2)}{\sec x - \cos x} & , \quad x \in \left(\frac{-\pi}{2}, \frac{\pi}{2}\right) - \{0\} \\ k & , \quad x = 0 \end{cases}$ is continuous at x = 0, then k is equal to:

(A) 1 (B) -1 (C) e (D) 0

Open solution

Q78. JEE Main 2022 (Online) 26th July Morning Shift

If $f(x) = \begin{cases} x + a & , \quad x \le 0 \\ |x-4| & , \quad x > 0 \end{cases}$ and $g(x) = \begin{cases} x+1 & , \quad x < 0 \\ (x-4)^2 + b & , \quad x \ge 0 \end{cases}$ are continuous on R, then $(gof)(2) + (fog)(-2)$ is equal to :

(A) -10 (B) 10 (C) 8 (D) -8

Open solution

Q79. JEE Main 2022 (Online) 25th July Evening Shift

$\lim\limits_{x\to\frac{\pi}{4}} \frac{8\sqrt{2} - (\cos x + \sin x)^7}{\sqrt{2} - \sqrt{2}\sin 2x}$ is equal to

(A) 14 (B) 7 (C) $14\sqrt{2}$ (D) $7\sqrt{2}$

Open solution

Q80. JEE Main 2022 (Online) 25th July Morning Shift

If $\lim\limits_{n\to\infty} \left(\sqrt{n^2 - n - 1} + n\alpha + \beta\right) = 0$, then $8(\alpha + \beta)$ is equal to :

(A) 4 (B) -8 (C) -4 (D) 8

Open solution

Q81. JEE Main 2022 (Online) 29th June Evening Shift

The value of $\lim\limits_{x\to 1} \frac{(x^2-1)\sin^2(\pi x)}{x^4 - 2x^3 + 2x - 1}$ is equal to:

(A) $\frac{\pi^2}{6}$ (B) $\frac{\pi^2}{3}$ (C) $\frac{\pi^2}{2}$ (D) π^2

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q82. JEE Main 2022 (Online) 28th June Evening Shift

Open solution

Let f, g : R $\to$ R be functions defined by

$f(x) = \begin{cases} [x] & , \quad x < 0 \\ |1-x| & , \quad x \geq 0 \end{cases}$ and $g(x) = \begin{cases} e^x - x & , \quad x < 0 \\ (x-1)^2 - 1 & , \quad x \geq 0 \end{cases}$ where [x] denote the greatest integer less than or equal to x. Then, the function fog is discontinuous at exactly :

(A) one point (B) two points (C) three points (D) four points

Q83. JEE Main 2022 (Online) 28th June Evening Shift

Open solution

The value of

$\lim\limits_{n\to\infty} 6\tan\left\{\sum\limits_{r=1}^{n}\tan^{-1}\left(\frac{1}{r^2+3r+3}\right)\right\}$ is equal to :

(A) 1 (B) 2 (C) 3 (D) 6

Q84. JEE Main 2022 (Online) 28th June Morning Shift

Open solution

Let f : R $\to$ R be defined as

$$f(x) = \begin{cases} [e^x], & x < 0 \\ ae^x + [x-1], & 0 \leq x < 1 \\ b + [\sin(\pi x)], & 1 \leq x < 2 \\ [e^{-x}] - c, & x \geq 2 \end{cases}$$

where a, b, c $\in$ R and [t] denotes greatest integer less than or equal to t. Then, which of the following statements is true?

(A) There exists a, b, c $\in$ R such that f is continuous on R.

(B) If f is discontinuous at exactly one point, then a + b + c = 1

(C) If f is discontinuous at exactly one point, then a + b + c $\neq$ 1

(D) f is discontinuous at at least two points, for any values of a, b and c

Q85. JEE Main 2022 (Online) 27th June Morning Shift

Open solution

Let a be an integer such that $\lim\limits_{x\to 7}\frac{18-[1-x]}{[x-3a]}$ exists, where [t] is greatest integer $\leq$ t. Then a is equal to :

(A) -6 (B) -2 (C) 2 (D) 6

Q86. JEE Main 2022 (Online) 26th June Evening Shift

Open solution

$\lim\limits_{x\to 0}\frac{\cos(\sin x)-\cos x}{x^4}$ is equal to :

(A) $\frac{1}{3}$ (B) $\frac{1}{4}$ (C) $\frac{1}{6}$ (D) $\frac{1}{12}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q87. JEE Main 2022 (Online) 26th June Evening Shift

Let f(x) = min {1, 1 + x sin x}, $0 \le x \le 2\pi$. If m is the number of points, where f is not differentiable and n is the number of points, where f is not continuous, then the ordered pair (m, n) is equal to

(A) (2, 0) (B) (1, 0) (C) (1, 1) (D) (2, 1)

Open solution

Q88. JEE Main 2022 (Online) 26th June Morning Shift

$\lim\limits_{x\to\frac{1}{\sqrt{2}}} \frac{\sin(\cos^{-1}x)-x}{1-\tan(\cos^{-1}x)}$ is equal to :

(A) $\sqrt{2}$ (B) $-\sqrt{2}$ (C) $\frac{1}{\sqrt{2}}$ (D) $-\frac{1}{\sqrt{2}}$

Open solution

Q89. JEE Main 2022 (Online) 26th June Morning Shift

Let f, g : R $\to$ R be two real valued functions defined as $f(x) = \begin{cases} -|x+3| & , \quad x < 0 \\ e^x & , \quad x \ge 0 \end{cases}$ and $g(x) = \begin{cases} x^2 + k_1 x & , \quad x < 0 \\ 4x + k_2 & , \quad x \ge 0 \end{cases}$, where k_1 and k_2 are real constants. If (gof) is differentiable at x = 0, then (gof) (-4) + (gof) (4) is equal to :

(A) $4(e^4+1)$ (B) $2(2e^4+1)$ (C) $4e^4$ (D) $2(2e^4-1)$

Open solution

Q90. JEE Main 2022 (Online) 25th June Evening Shift

$\lim\limits_{x\to\frac{\pi}{2}} \left(\tan^2 x \left((2\sin^2 x + 3\sin x + 4)^{\frac{1}{2}} - (\sin^2 x + 6\sin x + 2)^{\frac{1}{2}}\right)\right)$ is equal to

(A) $\frac{1}{12}$ (B) $-\frac{1}{18}$ (C) $-\frac{1}{12}$ (D) $\frac{1}{6}$

Open solution

Q91. JEE Main 2022 (Online) 25th June Morning Shift

Let f(x) be a polynomial function such that $f(x) + f'(x) + f''(x) = x^5 + 64$. Then, the value of $\lim\limits_{x\to 1} \frac{f(x)}{x-1}$ is equal to:

(A) -15 (B) -60 (C) 60 (D) 15

Open solution

Q92. JEE Main 2022 (Online) 24th June Evening Shift

Let $f(x) = \begin{cases} \frac{\sin(x-[x])}{x-[x]} & , x \in (-2,-1) \\ \max\{2x, 3[|x|]\} & , |x| < 1 \\ 1 & , otherwise \end{cases}$

where [t] denotes greatest integer $\le$ t. If m is the number of points where f is not continuous and n is the number of points where f is not differentiable, then the ordered pair (m, n) is :

(A) (3, 3) (B) (2, 4) (C) (2, 3) (D) (3, 4)

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q93. JEE Main 2021 (Online) 31st August Evening Shift

If $\alpha = \lim\limits_{x \to \frac{\pi}{4}} \frac{\tan^3 x - \tan x}{\cos\left(x + \frac{\pi}{4}\right)}$ and $\beta = \lim\limits_{x \to 0} (\cos x)^{\cot x}$ are the roots of the equation, $ax^2 + bx - 4 = 0$, then the ordered pair (a, b) is :

Open solution

(A) $(1, -3)$ (B) $(-1, 3)$ (C) $(-1, -3)$ (D) $(1, 3)$

Q94. JEE Main 2021 (Online) 31st August Morning Shift

The function

$f(x) = \left|x^2 - 2x - 3\right| . e^{\left|9x^2 - 12x + 4\right|}$ is not differentiable at exactly :

Open solution

(A) four points (B) three points (C) two points (D) one point

Q95. JEE Main 2021 (Online) 31st August Morning Shift

If the function

$$f(x) = \begin{cases} \frac{1}{x}\log_e\left(\frac{1+\frac{x}{a}}{1-\frac{x}{b}}\right) & , \quad x < 0 \\ k & , \quad x = 0 \\ \frac{\cos^2 x - \sin^2 x - 1}{\sqrt{x^2+1}-1} & , \quad x > 0 \end{cases}$$ is continuous

Open solution

at x = 0, then $\frac{1}{a} + \frac{1}{b} + \frac{4}{k}$ is equal to :

(A) -5 (B) 5 (C) -4 (D) 4

Q96. JEE Main 2021 (Online) 31st August Morning Shift

$\lim\limits_{x \to 0} \frac{\sin^2(\pi\cos^4 x)}{x^4}$ is equal to :

Open solution

(A) π^2 (B) $2\pi^2$ (C) $4\pi^2$ (D) 4π

Q97. JEE Main 2021 (Online) 27th August Evening Shift

If $\lim\limits_{x \to \infty}\left(\sqrt{x^2 - x + 1} - ax\right) = b$, then the ordered pair (a, b) is :

Open solution

(A) $\left(1, \frac{1}{2}\right)$ (B) $\left(1, -\frac{1}{2}\right)$

(C) $\left(-1, \frac{1}{2}\right)$ (D) $\left(-1, -\frac{1}{2}\right)$

Q98. JEE Main 2021 (Online) 27th August Morning Shift

If α, β are the distinct roots of $x^2 + bx + c = 0$, then

Open solution

$\lim\limits_{x \to \beta} \frac{e^{2(x^2+bx+c)} - 1 - 2(x^2+bx+c)}{(x-\beta)^2}$ is equal to :

(A) $b^2 + 4c$ (B) $2(b^2 + 4c)$ (C) $2(b^2 - 4c)$ (D) $b^2 - 4c$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q99. JEE Main 2021 (Online) 26th August Evening Shift

Open solution

Let [t] denote the greatest integer less than or equal to t. Let
f(x) = x − [x], g(x) = 1 − x + [x], and h(x) = min{f(x), g(x)}, x ∈ [−2, 2]. Then h is :

(A) continuous in [−2, 2] but not differentiable at more than four points in (−2, 2)

(B) not continuous at exactly three points in [−2, 2]

(C) continuous in [−2, 2] but not differentiable at exactly three points in (−2, 2)

(D) not continuous at exactly four points in [−2, 2]

Q100. JEE Main 2021 (Online) 26th August Evening Shift

Open solution

$\lim\limits_{x\to 2}\left(\sum\limits_{n=1}^{9}\frac{x}{n(n+1)x^2+2(2n+1)x+4}\right)$ is equal to :

(A) $\frac{9}{44}$ (B) $\frac{5}{24}$ (C) $\frac{1}{5}$ (D) $\frac{7}{36}$

Q101. JEE Main 2021 (Online) 27th July Evening Shift

Open solution

The value of

$\lim\limits_{x\to 0}\left(\frac{x}{\sqrt[8]{1-\sin x}-\sqrt[8]{1+\sin x}}\right)$ is equal to :

(A) 0 (B) 4 (C) −4 (D) −1

Q102. JEE Main 2021 (Online) 27th July Evening Shift

Open solution

Let $f:[0,\infty)\to[0,3]$ be a function defined by

$$f(x)=\begin{cases}\max\{\sin t: 0\le t\le x\}, & 0\le x\le\pi\\ 2+\cos x, & x>\pi\end{cases}$$

Then which of the following is true?

(A) f is continuous everywhere but not differentiable exactly at one point in (0, ∞)

(B) f is differentiable everywhere in (0, ∞)

(C) f is not continuous exactly at two points in (0, ∞)

(D) f is continuous everywhere but not differentiable exactly at two points in (0, ∞)

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q103. JEE Main 2021 (Online) 27th July Morning Shift

Let $f:\left(-\frac{\pi}{4},\frac{\pi}{4}\right)\to R$ be defined as $f(x)=\begin{cases}(1+|\sin x|)^{\frac{3a}{|\sin x|}} & , \quad -\frac{\pi}{4}<x<0\\ b & , \quad x=0\\ e^{\cot 4x/\cot 2x} & , \quad 0<x<\frac{\pi}{4}\end{cases}$

Open solution

If f is continuous at x = 0, then the value of 6a + b^2 is equal to :

(A) $1-e$ (B) $e-1$ (C) 1 + e (D) e

Q104. JEE Main 2021 (Online) 27th July Morning Shift

Let f : R $\to$ R be a function such that f(2) = 4 and f'(2) = 1. Then, the value of $\lim\limits_{x\to 2}\frac{x^2f(2)-4f(x)}{x-2}$ is equal to :

Open solution

(A) 4 (B) 8 (C) 16 (D) 12

Q105. JEE Main 2021 (Online) 25th July Morning Shift

Let f : R $\to$ R be defined as

$$f(x)=\begin{cases}\frac{\lambda|x^2-5x+6|}{\mu(5x-x^2-6)}, & x<2\\ e^{\frac{\tan(x-2)}{x-[x]}}, & x>2\\ \mu, & x=2\end{cases}$$

Open solution

where [x] is the greatest integer is than or equal to x. If f is continuous at x = 2, then $\lambda+\mu$ is equal to :

(A) e(−e + 1) (B) e(e − 2) (C) 1 (D) 2e − 1

Q106. JEE Main 2021 (Online) 22th July Evening Shift

Let f : R $\to$ R be defined as $f(x)=\begin{cases}\frac{x^3}{(1-\cos 2x)^2}\log_e\left(\frac{1+2xe^{-2x}}{(1-xe^{-x})^2}\right), & x\neq 0\\ \alpha, & x=0\end{cases}$

Open solution

If f is continuous at x = 0, then α is equal to :

(A) 1 (B) 3 (C) 0 (D) 2

Q107. JEE Main 2021 (Online) 20th July Morning Shift

Let a function f : R $\to$ R be defined as $f(x)=\begin{cases}\sin x-e^x & if \quad x\leq 0\\ a+[-x] & if \quad 0<x<1\\ 2x-b & if \quad x\geq 1\end{cases}$

Open solution

where [x] is the greatest integer less than or equal to x. If f is continuous on R, then (a + b) is equal to:

(A) 4 (B) 3 (C) 2 (D) 5

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q108. JEE Main 2021 (Online) 18th March Evening Shift

Let f : R $\rightarrow$ R be a function defined as

$$f(x) = \begin{cases} \frac{\sin(a+1)x+\sin 2x}{2x}, if\, x < 0 \\ b, if\, x = 0 \\ \frac{\sqrt{x+bx^3}-\sqrt{x}}{bx^{5/2}}, if\, x > 0 \end{cases}$$

If f is continuous at x = 0, then the value of a + b is equal to :

(A) -3 (B) -2 (C) $-\frac{5}{2}$ (D) $-\frac{3}{2}$

Open solution

Q109. JEE Main 2021 (Online) 18th March Morning Shift

If $\lim\limits_{x\to 0} \frac{\sin^{-1}x-\tan^{-1}x}{3x^3}$ is equal to L, then the value of (6L + 1) is

(A) $\frac{1}{6}$ (B) $\frac{1}{2}$ (C) 6 (D) 2

Open solution

Q110. JEE Main 2021 (Online) 18th March Morning Shift

If $f(x) = \begin{cases} \frac{1}{|x|} & ; |x| \geq 1 \\ ax^2 + b & ; |x| < 1 \end{cases}$ is differentiable at every point of the domain, then the values of a and b are respectively :

(A) $\frac{1}{2}, \frac{1}{2}$ (B) $\frac{1}{2}, -\frac{3}{2}$ (C) $\frac{5}{2}, -\frac{3}{2}$ (D) $-\frac{1}{2}, \frac{3}{2}$

Open solution

Q111. JEE Main 2021 (Online) 17th March Evening Shift

The value of the limit

$\lim\limits_{\theta\to 0} \frac{\tan(\pi\cos^2\theta)}{\sin(2\pi\sin^2\theta)}$ is equal to :

(A) 0 (B) $-\frac{1}{2}$ (C) $\frac{1}{4}$ (D) $-\frac{1}{4}$

Open solution

Q112. JEE Main 2021 (Online) 17th March Evening Shift

The value of $\lim\limits_{n\to\infty} \frac{[r]+[2r]+...+[nr]}{n^2}$, where r is a non-zero real number and [r] denotes the greatest integer less than or equal to r, is equal to :

(A) r (B) $\frac{r}{2}$ (C) 0 (D) 2r

Open solution

Q113. JEE Main 2021 (Online) 17th March Morning Shift

The value of

$\lim\limits_{x\to 0^+} \frac{\cos^{-1}(x-[x]^2).\sin^{-1}(x-[x]^2)}{x-x^3}$, where [x] denotes the greatest integer $\leq$ x is :

(A) π (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{2}$ (D) 0

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q114. JEE Main 2021 (Online) 16th March Evening Shift

Let f : S $\to$ S where S = (0, ∞) be a twice differentiable function such that f(x + 1) = xf(x). If g : S $\to$ R be defined as g(x) = $\log_e$ f(x), then the value of |g"(5) $-$ g"(1)| is equal to :

(A) 1 (B) $\frac{187}{144}$ (C) $\frac{197}{144}$ (D) $\frac{205}{144}$

Open solution

Q115. JEE Main 2021 (Online) 16th March Evening Shift

Let $\alpha \in$ R be such that the function $f(x) = \begin{cases} \frac{\cos^{-1}(1-\{x\}^2)\sin^{-1}(1-\{x\})}{\{x\}-\{x\}^3}, & x \neq 0 \\ \alpha, & x = 0 \end{cases}$ is continuous at x = 0,

where {x} = x $-$ [x], [x] is the greatest integer less than or equal to x. Then :

(A) no such α exists (B) α = 0

(C) $\alpha = \frac{\pi}{4}$ (D) $\alpha = \frac{\pi}{\sqrt{2}}$

Open solution

Q116. JEE Main 2021 (Online) 16th March Morning Shift

Let $S_k = \sum_{r=1}^{k} \tan^{-1}\left(\frac{6^r}{2^{2r+1}+3^{2r+1}}\right)$. Then $\lim_{k\to\infty} S_k$ is equal to :

(A) $\cot^{-1}\left(\frac{3}{2}\right)$ (B) $\frac{\pi}{2}$

(C) $\tan^{-1}(3)$ (D) $\tan^{-1}\left(\frac{3}{2}\right)$

Open solution

Q117. JEE Main 2021 (Online) 16th March Morning Shift

Let the functions f : R $\to$ R and g : R $\to$ R be defined as :

$f(x) = \begin{cases} x+2, & x < 0 \\ x^2, & x \geq 0 \end{cases}$ and

$g(x) = \begin{cases} x^3, & x < 1 \\ 3x-2, & x \geq 1 \end{cases}$

Then, the number of points in R where (fog) (x) is NOT differentiable is equal to :

(A) 0 (B) 3 (C) 1 (D) 2

Open solution

Q118. JEE Main 2021 (Online) 26th February Evening Shift

Let f(x) be a differentiable function at x = a with f'(a) = 2 and f(a) = 4.

Then $\lim_{x\to a} \frac{xf(a)-af(x)}{x-a}$ equals :

(A) 4 $-$ 2a (B) 2a + 4 (C) a + 4 (D) 2a $-$ 4

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q119. JEE Main 2021 (Online) 26th February Evening Shift

Let $f(x) = \sin^{-1}x$ and $g(x) = \frac{x^2-x-2}{2x^2-x-6}$. If $g(2) = \lim_{x\to 2} g(x)$, then the domain of the function fog is :

(A) $(-\infty, -2] \cup \left[-\frac{4}{3}, \infty\right)$

(B) $(-\infty, -2] \cup [-1, \infty)$

(C) $(-\infty, -2] \cup \left[-\frac{3}{2}, \infty\right)$

(D) $(-\infty, -1] \cup [2, \infty)$

Open solution

Q120. JEE Main 2021 (Online) 26th February Evening Shift

Let f : R $\to$ R be defined as

$$f(x) = \begin{cases} 2\sin\left(-\frac{\pi x}{2}\right), if\, x < -1 \\ |ax^2 + x + b|, \; if -1 \le x \le 1 \\ \sin(\pi x), \; if\, x > 1 \end{cases}$$ If f(x) is continuous on R, then a + b equals :

(A) -3

(B) 3

(C) -1

(D) 1

Open solution

Q121. JEE Main 2021 (Online) 26th February Morning Shift

The value of $\lim_{h\to 0} 2\left\{\frac{\sqrt{3}\sin\left(\frac{\pi}{6}+h\right)-\cos\left(\frac{\pi}{6}+h\right)}{\sqrt{3}h\left(\sqrt{3}\cosh - \sinh\right)}\right\}$ is :

(A) $\frac{4}{3}$

(B) $\frac{2}{3}$

(C) $\frac{3}{4}$

(D) $\frac{2}{\sqrt{3}}$

Open solution

Q122. JEE Main 2021 (Online) 25th February Morning Shift

$\lim_{n\to\infty}\left(1 + \frac{1+\frac{1}{2}+.....+\frac{1}{n}}{n^2}\right)^n$ is equal to :

(A) $\frac{1}{2}$

(B) 1

(C) 0

(D) $\frac{1}{e}$

Open solution

Q123. JEE Main 2021 (Online) 24th February Morning Shift

If f : R $\to$ R is a function defined by f(x)= [x - 1] $\cos\left(\frac{2x-1}{2}\right)\pi$, where [.] denotes the greatest integer function, then f is :

(A) continuous for every real x

(B) discontinuous at all integral values of x except at x = 1

(C) discontinuous only at x = 1

(D) continuous only at x = 1

Open solution

Q124. JEE Main 2020 (Online) 6th September Evening Slot

Let f : R $\to$ R be a function defined by
f(x) = max {x, x^2}. Let S denote the set of all points in R, where f is not differentiable. Then S is equal to :

(A) {0, 1}

(B) {0}

(C) ϕ(an empty set)

(D) {1}

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q125. JEE Main 2020 (Online) 6th September Evening Slot

For all twice differentiable functions f : R $\rightarrow$ R, with f(0) = f(1) = f'(0) = 0

(A) f"(x) $\neq$ 0, at every point x $\in$ (0, 1)

(B) f"(x) = 0, for some x $\in$ (0, 1)

(C) f"(0) = 0

(D) f"(x) = 0, at every point x $\in$ (0, 1)

Open solution

Q126. JEE Main 2020 (Online) 5th September Evening Slot

$$\lim_{x\to 0} \frac{x\left(e^{\left(\sqrt{1+x^2+x^4}-1\right)/x}-1\right)}{\sqrt{1+x^2+x^4}-1}$$

(A) is equal to 0.

(B) is equal to $\sqrt{e}$.

(C) is equal to 1.

(D) does not exist.

Open solution

Q127. JEE Main 2020 (Online) 5th September Morning Slot

If the function

$$f(x)=\begin{cases} k_1(x-\pi)^2-1, & x\le\pi \\ k_2\cos x, & x>\pi \end{cases}$$ is

twice differentiable, then the ordered pair (k_1, k_2) is equal to :

(A) $\left(\frac{1}{2},-1\right)$ (B) (1, 1) (C) (1, 0) (D) $\left(\frac{1}{2},1\right)$

Open solution

Q128. JEE Main 2020 (Online) 5th September Morning Slot

If α is positive root of the equation, p(x) = x^2 - x - 2 = 0, then

$\lim_{x\to\alpha^+} \frac{\sqrt{1-\cos(p(x))}}{x+\alpha-4}$ is equal to :

(A) $\frac{1}{\sqrt{2}}$ (B) $\frac{1}{2}$ (C) $\frac{3}{\sqrt{2}}$ (D) $\frac{3}{2}$

Open solution

Q129. JEE Main 2020 (Online) 4th September Evening Slot

Let $f:(0,\infty)\rightarrow(0,\infty)$ be a differentiable function such that f(1) = e and $\lim_{t\to x} \frac{t^2f^2(x)-x^2f^2(t)}{t-x}=0$. If f(x) = 1, then x is equal to :

(A) $\frac{1}{e}$ (B) e (C) $\frac{1}{2e}$ (D) 2e

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q130. JEE Main 2020 (Online) 4th September Evening Slot

The function

$$f(x)=\begin{cases}\frac{\pi}{4}+\tan^{-1}x, & |x|\le 1\\ \frac{1}{2}(|x|-1), & |x|>1\end{cases}$$ is :

(A) continuous on R−{−1} and differentiable on R−{−1, 1}

(B) both continuous and differentiable on R−{1}

(C) both continuous and differentiable on R−{−1}

(D) continuous on R−{1} and differentiable on R−{−1, 1}

Open solution

Q131. JEE Main 2020 (Online) 3rd September Evening Slot

$\lim\limits_{x\to a}\frac{(a+2x)^{\frac{1}{3}}-(3x)^{\frac{1}{3}}}{(3a+x)^{\frac{1}{3}}-(4x)^{\frac{1}{3}}}$ $(a\neq 0)$ is equal to :

(A) $\left(\frac{2}{9}\right)\left(\frac{2}{3}\right)^{\frac{1}{3}}$ (B) $\left(\frac{2}{3}\right)\left(\frac{2}{9}\right)^{\frac{1}{3}}$

(C) $\left(\frac{2}{3}\right)^{\frac{4}{3}}$ (D) $\left(\frac{2}{9}\right)^{\frac{4}{3}}$

Open solution

Q132. JEE Main 2020 (Online) 3rd September Morning Slot

Let [t] denote the greatest integer $\le$ t. If for some $\lambda\in$ R - {1, 0}, $\lim\limits_{x\to 0}\left|\frac{1-x+|x|}{\lambda-x+[x]}\right|$ = L, then L is equal to :

(A) 1 (B) 2 (C) 0 (D) $\frac{1}{2}$

Open solution

Q133. JEE Main 2020 (Online) 2nd September Evening Slot

$\lim\limits_{x\to 0}\left(\tan\left(\frac{\pi}{4}+x\right)\right)^{\frac{1}{x}}$ is equal to :

(A) 2 (B) 1 (C) e (D) e^2

Open solution

Q134. JEE Main 2020 (Online) 2nd September Morning Slot

If a function f(x) defined by

$$f(x)=\begin{cases}ae^{x}+be^{-x}, & -1\le x<1\\ cx^{2}, & 1\le x\le 3\\ ax^{2}+2cx, & 3<x\le 4\end{cases}$$

be continuous for some a, b, c $\in$ R and f'(0) + f'(2) = e, then the value of a is :

(A) $\frac{e}{e^2-3e-13}$ (B) $\frac{1}{e^2-3e+13}$ (C) $\frac{e}{e^2-3e+13}$ (D) $\frac{e}{e^2+3e+13}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q135. JEE Main 2020 (Online) 9th January Evening Slot

Let [t] denote the greatest integer $\leq$ t and $\lim_{x\to 0} x\left[\frac{4}{x}\right] = A$.

Then the function, f(x) = $[x^2]\sin(\pi x)$ is discontinuous, when x is equal to :

(A) $\sqrt{A+1}$ (B) $\sqrt{A+5}$ (C) $\sqrt{A+21}$ (D) $\sqrt{A}$

Open solution

Q136. JEE Main 2020 (Online) 9th January Morning Slot

Let f be any function continuous on [a, b] and twice differentiable on (a, b). If for all x $\in$ (a, b), f'(x) > 0 and f"(x) < 0, then for any c $\in$ (a, b), $\frac{f(c)-f(a)}{f(b)-f(c)}$ is greater than :

(A) 1 (B) $\frac{b-c}{c-a}$ (C) $\frac{b+a}{b-a}$ (D) $\frac{c-a}{b-c}$

Open solution

Q137. JEE Main 2020 (Online) 9th January Morning Slot

If $f(x) = \begin{cases} \frac{\sin(a+2)x+\sin x}{x}; & x<0 \\ b \quad ; & x=0 \\ \frac{\left(x+3x^2\right)^{\frac{1}{3}}-x^{\frac{1}{3}}}{x^{\frac{4}{3}}}; & x>0 \end{cases}$

is continuous at x = 0, then a + 2b is equal to :

(A) 0 (B) -1 (C) -2 (D) 1

Open solution

Q138. JEE Main 2020 (Online) 8th January Evening Slot

Let S be the set of all functions f : [0,1] $\to$ R, which are continuous on [0,1] and differentiable on (0,1). Then for every f in S, there exists a c $\in$ (0,1), depending on f, such that

(A) $|f(c)-f(1)| < |f'(c)|$ (B) $|f(c)+f(1)| < (1+c)\,|f'(c)|$

(C) $|f(c)-f(1)| < (1-c)\,|f'(c)|$ (D) None

Open solution

Q139. JEE Main 2020 (Online) 8th January Morning Slot

$\lim_{x\to 0}\left(\frac{3x^2+2}{7x^2+2}\right)^{\frac{1}{x^2}}$ is equal to

(A) e (B) e^2 (C) $\frac{1}{e^2}$ (D) $\frac{1}{e}$

Open solution

Q140. JEE Main 2019 (Online) 12th April Evening Slot

$\lim_{x\to 0}\frac{x+2\sin x}{\sqrt{x^2+2\sin x+1}-\sqrt{\sin^2 x-x+1}}$ is :

(A) 6 (B) 1 (C) 3 (D) 2

Open solution

Q141. JEE Main 2019 (Online) 12th April Evening Slot

Let f(x) = 5 − |x − 2| and g(x) = |x + 1|, x $\in$ R. If f(x) attains maximum value at α and g(x) attains minimum value at β, then $\lim_{x\to -\alpha\beta}\frac{(x-1)(x^2-5x+6)}{x^2-6x+8}$ is equal to :

(A) $\frac{1}{2}$ (B) $-\frac{1}{2}$ (C) $\frac{3}{2}$ (D) $-\frac{3}{2}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q142. JEE Main 2019 (Online) 12th April Morning Slot

If α and β are the roots of the equation $375x^2 - 25x - 2 = 0$, then $\lim\limits_{n\to\infty}\sum\limits_{r=1}^{n}\alpha^r + \lim\limits_{n\to\infty}\sum\limits_{r=1}^{n}\beta^r$ is equal to :

(A) $\frac{7}{116}$ (B) $\frac{29}{348}$ (C) $\frac{1}{12}$ (D) $\frac{21}{346}$

Open solution

Q143. JEE Main 2019 (Online) 10th April Evening Slot

If $\lim\limits_{x\to1}\frac{x^2-ax+b}{x-1} = 5$, then a + b is equal to :

(A) 1 (B) - 4 (C) - 7 (D) 5

Open solution

Q144. JEE Main 2019 (Online) 10th April Morning Slot

If $\lim\limits_{x\to1}\frac{x^4-1}{x-1} = \lim\limits_{x\to k}\frac{x^3-k^3}{x^2-k^2}$, then k is :

(A) $\frac{3}{2}$ (B) $\frac{8}{3}$ (C) $\frac{4}{3}$ (D) $\frac{3}{8}$

Open solution

Q145. JEE Main 2019 (Online) 10th April Morning Slot

If $f(x) = \begin{cases} \frac{\sin(p+1)x+\sin x}{x} & , x<0 \\ q & , x=0 \\ \frac{\sqrt{x+x^2}-\sqrt{x}}{x^{3/2}} & , x>0 \end{cases}$

is continuous at x = 0, then the ordered pair (p, q) is equal to

(A) $\left(-\frac{3}{2},-\frac{1}{2}\right)$ (B) $\left(-\frac{1}{2},\frac{3}{2}\right)$
(C) $\left(-\frac{3}{2},\frac{1}{2}\right)$ (D) $\left(\frac{5}{2},\frac{1}{2}\right)$

Open solution

Q146. JEE Main 2019 (Online) 10th April Morning Slot

Let f : R $\to$ R be differentiable at c $\in$ R and f(c) = 0. If g(x) = |f(x)| , then at x = c, g is :

(A) differentiable if f '(c) = 0 (B) differentiable if f '(c) $\neq$ 0
(C) not differentiable (D) not differentiable if f '(c) = 0

Open solution

Q147. JEE Main 2019 (Online) 9th April Evening Slot

If the function $f(x) = \begin{cases} a|\pi - x| + 1, & x \le 5 \\ b|x-\pi| + 3, & x > 5 \end{cases}$

is continuous at x = 5, then the value of a − b is :-

(A) $\frac{2}{\pi-5}$ (B) $\frac{2}{5-\pi}$ (C) $\frac{-2}{\pi+5}$ (D) $\frac{2}{\pi+5}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q148. JEE Main 2019 (Online) 9th April Evening Slot

If $f(x) = [x] - \left[\frac{x}{4}\right]$,x ∈ 4 , where [x] denotes the greatest integer function, then

(A) Both $\lim_{x\to 4-} f(x)$ and $\lim_{x\to 4+} f(x)$ exist but are not equal

(B) f is continuous at x = 4

(C) $\lim_{x\to 4+} f(x)$ exists but $\lim_{x\to 4-} f(x)$ does not exist

(D) $\lim_{x\to 4-} f(x)$ exists but $\lim_{x\to 4+} f(x)$ does not exist

Open solution

Q149. JEE Main 2019 (Online) 9th April Morning Slot

Let f(x) = 15 − |x − 10|; x ∈ R. Then the set of all values of x, at which the function, g(x) = f(f(x)) is not differentiable, is :

(A) {10,15} (B) {5,10,15,20} (C) {10} (D) {5,10,15}

Open solution

Q150. JEE Main 2019 (Online) 9th April Morning Slot

If the function f defined on , $\left(\frac{\pi}{6}, \frac{\pi}{3}\right)$ by

$$f(x) = \begin{cases} \frac{\sqrt{2}\cos x - 1}{\cot x - 1}, & x \neq \frac{\pi}{4} \\ k, & x = \frac{\pi}{4} \end{cases}$$

is continuous, then k is equal to

(A) 1 (B) 1 / $\sqrt{2}$ (C) $\frac{1}{2}$ (D) 2

Open solution

Q151. JEE Main 2019 (Online) 8th April Evening Slot

Let f : R → R be a differentiable function satisfying f'(3) + f'(2) = 0.

Then $\lim_{x\to 0} \left(\frac{1+f(3+x)-f(3)}{1+f(2-x)-f(2)}\right)^{\frac{1}{x}}$ is equal to

(A) e (B) e^2 (C) e^{-1} (D) 1

Open solution

Q152. JEE Main 2019 (Online) 8th April Evening Slot

Let f : [−1,3] → R be defined as

$$f(x) = \begin{cases} |x| + [x] & , \quad -1 \le x < 1 \\ x + |x| & , \quad 1 \le x < 2 \\ x + [x] & , \quad 2 \le x \le 3 \end{cases}$$

where [t] denotes the greatest integer less than or equal to t. Then, f is discontinuous at:

(A) only three points (B) four or more points

(C) only two points (D) only one point

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q153. JEE Main 2019 (Online) 8th April Morning Slot

$\lim_{x\to 0} \frac{\sin^2 x}{\sqrt{2}-\sqrt{1+\cos x}}$ equals:

(A) $\sqrt{2}$ (B) $2\sqrt{2}$ (C) 4 (D) $4\sqrt{2}$

Open solution

Q154. JEE Main 2019 (Online) 12th January Evening Slot

$\lim_{x\to 1^-} \frac{\sqrt{\pi}-\sqrt{2\sin^{-1}x}}{\sqrt{1-x}}$ is equal to :

(A) $\sqrt{\frac{2}{\pi}}$ (B) $\frac{1}{\sqrt{2\pi}}$ (C) $\sqrt{\frac{\pi}{2}}$ (D) $\sqrt{\pi}$

Open solution

Q155. JEE Main 2019 (Online) 12th January Evening Slot

Let f be a differentiable function such that f(1) = 2 and f '(x) = f(x) for all x $\in$ R R. If h(x) = f(f(x)), then h'(1) is equal to :

(A) 4e (B) $2e^2$ (C) $4e^2$ (D) 2e

Open solution

Q156. JEE Main 2019 (Online) 12th January Morning Slot

$\lim_{x\to \pi/4} \frac{\cot^3 x-\tan x}{\cos\left(x+\frac{\pi}{4}\right)}$ is :

(A) $8\sqrt{2}$ (B) 4 (C) $4\sqrt{2}$ (D) 8

Open solution

Q157. JEE Main 2019 (Online) 12th January Morning Slot

Let S be the set of all points in $(-\pi, \pi)$ at which the function, f(x) = min{sin x, cos x} is not differentiable. Then S is a subset of which of the following ?

(A) $\left\{-\frac{\pi}{2}, -\frac{\pi}{4}, \frac{\pi}{4}, \frac{\pi}{2}\right\}$

(B) $\left\{-\frac{3\pi}{4}, -\frac{\pi}{2}, \frac{\pi}{2}, \frac{3\pi}{4}\right\}$

(C) $\left\{-\frac{\pi}{4}, 0, \frac{\pi}{4}\right\}$

(D) $\left\{-\frac{3\pi}{4}, -\frac{\pi}{4}, \frac{3\pi}{4}, \frac{\pi}{4}\right\}$

Open solution

Q158. JEE Main 2019 (Online) 11th January Evening Slot

Let K be the set of all real values of x where the function f(x) = sin |x| − |x| + 2(x − π) cos |x| is not differentiable. Then the set K is equal to :

(A) {0, π} (B) ϕ (an empty set)

(C) { r } (D) {0}

Open solution

Q159. JEE Main 2019 (Online) 11th January Evening Slot

$\lim_{x\to 0} \frac{x\cot(4x)}{\sin^2 x\cot^2(2x)}$ is equal to :

(A) 0 (B) 4 (C) 1 (D) 2

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q160. JEE Main 2019 (Online) 11th January Morning Slot

Let $f(x) = \begin{cases} -1 & -2 \le x < 0 \\ x^2 - 1, & 0 \le x \le 2 \end{cases}$ and

$g(x) = |f(x)| + f(|x|)$.

Then, in the interval (−2, 2), g is :

(A) non continuous

(B) differentiable at all points

(C) not differentiable at two points

(D) not differentiable at one point

Open solution

Q161. JEE Main 2019 (Online) 11th January Morning Slot

Let [x] denote the greatest integer less than or equal to x. Then $\lim\limits_{x\to 0} \frac{\tan(\pi\sin^2 x) + (|x| - \sin([x]))^2}{x^2}$

(A) equals π + 1

(B) equals 0

(C) does not exist

(D) equals π

Open solution

Q162. JEE Main 2019 (Online) 10th January Evening Slot

Let f : $(-1, 1) \to$ R be a function defined by f(x) = max $\left\{-|x|, -\sqrt{1-x^2}\right\}$. If K be the set of all points at which f is not differentiable, then K has exactly -

(A) one element

(B) three elements

(C) five elements

(D) two elements

Open solution

Q163. JEE Main 2019 (Online) 10th January Morning Slot

For each t $\in$ R , let [t] be the greatest integer less than or equal to t

Then $\lim\limits_{x\to 1^+} \frac{(1-|x|+\sin|1-x|)\sin\left(\frac{\pi}{2}[1-x]\right)}{|1-x|.[1-x]}$

(A) equals $-$ 1

(B) equals 1

(C) equals 0

(D) does not exist

Open solution

Q164. JEE Main 2019 (Online) 10th January Morning Slot

Let $f(x) = \begin{cases} \max\{|x|, x^2\} & |x| \le 2 \\ 8 - 2|x| & 2 < |x| \le 4 \end{cases}$

Let S be the set of points in the interval (− 4, 4) at which f is not differentiable. Then S

(A) equals $\{-2, -1, 1, 2\}$

(B) equals $\{-2, -1, 0, 1, 2\}$

(C) equals $\{-2, 2\}$

(D) is an empty set

Open solution

Q165. JEE Main 2019 (Online) 9th January Evening Slot

For each x$\in$**R**, let [x] be the greatest integer less than or equal to x.

Then $\lim\limits_{x\to 0^-} \frac{x([x]+|x|)\sin[x]}{|x|}$ is equal to :

(A) $-$ sin 1

(B) 1

(C) sin 1

(D) 0

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q166. JEE Main 2019 (Online) 9th January Morning Slot

Open solution

Let f : R → R be a function defined as

$$f(x) = \begin{cases} 5 & ; \quad x \le 1 \\ a + bx & ; \quad 1 < x < 3 \\ b + 5x & ; \quad 3 \le x < 5 \\ 30 & ; \quad x \ge 5 \end{cases}$$

Then, f is

(A) continuous if a = 0 and b = 5

(B) continuous if a = −5 and b = 10

(C) continuous if a = 5 and b = 5

(D) not continuous for any values of a and b

Q167. JEE Main 2019 (Online) 9th January Morning Slot

Open solution

$\lim\limits_{y \to 0} \frac{\sqrt{1+\sqrt{1+y^4}}-\sqrt{2}}{y^4}$

(A) exists and equals $\frac{1}{2\sqrt{2}}$

(B) exists and equals $\frac{1}{4\sqrt{2}}$

(C) exists and equals $\frac{1}{2\sqrt{2}(1+\sqrt{2})}$

(D) does not exist

Q168. JEE Main 2018 (Online) 16th April Morning Slot

Open solution

$\lim\limits_{x \to 0} \frac{(27+x)^{\frac{1}{3}}-3}{9-(27+x)^{\frac{2}{3}}}$ equals.

(A) $\frac{1}{3}$ (B) $-\frac{1}{3}$ (C) $-\frac{1}{6}$ (D) $\frac{1}{6}$

Q169. JEE Main 2018 (Online) 16th April Morning Slot

Open solution

If the function f defined as

$f(x) = \frac{1}{x} - \frac{k-1}{e^{2x}-1}, x \neq 0$, is continuous at

x = 0, then the ordered pair (k, f(0)) is equal to :

(A) (3, 2) (B) (3, 1) (C) (2, 1) (D) $\left(\frac{1}{3}, 2\right)$

Q170. JEE Main 2018 (Offline)

Open solution

For each t ∈ R, let [t] be the greatest integer less than or equal to t.

Then $\lim\limits_{x \to 0^+} x\left(\left[\frac{1}{x}\right] + \left[\frac{2}{x}\right] + \ldots\ldots + \left[\frac{15}{x}\right]\right)$

(A) does not exist in R

(B) is equal to 0

(C) is equal to 15

(D) is equal to 120

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q171. JEE Main 2018 (Offline)

Let S = { t $\in R$: $f(x) = |x - \pi|.\left(e^{|x|} - 1\right)\sin|x|$ is not differentiable at t}, then the set S is equal to

(A) {0, π}
(B) ϕ (an empty set)
(C) {0}
(D) {π}

Open solution

Q172. JEE Main 2018 (Online) 15th April Evening Slot

$\lim\limits_{x\to 0} \frac{x\tan 2x - 2x\tan x}{(1-\cos 2x)^2}$ equals :

(A) $\frac{1}{4}$
(B) 1
(C) $\frac{1}{2}$
(D) $-\frac{1}{2}$

Open solution

Q173. JEE Main 2018 (Online) 15th April Evening Slot

Let f(x) = $\begin{cases} (x-1)^{\frac{1}{2-x}}, & x > 1, x \neq 2 \\ k & , x = 2 \end{cases}$

The value of k for which f is continuous at x = 2 is :

(A) 1
(B) e
(C) e^{-1}
(D) e^{-2}

Open solution

Q174. JEE Main 2018 (Online) 15th April Evening Slot

Let f(x) be a polynomial of degree 4 having extreme values at $x = 1$ and $x = 2$.

If $\lim\limits_{x\to 0}\left(\frac{f(x)}{x^2} + 1\right) = 3$ then f(−1) is equal to :

(A) $\frac{9}{2}$
(B) $\frac{5}{2}$
(C) $\frac{3}{2}$
(D) $\frac{1}{2}$

Open solution

Q175. JEE Main 2018 (Online) 15th April Morning Slot

Let S = {(λ, μ) $\in$ **R** $\times$ **R** : f(t) = ($|\lambda| e^{|t|} - \mu$). sin (2|t|), **t** $\in$ **R**, is a differentiable function}. Then S is a subset of :

(A) **R** $\times$ [0, ∞)
(B) [0, ∞) $\times$ **R**
(C) **R** $\times$ ($-\infty$, 0)
(D) ($-\infty$, 0) $\times$ **R**

Open solution

Q176. JEE Main 2017 (Online) 9th April Morning Slot

The value of k for which the function

$$f(x) = \begin{cases} \left(\frac{4}{5}\right)^{\frac{\tan 4x}{\tan 5x}}, & 0 < x < \frac{\pi}{2} \\ k + \frac{2}{5}, & x = \frac{\pi}{2} \end{cases}$$

is continuous at x = $\frac{\pi}{2}$, is :

(A) $\frac{17}{20}$
(B) $\frac{2}{5}$
(C) $\frac{3}{5}$
(D) $-\frac{2}{5}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q177. JEE Main 2017 (Online) 8th April Morning Slot

$\lim\limits_{x\to 3}\frac{\sqrt{3x}-3}{\sqrt{2x-4}-\sqrt{2}}$ is equal to :

(A) $\sqrt{3}$ (B) $\frac{1}{\sqrt{2}}$ (C) $\frac{\sqrt{3}}{2}$ (D) $\frac{1}{2\sqrt{2}}$

Open solution

Q178. JEE Main 2017 (Offline)

$\lim\limits_{x\to\frac{\pi}{2}}\frac{\cot x-\cos x}{(\pi-2x)^3}$ equals

(A) $\frac{1}{16}$ (B) $\frac{1}{8}$ (C) $\frac{1}{4}$ (D) $\frac{1}{24}$

Open solution

Q179. JEE Main 2016 (Online) 10th April Morning Slot

$\lim\limits_{x\to 0}\frac{(1-\cos 2x)^2}{2x\tan x-x\tan 2x}$ is :

(A) -2 (B) $-\frac{1}{2}$ (C) $\frac{1}{2}$ (D) 2

Open solution

Q180. JEE Main 2016 (Online) 10th April Morning Slot

Let a, b $\in$ **R**, (a $\neq$ 0). If the function *f* defined as

$$f(x)=\begin{cases}\frac{2x^2}{a}, & 0\le x<1\\ a, & 1\le x<\sqrt{2}\\ \frac{2b^2-4b}{x^3}, & \sqrt{2}\le x<\infty\end{cases}$$

is continuous in the interval [0, ∞), then an ordered pair (a, b) is :

(A) $\left(\sqrt{2},1-\sqrt{3}\right)$ (B) $\left(-\sqrt{2},1+\sqrt{3}\right)$

(C) $\left(\sqrt{2},-1+\sqrt{3}\right)$ (D) $\left(-\sqrt{2},1-\sqrt{3}\right)$

Open solution

Q181. JEE Main 2016 (Online) 9th April Morning Slot

If $\lim\limits_{x\to\infty}\left(1+\frac{a}{x}-\frac{4}{x^2}\right)^{2x}=e^3$, then 'a' is equal to :

(A) 2 (B) $\frac{3}{2}$ (C) $\frac{2}{3}$ (D) $\frac{1}{2}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q182. JEE Main 2016 (Online) 9th April Morning Slot

If the function

$$f(x) = \begin{cases} -x & x < 1 \\ a + \cos^{-1}(x+b), & 1 \le x \le 2 \end{cases}$$

is differentiable at x = 1, then $\frac{a}{b}$ is equal to :

(A) $\frac{\pi-2}{2}$ (B) $\frac{-\pi-2}{2}$

(C) $\frac{\pi+2}{2}$ (D) $-1-\cos^{-1}(2)$

Open solution

Q183. JEE Main 2016 (Offline)

For $x \in R,\ f(x) = |\log 2 - \sin x|$

and $g(x) = f(f(x)),$ then :

(A) g is not differentiable at $x = 0$

(B) $g'(0) = \cos(\log 2)$

(C) $g'(0) = -\cos(\log 2)$

(D) g is differentiable at $x = 0$ and $g'(0) = -\sin(\log 2)$

Open solution

Q184. JEE Main 2016 (Offline)

Let $p = \lim\limits_{x\to 0^+}\left(1+\tan^2\sqrt{x}\right)^{\frac{1}{2x}}$ then $\log p$ is equal to :

(A) $\frac{1}{2}$ (B) $\frac{1}{4}$ (C) 2 (D) 1

Open solution

Q185. JEE Main 2015 (Online) 11th April Morning Slot

Let k be a non-zero real number. If

$$f(x) = \begin{cases} \frac{(e^x-1)^2}{\sin\left(\frac{x}{k}\right)\log\left(1+\frac{x}{4}\right)}, & x \neq 0 \\ 12\ , & x = 0 \end{cases}$$

is a continuous function, then the value of k is :

(A) 1 (B) 2 (C) 3 (D) 4

Open solution

Q186. JEE Main 2015 (Online) 10th April Morning Slot

$\lim\limits_{x\to 0}\frac{e^{x^2}-\cos x}{\sin^2 x}$ is equal to :

(A) 3 (B) $\frac{3}{2}$ (C) $\frac{5}{4}$ (D) 2

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q187. JEE Main 2015 (Offline)

$\lim\limits_{x\to 0}\frac{(1-\cos 2x)(3+\cos x)}{x\tan 4x}$ is equal to

(A) 2 (B) $\frac{1}{2}$ (C) 4 (D) 3

Open solution

Q188. JEE Main 2015 (Offline)

If the function.

$$g(x)=\begin{cases}k\sqrt{x+1}, & 0\le x\le 3\\ mx+2, & 3<x\le 5\end{cases}$$

is differentiable, then the value of $k+m$ is :

(A) $\frac{10}{3}$ (B) 4 (C) 2 (D) $\frac{16}{5}$

Open solution

Q189. JEE Main 2014 (Online) 19th April Morning Slot

If the function

$$f(x)=\begin{cases}\frac{\sqrt{2+\cos x}-1}{(\pi-x)^2}, & x\neq\pi\\ k & ,x=\pi\end{cases}$$

is continuous at x = π, then k equals :

(A) 0 (B) $\frac{1}{2}$ (C) 2 (D) $\frac{1}{4}$

Open solution

Q190. JEE Main 2014 (Online) 19th April Morning Slot

Let f : **R** $\to$ **R** be a function such that $|f(x)|\le x^2$, for all x $\in$ **R**.
Then, at x = 0, f is :

(A) continuous but not differentiable (B) continuous as well as differentiable

(C) neither continuous nor differentiable (D) differentiable but not continuous.

Open solution

Q191. JEE Main 2014 (Online) 12th April Morning Slot

Let f, g : R $\to$ R be two functions defined by

$$f(x)=\begin{cases}x\sin\left(\frac{1}{x}\right), & x\neq 0\\ 0 & ,x=0\end{cases}\text{, and } g(x)=xf(x)$$

Statement I : f is a continuous function at x = 0,
Statement II : g is a differentiable function at x = 0.

(A) Both statements I and II are false.

(B) Both statements I and II are true.

(C) Statement I is true, statement II is false.

(D) Statement I is false, statement II is true .

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q192. JEE Main 2014 (Online) 11th April Morning Slot

If $\lim_{x\to 2} \frac{\tan(x-2)\{x^2+(k-2)x-2k\}}{x^2-4x+4} = 5$,
then k is equal to :

(A) 0 (B) 1 (C) 2 (D) 3

Open solution

Q193. JEE Main 2014 (Online) 11th April Morning Slot

Let f(x) = x $|x|$, g(x) = sin x and h(x) = (gof)(x). Then

(A) h(x) is not differentiable at x = 0.

(B) h(x) is differentiable at x = 0, but h'(x) is not continuous at x = 0.

(C) h'(x) is continuous at x = 0 but it is not differentiable at x = 0.

(D) h'(x) is differentiable at x = 0.

Open solution

Q194. JEE Main 2014 (Online) 9th April Morning Slot

If f(x) is continuous and $f\left(\frac{9}{2}\right) = \frac{2}{9}$, then $\lim_{x\to 0} f\left(\frac{1-\cos 3x}{x^2}\right)$ is equal to :

(A) 9/2 (B) 2/9 (C) 0 (D) 8/9

Open solution

Q195. JEE Main 2014 (Offline)

$\lim_{x\to 0} \frac{\sin(\pi\cos^2 x)}{x^2}$ is equal to :

(A) $-\pi$ (B) π (C) $\frac{\pi}{2}$ (D) 1

Open solution

Q196. JEE Main 2013 (Online) 25th April Morning Slot

Consider the function :
f(x) = [x] + $|1-x|$, -1 $\leq$ x $\leq$ 3 where [x] is the greatest integer function.

Statement 1: f is not continuous at x = 0, 1, 2 and 3.

Statement 2: f(x) = $\begin{cases} -x, & -1 \le x < 0 \\ 1-x & 0 \le x < 1 \\ 1+x & 1 \le x < 2 \\ 2+x & 2 \le x \le 3 \end{cases}$

(A) Statement 1 is true; Statement 2 is false

(B) Statement 1 is true; Statement 2 is true; Statement 2 is not correct explanation of Statement 1

(C) Statement 1 is true; Statement 2 is true; Statement 2 is correct explanation of Statement 1

(D) Statement 1 is false; Statement 2 is true

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q197. JEE Main 2013 (Online) 23th April Morning Slot

Let f be a composite function of x defined by
$f(u) = \frac{1}{u^2+u-2}, u(x) = \frac{1}{x-1}$
Then the number of points x where f is discontinuous is:

(A) 4 (B) 3 (C) 2 (D) 1

Open solution

Q198. JEE Main 2013 (Online) 22th April Morning Slot

Let $f(x) = -1 + |x-2|$, and $g(x) = 1 - |x|$; then the set of all points where fog is discontinuous is:

(A) {0, 2} (B) {0, 1, 2} (C) {0} (D) an empty set

Open solution

Q199. JEE Main 2013 (Online) 9th April Morning Slot

The value of $\lim_{x\to 0} \frac{1}{x}\left[\tan^{-1}\left(\frac{x+1}{2x+1}\right) - \frac{\pi}{4}\right]$ is :

(A) 1 (B) $-\frac{1}{2}$ (C) 2 (D) 0

Open solution

Q200. JEE Main 2013 (Offline)

$\lim\limits_{x\to 0} \frac{(1-\cos 2x)(3+\cos x)}{x \tan 4x}$ is equal to

(A) $-\frac{1}{4}$ (B) $\frac{1}{2}$ (C) 1 (D) 2

Open solution

Q201. AIEEE 2012 (Online) 26th May

$\lim_{x\to 0} \frac{\sin(\pi \cos^2 x)}{x^2}$ equals

(A) $-\pi$ (B) 1 (C) -1 (D) π

Open solution

Q202. AIEEE 2012 (Online) 26th May

If $f(x) = a|\sin x| + be^{|x|} + c|x|^3$, where $a, b, c \in R$, is differentiable at $x = 0$, then

(A) $a = 0$, b and c are any real numbers (B) $c = 0$, $a = 0$, b is any real number

(C) $b = 0$, $c = 0$, a is any real number (D) $a = 0$, $b = 0$, c is any real number

Open solution

Q203. AIEEE 2012 (Online) 19th May

If $f(x) = 3x^{10} - 7x^8 + 5x^6 - 21x^3 + 3x^2 - 7$, then $\lim\limits_{\alpha\to 0} \frac{f(1-\alpha)-f(1)}{\alpha^3+3\alpha}$ is

(A) $-\frac{53}{3}$ (B) $\frac{53}{3}$ (C) $-\frac{55}{3}$ (D) $\frac{55}{3}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q204. AIEEE 2012 (Online) 19th May

Let $f:[1,3]\to R$ be a function satisfying $\frac{x}{[x]}\le f(x)\le\sqrt{6-x}$, for all $x\neq 2$ and $f(2)=1$, where R is the set of all real numbers and $[x]$ denotes the largest integer less than or equal to x.
Statement 1: $\lim\limits_{x\to 2^-} f(x)$ exists.
Statement 2: f is continuous at $x=2$.

(A) Statement 1 is true, Statement 2 is true, Statement 2 is a correct explanation for Statement 1.
(B) Statement 1 is false, Statement 2 is true.
(C) Statement 1 is true, Statement 2 is true, Statement 2 is not a correct explanation for Statement 1.
(D) Statement 1 is true, Statement 2 is false.

Open solution

Q205. AIEEE 2012 (Online) 12th May

Statement 1: A function $f:R\to R$ is continuous at x_0 if and only if $\lim_{x\to x_0} f(x)$ exists and $\lim_{x\to x_0} f(x)=f(x_0)$.
Statement 2: A function $f:R\to R$ is discontinuous at x_0 if and only if, $\lim_{x\to x_0} f(x)$ exists and $\lim_{x\to x_0} f(x)\neq f(x_0)$.

(A) Statement 1 is true, Statement 2 is true; Statement 2 is not a correct explanation of Statement 1.
(B) Statement 1 is false, Statement 2 is true.
(C) Statement 1 is true, Statement 2 is true; Statement 2 is a correct explanation of Statement 1.
(D) Statement 1 is true, Statement 2 is false.

Open solution

Q206. AIEEE 2012 (Online) 7th May

$\lim_{x\to 0}\left(\frac{x-\sin x}{x}\right)\sin\left(\frac{1}{x}\right)$

(A) equals 1 (B) equals 0 (C) does not exist (D) equals -1

Open solution

Q207. AIEEE 2012 (Online) 7th May

If $x+|y|=2y$, then y as a function of x, at $x=0$ is

(A) differentiable but not continuous
(B) continuous but not differentiable
(C) continuous as well as differentiable
(D) neither continuous nor differentiable

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q208. AIEEE 2012

If $f : R \to R$ is a function defined by

$f(x) = [x] \cos\left(\frac{2x-1}{2}\right)\pi$,

where [x] denotes the greatest integer function, then f is

(A) continuous for every real x

(B) discontinuous only at $x = 0$

(C) discontinuous only at non-zero integral values of x

(D) continuous only at $x = 0$

Open solution

Q209. AIEEE 2012

Consider the function, $f(x) = |x-2| + |x-5|, x \in R$

Statement - 1 : $f'(4) = 0$

Statement - 2 : f is continuous in [2, 5], differentiable in (2, 5) and $f(2) = f(5)$

(A) Statement - 1 is false, statement - 2 is true

(B) Statement - 1 is true, statement - 2 is true; statement - 2 is a correct explanation for statement - 1

(C) Statement - 1 is true, statement - 2 is true; statement - 2 is not a correct explanation for statement - 1

(D) Statement - 1 is true, statement - 2 is false

Open solution

Q210. AIEEE 2011

$\lim_{x\to 2}\left(\frac{\sqrt{1-\cos\{2(x-2)\}}}{x-2}\right)$

(A) Equals $\sqrt{2}$

(B) Equals $-\sqrt{2}$

(C) Equals $\frac{1}{\sqrt{2}}$

(D) does not exist

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q211. AIEEE 2011

The value of p and q for which the function

$$f(x)=\begin{cases}\frac{\sin(p+1)x+\sin x}{x} & ,x<0\\ q & ,x=0\\ \frac{\sqrt{x+x^2}-\sqrt{x}}{x^{3/2}} & ,x>0\end{cases}$$

is continuous for all x in R, are

(A) $p=\frac{5}{2}, q=\frac{1}{2}$

(B) $p=-\frac{3}{2}, q=\frac{1}{2}$

(C) $p=\frac{1}{2}, q=\frac{3}{2}$

(D) $p=\frac{1}{2}, q=-\frac{3}{2}$

Open solution

Q212. AIEEE 2010

Let $f: R \to R$ be a positive increasing function with

$\lim_{x\to\infty}\frac{f(3x)}{f(x)}=1$. Then $\lim_{x\to\infty}\frac{f(2x)}{f(x)}=$

(A) $\frac{2}{3}$ (B) $\frac{3}{2}$ (C) 3 (D) 1

Open solution

Q213. AIEEE 2009

Let $f(x)=x|x|$ and $g(x)=\sin x$.

Statement-1: gof is differentiable at $x=0$ and its derivative is continuous at that point.

Statement-2: gof is twice differentiable at $x=0$.

(A) Statement-1 is true, Statement-2 is true; Statement-2 is not a correct explanation for Statement-1.

(B) Statement-1 is true, Statement-2 is false

(C) Statement-1 is false, Statement-2 is true

(D) Statement-1 is true, Statement-2 is true Statement-2 is a correct explanation for Statement-1

Open solution

Q214. AIEEE 2008

Let $f(x)=\begin{cases}(x-1)\sin\frac{1}{x-1} & if\ x\neq 1\\ 0 & if\ x=1\end{cases}$

Then which one of the following is true?

(A) f is neither differentiable at x = 0 nor at x = 1

(B) f is differentiable at x = 0 and at x = 1

(C) f is differentiable at x = 0 but not at x = 1

(D) f is differentiable at x = 1 but not at x = 0

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q215. AIEEE 2007

Let $f : R \to R$ be a function defined by

$f(x) = \min\{x+1, |x|+1\}$, then which of the following is true?

Open solution

(A) $f(x)$ is differentiale everywhere

(B) $f(x)$ is not differentiable at x = 0

(C) $f(x) > 1$ for all $x \in R$

(D) $f(x)$ is not differentiable at x = 1

Q216. AIEEE 2007

The function $f : R/\{0\} \to R$ given by

$f(x) = \frac{1}{x} - \frac{2}{e^{2x}-1}$

can be made continuous at x = 0 by defining $f(0)$ as

Open solution

(A) 0

(B) 1

(C) 2

(D) -1

Q217. AIEEE 2006

The set of points where $f(x) = \frac{x}{1+|x|}$ is differentiable is

Open solution

(A) $(-\infty, 0) \cup (0, \infty)$

(B) $(-\infty, -1) \cup (-1, \infty)$

(C) $(-\infty, \infty)$

(D) $(0, \infty)$

Q218. AIEEE 2005

Let α and β be the distinct roots of $ax^2 + bx + c = 0$, then

$\lim_{x\to\alpha} \frac{1-\cos(ax^2+bx+c)}{(x-\alpha)^2}$ is equal to

Open solution

(A) $\frac{a^2(\alpha-\beta)^2}{2}$

(B) 0

(C) $-\frac{a^2(\alpha-\beta)^2}{2}$

(D) $\frac{(\alpha-\beta)^2}{2}$

Q219. AIEEE 2005

Suppose $f(x)$ is differentiable at x = 1 and

$\lim_{h\to 0} \frac{1}{h} f(1+h) = 5$, then $f'(1)$ equals

Open solution

(A) 3

(B) 4

(C) 5

(D) 6

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q220. AIEEE 2005

If f is a real valued differentiable function satisfying

$|f(x) - f(y)| \leq (x-y)^2, x, y \in R$
and $f(0) = 0$, then $f(1)$ equals

(A) -1 (B) 0 (C) 2 (D) 1

Open solution

Q221. AIEEE 2004

If $\lim_{x\to\infty} \left(1 + \frac{a}{x} + \frac{b}{x^2}\right)^{2x} = e^2$, then the value of a and b, are

(A) a = 1 and b = 2 (B) a = 1 and $b \in R$ (C) $a \in R$ and b = 2 (D) $a \in R$ and $b \in R$

Open solution

Q222. AIEEE 2004

Let $f(x) = \frac{1-\tan x}{4x-\pi}$, $x \neq \frac{\pi}{4}$, $x \in \left[0, \frac{\pi}{2}\right]$.

If $f(x)$ is continuous in $\left[0, \frac{\pi}{2}\right]$, then $f\left(\frac{\pi}{4}\right)$ is

(A) -1 (B) $\frac{1}{2}$ (C) $-\frac{1}{2}$ (D) 1

Open solution

Q223. AIEEE 2003

If $\lim_{x\to 0} \frac{\log(3+x)-\log(3-x)}{x}$ = k, the value of k is

(A) $-\frac{2}{3}$ (B) 0 (C) $-\frac{1}{3}$ (D) $\frac{2}{3}$

Open solution

Q224. AIEEE 2003

Let $f(a) = g(a) = k$ and their nth derivatives
$f^n(a), g^n(a)$ exist and are not equal for some n. Further if

$\lim_{x\to a} \frac{f(a)g(x)-f(a)-g(a)f(x)+f(a)}{g(x)-f(x)} = 4$

then the value of k is

(A) 0 (B) 4 (C) 2 (D) 1

Open solution

Q225. AIEEE 2003

$\lim_{x\to\frac{\pi}{2}} \frac{\left[1-\tan\left(\frac{x}{2}\right)\right][1-\sin x]}{\left[1+\tan\left(\frac{x}{2}\right)\right][\pi-2x]^3}$ is

(A) ∞ (B) $\frac{1}{8}$ (C) 0 (D) $\frac{1}{32}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q226. AIEEE 2003

If $f(x) = \begin{cases} xe^{-\left(\frac{1}{|x|}+\frac{1}{x}\right)} & , x \neq 0 \\ 0 & , x = 0 \end{cases}$

then $f(x)$ is

(A) discontinuous everywhere

(B) continuous as well as differentiable for all x

(C) continuous for all x but not differentiable at x = 0

(D) neither differentiable nor continuous at x = 0

Open solution

Q227. AIEEE 2002

$\lim\limits_{x\to 0} \frac{\sqrt{1-\cos 2x}}{\sqrt{2}x}$ is

(A) 1 (B) -1 (C) zero (D) does not exist

Open solution

Q228. AIEEE 2002

$\lim\limits_{x\to\infty} \left(\frac{x^2+5x+3}{x^2+x+2}\right)^x$

(A) e^4 (B) e^2 (C) e^3 (D) 1

Open solution

Q229. AIEEE 2002

Let $f(2) = 4$ and $f'(x) = 4$.

Then $\lim\limits_{x\to 2} \frac{xf(2)-2f(x)}{x-2}$ is given by

(A) 2 (B) -2 (C) -4 (D) 3

Open solution

Q230. AIEEE 2002

$\lim\limits_{x\to 0} \frac{\log x^n - [x]}{[x]}$, $n \in N$, ([x] denotes the greatest integer less than or equal to x)

(A) has value -1 (B) has value 0 (C) has value 1 (D) does not exist

Open solution

Q231. AIEEE 2002

If $f(1) = 1, f'(1) = 2$, then

$\lim\limits_{x\to 1} \frac{\sqrt{f(x)}-1}{\sqrt{x}-1}$ is

(A) 2 (B) 4 (C) 1 (D) $\frac{1}{2}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q232. AIEEE 2002

f is defined in $[-5, 5]$ as

$f(x) = x$ if x is rational

$= -x$ if x is irrational. Then

(A) $f(x)$ is continuous at every x, except $x = 0$

(B) $f(x)$ is discontinuous at every x, except $x = 0$

(C) $f(x)$ is continuous everywhere

(D) $f(x)$ is discontinuous everywhere

Open solution

Q233. AIEEE 2002

f(x) and g(x) are two differentiable functions on [0, 2] such that

f"(x) - g"(x) = 0, f'(1) = 2, g'(1) = 4, f(2) = 3, g(2) = 9

then f(x) - g(x) at x = $\frac{3}{2}$ is

(A) 0 (B) 2 (C) 10 (D) -5

Open solution

Q234. AIEEE 2002

If f(x + y) = f(x).f(y) $\forall$ x, y and f(5) = 2, f'(0) = 3, then f'(5) is

(A) 0 (B) 1 (C) 6 (D) 2

Open solution

Numerical 62 questions

Q235. JEE Main 2026 (Online) 6th April Evening Shift

Let $f(x) = \begin{cases} x^3 + 8; & x < 0, \\ x^2 - 4; & x \geq 0, \end{cases}$ and $g(x) = \begin{cases} (x-8)^{1/3}; & x < 0, \\ (x+4)^{1/2}; & x \geq 0. \end{cases}$

Then the number of points, where the function $g \circ f$ is discontinuous, is ____ .

Open solution

Q236. JEE Main 2026 (Online) 4th April Evening Shift

Let $f(x) = \begin{cases} e^{x-1} & , x < 0 \\ x^2 - 5x + 6 & , x \geq 0 \end{cases}$ and $g(x) = f(|x|) + |f(x)|$. If the number of points where g is not continuous and is not differentiable are α and β respectively, then $\alpha + \beta$ is equal to ____

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q237. JEE Main 2026 (Online) 4th April Morning Shift

The number of points, at which the function $f(x) = \max\{6x, 2+3x^2\} + |x-1|\cos|x^2 - \frac{1}{4}|, x \in (-\pi, \pi)$, is not differentiable, is ____ .

Open solution

Q238. JEE Main 2026 (Online) 2nd April Evening Shift

The number of points in the interval $[2, 4]$, at which the function $f(x) = \left[x^2 - x - \frac{1}{2}\right]$, where $[\cdot]$ denotes the greatest integer function, is discontinuous, is _____.

Open solution

Q239. JEE Main 2025 (Online) 7th April Evening Shift

If the function $f(x) = \frac{\tan(\tan x) - \sin(\sin x)}{\tan x - \sin x}$ is continuous at $x = 0$, then $f(0)$ is equal to _____.

Open solution

Q240. JEE Main 2025 (Online) 7th April Evening Shift

For $t > -1$, let α_t and β_t be the roots of the equation

$\left((t+2)^{1/7} - 1\right)x^2 + \left((t+2)^{1/6} - 1\right)x + \left((t+2)^{1/21} - 1\right) = 0$

If $\lim\limits_{t \to -1^+} \alpha_t = a$ and $\lim\limits_{t \to -1^+} \beta_t = b$, then $72(a+b)^2$ is equal to _____.

Open solution

Q241. JEE Main 2025 (Online) 7th April Morning Shift

The number of points of discontinuity of the function $f(x) = \left[\frac{x^2}{2}\right] - [\sqrt{x}], x \in [0, 4]$, where $[\cdot]$ denotes the greatest integer function, is _____.

Open solution

Q242. JEE Main 2025 (Online) 4th April Morning Shift

Let m and n be the number of points at which the function $f(x) = \max\{x, x^3, x^5, \ldots x^{21}\}, x \in \mathbb{R}$, is not differentiable and not continuous, respectively. Then $m + n$ is equal to _____.

Open solution

Q243. JEE Main 2025 (Online) 3rd April Evening Shift

If $\lim\limits_{x \to 0} \left(\frac{\tan x}{x}\right)^{\frac{1}{x^2}} = p$, then $96 \log_e p$ is equal to _____

Open solution

Q244. JEE Main 2025 (Online) 29th January Morning Shift

Let [t] be the greatest integer less than or equal to t. Then the least value of p ∈ N for which $\lim\limits_{x \to 0^+} \left(x\left(\left[\frac{1}{x}\right] + \left[\frac{2}{x}\right] + \ldots + \left[\frac{p}{x}\right]\right) - x^2\left(\left[\frac{1}{x^2}\right] + \left[\frac{2^2}{x^2}\right] + \ldots + \left[\frac{9^2}{x^2}\right]\right)\right) \geq 1$ is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q245. JEE Main 2025 (Online) 28th January Evening Shift

Let $f(x)=\lim\limits_{n\to\infty}\sum\limits_{r=0}^{n}\left(\frac{\tan\left(x/2^{r+1}\right)+\tan^3\left(x/2^{r+1}\right)}{1-\tan^2\left(x/2^{r+1}\right)}\right)$. Then $\lim\limits_{x\to 0}\frac{e^x-e^{f(x)}}{(x-f(x))}$ is equal to _____.

Open solution

Q246. JEE Main 2025 (Online) 28th January Morning Shift

Let $f(x)=\begin{cases}3x, & x<0\\ \min\{1+x+[x],x+2[x]\}, & 0\le x\le 2\\ 5, & x>2\end{cases}$

where [.] denotes greatest integer function. If α and β are the number of points, where f is not continuous and is not differentiable, respectively, then $\alpha+\beta$ equals _____ .

Open solution

Q247. JEE Main 2025 (Online) 22nd January Morning Shift

Let the function,

$f(x)=\begin{cases}-3ax^2-2, & x<1\\ a^2+bx, & x\geqslant 1\end{cases}$

be differentiable for all $x\in\mathbf{R}$, where $a>1,\ b\in\mathbf{R}$. If the area of the region enclosed by $y=f(x)$ and the line $y=-20$ is $\alpha+\beta\sqrt{3},\alpha,\beta\in Z$, then the value of $\alpha+\beta$ is _____ .

Open solution

Q248. JEE Main 2024 (Online) 9th April Morning Shift

Let $f:(0,\pi)\to\mathbf{R}$ be a function given by $f(x)=\begin{cases}\left(\frac{8}{7}\right)^{\frac{\tan 8x}{\tan 7x}}, & 0<x<\frac{\pi}{2}\\ a-8, & x=\frac{\pi}{2}\\ (1+|\cot x|)^{\frac{b}{a}|\tan x|}, & \frac{\pi}{2}<x<\pi\end{cases}$

where $a,b\in\mathbf{Z}$. If f is continuous at $x=\frac{\pi}{2}$, then a^2+b^2 is equal to _____.

Open solution

Q249. JEE Main 2024 (Online) 8th April Evening Shift

If $\alpha=\lim\limits_{x\to 0^+}\left(\frac{e^{\sqrt{\tan x}}-e^{\sqrt{x}}}{\sqrt{\tan x}-\sqrt{x}}\right)$ and $\beta=\lim\limits_{x\to 0}(1+\sin x)^{\frac{1}{2}\cot x}$ are the roots of the quadratic equation $ax^2+bx-\sqrt{e}=0$, then $12\log_e(a+b)$ is equal to _____.

Open solution

Q250. JEE Main 2024 (Online) 8th April Morning Shift

The value of $\lim\limits_{x\to 0}2\left(\frac{1-\cos x\sqrt{\cos 2x}\sqrt[3]{\cos 3x}.\ldots..\sqrt[10]{\cos 10x}}{x^2}\right)$ is _____.

Open solution

Q251. JEE Main 2024 (Online) 6th April Evening Shift

Let $[t]$ denote the greatest integer less than or equal to t. Let $f:[0,\infty)\to\mathbf{R}$ be a function defined by $f(x)=\left[\frac{x}{2}+3\right]-[\sqrt{x}]$. Let S be the set of all points in the interval $[0,8]$ at which f is not continuous. Then $\sum\limits_{a\in S}a$ is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q252. JEE Main 2024 (Online) 5th April Evening Shift

Let $a > 0$ be a root of the equation $2x^2 + x - 2 = 0$. If $\lim\limits_{x \to \frac{1}{a}} \frac{16\left(1-\cos\left(2+x-2x^2\right)\right)}{(1-ax)^2} = \alpha + \beta\sqrt{17}$, where $\alpha, \beta \in Z$, then $\alpha + \beta$ is equal to _____.

Open solution

Q253. JEE Main 2024 (Online) 5th April Morning Shift

Let f be a differentiable function in the interval $(0, \infty)$ such that $f(1) = 1$ and $\lim\limits_{t \to x} \frac{t^2 f(x) - x^2 f(t)}{t - x} = 1$ for each $x > 0$. Then $2f(2) + 3f(3)$ is equal to _____.

Open solution

Q254. JEE Main 2024 (Online) 4th April Morning Shift

If $\lim\limits_{x \to 1} \frac{(5x+1)^{1/3} - (x+5)^{1/3}}{(2x+3)^{1/2} - (x+4)^{1/2}} = \frac{\mathrm{m}\sqrt{5}}{\mathrm{n}(2\mathrm{n})^{2/3}}$, where $\gcd(\mathrm{m}, \mathrm{n}) = 1$, then $8\,\mathrm{m} + 12\mathrm{n}$ is equal to _____.

Open solution

Q255. JEE Main 2024 (Online) 1st February Morning Shift

Let $\{x\}$ denote the fractional part of x and $f(x) = \frac{\cos^{-1}\left(1-\{x\}^2\right)\sin^{-1}(1-\{x\})}{\{x\}-\{x\}^3}$, $x \neq 0$. If L and R respectively denotes the left hand limit and the right hand limit of $f(x)$ at $x = 0$, then $\frac{32}{\pi^2}\left(\mathrm{L}^2 + \mathrm{R}^2\right)$ is equal to _____.

Open solution

Q256. JEE Main 2024 (Online) 31st January Evening Shift

If $\lim\limits_{x \to 0} \frac{ax^2 e^x - b\log_e(1+x) + cxe^{-x}}{x^2 \sin x} = 1$, then $16\left(a^2 + b^2 + c^2\right)$ is equal to _____.

Open solution

Q257. JEE Main 2024 (Online) 30th January Morning Shift

If the function

$$f(x) = \begin{cases} \frac{1}{|x|}, & |x| \geqslant 2 \\ \mathrm{a}x^2 + 2\,\mathrm{b}, & |x| < 2 \end{cases}$$

is differentiable on $\mathbf{R}$, then $48(a + b)$ is equal to _____.

Open solution

Q258. JEE Main 2024 (Online) 29th January Evening Shift

Let $f(x) = \sqrt{\lim\limits_{r \to x} \left\{ \frac{2r^2[(f(r))^2 - f(x)f(r)]}{r^2 - x^2} - r^3 e^{\frac{f(r)}{r}} \right\}}$ be differentiable in $(-\infty, 0) \cup (0, \infty)$ and $f(1) = 1$. Then the value of ea, such that $f(a) = 0$, is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q259. JEE Main 2023 (Online) 12th April Morning Shift

Let $[x]$ be the greatest integer $\leq x$. Then the number of points in the interval $(-2, 1)$, where the function $f(x) = |[x]| + \sqrt{x - [x]}$ is discontinuous, is _____.

Open solution

Q260. JEE Main 2023 (Online) 10th April Morning Shift

Let $f : (-2, 2) \to R$ be defined by $f(x) = \begin{cases} x[x], & -2 < x < 0 \\ (x-1)[x], & 0 \leq x < 2 \end{cases}$ where $[x]$ denotes the greatest integer function. If m and n respectively are the number of points in $(-2, 2)$ at which $y = |f(x)|$ is not continuous and not differentiable, then $m + n$ is equal to _____.

Open solution

Q261. JEE Main 2023 (Online) 8th April Evening Shift

Let k and m be positive real numbers such that the function $f(x) = \begin{cases} 3x^2 + k\sqrt{x+1}, & 0 < x < 1 \\ mx^2 + k^2, & x \geq 1 \end{cases}$ is differentiable for all $x > 0$. Then $\frac{8f'(8)}{f'(\frac{1}{8})}$ is equal to _____.

Open solution

Q262. JEE Main 2023 (Online) 6th April Morning Shift

Let $a \in \mathbb{Z}$ and [t] be the greatest integer $\leq$ t. Then the number of points, where the function $f(x) = [a + 13\sin x], x \in (0, \pi)$ is not differentiable, is _____.

Open solution

Q263. JEE Main 2022 (Online) 29th July Evening Shift

If $[t]$ denotes the greatest integer $\leq t$, then the number of points, at which the function $f(x) = 4|2x + 3| + 9\left[x + \frac{1}{2}\right] - 12[x + 20]$ is not differentiable in the open interval $(-20, 20)$, is _____.

Open solution

Q264. JEE Main 2022 (Online) 28th July Morning Shift

Let $f : [0, 1] \to \mathbf{R}$ be a twice differentiable function in $(0, 1)$ such that $f(0) = 3$ and $f(1) = 5$. If the line $y = 2x + 3$ intersects the graph of f at only two distinct points in $(0, 1)$, then the least number of points $x \in (0, 1)$, at which $f''(x) = 0$, is _____.

Open solution

Q265. JEE Main 2022 (Online) 28th July Morning Shift

$\lim\limits_{x \to 0} \left(\frac{(x+2\cos x)^3 + 2(x+2\cos x)^2 + 3\sin(x+2\cos x)}{(x+2)^3 + 2(x+2)^2 + 3\sin(x+2)} \right)^{\frac{100}{x}}$ is equal to _____.

Open solution

Q266. JEE Main 2022 (Online) 25th July Morning Shift

Let $f(x) = \begin{cases} |4x^2 - 8x + 5|, & \text{if } 8x^2 - 6x + 1 \geqslant 0 \\ [4x^2 - 8x + 5], & \text{if } 8x^2 - 6x + 1 < 0, \end{cases}$ where $[\alpha]$ denotes the greatest integer less than or equal to α. Then the number of points in $\mathbf{R}$ where f is not differentiable is _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q267. JEE Main 2022 (Online) 30th June Morning Shift

Suppose $\lim\limits_{x\to 0}\frac{F(x)}{x^3}$ exists and is equal to L, where

$$F(x)=\begin{vmatrix} a+\sin\frac{x}{2} & -b\cos x & 0 \\ -b\cos x & 0 & a+\sin\frac{x}{2} \\ 0 & a+\sin\frac{x}{2} & -b\cos x \end{vmatrix}.$$

Then, -112 L is equal to _____.

Open solution

Q268. JEE Main 2022 (Online) 28th June Evening Shift

If $\lim\limits_{x\to 1}\frac{\sin(3x^2-4x+1)-x^2+1}{2x^3-7x^2+ax+b}=-2$, then the value of (a − b) is equal to _____.

Open solution

Q269. JEE Main 2022 (Online) 27th June Evening Shift

Let [t] denote the greatest integer $\le$ t and {t} denote the fractional part of t. The integral value of α for which the left hand limit of the function

$f(x)=[1+x]+\frac{\alpha^{2[x]+\{x\}}+[x]-1}{2[x]+\{x\}}$ at x = 0 is equal to $\alpha-\frac{4}{3}$, is _____.

Open solution

Q270. JEE Main 2022 (Online) 25th June Evening Shift

Let $f(x)=\left[2x^2+1\right]$ and $g(x)=\begin{cases}2x-3, & x<0\\ 2x+3, & x\ge 0\end{cases}$, where [t] is the greatest integer $\le$ t. Then, in the open interval $(-1, 1)$, the number of points where fog is discontinuous is equal to _____.

Open solution

Q271. JEE Main 2022 (Online) 24th June Morning Shift

The number of points where the function

$$f(x)=\begin{cases}|2x^2-3x-7| & if \quad x\le -1\\ [4x^2-1] & if \quad -1<x<1\\ |x+1|+|x-2| & if \quad x\ge 1\end{cases}$$

[t] denotes the greatest integer $\le$ t, is discontinuous is _____.

Open solution

Q272. JEE Main 2021 (Online) 1st September Evening Shift

Let $f(x)=x^6+2x^4+x^3+2x+3$, x $\in$ R. Then the natural number n for which $\lim\limits_{x\to 1}\frac{x^n f(1)-f(x)}{x-1}=44$ is _____.

Open solution

Q273. JEE Main 2021 (Online) 1st September Evening Shift

Let [t] denote the greatest integer $\le$ t. The number of points where the function $f(x)=[x]\left|x^2-1\right|+\sin\left(\frac{\pi}{[x]+3}\right)-[x+1], x\in(-2,2)$ is not continuous is _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q274. JEE Main 2021 (Online) 26th August Morning Shift

Let a, b $\in$ R, b $\neq$ 0, Define a function

$$f(x) = \begin{cases} a\sin\frac{\pi}{2}(x-1), & for\ x \leq 0 \\ \frac{\tan 2x - \sin 2x}{bx^3}, & for\ x > 0 \end{cases}.$$

If f is continuous at x = 0, then $10 - ab$ is equal to _____.

Open solution

Q275. JEE Main 2021 (Online) 27th July Morning Shift

Let $f : [0, 3] \to R$ be defined by $f(x) = \min\{x - [x], 1 + [x] - x\}$ where [x] is the greatest integer less than or equal to x. Let P denote the set containing all x $\in$ [0, 3] where f i discontinuous, and Q denote the set containing all x $\in$ (0, 3) where f is not differentiable. Then the sum of number of elements in P and Q is equal to _____.

Open solution

Q276. JEE Main 2021 (Online) 25th July Evening Shift

Consider the function

where P(x) is a polynomial such that P" (x) is always a constant and P(3) = 9. If f(x) is continuous at x = 2, then P(5) is equal to _____.

$$f(x) = \begin{cases} \dfrac{P(x)}{\sin(x-2)}, & x \neq 2 \\ = 7, & x = 2 \end{cases}$$

Open solution

Q277. JEE Main 2021 (Online) 22th July Evening Shift

Let f : R $\to$ R be a function defined as $f(x) = \begin{cases} 3\left(1 - \frac{|x|}{2}\right) & if \quad |x| \leq 2 \\ 0 & if \quad |x| > 2 \end{cases}$

Let g : R $\to$ R be given by $g(x) = f(x + 2) - f(x - 2)$. If n and m denote the number of points in R where g is not continuous and not differentiable, respectively, then n + m is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q278. JEE Main 2021 (Online) 20th July Evening Shift

Let a function g : [0, 4] $\rightarrow$ R be defined as

$g(x) = \begin{cases} \max\limits_{0 \le t \le x} \{t^3 - 6t^2 + 9t - 3\}, & 0 \le x \le 3 \\ 4 - x, & 3 < x \le 4 \end{cases}$, then the number of points in the interval (0, 4) where g(x) is NOT differentiable, is _____.

Open solution

Q279. JEE Main 2021 (Online) 20th July Evening Shift

If $\lim\limits_{x \to 0} \frac{\alpha x e^x - \beta \log_e(1+x) + \gamma x^2 e^{-x}}{x \sin^2 x} = 10, \alpha, \beta, \gamma \in R$, then the value of $\alpha + \beta + \gamma$ is _____.

Open solution

Q280. JEE Main 2021 (Online) 20th July Morning Shift

If the value of $\lim\limits_{x \to 0} (2 - \cos x \sqrt{\cos 2x})^{\left(\frac{x+2}{x^2}\right)}$ is equal to e^a, then a is equal to _____.

Open solution

Q281. JEE Main 2021 (Online) 18th March Evening Shift

Let f : R $\rightarrow$ R satisfy the equation f(x + y) = f(x) . f(y) for all x, y $\in$R and f(x) $\neq$ 0 for any x$\in$R. If the function f is differentiable at x = 0 and f'(0) = 3, then

$\lim\limits_{h \to 0} \frac{1}{h}(f(h) - 1)$ is equal to _____.

Open solution

Q282. JEE Main 2021 (Online) 17th March Morning Shift

If the function $f(x) = \frac{\cos(\sin x) - \cos x}{x^4}$ is continuous at each point in its domain and $f(0) = \frac{1}{k}$, then k is _____.

Open solution

Q283. JEE Main 2021 (Online) 16th March Evening Shift

Let f : R $\rightarrow$ R and g : R $\rightarrow$ R be defined as

$f(x) = \begin{cases} x + a, & x < 0 \\ |x - 1|, & x \ge 0 \end{cases}$ and

$g(x) = \begin{cases} x + 1, & x < 0 \\ (x-1)^2 + b, & x \ge 0 \end{cases}$,

where a, b are non-negative real numbers. If (gof) (x) is continuous for all x $\in$ R, then a + b is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q284. JEE Main 2021 (Online) 16th March Morning Shift

If $\lim\limits_{x\to 0}\frac{ae^x-b\cos x+ce^{-x}}{x\sin x}=2$, then a + b + c is equal to _____.

Open solution

Q285. JEE Main 2021 (Online) 25th February Evening Shift

A function f is defined on $[-3, 3]$ as

$f(x)=\begin{cases}\min\{|x|,2-x^2\}, & -2\le x\le 2\\ [|x|], & 2<|x|\le 3\end{cases}$ where [x] denotes the greatest integer $\le$ x. The number of points, where f is not differentiable in $(-3, 3)$ is _____.

Open solution

Q286. JEE Main 2021 (Online) 25th February Evening Shift

If $\lim\limits_{x\to 0}\frac{ax-(e^{4x}-1)}{ax(e^{4x}-1)}$ exists and is equal to b, then the value of a $-$ 2b is _____.

Open solution

Q287. JEE Main 2021 (Online) 25th February Morning Shift

The number of points, at which the function

f(x) = | 2x + 1 | $-$ 3| x + 2 | + | x^2 + x $-$ 2 |, x$\in$R is not differentiable, is _____.

Open solution

Q288. JEE Main 2021 (Online) 24th February Morning Shift

$\lim\limits_{n\to\infty}\tan\left\{\sum\limits_{r=1}^{n}\tan^{-1}\left(\frac{1}{1+r+r^2}\right)\right\}$ is equal to _____.

Open solution

Q289. JEE Main 2020 (Online) 6th September Morning Slot

Let f : R $\to$ R be defined as

$f(x)=\begin{cases}x^5\sin\left(\frac{1}{x}\right)+5x^2, & x<0\\ 0, & x=0\\ x^5\cos\left(\frac{1}{x}\right)+\lambda x^2, & x>0\end{cases}$

The value of λ for which f "(0) exists, is _____.

Open solution

Q290. JEE Main 2020 (Online) 5th September Morning Slot

Let $f(x)=x.\left[\frac{x}{2}\right]$, for -10< x < 10, where [t] denotes the greatest integer function. Then the number of points of discontinuity of f is equal to _____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q291. JEE Main 2020 (Online) 4th September Morning Slot

Suppose a differentiable function f(x) satisfies the identity f(x+y) = f(x) + f(y) + $xy^2 + x^2y$, for all real x and y.
$\lim_{x\to 0} \frac{f(x)}{x} = 1$, then f'(3) is equal to _____.

Open solution

Q292. JEE Main 2020 (Online) 3rd September Morning Slot

If $\lim_{x\to 0} \left\{ \frac{1}{x^8} \left(1 - \cos\frac{x^2}{2} - \cos\frac{x^2}{4} + \cos\frac{x^2}{2}\cos\frac{x^2}{4} \right) \right\}$ = 2^{-k}

then the value of k is _____ .

Open solution

Q293. JEE Main 2020 (Online) 2nd September Morning Slot

If $\lim_{x\to 1} \frac{x+x^2+x^3+...+x^n-n}{x-1}$ = 820,
($n \in N$) then the value of n is equal to _____.

Open solution

Q294. JEE Main 2020 (Online) 7th January Evening Slot

If the function f defined on $\left(-\frac{1}{3}, \frac{1}{3}\right)$ by

$$f(x) = \begin{cases} \frac{1}{x}\log_e\left(\frac{1+3x}{1-2x}\right), & when\, x \neq 0 \\ k, & when\, x = 0 \end{cases}$$

is continuous, then k is equal to_____.

Open solution

Q295. JEE Main 2020 (Online) 7th January Morning Slot

Let S be the set of points where the function, f(x) = |2-|x-3||, $x \in R$ is not differentiable. Then $\sum_{x\in S} f(f(x))$ is equal to_____.

Open solution

Q296. JEE Main 2020 (Online) 7th January Morning Slot

$\lim_{x\to 2} \frac{3^x+3^{3-x}-12}{3^{-x/2}-3^{1-x}}$ is equal to_____.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Answer Key

MCQ (Single Correct Answer)

1. **D**	2. **B**	3. **C**	4. **B**	5. **C**
6. **B**	7. **A**	8. **D**	9. **C**	10. **C**
11. **A**	12. **D**	13. **B**	14. **C**	15. **D**
16. **C**	17. **D**	18. **C**	19. **C**	20. **C**
21. **C**	22. **C**	23. **B**	24. **D**	25. **D**
26. **D**	27. **A**	28. **C**	29. **D**	30. **D**
31. **B**	32. **B**	33. **C**	34. **A**	35. **D**
36. **D**	37. **D**	38. **C**	39. **C**	40. **C**
41. **D**	42. **B**	43. **D**	44. **D**	45. **A**
46. **B**	47. **D**	48. **C**	49. **A**	50. **C**
51. **D**	52. **C**	53. **A**	54. **B**	55. **D**
56. **C**	57. **D**	58. **B**	59. **D**	60. **B**
61. **D**	62. **A**	63. **C**	64. **B**	65. **B**
66. **D**	67. **B**	68. **A**	69. **D**	70. **D**
71. **C**	72. **B**	73. **B**	74. **B**	75. **C**
76. **B**	77. **A**	78. **D**	79. **A**	80. **C**
81. **D**	82. **B**	83. **C**	84. **C**	85. **A**
86. **C**	87. **B**	88. **D**	89. **D**	90. **A**
91. **A**	92. **C**	93. **D**	94. **C**	95. **A**
96. **C**	97. **B**	98. **C**	99. **A**	100. **A**
101. **C**	102. **B**	103. **C**	104. **D**	105. **A**
106. **A**	107. **B**	108. **D**	109. **D**	110. **D**
111. **B**	112. **B**	113. **C**	114. **D**	115. **A**
116. **A**	117. **C**	118. **A**	119. **A**	120. **C**
121. **A**	122. **B**	123. **A**	124. **A**	125. **B**
126. **C**	127. **D**	128. **C**	129. **A**	130. **D**
131. **B**	132. **B**	133. **D**	134. **C**	135. **A**
136. **D**	137. **A**	138. **D**	139. **C**	140. **D**
141. **A**	142. **B**	143. **C**	144. **B**	145. **C**
146. **A**	147. **B**	148. **B**	149. **D**	150. **C**

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

151. **D**	152. **A**	153. **D**	154. **A**	155. **A**
156. **D**	157. **D**	158. **B**	159. **C**	160. **D**
161. **C**	162. **B**	163. **C**	164. **B**	165. **A**
166. **D**	167. **B**	168. **C**	169. **B**	170. **D**
171. **B**	172. **C**	173. **C**	174. **A**	175. **A**
176. **C**	177. **B**	178. **A**	179. **A**	180. **A**
181. **B**	182. **C**	183. **B**	184. **A**	185. **C**
186. **B**	187. **A**	188. **C**	189. **D**	190. **B**
191. **B**	192. **D**	193. **C**	194. **B**	195. **B**
196. **A**	197. **B**	198. **D**	199. **B**	200. **D**
201. **D**	202. **D**	203. **B**	204. **D**	205. **D**
206. **B**	207. **B**	208. **A**	209. **C**	210. **D**
211. **B**	212. **D**	213. **B**	214. **C**	215. **A**
216. **B**	217. **C**	218. **A**	219. **C**	220. **B**
221. **B**	222. **C**	223. **D**	224. **B**	225. **D**
226. **C**	227. **D**	228. **A**	229. **C**	230. **D**
231. **A**	232. **B**	233. **D**	234. **C**	

Numerical

235. **3**	236. **4**	237. **3**	238. **10**	239. **2**
240. **98**	241. **8**	242. **3**	243. **32**	244. **24**
245. **1**	246. **5**	247. **34**	248. **81**	249. **6**
250. **55**	251. **17**	252. **170**	253. **24**	254. **100**
255. **18**	256. **81**	257. **15**	258. **2**	259. **2**
260. **4**	261. **309**	262. **25**	263. **79**	264. **2**
265. **1**	266. **3**	267. **14**	268. **11**	269. **3**
270. **62**	271. **7**	272. **7**	273. **2**	274. **14**
275. **5**	276. **39**	277. **4**	278. **1**	279. **3**
280. **3**	281. **3**	282. **6**	283. **1**	284. **4**
285. **5**	286. **5**	287. **2**	288. **1**	289. **5**
290. **8**	291. **10**	292. **8**	293. **40**	294. **5**
295. **3**	296. **36**			

For numerical answers given as a range, any value within the range is correct. Full solutions: scan the QR code next to each question, or visit questions.examside.com.

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

ExamGOAL
PYQ PDF

Your Exam. Our Responsibility.

From first chapter to exam day.

Practise with the questions your exam has already asked, then test yourself under real exam conditions.

PAST YEAR QUESTIONS

Every past paper, chapter-wise.

Previous year questions sorted the way you study, each with a full solution.

- Chapter-wise and year-wise
- Weightage and topic tags on every chapter
- Your progress on each one, at a glance

MOCK TESTS

Mock tests with a real exam feel.

Practise under the same pressure as the real paper, on your phone or on the big screen.

- Timed sections, just like exam day
- Mark questions for review and come back

ANALYSIS

Know your score. Know your rank.

A full report after every test, so you know exactly where you stand.

- Your percentile among everyone who took it
- Predicted real-exam percentile and All India Rank
- Subject breakdown and time analysis

PRACTICE

Solve. Check. Understand.

Learn from every question, not just the ones you get right.

- Instant answers as you practise
- Step-by-step explanations
- See how everyone else answered

NCERT Exemplar
With full solutions

Exam syllabus
Subject and chapter-wise

Bookmarks
Organised in groups

Your notes
On any question

ExamGOAL
LEARNING APP

ExamGOAL: Exam Preparation App

One app for every exam. Free trial for new users, on Android and the web.

GET IT ON Google Play

Scan to get the app

www.examgoal.com
Practice and mock tests

questions.examside.com
Free solutions

pdf.questions.examgoal.com
Free PYQ PDFs