ExamGOAL
PYQ PDF

Free · With answer key

NEET · CHEMISTRY · CHAPTER-WISE PYQS

Solutions

NEET Chemistry · Previous Year Questions with Answer Key

70	40	1988 – 2026	English
QUESTIONS	PAPERS COVERED	YEARS	LANGUAGE

INSIDE THIS PDF

MCQ (Single Correct Answer)	69 questions
MCQ (More than One Correct Answer)	1 question
Answer key	at the end

Practise online

Every question with its full solution, free on **questions.examside.com**

1. **Solve**
   Print it, or work through it on screen.
2. **Check**
   The answer key is on the last pages.
3. **Understand**
   Scan the QR code beside any question for its full solution.

MCQ (Single Correct Answer) 69 questions

Q1. RE-NEET 2026

Consider the following statements about the solutions formed by mixing two liquids.

A. An ideal solution thus formed obeys Raoult's law throughout the composition range.

B. Mixture of chloroform and acetone shows negative deviation from Raoult's law.

C. Mixture of aniline and phenol shows positive deviation from Raoult's law.

Choose the correct statements from the options given below:

(A) A and C only (B) A and B only (C) B and C only (D) A only

Open solution

Q2. RE-NEET 2026

Assertion A: For an ideal solution formed by mixing liquids $\mathbf{P}$ and $\mathbf{Q}$, $\Delta_{mix} H = 0$ and $\Delta_{mix} V = 0$

Reason R: No interactions occur between P and Q

In the light of the above statements, choose the most appropriate answer from the options given below.

(A) $\mathbf{A}$ is not correct but $\mathbf{R}$ is correct

(B) Both $\mathbf{A}$ and $\mathbf{R}$ are correct and $\mathbf{R}$ is the correct explanation of $\mathbf{A}$

(C) Both $\mathbf{A}$ and $\mathbf{R}$ are correct but $\mathbf{R}$ is NOT the correct explanation of $\mathbf{A}$

(D) $\mathbf{A}$ is correct but $\mathbf{R}$ is not correct

Open solution

Q3. NEET 2026

Identify the correct statements :

(A) The molality of 2.5 g of ethanoic acid (Molar mass : $60\ \text{g mol}^{-1}$) in 75 g of benzene solution is 0.556 m .

(B) The molarity of a solution containing 5 g of NaOH (molar mass : $40\ \text{g mol}^{-1}$) in 450 mL of solution is 0.278 M at 298 K .

(C) Aquatic species are more comfortable in cold water.

(D) The solubility of gas increases with decrease in pressure.

(E) For a binary mixture of A and B, the number of moles of A and B are n_A and n_B respectively. The mole fraction of B will be $x_B = \frac{n_A}{n_A + n_B}$.

Choose the correct answer from the options given below :

(A) A, B and C only (B) A and B only (C) A and C only (D) A, D and E only

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q4. NEET 2026

Mixture of chloroform and acetone forms a solution with negative deviation from Raoult's law due to :

Open solution

(A) Increase in escaping tendency of molecules of each component.

(B) Formation of hydrogen bonding between acetone and chloroform

(C) Stronger intermolecular forces between chloroform molecules than those between chloroform and acetone molecules.

(D) Repulsive forces.

Q5. NEET 2025

Match List-I with List-II

Open solution

	List - I (Example)		List - II (Type of Solution)
A	Humidity	I	Solid in solid
B	Alloys	II	Liquid in gas
C	Amalgams	III	Solid in gas
D	Smoke	IV	Liquid in solid

Choose the correct answer from the options given below:

(A) A-III, B-I, C-IV, D-II

(B) A-III, B-II, C-I, D-IV

(C) A-II, B-IV, C-I,D-III

(D) A-II, B-I, C-IV, D-III

Q6. NEET 2025

5 moles of liquid X and 10 moles of liquid Y make a solution having a vapour pressure of 70 torr. The vapour pressures of pure X and Y are 63 torr and 78 torr respectively. Which of the following is true regarding the described solution?

Open solution

(A) The solution is ideal.

(B) The solution has volume greater than the sum of individual volumes.

(C) The solution shows positive deviation.

(D) The solution shows negative deviation.

Q7. NEET 2024 (Re-Examination)

Mass of glucose $(C_6H_{12}O_6)$ required to be dissolved to prepare one litre of its solution which is isotonic with $15\ g\ L^{-1}$ solution of urea (NH_2CONH_2) is (Given: Molar mass in $gmol^{-1} C : 12, H : 1, O : 16, N : 14$)

Open solution

(A) 55 g (B) 15 g (C) 30 g (D) 45 g

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q8. NEET 2024

The Henry's law constant (K_H) values of three gases (A, B, C) in water are $145, 2 \times 10^{-5}$ and $35\ \text{kbar}$ respectively. The solubility of these gases in water follows the order:

(A) B > A > C (B) B > C > A (C) A > C > B (D) A > B > C

Open solution

Q9. NEET 2024

The plot of osmotic pressure (Π) vs concentration (mol L^{-1}) for a solution gives a straight line with slope $25.73\ \text{L bar mol}^{-1}$. The temperature at which the osmotic pressure measurement is done is

(Use $R = 0.083\ \text{L bar mol}^{-1}\ \text{K}^{-1}$)

(A) 37°C (B) 310°C (C) 25.73°C (D) 12.05°C

Open solution

Q10. NEET 2023 Manipur

Which amongst the following aqueous solution of electrolytes will have minimum elevation in boiling point? Choose the correct option :-

(A) 0.05 M NaCl (B) 0.1 M KCl (C) 0.1 M $MgSO_4$ (D) 1 M NaCl

Open solution

Q11. NEET 2022 Phase 2

K_H value for some gases at the same temperature 'T' are given :

Gas	K_H/k bar
Ar	40.3
CO_2	1.67
HCHO	1.83×10^{-5}
CH_4	0.413

where K_H is Henry's Law constant in water. The order of their solubility in water is :

(A) HCHO < CH_4 < CO_2 < Ar (B) Ar < CO_2 < CH_4 < HCHO

(C) Ar < CH_4 < CO_2 < HCHO (D) HCHO < CO_2 < CH_4 < Ar

Open solution

Q12. NEET 2021

The following solutions were prepared by dissolving 10 g of glucose ($C_6H_{12}O_6$) in 250 ml of water (P_1), 10 g of urea (CH_4N_2O) in 250 ml of water (P_2) and 10 g of sucrose ($C_{12}H_{22}O_{11}$) in 250 ml of water (P_3). The right option for the decreasing order of osmotic pressure of these solutions is :

(A) $P_3 > P_1 > P_2$ (B) $P_2 > P_1 > P_3$ (C) $P_1 > P_2 > P_3$ (D) $P_2 > P_3 > P_1$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q13. NEET 2021

The correct option for the value of vapour pressure of a solution at 45°C with benzene to octane in molar ratio 3 : 2 is :

[At 45°C vapour pressure of benzene is 280 mm Hg and that of octane is 420 mm Hg. Assume Ideal gas]

(A) 350 mm of Hg (B) 160 mm of Hg (C) 168 mm of Hg (D) 336 mm of Hg

Open solution

Q14. NEET 2020 Phase 2

If 8 g of a non-electrolyte solute is dissolved in 114 g of n-octane to reduce its vapour pressure to 80%, the molar mass (in $\mathrm{g\,mol^{-1}}$) of the solute is

[Given that molar mass of n-octane is $114\ \mathrm{g\,mol^{-1}}$]

(A) 20 (B) 40 (C) 60 (D) 80

Open solution

Q15. NEET 2020 Phase 2

Isotonic solutions have same

(A) boiling temperature (B) vapour pressure

(C) freezing temperature (D) osmotic pressure

Open solution

Q16. NEET 2020 Phase 1

The freezing point depression constant (K_f) of benzene is 5.12 K kg mol^{-1}. The freezing point depression for the solution of molality 0.078 m containing a non-electrolyte solute in benzene is rounded off upto two decimal places :

(A) 0.80 K (B) 0.40 K (C) 0.60 K (D) 0.20 K

Open solution

Q17. NEET 2020 Phase 1

The mixture which shows positive deviation from Raoult's law is :

(A) Benzene + Toluene (B) Acetone + Chloroform

(C) Chloroethane + Bromoethane (D) Ethanol + Acetone

Open solution

Q18. NEET 2019 (Odisha)

Which of the following statements is correct regarding a solution of two components A and B exhibiting positive deviation from ideal behaviour?

(A) Intermolecular attractive forces between A-A and B-B are equal to those between A-B.

(B) Intermolecular attractive forces between A-A and B-B are stronger than those between A-B.

(C) $\Delta_{\mathrm{mix}}H = 0$ at constant T and P.

(D) $\Delta_{\mathrm{mix}}V = 0$ at constant T and P.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q19. NEET 2019

For an ideal solution, the **correct** option is :

(A) $\Delta_{mix}H = 0$ at constant T and P

(B) $\Delta_{mix}G = 0$ at constant T and P

(C) $\Delta_{mix}S = 0$ at constant T and P

(D) $\Delta_{mix}V \neq 0$ at constant T and P

Open solution

Q20. NEET 2019

The mixture that forms maximum boiling azeotrope is :

(A) Ethanol + Water

(B) Acetone + Carbon disulphide

(C) Heptane + Octane

(D) Water + Nitric acid

Open solution

Q21. NEET 2017

If molality of the dilute solution is doubled, the value of molal depression constant (K_f) will be

(A) halved

(B) tripled

(C) unchanged

(D) doubled.

Open solution

Q22. NEET 2016 Phase 2

The van't Hoff factor (i) for a dilute aqueous solution of the strong electrolyte barium hydroxide is

(A) 0

(B) 1

(C) 2

(D) 3

Open solution

Q23. NEET 2016 Phase 2

Which one of the following is incorrect for ideal solution?

(A) $\Delta H_{mix} = 0$

(B) $\Delta U_{mix} = 0$

(C) $\Delta P = P_{obs} - P_{\text{calculated by Raoult's law}}$

(D) $\Delta G_{mix} = 0$

Open solution

Q24. NEET 2016 Phase 1

Which of the following statements about the composition of the vapour over an ideal 1 : 1 molar mixture of benzene and toluene is correct? Assume that the temperature is constant at 25°C. (Given, vapour pressure data at 25°C, benzene = 12.8 kPa, toluene = 3.85 kPa)

(A) The vapour will contain equal amounts of benzene and toluene.

(B) Not enough information is given to make a prediction.

(C) The vapour will contain a higher percentage of benzene.

(D) The vapour will contain a higher percentage of toluene

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q25. NEET 2016 Phase 1

At 100°C the vapour pressure of a solution of 6.5 g of a solute in 100 g water is 732 mm. If K_b = 0.52, the boiling point of this solution will be

(A) 102°C (B) 103°C (C) 101°C (D) 100°C

Open solution

Q26. AIPMT 2015 Cancelled Paper

The boiling point of 0.2 mol kg^{-1} solution of *X* in water is greater than equimolal solution of *Y* in water. Which one of the following statements is true in this case?

(A) Molecular mass of *X* is less than the molecular mass of *Y*.

(B) *Y* is undergoing dissociation in water while *X* undergoes no change.

(C) *X* is undergoing dissociation in water.

(D) Molecular mass of *X* is greater than the molecular mass of *Y*.

Open solution

Q27. AIPMT 2015 Cancelled Paper

Which one of the following electrolytes has the same value of van't Hoff factor (i) as that of $Al_2(SO_4)_3$ (if all are 100% ionised)?

(A) $Al(NO_3)_3$ (B) $K_4[Fe(CN)_6]$ (C) K_2SO_4 (D) $K_3[Fe(CN)_6]$

Open solution

Q28. AIPMT 2015 Cancelled Paper

Which of them is not equal to zero for an ideal solution?

(A) ΔV_{mix} (B) $\Delta P = P_{observed} - P_{Raoult}$

(C) ΔH_{mix} (D) ΔS_{mix}

Open solution

Q29. AIPMT 2014

Of the following 0.10 m aqueous solutions, which one will exhibit the largest freezing point depression?

(A) KCl (B) $C_6H_{12}O_6$ (C) $Al_2(SO_4)_3$ (D) K_2SO_4

Open solution

Q30. NEET 2013 (Karnataka)

Which condition is not satisfied by an ideal solution?

(A) $\Delta_{mix} V = 0$ (B) $\Delta_{mix} S = 0$

(C) Obeyance to Raoult's Law (D) $\Delta_{mix} H = 0$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q31. AIPMT 2012 Mains

Vapour pressure of chloroform ($CHCl_3$) and dichloromethane (CH_2Cl_2) at 25°C are 200 mm Hg and 41.5 mm Hg respectively. Vapour pressure of the solution obtained by mixing 25.5 g of $CHCl_3$ and 40 g of CH_2Cl_2 at the same temperature will be
(Molecular mass of $CHCl_3$ = 119.5 u and molecular mass of CH_2Cl_2 = 85 u)

(A) 173.9 mm Hg (B) 615.0 mm Hg (C) 90.6 mm Hg (D) 285.5 mm Hg

Open solution

Q32. AIPMT 2012 Prelims

p_A and p_B are the vapour pressure of pure liquid components, *A* and *B*, respectively of an ideal binary solution. If x_A represents the mole fraction of component *A*, the total pressure of the solution will be

(A) $p_A + x_A(p_B - p_A)$ (B) $p_A + x_A(p_A - p_B)$
(C) $p_B + x_A(p_B - p_A)$ (D) $p_B + x_A(p_A - p_B)$

Open solution

Q33. AIPMT 2011 Mains

A 0.1 molal aqueous solution of a weak acid is 30% ionized. If K_f for water is 1.86°C/m, the freezing point of the solution will be

(A) -0.18°C (B) -0.54°C (C) -0.36°C (D) -0.24°C

Open solution

Q34. AIPMT 2011 Mains

200 mL of an aqueous solution of a protein contains its 1.26 g. The osmotic pressure of this solution at 300 K is found to be 2.57×10^{-3} bar. The molar mass of protein will be (R = 0.083 L bar mol^{-1} K^{-1})

(A) 51022 g mol^{-1} (B) 122044 g mol^{-1} (C) 31011 g mol^{-1} (D) 61038 g mol^{-1}

Open solution

Q35. AIPMT 2011 Prelims

The freezing point depression constant for water is -1.86^o C m^{-1}. If 5.00 g Na_2SO_4 is dissolved in 45.0 g H_2O, the freezing point is changed by -3.82°C. Calculate the van't Hoff factor for Na_2SO_4.

(A) 2.05 (B) 2.63 (C) 3.11 (D) 0.381

Open solution

Q36. AIPMT 2011 Prelims

The van't Hoff factor i for a compound which undergoes dissociation in one solvent and association in other solvent is respectively

(A) less than one and greater than one (B) less than one and less than one
(C) greater than one and less than one (D) greater than one and greater than one.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q37. AIPMT 2010 Prelims

An aqueous solution is 1.00 molal in KI. Which change will cause the vapour pressure of the solution to increase?

(A) Addition of NaCl
(B) Addition of Na_2SO_4
(C) Addition of 1.00 molal KI
(D) Addition of water

Open solution

Q38. AIPMT 2010 Prelims

A solution of sucrose (molar mass = 342 g mol^{-}) has been prepared by dissolving 68.5 g of sucrose in 1000 g of water. The freezing point of the solution obtained will be (K_f for water = 1.86 K kg mol $^{-1}$)

(A) -0.372^oC (B) -0.520^oC (C) $+0.372^oC$ (D) -0.570^oC

Open solution

Q39. AIPMT 2009

A 0.0020 m aqueous solution of an ionic compound $[Co(NH_3)_5(NO_2)]Cl$ freezes at -0.00732^oC. Number of moles of ions which 1 mol of ionic compound produces on being dissolved in water will be ($K_f = -1.86^oC/m$)

(A) 3 (B) 4 (C) 1 (D) 2

Open solution

Q40. AIPMT 2007

0.5 molal aqueous solution of a weak acid (HX) is 20% ionised. If K_f for water is 1.86 K kg mol^{-1}, the lowering in freezing point of the solution is

(A) 0.56 K (B) 1.12 K (C) -0.56 K (D) -1.12 K

Open solution

Q41. AIPMT 2006

During osmosis, flow of water through a semipermeable membrane is

(A) from solution having lower concentration only
(B) from solution having higher concentration only
(C) from both sides of semipermeable membrane with equal flow rates
(D) from both sides of semipermeable membrane with unequal flow rates.

Open solution

Q42. AIPMT 2006

A solution of acetone in ethanol

(A) obeys Raoult's law
(B) shows a negative deviation from Raoult's law
(C) shows a positive deviation from Raoult's law
(D) behaves like a near ideal solution.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q43. AIPMT 2006

1.00 g of a non-electrolyte solute (molar mass 250 g mol^{-1}) was dissolved in 51.2 g of benzene. If the freezing point depression constant, K_f of benzene is 5.12 K kg mol^{-1}, the freezing point of benzene will be lowered by

(A) 0.2 K (B) 0.4 K (C) 0.3 K (D) 0.5 K

Open solution

Q44. AIPMT 2006

A solution containing 10 g per dm^3 of urea (molecular mass = 60 g mol^{-1}) is isotonic with a 5% solution of a nonvolatile solute. The molecular mass of this nonvolatile solute is

(A) 200 g mol^{-1} (B) 250 g mol^{-1} (C) 300 g mol^{-1} (D) 350 g mol^{-1}

Open solution

Q45. AIPMT 2005

The vapour pressure of two liquids P and Q are 80 and 60 torr, respectively. The total vapour pressure of solution obtained by mixing 3 mole of P and 2 mole of Q would be

(A) 72 torr (B) 140 torr (C) 68 torr (D) 20 torr

Open solution

Q46. AIPMT 2005

A solution has a 1 : 4 mole ratio of pentane to hexane. The vapour pressures of the pure hydrocarbons at 20°C are 440 mm Hg for pentane and 120 mm Hg for hexane. The mole fraction of pentane in the vapour phase would be

(A) 0.200 (B) 0.549 (C) 0.786 (D) 0.478

Open solution

Q47. AIPMT 2005

A solution of urea (mol. mass 56 g mol^{-1}) boils at 100.18°C at the atmospheric pressure. If K_f and K_b for water are 1.86 and 0.512 K kg mol^{-1} respectively, the above solution will freeze at

(A) 0.654°C (B) − 0.654°C (C) 6.54°C (D) −6.54°C

Open solution

Q48. AIPMT 2004

Camphor is often used in molecular mass determination because

(A) it is readily available (B) it has a very high cryoscopic constant
(C) it is volatile (D) it is solvent for organic substances

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q49. AIPMT 2002

A solution containing components A and B follows Raoult's law

(A) A - B attraction force is greater than A - A and B - B

(B) A - B attraction force is less than A - A and B - B

(C) A - B attraction force remains same as A - A and B - B

(D) volume of solution is different from sum of volume of solute and solvent.

Open solution

Q50. AIPMT 2002

A solution contains non volatile solute of molecular mass M_2. Which of the following can be used to calculate the molecular mass of solute in terms of osmotic pressure?

(A) $M_2 = \left(\frac{m_2}{\pi}\right)VRT$

(B) $M_2 = \left(\frac{m_2}{V}\right)\frac{RT}{\pi}$

(C) $M_2 = \left(\frac{m_2}{V}\right)\pi RT$

(D) $M_2 = \left(\frac{m_2}{V}\right)\frac{\pi}{RT}$

Open solution

Q51. AIPMT 2001

Pure water can be obtained from sea water by

(A) centrifugation (B) plasmolysis (C) reverse osmosis (D) sedimentation

Open solution

Q52. AIPMT 2000

From the colligative properties of solution, which one is the best method for the determination of molecular weight of proteins and polymers?

(A) Osmotic pressure

(B) Lowering in vapour pressure

(C) Lowering in freezing point

(D) Elevation in boiling point

Open solution

Q53. AIPMT 1999

The vapour pressure of benzene at a certain temperature is $640\ \text{mm}$ of Hg. A non-volatile and non-electrolyte solid, weighing $2.175\ \text{g}$ is added to $39.08\ \text{g}$ of benzene. If the vapour pressure of the solution is $600\ \text{mm}$ of Hg, what is the molecular weight of solid substance?

(A) 49.50 (B) 59.60 (C) 69.40 (D) 79.82

Open solution

Q54. AIPMT 1999

If $0.15\ \text{g}$ of solute, dissolved in $15\ \text{g}$ of solvent, is boiled at a temperature higher by 0.216°C, than that of the pure solvent, the molecular weight of the substance is (molal elevation constant for the solvent is 2.16°C)

(A) 1.01 (B) 10 (C) 10.1 (D) 100

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q55. AIPMT 1998

The vapour pressure of a solvent decreased by $10\ \text{mm}$ of mercury when a non-volatile solute was added to the solvent. The mole fraction of the solute in the solution is 0.2. What should be the mole fraction of the solvent if the decrease in the vapour pressure is to be $20\ \text{mm}$ of mercury?

(A) 0.8 (B) 0.6 (C) 0.4 (D) 0.2

Open solution

Q56. AIPMT 1998

A 5% solution of cane sugar (mol. wt. = 342) is isotonic with 1% solution of a substance X. The molecular weight of X is

(A) 34.2 (B) 171.2 (C) 68.4 (D) 136.8

Open solution

Q57. AIPMT 1997

Which of the following $0.10\ \text{m}$ aqueous solution will have the lowest freezing point?

(A) KI (B) $\text{C}_{12}\text{H}_{22}\text{O}_{11}$ (C) $\text{Al}_2(\text{SO}_4)_3$ (D) $\text{C}_5\text{H}_{10}\text{O}_5$

Open solution

Q58. AIPMT 1996

The vapour pressure of CCl_4 at $25\ ^\circ\text{C}$ is $143\ \text{mm Hg}$. If $0.5\ \text{g}$ of a non-volatile solute (mol. weight $= 65$) is dissolved in $100\ \text{mL}$ CCl_4, the vapour pressure of the solution will be

(A) $199.34\ \text{mm Hg}$ (B) $143.99\ \text{mm Hg}$ (C) $141.43\ \text{mm Hg}$ (D) $94.39\ \text{mm Hg}$

Open solution

Q59. AIPMT 1996

The relationship between osmotic pressure at $273\ \text{K}$ when 10 g glucose (p_1), 10 g urea (p_2), and 10 g sucrose (p_3) are dissolved in 250 mL of water is

(A) $p_2 > p_1 > p_3$ (B) $p_2 > p_3 > p_1$ (C) $p_1 > p_2 > p_3$ (D) $p_3 > p_1 > p_2$

Open solution

Q60. AIPMT 1995

According to Raoult's law, the relative lowering of vapour pressure for a solution is equal to

(A) mole fraction of solute (B) mole fraction of solvent
(C) moles of solute (D) moles of solvent

Open solution

Q61. AIPMT 1994

At $25\ ^\circ\text{C}$, the highest osmotic pressure is exhibited by 0.1 M solution of

(A) glucose (B) urea (C) CaCl_2 (D) KCl

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q62. AIPMT 1994

Which of the following salt has the same value of vant's Hoff factor (i) as that of $K_3[Fe(CN)_6]$?

(A) Na_2SO_4 (B) $Al(NO_3)_3$ (C) $Al_2(SO_4)_3$ (D) NaCl

Open solution

Q63. AIPMT 1992

If 0.1 M solution of glucose and 0.1 M solution of urea are placed on two sides of the semipermeable membrane to equal heights, then it will be correct to say that

(A) there will be no net movement across the membrane

(B) glucose will flow towards glucose solution

(C) urea will flow towards glucose solution

(D) water will flow from urea solution to glucose

Open solution

Q64. AIPMT 1992

Which one is a colligative property?

(A) Boiling point (B) Vapour pressure (C) Osmotic pressure (D) Freezing point

Open solution

Q65. AIPMT 1991

Blood cells retain their normal shape in solution which are

(A) hypotonic to blood (B) isotonic to blood

(C) hypertonic to blood (D) equinormal to blood

Open solution

Q66. AIPMT 1991

Which of the following aqueous solution has minimum freezing point?

(A) $0.01\ m$ NaCl (B) $0.005\ m\ C_2H_5OH$

(C) $0.005\ m\ MgI_2$ (D) $0.005\ m\ MgSO_4$

Open solution

Q67. AIPMT 1991

The relative lowering of the vapour pressure is equal to the ratio between the number of

(A) solute molecules to the solvent molecules

(B) solute molecules to the total molecules in the solution

(C) solvent molecules to the total molecules in the solution

(D) solvent molecules to the total number of ions of the solute

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q68. AIPMT 1988

All form ideal solution except

(A) C_6H_6 and $C_6H_5CH_3$

(B) C_2H_6 and C_2H_5I

(C) C_6H_5Cl and C_6H_5Br

(D) C_2H_5I and C_2H_5OH

Open solution

Q69. AIPMT 1988

An ideal solution is formed when its components

(A) have no volume change on mixing

(B) have no enthalpy change on mixing

(C) have both the above characteristics

(D) have high solubility

Open solution

MCQ (More than One Correct Answer) 1 questions

Q70. NEET 2025

Which of the following aqueous solution will exhibit highest boiling point?

(A) $0.01\ M\ Na_2SO_4$

(B) $0.015\ M\ C_6H_{12}O_6$

(C) 0.01 M Urea

(D) $0.01M\ KNO_3$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Answer Key

MCQ (Single Correct Answer)

1. **B**	2. **D**	3. **A**	4. **B**	5. **D**
6. **D**	7. **D**	8. **B**	9. **A**	10. **A**
11. **B**	12. **B**	13. **D**	14. **B**	15. **D**
16. **B**	17. **D**	18. **B**	19. **A**	20. **D**
21. **C**	22. **D**	23. **D**	24. **C**	25. **C**
26. **C**	27. **B**	28. **D**	29. **C**	30. **B**
31. **C**	32. **D**	33. **D**	34. **D**	35. **B**
36. **C**	37. **D**	38. **A**	39. **D**	40. **B**
41. **D**	42. **C**	43. **B**	44. **C**	45. **A**
46. **D**	47. **B**	48. **B**	49. **C**	50. **B**
51. **C**	52. **A**	53. **C**	54. **D**	55. **B**
56. **C**	57. **C**	58. **C**	59. **A**	60. **A**
61. **C**	62. **B**	63. **A**	64. **C**	65. **B**
66. **A**	67. **B**	68. **D**	69. **C**	

MCQ (More than One Correct Answer)

70. **A**

For numerical answers given as a range, any value within the range is correct. Full solutions: scan the QR code next to each question, or visit questions.examside.com.

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

ExamGOAL
PYQ PDF

Your Exam. Our Responsibility.

From first chapter to exam day.

Practise with the questions your exam has already asked, then test yourself under real exam conditions.

PAST YEAR QUESTIONS

Every past paper, chapter-wise.

Previous year questions sorted the way you study, each with a full solution.

- Chapter-wise and year-wise
- Weightage and topic tags on every chapter
- Your progress on each one, at a glance

MOCK TESTS

Mock tests with a real exam feel.

Practise under the same pressure as the real paper, on your phone or on the big screen.

- Timed sections, just like exam day
- Mark questions for review and come back

ANALYSIS

Know your score. Know your rank.

A full report after every test, so you know exactly where you stand.

- Your percentile among everyone who took it
- Predicted real-exam percentile and All India Rank
- Subject breakdown and time analysis

PRACTICE

Solve. Check. Understand.

Learn from every question, not just the ones you get right.

- Instant answers as you practise
- Step-by-step explanations
- See how everyone else answered

NCERT Exemplar
With full solutions

Exam syllabus
Subject and chapter-wise

Bookmarks
Organised in groups

Your notes
On any question

ExamGOAL
LEARNING APP

ExamGOAL: Exam Preparation App

One app for every exam. Free trial for new users, on Android and the web.

GET IT ON Google Play

Scan to get the app

www.examgoal.com
Practice and mock tests

questions.examside.com
Free solutions

pdf.questions.examgoal.com
Free PYQ PDFs