ExamGOAL
PYQ PDF

Free · With answer key

NEET · PHYSICS · CHAPTER-WISE PYQS

Gravitation

NEET Physics · Previous Year Questions with Answer Key

66	35	2000 – 2026	English
QUESTIONS	PAPERS COVERED	YEARS	LANGUAGE

INSIDE THIS PDF

MCQ (Single Correct Answer)	66 questions
Answer key	at the end

Practise online

Every question with its full solution, free on **questions.examside.com**

1. **Solve**
   Print it, or work through it on screen.
2. **Check**
   The answer key is on the last pages.
3. **Understand**
   Scan the QR code beside any question for its full solution.

MCQ (Single Correct Answer) 66 questions

Q1. RE-NEET 2026

Two planets P_1 and P_2 with equal mass have radii R_1 and R_2, respectively, where $R_2 = \frac{R_1}{2}$. The escape speeds of P_1 and P_2 are v_1 and v_2, respectively. Then $\frac{v_2}{v_1}$ is:

(A) 2 (B) $\frac{1}{\sqrt{2}}$ (C) 1 (D) $\sqrt{2}$

Open solution

Q2. RE-NEET 2026

In a solar system, the time-period of revolution of a planet tracing a circular orbit of radius R is proportional to:

(A) R^3 (B) $R^{1/2}$ (C) $R^{3/2}$ (D) R^2

Open solution

Q3. NEET 2026

The amount of work done to raise a mass ' m ' from the surface of the Earth to a height equal to the radius of the Earth ' R ' will be

(A) $2mgR$ (B) $mg\frac{R}{4}$ (C) mgR (D) $mg\frac{R}{2}$

Open solution

Q4. NEET 2025

The radius of Martian orbit around the Sun is about 4 times the radius of the orbit of Mercury. The Martian year is 687 Earth days. Then which of the following is the length of 1 year on Mercury?

(A) 172 earth days (B) 124 earth days (C) 88 earth days (D) 225 earth days

Open solution

Q5. NEET 2025

A body weighs 48 N on the surface of the earth. The gravitational force experienced by the body due to the earth at a height equal to one-third the radius of the earth from its surface is:

(A) 32 N (B) 36 N (C) 16 N (D) 27 N

Open solution

Q6. NEET 2024 (Re-Examination)

The escape velocity for earth is v. A planet having 9 times mass that of earth and radius, 16 times that of earth, has the escape velocity of:


(A) $\frac{v}{3}$ (B) $\frac{2v}{3}$ (C) $\frac{3v}{4}$ (D) $\frac{9v}{4}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q7. NEET 2024 (Re-Examination)

An object of mass $100\,\text{kg}$ falls from point A to B as shown in figure. The change in its weight, corrected to the nearest integer is (R_E is the radius of the earth)



Open solution

(A) 49 N (B) 89 N (C) 5 N (D) 10 N

Q8. NEET 2024

The mass of a planet is $\frac{1}{10}$th that of the earth and its diameter is half that of the earth. The acceleration due to gravity on that planet is:

(A) $19.6\,\text{m s}^{-2}$ (B) $9.8\,\text{m s}^{-2}$

(C) $4.9\,\text{m s}^{-2}$ (D) $3.92\,\text{m s}^{-2}$

Open solution

Q9. NEET 2024

The minimum energy required to launch a satellite of mass m from the surface of earth of mass M and radius R in a circular orbit at an altitude of $2R$ from the surface of the earth is:

(A) $\frac{5GmM}{6R}$ (B) $\frac{2GmM}{3R}$ (C) $\frac{GmM}{2R}$ (D) $\frac{GmM}{3R}$

Open solution

Q10. NEET 2023 Manipur

The escape velocity of a body on the earth surface is $11.2\,\text{km/s}$. If the same body is projected upward with velocity $22.4\,\text{km/s}$, the velocity of this body at infinite distance from the centre of the earth will be:

(A) $11.2\sqrt{2}\,\text{km/s}$ (B) Zero

(C) $11.2\,\text{km/s}$ (D) $11.2\sqrt{3}\,\text{km/s}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q11. NEET 2023 Manipur

If R is the radius of the earth and g is the acceleration due to gravity on the earth surface. Then the mean density of the earth will be :

(A) $\frac{\pi \mathrm{RG}}{12\ \mathrm{g}}$ (B) $\frac{3\pi R}{4gG}$

(C) $\frac{3g}{4\pi RG}$ (D) $\frac{4\pi \mathrm{G}}{3gR}$

Open solution

Q12. NEET 2023

Two bodies of mass m and $9m$ are placed at a distance R. The gravitational potential on the line joining the bodies where the gravitational field equals zero, will be (G = gravitational constant) :

(A) $-\frac{12Gm}{R}$ (B) $-\frac{16Gm}{R}$ (C) $-\frac{20Gm}{R}$ (D) $-\frac{8Gm}{R}$

Open solution

Q13. NEET 2023


A satellite is orbiting just above the surface of the earth with period T. If d is the density of the earth and G is the universal constant of gravitation, the quantity $\frac{3\pi}{Gd}$ represents :

(A) T^2 (B) T^3 (C) $\sqrt{T}$ (D) T

Open solution

Q14. NEET 2022 Phase 2

A gravitational field is present in a region and a mass is shifted from A to B through different paths as shown. If W_1, W_2 and W_3 represent the work done by the gravitational force along the respective paths, then :



(A) $W_1 < W_2 < W_3$ (B) $W_1 = W_2 = W_3$ (C) $W_1 > W_2 > W_3$ (D) $W_1 > W_3 > W_2$

Open solution

Q15. NEET 2022 Phase 2

In a gravitational field, the gravitational potential is given by, $V = -\frac{K}{x}$ (J/Kg). The gravitational field intensity at point (2, 0, 3) m is

(A) $+\frac{K}{4}$ (B) $+\frac{K}{2}$ (C) $-\frac{K}{2}$ (D) $-\frac{K}{4}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q16. NEET 2022 Phase 1

A body of mass 60 g experiences a gravitational force of 3.0 N, when placed at a particular point. The magnitude of the gravitational field intensity at that point is

(A) 0.05 N/kg (B) 50 N/kg (C) 20 N/kg (D) 180 N/kg

Open solution

Q17. NEET 2022 Phase 1

Match List - I with List - II

	List - I		List - II
(a)	Gravitational constant (G)	(i)	$[L^2T^{-2}]$
(b)	Gravitational potential energy	(ii)	$[M^{-1}L^3T^{-2}]$
(c)	Gravitational potential	(iii)	$[LT^{-2}]$
(d)	Gravitational intensity	(iv)	$[ML^2T^{-2}]$

Choose the correct answer from the options given below

(A) (a)-(ii), (b)-(i), (c)-(iv), (d)-(iii) (B) (a)-(ii), (b)-(iv), (c)-(i), (d)-(iii)

(C) (a)-(ii), (b)-(iv), (c)-(iii), (d)-(i) (D) (a)-(iv), (b)-(ii), (c)-(i), (d)-(iii)

Open solution

Q18. NEET 2021

The escape velocity from the Earth's surface is v. The escape velocity from the surface of another planet having a radius, four times that of Earth and same mass density is :

(A) $4v$ (B) v (C) $2v$ (D) $3v$

Open solution

Q19. NEET 2021

A particle of mass 'm' is projected with a velocity $v = kV_e = (k < 1)$ from the surface of the earth. (V_e = escape velocity)

The maximum height above the surface reached by the particle is :

(A) $\frac{Rk^2}{1-k^2}$ (B) $R\left(\frac{k}{1-k^2}\right)^2$

(C) $R\left(\frac{k}{1+k^2}\right)^2$ (D) $\frac{R^2k}{1+k}$

Open solution

Q20. NEET 2020 Phase 1

A body weighs 72 N on the surface of the earth. What is the gravitational force on it, at a height equal to half the radius of the earth?

(A) 32 N (B) 30 N (C) 24 N (D) 48 N

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q21. NEET 2019

The work done to raise a mass m from the surface of the earth to a height h, which is equal to the radius of the earth, is :

(A) mgR (B) 2mgR (C) $\frac{1}{2}$mgR (D) $\frac{3}{2}$mgR

Open solution

Q22. NEET 2019


A body weighs 200 N on the surface of earth. How much will it weigh half way down to the centre of the earth?

(A) 200 N (B) 250 N (C) 100 N (D) 150 N

Open solution

Q23. NEET 2018

The kinetic energies of a planet in an elliptical orbit about the Sun, at positions A, B and C are K_A, K_B and K_C, respectively. AC is the major axis and SB is perpendicular to AC at the position of the Sun S as shown in the figure. Then



NEETCARE.com

(A) $K_A < K_B < K_C$ (B) $K_A > K_B > K_C$ (C) $K_B < K_A < K_C$ (D) $K_B > K_A > K_C$

Open solution

Q24. NEET 2018

If the mass of the Sun were ten times smaller and the universal gravitational constant were ten times larger in magnitude, which of the following is not correct?

(A) Raindrops will fall faster.

(B) Walking on the ground would become more difficult.

(C) Time period of a simple pendulum on the Earth would decrease.

(D) g on the Earth will not change.

Open solution

Q25. NEET 2017

The acceleration due to gravity at a height 1 km above the earth is the same as at a depth d below the surface of earth. Then

(A) d = 1 km (B) d = $\frac{3}{2}$ km (C) d = 2 km (D) d = $\frac{1}{2}$ km

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q26. NEET 2017

Two astronauts are floating in gravitational free space after having lost contact with their spaceship. The two will

Open solution

(A) move towards each other.

(B) move away from each other.

(C) will become stationary.


(D) keep floating at the same distance between them.

Q27. NEET 2016 Phase 2


Starting from the centre of the earth having radius R, the variation of g (acceleration due to gravity) is shown by

Open solution


(A)




(B)



(C)



(D)



Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q28. NEET 2016 Phase 2

A satellite of mass m is orbiting the earth (of radius R) at a height h from its surface. The total energy of the satellite in terms of g_0, the value of acceleration due to gravity at the earth's surface, is

(A) $\frac{mg_0R^2}{2(R+h)}$ (B) $-\frac{mg_0R^2}{2(R+h)}$

(C) $\frac{2mg_0R^2}{R+h}$ (D) $-\frac{2mg_0R^2}{R+h}$

Open solution

Q29. NEET 2016 Phase 1

At what height from the surface of earth the gravitation potential and the value of g are -5.4×10^7 J kg^{-1} and 6.0 m s^{-2} respectively? Take the radius of earth as 6400 km.

(A) 1400 km (B) 2000 km (C) 2600 km (D) 1600 km

Open solution

Q30. NEET 2016 Phase 1

The ratio of escape velocity at earth (v_e) to the escape velocity at a planet (v_p) whose radius and mean density are twice as that of earth is

(A) 1 : 4 (B) $1 : \sqrt{2}$ (C) 1 : 2 (D) $1 : 2\sqrt{2}$

Open solution

Q31. AIPMT 2015

A remote-sensing satellite of earth revolves in a circular orbit at a height of 0.25×10^6 m above the surface of earth. If earth's radius is 6.38×10^6 m and g = 9.8 ms^{-2}, then the orbital speed of the satellite is

(A) 9.13 km s^{-1} (B) 6.67 km s^{-1} (C) 7.76 km s^{-1} (D) 8.56 km s^{-1}

Open solution

Q32. AIPMT 2015

A satellite S is moving in an elliptical orbit around the earth. The mass of the satellite is very small compared to the mass of the earth. Then,

(A) the linear momentum of S remains constant in magnitude.

(B) the acceleration of S is always directed towards the centre of the earth.

(C) the angular momentum of S about the centre of the earth changes in direction, but its magnitude remains constant.

(D) the total mechanical energy of S varies periodically with time.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q33. AIPMT 2015 Cancelled Paper

Open solution

Kepler's third law states that square of period of revoluation (T) of a planet around the sun, is proportional to third power of average distance r between sun and planet i.e. $T^2 = Kr^3$ here K is constant. If the masses of sun and planet are M and m respectively then as per Newton's law of gravitation force of attraction between them is $F = \frac{GMm}{r^2}$, here G is gravitational constant. The relation between G and K is described as

(A) K = G (B) $K = \frac{1}{G}$ (C) $GK = 4\pi^2$ (D) $GMK = 4\pi^2$

Q34. AIPMT 2014

Open solution

A black hole is an object whose gravitational field is so strong that even light cannot escape from it. To what approximate radius would earth (mass = 5.98×10^{24} kg) have to be compressed to be a black hole?

(A) 10^{-9} m (B) 10^{-6} m (C) 10^{-2} m (D) 100 m


Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q35. AIPMT 2014


Dependence of intensity of gravitational field (E) of earth with distance (r) from centre of earth is correctly represented by

Open solution


(A)




(B)



(C)



(D)



Q36. NEET 2013 (Karnataka)

The radius of a planet is twice the radius of earth. Both have almost equal average mass densities. V_P and V_E are escape velocities of the planet and the earth, respectively, then

Open solution

(A) $V_P = 1.5\ V_E$ (B) $V_P = 2\ V_E$ (C) $V_E = 3\ V_P$ (D) $V_E = 1.5\ V_P$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q37. NEET 2013 (Karnataka)

A particle of mass 'm' is kept at rest at a height 3R from the surface of earth, where 'R' is radius of earth and 'M' is mass of earth. The minimum speed with which it should be projected , so that it does not return back, is
(g is acceleration due to gravity on the surface of earth)

Open solution

(A) $\left(\frac{GM}{2R}\right)^{1/2}$ (B) $\left(\frac{gR}{4}\right)^{1/2}$
(C) $\left(\frac{2g}{R}\right)^{1/2}$ (D) $\left(\frac{GM}{R}\right)^{1/2}$

Q38. NEET 2013

Infinite number of bodies, each of mass 2 kg are situated on x-axis at distance 1 m, 2 m, 4 m, 8 m, . . . , respectively, from the origin. The resulting gravitational potential due to this system at the origin will be

Open solution

(A) $-\frac{4}{3}$ G (B) $-$ 4G (C) $-$ G (D) $-\frac{8}{3}$ G

Q39. NEET 2013

A body of mass 'm' is taken from the earth's surface to the height equal to twice the radius (R) of the earth. The change in potential energy of body will be

Open solution

(A) 3mgR (B) $\frac{1}{3}$mgR (C) mg2R (D) $\frac{2}{3}$ mgR

Q40. AIPMT 2012 Mains

If v_e is escape velocity and v_o is orbital velocity of a satellite for orbit close to the earth's surface, then these are related by

Open solution

(A) $v_o = \sqrt{2v_e}$ (B) $v_o = v_e$ (C) $v_e = \sqrt{2v_o}$ (D) $v_e = \sqrt{2}v_o$


Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q41. AIPMT 2012 Mains


Which one of the following plots represents the variation of gravitiational field on a particle with distance r due to a thin spherical shell of radius R? (r is measured from the centre of the spherical shell)

Open solution


(A)




(B)



(C)



(D)



Q42. AIPMT 2012 Prelims

The height at which the weight of a body becomes $\left(\frac{1}{16}\right)^{th}$, its weight on the surface of earth (radius R), is

Open solution

(A) 5R (B) 15R (C) 3R (D) 4R

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q43. AIPMT 2012 Prelims

A spherical planet has a mass M_P and diameter D_P. A particle of mass m falling freely near the surface of this planet will experience an acceleration due to gravity, equal to

(A) $\frac{4GM_P}{D_P^2}$ (B) $\frac{GM_Pm}{D_P^2}$ (C) $\frac{GM_P}{D_P^2}$ (D) $\frac{4GM_Pm}{D_P^2}$

Open solution

Q44. AIPMT 2012 Prelims

A geostationary satellite is orbiting the earth at a height of 5R above the surface of the earth, R being the radius of the earth. The time period of another satellite in hours at a height of 2R from the surface of the earth is

(A) 5 (B) 10 (C) $6\sqrt{2}$ (D) $\frac{6}{\sqrt{2}}$

Open solution

Q45. AIPMT 2011 Mains

A particle of mass m is thrown upwards from the surface of the earth, with a velocity u. The mass and the radius of the earth are, respectively, M and R. G is gravitational constant and g is acceleration due to gravity on the surface of the earth. The minimum value of u so that the particle does not return back to earth, is

(A) $\sqrt{\frac{2GM}{R^2}}$ (B) $\sqrt{\frac{2GM}{R}}$ (C) $\sqrt{\frac{2gM}{R^2}}$ (D) $\sqrt{2gR^2}$

Open solution

Q46. AIPMT 2011 Mains

A particle of mass M is situated at the centre of a spherical shell of same mass and radius a. The magnitude of the gravitational potential at a point situated at a/2 distance from the centre, will be :

(A) $\frac{GM}{a}$ (B) $\frac{2GM}{a}$ (C) $\frac{3GM}{a}$ (D) $\frac{4GM}{a}$

Open solution

Q47. AIPMT 2011 Prelims

A planet moving along an elliptical orbit is closest to the sun at a distance r_1 and farthest away at a distance of r_2. If v_1 and v_2 are the linear velocities at these points respectively, then the ratio $\frac{v_1}{v_2}$ is

(A) $(r_1/r_2)^2$ (B) r_2/r_1 (C) $(r_2/r_1)^2$ (D) r_1/r_2

Open solution

Q48. AIPMT 2010 Mains

The additional kinetic energy to be provided to a satellite of mass m revolving around a planet of mass M, to transfer it from a circular orbit of radius R_1 to another of radius $R_2 (R_2 > R_1)$ is

(A) $GmM\left(\frac{1}{R_1^2}-\frac{1}{R_2^2}\right)$ (B) $GmM\left(\frac{1}{R_1}-\frac{1}{R_2}\right)$

(C) $2GmM\left(\frac{1}{R_1}-\frac{1}{R_2}\right)$ (D) $\frac{1}{2}GmM\left(\frac{1}{R_1}-\frac{1}{R_2}\right)$


Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q49. AIPMT 2010 Mains

The dependence of acceleration due to gravity g on the distance r from the centre of the earth, assumed to be a sphere of radius R of uniform density is as shown in figures below

Open solution



The correct figure is

(A) (4) (B) (1) (C) (2) (D) (3)

Q50. AIPMT 2010 Mains

Open solution

(1) Centre of gravity (C.G) of a body is the point at which the weight of the body acts
(2) Centre of mass coincides with the centre of gravity if the earth is assumed to have infinitely large radius.
(3) To evaluate the gravitational field intensity due to any body at an external point, the entire mass of the body can be considered to be concentrated at its C.G.
(4) The radius of gyration of any body rotating about an axis is the length of the perpendicular dropped from the C.G. of the body to the axis.

Which one of the following pairs of statements is correct ?

(A) (4) and (1) (B) (1) and (2) (C) (2) and (3) (D) (3) and (4)

Q51. AIPMT 2010 Prelims

A particle of mass M is situated at the centre of a spherical shell of same mass and radius a. The magnitude of the gravitational potential at a point sutuated at a/2 distance from the centre, will be :

Open solution

(A) $\frac{GM}{a}$ (B) $\frac{2GM}{a}$ (C) $\frac{3GM}{a}$ (D) $\frac{4GM}{a}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q52. AIPMT 2010 Prelims

The radii of circular orbits of two satellites A and B of the earth, are 4R and R, respectively. If the speed of satellite A is 3V, then the speed of satellite B will be

(A) $\frac{3V}{4}$ (B) $6V$ (C) $12V$ (D) $\frac{3V}{2}$

Open solution

Q53. AIPMT 2010 Prelims


A man of 50 kg mass is standing in a gravity free space at a height of 10 m above the floor. He throws a stone of 0.5 kg mass downwards with a speed 2 m/s. When the stone reaches the floor, the distance of the man above the floor will be

(A) 9.9 m (B) 10.1 m (C) 10 m (D) 20 m

Open solution

Q54. AIPMT 2009

The figure shows elliptical orbit of a planet m about the sun S. The shaded area SCD is twice the shaded area SAB. If t_1 is the time for the planet to move from C to D and t_2 is the time to move from A to B then



(A) $t_1 = 4t_2$ (B) $t_1 = 2t_2$ (C) $t_1 = t_2$ (D) $t_1 > t_2$

Open solution

Q55. AIPMT 2007

Two satellites of earth, S_1 and S_2 are moving in the same orbit. The mass of S_1 is four times the mass of S_2. Which one of the following statements is true ?

(A) The potential energies of earth and satellite in the two cases are equal.

(B) S_1 and S_2 are moving with the same speed.

(C) The kinetic energies of the two satellites are equal.

(D) The time period of S_1 is four times that of S_2.

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q56. AIPMT 2006

The earth is assumed to be a sphere of radius R. A platform is arranged at a height R from the surface of the earth. The escape velocity of a body from this platform is fv, where v is its escape velocity from the surface of the Earth. The value of f is

(A) 1/2 (B) $\sqrt{2}$ (C) $1/\sqrt{2}$ (D) 1/3

Open solution

Q57. AIPMT 2005

Imagine a new planet having the same density as that of earth but it is 3 times bigger than the earth in size. If the acceleration due to gravity on the surface of earth is g and that on the surface of the new planet is g', then

(A) g' = g/9 (B) g' = 27g (C) g' = 9g (D) g' = 3g

Open solution

Q58. AIPMT 2005

For a satellite moving in an orbit around the earth, the ratio of kinetic energy to potential energy is

(A) 1/2 (B) $1/\sqrt{2}$ (C) 2 (D) $\sqrt{2}$

Open solution

Q59. AIPMT 2004

The density of a newly discovered planet is twice that of earth. The acceleration due to gravity at the surface of the planet is equal to that at the surface of the earth. If the radius of the earth is R, the radius of the planet would be

(A) 2R (B) 4R (C) $\frac{1}{4}$R (D) $\frac{1}{2}$R

Open solution

Q60. AIPMT 2003

Two spheres of masses m and M are situated in air and the gravitational force between them is F. The space around the masses is now filled with a liquid of specific gravity 3. The gravitational force will now be

(A) 3 F (B) F (C) F/3 (D) F/9

Open solution

Q61. AIPMT 2003

The acceleration due to gravity on the planet A is 9 times the acceleration due to gravity on planet B. A man jumps to a height of 2 m on the surface of A. What is the height of jump by the same person on the planet B?

(A) (2/9) m (B) 18 m (C) 6 m (D) (2/3) m

Open solution

Q62. AIPMT 2003

A body of mass m is placed on earth's surface which is taken from earth surface to a height of h = 3R, then change in gravitational potential energy is

(A) $\frac{mgR}{4}$ (B) $\frac{2}{3}mgR$ (C) $\frac{3}{4}mgR$ (D) $\frac{mgR}{2}$

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q63. AIPMT 2001

With what velocity should a particle be projected so that its height becomes equal to radius of earth?

(A) $\left(\frac{GM}{R}\right)^{1/2}$ (B) $\left(\frac{8GM}{R}\right)^{1/2}$

(C) $\left(\frac{2GM}{R}\right)^{1/2}$ (D) $\left(\frac{4GM}{R}\right)^{1/2}$

Open solution

Q64. AIPMT 2000

For a planet having mass equal to mass of the earth but radius is one fourth of radius of the earth. Then escape velocity for this planet will be

(A) 11.2 km/sec (B) 22.4 km/sec (C) 5.6 km/sec (D) 44.8 km/sec.

Open solution

Q65. AIPMT 2000

Gravitational force is required for

(A) stirring of liquid (B) convection

(C) conduction (D) radiation.

Open solution

Q66. AIPMT 2000

A body of weight 72 N moves from the surface of earth at a height half of the radius of earth, then gravitational force exerted on it will be

(A) 36 N (B) 32 N (C) 144 N (D) 50 N

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Answer Key

1. **D**	2. **C**	3. **D**	4. **C**	5. **D**
6. **C**	7. **A**	8. **D**	9. **A**	10. **D**
11. **C**	12. **B**	13. **A**	14. **B**	15. **D**
16. **B**	17. **B**	18. **A**	19. **A**	20. **A**
21. **C**	22. **C**	23. **B**	24. **D**	25. **C**
26. **A**	27. **B**	28. **B**	29. **C**	30. **D**
31. **C**	32. **B**	33. **D**	34. **C**	35. **A**
36. **B**	37. **A**	38. **B**	39. **D**	40. **D**
41. **B**	42. **C**	43. **A**	44. **C**	45. **B**
46. **C**	47. **B**	48. **D**	49. **A**	50. **A**
51. **C**	52. **B**	53. **B**	54. **B**	55. **B**
56. **C**	57. **D**	58. **A**	59. **D**	60. **B**
61. **B**	62. **C**	63. **A**	64. **B**	65. **B**
66. **B**				

For numerical answers given as a range, any value within the range is correct. Full solutions: scan the QR code next to each question, or visit questions.examside.com.

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

ExamGOAL
PYQ PDF

Your Exam. Our Responsibility.

From first chapter to exam day.

Practise with the questions your exam has already asked, then test yourself under real exam conditions.

PAST YEAR QUESTIONS

Every past paper, chapter-wise.

Previous year questions sorted the way you study, each with a full solution.

- Chapter-wise and year-wise
- Weightage and topic tags on every chapter
- Your progress on each one, at a glance

MOCK TESTS

Mock tests with a real exam feel.

Practise under the same pressure as the real paper, on your phone or on the big screen.

- Timed sections, just like exam day
- Mark questions for review and come back

ANALYSIS

Know your score. Know your rank.

A full report after every test, so you know exactly where you stand.

- Your percentile among everyone who took it
- Predicted real-exam percentile and All India Rank
- Subject breakdown and time analysis

PRACTICE

Solve. Check. Understand.

Learn from every question, not just the ones you get right.

- Instant answers as you practise
- Step-by-step explanations
- See how everyone else answered

NCERT Exemplar
With full solutions

Exam syllabus
Subject and chapter-wise

Bookmarks
Organised in groups

Your notes
On any question

ExamGOAL
LEARNING APP

ExamGOAL: Exam Preparation App

One app for every exam. Free trial for new users, on Android and the web.

GET IT ON Google Play

Scan to get the app

www.examgoal.com
Practice and mock tests

questions.examside.com
Free solutions

pdf.questions.examgoal.com
Free PYQ PDFs