ExamGOAL
PYQ PDF

Free · With answer key

NEET · PHYSICS · CHAPTER-WISE PYQS

Laws of Motion

NEET Physics · Previous Year Questions with Answer Key

53	30	2000 – 2026	English
QUESTIONS	PAPERS COVERED	YEARS	LANGUAGE

INSIDE THIS PDF

MCQ (Single Correct Answer)	53 questions
Answer key	at the end

Practise online

Every question with its full solution, free on **questions.examside.com**

1. **Solve**
   Print it, or work through it on screen.
2. **Check**
   The answer key is on the last pages.
3. **Understand**
   Scan the QR code beside any question for its full solution.

MCQ (Single Correct Answer) 53 questions

Q1. RE-NEET 2026

A car travels on a circular racetrack of radius 50 m , which is banked at an angle θ. If the car travels at a speed $10\ \mathrm{ms}^{-1}$, then the wear and tear on its tyres is minimum. Taking the acceleration due to gravity to be $10\ \mathrm{ms}^{-2}$, the value of θ is:

(A) $\tan^{-1}(2\sqrt{3})$ (B) $\tan^{-1}\left(\frac{1}{5}\right)$

(C) $\tan^{-1}\left(\frac{2}{5}\right)$ (D) $\tan^{-1}(\sqrt{3}/2)$

Open solution

Q2. NEET 2026

A box of mass 15 kg is kept on the floor of a stationary trolley. The coefficient of static friction between the box and the trolley is 0.12 . Keeping the box in stationary state over the trolley, the maximum acceleration with which the trolley can be moved horizontally in ms^{-2} is:

$\left(g = 10\ \mathrm{m/s}^2\right)$

(A) 2.1 (B) 1.8 (C) 1.5 (D) 1.2

Open solution

Q3. NEET 2026

The magnitude and direction of the acceleration produced in a body of mass 5 kg when two mutually perpendicular forces 8 N and 6 N act on it, are respectively:

(A) $20\ \mathrm{m\,s}^{-2}; \tan^{-1}(4/3)$ with 8 N force

(B) $2\ \mathrm{m\,s}^{-2}; \tan^{-1}(3/4)$ with 6 N force

(C) $2\ \mathrm{m\,s}^{-2}; \tan^{-1}(4/3)$ with 8 N force

(D) $2\ \mathrm{m\,s}^{-2}; \tan^{-1}(3/4)$ with 8 N force

Open solution

Q4. NEET 2025

There are two inclined surfaces of equal length (L) and same angle of inclination 45° with the horizontal. One of them is rough and the other is perfectly smooth. A given body takes 2 times as much time to slide down on rough surface than on the smooth surface. The coefficient of kinetic friction (μ_k) between the object and the rough surface is close to

(A) 0.5 (B) 0.75 (C) 0.25 (D) 0.40

Open solution

Q5. NEET 2024 (Re-Examination)

A box of mass $5\ \mathrm{kg}$ is pulled by a cord, up along a frictionless plane inclined at 30° with the horizontal. The tension in the cord is $30\ \mathrm{N}$. The acceleration of the box is (Take $g = 10\ \mathrm{m\,s}^{-2}$)

(A) $2\ \mathrm{m\,s}^{-2}$ (B) Zero (C) $0.1\ \mathrm{m\,s}^{-2}$ (D) $1\ \mathrm{m\,s}^{-2}$


Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q6. NEET 2024

A horizontal force $10\ \text{N}$ is applied to a block A as shown in figure. The mass of blocks A and B are $2\ \text{kg}$ and $3\ \text{kg}$ respectively. The blocks slide over a frictionless surface. The force exerted by block A on block B is :

Open solution



(A) Zero (B) 4 N (C) 6 N (D) 10 N

Q7. NEET 2024

A particle moving with uniform speed in a circular path maintains:

(A) Constant velocity

(B) Constant acceleration

(C) Constant velocity but varying acceleration


(D) Varying velocity and varying acceleration

Open solution

Q8. NEET 2023 Manipur

A block of mass 2 kg is placed on inclined rough surface AC (as shown in figure) of coefficient of friction μ. If g = 10 m s^{-2}, the net force (in N) on the block will be:

Open solution



(A) $10\sqrt{3}$ (B) zero (C) 10 (D) 20

Q9. NEET 2023

A football player is moving southward and suddenly turns eastward with the same speed to avoid an opponent. The force that acts on the player while turning is :

(A) along northward (B) along north-east (C) along south-west (D) along eastward

Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q10. NEET 2023

Calculate the maximum acceleration of a moving car so that a body lying on the floor of the car remains stationary. The coefficient of static friction between the body and the floor is 0.15 $\left(g = 10\ \mathrm{m\,s^{-2}}\right)$.

Open solution

(A) $150\ \mathrm{m\,s^{-2}}$ (B) $1.5\ \mathrm{m\,s^{-2}}$ (C) $50\ \mathrm{m\,s^{-2}}$ (D) $1.2\ \mathrm{m\,s^{-2}}$

Q11. NEET 2023

A bullet from a gun is fired on a rectangular wooden block with velocity u. When bullet travels $24\ \mathrm{cm}$ through the block along its length horizontally, velocity of bullet becomes $\frac{u}{3}$. Then it further penetrates into the block in the same direction before coming to rest exactly at the other end of the block. The total length of the block is :

Open solution


(A) $24\ \mathrm{cm}$ (B) $28\ \mathrm{cm}$ (C) $30\ \mathrm{cm}$ (D) $27\ \mathrm{cm}$

Q12. NEET 2022 Phase 2

In the diagram shown, the normal reaction force between 2 kg and 1 kg is (Consider the surface, to be smooth) :

(Given g = 10 $\mathrm{ms^{-2}}$)

Open solution




(A) 10 N (B) 25 N (C) 39 N (D) 6 N

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q13. NEET 2020 Phase 1

Two bodies of mass 4 kg and 6 kg are tied to the ends of a massless string. The string passes over a pulley which is frictionless (see figure). The acceleration of the system in terms of acceleration due to gravity (g) is :

Open solution



(A) g/2 (B) g/5 (C) g/10 (D) g

Q14. NEET 2019

A block of mass 10 kg is in contact against the inner wall of a hollow cylindrical drum of radius 1 m. The coefficient of friction between the block and the inner wall of the cylinder is 0.1. The minimum angular velocity needed for the cylinder to keep the block stationary when the cylinder is vertical and rotating about its axis, will be : (g = 10 m/s^2)

Open solution

(A) $\frac{10}{2\pi}$ rad/s (B) 10 rad/s (C) 10 π rad/s (D) $\sqrt{10}$ rad/s

Q15. NEET 2019

A mass m is attached to a thin wire and whirled in a vertical circle. The wire is most likely to break when :

Open solution

(A) the wire is horizontal

(B) inclined at an angle of 60° from vertical

(C) the mass is at the lowest point

(D) the mass is at the highest point

Q16. NEET 2018

Which one of the following statements is incorrect?

Open solution

(A) Rolling friction is smaller than sliding friction.

(B) Limiting value of static friction is directly proportional to normal reaction.

(C) Frictional force opposes the relative motion.


(D) Coefficient of sliding friction has dimensions of length.

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q17. NEET 2018

A block of mass m is placed on a smooth inclined wedge ABC of inclination θ as shown in the figure. The wedge is given an acceleration 'a' towards the right. The relation between a and θ for the block to remain stationary on the wedge is

Open solution



(A) $a = \frac{g}{\cos ec\theta}$ (B) $a = \frac{g}{\sin\theta}$ (C) $a = g\cos\theta$ (D) $a = g\tan\theta$

Q18. NEET 2017

One end of string of length l is connected to a particle of mass 'm' and the other end is connected to a small peg on a smooth horizontal table. If the particle moves in circle with speed 'v', the net force on the particle (directed towards centre) will be (T represents the tension in the string)


Open solution

(A) $T + \frac{mv^2}{l}$ (B) $T - \frac{mv^2}{l}$ (C) zero (D) T

Q19. NEET 2017

Two blocks A and B of masses 3m and m respectively are connected by a massless and inextensible string. The whole system is suspended by a massless spring as shown in figure. The magnitudes of acceleration of A and B immediately after the string is cut, are respectively

Open solution



(A) $\frac{g}{3}, g$ (B) g, g (C) $\frac{g}{3}, \frac{g}{3}$ (D) $g, \frac{g}{3}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q20. NEET 2016 Phase 1

A car is negotiating a curved road of radius R. The road is banked at an angle θ. The coefficient of friction between the tyres of the car and the road is μ_s. The maximum safe velocity on this road is

Open solution

(A) $\sqrt{\frac{g}{R}\frac{\mu_s+\tan\theta}{1-\mu_s\tan\theta}}$

(B) $\sqrt{\frac{g}{R^2}\frac{\mu_s+\tan\theta}{1-\mu_s\tan\theta}}$

(C) $\sqrt{gR^2\frac{\mu_s+\tan\theta}{1-\mu_s\tan\theta}}$

(D) $\sqrt{gR\frac{\mu_s+\tan\theta}{1-\mu_s\tan\theta}}$

Q21. AIPMT 2015

Two stones of masses m and 2m are whirled in horizontal circles, the heavier one in a radius $\frac{r}{2}$ and the lighter one in radius r. The tangential speed of lighter stone is n times that of the value of heavier stone when they exprience same centripetal forces. The value of n is


Open solution

(A) 4 (B) 1 (C) 2 (D) 3

Q22. AIPMT 2015

A plank with a box on it at one end is gradually raised about the other end. As the angle of inclination with the horizontal reaches 30°, the box starts to slip and slides 4.0 m down the plank in 4.0 s.

Open solution



The coefficients of static and kinetic friction between the box and the plank will be, respectively

(A) 0.5 and 0.6 (B) 0.4 and 0.3 (C) 0.6 and 0.6 (D) 0.6 and 0.5

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q23. AIPMT 2015 Cancelled Paper

A block A of mass m_1 rests on a horizontal table. A light string connected to it passes over a frictionless pully at the edge of table and from its other end another block B of mass m_2 is suspended. The coefficient of kinetic friction between the block and the table is μ_k. When the block A is sliding on the table, the tension in the string is

Open solution

(A) $\frac{m_1 m_2(1+\mu_k)g}{(m_1+m_2)}$

(B) $\frac{m_1 m_2(1-\mu_k)g}{(m_1+m_2)}$

(C) $\frac{(m_2+\mu_k m_1)g}{(m_1+m_2)}$

(D) $\frac{(m_2-\mu_k m_1)g}{(m_1+m_2)}$

Q24. AIPMT 2015 Cancelled Paper

Three blocks A, B and C, of masses 4 kg, 2 kg and 1 kg respectively, are in contact on a frictionless surface, as shown. If a force of 14 N is applied on the 4 kg block, then the contact force between A and B is

Open solution



(A) 8 N (B) 18 N (C) 2 N (D) 6 N

Q25. AIPMT 2014

A balloon with mass m is descending down with an acceleration a (where a < g). How much mass should be removed from it so that it starts moving up with an acceleration a ?

Open solution


(A) $\frac{2ma}{g+a}$ (B) $\frac{2ma}{g-a}$ (C) $\frac{ma}{g+a}$ (D) $\frac{ma}{g-a}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q26. AIPMT 2014

Open solution

A system consists of three masses m_1, m_2 and m_3 connected by a string passing over a pulley P. The mass m_1 hangs freely and m_2 and m_3 are on a rough horizontal table (the coefficient of friction = μ). The pulley is frictionless and of negligible mass. The downward acceleration of mass m_1 is (Assume $m_1 = m_2 = m_3 = m$)



NEETCARE.Com

(A) $\frac{g(1-g\mu)}{9}$

(B) $\frac{2g\mu}{3}$

(C) $\frac{g(1-2\mu)}{3}$

(D) $\frac{g(1-2\mu)}{2}$

Q27. NEET 2013 (Karnataka)

Open solution

A car is moving in a circular horizontal track of radius 10 m with a constant speed of 10 m/s. A bob is suspended from the roof of the car by a light wire of length 1.0 m. The angle made by the wire with the vertical is

(A) $\frac{\pi}{3}$

(B) $\frac{\pi}{6}$

(C) $\frac{\pi}{4}$

(D) 0°

Q28. NEET 2013

Open solution

The upper half of an inclined plane of inclination θ is perfectly smooth while lower half is rough. A block starting from rest at the top of the plane will again come to rest at the bottom, if the coefficient of friction between the block and lower half of the plane is given by

(A) $\mu = 2\tan\theta$

(B) $\mu = \tan\theta$

(C) $\mu = \frac{1}{\tan\theta}$


(D) $\mu = \frac{2}{\tan\theta}$

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q29. NEET 2013

Three blocks with masses m, 2m and 3m are connected by strings, as shown in the figure. After an upward force F is applied on block m, the masses move upwards at constant speed v. What is the net force on the block of mass 2m ? (g is the acceleration due to gravity)

Open solution



(A) 3mg (B) 6mg (C) zero (D) 2mg

Q30. AIPMT 2012 Mains

A stone is dropped from a height h. It hits the ground with a certain momentum P. If the same stone is dropped from a height 100% more than the previous height, the momentum when it hits the ground will change by

Open solution

(A) 68% (B) 41% (C) 200% (D) 100%

Q31. AIPMT 2011 Mains

A conveyor belt is moving at a constant speed of 2 ms^{-1}. A box is gently dropped on it. The coefficient of friction between them is μ = 0.5. The distance that the box will move relativce to belt before coming to rest on it, taking g = 10 m s^{-2}, is

Open solution

(A) 0.4 m (B) 1.2 m (C) 0.6 m (D) zero

Q32. AIPMT 2011 Prelims

A person of mass 60 kg is inside a lift of mass 940 kg and presses the button on control panel. The lift starts moving upwards with an acceleration 1.0 m/s^2. If g = 10 ms^{-2}, the tension in the supporting cable is

Open solution

(A) 8600 N (B) 9680 N (C) 11000 N (D) 1200 N

Q33. AIPMT 2011 Prelims

A body of mass M hits normally a rigid wall with velocity V and bounces back with the same velocity. The impulse experienced by the body is

Open solution

(A) MV (B) 1.5 MV (C) 2MV (D) zero


Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q34. AIPMT 2010 Prelims

A block of mass m is in contact with the cart C as shown in the figure.

The coefficient of static friction between the block and the cart is μ. The acceleration α of the cart that will prevent the block from falling satisfies

Open solution



(A) $\alpha > \frac{mg}{\mu}$ (B) $\alpha > \frac{g}{\mu m}$ (C) $\alpha \geq \frac{g}{\mu}$ (D) $\alpha < \frac{g}{\mu}$

Q35. AIPMT 2009

The mass of a lift is 2000 kg. When the tension in the supporting cable is 28000 N, then its acceleration is :

Open solution

(A) 4 m s^{-2} upwards (B) 4 m s^{-2} downwards

(C) 14 m s^{-2} upwards (D) 30 m s^{-2} downwards

Q36. AIPMT 2009

A body, under the action of a force $\vec{F} = 6\hat{i} - 8\hat{j} + 10\hat{k}$, acquires an accelerations of 1 m/s^2. The mass of this body must be

Open solution

(A) 10 kg (B) 20 kg (C) $10\sqrt{2}$ kg (D) $2\sqrt{10}$ kg

Q37. AIPMT 2008

A roller coaster is designed such that riders experience "weightlessness" as they go round the top of a hill whose radius of curvature is 20 m. The speed of the car at the top of the hill is between

Open solution

(A) 16 m/s and 17 m/s (B) 13 m/s and 14 m/s

(C) 14 m/s and 15 m/s (D) 15 m/s and 16 m/s

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play


Q38. AIPMT 2008

Sand is being dropped on a conveyer belt at the rate of M kg/s. The force necessary to keep the belt moving with a constant velocity of v m/s will be :

(A) $\frac{Mv}{2}$ newton (B) zero (C) Mv newton (D) 2 Mv newton

Open solution

Q39. AIPMT 2008




Open solution

Three forces acting on a body are shown in the figure. To have the resultant force only along the y-direction, the magnitude of the minimum additional force needed is

(A) $\frac{\sqrt{3}}{4}N$ (B) $\sqrt{3}$ N (C) 0.5 N (D) 1.5 N

Q40. AIPMT 2007

A block B is pushed momentarily along a horizontaly surface with an initial velocity V. If μ is the coefficient of sliding friction between B and the surface, block B will come to rest after a time



Open solution

(A) gμ/V (B) g/V (C) V/g (D) V/(gμ).

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q41. AIPMT 2004

A block of mass m is placed on a smooth wedge of inclination θ. The whole system is accelerated horizontally so that the block does not slip on the wedge. The force exerted by the wedge on the block will be (g is acceleration due to gravity)


Open solution

(A) mg cosθ (B) mg sinθ (C) mg (D) mg/cos θ

Q42. AIPMT 2004

The coefficient of static friction, μ_s, between block A of mass 2 kg and the table as shown in the figure is 0.2. What would be the maximum mass value of block B so that the two blocks do not move? The string and the pulley are assumed to be smooth and massless. (g = 10 m/s^2)

Open solution



(A) 2.0 kg (B) 4.0 kg (C) 0.2 kg (D) 0.4 kg

Q43. AIPMT 2003

A man weight 80 kg. He stands on a weighting scale in a lift which is moving upwards with a uniform acceleration of 5 m/s^2. What would be the reading on the scale ? (g = 10 m/s^2)

Open solution

(A) zero (B) 400 N (C) 800 N (D) 1200 N

Q44. AIPMT 2003

A monkey of mass 20 kg is holding a vertical rope. The rope will not break when a mass of 25 kg is suspended from it but will break if the mass exceeds 25 kg. What is the maximum acceleration with which the monkey can climb up along the rope ? (g = 10 m/s^2)

Open solution

(A) 5 m/s^2 (B) 10 m/s^2 (C) 25 m/s^2 (D) 2.5 m/s^2

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q45. AIPMT 2002

An object of mass 3 kg is at rest. Now a force of $\vec{F} = 6t^2\,\hat{i} + 4t\,\hat{j}$ is applied on the object then velocity of object at t = 3 sec. is

(A) $18\hat{i} + 3\hat{j}$ (B) $18\hat{i} + 6\hat{j}$ (C) $3\hat{i} + 18\hat{j}$ (D) $18\hat{i} + 4\hat{j}$

Open solution

Q46. AIPMT 2002

A lift of mass 1000 kg which is moving with acceleration of 1 m/s^2 in upward direction, then the tension developed in string which is connected to lift is

(A) 9800 N (B) 10,800 N (C) 11,000 N (D) 10,000 N.

Open solution

Q47. AIPMT 2002

A block of mass 10 kg placed on rough horizontal surface having coefficient of friction μ = 0.5, if a horizontal force of 100 N acting on it then acceleration of the block will be

(A) 10 m/s^2 (B) 5 m/s^2 (C) 15 m/s^2 (D) 0.5 m/s^2.

Open solution

Q48. AIPMT 2001

250 N force is required to raise 75 kg mass from a pulley. If rope is pulled 12 m then the load is lifted to 3 m. the efficiency of pulley system will be

(A) 25% (B) 33.3% (C) 75% (D) 90%

Open solution

Q49. AIPMT 2001

On the horizontal surface of a truck a block of mass 1 kg is placed (μ = 0.6) and truck is moving with acceleration 5 m/sec^2 then the frictional force on the block will be

(A) 5 N (B) 6 N (C) 5.88 N (D) 8 N

Open solution

Q50. AIPMT 2001

A cricketer catches a ball of mass 150 gm in 0.1 sec moving with speed 20 m/s, then he experiences force of

(A) 300 N (B) 30 N (C) 3 N (D) 0.3 N.

Open solution

Q51. AIPMT 2001

A 1 kg stationary bomb is exploded in three parts having mass 1 : 1 : 3 respectively. Parts having same mass move in perpendicular direction with velocity 30 m/s, then the velocity of bigger part will be

(A) $10\sqrt{2}$ m/sec (B) $\frac{10}{\sqrt{2}}$ m/sec

(C) $15\sqrt{2}$ m/sec (D) $\frac{15}{\sqrt{2}}$ m/sec


Open solution

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Q52. AIPMT 2000

A body of mass 3 kg hits a wall at an angle of 60° and returns at the same angle. The impact time was 0.2 sec. The force exerted on the wall

Open solution




(A) $150\sqrt{3}$ N (B) $50\sqrt{3}$ N (C) 100 N (D) $75\sqrt{3}$ N.

Q53. AIPMT 2000

Two masses as shown in the figure are suspended from a massless pulley. The acceleration of the system when masses are left free is

Open solution



(A) $\frac{2g}{3}$ (B) $\frac{g}{3}$ (C) $\frac{g}{9}$ (D) $\frac{g}{7}$.

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

Answer Key

1. **B**	2. **D**	3. **D**	4. **B**	5. **D**
6. **C**	7. **D**	8. **B**	9. **B**	10. **B**
11. **D**	12. **B**	13. **B**	14. **B**	15. **C**
16. **D**	17. **D**	18. **D**	19. **A**	20. **D**
21. **C**	22. **D**	23. **A**	24. **D**	25. **A**
26. **C**	27. **C**	28. **A**	29. **C**	30. **B**
31. **A**	32. **C**	33. **C**	34. **C**	35. **A**
36. **C**	37. **C**	38. **C**	39. **C**	40. **D**
41. **D**	42. **D**	43. **D**	44. **D**	45. **B**
46. **B**	47. **B**	48. **C**	49. **A**	50. **B**
51. **A**	52. **A**	53. **B**		

For numerical answers given as a range, any value within the range is correct. Full solutions: scan the QR code next to each question, or visit questions.examside.com.

Keep practising on the ExamGOAL app
Chapter-wise PYQs with solutions, full mock tests and analysis for JEE, NEET, GATE and more.
GET IT ON Google Play

ExamGOAL
PYQ PDF

Your Exam. Our Responsibility.

From first chapter to exam day.

Practise with the questions your exam has already asked, then test yourself under real exam conditions.

PAST YEAR QUESTIONS

Every past paper, chapter-wise.

Previous year questions sorted the way you study, each with a full solution.

- Chapter-wise and year-wise
- Weightage and topic tags on every chapter
- Your progress on each one, at a glance

MOCK TESTS

Mock tests with a real exam feel.

Practise under the same pressure as the real paper, on your phone or on the big screen.

- Timed sections, just like exam day
- Mark questions for review and come back

ANALYSIS

Know your score. Know your rank.

A full report after every test, so you know exactly where you stand.

- Your percentile among everyone who took it
- Predicted real-exam percentile and All India Rank
- Subject breakdown and time analysis

PRACTICE

Solve. Check. Understand.

Learn from every question, not just the ones you get right.

- Instant answers as you practise
- Step-by-step explanations
- See how everyone else answered

NCERT Exemplar
With full solutions

Exam syllabus
Subject and chapter-wise

Bookmarks
Organised in groups

Your notes
On any question

ExamGOAL
LEARNING APP

ExamGOAL: Exam Preparation App

One app for every exam. Free trial for new users, on Android and the web.

GET IT ON Google Play

Scan to get the app

www.examgoal.com
Practice and mock tests

questions.examside.com
Free solutions

pdf.questions.examgoal.com
Free PYQ PDFs