



---

### A Simple Proof of Descartes's Rule of Signs

Author(s): Xiaoshen Wang

Source: *The American Mathematical Monthly*, Vol. 111, No. 6 (Jun. - Jul., 2004), pp. 525-526

Published by: Mathematical Association of America

Stable URL: <http://www.jstor.org/stable/4145072>

Accessed: 07/09/2009 22:01

---

Your use of the JSTOR archive indicates your acceptance of JSTOR's Terms and Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>. JSTOR's Terms and Conditions of Use provides, in part, that unless you have obtained prior permission, you may not download an entire issue of a journal or multiple copies of articles, and you may use content in the JSTOR archive only for your personal, non-commercial use.

Please contact the publisher regarding any further use of this work. Publisher contact information may be obtained at <http://www.jstor.org/action/showPublisher?publisherCode=maa>.

Each copy of any part of a JSTOR transmission must contain the same copyright notice that appears on the screen or printed page of such transmission.

JSTOR is a not-for-profit organization founded in 1995 to build trusted digital archives for scholarship. We work with the scholarly community to preserve their work and the materials they rely upon, and to build a common research platform that promotes the discovery and use of these resources. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).



Mathematical Association of America is collaborating with JSTOR to digitize, preserve and extend access to *The American Mathematical Monthly*.

<http://www.jstor.org>

We can now recover the three classical inequalities cited in section 2 from the results of section 3 by taking the limit as  $\alpha \rightarrow -1^+$ . In fact, it is clear that the Fejér-Riesz inequality (Theorem 1) follows from Theorem 4, Proposition 8, and Fatou's lemma; Hardy's inequality (Theorem 2) is a consequence of Theorem 5, Proposition 8, and Fatou's lemma; and the Hardy-Littlewood inequality (Theorem 3) can be deduced from Theorem 6, Proposition 8, and Fatou's lemma.

## REFERENCES

1. P. Duren, *Theory of  $H^p$  Spaces*, Academic Press, New York, 1970.
2. J. Garnett, *Bounded Analytic Functions*, Academic Press, New York, 1981.
3. H. Hedenmalm, B. Korenblum, and K. Zhu, *Theory of Bergman Spaces*, Springer-Verlag, New York, 2000.
4. K. Zhu, *Operator Theory in Function Spaces*, Marcel Dekker, New York, 1990.

Department of Mathematics, SUNY, Albany, NY 12222  
kzhu@math.albany.edu

---

# A Simple Proof of Descartes's Rule of Signs

---

Xiaoshen Wang

---

"Descartes' Rule of Signs is a staple of high school algebra, but a proof is seldom seen, even at the college level" [2]. A proof of the theorem is usually several pages long [2]. In this note we give a simple proof.

**Theorem (Descartes's Rule of Signs).** *Let  $p(x) = a_0x^{b_0} + a_1x^{b_1} + \cdots + a_nx^{b_n}$  denote a polynomial with nonzero real coefficients  $a_i$ , where the  $b_i$  are integers satisfying  $0 \leq b_0 < b_1 < b_2 < \cdots < b_n$ . Then the number of positive real zeros of  $p(x)$  (counted with multiplicities<sup>1</sup>) is either equal to the number of variations in sign in the sequence  $a_0, \dots, a_n$  of the coefficients or less than that by an even whole number. The number of negative zeros of  $p(x)$  (counted with multiplicities) is either equal to the number of variations in sign in the sequence of the coefficients of  $p(-x)$  or less than that by an even whole number.*

In the following we denote the number of variations in the signs of the sequence of the coefficients of  $p$  by  $v(p)$  and the number of positive zeros of  $p$  counting multiplicities by  $z(p)$ . We need the following simple lemma.

**Lemma.** *Let  $p(x) = a_0x^{b_0} + a_1x^{b_1} + \cdots + a_nx^{b_n}$  be a polynomial as in the theorem. If  $a_0a_n > 0$ , then  $z(p)$  is even; if  $a_0a_n < 0$ , then  $z(p)$  is odd.*

*Proof.* We consider only the case when  $a_0 > 0$  and  $a_n > 0$ . The other cases can be handled similarly. Because  $p(0) \geq 0$  and  $p(x) \rightarrow \infty$  as  $x \rightarrow \infty$ , it is clear that the graph of  $p$  crosses the positive  $x$ -axis an even number of times, where we count crossings without regard to multiplicity. If  $x = a$  is a point at which the graph of  $p$  touches but does not cross the positive  $x$ -axis, then the multiplicity of  $a$  is even. If the graph

---

<sup>1</sup>The theorem would be false if the positive real roots were counted without multiplicity, as the example  $x^2 - 2x + 1$  shows.

of  $p$  crosses the positive  $x$ -axis at a multiple zero  $x = a$ , then  $x = a$  is a zero of odd multiplicity. Thus the multiple zeros contribute an additional even number to  $z(p)$ , whence  $z(p)$  is even. ■

*Proof of the theorem.* We need prove only the first part of the theorem. Without loss of generality we may assume that  $b_0 = 0$ . We argue by induction on  $n$ . It is obvious that for  $n = 1$ ,  $v(p) = z(p)$ . Assume that for  $n \leq k - 1$ ,  $v(p) \geq z(p)$  and  $z(p) \equiv v(p) \pmod{2}$ , and consider the situation for  $n = k$ . We need to treat the following cases.

Case 1:  $a_0 a_1 > 0$ . In this instance  $v(p) = v(p')$ . By the lemma,  $z(p) \equiv z(p') \pmod{2}$ . By the induction hypothesis,  $z(p') \equiv v(p') \pmod{2}$  and  $z(p') \leq v(p')$ . Thus  $z(p) \equiv v(p) \pmod{2}$ . An appeal to Rolle's theorem gives  $z(p') \geq z(p) - 1$ . It follows that

$$v(p) = v(p') \geq z(p') \geq z(p) - 1 > z(p) - 2,$$

so  $z(p) \leq v(p)$ .

Case 2:  $a_0 a_1 < 0$ . Here  $v(p') + 1 = v(p)$  and  $z(p) - z(p') \equiv 1 \pmod{2}$  in view of the lemma. By the induction hypothesis,  $z(p') \equiv v(p') \pmod{2}$  and  $z(p') \leq v(p')$ . Again we infer that  $z(p) \equiv v(p) \pmod{2}$ . By Rolle's theorem,  $z(p') \geq z(p) - 1$ . Thus  $v(p) = v(p') + 1 \geq z(p') + 1 \geq z(p)$ . ■

**Remark.** All steps of the foregoing proof remain valid if we drop the requirement that the  $b_i$  be nonnegative integers and instead allow them to be arbitrary real numbers (possibly negative) satisfying  $b_0 < b_1 < \cdots < b_n$ . As a result, the first part of the theorem can be generalized to polynomials with real exponents.

**Example.** Let  $p(t) = a_0 + a_1 e^{b_1 t} + a_2 e^{b_2 t}$ . Letting  $x = e^{b_1 t}$  yields  $p(t) = a_0 + a_1 x + a_2 x^{b_2/b_1} = Q(x)$ . Since  $Q$  has at most two positive zeros,  $p$  has at most two real zeros.

**ACKNOWLEDGMENT.** The author would like to thank the referee for his valuable suggestions. The referee also pointed out that [1] refers to  $z(p) \leq v(p)$  as "Descartes's Lemma," with no mention of the additional fact that  $z(p) \equiv v(p) \pmod{2}$ . The proof that  $z(p) \leq v(p)$  given in [1] is essentially the same as our proof of that fact.

## REFERENCES

1. J. Bochnak, M. Coste, and M.-F. Roy, *Real Algebraic Geometry*, Springer-Verlag, New York, 1998.
2. S. E. Brodie, *Descartes' Rule of Signs*, <http://www.cut-the-knot.com/fta/ROS2.shtml>.
3. R. Descartes, *The Geometry of René Descartes with a Facsimile of the First Edition* (trans. D. E. Smith and M. L. Latham), Dover, New York, 1954.
4. B. E. Meserve, *Fundamental Concepts of Algebra*, Dover, New York, 1982.

*Department of Mathematics and Statistics, University of Arkansas at Little Rock, Little Rock, AR 72204*  
 xxwang@ualr.edu