

# External priors for the next generation of CMB experiments.

Alessandro Manzotti<sup>1,2,\*</sup> and Scott Dodelson<sup>1,2,3</sup>

<sup>1</sup>*Department of Astronomy & Astrophysics, University of Chicago, Chicago IL 60637*

<sup>2</sup>*Kavli Institute for Cosmological Physics, Enrico Fermi Institute, University of Chicago, Chicago, IL 60637*

<sup>3</sup>*Fermilab Center for Particle Astrophysics, Fermi National Accelerator Laboratory, Batavia, IL 60510-0500*

(Dated: October 29, 2015)

The next generation of cosmic microwave background (CMB) experiments can dramatically improve what we know about neutrino physics, inflation, Dark Matter and Dark Energy. Indeed the low level of noise, together with improved angular resolution, will allow a precise measurement of the CMB polarization anisotropies as well as an improved reconstruction of the lensing potential of high redshift large scale structure. Projected constraints on cosmological parameters are extremely tight, but these can be improved even further with information from external experiments. Here, we examine quantitatively, with a Fisher matrix approach, the extent to which external priors can lead to improvement in projected constraints from a CMB-Stage 4 (S4) experiment. We find that CMB S4 constraints on neutrino mass will be strongly enhanced by cold dark matter  $\Omega_c h^2$  and Hubble constant  $H_0$  external informations. If the largest scales ( $\ell < 50$ ) will not be measured an external prior on the primordial amplitude  $A_s$  or the optical depth  $\tau$  will also be important. A CMB constraint on  $N_{\text{eff}}$  will benefit from external prior on the spectral index  $n_s$  and baryons energy density  $\Omega_b h^2$ . Finally, external prior on  $H_0$  will be crucial to constraint a constant dark energy equation of state( $w_0$ ).

## I. INTRODUCTION

Since their early stages, Cosmic Microwave Background (CMB) experiments have been crucial in our understanding of the universe. In the near future, they have the potential to maintain this important role thanks to the unprecedented low level of noise and high resolution of their next Stage IV (S4) generation. Indeed, the next generation of CMB experiments (S4), now in its planning stage, will measure the E-mode polarization with almost cosmic variance limited precision together with an order of magnitude improvement in B-mode measurement and lensing reconstruction. This new sensitivity, in particular if supported with the essential synergy with other cosmological probes, will be a fundamental step forward in our understanding of dark matter, inflation, Dark Energy and neutrinos.

Neutrinos properties can be effectively constrained with CMB measurements because they influence both the early history of the universe and the later stage of structure formation. In the early Universe relativistic species, like neutrinos, are the main drivers of the cosmic expansion and the expansion rate depends on their effective number  $N_{\text{eff}}$ . This can be powerfully tested using the CMB, by comparing the sound horizon scale, obtained from the CMB peaks positions, and the Silk damping scale (see [1], [2] and references therein). S4 experiments, with their high resolution, will be able to measure the CMB power spectrum, at small scales, deeply into the damping tale, testing possible deviations from  $N_{\text{eff}} = 3.046$ , the value predicted by the standard model. The total mass of neutrinos, on the other hand, has a

modest effect on the primary CMB because, for the range of masses allowed by recent constraints ( $\sum m_\nu < 230$  meV from [3]), neutrinos are still relativistic at the last scattering surface. However massive neutrinos influence the growth of structure at scales shorter than their free streaming one. Different neutrino masses consequently lead to different CMB lensing. Stage IV experiment will measure small scales temperature and polarization anisotropies with low noise, drastically improving the lensing reconstruction. This will turn into such a precise constraints of the sum of neutrino masses to get a possible hint of the hierarchical structure of the individual masses. Quantitatively, CMB alone will constrain the number of relativistic degree of freedom  $N_{\text{eff}}$  with a 1% precision and the total mass of neutrinos at 45% with a factor of two improvement on  $\sum m_\nu$  when priors from baryon acoustic oscillations (BAO) data are introduced [4]. The future generation of CMB experiments will be a big step in the understanding of the neutrino sector.

As previously mentioned, CMB is also a powerful probe of the dark energy properties (see [5]). The challenge for future experiments is to go beyond the measurement of the energy density and to determine other dark energy properties. Among these, a crucial step would be to constraint the equation of state, the ratio of pressure and energy density, looking for possible deviations from the value  $w = -1$  expected in a cosmological constant scenario. CMB is sensitive to Dark energy because this component alters the late universe's expansion and it consequently changes the distance to the last scattering surface. Furthermore different DE models lead to a different growth of large scale structure which are tested by CMB lensing. However, dark energy properties are strongly degenerate with other geometrical parameters like  $H_0$  and  $\Omega_k$ . As a consequence, similarly to the neutrino sector, CMB experiments will have to rely on external prior com-

---

\* manzotti.alessandro@gmail.com

ing from BAO and supernova experiments to fully exploit their potential.

The crucial importance of the feedback from external experiments is true not only for the dark energy and the neutrino sector. In the future, cosmology will hinge on the ability of combining different probes. In this regard CMB makes no exception: external priors will be fundamental to improve the already tight CMB constraints on cosmological parameters. In particular CMB will really benefit from experiments like large scale structure clustering and weak lensing, BAO targeted experiments, and supernovae. To improve the synergy between different experiments and to guide the plan of future experiments, in this work we study the dependance of CMB future constraints on the external priors assumed. For all the parameters of the standard  $\Lambda$ CDM plus massive neutrinos ( $\Omega_c h^2, \Omega_b h^2, A_s, n_s, \tau, H_0, \sum m_\nu$ ) together with possible non-standard extension ( $N_{\text{eff}}, w$ ) we quantify, using a Fisher matrix approach, the CMB constraints as a function of external priors in each of the cosmological parameters.

This paper is organized as follow: in §II we introduce the technique and assumptions we used to derive the effect of external priors on the CMB parameter constraints. In §III we will describe our results and we then conclude with a discussion of them in §IV.

## II. ASSUMPTIONS AND METHODS

To measure quantitatively the impact of external priors on the CMB ability to constrain cosmological parameters we use a Fisher matrix formalism. In this section we will quickly review the technique and we will then present our fiducial cosmological model together with the experimental specifications we used.

As usual, we define the Fisher matrix elements as the curvature of the likelihood:

$$F_{ij} \equiv - \left\langle \frac{\partial^2 \log \mathcal{L}}{\partial \theta_i \partial \theta_j} \bigg|_{\theta=\theta_0} \right\rangle, \quad (1)$$

where  $\theta_{i,j}$  represents two of the cosmological parameters and  $\theta_0$  is the fiducial values array that, by definition, maximizes the likelihood.

For CMB experiments the Fisher matrix can be related to the power spectrum  $\mathbf{C}_\ell$  by:

$$F_{ij} = \sum_\ell \frac{2\ell+1}{2} f_{\text{sky}} \text{Tr} \left( \mathbf{C}_\ell^{-1}(\theta) \frac{\partial \mathbf{C}_\ell}{\partial \theta_i} \mathbf{C}_\ell^{-1}(\theta) \frac{\partial \mathbf{C}_\ell}{\partial \theta_j} \right). \quad (2)$$

In this work we constrain cosmological parameters with CMB temperature and E mode polarization together with the reconstructed lensing potential of large scale structure. For this reason,  $\mathbf{C}_\ell$  in Eq. (2) is:

$$\mathbf{C}_\ell \equiv \begin{pmatrix} C_\ell^{TT} + N_\ell^{TT} & C_\ell^{TE} & C_\ell^{T\phi} \\ C_\ell^{TE} & C_\ell^{EE} + N_\ell^{EE} & 0 \\ C_\ell^{T\phi} & 0 & C_\ell^{\phi\phi} + N_\ell^{\phi\phi} \end{pmatrix}. \quad (3)$$

Note that we are neglecting the term  $C_\ell^{E\phi}$ . As also noticed in previous literature [2, 4] this term contains very little information while adding possible numerical issues. The terms  $N_\ell^X$  represents the instrumental noise power of the specific experiment and will be described at the end of this section.

A lower bound on the error on the parameter  $i$ , marginalized over all the other parameters,  $\sigma_i$ , can be easily obtained as:

$$\sigma_i \equiv \sigma(\theta_i) \geq \sqrt{(\mathbf{F}^{-1})_{ii}}. \quad (4)$$

Furthermore it is easy to introduce external priors on cosmological parameters. Indeed we simply need to add to the Fisher matrixes elements the external priors (before we perform the matrix inversion of Eq. (4)). For example, a 1% prior on the parameter  $i$  can be added by:

$$F_{ii} \rightarrow F_{ii} + \frac{1}{(1\% \times \theta_{i,\text{fid}})^2}. \quad (5)$$

We carefully try to avoid any possible source of errors in computing the elements of Eq. (2) and in the matrix inversion of Eq. (4). We compute the derivatives in Eq. (2) using a 5 points finite difference formula: this high order approach allows us to use bigger steps around the fiducial parameters to compute derivatives. As a consequence the differences of power spectra corresponding to different values are big enough to make possible numerical accuracy issues in computing  $\mathbf{C}_\ell$  negligible. We also test the robustness of this calculation by changing the derivative steps in the range 2 – 7% of the correspondent  $\theta_0$ . The constraints change by at most 10% that should then be considered as a conservative numerical uncertainties.

We compute the power spectra  $C_\ell$  and the noise power  $N_\ell$  in Eq. (2) using CAMB. Our base cosmology is a flat  $\nu\Lambda$ CDM. We then extend the parameter space by introducing  $N_{\text{eff}}$  and  $w$  as free parameters. We chose our fiducial parameters following Table 2 of *Planck* best fit [6], i.e.  $\Omega_c h^2 = 0.12029$ ,  $\Omega_b h^2 = 0.022068$ ,  $A_s = 2.215 \times 10^{-9}$  at  $k_0 = 0.05 \text{ Mpc}^{-1}$ ,  $n_s = 0.9624$ ,  $\tau = 0.0925$ ,  $H_0 = 67.11 \text{ km/s/Mpc}$  and a neutrino total mass  $\sum m_\nu \simeq 85 \text{ meV}$ . Regarding the  $\Lambda$ CDM extensions, we chose as fiducial, a standard  $N_{\text{eff}} = 3.046$  and a cosmological constant equation of state,  $w = -1, w_a = 0$ . It is important to notice that while we vary one parameter to compute derivatives in Eq. (2) we keep all the others fixed with the exception of  $\Omega_\Lambda$  which is always changed in order to keep the universe flat ( $\Omega_k = 0$ ).

The instrumental noise power  $N_\ell^{T,E}$  and the lensing reconstruction  $N_\ell^\phi$  in Eq. (2) correspond to the expected level for the next generation of CMB experiments (S4). For the temperature and E-mode polarization of the CMB, together with the improved depth and resolution we also assume that large scale foregrounds, like dust, are under control or negligible. This allow us to use all the polarization power spectrum multipoles all the way up to  $\ell_{\text{E,max}} = 5000$ . We deal with the poisson noise from point

sources in the temperature signal by simply discarding all the small scales modes with  $\ell > \ell_{T,\max} = 3000$ . The remaining source of noise, the instrumental noise, is added to the power spectrum in the usual way:

$$N_\ell^X = s^2 \exp\left(\ell(\ell+1)\frac{\theta_{\text{FWHM}}^2}{8\log 2}\right), \quad (6)$$

where  $\theta_{\text{FWHM}}^2$  is the FWHM of the experiment's beam and  $s$  represents the instrumental white noise. We decide to use a level of noise  $s = 0.58 \mu\text{K-arcmin}$  for  $X = T$  and a beam of  $\theta_{\text{FWHM}} = 1 \text{ arcmin}$ . This correspond to the  $N_{\text{det}} = 10^6$  case of [4]. Note that the quoted noise in temperature and we assume that  $s \rightarrow s \times \sqrt{2}$  in the case of polarization  $XX' = \{EE, BB\}$ .

The noise  $N_\ell^{\phi\phi}$  associated to the reconstructed  $\phi$  is modeled assuming an iterative reconstruction technique [7].

### III. RESULTS

In this section we present our main results. We will start from our base  $\nu\Lambda\text{CDM}$  and we then explore extensions to this model ( $N_{\text{eff}}, w$ ).

#### A. Neutrino Masses, $\sum m_\nu$

Small scale structure formation is suppressed if fast-moving neutrinos comprise a significant part of the matter budget. This effect, probed by CMB lensing, will allow us to determine the contribution of neutrinos to the total matter density. Since the active neutrino number densities are known in the standard model, this can be directly transformed into a constraint on the the sum of the neutrino masses,  $\sum m_\nu$ . This is particularly exciting because we now know that neutrinos are massive, with a lower limit  $\sum m_\nu > 0.05 \text{ eV}$  that emerges from oscillation experiments.

Fig. 1 shows the errors on  $\sum m_\nu$  with a CMB S4 experiment ( $N_{\text{det}} = 10^6$ ,  $f_{\text{sky}} = 0.75$ ,  $2 < \ell < 5000$ ) as a function of external priors. Without any external priors, upcoming CMB experiments are poised to obtain 1-sigma limits on  $\sum m_\nu$  of order 30 meV, corresponding to close to a 2-sigma detection even in the worst case scenario. Adding in external priors helps this significantly though, as discussed in [1, 8, 9]. There, the prior was described as ‘‘DESI BAO’’, that is a measurement of distances as a function of redshift from the baryon acoustic oscillation feature probed by DESI. Here, the same improvement is parametrized by a strong prior on  $\Omega_c h^2$  and  $H_0$ , which indeed determines these low-redshift distances. Notice that external prior effect are not completely independent. An external prior on  $H_0$  for example, improve the internal reconstruction of  $\Omega_c h^2$  as well. As found in [1], the projected error falls below 20 meV with this prior. Adding in other priors would help further.




FIG. 1. In the figure, the constraints on the sum of the neutrino masses as a function of priors on cosmological parameters are shown ( $N_{\text{det}} = 10^6$ ,  $f_{\text{sky}} = 0.75$ ,  $2 < \ell < 5000$ ). Each prior is expressed in units of the error that would be obtained internally from our suite of CMB experiments; e.g.,  $\sigma(H_0)_{\text{CMB}} = 0.6\%$ .  $H_0$  and  $\Omega_c h^2$  will be the more useful external prior to improve neutrino masses constraint in this regime. We chose a fiducial value  $\sum m_\nu = 85 \text{ meV}$  and we impose a Planck-like prior. The blue horizontal line correspond to the CMB S4+BAO constraint of [4].




FIG. 2. Some as Fig. 1 but with  $\ell_{\min} > 50$  as a function of priors on cosmological parameters, as in Fig. 3.

In Fig. 2 we limit CMB accessible modes to  $\ell_{\min} > 50$ . Together with the importance of  $H_0$  and  $\Omega_c h^2$  that we also observed in Fig. 1, in this case external priors on  $A_s$  and  $\tau$  will be helpful. Indeed as recently pointed out by [9], a degeneracy exists between  $\sum m_\nu$  and  $\tau$ . A measurement of the optical depth will help to constraint the primordial amplitude  $A_s$  which is degenerate with the mass of neutrinos. Indeed, increasing the mass of

neutrinos will decrease the CMB lensing power spectrum and this can be compensating by increasing  $A_s$ .

### B. Relativistic Degrees of Freedom, $N_{\text{eff}}$

In the standard cosmology, three active neutrinos are thermally produced in the early universe. Were they to decouple well before the epoch of electron-positron annihilation, their energy density after their decoupling would be equal to  $3 \times (7/8) \times (4/11)^{4/3} \rho_{\text{cmb}}$ , with the first factor capturing the contributions from the 3 active species; the second the difference between fermions and bosons; and the last the relative heating of the photons in the CMB by electron-positron annihilation. However decoupling is not a discrete event and occurs close to the time of electron-positron annihilation, so the neutrinos share a bit in the heating, with the factor of 3 replaced by  $N_{\text{eff}} = 3.046$ . The additional fraction depends not only on well-known neutrino scattering rates but also finite temperature quantum corrections. Upcoming experiment have the potential to measure this tiny deviation of  $N_{\text{eff}}$  from 3. This will be an amazing test of our understanding of the Universe when it was about a second old. Furthermore any possible significant deviation from this value could be a hint of a different scenario not predicted by the standard model.




FIG. 3. Projected 1-sigma error on  $N_{\text{eff}}$  as a function of the strength of the prior on the other 6 cosmological parameters of the standard  $\Lambda$ CDM model. The parameters that would be most useful to constrain externally for the purposes of determining  $N_{\text{eff}}$  are  $n_s$  and  $\Omega_b h^2$ . The bottom-most (grey) curve shows the impact of imposing the given prior on *all* the other parameters simultaneously.

Fig. 3 shows projections for how well our suite of CMB experiments will do at measuring  $N_{\text{eff}}$  as a function of priors on the other 6 parameters in the standard cosmological model ( $H_0, \Omega_m h^2, \Omega_b h^2, n_s, A_s$ , and  $\tau$ ). With no

priors on the other parameters, the projected 1-sigma error is about 0.02, suggesting a 2-sigma detection of the deviation from  $N_{\text{eff}} = 3$ . The sensitivity comes from the effect of extra species on the damping tail of the CMB anisotropies [2], both in temperature and polarization, so parameters that also affect the shape of the damping tail, like the slope  $n_s$  and the baryon density  $\Omega_b h^2$ , are most degenerate with  $N_{\text{eff}}$ . As such, Fig. 3 shows that obtaining external priors on either of these would reduce the errors on  $N_{\text{eff}}$  to ensure a 3-sigma detection of the partial decoupling prediction. The bottom-most curve shows that if all parameters were constrained externally, then the projected error on  $N_{\text{eff}}$  would improve significantly.

### C. Dark Energy Equation of State, $w$

The CMB constrained the late-time dark energy equation of state in two ways. The primordial CMB is very sensitive to the distance to the last scattering surface which depends on  $w$ . Furthermore the CMB lensing probes the growth of structure at late times which is different for different dark matter equations of state.




FIG. 4. Constraints on the dark energy equation of state  $w$  as a function of external priors on cosmological parameters. An external prior on  $H_0$  will be crucial to improve the constraint.

Fig. 4 shows that without any priors, CMB experiments will not do much better than current constraints, which hover around 10%. However, an external constraint on the Hubble constant would improve the CMB constraints on  $w$  considerably. This primarily due to the fact that CMB constrained  $w$  through a measurement of the last scattering surface. However that distance can be kept fixed while varying  $w$  by accordingly changing the value of  $H_0$  (and  $\Omega_\Lambda$  to keep the universe flat). Indeed in a two dimensional ( $H_0, w$ ) the CMB constrained approximately lies on the region of constant large scattering surface as shown in Fig. 5. Notice that an external




FIG. 5. Two dimensional constraint  $1\sigma$  ellipse for  $w, H_0$  space. The dashed (red) line correspond to values in the plane corresponding to the same distance to the last scattering surface.

prior on  $H_0$  three times more accurate than CMB S4 can improve the CMB constraint almost by a factor of three from a 8% error to a 3% level.

#### IV. CONCLUSIONS

The design of the next CMB Stage IV experiments represents an exciting challenge. The possible reward is remarkable. We can finally put a tight constraint on the total mass of neutrinos with deeply implications on particle physics like, for example, the solution of the mass hierarchy problem. We will measure with great accuracy the number of relativistic species, both testing standard model prediction and, eventually, probing new physics. The same is true for dark energy where we will tighten the constraint on  $w, w_a$  getting closer to test the nature of the accelerated expansion of the universe. These results, however, will not be possible without an optimal synergy with the other experiments observing a variety of others

cosmological probes. Indeed this should be taken into account, especially during the CMB Stage IV planning phase. Understanding CMB weaknesses and the impact of external data can have a deep influence in the needed experiment specifications. For this reason in this work we explored the impact of external priors on CMB S4 cosmological constraints and in particular on the neutrino sector and dark energy parameters. We find:

$\sum m_\nu$ : To improve the CMB constraint by a factor of two, low-redshift measurements will be needed... If the final design of CMB S4 will not include measurement at large scale ( $\ell < 50$ ) external constraint on the primordial amplitude will be needed. Moreover, an external prior on  $\tau$  will allow the CMB to constraint  $A_s$  internally thus strongly alleviating the problem.

$N_{\text{eff}}$ : The effective neutrino numbers will benefit from external prior on parameters that affect the shape of the damping tail.

$w$ : In this case the improvement will mainly come by external prior on  $H_0$ .

*Comment on issue and future. This will be even more relevant when we open the parameter space. Will CMB + others be able to discover something CMB can't alone?  $N_{\text{eff}}$  extensions, mass splitting etc*

#### ACKNOWLEDGMENTS

We thank Youngsoo Park for his important contribution in the early stage of this work. AM wants to thank Zhen Pan who allowed a careful cross-check of our results. This work was partially supported by the Kavli Institute for Cosmological Physics at the University of Chicago through grants NSF PHY-1125897 and an endowment from the Kavli Foundation and its founder Fred Kavli. The work of SD is supported by the U.S. Department of Energy, including grant DE-FG02-95ER40896.

- 
- [1] K. N. Abazajian, K. Arnold, J. Austermann, B. A. Benson, C. Bischoff, J. Bock, J. R. Bond, J. Borrill, E. Calabrese, J. E. Carlstrom, C. S. Carvalho, C. L. Chang, H. C. Chiang, S. Church, A. Cooray, T. M. Crawford, K. S. Dawson, S. Das, M. J. Devlin, M. Dobbs, S. Dodelson, O. Dore, J. Dunkley, J. Errard, A. Fraisse, J. Gallicchio, N. W. Halverson, S. Hanany, S. R. Hildebrandt, A. Hincks, R. Hlozek, G. Holder, W. L. Holzapfel, K. Honscheid, W. Hu, J. Hubmayr, K. Irwin, W. C. Jones, M. Kamionkowski, B. Keating, R. Keisler, L. Knox, E. Komatsu, J. Kovac, C.-L. Kuo, C. Lawrence, A. T. Lee, E. Leitch, E. Linder, P. Lubin, J. McMahon, A. Miller, L. Newburgh, M. D. Niemack, H. Nguyen, H. T. Nguyen, L. Page, C. Pryke, C. L. Reichardt, J. E. Ruhl, N. Sehgal, U. Seljak, J. Sievers, E. Silverstein, A. Slosar, K. M. Smith, D. Spergel, S. T. Staggs, A. Stark, R. Stompor, A. G. Vieregg, G. Wang, S. Watson, E. J. Wollack, W. L. K. Wu, K. W. Yoon, and O. Zahn, ArXiv e-prints (2013), arXiv:1309.5383.
- [2] Z. Hou, R. Keisler, L. Knox, M. Millea, and C. Reichardt, Phys. Rev. D **87**, 083008 (2013), arXiv:1104.2333 [astro-ph.CO].
- [3] Planck Collaboration, P. A. R. Ade, N. Aghanim, C. Armitage-Caplan, M. Arnaud, M. Ashdown, F. Atrio-Barandela, J. Aumont, C. Baccigalupi, A. J. Banday, and et al., A&A **571**, A16 (2014), arXiv:1303.5076.
- [4] W. L. K. Wu, J. Errard, C. Dvorkin, C. L. Kuo, A. T. Lee, P. McDonald, A. Slosar, and O. Zahn, Astrophys. J. **788**, 138 (2014), arXiv:1402.4108.

- [5] H. K. Jassal, J. S. Bagla, and T. Padmanabhan, MNRAS **405**, 2639 (2010), astro-ph/0601389.
- [6] Planck Collaboration, P. A. R. Ade, N. Aghanim, C. Armitage-Caplan, M. Arnaud, M. Ashdown, F. Atrio-Barandela, J. Aumont, C. Baccigalupi, A. J. Banday, and et al., A&A **571**, A16 (2014), arXiv:1303.5076.
- [7] U. Seljak and C. M. Hirata, Phys. Rev. D **69**, 043005 (2004), astro-ph/0310163.
- [8] Z. Pan and L. Knox, ArXiv e-prints (2015), arXiv:1506.07493.
- [9] R. Allison, P. Caucal, E. Calabrese, J. Dunkley, and T. Louis, ArXiv e-prints (2015), arXiv:1509.07471.