

The importance of external priors for the next generation of CMB experiments.

A. Manzotti and S. Dodelson

(Dated: August 11, 2015)

Abstract

The next generation of cosmic microwave background (CMB) experiments can potentially improve drastically what we know about neutrino physics, inflation, Dark Matter and Dark Energy. Indeed they will likely exploit the great sensitivity of the CMB to the details of physical processes in the early Universe (through temperature and polarization) as well as in the late universe (through CMB secondary effects like lensing). They will be possible because the temperature anisotropies and E mode polarization of the CMB will be measured with cosmic variance limited precision. Such a low level of noise, together with the improved resolution, will also allow CMB experiment to reconstruct the lensing potential of high redshift large scale structure with a signal to noise bigger than one on several scales. Indeed recently, forecasts papers validate these possible outcomes and help to plan and motivate the next generation of CMB experiments. However these results also indirectly disclose the needed synergy between CMB and other experiments, for example the need of imposing external prior on some of the cosmological parameters. We think that the extent to which CMB constraints depend on external priors needs to be explored, for instance to guide the combined effort of different experimental fields. For this reason, in this work we adopt a fisher-matrix technique to test how external priors can lead to improvement in the S4 generation of CMB parameters constraint. We find that quite surprisingly CMB S4 are very powerful even on .. There are few exceptions.. like w .

I. INTRODUCTION

Since their early stages, Cosmic Microwave Background (CMB) experiments have been crucial in our understanding of the universe. Moreover they also represent an amazing success in instrumental physics. Indeed every new generation of satellite experiments improved its sensitivity gaining almost a factor of ten in sensitivity compared to its predecessor, from the first generation instrument COBE to WMAP all the way to the current state of the art represented by Planck [1–5]. Experimentally we went from observing the first peak of the CMB temperature power spectrum with COBE to a cosmic variance limited measurement of several peaks with Planck. With the improved sensitivity we managed to extend the scientific outcome of these experiments from a measurement of the universe flatness to a percent level constraint on the parameters of the standard Λ CDM model. The same is true for ground experiment where the progression from DASI [6] to SPT and ACT [7] [8] allowed us to measure the small scale damping tail of the CMB spectrum with increasing accuracy. Ground experiments have already reached the maximum possible sensitivity per detector but will significantly increase the number of detectors in future generation. These series of successes is far from its end. Indeed the next generation of CMB experiments (S4) is now in its planning stage and it promises to measure with cosmic variance limited precision the E-mode polarization together with an order of magnitude improvement in B-mode measurement and lensing reconstruction. As happened in the past this new sensitivity will improve our understanding of several areas like neutrinos, inflation, Dark Energy and Dark matter. In this work we will mainly focus on two of these: neutrinos and Dark Energy.

The main aspects of neutrinos physics that CMB can access are the number of relativistic degree of freedom in the early universe N_{eff} and the total mass of the three neutrino species M_ν . Indeed the number of relativistic species, like neutrinos, in the early Universe alter the expansion history. This effect can be probed with CMB measurement by comparing the sound horizon scale, obtained from the CMB peaks position, with the silk damping scale. Indeed because they scale differently with the expansion rate $H(z)$ their ratio $r_d/r_s \propto \sqrt{H}$ (see [9], [10] and references therein). This allow us to constrain N_{eff} and any deviation from the predicted value $N_{\text{eff}} = 3.046$ will be a sign of possible extensions of the Standard Model. The mass of neutrinos, on the other hand, has a modest effect on the CMB because, for the range of masses allowed by recent constraints ($M_\nu < 230$ meV from [2]), neutrinos

are still relativistic at the last scattering surface. As other low redshift effects however, massive neutrinos modify the CMB through the lensing effect of large scale structure on the CMB photons. Different neutrino masses result into different growth of structure which consequently lead to different CMB lensing. The lensing potential can be reconstructed from the CMB itself, opening the possibility of constraining the neutrino mass. In the future, according for example to [11], CMB alone will constrain N_{eff} with a 1% precision and the total mass of neutrinos at 60% with a factor of two improvement when BAO data are also used. The future generation of CMB experiments will be a big step in the understanding of the neutrino sector.

As previously mentioned, CMB is also sensitive to Dark energy (see [12]). The current generation of experiments measured with percent accuracy the energy density of Dark Energy. The next big challenge is to identify the nature of this mysterious component. The first step will be to measure any deviation from the equation of state predicted by a cosmological constant model. Furthermore CMB will also be a powerful probe of any time dependence of the dark energy equation of state. Indeed dark energy mainly affect the CMB because of its effect on the universe's expansion that modifies the distance to the last scattering surface. Furthermore the influence of Dark Energy on the low redshift universe modify the energy of CMB photons through the Integrated Sachs Wolfe Effect. Also, as it was true in the neutrino case, CMB lensing can probe the growth of structure that depends on the dark energy properties and in particular on its equation of state. However dark energy properties are strongly degenerate with other geometrical parameters like H_0 and Ω_k . To get competitive constraints, CMB experiments will have to rely on external prior coming from BAO and supernova experiments.

The importance of external priors in the dark energy case is actually more general issue. Even if it is certainly true that CMB S4 will open an all new window on several area of physics and cosmology the synergy with other experiments is crucial. In particular CMB will really benefit from experiments like large scale structure clustering and weak lensing, BAO targeted experiments, and supernovae. For this reason forecasted constraints very often assume external realistic priors. To improve the synergy between different experiments and to guide the plan of future experiments, in this work we study the dependance of CMB future constraints on the external priors assumed. This can tell us for example which CMB parameter constraints will benefit more from improvement on external priors or if proposed

experiments like PIXIE [13] can indirectly also benefit CMB constraints.

This paper is organized as follow: in section II we introduce the technique and assumptions we use to derive the effect of external priors on the CMB parameter constraints. In section III we will describe our findings and we then conclude with a discussion of our results in section IV.

II. ASSUMPTION AND METHODS

The main goal of this paper is to measure the effect of external priors on the statistical errors of cosmological parameters derived from future CMB experiments. We start in section II A by introducing the Fisher matrix formalism, a simple but powerful technique widely used to forecast future experimental constraints on the parameters of the assumed cosmological model. To apply this technique, we need to choose a model (Λ CDM plus neutrinos and dark energy extensions in our case) as well as a set of fiducial parameters and the details of the CMB experiments we have in mind. These will be presented in section II B.

A. Fisher Matrix formalism

The natural starting point to estimate errors on cosmological parameters is the Bayes theorem that relates the likelihood to measure a set of data given the parameters of the model $\mathcal{L}(d|\theta)$, to what we want: the posterior probability of those parameters given the data, $\mathcal{L}(\theta|d)$. These two are related by the prior probability of the parameters $P(\theta)$ through:

$$\mathcal{L}(\theta|d) \propto \mathcal{L}(d|\theta)P(\theta), \quad (1)$$

where, as usual, we have neglected the probability of the data itself. These general framework can be simplified in our case. We will focus not on an entirely general posterior distribution, but we will assume that the likelihoods are gaussian. Furthermore we will not derive the global shape of the likelihood but we will obtain parameters errors by studying small perturbations around its maximum. These are the two basic assumptions of the Fisher approach.

Regarding the first assumption, we recognize the fact that, even if the gaussian approximation has been shown to be appropriate most of the time, some problems have been found

in other cases [14]. Despite this, we do not need the level of accuracy that a careful modeling of the likelihoods will give us, because we are looking for the general behavior of parameters errors as a function of external priors. Moreover the gaussian approximation gets better at smaller scales (high ℓ in Fourier space) which, a part for the exception of the optical depth parameters τ , is where most of the constraining power of the CMB is coming from.

Secondly we will assume to know the true “fiducial” parameters that maximize the likelihood $\mathcal{L}(\theta|d)$ and we will get the errors on those by a simple analysis of the likelihood curvature around the fiducial values. Indeed, as usual, we define the fisher matrix elements as the curvature:

$$F_{ij} \equiv - \left\langle \frac{\partial^2 \log \mathcal{L}}{\partial \theta_i \partial \theta_j} \bigg|_{\theta=\theta_0} \right\rangle, \quad (2)$$

where $\theta_{i,j}$ represents two of the parameters and θ_0 is the parameters array that, by definition, maximize the likelihood. We refer the reader to [] for a detailed proof that with the usual definition of power spectrum $\langle a_{\ell m}^X a_{\ell' m'}^Y \rangle = \delta_{\ell\ell'} \delta_{mm'} C^{X,Y}$, the Fisher matrix of a CMB experiment can be rewritten as:

$$F_{ij} = \sum_{\ell} \frac{2\ell+1}{2} f_{sky} \text{Tr} \left(\mathbf{C}_{\ell}^{-1}(\theta) \frac{\partial \mathbf{C}_{\ell}}{\partial \theta_i} \mathbf{C}_{\ell}^{-1}(\theta) \frac{\partial \mathbf{C}_{\ell}}{\partial \theta_j} \right). \quad (3)$$

In this work we will use the CMB temperature and E mode polarization together with lensing reconstruction to constrain parameters. For this reason, \mathbf{C}_{ℓ} that encapsulate the used power spectra in a matrix structure is:

$$\mathbf{C}_{\ell} \equiv \begin{pmatrix} C_{\ell}^{TT} + N_{\ell}^{TT} & C_{\ell}^{TE} & C_{\ell}^{Td} \\ C_{\ell}^{TE} & C_{\ell}^{EE} + N_{\ell}^{EE} & 0 \\ C_{\ell}^{Td} & 0 & C_{\ell}^{dd} + N_{\ell}^{dd} \end{pmatrix}. \quad (4)$$

Note that we are neglecting the term $C_{\ell}^{E\phi}$. As also noticed in previous literature like ([10, 11]) this term contains very little information while adding possible numerical issues. Furthermore the terms N_{ℓ}^X represents the instrumental noise power of the specific experiment and will be discussed in section II B. The power of the Fisher approach descend from the Cramer-Rao inequality that relates the errors on parameter i , marginalized over all the other parameters, σ_i , to the Fisher matrix as:

$$\sigma_i \equiv \sigma(\theta_i) = \sqrt{(\mathbf{F}^{-1})_{ii}}. \quad (5)$$

Once F_{ij} is computed following Eq. (3) it is straightforward to get the error σ_i from Eq. (5). Furthermore in this context it is easy to introduce external priors on cosmological parameters, which is the main focus of our work. Indeed to add priors we simply need to add external experiments Fisher matrixes before performing the matrix inversion of Eq. (5), i.e.:

$$F_{total} = F_{CMB} + F_{external}. \quad (6)$$

In the same way we can add priors on a single cosmological parameter just by adding the prior to the matrix element. For example, a 1% prior on H_0 can be obtained by:

$$F_{H_0 H_0} \rightarrow F_{H_0 H_0} + \frac{1}{(1\% \times H_{0,\text{fid}})^2}. \quad (7)$$

We chose a Fisher matrix approach because of its ability to rapidly forecast future experiments performances without generating mock data together with the ease of including external priors. This technique however introduces some technical difficulties together with the previously mentioned assumption that the likelihood is gaussian. Indeed it is known [15] that increasing the number of parameters used in the analysis can lead to numerical issues (see also [16] in the gravitational waves context where a lot of parameters are needed). Fisher matrix indeed can become ill-conditioned: a small change in the Fisher matrix led to a big change in its inverse. Because we use Eq. (5) this can be a problem for error estimation. Even if other methods of analysis have been proposed [15, 17] this is still the standard method used to forecast future constraints [11]. While we know these issues can be a possible concern we carefully try to avoid any possible source of errors in computing the elements of Eq. (3) and in the matrix inversion of Eq. (5). We compute the derivatives in Eq. (3) using a 5 points formula:

$$\left. \frac{\partial C}{\partial \theta} \right|_{\theta_0} \sim \frac{-C(\theta_0 + 2h) + 8C(\theta_0 + h) - 8C(\theta_0 - h) + C(\theta_0 - 2h)}{12h}. \quad (8)$$

This high order definition allows us to use a bigger gap h around the fiducial parameters θ_0 . As a consequence the differences of power spectra corresponding to different values are big enough to make possible numerical accuracy issues in computing C negligible. We also test the robustness of this calculation by changing the gap h in the range 2 – 7% of the correspondent θ_0 without noticing any significant change in the results. Furthermore we compare our results to similar previous work in the literature [18] obtaining a perfect agreement. Lastly we implement the same technique of [15] to avoid possible numerical

162 instability in the marginalizing procedure. This allow us not to invert the entire matrix
 163 when we want to marginalize over a set of parameters, in order to minimize numerical issues
 164 and conserve parameters degeneracies as much as possible.




FIG. 1. Lensing potential power spectrum use in this work has a signal to noise bigger than one up to $\ell \simeq 800 - 1000$. In the figure: the deflection power spectrum for our fiducial cosmology together with the two examples of lensing reconstruction noise N^ϕ used in this work.

165 B. CMB S4 experiments: level of noise.

166 In this subsection we will describe the assumptions we made in calculating the elements
 167 that goes in Eq. (3) and, in particular, the power spectra C and the noise power N . The
 168 first step is the cosmological model we used to compute $C^{X,Y}$ with $X, Y \in \{T, E, \phi\}$. We
 169 parametrize our cosmology using a flat $\nu\Lambda$ CDM universe. We also allow different Dark




FIG. 2. The next generation $C^{T,E}$ used in this work are almost cosmic variance limited. In the figure: CMB power spectrum for our fiducial cosmology together with the instrumental noise used in this work.

Energy model by introducing the equations of state parameter w as a varying parameter. We chose our fiducial parameters following Table 2 of *Planck* best fit [19], i.e. $\Omega_c h^2 = 0.12029$, $\Omega_b h^2 = 0.022068$, $A_s = 2.215 \times 10^{-9}$ at $k_0 = 0.05 \text{ Mpc}^{-1}$, $n_s = 0.9624$, $\tau = 0.0925$, and $H_0 = 67.11 \text{ km/s/Mpc}$. Regarding the Λ CDM extension, we chose a neutrino energy densities $\Omega_\nu h^2 = 0.0009$, which corresponds to $M_\nu \simeq 85 \text{ meV}$ a standard $N_{\text{eff}} = 3.046$ and a dark energy equation of state, $w = -1$. Given the model, we use the CAMB software to compute the power spectra C_ℓ at the fiducial values and at those needed to compute derivatives using Eq. (8). Notice that while we vary one parameter in Eq. (8) we keep all the others fixed with the exception of Ω_Λ which is always changed in order to keep the universe flat ($\Omega_k = 0$).

We chose the second element of Eq. (3), the noise power in the CMB $N_\ell^{T,E}$ and in the lensing reconstruction N_ℓ^ϕ with the next generation of CMB experiments (S4) in mind. For the temperature and E-mode polarization of the CMB, together with the an improved depth and resolution we will also assume that large scale foregrounds, like dust, are under control or negligible. We will deal with the presence of point sources poisson noise in the temperature signal by simply discarding all the small scales modes with $\ell > \ell_{T,\max} = 3000$. The remaining source of noise, the instrumental noise is added to the power spectrum in the usual way:

$$N_\ell^X = s^2 \exp \left(\ell(\ell+1) \frac{\theta_{\text{FWHM}}^2}{8 \log 2} \right), \quad (9)$$

where θ_{FWHM}^2 is the FWHM of the experiment's beam and s represent the instrumental white noise. We decide to use a level of noise $s = 1.5 \mu\text{K-arcmin}$ for $X = T$ and a beam of 1 arcmin (PRELIMINARY). Note that $1.5 \mu\text{K-arcmin}$ is the noise in temperature and we need $s \rightarrow s \times \sqrt{2}$ in the case of polarization $XX' = \{EE, BB\}$.

Together with E and T we will use the information contained in the lensing potential ϕ as it is reconstructed from CMB experiments. The lensing potential represents the integration along the line of site of the gravitational potential and it leaves its signature in the CMB, both in temperature and polarization, by bending the trajectory of CMB photons. This introduce non gaussianities that couple different modes in the otherwise independent CMB modes and it can be reconstructed using a quadrature estimator technique [20, 21]. For the noise $N_\ell^{\phi\phi}$ associated to the reconstructed ϕ we follow [20, 21] without using more advanced iterative techniques.

III. RESULTS

H_0	67.11 km/s/Mpc
τ	0.0925
A_s	2.215×10^{-9}
n_s	0.9624
N_{eff}	3.046

TABLE I. Fiducial values of cosmological parameters used in this work.

H_0	M_ν	$\Omega_{bc}h^2$	$\Omega_b h^2$	τ	A_s	n_s
0.80%	40.92%	0.44%	0.11%	3.075%	0.53%	0.18%

TABLE II. How well we do constrain separate parameters with this data without any external prior? Things to notice: this is done with the Zhen test parameters ($\ell < 3000$ and no error on lensing). Now if we trust it CMB alone can get a 0.8% error on H_0 thus I am not surprised if a prior on H would not help. However what do we think about it? is it really CMB better than SN. People will not agree on that. τ will probably improve and also M_ν from BAO may help improving parameters.

IV. CONCLUSIONS

–We find that...

–This means that supernova experiments CMB spatial distortions and large scale structure can/will/should

–Future works

Appendix A: Marginalization

$$G = F^{\phi\phi} - F^{\phi\psi} U \Lambda^{-1} U^T F^{\psi\psi}, \quad (\text{A1})$$

where we define $\phi = \{N_{eff}, \dots\}$ and $\psi = \{\Omega_m \dots marginal\}$; therefore, $F^{\phi\phi}$ is the block of the total Fisher matrix containing the parameters we want to constrain, whilst $F^{\psi\psi}$ is the nuisance-parameter Fisher sub-matrix. Here, Λ is the diagonal matrix whose elements are the eigenvalues of $F^{\psi\psi}$, whilst U is the orthogonal matrix diagonalising $F^{\psi\psi}$. By using Eq., our marginalizing procedure is more stable, since degeneracies in $F^{\phi\phi}$ are properly propagated to G with no instabilities, and we do not even worry about a possibly ill-conditioned $F^{\phi\phi}$ sub-matrix, since we check its stability on the fly by the diagonalisation.

ACKNOWLEDGMENTS

We thank Youngsoo Park for his contribution in the early stage of this work. AM want to thank Zhen Pan who allows a careful cross-check of our results. This work was partially




FIG. 3.

supported by the Kavli Institute for Cosmological Physics at the University of Chicago through grants NSF PHY-1125897 and an endowment from the Kavli Foundation and its founder Fred Kavli. The work of SD is supported by the U.S. Department of Energy, including grant DE-FG02-95ER40896.

-
- [1] Planck Collaboration, P. A. R. Ade, N. Aghanim, M. Arnaud, M. Ashdown, J. Aumont, C. Baccigalupi, A. J. Banday, R. B. Barreiro, J. G. Bartlett, and et al., ArXiv e-prints (2015), arXiv:1502.01589.
- [2] Planck Collaboration, P. A. R. Ade, N. Aghanim, C. Armitage-Caplan, M. Arnaud, M. Ashdown, F. Atrio-Barandela, J. Aumont, C. Baccigalupi, A. J. Banday, and et al., A&A **571**, A16 (2014), arXiv:1303.5076.




FIG. 4.

- [3] D. N. Spergel, L. Verde, H. V. Peiris, E. Komatsu, M. R. Nolta, C. L. Bennett, M. Halpern, G. Hinshaw, N. Jarosik, A. Kogut, M. Limon, S. S. Meyer, L. Page, G. S. Tucker, J. L. Weiland, E. Wollack, and E. L. Wright, *ApJS* **148**, 175 (2003), astro-ph/0302209.
- [4] S. Hanany, P. Ade, A. Balbi, J. Bock, J. Borrill, A. Boscaleri, P. de Bernardis, P. G. Ferreira, V. V. Hristov, A. H. Jaffe, A. E. Lange, A. T. Lee, P. D. Mauskopf, C. B. Netterfield, S. Oh, E. Pascale, B. Rabbii, P. L. Richards, G. F. Smoot, R. Stompor, C. D. Winant, and J. H. P. Wu, *ApJL* **545**, L5 (2000), astro-ph/0005123.
- [5] P. de Bernardis, P. A. R. Ade, J. J. Bock, J. R. Bond, J. Borrill, A. Boscaleri, K. Coble, B. P. Crill, G. De Gasperis, P. C. Farese, P. G. Ferreira, K. Ganga, M. Giacometti, E. Hivon, V. V. Hristov, A. Iacoangeli, A. H. Jaffe, A. E. Lange, L. Martinis, S. Masi, P. V. Mason, P. D. Mauskopf, A. Melchiorri, L. Miglio, T. Montroy, C. B. Netterfield, E. Pascale, F. Piacentini, D. Pogosyan, S. Prunet, S. Rao, G. Romeo, J. E. Ruhl, F. Scaramuzzi, D. Sforza, and




FIG. 5.

N. Vittorio, Nature (London) **404**, 955 (2000), astro-ph/0004404.

[6] N. W. Halverson, E. M. Leitch, C. Pryke, J. Kovac, J. E. Carlstrom, W. L. Holzapfel, M. Dragovan, J. K. Cartwright, B. S. Mason, S. Padin, T. J. Pearson, A. C. S. Readhead, and M. C. Shepherd, Astrophys. J. **568**, 38 (2002), astro-ph/0104489.

[7] J. Dunkley, R. Hlozek, J. Sievers, V. Acquaviva, P. A. R. Ade, P. Aguirre, M. Amiri, J. W. Appel, L. F. Barrientos, E. S. Battistelli, J. R. Bond, B. Brown, B. Burger, J. Chervenak, S. Das, M. J. Devlin, S. R. Dicker, W. Bertrand Doriese, R. Dünner, T. Essinger-Hileman, R. P. Fisher, J. W. Fowler, A. Hajian, M. Halpern, M. Hasselfield, C. Hernández-Monteagudo, G. C. Hilton, M. Hilton, A. D. Hincks, K. M. Huffenberger, D. H. Hughes, J. P. Hughes, L. Infante, K. D. Irwin, J. B. Juin, M. Kaul, J. Klein, A. Kosowsky, J. M. Lau, M. Limon, Y.-T. Lin, R. H. Lupton, T. A. Marriage, D. Marsden, P. Mauskopf, F. Menanteau, K. Moodley, H. Moseley, C. B. Netterfield, M. D. Niemack, M. R. Nolta, L. A. Page, L. Parker, B. Partridge, B. Reid,

- N. Sehgal, B. Sherwin, D. N. Spergel, S. T. Staggs, D. S. Swetz, E. R. Switzer, R. Thornton,
H. Trac, C. Tucker, R. Warne, E. Wollack, and Y. Zhao, *Astrophys. J.* **739**, 52 (2011),
arXiv:1009.0866.
- [8] R. Keisler, C. L. Reichardt, K. A. Aird, B. A. Benson, L. E. Bleem, J. E. Carlstrom, C. L.
Chang, H. M. Cho, T. M. Crawford, A. T. Crites, T. de Haan, M. A. Dobbs, J. Dudley,
E. M. George, N. W. Halverson, G. P. Holder, W. L. Holzapfel, S. Hoover, Z. Hou, J. D.
Hrubes, M. Joy, L. Knox, A. T. Lee, E. M. Leitch, M. Lueker, D. Luong-Van, J. J. McMahon,
J. Mehl, S. S. Meyer, M. Millea, J. J. Mohr, T. E. Montroy, T. Natoli, S. Padin, T. Plagge,
C. Pryke, J. E. Ruhl, K. K. Schaffer, L. Shaw, E. Shirokoff, H. G. Spieler, Z. Staniszewski,
A. A. Stark, K. Story, A. van Engelen, K. Vanderlinde, J. D. Vieira, R. Williamson, and
O. Zahn, *Astrophys. J.* **743**, 28 (2011), arXiv:1105.3182.
- [9] K. N. Abazajian, K. Arnold, J. Austermann, B. A. Benson, C. Bischoff, J. Bock, J. R. Bond,
J. Borrill, E. Calabrese, J. E. Carlstrom, C. S. Carvalho, C. L. Chang, H. C. Chiang, S. Church,
A. Cooray, T. M. Crawford, K. S. Dawson, S. Das, M. J. Devlin, M. Dobbs, S. Dodelson,
O. Dore, J. Dunkley, J. Errard, A. Fraisse, J. Gallicchio, N. W. Halverson, S. Hanany, S. R.
Hildebrandt, A. Hincks, R. Hlozek, G. Holder, W. L. Holzapfel, K. Honscheid, W. Hu, J. Hub-
mayr, K. Irwin, W. C. Jones, M. Kamionkowski, B. Keating, R. Keisler, L. Knox, E. Komatsu,
J. Kovac, C.-L. Kuo, C. Lawrence, A. T. Lee, E. Leitch, E. Linder, P. Lubin, J. McMahon,
A. Miller, L. Newburgh, M. D. Niemack, H. Nguyen, H. T. Nguyen, L. Page, C. Pryke, C. L.
Reichardt, J. E. Ruhl, N. Sehgal, U. Seljak, J. Sievers, E. Silverstein, A. Slosar, K. M. Smith,
D. Spergel, S. T. Staggs, A. Stark, R. Stompor, A. G. Vieregg, G. Wang, S. Watson, E. J.
Wollack, W. L. K. Wu, K. W. Yoon, and O. Zahn, *ArXiv e-prints* (2013), arXiv:1309.5383.
- [10] Z. Hou, R. Keisler, L. Knox, M. Millea, and C. Reichardt, *Phys. Rev. D* **87**, 083008 (2013),
arXiv:1104.2333 [astro-ph.CO].
- [11] W. L. K. Wu, J. Errard, C. Dvorkin, C. L. Kuo, A. T. Lee, P. McDonald, A. Slosar, and
O. Zahn, *Astrophys. J.* **788**, 138 (2014), arXiv:1402.4108.
- [12] H. K. Jassal, J. S. Bagla, and T. Padmanabhan, *MNRAS* **405**, 2639 (2010), astro-ph/0601389.
- [13] A. Kogut, D. J. Fixsen, D. T. Chuss, J. Dotson, E. Dwek, M. Halpern, G. F. Hinshaw, S. M.
Meyer, S. H. Moseley, M. D. Seiffert, D. N. Spergel, and E. J. Wollack, *JCAP* **7**, 025 (2011),
arXiv:1105.2044.
- [14] L. Wolz, M. Kilbinger, J. Weller, and T. Giannantonio, *JCAP* **9**, 009 (2012), arXiv:1205.3984.

- [15] A. Albrecht, G. Bernstein, R. Cahn, W. L. Freedman, J. Hewitt, W. Hu, J. Huth,
M. Kamionkowski, E. W. Kolb, L. Knox, J. C. Mather, S. Staggs, and N. B. Suntzeff,
ArXiv Astrophysics e-prints (2006), astro-ph/0609591.
- [16] M. Vallisneri, Phys. Rev. D **77**, 042001 (2008), gr-qc/0703086.
- [17] L. Perotto, J. Lesgourgues, S. Hannestad, H. Tu, and Y. Y Y Wong, *JCAP* **10**, 013 (2006),
astro-ph/0606227.
- [18] We thank the authors of [?] for their collaboration.
- [19] Planck Collaboration, P. A. R. Ade, N. Aghanim, C. Armitage-Caplan, M. Arnaud, M. Ash-
down, F. Atrio-Barandela, J. Aumont, C. Baccigalupi, A. J. Banday, and et al., *A&A* **571**,
A16 (2014), arXiv:1303.5076.
- [20] T. Okamoto and W. Hu, Phys. Rev. D **67**, 083002 (2003), astro-ph/0301031.
- [21] W. Hu and T. Okamoto, *Astrophys. J.* **574**, 566 (2002), arXiv:astro-ph/0111606.