

# Toric Code

University of Florida, August 26, 2020

## Recap of last time

- ▶ The toric code:

$$\mathcal{C} = \text{span}\{|\Omega_0\rangle \in \mathcal{H} : A(\mathbf{R}_i)|\Omega_0\rangle = |\Omega_0\rangle, B(\mathbf{R}_i)|\Omega_0\rangle = |\Omega_0\rangle\}$$

- ▶ A *physical* (local) Hamiltonian can have the codeword space as its ground state:  $H = -\sum_{\mathbf{R}_i} (A(\mathbf{R}_i) + B(\mathbf{R}_i))$ .
- ▶ On a torus, the toric code encode 2 qubits.
- ▶ The 2 qubits are manipulated using the non-local operators  $Z_1, Z_2, X_1, X_2$ .
- ▶ The toric codes has an  $e^{-\alpha N}$  probability of making an error.
- ▶ The toric code is said to store the information encoded in the 2 qubits non-locally.
- ▶ Excited states:  $S^z(t)|\Omega_0\rangle$  and  $S^x(t)|\Omega_0\rangle$ . (Open strings.)
- ▶ The end of open strings can be thought of as particles. The toric code have an  $x$ -type particle and a  $z$ -type particle.

# Outline

1. Anyonic nature of the excitations

## The toric code in different geometries


A (surprising) aspect about the toric code is that the ground state degeneracy depends on the genus,  $g$ , of the manifold. The toric code is  $4^g$  degenerate.

## The toric code in different geometries

A (surprising) aspect about the toric code is that the ground state degeneracy depends on the genus,  $g$ , of the manifold. The toric code is  $4^g$  degenerate.

On a *sphere* there are no non-contractible loops.  $A(\mathbf{R}_i)$  and  $B(\mathbf{R}_i)$  can label the entire Hilbert space.

- ▶ Hilbert space is  $2^{12N^2}$  dimensional.
- ▶  $6N^2$   $B(\mathbf{R}_i)$  operators.
- ▶  $6N^2 + 2$   $A(\mathbf{R}_i)$  operators.




## The toric code in different geometries

A (surprising) aspect about the toric code is that the ground state degeneracy depends on the genus,  $g$ , of the manifold. The toric code is  $4^g$  degenerate.

On a *sphere* there are no non-contractible loops.  $A(\mathbf{R}_i)$  and  $B(\mathbf{R}_i)$  can label the entire Hilbert space.


- ▶ Hilbert space is  $2^{12N^2}$  dimensional.
- ▶  $6N^2$   $B(\mathbf{R}_i)$  operators.
- ▶  $6N^2 + 2$   $A(\mathbf{R}_i)$  operators.



The dependence of the ground state degeneracy on the geometry of the manifold is one of the defining features of topological order.


# Particle content of the toric code

- ▶ No particles, 1.
- ▶  $z$ -type particle, referred to as electric charge,  $e$ .
- ▶  $x$ -type particle, referred to as a magnetic vortex,  $m$ .
- ▶ A combinations of an  $e$  and an  $m$  particle,  
 $\psi = e \times m$
- ▶ For convince we drop the lattice from the background, and distinguish different strings by different colors.



# Particle content of the toric code

- ▶ No particles, 1.
- ▶  $z$ -type particle, referred to as electric charge,  $e$ .
- ▶  $x$ -type particle, referred to as a magnetic vortex,  $m$ .
- ▶ A combinations of an  $e$  and an  $m$  particle,  
 $\psi = e \times m$
- ▶ For convince we drop the lattice from the background, and distinguish different strings by different colors.



Next we ask what are the statistics of these particles.

That is we ask What happen when we exchange two  $e$ ,  $m$  or  $\psi$  particles.



## A braid group

To answer the question about statistics we consider processes that physically exchange the two particles and see how the wavefunction changes under such processes.

# A braid group

To answer the question about statistics we consider processes that physically exchange the two particles and see how the wavefunction changes under such processes.

- ▶ It's natural to consider a braid group because of the strings attached to the particles.
- ▶ There are two kinds of strings. The group is then said to be a colored braid group.
- ▶ A state with  $2n_e$ , and  $2n_m$  of  $e$  and  $m$  particles has  $n_e$ , and  $n_m$  red and blue strings.
- ▶ Strings can go above or under each other depending on which operator act first.



# Brading and fusion rules



► The electric charge,  $e$  is a boson.


► The magnetic vortex,  $m$  is a boson.



►  $e$  going around  $m$  gives a  $-1$ .

► These phases can be explained as Aharonov-Bohm effect

►  $\psi$  is a fermion.



Anyons implies the ground state degeneracy.

- ▶ The ground state(s) is a state with no particles in it. If  $|\Omega_0\rangle$  is a ground state, so are  $Z_i |\Omega_0\rangle$  and  $X_i |\Omega_0\rangle$ .
- ▶ Braiding rules implies
  1.  $Z_1^2 = Z_2^2 = X_1^2 = X_2^2 = 1$
  2.  $[Z_1, Z_2] = [X_1, X_2] = 0$
  3.  $\{Z_1, X_1\} = \{Z_2, X_2\} = 0$ .
- ▶ This implies the ground state is fourfold degenerate.

8 / 9

# Summary

- ▶ Quantum codes encode  $k$  qubits into  $n$  qubits.
- ▶ Quantum codes allow for error correction
- ▶ The Kitaev toric code encode  $2g$  qubits into a spin lattice.
- ▶ The Kitaev code has  $e^{-\alpha N}$  probability of missing errors.
- ▶ Toric code encodes the information non-locally.
- ▶ The anyon content of a theory is enough to define its topological order.