

# Antiferromagnetic Spin Chains

University of Florida, April 2020

This talk follows the review by I Affleck 1989 J. Phys.: Condens. Matter 1 3047

# Introduction

The problem at hand is to study whether a 1D antiferromagnetic system is gapped or not.

# Introduction

The problem at hand is to study whether a 1D antiferromagnetic system is gapped or not.

Review of what we know about spin systems and long range order that can help us answer this problem:

# Introduction

The problem at hand is to study whether a 1D antiferromagnetic system is gapped or not.

Review of what we know about spin systems and long range order that can help us answer this problem:

1. Mermin-Wagner theorem, no long range order in 1D.

# Introduction

The problem at hand is to study whether a 1D antiferromagnetic system is gapped or not.

Review of what we know about spin systems and long range order that can help us answer this problem:

1. Mermin-Wagner theorem, no long range order in 1D.
  - ▶ Doesn't imply the existence of a gap. Something like a BKT phase might occur.

# Introduction

The problem at hand is to study whether a 1D antiferromagnetic system is gapped or not.

Review of what we know about spin systems and long range order that can help us answer this problem:

1. Mermin-Wagner theorem, no long range order in 1D.
  - ▶ Doesn't imply the existence of a gap. Something like a BKT phase might occur.
  - ▶ Can only tell us about long range order at finite temperature. Not much about the ground state. In a sense our question is more elementary, it's about the nature of the ground state itself.

# Introduction

The problem at hand is to study whether a 1D antiferromagnetic system is gapped or not.

Review of what we know about spin systems and long range order that can help us answer this problem:

1. Mermin-Wagner theorem, no long range order in 1D.
  - ▶ Doesn't imply the existence of a gap. Something like a BKT phase might occur.
  - ▶ Can only tell us about long range order at finite temperature. Not much about the ground state. In a sense our question is more elementary, it's about the nature of the ground state itself.
2. Goldstone modes, and spin-wave theory.

# Introduction

The problem at hand is to study whether a 1D antiferromagnetic system is gapped or not.

Review of what we know about spin systems and long range order that can help us answer this problem:

1. Mermin-Wagner theorem, no long range order in 1D.
  - ▶ Doesn't imply the existence of a gap. Something like a BKT phase might occur.
  - ▶ Can only tell us about long range order at finite temperature. Not much about the ground state. In a sense our question is more elementary, it's about the nature of the ground state itself.
2. Goldstone modes, and spin-wave theory.
  - ▶ Using the Neel state as the broken symmetry state one might think we should be able to find spin-wave excitations which we know are gapless, and lead to power law correlations.



# Outline

1. Spin-waves for 1D AFM
2. Lieb-Shultz-Mattis theorem
3. Solvable models and valence bond solid states
4. Symmetry of the VBS state

# Outline

1. Spin-waves for 1D AFM
2. Lieb-Shultz-Mattis theorem
3. Solvable models and valence bond solid states
4. Symmetry of the VBS state

# Heisenberg model

For a 1D chain with local spin- $s$  on each site the Heisenberg Hamiltonian can be written as:

$$\begin{aligned} H &= J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} \\ &= J \sum_i S_i^+ S_{i+1}^- + S_i^- S_{i+1}^+ + S_i^z S_{i+1}^z, \end{aligned}$$

with

$$\begin{aligned} S^+ &= \frac{1}{\sqrt{2}} (S^x + iS^y) & S^- &= \frac{1}{\sqrt{2}} (S^x - iS^y), \\ [S^i, S^j] &= i\epsilon^{ijk} S^k & [S^+, S^-] &= S^z \end{aligned}$$

# Heisenberg model

For a 1D chain with local spin- $s$  on each site the Heisenberg Hamiltonian can be written as:

$$\begin{aligned} H &= J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} \\ &= J \sum_i S_i^+ S_{i+1}^- + S_i^- S_{i+1}^+ + S_i^z S_{i+1}^z, \end{aligned}$$

with

$$\begin{aligned} S^+ &= \frac{1}{\sqrt{2}} (S^x + iS^y) & S^- &= \frac{1}{\sqrt{2}} (S^x - iS^y), \\ [S^i, S^j] &= i\epsilon^{ijk} S^k & [S^+, S^-] &= S^z \end{aligned}$$

Ferromagnetism:  $J < 0$

$$|\Psi_{\text{FM}}\rangle = | +s \ +s \ +s \dots \rangle$$

$$H |\Psi_{\text{FM}}\rangle = E_0 |\Psi_{\text{FM}}\rangle$$

$$|\Psi_0\rangle = |\Psi_{\text{FM}}\rangle$$

# Heisenberg model

For a 1D chain with local spin- $s$  on each site the Heisenberg Hamiltonian can be written as:

$$\begin{aligned} H &= J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} \\ &= J \sum_i S_i^+ S_{i+1}^- + S_i^- S_{i+1}^+ + S_i^z S_{i+1}^z, \end{aligned}$$

with

$$\begin{aligned} S^+ &= \frac{1}{\sqrt{2}} (S^x + iS^y) \\ [S^i, S^j] &= i\epsilon^{ijk} S^k \end{aligned}$$

$$\begin{aligned} S^- &= \frac{1}{\sqrt{2}} (S^x - iS^y), \\ [S^+, S^-] &= S^z \end{aligned}$$

Ferromagnetism:  $J < 0$

$$\begin{aligned} |\Psi_{\text{FM}}\rangle &= | +s \ +s \ +s \dots \rangle \\ H |\Psi_{\text{FM}}\rangle &= E_0 |\Psi_{\text{FM}}\rangle \\ |\Psi_0\rangle &= |\Psi_{\text{FM}}\rangle \end{aligned}$$

Anti-ferromagnetism:  $J > 0$

$$\begin{aligned} |\Psi_{\text{Neel}}\rangle &= | +s \ -s \ +s \dots \rangle \\ H |\Psi_{\text{Neel}}\rangle &\neq E_0 |\Psi_{\text{Neel}}\rangle \\ |\Psi_0\rangle &\neq |\Psi_{\text{Neel}}\rangle \end{aligned}$$

# Classical limit in the AFM case

The question then becomes: How close is the Neel state to the ground state?

# Classical limit in the AFM case

The question then becomes: How close is the Neel state to the ground state?

We notice that,

$$\begin{aligned}\langle \Psi_{\text{Neel}} | S_i^+ S_{i+1}^- | \Psi_{\text{Neel}} \rangle &\propto s \\ \langle \Psi_{\text{Neel}} | S_i^z S_{i+1}^z | \Psi_{\text{Neel}} \rangle &\propto s^2.\end{aligned}$$

Hence in the classical limit,  $s \rightarrow \infty$ , one might expect the Neel state to approach the ground state.

## Classical limit in the AFM case

The question then becomes: How close is the Neel state to the ground state?

We notice that,

$$\begin{aligned}\langle \Psi_{\text{Neel}} | S_i^+ S_{i+1}^- | \Psi_{\text{Neel}} \rangle &\propto s \\ \langle \Psi_{\text{Neel}} | S_i^z S_{i+1}^z | \Psi_{\text{Neel}} \rangle &\propto s^2.\end{aligned}$$

Hence in the classical limit,  $s \rightarrow \infty$ , one might expect the Neel state to approach the ground state.

Neel state spontaneously breaks the symmetry and hence would result in Goldstone modes leading to gapless excitations.



# Holstien-Primakov transformation in AFM case

$$S_A^z = s - a^\dagger a \quad S_A^- = a^\dagger \left[ s - \frac{a^\dagger a}{2} \right]^{1/2}$$
$$S_B^z = -s + b^\dagger b \quad S_B^+ = b^\dagger \left[ s - \frac{b^\dagger b}{2} \right]^{1/2}$$

with

$$[a, a^\dagger] = 1 \quad [b, b^\dagger] = 1 \quad [a, b] = 0 \quad [a, b^\dagger] = 0$$



Figure: AB lattice, artificially doubling the unit cell. We put the distance between 2 A sub-lattices to be 1. Let length of the entire chain be  $L$ .

## Classical limit in the AFM case

Expanding around a Neel state, using  $1/s$  as a small parameter.

$$\begin{aligned} S_A^z &= s - a^\dagger a & S_A^- &= \sqrt{s} a^\dagger \\ S_B^z &= -s + b^\dagger b & S_B^+ &= \sqrt{s} b^\dagger \end{aligned}$$

Using this and after Fourier transforming, dropping a constant term, and keeping only terms of order  $O(s)$ ,

$$a_i = \sqrt{\frac{1}{L}} \sum_k e^{ikR_i} a_k \quad b_i = \sqrt{\frac{1}{L}} \sum_k e^{ik(R_i+1/2)} b_k$$

$$H = 2Js \sum_k a_k^\dagger a_k + b_k^\dagger b_k + \cos(k/2) a_k b_{-k} + \cos(k/2) b_{-k}^\dagger a_k^\dagger$$

# Bogoliubov transformation

The Bogoliubov transformation,


$$c_k = u_k a_k - v_k b_{-k}^\dagger$$

$$d_k = u_k b_k - v_k a_{-k}^\dagger$$

$$|u_k|^2 - |v_k|^2 = 1$$

diagonalize the Hamiltonian,

$$H = 2J_s [1 - \cos(k/2)]^{1/2} (c_k^\dagger c_k + d_k^\dagger d_k)$$



**Have we shown that the chain is gapless?**

# What goes wrong?

## What goes wrong?

This approximation only works if the ground state is close to the Neel state. The ground state defined as,

$$c_k |\Psi_0\rangle = 0$$

$$d_k |\Psi_0\rangle = 0$$

## What goes wrong?

This approximation only works if the ground state is close to the Neel state. The ground state defined as,

$$c_k |\Psi_0\rangle = 0$$

$$d_k |\Psi_0\rangle = 0$$

This can be checked by comparing  $\langle S^z \rangle$  in the ground state to that of the Neel state,

$$\langle \Psi_{\text{Neel}} | S_i^z | \Psi_{\text{Neel}} \rangle = \pm s \qquad \langle \Psi_0 | S_i^z | \Psi_0 \rangle = \pm s + \int dk \frac{1}{2} \left[ \frac{1}{|\sin(k)|} - 1 \right]$$

Remember,

**The correction is infinite for any  $s$ .**

$$S_A^- = a^\dagger \left[ s - \frac{a^\dagger a}{2} \right]^{1/2}$$

# Outline

1. Spin-waves for 1D AFM
2. Lieb-Shultz-Mattis theorem
3. Solvable models and valence bond solid states
4. Symmetry of the VBS state

# LSM theory

## Statement of the theorem:

Any half integer spin chain that is symmetric under parity is either gapless or has a degenerate ground state that breaks parity.



# LSM theory


## Statement of the theorem:

Any half integer spin chain that is symmetric under parity is either gapless or has a degenerate ground state that breaks parity.

Construct a state  $|\Psi_1\rangle = \hat{\eta} |\Psi_0\rangle$  with,

$\hat{\eta} = R^z(\{\theta_i\}) = e^{i \sum_{i=-l}^l \theta_i S_i^z}$  (rotation about odd number of local  $z$ -axes)

$$\theta_i = \frac{R_i + l}{l} \pi$$



**Figure:** LSM unitary operator act on only on the part of the chain from  $-l$  to  $l$ .

# LSM theory

We need to show 2 things as  $l \rightarrow \infty$  to show the system is gapless:

1.  $\langle \Psi_1 | H - E_0 | \Psi_1 \rangle \rightarrow 0$  shows it's a low energy excitation. ( $E_0$  is the ground state energy.)
2.  $\langle \Psi_1 | \Psi_0 \rangle \rightarrow 0$  shows that  $|\Psi_1\rangle \neq |\Psi_0\rangle$


$$\langle \Psi_1 | H - E_0 | \Psi_1 \rangle \rightarrow 0$$

First we notice that,

$$R^{z-1}(\theta_i) S_i^\pm R^z(\theta_i) = e^{\pm i\theta_i} S^\pm.$$

Remember

$$H = J \sum_i S_i^+ S_{i+1}^- + S_i^- S_{i+1}^+ + S_i^z S_{i+1}^z$$





$$\langle \Psi_1 | \Psi_0 \rangle \rightarrow 0$$

We study the parity of the state  $|\Psi_1\rangle$  as compared to the parity of the ground state  $|\Psi_0\rangle$ . Let the parity operator be  $\mathcal{P}$ .

$$\mathcal{P}H\mathcal{P}^{-1} = H$$

$$\mathcal{P}|\Psi_0\rangle = \pm |\Psi_0\rangle$$

$$\mathcal{P}|\Psi_1\rangle = \pm \mathcal{P}\eta\mathcal{P}^{-1}|\Psi_0\rangle$$

$$\langle \Psi_1 | \Psi_0 \rangle \rightarrow 0$$

We study the parity of the state  $|\Psi_1\rangle$  as compared to the parity of the ground state  $|\Psi_0\rangle$ . Let the parity operator be  $\mathcal{P}$ .

$$\mathcal{P}H\mathcal{P}^{-1} = H$$

$$\mathcal{P}|\Psi_0\rangle = \pm |\Psi_0\rangle$$

$$\mathcal{P}|\Psi_1\rangle = \pm \mathcal{P}\eta\mathcal{P}^{-1}|\Psi_0\rangle$$

Using  $\mathcal{P}S_i^z\mathcal{P}^{-1} = -S_{-i}^z$ ,

$$\mathcal{P}\eta\mathcal{P}^{-1} = e^{i2\pi\sum_{i=-l}^l S_i^z} \eta = \begin{cases} \eta & s \text{ integer} \\ -\eta & s \text{ half integer} \end{cases}$$

# Outline

1. Spin-waves for 1D AFM
2. Lieb-Shultz-Mattis theorem
3. Solvable models and valence bond solid states
4. Symmetry of the VBS state

# Majumdar-Gosh Model (a warm up)

For a spin-1/2 chain,

$$H = J \sum_i (\vec{S}_i + \vec{S}_{i+1} + \vec{S}_{i+2})^2 = J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} + \frac{1}{2} \vec{S}_i \cdot \vec{S}_{i+2}$$



## Majumdar-Gosh Model (a warm up)

For a spin-1/2 chain,

$$H = J \sum_i (\vec{S}_i + \vec{S}_{i+1} + \vec{S}_{i+2})^2 = J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} + \frac{1}{2} \vec{S}_i \cdot \vec{S}_{i+2}$$

Adding 3 spin-1/2 gives either spin-3/2 or spin-1/2.

## Majumdar-Gosh Model (a warm up)

For a spin-1/2 chain,

$$H = J \sum_i (\vec{S}_i + \vec{S}_{i+1} + \vec{S}_{i+2})^2 = J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} + \frac{1}{2} \vec{S}_i \cdot \vec{S}_{i+2}$$

Adding 3 spin-1/2 gives either spin-3/2 or spin-1/2.

The ground state must have any adjacent 3 sites to have total spin of 1/2.

# Majumdar-Gosh Model (a warm up)

For a spin-1/2 chain,

$$H = J \sum_i (\vec{S}_i + \vec{S}_{i+1} + \vec{S}_{i+2})^2 = J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} + \frac{1}{2} \vec{S}_i \cdot \vec{S}_{i+2}$$

Adding 3 spin-1/2 gives either spin-3/2 or spin-1/2.

The ground state must have any adjacent 3 sites to have total spin of 1/2.

This can be achieved by paring adjacent spins into singlets.

$$\begin{array}{c} \bullet - \bullet \quad \bullet - \bullet \quad \bullet - \bullet \\ |\Psi_0\rangle = |\phi_{\alpha_1} \phi_{\alpha_2} \phi_{\alpha_3} \phi_{\alpha_4} \dots\rangle \epsilon^{\alpha_1 \alpha_2} \epsilon^{\alpha_3 \alpha_4} \dots \end{array}$$

(a)

$$\begin{array}{c} - \bullet \quad \bullet - \bullet \quad \bullet - \bullet \quad \bullet - \\ |\Psi_0\rangle = |\phi_{\alpha_1} \phi_{\alpha_2} \phi_{\alpha_3} \phi_{\alpha_4} \phi_{\alpha_5} \dots\rangle \epsilon^{\alpha_2 \alpha_3} \epsilon^{\alpha_4 \alpha_5} \dots \end{array}$$

(b)

Figure: Two ways adjacent spins can be paired into singlets.

# Majumdar-Gosh Model

Remarks about the ground states:

- ▶ A contraction between 2 spin-1/2 into a spin singlet is called a valence bond. These ground states of the Majumdar-Gosh model are valence bond states with all the spins contracted.

# Majumdar-Gosh Model

Remarks about the ground states:

- ▶ A contraction between 2 spin-1/2 into a spin singlet is called a valence bond. These ground states of the Majumdar-Gosh model are valence bond states with all the spins contracted.
- ▶ There are two ground states in consistency with LSM theorem.

# Majumdar-Gosh Model

Remarks about the ground states:

- ▶ A contraction between 2 spin-1/2 into a spin singlet is called a valence bond. These ground states of the Majumdar-Gosh model are valence bond states with all the spins contracted.
- ▶ There are two ground states in consistency with LSM theorem.
- ▶ The ground states breaks translational symmetry. These states are fully dimerized.

# Majumdar-Gosh Model

Remarks about the ground states:

- ▶ A contraction between 2 spin-1/2 into a spin singlet is called a valence bond. These ground states of the Majumadar-Gosh model are valence bond states with all the spins contracted.
- ▶ There are two ground states in consistency with LSM theorem.
- ▶ The ground states breaks translational symmetry. These states are fully dimerized.
- ▶ These states has a very short correlation length,

$$\langle \vec{S}_i \cdot \vec{S}_j \rangle = 0 \quad \text{for } |R_i - R_j| > 1$$

# Majumdar-Gosh Model

Remarks about the ground states:

- ▶ A contraction between 2 spin-1/2 into a spin singlet is called a valence bond. These ground states of the Majumadar-Gosh model are valence bond states with all the spins contracted.
- ▶ There are two ground states in consistency with LSM theorem.
- ▶ The ground states breaks translational symmetry. These states are fully dimerized.
- ▶ These states has a very short correlation length,

$$\langle \vec{S}_i \cdot \vec{S}_j \rangle = 0 \quad \text{for } |R_i - R_j| > 1$$

- ▶ This suggest the existence of a gap.



# Valence bond states

The motivation for the next model came from trying to find a spin-1 Hamiltonian that has a valence bond state as its ground state.

# Valence bond states

The motivation for the next model came from trying to find a spin-1 Hamiltonian that has a valence bond state as its ground state.

We need a way to generalize valence bond states to higher spins

# Valence bond states

The motivation for the next model came from trying to find a spin-1 Hamiltonian that has a valence bond state as its ground state.

We need a way to generalize valence bond states to higher spins  
For spin-1 we think of each spin site as being made of 2 spin-1/2s.  
And always construct states that are symmetric in same site spin-1/2s.

# Valence bond states

The motivation for the next model came from trying to find a spin-1 Hamiltonian that has a valence bond state as its ground state.

We need a way to generalize valence bond states to higher spins  
For spin-1 we think of each spin site as being made of 2 spin-1/2s.  
And always construct states that are symmetric in same site spin-1/2s.

The following can be used as a basis on each site,

$$|\phi_{\alpha_1\alpha_2}\rangle = |\phi_{\alpha_2\alpha_1}\rangle = \frac{1}{\sqrt{2}}(|\phi_{\alpha_1}\phi_{\alpha_2}\rangle + |\phi_{\alpha_2}\phi_{\alpha_1}\rangle)$$
$$\langle\phi_{\alpha_1\alpha_2}|\phi_{\beta_1\beta_2}\rangle = \delta_{\alpha_1\beta_1}\delta_{\alpha_2\beta_2} + \delta_{\alpha_1\beta_2}\delta_{\alpha_2\beta_1}$$

# Valence bond states

Let's look at the possible valence bond states of spin-1 chain.

$$\begin{array}{c}
 \bullet - \bullet - \bullet - \bullet - \bullet - \bullet - \bullet - \bullet - \bullet \\
 |\Psi_{\alpha_1 \alpha_{2L}}^{\text{VBS}}\rangle = |\phi_{\alpha_1 \alpha_2} \phi_{\alpha_3 \alpha_4} \phi_{\alpha_5 \alpha_6} \phi_{\alpha_7 \alpha_8} \dots\rangle \epsilon^{\alpha_2 \alpha_3} \epsilon^{\alpha_4 \alpha_5} \dots
 \end{array}
 \quad (a)$$

$$\begin{array}{c}
 \bullet \frown \bullet \quad \bullet \frown \bullet \quad \bullet \frown \bullet \\
 |\Psi^{\text{VB1}}\rangle = |\phi_{\alpha_1 \alpha_2} \phi_{\alpha_3 \alpha_4} \phi_{\alpha_5 \alpha_6} \phi_{\alpha_7 \alpha_8} \dots\rangle \epsilon^{\alpha_1 \alpha_3} \epsilon^{\alpha_2 \alpha_4} \dots
 \end{array}
 \quad (b)$$

$$\begin{array}{c}
 \smile \bullet \quad \bullet \frown \bullet \quad \bullet \frown \bullet \quad \bullet \smile \\
 |\Psi_{\alpha_1 \alpha_2 \alpha_{2L-1} \alpha_{2L}}^{\text{VB2}}\rangle = |\phi_{\alpha_1 \alpha_2} \phi_{\alpha_3 \alpha_4} \phi_{\alpha_5 \alpha_6} \phi_{\alpha_7 \alpha_8} \dots\rangle \epsilon^{\alpha_3 \alpha_5} \epsilon^{\alpha_4 \alpha_6} \dots
 \end{array}
 \quad (c)$$

**Figure:** 3 possible valence bond states. (a) is the VBS partially dimerized state. (b) and (c) are the fully dimerized states.

# AKLT model

The AKLT model is a spin-1 chain model with the VBS state being the ground state,

$$\begin{aligned} H &= 2J \sum_i P^{(2)}(\vec{S}_i + \vec{S}_{i+1}) \\ &= J \sum_i \frac{1}{12} (\vec{S}_i + \vec{S}_{i+1})^2 ((\vec{S}_i + \vec{S}_{i+1})^2 - 2) \\ &= J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} + \frac{1}{3} (\vec{S}_i \cdot \vec{S}_{i+1})^2 + \text{const.} \end{aligned}$$

## AKLT model

The AKLT model is a spin-1 chain model with the VBS state being the ground state,

$$\begin{aligned} H &= 2J \sum_i P^{(2)}(\vec{S}_i + \vec{S}_{i+1}) \\ &= J \sum_i \frac{1}{12} (\vec{S}_i + \vec{S}_{i+1})^2 ((\vec{S}_i + \vec{S}_{i+1})^2 - 2) \\ &= J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} + \frac{1}{3} (\vec{S}_i \cdot \vec{S}_{i+1})^2 + \text{const.} \end{aligned}$$

This Hamiltonian is positive definite, the ground state will have  $E_0 = 0$ .

# AKLT model

The AKLT model is a spin-1 chain model with the VBS state being the ground state,

$$\begin{aligned} H &= 2J \sum_i P^{(2)}(\vec{S}_i + \vec{S}_{i+1}) \\ &= J \sum_i \frac{1}{12} (\vec{S}_i + \vec{S}_{i+1})^2 ((\vec{S}_i + \vec{S}_{i+1})^2 - 2) \\ &= J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} + \frac{1}{3} (\vec{S}_i \cdot \vec{S}_{i+1})^2 + \text{const.} \end{aligned}$$

This Hamiltonian is positive definite, the ground state will have  $E_0 = 0$ .

The VBS state has for every 2 adjacent spin sites 2 of the 4 spin-1/2s to be contracted into a spin singlet.



## AKLT model

The AKLT model is a spin-1 chain model with the VBS state being the ground state,

$$\begin{aligned} H &= 2J \sum_i P^{(2)}(\vec{S}_i + \vec{S}_{i+1}) \\ &= J \sum_i \frac{1}{12} (\vec{S}_i + \vec{S}_{i+1})^2 ((\vec{S}_i + \vec{S}_{i+1})^2 - 2) \\ &= J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} + \frac{1}{3} (\vec{S}_i \cdot \vec{S}_{i+1})^2 + \text{const.} \end{aligned}$$

This Hamiltonian is positive definite, the ground state will have  $E_0 = 0$ .

The VBS state has for every 2 adjacent spin sites 2 of the 4 spin-1/2s to be contracted into a spin singlet.

Hence the total spin of any 2 adjacent spin sites cannot be 2 with only 2 spin-1/2s left. This shows the VBS state to be a ground state.

# AKLT model

$$H |\Psi_{\text{VBS}}\rangle = 0$$

# AKLT model

$$H |\Psi_{\text{VBS}}\rangle = 0$$

What is more subtle is to show that this is the only ground state!

# AKLT model

$$H |\Psi_{\text{VBS}}\rangle = 0$$

What is more subtle is to show that this is the only ground state!  
It is the unique ground state up to some boundary modification.

# AKLT model

$$H |\Psi_{\text{VBS}}\rangle = 0$$


What is more subtle is to show that this is the only ground state!  
It is the unique ground state up to some boundary modification.

$$\begin{array}{c} -\bullet- \bullet- \bullet- \bullet- \bullet- \bullet- \\ |\Psi_{\alpha_1 \alpha_{2L}}^{\text{VBS}}\rangle \end{array}$$

# AKLT model

$$H |\Psi_{\text{VBS}}\rangle = 0$$

What is more subtle is to show that this is the only ground state!  
It is the unique ground state up to some boundary modification.



A horizontal line with six black dots representing sites. The line starts and ends with a short horizontal segment, indicating open boundaries.

$$|\Psi_{\alpha_1 \alpha_{2L}}^{\text{VBS}}\rangle$$

One can get around this by considering an infinite system, or to be more concrete, study the system under periodic boundary conditions.



A horizontal line with six black dots representing sites. A curved line connects the first and last dots, representing periodic boundary conditions.

$$|\Psi^{\text{VBS}}\rangle$$

## Correlation length of the VBS state

$$\vec{S}|\phi_{\alpha_1\alpha_2}\rangle = \frac{1}{2}(\vec{\sigma}_{\beta\alpha_1}|\phi_{\beta\alpha_2}\rangle + \vec{\sigma}_{\beta\alpha_2}|\phi_{\alpha_1\beta}\rangle)$$

$$\langle\vec{S}_i\cdot\vec{S}_j\rangle = 4(-1)^{R_j-R_i}3^{-|R_j-R_i|}$$

$$\xi = \frac{1}{\log 3} = 0.9$$

## Correlation length of the VBS state

$$\vec{S}|\phi_{\alpha_1\alpha_2}\rangle = \frac{1}{2}(\vec{\sigma}_{\beta\alpha_1}|\phi_{\beta\alpha_2}\rangle + \vec{\sigma}_{\beta\alpha_2}|\phi_{\alpha_1\beta}\rangle)$$

$$\langle\vec{S}_i\cdot\vec{S}_j\rangle = 4(-1)^{R_j-R_i}3^{-|R_j-R_i|}$$

$$\xi = \frac{1}{\log 3} = 0.9$$

Finite length correlation function suggests the existence of a gap.



## Correlation length of the VBS state

$$\vec{S}|\phi_{\alpha_1\alpha_2}\rangle = \frac{1}{2}(\vec{\sigma}_{\beta\alpha_1}|\phi_{\beta\alpha_2}\rangle + \vec{\sigma}_{\beta\alpha_2}|\phi_{\alpha_1\beta}\rangle)$$

$$\langle\vec{S}_i \cdot \vec{S}_j\rangle = 4(-1)^{R_j-R_i}3^{-|R_j-R_i|}$$

$$\xi = \frac{1}{\log 3} = 0.9$$

Finite length correlation function suggests the existence of a gap. Numerical results shows that the VBS state has energy that is only 5% higher than the ground state of the Heisenberg Hamiltonian.

## Correlation length of the VBS state

$$\vec{S}|\phi_{\alpha_1\alpha_2}\rangle = \frac{1}{2}(\vec{\sigma}_{\beta\alpha_1}|\phi_{\beta\alpha_2}\rangle + \vec{\sigma}_{\beta\alpha_2}|\phi_{\alpha_1\beta}\rangle)$$

$$\langle\vec{S}_i \cdot \vec{S}_j\rangle = 4(-1)^{R_j-R_i}3^{-|R_j-R_i|}$$

$$\xi = \frac{1}{\log 3} = 0.9$$

Finite length correlation function suggests the existence of a gap. Numerical results shows that the VBS state has energy that is only 5% higher than the ground state of the Heisenberg Hamiltonian. Correlation length does agree as much. Numerical results suggests  $\xi = 5$ .

## Trivia

Looking on the more general Hamiltonian,

$$H = J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} - \beta (\vec{S}_i \cdot \vec{S}_{i+1})^2$$

there is another solvable point at  $\beta = 1$ , which is gapless.

## Trivia

Looking on the more general Hamiltonian,

$$H = J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} - \beta (\vec{S}_i \cdot \vec{S}_{i+1})^2$$

there is another solvable point at  $\beta = 1$ , which is gapless.  
The model with  $\beta = 1$  was solved by Bethe in the 1930s.

# Trivia

Looking on the more general Hamiltonian,

$$H = J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} - \beta (\vec{S}_i \cdot \vec{S}_{i+1})^2$$

there is another solvable point at  $\beta = 1$ , which is gapless.

The model with  $\beta = 1$  was solved by Bethe in the 1930s.

A phase transition  
occur at  $\beta = 1$ .

# Trivia

Looking on the more general Hamiltonian,


$$H = J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} - \beta (\vec{S}_i \cdot \vec{S}_{i+1})^2$$

there is another solvable point at  $\beta = 1$ , which is gapless.

The model with  $\beta = 1$  was solved by Bethe in the 1930s.

A phase transition occur at  $\beta = 1$ .

The graphs suggests that the phase transition occur between the VBS and the dimerized state.



# Outline

1. Spin-waves for 1D AFM
2. Lieb-Shultz-Mattis theorem
3. Solvable models and valence bond solid states
4. Symmetry of the VBS state

## Symmetry of VBS state

No local order parameter characterizes the transition to the VBS state.



# Symmetry of VBS state

No local order parameter characterizes the transition to the VBS state.

The periodic VBS chain preserve all local symmetries of the Hamiltonian like rotation, inversion, translation, and time reversal symmetry.

## Symmetry of VBS state

No local order parameter characterizes the transition to the VBS state.

The periodic VBS chain preserve all local symmetries of the Hamiltonian like rotation, inversion, translation, and time reversal symmetry.

This here is an example of a symmetry protected topology. Let's study the VBS state with open boundary condition.

$$\text{spin-1/2} \quad -\bullet-\bullet-\bullet-\bullet-\bullet-\bullet-\bullet-\quad \text{spin-1/2}$$

The state has 2 "free" spin-1/2, one at each end of the chain, that transforms differently as compared to the bulk.

# Symmetry of VBS state

No local order parameter characterizes the transition to the VBS state.

The periodic VBS chain preserve all local symmetries of the Hamiltonian like rotation, inversion, translation, and time reversal symmetry.

This here is an example of a symmetry protected topology. Let's study the VBS state with open boundary condition.

$$\text{spin-1/2} \quad -\bullet-\bullet-\bullet-\bullet-\bullet-\bullet-\bullet-\quad \text{spin-1/2}$$

The state has 2 "free" spin-1/2, one at each end of the chain, that transforms differently as compared to the bulk.

Such a weird situation at the boundaries can only be changed by a transformations that mixes the boundaries.

# Symmetry of VBS state

No local order parameter characterizes the transition to the VBS state.

The periodic VBS chain preserve all local symmetries of the Hamiltonian like rotation, inversion, translation, and time reversal symmetry.

This here is an example of a symmetry protected topology. Let's study the VBS state with open boundary condition.

$$\text{spin-1/2} \quad -\bullet-\bullet-\bullet-\bullet-\bullet-\bullet-\bullet-\quad \text{spin-1/2}$$

The state has 2 "free" spin-1/2, one at each end of the chain, that transforms differently as compared to the bulk.

Such a weird situation at the boundaries can only be changed by a transformations that mixes the boundaries.

Such along range correlation between the ends of the chain can only happen if the bulk closes a gap.