



Puberty-Related Maturation of Adolescent Fronto-Striatal Resting-State Functional Connectivity is Implicated in the Development of Inhibitory Control

Amar Ojha^{1,2}; Ashley Parr^{2,3}; Will Foran³; Finnegan Calabro^{2,3,4}; Beatriz Luna^{1,2,3,5}

1. Center for Neuroscience; 2. Center for Neural Basis of Cognition; 3. Department of Psychiatry;
4. Department of Bioengineering; 5. Department of Psychology, University of Pittsburgh.



Background & Motivation

Adolescent Biobehavioral Development


- Puberty defines adolescence, the transition to adulthood characterized by widespread biological, cognitive, and behavioral maturation.
- By the end of adolescence, individuals are better able to reliably engage brain circuitry supporting goal-directed cognitive processes¹.
- Central to these cognitive processes is specialization of fronto-striatal systems².
- However, the extent to which puberty contributes to maturation of this circuitry beyond age-related effects is unclear³⁻⁶.

Hypothesis

Based on the literature, we expect that puberty will be associated with NAcc medial prefrontal cortex (PFC) resting-state functional connectivity (rsFC). It is unclear how puberty will be associated with lateral PFC.


Study Design & Analyses

- We harmonized two datasets of 110 adolescents (52 females, 266 total scans, ages 10-18, 1-3 visits) imaged on 3T MRI scanners who had started puberty.
- Puberty was assessed using the Petersen Puberty Development Scale (PDS)⁷, a self-report puberty measure; PDS scores were transformed to a 5-point Tanner scale and participants were only included in analyses if scores were ≥ 2 .
- Response inhibition was assessed using mean latencies and accuracy rates from correct antisaccade task trials.
- AIC values were used to determine the best functional form of puberty (along with linear, inverse, and quadratic models) to test fronto-striatal rsFCs associated with puberty beyond age effects separately in males and females.
- We applied Bonferroni corrections ($p < .0083$) and covaried for age across models.




Results

Puberty and Inhibitory Control




- Inhibitory responses improved with age (accuracy: $\beta = .43$, $t = 5.20$, $p < .001$; latency: $\beta = -.38$, $t = -4.12$, $p < .001$) but not with puberty when controlling for age ($p > .05$).

Puberty and PFC-NAcc rsFC



- Puberty was associated with greater dIPFC-NAcc rsFC in males ($\beta = .18$, $t = 3.46$, $p_{\text{Bonf}} = .007$) and females ($\beta = 1.55$, $t = 3.26$, $p_{\text{Bonf}} = .015$); in females, puberty was also associated with vIPFC-NAcc rsFC ($\beta = .30$, $t = 4.99$, $p_{\text{Bonf}} < .001$).

Puberty and PFC-NAcc rsFC and Inhibitory Control



- Puberty and antisaccade accuracy interacted to predict dIPFC-NAcc rsFC ($\beta = .10$, $t = 2.04$, $p = .041$), such that stronger rsFC was associated with worse performance before mid-puberty.
- Puberty and correct AS trial latency interacted to predict vIPFC-NAcc rsFC ($\beta = -.14$, $t = -2.51$, $p = .012$), such that in early puberty, greater connectivity was associated with longer/slower latency.

Discussion

- Puberty was uniquely associated with increases in dIPFC-NAcc rsFC but not with medial PFC subregions.
- Females showed an additional association between puberty and vIPFC-NAcc rsFC, which may reflect their earlier pubertal maturation or changes in unique sex hormones.
- Maturation of these two fronto-striatal connections were related to antisaccade accuracy and correct trial latency, respectively.
- Taken together, these results show that beyond chronological age, pubertal maturation may play a significant role in strengthening critical fronto-striatal connectivity shown to support the maturation of cognitive control through adolescence and into adulthood.

Future Directions

- Analyzing hormonal assays can test possible mechanisms underlying the puberty-related neurobiological changes in functional development of fronto-striatal systems.
- Including younger ages (in females) and older ages (in males) will help to better characterize the effects of pubertal maturation on neurocognitive development.
- Assessing pubertal tempo—one's pubertal maturation relative to same-age, same-sex peers—alongside pubertal timing may yield additional important inter-individual differences in relative rates of development.
- Characterizing the effects of puberty on additional mesocorticolimbic ROIs (e.g., amygdala nuclei) supporting goal-directed (motivated) behaviors can inform neurobiological markers associated with risk of adolescent psychopathologies⁸⁻¹¹.

References

1. Ordaz, S. J., Foran, W., Velanova, K., & Luna, B. (2013). Longitudinal growth curves of brain function underlying inhibitory control through adolescence. *Journal of Neuroscience*, 33(46), 18109-18124.
2. Parr, A. C., Calabro, F., Larsen, B., Tervo-Clemmens, B., Elliot, S., Foran, W., ... & Luna, B. (2021). Dopamine related striatal neurophysiology is associated with specialization of frontostriatal reward circuitry through adolescence. *Progress in Neurobiology*, 201, 101997.
3. van den Bos, W., Cohen, M. X., Kahnt, T., Crone, E. A., 2012. Striatum—Medial Prefrontal Cortex Connectivity Predicts Developmental Changes in Reinforcement Learning. *Cerebral Cortex*, 22(6), 1247-1255.
4. Van Duijvenvoorde, A. C., Westhoff, B., de Vos, F., Wierenga, L. M., & Crone, E. A. (2019). A three-wave longitudinal study of subcortical-cortical resting-state connectivity in adolescence: Testing age- and puberty-related changes. *Human Brain Mapping*, 40(13), 3769-3783.
5. Faraoni, D. S., Gabard-Durnam, L., Goff, B., Flannery, J., Gee, D. G., Lumian, D. S., Caldera, C., Tottenham, N. (2015). Normative development of ventral striatal resting state connectivity in humans. *NeuroImage*, 118, 422-437.
6. Harsay, H. A., Cohen, M. X., Oosterhof, N. N., Forstmann, B. U., Mars, R. B., Ridderinkhof, K. R. (2011). Functional connectivity of the striatum links motivation to action control in humans. *Journal of Neuroscience*, 31(29), 10701-11.
7. Petersen, A. C., Crockett, L., Richards, M., & Boxer, A. (1988). A self-report measure of pubertal status: Reliability, validity, and initial norms. *Journal of Youth and Adolescence*, 17(2), 117-133.
8. Green, I. W., Pizzagalli, D. A., Admon, R., & Kumar, P. (2019). Anhedonia modulates the effects of positive mood induction on reward-related brain activation. *NeuroImage*, 193, 115-125.
9. Weissman, D. G., Schriber, R. A., Fassbender, C., Atherton, O., Krafft, C., Robins, R. W., ... & Guyer, A. E. (2015). Earlier adolescent substance use onset predicts stronger connectivity between reward and cognitive control brain networks. *Developmental Cognitive Neuroscience*, 16, 121-129.
10. Wang, C., Huang, P., Shen, Z., Qian, W., Wang, S., Jiaerken, Y., ... & Zhang, M. (2021). Increased striatal functional connectivity is associated with improved smoking cessation outcomes: A preliminary study. *Addiction Biology*, 26(2), e12919.
11. Ballard, I. C., Murty, V. P., Carter, R. M., MacInnes, J. J., Huettel, S. A., & Adcock, R. A. (2011). Dorsolateral prefrontal cortex drives mesolimbic dopaminergic regions to initiate motivated behavior. *Journal of Neuroscience*, 31(28), 10340-10346.

Acknowledgements

The authors thank the participants and their families for participating in the studies and the staff and students at the Laboratory of Neurocognitive Development (LNCD) for making this research possible. This research was supported by funding from Staunton Farm Foundation (BL), R01MH080243 (BL), R01MH067924 (BL), and T32GM081760 (AO).