



# Pubertal Maturation Influences Mesocorticolimbic Development in Youth: Implications for Sensitivity to Rewards and Punishment

Amar Ojha<sup>1,2</sup>; Neil P. Jones<sup>3</sup>; Elizabeth A. Shirtcliff<sup>4</sup>; Cecile D. Ladouceur<sup>1,2,3,5</sup>

1. Center for Neuroscience; 2. Center for Neural Basis of Cognition; 3. Department of Psychiatry, University of Pittsburgh;  
4. Human Development & Family Studies, Iowa State University; 5. Department of Psychology, University of Pittsburgh




## BACKGROUND

- Puberty initiates neuroendocrinological changes in adolescent socio-affective processing and behavior<sup>1</sup>.
- Puberty-related shifts in sensitivity to reward or punishment might underlie heightened risk for psychopathology<sup>2</sup>.
- It remains unknown how various aspects of puberty influence the maturation of neural circuitry underlying reward/punishment sensitivity.

**Hypothesis:** We predict that fronto-striatal connectivity will support reward sensitivity<sup>3</sup> whereas fronto-amygdala connectivity will support punishment sensitivity<sup>4</sup>. Pubertal maturation will moderate the association.

## METHODS

- Sample: 126 adolescents studied longitudinally (216 scans; approx. 2-year interval).
- Acquired resting-state fMRI scan (two 6-min sessions).
- Indexed resting-state functional connectivity (RSFC) by taking the PCA across all ROI voxels; Pearson correlation coefficients were computed between ROIs and normalized using Fisher's Z transformation.
- Puberty was assessed via the Puberty Development Scale (PDS)<sup>5</sup>, which was transformed to 5-scale Tanner ratings.
- Parent-report SPSRQ (max subscore = 24) was used to assess sensitivity to rewards/punishments<sup>6</sup>.
- Statistical tests were FDR-corrected to account for multiple comparisons and control for age effects.



## RESULTS

### Reward sensitivity is supported by fronto-striatal connectivity.


Weaker ant.vmpFC-NAcc rsFC is associated with greater reward sensitivity ( $\beta = -0.03$ ,  $p_{FDR} = .007$ ).

### Punishment sensitivity is supported by both fronto-striatal and fronto-amygdala connectivity.

Stronger fronto-striatal/amygdala rsFC is associated with punishment sensitivity (ant.vmpFC-NAcc:  $\beta = 0.03$ ,  $p_{FDR} = .003$ ; ant.vmpFC-amyg:  $\beta = 0.03$ ,  $p_{FDR} = .015$ ; dlPFC-NAcc:  $\beta = 0.05$ ,  $p_{FDR} = .0002$ ; dlPFC-amyg:  $\beta = 0.03$ ,  $p_{FDR} = .007$ ; rACC-NAcc:  $\beta = 0.03$ ,  $p_{FDR} = .003$ ; rACC-amyg:  $\beta = 0.02$ ,  $p_{FDR} = .046$ ; vACC-NAcc:  $\beta = 0.02$ ,  $p_{FDR} = .045$ ; viPFC-NAcc:  $\beta = 0.03$ ,  $p_{FDR} = .003$ ; viPFC-amyg:  $\beta = 0.03$ ,  $p_{FDR} = .003$ ).

### Puberty significantly moderates the relationship between reward/punishment sensitivity and rsFC in boys but not girls.

Pubertal maturation is associated with a weakening of the relationship between ant.vmpFC-NAcc RSFC and reward sensitivity in boys ( $\beta = 0.06$ ,  $p_{FDR} < .001$ ). In contrast, pubertal maturation is associated with a strengthening of the relationship between rACC/viPFC-amygdala RSFC and punishment sensitivity in boys (rACC:  $\beta = 0.04$ ,  $p_{FDR} = .006$ ; viPFC:  $\beta = 0.05$ ,  $p_{FDR} < .001$ ).



## CONCLUSIONS

- Puberty significantly influences functional mesocorticolimbic circuitry underlying sensitivity to rewards/punishments, particularly in boys, which has been relatively understudied in the literature.
- Such pubertal influences on mesocorticolimbic circuitry may underscore a neurobiological process by which puberty confers risk for affective disorders differently in boys and girls.
- Future work should investigate hormonal changes as they relate to sensitivity to reward/punishment and increasing risk of anxiety/depression.

## REFERENCES

1. Vijayakumar, N., de Macks, Z. O., Shirtcliff, E. A., & Pfeifer, J. H. (2018). Puberty and the human brain: Insights into adolescent development. *Neuroscience & Biobehavioral Reviews*, 92, 417-436.
2. Ladouceur, C. D. (2012). Neural systems supporting cognitive-affective interactions in adolescence: the role of puberty and implications for affective disorders. *Frontiers in Integrative Neuroscience*, 6, 65.
3. Parr, A. C., Calabro, F., Larsen, B., Tervo-Clemmens, B., Elliot, S., Foran, W., ... & Luna, B. (2021). Dopamine-related striatal neurophysiology is associated with specialization of frontostriatal reward circuitry through adolescence. *Progress in Neurobiology*, 201, 101997.
4. Sequeira, S. L., Rosen, D. K., Silk, J. S., Hutchinson, E., Allen, K. B., Jones, N. P., ... & Ladouceur, C. D. (2021). "Don't judge me!": Links between in vivo attention bias toward a potentially critical judge and fronto-amygdala functional connectivity during rejection in adolescent girls. *Developmental Cognitive Neuroscience*, 49, 100960.
5. Petersen, A. C., Crockett, L., Richards, M., & Boxer, A. (1988). A self-report measure of pubertal status: Reliability, validity, and initial norms. *Journal of Youth and Adolescence*, 17(2), 117-133.
6. Torrubia, R., Ávila, C., Moltó, J., & Caseras, X. (2001). The Sensitivity to Punishment and Sensitivity to Reward Questionnaire (SPSRQ) as a measure of Gray's anxiety and impulsivity dimensions. *Personality and Individual Differences*, 31(6), 837-862.

## FUNDING

Research funded by NIMH R01MH099007 (PI: Ladouceur)

## ACKNOWLEDGEMENTS

The authors would like to thank the study participants and the staff of the CAN-D Lab who made this research possible.

