

Solving PDEs with Denoising Diffusion Restoration Models

Application to the Poisson Equation

Marty Mukherjee^{*1} Melissa Stadt^{*1} Jun Liu¹ Anita Layton¹

Abstract

This document provides a basic paper template and submission guidelines. Abstracts must be a single paragraph, ideally between 4–6 sentences long. Gross violations will trigger corrections at the camera-ready phase.

1. Introduction

2. Background

3. Theoretical Findings

3.1. Distribution of u

Assumption 3.1. f is Lipschitz in the domain Ω with Lipschitz constant L_f .

Consider the following blind inverse problem

$$f = \Delta u + z \quad (1)$$

where $z \sim N(0, \sigma_f^2 L_f^2)$ where σ_f is a known quantity and L_f is the Lipschitz constant of f .

Theorem 3.2. If $f = \Delta u + z$ where $z \sim N(0, \sigma_f^2 L_f^2)$, then

$$u(x, y) \sim N(u_0, \sigma_f^2 L_f^2 K(x, y)), \quad (2)$$

where

$$K(x, y) = \iint_{\Omega} (\psi((x', y') - (x, y)))^2 dx' dy', \quad (3)$$

where $\psi(\cdot)$ is the Green's function in two dimensions

$$\psi(x, y) = \frac{\ln(\|(x, y)\|)}{2\pi} \quad (4)$$

^{*}Equal contribution ¹Department of Applied Mathematics, University of Waterloo, Waterloo, Ontario, Canada. Correspondence to: Marty Mukherjee <a29mukhe@uwaterloo.ca>, Melissa Stadt <mstadt@uwaterloo.ca>.

3.2. Eigenvalues and Eigenfunctions of the Laplacian operator

Theorem 3.3. The eigenfunction and eigenvalue pairs of the Laplacian operator Δ in a domain $\Omega = [0, 1]^2$ subject to the boundary conditions $u = 0$ on $\partial\Omega$ are of the form

$$u_{n,m}(x, y) = \sin(n\pi x) \sin(m\pi y) \quad (5)$$

$$\lambda_{n,m} = -(n\pi)^2 - (m\pi)^2 \quad (6)$$

4. DDRM for solving PDEs

Sampling of x_T .

We consider the fast Fourier transform (FFT) of x_t and f and perform the diffusion in its spectral space. Define $\bar{x}_t^{(n,m)}$ and $\bar{f}^{(n,m)}$ as follows:

$$\bar{x}_t^{(n,m)} = \langle x_t, \sin(n\pi x) \sin(m\pi y) \rangle, \quad (7)$$

$$\bar{f}^{(n,m)} = \langle f, \sin(n\pi x) \sin(m\pi y) \rangle / \lambda_{n,m}. \quad (8)$$

Since none of the eigenvalues $\lambda_{m,n}$ are zero, our DDRM method for sampling x_T is as follows:

$$p_{\theta}^{(T)}(\bar{x}_t^{(n,m)} \mid \mathbf{f}) = \mathcal{N}(\bar{\mathbf{f}}^{(n,m)}, \sigma_T^2 - L_f^2 \sigma_y^2 / \lambda_{n,m}^2) \quad (9)$$

Choose $\sigma_T > \sigma_y / (\pi\sqrt{2})$.

Sampling of x_t .

$$p_{\theta}^{(t)}(\bar{x}_t^{(n,m)} \mid \mathbf{x}_{t+1}, \mathbf{f}) = \quad (10)$$

$$\begin{cases} \mathcal{N}\left(\bar{x}_{\theta,t}^{(n,m)} + \sqrt{1 - \eta^2} \sigma_t \frac{\bar{\mathbf{f}}^{(n,m)} - \bar{x}_{\theta,t}^{(n,m)}}{\sigma_{\mathbf{f}}/s_i}, \eta^2 \sigma_t^2\right), & \sigma_t < \frac{\sigma_{\mathbf{f}}}{s_i} \\ \mathcal{N}\left((1 - \eta_b) \bar{x}_{\theta,t}^{(n,m)} + \eta_b \bar{\mathbf{f}}^{(n,m)}, \sigma_t^2 - \frac{\sigma_{\mathbf{f}}^2}{s_i^2} \eta_b^2\right), & \sigma_t \geq \frac{\sigma_{\mathbf{f}}}{s_i} \end{cases} \quad (11)$$

5. Numerical Results

6. Conclusion

Acknowledgements

The authors would like to thank Professor Mohammad Kohandel from the Department of Applied Mathematics, University of Waterloo, and Lena Podina from the Department of Computer Science, University of Waterloo for providing us with feedback for this paper.

References

Gilbarg, D. and Trudinger, N. S. *Elliptic Partial Differential Equations of Second Order, Second Edition*. Springer-Verlag, 1983.

A. Proof of theorem 3.2

Proof. (Gilbarg & Trudinger, 1983) In a bounded domain $\Omega \subset \mathbb{R}^2$ with smooth boundary, any function $u \in C^2(\bar{\Omega})$ satisfies

$$\begin{aligned} u(x, y) &= \iint_{\Omega} \psi((x', y') - (x, y)) \Delta u(x', y') dx' dy' \\ &+ \int_{\partial\Omega} u(x', y') \frac{\partial \psi}{\partial n}((x', y') - (x, y)) dS_{x', y'} \\ &- \int_{\partial\Omega} \psi((x', y') - (x, y)) \frac{\partial u}{\partial n}(x', y') dS_{x', y'}. \end{aligned}$$

Given that $\Delta u = f + z$ with $z \sim N(0, \sigma_y^2 L_f^2)$, the first integrand can be written as

$$\begin{aligned} &\iint_{\Omega} \psi((x', y') - (x, y)) \Delta u(x', y') dx' dy' \\ &= \iint_{\Omega} \psi((x', y') - (x, y)) f dx' dy' \\ &+ \iint_{\Omega} \psi((x', y') - (x, y)) z dx' dy'. \end{aligned}$$

And so we have: $E[u(x, y)] = u_0$ where $\Delta u_0 = f$. And $Var[u(x, y)] = \sigma_f^2 L_f^2 K(x, y)$ where

$$K(x, y) = \iint_{\Omega} (\psi((x', y') - (x, y)))^2 dx' dy',$$

□

B. Proof of theorem 3.3

Proof. To find the eigenvalues and eigenfunctions of the Laplacian operator, we are essentially solving the following PDE

$$\Delta u = \lambda u$$

We use separation of parts to split u as $u(x, y) = X(x)Y(y)$.

$$\Delta u = X''(x)Y(y) + X(x)Y''(y) = \lambda X(x)Y(y)$$

$$\frac{X''}{X} + \frac{Y''}{Y} = \lambda$$

Here, we solve two eigenvalue problems, $X''/X = \lambda_1$ and $Y''/Y = \lambda_2$, which means $\lambda = \lambda_1 + \lambda_2$. Consider the boundary value problem for X .

$$\frac{X''}{X} = \lambda_1, X(0) = X(1) = 0$$

The eigenvalue and eigenfunction pairs are trivially of the form

$$X_n(x) = \sin(n\pi x), \lambda_{1,n} = -(n\pi)^2$$

By symmetry, we have

$$Y_m(y) = \sin(m\pi y), \lambda_{2,m} = -(m\pi)^2$$

Thus, the eigenvalue and eigenfunction pairs of the Laplacian operator are of the form

$$\begin{aligned} u_{n,m}(x, y) &= \sin(n\pi x) \sin(m\pi y) \\ \lambda_{n,m} &= -(n\pi)^2 - (m\pi)^2 \end{aligned}$$

□