

# Microbial Communities Collectively Recycle Cadavers Over One Year of Human Decomposition

Allison R. Mason<sup>1</sup>, Lois S. Taylor<sup>2</sup>, Naomi E. Gilbert<sup>1</sup>,  
Steven W. Wilhelm<sup>1</sup>, Jennifer M. DeBruyn<sup>1,2\*</sup>

<sup>1</sup>Department of Microbiology, University of Tennessee-Knoxville, 1311  
Cumberland Avenue, Knoxville, 37996.

<sup>2</sup>Department of Biosystems Engineering and Soil Science, University of  
Tennessee-Knoxville, 2506 E.J. Chapman Drive, Knoxville, 37996.

\*Corresponding author(s). E-mail(s): [jdebruyn@utk.edu](mailto:jdebruyn@utk.edu);

## Abstract

During terrestrial vertebrate decomposition, host and environmental microbial communities work together to drive biogeochemical cycling of carbon and nutrients. These mixed communities undergo dramatic restructuring in the resulting decomposition hotspots. To reveal the succession of both the active microbial members and the metabolic pathways they use, we generated metatranscriptomes from soil samples collected over one year from below three decomposing human bodies. Soil microbes increased expression of “heat shock” proteins in response to decomposition products changing physiochemical conditions (*i.e.*, reduced oxygen, high salt). Increased fungal lipase expression implicated fungi as key decomposers of fat tissue. Expression of nitrogen cycling genes was phased with soil oxygen concentrations: during hypoxic soil conditions, genes catalyzing N-reducing processes (*e.g.*, hydroxylamine to nitric oxide and nitrous oxide to nitrogen gas during reduced oxygen conditions) were increased, followed by increased expression of nitrification genes once oxygen diffused back into the soil. Increased expression of bile salt hydrolases implicated a microbial source for

the high concentrations of taurine typically observed during vertebrate decomposition. Collectively, microbial gene expression profiles remained altered even after one year. Together, we show how human decomposition alters soil microbial gene expression, revealing both ephemeral and lasting effects on soil microbial communities.

**Keywords:** Human Decomposition, Microbial Succession, Metatranscriptomics, Soil Microbial Ecology

## Introduction

Soil microbial communities are important drivers of ecosystem processes in terrestrial environments. Many soil microbes are decomposers that degrade complex organic matter and drive nutrient cycling in terrestrial ecosystems. Environmental disturbances can impact the presence and/or activity of soil microorganisms involved in these cycles, ultimately affecting nutrient availability and greenhouse gas emissions, such as CO<sub>2</sub> and N<sub>2</sub>O [1, 2]. Vertebrate death and subsequent carcass deposition in terrestrial ecosystems is one disturbance resulting in the deposition of large quantities of organic C and N [3–10], along with other elements (P, K, S, *etc*) [11], which collectively contribute to microbially-mediated biogeochemical cycling. In addition to this, changes in pH, temperature, and fluctuations in soil oxygen provide abiotic filtering further impacting microbial metabolic strategies [7–9, 11–13]. Vertebrate decomposition also results in mixing of host and environmental microbes: the animal’s microflora are flushed into the soil along with decomposition products where they further contribute to decomposition processes (*e.g.*, organic nitrogen mineralization) [14].

While C and N transformations have been documented during decomposition, the functional response of microbes and their roles in nutrient cycles remain unclear. The composition and structure of decomposition-impacted soil microbial communities have been investigated using sequencing of marker genes amplicons (*i.e.*, 16S rRNA, 18S

rRNA, ITS). This has allowed for the identification of changes in microbial biodiversity and taxonomic succession in response to vertebrate decomposition, revealing patterns that include increases in the anaerobic taxa *Firmicutes* and *Bacteroidetes* [15]. However, few studies have integrated soil biogeochemistry with microbial community composition, which can further help to describe microbial ecology in these decomposition systems. Taylor et al. (2024) [13] showed that fungal community shifts were linked to changes in soil dissolved oxygen, highlighting interactions between soil microbes and changes in the surrounding environment. While insightful for making potential connections between taxa and environment, these analyses do not inform which taxa are active members of the community, which functional pathways/genes are expressed, and how these pathways facilitate decomposition processes.

RNA sequencing (*i.e.*, metatranscriptomics) and metabolomics can be used to investigate microbial community functional succession during decomposition. They can identify how ecological functions, including C and N cycling, are impacted by decomposition events in terrestrial ecosystems. To date, applications of metatranscriptomics to vertebrate decomposition samples have been limited to internal host communities [16, 17]: Burcham et al. (2019) [16] revealed differential expression of amino acid and carbohydrate metabolism in the heart during mouse decomposition, while Ashe et al. (2021) [17] documented taxonomic shifts in gene expression of oral microbial communities during human decomposition.

We expected that the impacted soil microbial community, which includes a mix of host and environmental taxa, would also have altered gene expression profiles, given the release of decomposition byproducts into the soil during terrestrial decomposition. We previously assessed the decomposition-impacted soil metabolome [18], demonstrating a prevalence of amino acids and suggesting upregulation of organic nitrogen metabolic pathways. Additionally, DeBruyn et al. (2021) [18] showed the soil metabolome was

139 surprisingly still altered compared to starting conditions at the end of that 21-week  
 140  
 141 study, suggesting long-term impacts of decomposition on soil microbial functioning.  
 142  
 143 Here, we investigated soil microbial gene expression during a one-year period of human  
 144  
 145 decomposition. The overarching goal of this work was to assess the effects of vertebrate  
 146  
 147 decomposition on ecosystem function by characterizing community-level shifts in soil  
 148  
 149 microbial function. We hypothesized that: (i) gene expression would shift over time as  
 150  
 151 resources were consumed and transformed and soil chemical and physical conditions  
 152  
 153 changed due to the influx of decomposition products during soft tissue degradation  
 154  
 155 [8, 9, 18]; (ii) gene expression for enzymes involved in nitrogen cycling would be  
 156  
 157 altered, as changes in nitrogen pools have been previously described in decomposition  
 158  
 159 soils [8]; (iii) expression of genes involved in lipid metabolism would increase, as lipids  
 160  
 161 from the body entered the soil during decomposition and previous studies identified  
 162  
 163 lipolytic organisms in decomposition soils [12, 19]; (iv) microbial expression profiles  
 164  
 165 in the impacted soil would remain altered even after a year, as previous studies have  
 166  
 167 shown that community composition [20, 21] can remain altered longer than a year.  
 168  
 169 We analyzed metatranscriptomes of soil samples collected at six key timepoints over  
 170  
 171 one year of human decomposition to determine the identity of active populations and  
 172  
 173 the expression of genes and pathways relevant to the enhanced biogeochemical cycling  
 174  
 175 observed in decomposition hotspots. We compared gene expression between decom-  
 176  
 177 position timepoints and control soils that were unexposed to decomposition products  
 178  
 179 to identify functions or functional pathways of interest. We show: (i) decomposition  
 180  
 181 shifts soil microbial community gene expression, with the effects still measurable after  
 182  
 183 one year; (ii) expression of genes related to stress response are elevated in decompo-  
 184  
 185 sition soils; (iii) expression of genes encoding triacylglycerol lipase differed between  
 186  
 187 fungi (increased) and bacteria (decreased); (iv) evidence for phased nitrification and  
 188  
 189 denitrification, driven by changes in soil dissolved oxygen; (v) evidence for organic

sulfur processing (taurine) via bile salt hydrolases. This direct assessment of function expands the fundamental understanding of terrestrial vertebrate decomposition, providing insight into pathways of biogeochemical cycling within these hotspots.

# Results

## Soil Physiochemistry

Soil chemistry was altered in response to the presence of a decomposing human cadaver, with multiple parameters still impacted after one year [13]. Generally, soil pH decreased and remained low in decomposition soils of all but one individual. Soil electrical conductivity (EC) increased in response to decomposition, remaining elevated through approximately day 58 before gradually decreasing throughout the remainder of the study (Supplementary Fig. 1). Respiration (evolved CO<sub>2</sub>) increased by an order of magnitude beginning at day 12, which corresponded to a reduction in soil dissolved oxygen (DO) to 29% - 48.9%. Ammonium concentrations increased 78-fold, reaching maximum concentrations between days 12 and 58. This was followed by decreased ammonium and increased nitrate concentrations at day 86, with nitrate concentrations reaching a maximum at day 168 (Supplementary Fig. 1).

## Microbial gene expression in response to human decomposition

Gene expression profiles in decomposition-impacted soils shifted away from controls and day zero samples as decomposition progressed (Fig 1A). Expression was most different from controls on study days 58, 86, 168 (Supplementary Fig. 2), before returning toward control conditions on study day 376. After one year of decomposition, soil gene expression profiles had not returned to pre-decomposition conditions, as evidenced by their clustering away from controls and day zero samples in the MDS plot (Fig 1A).

231 Some correlations were observed between gene expression shifts and soil physiochemi-  
 232 cal data at decomposition timepoints. Canonical correspondence analysis (CCA) was  
 233 used to constrain gene expression data with soil physiochemical data (Fig 1B). CCA1  
 234 and CCA2 explained 29.2% and 18.1% of the variance in gene expression, respectively.  
 235 Transcript profiles at day 12 were associated with an increase in soil carbon to nitrogen  
 236 ratio (C:N). Gene expression profiles at days 58 to 86 were positively correlated with  
 237 increased soil temperature, EC, and evolved CO<sub>2</sub>, while study day 168 was associated  
 238 with elevated levels of soil NO<sub>3</sub>. Further, Permutational Analysis of Variance (PER-  
 239 MANOVA) revealed that internal accumulated degree hours (ADH) ( $p = 0.001$ ), soil  
 240 temperature ( $p = 0.039$ ), pH ( $p = 0.033$ ), and EC ( $p = 0.031$ ) significantly explained  
 241 some of the variation in gene expression profiles ( $p < 0.05$ ). No other soil chemical  
 242 variables were significant at  $\alpha = 0.05$  (Supplementary Table 1).  
 243  
 244 Overall, decomposition changed soil gene expression profiles over the one-year study  
 245 relative to control soils. Differential expression analysis between decomposition and  
 246 control soils identified 7,047 down-regulated and 38,425 up-regulated genes. Gene  
 247 transcripts that were associated with control soils belonged to a wide variety of clus-  
 248 ters of orthologous genes (COG) functional categories. Specifically, the top 20 genes  
 249 whose expression was higher in control soils belonged to ten unique COG categories,  
 250 including signal transduction mechanisms, transcription, and those of unknown func-  
 251 tion. In contrast, the top 20 genes whose expression was higher in decomposition soils  
 252 only fell into four COG categories (Supplementary Fig. 3 A): 1) post-translational  
 253 modification, protein turnover, and chaperones; 2) energy production and conversion;  
 254 3) cell motility; and 4) carbohydrate transport and metabolism. The most common  
 255 COG category represented in decomposition soils (80% of the top 20 genes) was post-  
 256 translational modification, protein turnover, and chaperones. Within this category,  
 257 several heat shock stress response genes were identified, including clpB, dnaK, groL2,  
 258 SSA2, HSP82, and clpB (Supplementary Table 2). Further investigation of these genes  
 259  
 260  
 261  
 262  
 263  
 264  
 265  
 266  
 267  
 268  
 269  
 270  
 271  
 272  
 273  
 274  
 275  
 276

over time shows that their expression increased, typically reaching maximum transcript levels around study days 58 and 86 (Fig 2). This corresponded to elevated soil temperatures below decomposing bodies between study days 12-80, with soil temperatures increasing to approximately 43°C [13], as well as maximum soil EC and minimum dissolved oxygen measurements between days 12 and 58 (Supplementary Fig. 1).

Taxonomy associated with top differentially expressed gene transcripts also differed between control and decomposition soils. The top 40 most significant differentially expressed gene transcripts in decomposition soils were associated with Fungi, *Actinobacteria*, and *Xanthomonadales*, while gene transcripts in controls were associated with *Acidobacteria*, *Cyanobacteria*, *Proteobacteria* ( $\alpha$ ,  $\delta$ ,  $\gamma$ ), and *Planctomycetes* (Supplementary Fig. 3 B). The greatest number of differentially expressed genes relative to control samples was observed at day 86, where we saw 145,460 and 124,883 up- and down-regulated genes, respectively.

## Temporal gene expression shows shifted in decomposer functions

Differential expression analysis between sequential study days revealed which genes were altered during decomposition time. The top ten transcripts that changed in representation (increased/decreased), determined by the lowest p-values from differential expression analysis, are reported in Supplementary Table 3 and Fig 3.

Expression of genes annotated with the COG categories cell wall/membrane/envelope biogenesis, inorganic ion transport and metabolism, and carbohydrate transport and metabolism increased proportionally from day 0 to 12. In contrast, expression of secondary metabolite biosynthesis, transport, and catabolism genes decreased during this period (Fig 3A). Transcripts from *Bacilli* and *Clostridia* increased, while transcripts from *Actinobacteria* decreased between study days zero and 12 (Fig 3).

323 Between days 12 and 58, 90% of the top 10 upregulated genes were associated with  
 324 the translation, ribosomal structure and biogenesis COG and all were taxonomically  
 325 associated with *Betaproteobacteria* (Fig 3A,B). Many of these genes were annotated  
 326 as ribosomal protein large (RPL), involved in ribosomal binding. Genes across multi-  
 327 ple COG categories with taxonomic associations to *Bacilli* and *Clostridia* decreased  
 328 between study days 12 and 58, six of which were transcripts that previously increased  
 329 between days zero and 12 (Fig 3B, Supplementary Table 3).  
 330  
 331 Multiple transcripts associated with the energy production and conversion COG, as  
 332 well as transcripts annotated with the COGs inorganic transport and metabolism,  
 333 and translation, ribosomal structure and biogenesis, increased between days 58 and  
 334 86 (Fig 3A). Two of the upregulated energy and production and conservation tran-  
 335 scripts were associated with cytochrome c oxidase subunits in *Betaproteobacteria*,  
 336 while another was annotated as *hao*, encoding the enzyme hydroxylamine dehydroge-  
 337 nase which is involved in conversion of hydroxylamine to nitrite during nitrification  
 338 (Supplementary Table 3). Further investigation into hydroxylamine dehydrogenase  
 339 showed a significant increase in *hao* transcripts at day 86 followed by subsequent  
 340 decreases at days 168 and 376 ( $F = 4.183$ ;  $p = 0.02$ ). This increase corresponded to  
 341 decreased soil ammonium levels and subsequent accumulation of nitrate (Supplemen-  
 342 tary Fig. 1). Half of the topmost downregulated genes between days 58 and 86 were  
 343 not assigned to a COG (*i.e.*, unclassified) or were of unknown function.  
 344  
 345 Differential expression comparing study days 86 with 168 and 168 with 376 identified  
 346 genes across a variety of functional categories, with many unclassified in the COG  
 347 database or with unknown function (Fig 3A). Expression of carbohydrate transport  
 348 and metabolism genes associated with *Bacilli* decreased between day 168 and 376.  
 349 *Acidobacteria* transcripts increased in decomposition-impacted soils between study  
 350 day 168 and 376, but were not associated with any single COG category (Fig 3B).  
 351  
 352  
 353  
 354  
 355  
 356  
 357  
 358  
 359  
 360  
 361  
 362  
 363  
 364  
 365  
 366  
 367  
 368



## Organic carbon metabolism

We expected to observe increased expression of lipid metabolizing genes during active and advanced decomposition as microbes degraded lipids deposited in the soil [19]. Therefore, we investigated changes in triacylglycerol lipase (enzyme commission number: 3.1.1.3) gene transcription in our soils. Generally, lipase transcripts decreased as decomposition progressed (HLM  $F = 6.564$ ,  $p < 0.001$ ), however we also observed a significant interaction between study day and taxonomic annotation ( $F = 8.786$ ;  $p < 0.001$ ). Specifically, lipase gene transcripts annotated as bacteria decreased with decomposition time ( $F = 10.392$ ;  $p = 0.001$ ), while fungal lipase transcripts increased, reaching a maximum at study day 58 ( $F = 4.509$ ;  $p = 0.015$ ) (Fig 4).

## Nitrogen- and sulfur compound transformations

Expression of nitrogen cycling genes was impacted in response to human decomposition. Due to the detection of hydroxylamine oxidoreductase (*hao*) transcripts in our differential expression analysis, and our hypotheses predicting changes to nitrogen transformation processes, the expression of genes encoding common enzymes involved in nitrogen cycling (*nifH*, *nirB*, *nirK*, *norB*, *nosZ*, *nrfA*, *nxrA*, and *amoA*) were assessed using their enzyme commission numbers (Fig 5A,B). *nifH*, encoding a subunit of nitrogenase which is involved in nitrogen fixation, displayed little to no changes in gene expression between control and decomposition soils. Transcripts for two genes encoding enzymes contributing to the last two steps of denitrification, *norB* (nitric oxide reductase) and *nosZ* (nitrous oxide reductase), increased between study days 12 and 86, and decreased at study day 168 before increasing again at day 376. In contrast, expression of genes encoding nitrate reductase, *narG*, and NO-forming nitrite reductase, *nirK*, remained low until day 376 when transcripts for both genes increased. As noted above, expression of *hao*, encoding hydroxylamine dehydrogenase, increased at study day 86 before decreasing at remaining timepoints (Fig 3A, Fig 5B). Expression

415 of *amoA*, encoding a subunit of ammonia monooxygenase, and *nxrA*, encoding a sub-  
 416 unit of nitrite oxidoreductase, which are involved in nitrification, changed in response  
 417 to decomposition. *amoA* transcripts initially decreased at day 12, remaining reduced  
 418 until study day 376. Similarly, abundance of genes that encode for enzymes involved  
 419 in dissimilatory nitrate reduction, *nirB*, and *nrfA*, was low for the first 168 days, with  
 420 *nrfA* expression increasing at day 376 (Fig 5B).  
 421  
 422 Expression of genes involved in metabolism of nitrogen and sulfur-containing com-  
 423 pounds were also impacted by human decomposition. Specifically, four of the top  
 424 ten genes whose expression decreased at day 12 were related to taurine metabolism,  
 425 with their annotations associated with *tauD*, encoding taurine dioxygenase. (Supple-  
 426 mentary Table 3). Further investigation into *tauD* showed that mean expression of  
 427 these genes decreased steadily over one year, beginning at day 12 (Fig 6B); however,  
 428 *tauD* expression in response to human decomposition was variable across taxonomic  
 429 associations. Most *tauD* transcripts were associated with *Gammaproteobacteria*, *Acti-*  
 430 *nobacteria*, *Betaproteobacteria*, *Alphaproteobacteria*, and fungi. While a majority of  
 431 the *tauD* gene queries displayed reduced expression over time, expression of fungal-  
 432 associated and a few *Betaproteobacteria*-associated *tauD* genes increased at day  
 433 58 (Supplementary Fig. 4). Sources of taurine in the human body include taurine  
 434 absorbed from the diet and taurine produced from anaerobic microbial deconjugation  
 435 of bile salts via bile salt hydrolase (BSH) enzymes [22]. Therefore, we examined expres-  
 436 sion for genes encoding BSH enzymes in decomposition soils. Expression of these genes  
 437 was elevated at days 12, 58, and 86 before converging toward pre-decomposition levels  
 438 at days 168 and 376 (Fig 6A). Hierarchical liner mixed effects (HLM) models showed  
 439 that both *tauD* (HLM  $F = 7.356$ ,  $p = 0.002$ ) and BSH ( $F = 13.768$ ,  $p < 0.001$ ) gene  
 440 expression was significantly different over time (Fig 6A,B).  
 441  
 442  
 443  
 444  
 445  
 446  
 447  
 448  
 449  
 450  
 451  
 452  
 453  
 454  
 455  
 456  
 457  
 458  
 459  
 460

## Discussion

The goal of this study was to assess microbial gene expression in soils responding to human decomposition. Metatranscriptomics were applied to soil samples collected over one year from below three decomposing human bodies. From this, we found that microbial gene expression reproducibly shifted over time. Additionally, we showed that gene expression profiles had not recovered to pre-decomposition conditions after one year. Comparison of control and decomposition expression profiles revealed that heat-shock proteins were elevated in response to decomposition. We also described expression patterns between decomposition timepoints, noting changes in functional gene categories at certain timepoints, in particular with respect to lipid, nitrogen and sulfur metabolism.

### Decomposition impacted soil community gene expression, for at least one year

Gene expression profiles remained altered after one year of decomposition. It is unclear if soil microbial communities, in terms of gene expression profiles, have reached a new steady state as a result of decomposition, or if they would eventually return to pre-decomposition conditions. The soil pH, EC,  $\text{NH}_4^+$ ,  $\text{NO}_3^-$ , and total nitrogen (TN) exhibited differences (although not statistically significant) in these soils following a year of decomposition, however bacterial and fungal community structures, as assessed by rRNA amplicon libraries, were still altered [13]. This indicates that decomposition can continue to structure microbial communities and impact their function for extended periods of time. While nutrient pools and communities both demonstrate less rapid change at later time points in the study, there is no evidence suggesting an arrival at a steady-state post-disturbance microbial community within our study. In some studies, human decomposition can result in elevated carbon and nutrients (organic nitrogen, ammonium, nitrate, and phosphate) for longer than a year [3], suggesting

decomposition events have long-lasting effects on the local ecosystem. Together, this has implications for terrestrial ecosystem processing (*e.g.*, nutrient cycling, emission of greenhouse gasses, etc.), as we show that decomposition alters functional metabolism pathways within soil microbial communities. It is clear that extended sample collections beyond a single year are needed to address how long microbial communities are effected, and whether there is a return to the original state or some new altered community condition.

Bacteria, fungi, and archaea were all represented by expressed genes throughout decomposition, suggesting that members of all three domains have the potential to contribute to decomposition processes and nutrient cycling. While a majority of annotated transcripts were identified as bacterial, fungal transcripts were the second most abundant group. Fungal transcripts made up almost half (*e.g.*, seven of the top fifteen) of the significantly differentially expressed genes associated with decomposition-impacted soils. Additionally, with respect to expression shifts between decomposition timepoints, fungal transcripts were among the topmost upregulated genes at study day 86. This is not surprising as fungi are key decomposers, involved in the degradation of organic matter in terrestrial ecosystems [23]. It was interesting to see an increase in certain fungal transcripts, such as lipase, at study days 58 and 86 when soil oxygen began to recover. We would expect lipids to enter the soil as tissues are broken down during decomposition, so we were surprised to see bacterial lipase genes decrease during decomposition. This suggests that microbial activity in decomposition soils may be constrained by the changing chemical environment, potentially altered oxygen levels in the case of bacterial lipase gene expression. Prior work with these same soils showed that soil oxygen concentration was a key driver of changes in both bacterial and fungal community composition [13].

<b>Increased stress responses during decomposition</b>	553
	554
Soil microbial communities expressed stress response genes in response to human	555
decomposition. Differential expression analysis identified increased expression of mul-	556
multiple heat shock proteins associated with the taxa <i>Xanthomonadales</i> , <i>Actinobacteria</i> ,	557
and fungi. Upon further investigation, expression of these genes increased through	558
day 58 and remained high for the remainder of the year. Soil temperature was ele-	559
vated relative to controls between study days 8 and 80, with maximum temperatures	560
>40°C, while soil electrical conductivity increased up to 663 µS/cm (16X higher than	561
background) through day 58 before slowly decreasing through the remainder of the	562
study. Soil electrical conductivity correlates with ionic strength and can be an indica-	563
tor of increased salinity [24]. With regard to vertebrate decomposition, early elevated	564
conductivity in impacted soils is attributable to sodium (Na), potassium (K), and	565
ammonium (NH <sub>4</sub> ) [8–11, 13]. As a result, we would expect these microbes to be expe-	566
riencing both heat and osmotic stress during this period. Prior work has observed	567
increased heat shock gene expression during salt stress in paddy soils [25] and the	568
presence of both heat and osmotic stress genes in desert soils along a salt gradient [26],	569
suggesting saline conditions can alter the expression of heat and/or osmotic stress	570
genes. In our study we observed the stress response within soil microbial communities	571
was stimulated during human decomposition. At this time, however, it is unclear if	572
expression of these genes is in response to heat stress alone, or in combination with	573
osmotic stress.	574
	575
	576
	577
	578
	579
<b>Increased expression of fungal lipase genes during</b>	580
<b>decomposition</b>	581
	582
Human fat tissue contains lipids that are broken down during decomposition. There-	583
fore, we assessed expression of triacylglycerol lipase genes in decomposition soils. Our	584
results show that expression of triacylglycerol lipase genes was altered in response	585
	586
	587
	588
	589
	590
	591
	592
	593
	594
	595
	596
	597
	598

599 to decomposition, and these shifts differed between bacterial and fungal transcripts.  
600 Specifically, bacterial triacylglycerol lipase transcripts decreased in response to decom-  
601 position, while fungal triacylglycerol lipase transcripts increased. Further, expression  
602 of these genes corresponded to changes in relative abundance of the fungal classes  
603 *Saccharomycetes*, *Sordariomycetes*, and *Eurotiomycetes* [13]. These fungi have been  
604 previously associated with decomposition soils [27, 28] and are known to contain tri-  
605 acylglycerol lipase genes in their genomes [29, 30], suggesting that they play a role in  
606 lipid degradation in decomposition soils.  
607

612  
613 Our observation of an overall decrease in triacylglycerol lipase transcripts contrasts  
614 with previous work by Howard et al. (2010) [19], who observed increased gene copy  
615 number of Group 1 lipase genes via qPCR during swine decomposition. Fatty acid  
616 composition differs in human compared to pig tissue [31], potentially altering the  
617 lipids profile available for microbes, leading to differences in decomposition products  
618 within the soil [18]. These products can then directly or indirectly alter commu-  
619 nity composition and/or activity of functional proteins via substrate availability or  
620 the chemical environment. Further, decomposition of humans and pigs resulted in  
621 increased pH in soils below pigs, and decreased pH below humans [18]. Altered pH  
622 and soil chemistry could result in a different functional potential and/or gene expres-  
623 sion in decomposition-impacted soils. Many triacylglycerol lipases have a pH optimum  
624 that is neutral to basic [32–34], so cells may be decreasing expression under acidic  
625 conditions in human decomposition soils. Availability of lipid species and changes to  
626 pH may select for taxa that favor these substrates/pH conditions; for example, Mason  
627 et al. (2022) [12] suggested the abundance of the fungal taxa *Saccharomycetes* was  
628 related to antemortem BMI due to relative proportions of fat and muscle tissue.  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644

## Evidence for phased denitrification and nitrification

The human body is a concentrated source of nitrogen that is released into the surrounding soil during decomposition. Expression of common marker genes for nitrogen cycling was altered in decomposition soil and suggested nitrogen transformations during human decomposition are driven by soil oxygen concentrations with hydroxylamine as an important intermediate. We observed low or reduced expression of the nitrification genes *nrrA* and *amoA* between days 12 and 86, during a period when oxygen was reduced to 39% - 85%. This was concomitant with an accumulation of ammonium, which reached a maximum on day 12, and low nitrate conditions indicating that nitrification was inhibited. This period of reduced soil oxygen constraining nitrification was also described in a decomposition experiment with beaver carcasses Keenan et al. (2018) [8].

We observed increased gene expression for the enzyme hydroxylamine dehydrogenase (HAO) at day 86 while oxygen was reduced (~85%). This corresponded to simultaneous increases in expression of genes encoding nitric oxide reductase (*norB*) and nitrous oxide reductase (*nosZ*). Traditionally HAO has been thought to process hydroxylamine to nitrite during nitrification, while NorB and NosZ are enzymes involved in the last two steps of denitrification converting nitric oxide (NO) to dinitrogen gas (N<sub>2</sub>). However, recent work suggested hydroxylamine can be converted to nitric oxide (NO), and can interact with multiple phases of the nitrogen cycle [35]. Even though *amoA* expression was shown to decrease during reduced oxygen conditions, *amoA* transcripts were still present and likely able to convert ammonium to hydroxylamine as soil oxygen was not completely depleted during decomposition. Additionally, a previous study reported that the growth of the ammonia oxidizing bacteria *Nitrosomonas europaea* under anoxic conditions lead to accumulation of hydroxylamine in a chemostat bioreactor [36], suggesting anaerobic ammonium oxidation (anammox) may also be occurring in decomposition soils. However, we did not observe increases in *nirK*

expression, which might suggest conversion of nitrite to NO for use in the anammox pathway. NO produced via HAO activity may be used for anammox in these soils; however, the role of hydroxylamine as an intermediate in anammox is still debated [35]. Therefore, our current hypothesis is that hydroxylamine accumulates under anaerobic conditions during decomposition, which can then be converted to NO by HAO. This NO would then be present for anaerobic denitrifying bacteria to convert to nitrous oxide (N<sub>2</sub>O) by NorB and finally to N<sub>2</sub> by NosZ. Keenan et al. (2018) [8] noted a brief increase in N<sub>2</sub>O emissions, which suggests denitrification was occurring during this phase of reduced soil oxygen concentrations.

As soils fully reoxygenated by day 168, we observed increased expression of genes encoding enzymes involved in aerobic nitrification, *amoA* and *nxrR*. Nitrification is an oxygen-dependent process which would convert accumulated ammonium to nitrate; the increase in nitrate concentrations may then serve as a substrate for denitrification. We observed increased expression of marker genes encoding all four enzymes in the complete dissimilatory denitrification pathway (*narG*, *nirK*, *norB*, and *nosZ*) at day 376. Increased expression of nitrification and denitrification marker genes is consistent with the accumulation of nitrite, nitrate, and N<sub>2</sub>O after oxygen is reintroduced to soils described in Keenan et al. (2018) [3, 8]. Together, gene expression patterns in our study provide further insight into nitrogen transformations in during vertebrate decomposition, suggesting an important role of hydroxylamine.

## Increased expression of bile salt hydrolases

Sulfur is present in various organic molecules, including taurine, a sulfur- and nitrogen-rich compound involved in bile acid formation [22]. Taurine in the human body can be absorbed from the diet or synthesized in the liver [37]. However, taurine is also produced as a byproduct of the deconjugation of bile salts via bile salt hydrolases (BSHs) present in the anaerobic gut taxa *Lactobacillus* and *Clostridium* [22]. We



observed increased expression of genes encoding BSH enzymes between days 12 and 86. Given that increased expression of BSH genes corresponded to the beginning of active decomposition, when decomposition products were observed to enter the soil, and the period of reduced dissolved oxygen in our study, it is likely that taurine accumulation is the result of BSH enzyme activity by anaerobic microorganisms. While we did not measure taurine concentrations in the present study, our results correspond to previous decomposition studies that report accumulation of taurine in various organs and body regions [38–40] and soils [18, 41] during decomposition via metabolomics, and increased relative abundance of *Clostridium* and *Lactobacillus* within the body [42–44] and in decomposition soils [20] via DNA sequencing methods, including in these soils [13].

Taurine can be metabolised through desulfurization via the  $\alpha$ -ketoglutarate-dependent enzyme taurine dioxygenase (TauD). Specifically, this enzyme, encoded by the gene *tauD*, converts 2-oxoglutarate and taurine to produce aminoacetaldehyde, succinate, sulfite, and CO<sub>2</sub> [45]. Succinate and sulfite from this reaction can then be used for the citric acid cycle and sulfur metabolism, respectively. Given increased BSH expression in our study and reported taurine accumulation in others, we would expect taurine to be present for microbial metabolism by TauD. However, we observed a general decrease in *tauD* expression between days 12 through 376. This trend was driven by reduced expression of *tauD* transcripts associated with *Proteobacteria*, *Gammaproteobacteria*, and *Actinobacteria* whose relative abundance have been shown to remain consistent or increase during human decomposition [20], suggesting that *tauD* expression is downregulated under decomposition conditions. However, we noted that expression of *tauD* genes associated with fungi and a few *Betaproteobacteria* displayed increased representation at day 58, corresponding to increased expression of bile salt hydrolases (BSH) between days 12 and 86. The reduction in *tauD* expression may be due to increased sulfur availability. We did not measure sulfur species

783 in this experiment; however, others have observed increased sulfur concentrations in  
784 decomposition-impacted soils [3, 7, 11]. Thus, sulfur scavenging pathways such as tau-  
785 rine desulfurization by TauD [46], whose genes are expressed under sulfur-limiting  
786 conditions, likely display reduced expression under sulfur replete conditions. Addition-  
787 ally, taurine may be processed through other pathways. For example, taurine can be  
788 deaminated by taurine dehydrogenase to produce sulfite and acetyl-CoA for carbon  
789 metabolism [45, 47]. Overall, our results suggest that human decomposition has poten-  
790 tial impacts on soil sulfur biogeochemistry through deposition of inorganic (sulfate)  
791 and organic (sulfur-containing amino acids) sulfur compounds.  
792  
793  
794  
795  
796  
797  
798

## 799 Conclusion

800  
801 This study investigated soil microbial gene expression during human decomposition.  
802 Metatranscriptomic analysis of soils from three human individuals shows that decom-  
803 position impacted microbial community gene expression profiles, exhibiting functional  
804 shifts over time for over one year. This included altered expression of genes involved in  
805 lipid, N and S metabolism as microbes processed the nutrient-rich tissues of the human  
806 body. Additionally, we noted that functionality within decomposition-impacted soils  
807 was still affected after one year and had not returned to starting or background condi-  
808 tions. Together, these results show that vertebrate decomposition has lasting impacts  
809 on local soil ecosystems, including soil microbial communities. These results have  
810 important implications for understanding biogeochemical changes due to vertebrate  
811 mortality events in terrestrial ecosystems.  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828

## Materials and Methods

### Study design

In February 2018, three deceased male human subjects (hereafter, “donors”) were placed supine on the soil surface at the University of Tennessee Anthropology Research Facility (ARF) and allowed to decompose. Located in Knoxville, TN (35° 56’ 28” N, 83° 56’ 25” W) the ARF is a roughly 2-acre outdoor facility dedicated to studying human decomposition [48]. The soils at the ARF are comprised of the Loyston-Talbott-Rock outcrop (LtD) and Coghill-Corryton (CcD) complexes. LtD soils are a silty clay loam and channery clay overlaying lithic bedrock, while CcD soils are comprised of clay from weathered quartz limestone [13, 48]. A site that had not been previously exposed to decomposition was used for this study.

The decomposition field experiment is fully described in Taylor et al. (2024) [13]. Briefly, experiments were conducted in a block design, where each block consisted of one decomposition site and one control site [13]. In total three blocks, *i.e.*, three donors paired with three respective control sites, were included in the study. Each control site was chosen in a manner to ensure their location was uphill and roughly 2 m away from decomposition sites [13]. Donor internal temperatures were recorded by probes located in the abdomen, while ambient air temperatures were monitored via sensors located roughly 50 cm above the soil surface. Soil temperature and salinity were measured with sensors placed directly underneath each individual (Decagon Devices, GS3) [13]. Donor ages ranged from 65 to 86 and were within 1 kg of each other with regard to weight (90.7 to 91.6 kg); donor BMI varied between 27.7 to 29.6 [13].

### Sampling and physiochemistry

Decomposition of all subjects was observed for one year. During the one-year study period, soils were sampled at 20 timepoints chosen to correspond with morphological

875 stages of decomposition as described by [49]. Once advanced decay was reached, soils  
876 were collected at intervals of 350 accumulated degree days (ADD), calculated using  
877 ambient air temperatures, up to one year. All soil cores were taken using a 1.9 cm  
878 (3/4 inch) diameter soil auger to a depth of 16 cm. Soils were divided into two depth  
880 fractions: 0-1 cm (interface) and 1-16 cm (core) for the analyses reported in Taylor et  
881 al. (2024) [13]; the entire 0 to 16 cm core was used for this current study. Decomposi-  
882 tion soils were taken from directly beneath the cadavers, taking care to not re-sample  
883 the same location more than once. At the time of sampling, soil dissolved oxygen was  
884 measured in triplicate using an Orion Star™ A329 pH/ISE/Conductivity/Dissolved  
885 Oxygen portable multiparameter meter (ThermoFisher) [13].

891  
892 A subset of 6 study timepoints were chosen for metatranscriptomics analysis. Study  
893 days 0, 12, 58, 86, 168, and 376 were chosen as they represented distinct morphologi-  
894 cal and soil biogeochemical stages during decomposition. Study day 0 was chosen as  
895 a baseline sample prior to cadaver placement. Study day 12 was the start of active  
896 decomposition and corresponded to maximum soil ammonium concentrations and  
897 minimum soil oxygen (approximately 39%). Study day 58 was chosen as this sample  
898 represented the pH minimum, and respiration and soil temperature were at a maxi-  
899 mum [13]. Additionally, ammonium concentrations began to decrease around day 58.  
900 Study day 86 was when soil oxygen started to recover and nitrate levels began to  
901 increase. Study day 168 was chosen as nitrate was at its maximum and soil dissolved  
902 oxygen had returned to 99%. Finally, day 376 was chosen to represent the end of the  
903 study, 1 year since cadaver placement. Each study day was represented by four soil  
904 samples for RNA extraction: one pooled control sample which was a mix of the three  
905 control locations, plus one sample from each of the three donors, yielding a total of  
906 24 samples for this study.

Soil samples were transported back to the University of Tennessee (Knoxville, TN) and processed within 24 hours of collection. Soils were homogenized by hand to remove insect larvae, roots, rocks, and other debris ( $> 2$  mm). A subset of soils were used to measure pH, electrical conductivity (EC), and evolved  $\text{CO}_2$  as described in Taylor (2024). Soil nitrogen species ( $\text{NH}_4^+$ ,  $\text{NO}_3^-$ ) and total carbon (TC) and nitrogen (TN) were measured in all soil samples as described in [13]. Reported values for soil physiochemistry represent the full 16 cm core; estimated by summing interface and core values reported by Taylor et. al, (2024) [13] in 1:16 and 15:16 ratios, respectively. Control reported here are means of the three experimental controls that were unimpacted by decomposition.

Roughly 10 g of soil was reserved for nucleic acid extraction, placed in a 4 oz. Whirl-Pak™ bag (Nasco), and flash frozen in liquid nitrogen. All samples were stored at  $-80^\circ\text{C}$  until further analysis. Bacterial and fungal community composition was assessed via amplicon sequencing of the 16S rRNA gene and ITS2 region as described in Taylor et al. (2024).

## RNA Extraction and Sequencing

RNA was extracted from 2 g of soil using Qiagen's RNeasy® PowerSoil® Total RNA kit. Manufacturer's instructions were followed with a few modifications. Soils became saline during decomposition; therefore, we followed the manufacturer's suggestion and incubated all extracts at  $-20^\circ\text{C}$  following addition of solution SR4 (step 9) to decrease salt precipitation. All RNA samples were resuspended in 40  $\mu\text{l}$  of Solution SR7. RNA concentrations were assessed fluorometrically using the Qubit® RNA HS assay (catalog no. Q32852) with 1  $\mu\text{l}$  of RNA. DNA contamination was removed by DNase treating RNA extracts twice using Qiagen's DNase Max® kit in 50  $\mu\text{l}$  reactions. RNA concentrations were remeasured after DNase treatment. PCR with V4 16S rRNA gene primers [50, 51] was conducted using RNA extracts as the template to confirm removal

of all DNA prior to sequencing. RNA aliquots were shipped to HudsonAlpha Discovery (Huntsville, AL) for library preparation and RNA sequencing. Dual-indexed libraries were prepared using the Illumina® Stranded Total RNA prep with ribosomal RNA depletion via ligation with Ribo-Zero Plus. Libraries were then pooled and sequenced on Illumina’s NovaSeq 6000 v4 platform, resulting in demultiplexed fastq files for each sample.

## Bioinformatics

Illumina sequencing of the 24 libraries yielded a total of 5,073,476,730 reads, or 2,536,738,365 paired reads, with a mean of 105,697,432 paired reads per sample. Read quality control (QC) was conducted in KBase [52] using Trimmomatic [53]. Paired fastq files were imported to KBase through Globus. Poor quality reads were removed (4.7% of all reads), and adapters trimmed via Trimmomatic (v0.36) using default settings and the TruSeq3-PE-2 adapter file, resulting in 4,834,123,062 total reads. After QC check with FastQC, trimmed libraries were exported as fastq files from KBase through Globus. Remaining ribosomal RNA was filtered using bbmap (maxindel = 20, minid = 0.93) from the Joint Genome Institute’s (JGI) bbtools suite [54]. Filtering of ribosomal RNA further removed 7.3% of reads, leaving 4,479,804,360 reads for assembly. Following this step, all non-ribosomal reads from all 24 samples were merged into one file. Reads were then co-assembled into contigs using the de novo assembler MEGAHIT (v1.2.9) [55] (–12 –k-min 23, –k-max 123, –k-step 10).

Gene identification and annotation from co-assembled contigs was performed using Prodigal [56] and eggNOG mapper [57], respectively. Briefly, the DNA fasta containing all contigs was submitted to Prodigal (v2.6.3) for protein coding gene predication for a meta-sample (–p meta –f gff). After co-assembly, a total of 6,257,674 gene calls were identified by Prodigal. Next, predicated genes were functionally and taxonomically annotated using eggNOG mapper (v2.1.6) using basic settings to perform a

diamond blastp search [58]. From this, 1,048,573 proteins were annotated by eggNOG-mapper (16.7%). Most of the annotated proteins were taxonomically annotated as bacteria (91.3%), followed by eukaryotes (7.6 %), and archaea (0.81 %). Of the 7.6% of eukaryotic proteins, 64.4% (4.9% of all proteins) were annotated as fungi. For this study, genes of interest included all bacterial, archaeal, and fungal proteins, therefore all non-fungal eukaryotic proteins (32,004) were removed prior to downstream analysis. Transcript counts for all genes of interest were obtained by mapping reads from each respective sample to genes of interest obtained from co-assembly using QIAGEN CLC Genomics Workbench 20.0 (<https://digitalinsights.qiagen.com/>). The percent of reads mapped to genes of interest ranged from 21% to 38% between samples, with an average of 31% reads mapped. Gene counts were then combined in a single file and used for downstream analyses in R.

## Differential Expression

Transcript counts from all samples were combined in a single workable data file and imported into R for differential expression analysis using the R packages edgeR [59] and limma [60] following a modified pipeline by Phipson et al. (2020) [61]. The transcript count table was imported into R and converted to a DGElist object. Genes without sufficient counts for statistical analysis were removed to increase power using the edgeR function filterByExpr(), using study day as the comparison group.

Raw counts were then log2 normalized and gene expression profiles compared via multidimensional scaling (MDS) and hierarchical clustering. Multidimensional scaling (MDS) was conducted using plotMDS() from the limma package to assess differences between samples. MDS values were extracted from the MDS object, and the first two dimensions plotted using ggplot2 [62]. We also assessed the relationship between gene expression profiles and changes in the soil environment using canonical correspondence analysis (CCA). Environmental variables of interest included decomposition time in

1059 accumulated degree hours (ADH) based on ambient temperatures, ADH based on  
 1060 internal gut temperatures, ADH based on soil temperatures, gravimetric moisture,  
 1061 pH, electrical conductivity (EC), dissolved oxygen (DO), CO<sub>2</sub> (μmol gdw<sup>-1</sup>), NH<sub>4</sub> (mg  
 1062 gdw<sup>-1</sup>), NO<sub>3</sub> (mg gdw<sup>-1</sup>), N %, C %, and CN ratio. First, permutational multivariate  
 1063 analysis of variance (PERMANOVA) with `adonis()` (vegan v2.6.7) [63] was used to  
 1064 identify significant soil parameters. Then the vegan functions `cca()` and `scores()` were  
 1065 applied to run the CCA and extract scores, respectively. Scores for the first two  
 1066 dimension were plotted using `ggplot2`, with loadings extracted from the CCA biplot.  
 1067 For differential expression analysis, raw filtered reads were normalized using edgeR's  
 1068 trimmed mean of M values (TMM) normalization using the function `calcNormFac-`  
 1069 `tors()`. TMM normalized reads were then log2 transformed using limma's `voom()` and  
 1070 differential expression assessed. Empirical Bayes shrinkage was used correct to p-  
 1071 values for false discovery rates. The topmost up and down regulated genes for each  
 1072 comparison, determined by log2 fold change and adjusted p-values, were then reported.  
 1073 Expression of certain genes were assessed after performing transcripts per million  
 1074 (TPM) normalization and statistical analyses with a combination of analysis of vari-  
 1075 ance (ANOVA) and post-hoc Tukey tests. ANOVA across all timepoints were applied  
 1076 to hierarchical linear mixed effects models to account for repeated sampling within  
 1077 each donor block.

## 1093 Data availability

1094 Raw RNA sequence files from the Illumina Novaseq are available at the National Cen-  
 1095 ter for Biotechnology Information's (NCBI) Sequence Read Archive (SRA) as a part  
 1096 of [BioProject PRJNA1066312](#) under BioSample accession numbers SAMN45195141-  
 1097 SAMN45195164. Additional datasets supporting the conclusions of this article are  
 1098 available on [GitHub](#).



<b>Code availability</b>	1105
	1106
The code used for analysis and to generate figures are available on <a href="#">GitHub</a> .	1107
	1108
	1109
<b>References</b>	1110
	1111
	1112
[1] Benninger, L. A., Carter, D. O. & Forbes, S. L. The biochemical alteration of soil	1113
beneath a decomposing carcass. <i>Forensic Science International</i> <b>180</b> , 70–5 (2008).	1114
	1115
	1116
[2] Towne, E. G. Prairie vegetation and soil nutrient responses to ungulate carcasses.	1117
<i>Oecologia</i> <b>122</b> , 232–239 (2000). URL <a href="https://doi.org/10.1007/PL00008851">https://doi.org/10.1007/PL00008851</a> .	1118
	1119
	1120
[3] DeBruyn, J. M., Keenan, S. W. & Taylor, L. S. From carrion to soil: microbial	1121
recycling of animal carcasses. <i>Trends in Microbiology</i> (2024). URL <a href="https://doi.org/10.1016/j.tim.2024.09.003">https://doi.org/10.1016/j.tim.2024.09.003</a> . Publisher: Elsevier.	1122
	1123
	1124
	1125
	1126
[4] Parmenter, R. R. & MacMahon, J. A. Carrion decomposition and nutrient cycling	1127
in a semiarid shrub–steppe ecosystem. <i>Ecological Monographs</i> <b>79</b> , 637–661 (2009).	1128
	1129
	1130
[5] Macdonald, B. C. T. <i>et al.</i> Carrion decomposition causes large and lasting effects	1131
on soil amino acid and peptide flux. <i>Soil Biology and Biochemistry</i> <b>69</b> , 132–140	1132
(2014).	1133
	1134
	1135
	1136
[6] Bump, J. K. <i>et al.</i> Ungulate carcasses perforate ecological filters and create	1137
biogeochemical hotspots in forest herbaceous layers allowing trees a competitive	1138
advantage. <i>Ecosystems</i> <b>12</b> , 996–1007 (2009).	1139
	1140
	1141
	1142
[7] Aitkenhead-Peterson, J. A., Owings, C. G., Alexander, M. B., Larison, N. &	1143
Bytheway, J. A. Mapping the lateral extent of human cadaver decomposition	1144
with soil chemistry. <i>Forensic Science International</i> <b>216</b> , 127–34 (2012).	1145
	1146
	1147
	1148
	1149
	1150

- 1151 [8] Keenan, S. W., Schaeffer, S. M., Jin, V. L. & DeBruyn, J. M. Mortality hotspots:  
1152 nitrogen cycling in forest soils during vertebrate decomposition. *Soil Biology and*  
1153 *Biochemistry* **121**, 165–176 (2018).  
1154  
1155  
1156  
1157 [9] Fancher, J. P. *et al.* An evaluation of soil chemistry in human cadaver decom-  
1158 position islands: Potential for estimating postmortem interval (PMI). *Forensic*  
1159 *Science International* **279**, 130–139 (2017).  
1160  
1161  
1162  
1163 [10] Quaggiotto, M.-M., Evans, M. J., Higgins, A., Strong, C. & Barton, P. S.  
1164 Dynamic soil nutrient and moisture changes under decomposing vertebrate  
1165 carcasses. *Biogeochemistry* **146**, 71–82 (2019).  
1166  
1167  
1168 [11] Taylor, L. S. *et al.* Soil elemental changes during human decomposition.  
1169 *PLoS ONE* **18**, 1–24 (2023). URL <https://doi.org/10.1371/journal.pone.0287094>.  
1170  
1171 Publisher: Public Library of Science.  
1172  
1173  
1174 [12] Mason, A. R. *et al.* Body mass index (BMI) impacts soil chemical and microbial  
1175 response to human decomposition. *mSphere* e0032522 (2022).  
1176  
1177  
1178 [13] Taylor, L. S. *et al.* Transient hypoxia drives soil microbial community dynamics  
1179 and biogeochemistry during human decomposition. *FEMS Microbiology Ecology*  
1180 **100**, fae119 (2024). URL <https://doi.org/10.1093/femsec/fae119>.  
1181  
1182  
1183  
1184 [14] Keenan, S. W., Emmons, A. L. & DeBruyn, J. M. Microbial community coa-  
1185 lescence and nitrogen cycling in simulated mortality decomposition hotspots.  
1186 *Ecological Processes* **12**, 45 (2023). URL [https://doi.org/10.1186/s13717-023-](https://doi.org/10.1186/s13717-023-00451-y)  
1187 [00451-y](https://doi.org/10.1186/s13717-023-00451-y).  
1188  
1189  
1190  
1191  
1192 [15] Mason, A. R., Taylor, L. S. & DeBruyn, J. M. Microbial ecology of vertebrate  
1193 decomposition in terrestrial ecosystems. *FEMS Microbiology Ecology* **99**, fiad006  
1194 (2023). URL <https://doi.org/10.1093/femsec/fiad006>.  
1195  
1196

- [16] Burcham, Z. M. *et al.* Total RNA analysis of bacterial community structural and functional shifts throughout vertebrate decomposition. *Journal of Forensic Sciences* **64**, 1707–1719 (2019).
- [17] Ashe, E. C., Comeau, A. M., Zejdlik, K. & O’Connell, S. P. Characterization of bacterial community dynamics of the human mouth throughout decomposition via metagenomic, metatranscriptomic, and culturing techniques. *Frontiers in Microbiology* **12**, 689493 (2021).
- [18] DeBruyn, J. M. *et al.* Comparative decomposition of humans and pigs: soil biogeochemistry, microbial activity and metabolomic profiles. *Frontiers in Microbiology* **11**, 608856 (2021).
- [19] Howard, G. T., Duos, B. & Watson-Horzelski, E. J. Characterization of the soil microbial community associated with the decomposition of a swine carcass. *International Biodeterioration & Biodegradation* **64**, 300–304 (2010).
- [20] Cobaugh, K. L., Schaeffer, S. M. & DeBruyn, J. M. Functional and structural succession of soil microbial communities below decomposing human cadavers. *Plos One* **10**, e0130201 (2015).
- [21] Singh, B. *et al.* Temporal and spatial impact of human cadaver decomposition on soil bacterial and arthropod community structure and function. *Frontiers in Microbiology* **8**, 2616 (2018).
- [22] Urdaneta, V. & Casadesús, J. Interactions between Bacteria and Bile Salts in the Gastrointestinal and Hepatobiliary Tracts. *Frontiers in Medicine* **4** (2017).
- [23] van der Wal, A., Geydan, T. D., Kuyper, T. W. & de Boer, W. A thready affair: linking fungal diversity and community dynamics to terrestrial decomposition processes. *FEMS Microbiology Reviews* **37**, 477–494 (2013).

1243 [24] Essington, M. E. *Soil and water chemistry: an integrative approach* (CRC press,  
1244 2015).  
1245  
1246

1247 [25] Peng, J., Wegner, C.-E. & Liesack, W. Short-term exposure of paddy soil micro-  
1248 bial communities to salt stress triggers different transcriptional responses of key  
1249 taxonomic groups. *Frontiers in Microbiology* **8** (2017).  
1250  
1251  
1252

1253 [26] Pandit, A. S. *et al.* A snapshot of microbial communities from the Kutch: one of  
1254 the largest salt deserts in the World. *Extremophiles* **19**, 973–987 (2015).  
1255  
1256

1257 [27] Metcalf, J. L. *et al.* Microbial community assembly and metabolic function during  
1258 mammalian corpse decomposition. *Science* **351**, 158–62 (2016).  
1259  
1260

1261 [28] Fu, X. *et al.* Fungal succession during mammalian cadaver decomposition and  
1262 potential forensic implications. *Scientific Reports* **9**, 12907 (2019).  
1263  
1264

1265 [29] Dujon, B. *et al.* Genome evolution in yeasts. *Nature* **430**, 35–44 (2004).  
1266  
1267

1268 [30] Haridas, S. *et al.* The genome and transcriptome of the pine saprophyte *Ophios-*  
1269 *toma piceae*, and a comparison with the bark beetle-associated pine pathogen  
1270 *Grosmannia clavigera*. *BMC Genomics* **14**, 373 (2013).  
1271  
1272

1273 [31] Notter, S. J., Stuart, B. H., Rowe, R. & Langlois, N. The initial changes of fat  
1274 deposits during the decomposition of human and pig remains. *Journal of Forensic*  
1275 *Sciences* **54**, 195–201 (2009).  
1276  
1277  
1278

1279 [32] Kok, R. G. *et al.* Characterization of the extracellular lipase, LipA, of *Acineto-*  
1280 *bacter calcoaceticus* BD413 and sequence analysis of the cloned structural gene.  
1281 *Molecular Microbiology* **15**, 803–818 (1995).  
1282  
1283  
1284

1285 [33] Hasan, F., Shah, A. A. & Hameed, A. Influence of culture conditions on lipase  
1286 production by *Bacillus* sp. FH5. *Annals of Microbiology* **56**, 247–252 (2006).  
1287  
1288

- [34] Zouaoui, B. & Bouziane, A. Production, optimization and characterization of the lipase from *Pseudomonas aeruginosa*. *Romanian biotechnological letters* **17**, 7187–7193 (2012).
- [35] Soler-Jofra, A., Pérez, J. & van Loosdrecht, M. C. M. Hydroxylamine and the nitrogen cycle: A review. *Water Research* **190**, 116723 (2021).
- [36] Yu, R., Perez-Garcia, O., Lu, H. & Chandran, K. *Nitrosomonas europaea* adaptation to anoxic-oxic cycling: Insights from transcription analysis, proteomics and metabolic network modeling. *Science of the Total Environment* **615**, 1566–1573 (2018).
- [37] Seidel, U., Huebbe, P. & Rimbach, G. Taurine: A regulator of cellular redox homeostasis and skeletal muscle function. *Molecular Nutrition & Food Research* **63**, 1800569 (2019).
- [38] Mora-Ortiz, M., Trichard, M., Oregioni, A. & Claus, S. P. Thanatometabolomics: introducing NMR-based metabolomics to identify metabolic biomarkers of the time of death. *Metabolomics* **15**, 37 (2019).
- [39] Locci, E. *et al.* A  $^1\text{H}$  NMR metabolomic approach for the estimation of the time since death using aqueous humour: an animal model. *Metabolomics* **15**, 76 (2019).
- [40] Zelentsova, E. A. *et al.* Post-mortem changes in the metabolomic compositions of rabbit blood, aqueous and vitreous humors. *Metabolomics* **12**, 172 (2016).
- [41] Hoeland Katharina, M. *Investigating the potential of postmortem metabolomics in mammalian decomposition studies in outdoor settings*. Ph.D. thesis, University of Tennessee-Knoxville, [https://trace.tennessee.edu/utk\\_graddiss/7000](https://trace.tennessee.edu/utk_graddiss/7000) (2021).

1335 [42] Javan, G. T. *et al.* Human thanatobiome succession and time since death.  
1336 *Scientific Reports* **6**, 29598 (2016).  
1337  
1338  
1339 [43] Javan, G. T., Finley, S. J., Smith, T., Miller, J. & Wilkinson, J. E. Cadaver  
1340 thanatobiome signatures: the ubiquitous nature of *Clostridium* species in  
1341 human decomposition. *Frontiers in Microbiology* **8**, 2096 (2017).  
1342  
1343  
1344  
1345 [44] DeBruyn, J. M. & Hauther, K. A. Postmortem succession of gut microbial  
1346 communities in deceased human subjects. *Peerj* **5**, e3437 (2017).  
1347  
1348  
1349 [45] Cook, A. M. & Denger, K. Metabolism of taurine in microorganisms. *Taurine* **6**  
1350 3–13 (2006).  
1351  
1352  
1353 [46] Kertesz, M. A. Riding the sulfur cycle – metabolism of sulfonates and sul-  
1354 fate esters in Gram-negative bacteria. *FEMS Microbiology Reviews* **24**, 135–175  
1355 (2000).  
1356  
1357  
1358  
1359 [47] Brüggemann, C., Denger, K., Cook, A. M. & Ruff, J. Enzymes and genes of  
1360 taurine and isethionate dissimilation in *Paracoccus denitrificans*. *Microbiology*  
1361 (*Reading, England*) **150**, 805–816 (2004).  
1362  
1363  
1364  
1365 [48] Keenan, S. W. *et al.* Spatial impacts of a multi-individual grave on microbial  
1366 and microfaunal communities and soil biogeochemistry. *PLoS One* **13**, e0208845  
1367 (2018).  
1368  
1369  
1370  
1371 [49] Payne, J. A. A summer carrion study of the baby pig *Sus Scrofa* Linnaeus.  
1372 *Ecology* **46**, 592–602 (1965).  
1373  
1374  
1375 [50] Apprill, A., McNally, S., Parsons, R. & Weber, L. Minor revision to V4 region SSU  
1376 rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton.  
1377 *Aquatic Microbial Ecology* **75**, 129–137 (2015).  
1378  
1379  
1380

[51] Parada, A. E., Needham, D. M. & Fuhrman, J. A. Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. *Environmental Microbiology* **18**, 1403–14 (2016).

[52] Arkin, A. P. *et al.* KBase: The United States Department of Energy Systems Biology Knowledgebase. *Nature Biotechnology* **36**, 566–569 (2018).

[53] Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

[54] Bushnell, B. Bbtools software packag. *e* (2014).

[55] Li, D., Liu, C.-M., Luo, R., Sadakane, K. & Lam, T.-W. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* **31**, 1674–1676 (2015).

[56] Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**, 119 (2010).

[57] Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas, J. eggNOG-mapper v2: functional annotation, orthology assignments, and domain prediction at the metagenomic scale. *Molecular Biology and Evolution* **38**, 5825–5829 (2021).

[58] Buchfink, B., Reuter, K. & Drost, H.-G. Sensitive protein alignments at tree-of-life scale using DIAMOND. *Nature Methods* **18**, 366–368 (2021).


[59] Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140 (2010).

1427 [60] Smyth, G. K. in *limma: Linear Models for Microarray Data* (eds Gentleman,  
1428 R., Carey, V. J., Huber, W., Irizarry, R. A. & Dudoit, S.) *Bioinformatics and*  
1429 *Computational Biology Solutions Using R and Bioconductor* 397–420 (Springer  
1430 New York, New York, NY, 2005).  
1431  
1432  
1433  
1434 [61] Phipson, B. *et al.* Differential expression analysis (2020). URL [https://combine-](https://combine-australia.github.io/RNAseq-R/06-rnaseq-day1.html#References)  
1435 [australia.github.io/RNAseq-R/06-rnaseq-day1.html#References](https://combine-australia.github.io/RNAseq-R/06-rnaseq-day1.html#References).  
1436  
1437  
1438 [62] Wickham, H. *ggplot2: Elegant Graphics for Data Analysis* (Springer-Verlag New  
1439 York, 2016). URL <https://ggplot2.tidyverse.org>.  
1440  
1441  
1442 [63] Oksanen, J. *et al.* *vegan: Community Ecology Package* (2024). URL [https://](https://vegandevs.github.io/vegan/)  
1443 [vegandevs.github.io/vegan/](https://vegandevs.github.io/vegan/).  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472




## Figures

**Figure 1: Microbial gene expression profiles are altered during human decomposition.** Multidimensional scaling (MDS) shows gene expression within soils changed as decomposition progressed (A). Additionally, canonical correspondence analysis (CCA) shows that environmental variables explained 47.3% of the variation in gene expression profiles (B). Variables in bold red type significantly ( $p < 0.05$ ) explained some of the variation in gene expression profiles as assessed by Permutational Analysis of Variance (PERMANOVA). In both panels soils from controls (CON) and the three donors (SP1, SP2, SP3) are denoted by symbol shape, while color represents study day. In B, soil physiochemical variable loadings are represented by arrows: Gravimetric water content (GWC), electrical conductivity (EC), pH (pH), dissolved oxygen (DO), respiration (evolved  $\text{CO}_2$   $\mu\text{mol gdw}^{-1}$ ), ammonium ( $\text{NH}_4$ ), and nitrate ( $\text{NO}_3$ ) concentrations ( $\text{mg gdw}^{-1}$ ), percent carbon (%C), percent nitrogen (%N), carbon:nitrogen ratio (C:N), ambient temperature ( $T_A$ ), and soil temperature ( $T_S$ ).




**Figure 2: Mean normalized log2 expression of heat shock proteins identified by differential expression analysis comparing decomposition and control soils.** Each panel represents a single heat shock gene, labeled with gene names, identified via Prodigal. Symbol color denotes if the sample is a control (CON, green), or one of three individuals: SP1 (orange), SP2 (purple), or SP3 (pink). Error bars are standard error of individual query genes in the top 20 transcripts associated with decomposition soils.


1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610




**Figure 3: Top twenty up- and down-regulated genes in decomposition soils comparing sequential study days (0, 12, 58, 86, 168, 376) colored by COG functional category (A) and taxonomic annotation (B). Positive values denote increased expression compared to the preceding timepoint, while negative values denote a decrease.**



1703 **Figure 4: Mean transcript abundance, in transcripts per million (TPM),**  
 1704 **of all bacterial (A) and fungal (B) triacylglycerol lipase (EC 3.1.1.3) genes**  
 1705 **over time.** Abundance of both bacterial (Anova  $p = 0.001$ ) and fungal (Anova  $p$   
 1706  $= 0.015$ ) lipase transcripts change significantly over time. Black lines (A, B) report  
 1707 mean and standard deviation of TPM from three individuals (black line), while gold  
 1708 stars denote mean TPM in control soils. Letters are the result of post-hoc Tukey  
 1709 tests between decomposition timepoints. In B, bars show the relative abundance of  
 1710 the fungal classes *Saccharomycetes*, *Sordariomycetes*, and *Eurotiomycetes*, reported in  
 1711 Taylor et al. (2024).  
 1712





**Figure 5: Mean gene expression, in transcripts per million (TPM), of commonly used marker genes for enzymes involved in nitrogen cycling over time in controls (A) and decomposition (B) soils. Data in B represent mean and standard deviation of TPM from three individuals.**



1795 **Figure 6: Mean bile salt hydrolase, BSH, (A) and *tauD*, taurine dioxy-**  
1796 **genase, (B) log2 normalized expression in controls (gold stars) and**  
1797 **decomposition (boxplots) soils.** Boxplots display the 25th and 75th quartiles  
1798 and median log2 normalized values between all three individuals at each timepoint.  
1799 ANOVA p-value is the result of a hierarchical linear mixed effects model accounting  
1800 for repeated measures of each donor block, while letters denote the results of *post-hoc*  
1801 Tukey test.

1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840





## 1887 Acknowledgements

1888

1889 We would like to thank the Forensic Anthropology Center at the University of  
1890 Tennessee-Knoxville for their help in setting up field experiments. We would like to  
1891 thank Mary Davis for her help in managing the field site and helping to obtain donors  
1892 for this work. This research was funded by a National Institute of Justice Award  
1893 (DOJ-NIJ-2017-R2-CX-0008) to LST and JMD.  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932