


## Local P-adaptivity with the Discontinuous Galerkin Method

Devin Light  
Scott Moe

March 13<sup>th</sup>, 2015

# 1D DG-FEM




**Conservation law weak form:**

- Multiply by test function:

$$\frac{1}{2} \int_{-1}^1 \varphi^{(k)} \left\{ q_{,t} + f(q)_{,x} \right\} d\xi = 0$$

# 1D DG-FEM



**Conservation law weak form:**


- Multiply by test function:

$$\frac{1}{2} \int_{-1}^1 \varphi^{(k)} \{q_{,t} + f(q)_{,x}\} d\xi = 0$$

- Integrate by parts

$$\frac{1}{2} \int_{-1}^1 \varphi^{(k)} q_{,t} d\xi = \frac{1}{\Delta x} \left( \int_{-1}^1 f(q^h) \varphi_{\xi}^{(k)} d\xi - \left[ (\varphi^{(k)} f(q))|_{\xi=1} - (\varphi^{(k)} f(q))|_{\xi=-1} \right] \right)$$

# 1D DG-FEM



**Conservation law weak form:**

- Multiply by test function:

$$\frac{1}{2} \int_{-1}^1 \varphi^{(k)} \{q_{,t} + f(q)_{,x}\} d\xi = 0$$


- Integrate by parts

$$\frac{1}{2} \int_{-1}^1 \varphi^{(k)} q_{,t} d\xi = \frac{1}{\Delta x} \left( \int_{-1}^1 f(q^h) \varphi_{\xi}^{(k)} d\xi - \left[ (\varphi^{(k)} f(q))|_{\xi=1} - (\varphi^{(k)} f(q))|_{\xi=-1} \right] \right)$$

- Galerkin ansatz:

$$q(t, \xi) \sim \sum_{k=0}^N Q^{(k)}(t) \varphi^{(k)}(\xi)$$

# 1D DG-FEM




**Conservation law weak form:**

- Use an Orthogonal Basis:

$$\frac{1}{2} \int_{-1}^1 \varphi^{(\ell)} q_{,t} = \frac{1}{2k+1} \dot{Q}^{(k)}$$

# 1D DG-FEM



**Conservation law weak form:**


- Use an Orthogonal Basis:

$$\frac{1}{2} \int_{-1}^1 \varphi^{(\ell)} q_{,t} = \frac{1}{2k+1} \dot{Q}^{(k)}$$

- Semi-discrete:

$$\dot{Q}_i^{(k)} = \frac{2k+1}{\Delta x} \int_{-1}^1 f(q^h) \varphi_\xi^{(k)} d\xi - \frac{2k+1}{\Delta x} \left[ \varphi^{(k)}(1) F_{i+\frac{1}{2}} - \varphi^{(k)}(-1) F_{i-\frac{1}{2}} \right]$$

# 1D DG-FEM



**Conservation law weak form:**

- Use an Orthogonal Basis:

$$\frac{1}{2} \int_{-1}^1 \varphi^{(\ell)} q_{,t} = \frac{1}{2k+1} \dot{Q}^{(k)}$$

- Semi-discrete:


$$\dot{Q}_i^{(k)} = \frac{2k+1}{\Delta x} \int_{-1}^1 f(q^h) \varphi_\xi^{(k)} d\xi - \frac{2k+1}{\Delta x} \left[ \varphi^{(k)}(1) F_{i+\frac{1}{2}} - \varphi^{(k)}(-1) F_{i-\frac{1}{2}} \right]$$

- For  $k = 0$ :

$$\dot{Q}_i^{(0)} = -\frac{1}{\Delta x} \left[ F_{i+\frac{1}{2}} - F_{i-\frac{1}{2}} \right]$$

# Orthogonal Series


- For continuous functions series coefficients converge.
- Coefficients give us a proxy for error.





# Orthogonal Series


- For continuous functions series coefficients converge.
- Coefficients give us a proxy for error.
- Small coefficients have little effect on our solution.



# Orthogonal Series

- For continuous functions series coefficients converge.
- Coefficients give us a proxy for error.
- Small coefficients have little effect on our solution.
- If  $f$  has a derivative  $\nu$  of bounded variation

$$\|f - f_p\| \leq \sum_{i=p+1}^{\infty} |c_k \varphi^{(k)}(\xi)| \leq \frac{1}{\nu(p - \nu)^\nu}$$



# hp-Adaptivity

- Discretize domain with cells of size  $h$
- h-adaptivity  $\rightarrow$  Adaptive Mesh Refinement

# hp-Adaptivity

- Discretize domain with cells of size  $h$
- h-adaptivity  $\rightarrow$  Adaptive Mesh Refinement
- Discretize using piecewise continuous polynomials of order  $p$ .
- p-adaptivity  $\rightarrow$  Add higher order polynomials:

$$q(t, \xi) = \sum_{k=0}^p Q^{(k)}(t) \varphi^{(k)}(\xi)$$

# hp-Adaptivity

- Discretize domain with cells of size  $h$
- h-adaptivity  $\rightarrow$  Adaptive Mesh Refinement
- Discretize using piecewise continuous polynomials of order  $p$ .
- p-adaptivity  $\rightarrow$  Add higher order polynomials:

$$q(t, \xi) = \sum_{k=0}^p Q^{(k)}(t) \varphi^{(k)}(\xi)$$

- For smooth functions:  
higher order polynomials  $\rightarrow$  **MUCH** better approximation

# Constructing an Algorithm

- For DG  $p$  can be different on each cell

# Constructing an Algorithm

- For DG  $p$  can be different on each cell
- The highest coefficient gives an indication of error on each cell
- Each coefficient requires extra work

# Constructing an Algorithm


- For DG  $p$  can be different on each cell
- The highest coefficient gives an indication of error on each cell
- Each coefficient requires extra work
- Idea:
  - 1 Very small coefficients  $\rightarrow$  can be ignored
  - 2 Large coefficients  $\rightarrow$  need more resolution




# Constructing an Algorithm

- For DG  $p$  can be different on each cell
- The highest coefficient gives an indication of error on each cell
- Each coefficient requires extra work
- Idea:
  - 1 Very small coefficients  $\rightarrow$  can be ignored
  - 2 Large coefficients  $\rightarrow$  need more resolution
- This is complicated by the PDE
  - 1 All PDEs propagate information
  - 2 Hyperbolic PDEs  $\rightarrow$  propagation speed is finite
  - 3 Cells with high-order neighbors may soon become high-order

# Buffer Regions



# Buffer Regions



# Outline of the Algorithm

Pick a maximum order  $p_M$  and project initial conditions

Our p-adaptive algorithm:

# Outline of the Algorithm

Pick a maximum order  $p_M$  and project initial conditions

Our p-adaptive algorithm:

- 1 Eliminate coefficients below a tolerance  $\epsilon$

# Outline of the Algorithm

Pick a maximum order  $p_M$  and project initial conditions

Our p-adaptive algorithm:

- 1 Eliminate coefficients below a tolerance  $\epsilon$
- 2 Determine time-interval between local order modifications
- 3 Set buffer region (determined from characteristic speeds)

# Outline of the Algorithm

Pick a maximum order  $p_M$  and project initial conditions

Our p-adaptive algorithm:

- 1 Eliminate coefficients below a tolerance  $\epsilon$
- 2 Determine time-interval between local order modifications
- 3 Set buffer region (determined from characteristic speeds)
- 4 Evolve coefficients over the interval

# Outline of the Algorithm

Pick a maximum order  $p_M$  and project initial conditions

Our p-adaptive algorithm:


- 1 Eliminate coefficients below a tolerance  $\epsilon$
- 2 Determine time-interval between local order modifications
- 3 Set buffer region (determined from characteristic speeds)
- 4 Evolve coefficients over the interval
- 5 Repeat until we have reached the final time



# Cosbell Advection


# Cosbell Advection

Both methods similarly accurate




# Cosbell Advection

Both methods similarly accurate  
p-Adaptive uses less DOFs!




# Cosbell Advection

Both methods similarly accurate  
p-Adaptive uses less DOFs!  
Less DOFs  $\rightarrow$  Less CPU time



# Burger's Equation



# Conclusions

- 1 Modifying local polynomial degree can be an efficient method of adding/removing DOFs in problems which demand efficiency
  - Atmospheric modeling, Seismology, etc..

# Conclusions

- 1 Modifying local polynomial degree can be an efficient method of adding/removing DOFs in problems which demand efficiency
  - Atmospheric modeling, Seismology, etc..
- 2 Works best where solution is smooth
  - Coefficients decay quickly

# Conclusions

- 1 Modifying local polynomial degree can be an efficient method of adding/removing DOFs in problems which demand efficiency
  - Atmospheric modeling, Seismology, etc..
- 2 Works best where solution is smooth
  - Coefficients decay quickly
- 3 Not as great where solutions are discontinuous
  - Coefficients decay slowly



# The Future

- 1 Local high order time stepping (Local Lax-Wendroff method)
- 2 Simultaneous  $h$ - $p$  adaptivity
- 3 Limiting?
- 4 Adaptive quadrature

# References

- [1] J.-F. Remacle, J. E. Flaherty, and M. S. Shephard, “An adaptive discontinuous galerkin technique with an orthogonal basis applied to compressible flow problems,” *SIAM review*, vol. 45, no. 1, pp. 53–72, 2003.
- [2] M. J. Berger and R. J. LeVeque, “Adaptive mesh refinement using wave-propagation algorithms for hyperbolic systems,” *SIAM Journal on Numerical Analysis*, vol. 35, no. 6, pp. 2298–2316, 1998.
- [3] C. Eskilsson, “An hp-adaptive discontinuous galerkin method for shallow water flows,” *International Journal for Numerical Methods in Fluids*, vol. 67, no. 11, pp. 1605–1623, 2011.
- [4] G. Tumolo, L. Bonaventura, and M. Restelli, “A semi-implicit, semi-lagrangian, p-adaptive discontinuous galerkin method for the shallow water equations,” *Journal of Computational Physics*, vol. 232, no. 1, pp. 46–67, 2013.

# Thank you!

Questions?