

# Longitudinal analysis of AF in NHP baseline *in vivo* retinal images

Alex Matov  
January 2023

## **Computer vision objectives**


- Utilize NHP eye imaging data to develop an automated computer vision software algorithm for the automated detection and quantification of hyper-fluorescent signals on fundus fluorescence images
- Identify the presence of auto-fluorescent signal at prior to injection
- Compare pre- and post-injection images for Visit 1 (prior to surgery-induced disease) to determine if there are increased number of hyper-fluorescent puncta after injection (determine specificity of methodology)
- Compare pre- and post-injection images for Visit 2 (~1 month after surgery-induced disease) to determine if there are increased number of hyper-fluorescent puncta after injection
- Compare Visit 1 and Visit 2 post-injection images (by same time point(s), by eye)

## **Summary of data & findings**


- 24 eyes from 12 NHP over 2 visits 3 months apart
  - AF channel images were acquired prior to and 2 min, 5 min, 15 min & 20 min post dye injection
  - 2 of 13 eyes exhibit no auto-fluorescent (AF) puncta prior to injection
- 
- Multiple analysis parameters to optimize:
    1. Detection -> contrast threshold, edge threshold, #layers in SIFT octave, Gauss filter sigma
    2. Matching -> method, match threshold, max. ratio, distance metric, uniqueness

## Number of detected AF puncta per eye


Cyno 191800 OS, V1 Pre-inj vs V1 Post 2min




## Cyno 191800 OS, V1 Pre-inj vs V1 Post 5min



## Cyno 191800 OS, V1 Pre-inj vs V1 Post 15min



## Cyno 191800 OS, V1 Pre-inj vs V1 Post 30min




## Cyno 191800 OS, V2 Pre-inj vs V2 Post 2min




Image n.a.

Cyno 191800 OS, V2 Pre-inj vs V2 Post 5min



## Cyno 191800 OS, V2 Pre-inj vs V2 Post 15min



Cyno 191800 OS, V2 Pre-inj vs V2 Post 30min

