

## Contents

<b>1</b>	<b>Main Theorem</b>	<b>1</b>
<b>2</b>	<b>Interpreting our Derivations</b>	<b>9</b>
2.1	Properties of continents . . . . .	15
2.2	Generating Examples . . . . .	15
<b>3</b>	<b>Community Detection in a hypergraph</b>	<b>16</b>

## List of Tables

## List of Figures

1	Mountain Ranges . . . . .	9
2	Continents . . . . .	9
3	Weights on hyperedges in a mountain range . . . . .	10
4	Peninsulas off the Mainland . . . . .	11
5	Weights on hyperedges in a continent with peninsulas . . . . .	12
6	Combination of Continents, Mountain Ranges, and Peninsulas . .	13

## 1 Main Theorem

This theory builds on work done by [2] and [1] in studying derivation algebras.

**Definition 1.1.** Let  $V$  be a finite set of size  $m$  and  $\mathcal{H} \subset \binom{V}{n}$ , where  $\binom{V}{n}$  denotes subsets of size  $n$ . Label the entries of  $V$  by  $\{1, 2, \dots, m\}$ . For a given field  $\mathbb{F}$ , define

$$\begin{aligned}
 \varphi : V &\rightarrow \mathbb{F}^m \\
 i &\mapsto e_i \\
 \tilde{\varphi} : V^n &\rightarrow (\mathbb{F}^m)^n \\
 (i_1, i_2, \dots, i_n) &\mapsto (\varphi(i_1), \varphi(i_2), \dots, \varphi(i_n))
 \end{aligned}$$

**Definition 1.2.** Let  $V$  be a finite set of size  $m$  and  $\mathcal{H} \subset \binom{V}{n}$ . We define the *adjacency tensor* corresponding to  $\mathcal{H}$  as

$$\Gamma_{\mathcal{H}} : (\mathbb{F}^m)^n \rightarrow \mathbb{F} \tag{1}$$

$$\Gamma_{\mathcal{H}}(\tilde{\varphi}(X)) := \begin{cases} 1 & X \in \mathcal{H} \\ 0 & \text{else} \end{cases} \tag{2}$$

We will write  $\Gamma := \Gamma_{\mathcal{H}}$  if the hypergraph is clear by context.

**Definition 1.3.** Let  $\langle \Gamma \mid V_l \times \cdots \times V_1 \rightharpoonup V_0$  be a tensor. Its *derivation algebra*  $\text{Der}(\Gamma)$  is

$$\text{Der}(\Gamma) := \left\{ D \in \text{End}(V_0) \times \prod_{i=1}^l \text{End}(V_i) \mid \sum \langle \Gamma \mid D_\alpha v_\alpha, v_{\bar{\alpha}} \rangle = D_0 \langle \Gamma \mid v \rangle \right\}$$

**Definition 1.4.** For a set  $A \in V$ , define

$$\{x, y, A\} := \left\{ \{x, y, a\} \in \binom{V}{3} \mid a \in A \right\}.$$

If  $A \subset \binom{V}{2}$ , and our sets are unordered, define

$$\{x, A\} := \left\{ \{x, a, b\} \in \binom{V}{3} \mid \{a, b\} \in A \right\}$$

If an ordering is necessary, we will write such elements out more explicitly.

**Definition 1.5.** The following notation will be used to define our tensors

- $E_{(i,j,k)}$  is the  $n \times n \times n$  tensor (n to be determined explicitly or by context) with a 1 at position  $(i, j, k)$ , and a zero elsewhere.
- $E_{\{i,j,k\}} := \sum_{\sigma \in \text{Sym}(i,j,k)} E_{(i^\sigma, j^\sigma, k^\sigma)}$  is the  $n \times n \times n$  tensor with a 1 at position  $(a, b, c)$ , where  $(a, b, c)$  is a permutation of  $(i, j, k)$ , for all possible permutations of the three elements
- $E_{\{A,j,k\}} := \sum_{a \in A} \sum_{\sigma \in \text{Sym}(a,j,k)} E_{(a^\sigma, j^\sigma, k^\sigma)}$  is the sum of the tensors  $E_{\{a,j,k\}}$ , indexed by the elements  $a \in A$ . Similarly,  $E_{\{A,j,k\}} := \sum_{a \in A} E_{(a,j,k)}$ .

**Definition 1.6.** Let  $\mathcal{H}$  be an arbitrary hypergraph with vertices  $V$ . Let  $X = \{x_1, x_2, \dots, x_a\} \subset V$ . Define the *neighborhood* of  $X$  to be

$$N_{\mathcal{H}}(X) := \{\{y_1, y_2, \dots, y_b\} \subset V \mid \{x_1, x_2, \dots, x_a, y_1, y_2, \dots, y_b\} \in \mathcal{H}\}$$

**Theorem 1.7.** Consider a hypergraph  $\mathcal{H}$  with  $n$  vertices (labeled 1-n, collectively in the set  $V$ ) and its associated tensor  $\Gamma$ . Let  $\mathcal{D}$  denote the derivation algebra of  $\Gamma$ . Define  $D$  to be the matrix

$$D_0 = \begin{bmatrix} 1 & & \\ & -1 & \\ & & 0_{(n-2) \times (n-2)} \end{bmatrix},$$

where  $0_{m \times m}$  is the  $m \times m$  all zero matrix. Let  $(a, b, c)$  is any hyperedge in  $\mathcal{H}$ . Then after renaming vertices so that this hyperedge is labeled  $(1, 2, c)$ ,  $(D_0, D_0, D_0)$  is a derivation in  $\mathcal{D}$ .

*Proof.* We shall utilize the notation of the Tucker product in our proof. As a 3-valent tensor, there are three modes in which to apply the product. The translation of the derivation conditions in terms of the Tucker product is as follows:

$$\Gamma(E_1 u_i, u_j, u_k) + \Gamma(u_i, E_2 u_j, u_k) + \Gamma(u_i, u_j, E_3 u_k) = 0, \quad \forall (u_i, u_j, u_k) \quad (3)$$

$$(\Gamma \times_1 E_1) + (\Gamma \times_2 E_2) + (\Gamma \times_3 E_3) = \sum_{m=1}^3 \Gamma \times_m E_m = 0 \quad (4)$$

Furthermore, we will rewrite our tensor  $\Gamma$  more explicitly in terms of the previously defined  $E_{\{i,j,k\}}$ . To visualize how to read our derivation, we can imagine each vertex with label  $i$  get assigned a weight given by  $(D_0)_{ii}$ . So the vertex 1 gets weight 1, the vertex 2 gets weight  $-1$ , and all other vertices get weight 0.

Define the sets  $A = N_{\mathcal{H}}(\{1, 2\})$ , and  $X := \{\{i, j, k\} \in \mathcal{H} \mid \{1, 2\} \not\subset \{i, j, k\}\}$ . Clearly,  $\mathcal{H} = X \cup \{a, 1, 2\}$ . We can use these two sets to rewrite  $\Gamma$  as

$$\Gamma = E_{\{A, 1, 2\}} + \sum_{\{i,j,k\} \in X} E_{\{i,j,k\}}.$$

This allows us to write our Tucker product as

$$\begin{aligned} \sum_{m=1}^3 \Gamma \times_m D_0 &= \sum_{m=1}^3 \left( E_{\{A, 1, 2\}} + \sum_{\{i,j,k\} \in X} E_{\{i,j,k\}} \right) \times_m (E_{11} - E_{22}) \\ &= \left( \sum_{m=1}^3 E_{\{A, 1, 2\}} \times_m (E_{11} - E_{22}) \right) + \left( \sum_{m=1}^3 \sum_{\{i,j,k\} \in X} E_{\{i,j,k\}} \times_m (E_{11} - E_{22}) \right) \end{aligned}$$

Now, by construction,

$$\begin{aligned} E_{\{x,y,z\}} \times_1 E_{rr} &= \begin{cases} E_{(r,y,z)} + E_{(r,z,y)} & x = r \\ 0 & \text{else} \end{cases} \\ E_{\{x,y,z\}} \times_2 E_{rr} &= \begin{cases} E_{(y,r,z)} + E_{(z,r,y)} & x = r \\ 0 & \text{else} \end{cases} \\ E_{\{x,y,z\}} \times_3 E_{rr} &= \begin{cases} E_{(y,z,r)} + E_{(z,y,r)} & x = r \\ 0 & \text{else} \end{cases}, \end{aligned}$$

where the condition of  $x = r$  is to say that  $r \in \{x, y, z\}$ . While the specific equality is unknown (as in is it that  $x = r$ ,  $y = r$ , or  $z = r$ ),  $\{x, y, z\}$  is a generic

hyperedge.  $E_{\{x,y,z\}}$  has nonzero values only at coordinates corresponding to permutations of  $\{x,y,z\}$ , and  $(\times_m E_{rr})$  annihilates anything in the  $m^{th}$ -axis which is not in the  $r^{th}$  coordinate. This allows us to simplify our previous sum

$$\sum_{m=1}^3 \Gamma \times_m D_0 = \sum_{m=1}^3 E_{\{A,1,2\}} \times_m (E_{11} - E_{22})$$

From here, we can evaluate the sums over  $m$  and  $\sigma$  explicitly.

$$\begin{aligned} \sum_{m=1}^3 \Gamma \times_m D_0 &= \sum_{m=1}^3 E_{\{A,1,2\}} \times_m (E_{11} - E_{22}) \\ &= ((E_{(1,A,2)} + E_{1,2,A}) - (E_{(2,1,A)} + E_{(2,A,1)})) \\ &\quad + ((E_{(A,1,2)} + E_{2,1,A}) - (E_{(1,2,A)} + E_{(A,2,1)})) \\ &\quad + ((E_{(A,2,1)} + E_{2,A,1}) - (E_{(1,A,2)} + E_{(A,1,2)})) \end{aligned}$$

This sum is separated vertically into the three modes. We only need to utilize four permutations from  $Sym(a,1,2)$  for each mode as the only permutations which contribute non trivially are the ones which have either 1 or 2 in the coordinate corresponding to the particular mode. Now, we can reorganize our sum (and distribute negatives)

$$\begin{aligned} &E_{(1,A,2)} + E_{1,2,A} - E_{(2,1,A)} - E_{(2,A,1)} \\ &+ E_{(A,1,2)} + E_{2,1,A} - E_{(1,2,A)} - E_{(A,2,1)} \\ &+ E_{(A,2,1)} + E_{2,A,1} - E_{(1,A,2)} - E_{(A,1,2)} \\ &= (E_{(1,A,2)} - E_{(1,A,2)}) + (E_{1,2,A} - E_{(1,2,A)}) \\ &\quad + (E_{(A,1,2)} - E_{(A,1,2)}) + (E_{2,1,A} - E_{(2,1,A)}) \\ &\quad + (E_{(A,2,1)} - E_{(A,2,1)}) + (E_{2,A,1} - E_{(2,A,1)}) \\ &= 0 \end{aligned}$$

As  $\sum_{m=1}^3 \Gamma \times_m D = 0$ ,  $(D_0, D_0, D_0)$  is indeed a derivation in  $\mathcal{D}$ .  $\square$

We can now create a family of derivations utilizing this theorem.

**Definition 1.8.** Let  $\vec{x} = (x_1, x_2, \dots, x_{k+1}) \in \mathbb{F}^{k+1}$  denote an arbitrary vector where for all  $i$ ,  $x_i \neq 0$ . Define

$$\begin{aligned} E_k(\vec{x}) &:= \sum_{i=1}^{k+1} x_i E_{2i-1} - x_i E_{2i}, \quad E_{jj} \in M_n(\mathbb{Q}), \\ E_k(\vec{x}) &:= \text{Diag}(x_1, -x_1, x_2, -x_2, \dots, x_{k+1}, -x_{k+1}, 0, 0, \dots, 0) \in M_n(\mathbb{Q}) \\ F_k &:= \sum_{i=1}^{k+2} (-1)^{i-1} E_{ii}, \quad E_{ii} \in M_n(\mathbb{Q}) \end{aligned}$$

$$F_k := \text{Diag}((-1)^0, (-1)^1, (-1)^2, \dots, (-1)^k, (-1)^{k+1}, 0, 0, \dots, 0) M_n(\mathbb{Q}),$$

Note that  $D_0 = \frac{1}{x_1} E_0(\vec{x}) = F_0$ .

**Lemma 1.9.** After renaming the vertices of  $\mathcal{H}$ , choose  $X := \{\{2i-1, 2i, A_i\}\}_{i=1}^{k+1} \subset \mathcal{H}$  such that

$$\begin{aligned} A_i &:= N_{\mathcal{H}}(\{2i-1, 2i\}), \\ \forall j \neq i, 2j, 2j-1 &\notin A_i \\ \forall \{x, y, z\} \in \mathcal{H}, \forall i, 2i \in \{x, y, z\} &\iff 2i-1 \in \{x, y, z\} \end{aligned}$$

With a different renaming of the vertices, choose  $Y := \{\{i, i+1, A_i\}\}_{i=1}^{k+1} \subset \mathcal{H}$  such that

$$\begin{aligned} A_i &:= N_{\mathcal{H}}(\{i, i+1\}), \\ \forall \{x, y, z\} \in \mathcal{H}, i \in \{x, y, z\} &\implies (i+1 \in \{x, y, z\}) \text{ XOR } (i-1 \in \{x, y, z\}) \\ \forall i, j, \{i, i+1, j\}, \{i, i+1, j+1\} &\notin \mathcal{H}, \end{aligned}$$

Let  $\Gamma_X$  ( $\Gamma_Y$ ) be the associated tensor of  $\mathcal{H}$  with respect to the labeling corresponding to the set  $X$  (the set  $Y$ ), and  $\mathcal{D}_X$  ( $\mathcal{D}_Y$ ) the derivation algebra of  $\Gamma_X$  ( $\Gamma_Y$ ). Then  $(E_k(\vec{x}), E_k(\vec{x}), E_k(\vec{x})) \in \mathcal{D}_X$ , and  $(F_k, F_k, F_k) \in \mathcal{D}_Y$ .

*Proof.* First, choices of  $X$  and  $Y$  are always possible by setting  $k = 0$  and choosing a single hyperedge in  $\mathcal{H}$ .

We will first show that  $(F_k, F_k, F_k) \in \mathcal{D}_Y$ . This requires strong induction with the two previous cases, and so a direct computation of the second base case of  $k = 1$  is necessary. For our second base case, our derivation is  $(F_1, F_1, F_1)$ , with

$$F_1 := \begin{bmatrix} 1 & & & \\ & -1 & & \\ & & 1 & \\ & & & 0_{(n-3) \times (n-3)} \end{bmatrix}$$

We factor our tensor as

$$\Gamma_Y := (E_{\{1,2,A_1\}}) + (E_{\{2,3,A_2\}}) + \left( \sum_{\substack{\{x,y,z\} \in \mathcal{H} \\ \{1,2\} \notin \{x,y,z\} \\ \{2,3\} \notin \{x,y,z\}}} E_{\{x,y,z\}} \right)$$

By the conditions used to rename our vertices, these three sets partition all hyperedges in our tensor. Our Tucker product now evaluates to

$$\sum_{m=1}^3 \Gamma_Y \times_m F_k =$$

$$\begin{aligned}
& \sum_{m=1}^3 \left( E_{\{1,2,A_1\}} + E_{\{2,3,A_2\}} + \left( \sum_{\substack{\{x,y,z\} \in \mathcal{H} \\ \{1,2\} \not\subset \{x,y,z\} \\ \{2,3\} \not\subset \{x,y,z\}}} E_{\{x,y,z\}} \right) \right) \times_m F_k \\
&= \sum_{m=1}^3 (E_{\{1,2,A_1\}} \times_m (E_{11} - E_{22} + E_{33})) + (E_{\{2,3,A_2\}} \times_m (E_{11} - E_{22} + E_{33})) \\
&\quad + \left( \sum_{\substack{\{x,y,z\} \in \mathcal{H} \\ \{1,2\} \not\subset \{x,y,z\} \\ \{2,3\} \not\subset \{x,y,z\}}} E_{\{x,y,z\}} \right) \times_m (E_{11} - E_{22} + E_{33}) \\
&= \sum_{m=1}^3 (E_{\{1,2,A_1\}} \times_m (E_{11} - E_{22} + E_{33})) + (E_{\{2,3,A_2\}} \times_m (E_{11} - E_{22} + E_{33})) \\
&= \sum_{m=1}^3 (E_{\{1,2,A_1\}} \times_m (E_{11} - E_{22})) + (E_{\{2,3,A_2\}} \times_m (-E_{22} + E_{33})) \\
&= (E_{(1,2,A_1)} + E_{(1,A_1,2)} - E_{(2,1,A_1)} - E_{(2,A_1,1)} + E_{(2,1,A_1)} + E_{(A_1,1,2)} - E_{(1,2,A_1)} - E_{(A_1,2,1)} \\
&\quad + E_{(A_1,2,1)} + E_{(2,A_1,1)} - E_{(A_1,1,2)} - E_{(1,A_1,2)}) \\
&\quad + (-E_{(2,3,A_2)} - E_{(2,A_2,3)} + E_{(3,2,A_2)} + E_{(3,A_2,2)} - E_{(3,2,A_2)} - E_{(A_2,2,3)} + E_{(2,3,A_2)} + E_{(A_2,3,2)} \\
&\quad - E_{(A_2,3,2)} - E_{(A_2,2,3)} + E_{(A_2,2,3)} + E_{(2,A_2,3)}) \\
&= 0 + 0 = 0
\end{aligned}$$

Now that both base cases have been verified, we may proceed with the induction. Let  $A := N_{\mathcal{H}}$ , i.e. the pairs of vertices which lie in a hyperedge with  $k+2$ , and  $B := \binom{V - \{k+2\}}{3} \cap \mathcal{H}$ , i.e. the hyperedges which are not adjacent to  $k+2$ . This allows us to partition our tensor via adjacency with  $k+2$ . We factor our tensor as

$$\Gamma_Y = \left( \sum_{\{r,s\} \in A} E_{\{r,s,k+3\}} \right) + \left( \sum_{\{x,y,z\} \in \binom{V - \{k+3\}}{3} \cap \mathcal{H}} E_{\{x,y,z\}} \right)$$

We will now continue on with evaluating the Tucker product for the derivation.

$$\begin{aligned}
& \sum_{m=1}^3 \Gamma_Y \times_m F_k \\
&= \sum_{m=1}^3 \Gamma_Y \times_m \left( \sum_{i=1}^{k+2} (-1)^{i-1} E_{ii} \right)
\end{aligned}$$

$$\begin{aligned}
&= \sum_{m=1}^3 \Gamma_Y \times_m ((-1)^{k+1} E_{k+2,k+2} + (-1)^k E_{k+1,k+1}) \\
&+ \sum_{m=1}^3 \Gamma_Y \times_m \sum_{i=1}^k (-1)^{i-1} E_{ii} \\
&= \sum_{m=1}^3 \left( \sum_{\{r,s\} \in A} E_{\{r,s,k+2\}} + \sum_{\{x,y,z\} \in B} E_{\{x,y,z\}} \right) \times_m (-1)^k (E_{k+1,k+1} - E_{k+2,k+2}) \\
&+ \sum_{m=1}^3 \Gamma \times_m F_{k-2,k-2} \\
&= (-1)^k \left( \sum_{m=1}^3 \left( \sum_{\{r,s\} \in A} E_{\{r,s,k+2\}} + \sum_{\{x,y,z\} \in B} E_{\{x,y,z\}} \right) \times_m (E_{k+1,k+1} - E_{k+2,k+2}) \right) + 0 \\
&= (-1)^k \sum_{m=1}^3 \left( \sum_{\{r,s\} \in A} E_{\{r,s,k+2\}} \right) \times_m (E_{k+1,k+1} - E_{k+2,k+2}) \\
&+ (-1)^k \sum_{m=1}^3 \left( \sum_{\{x,y,z\} \in B} E_{\{x,y,z\}} \right) \times_m (E_{k+1,k+1} - E_{k+2,k+2}) \\
&= (-1)^k \sum_{m=1}^3 \left( \sum_{\{r,s\} \in A} E_{\{r,s,k+2\}} \right) \times_m (E_{k+1,k+1} - E_{k+2,k+2})
\end{aligned}$$

Where  $\sum_{m=1}^3 \left( \sum_{\{x,y,z\} \in B} E_{\{x,y,z\}} \right) \times_m (E_{k+1,k+1} - E_{k+2,k+2}) = 0$  because by construction neither  $k+1$  nor  $k+2$  are in any  $\{x,y,z\} \in B$ , and so the tucker product will always evaluate to zero. We proceed with our calculations. For the following we factor out  $(-1)^k$  for brevity, as our end result is to show that the sum evaluates to zero.

$$\begin{aligned}
&\sum_{m=1}^3 \left( \sum_{\{r,s\} \in A} E_{\{r,s,k+2\}} \right) \times_m (E_{k+1,k+1} - E_{k+2,k+2}) \\
&= \sum_{m=1}^3 \sum_{\{r,s\} \in A} (E_{\{r,s,k+2\}} \times_m (E_{k+1,k+1} - E_{k+2,k+2})) \\
&= \sum_{m=1}^3 \sum_{\substack{\{r,s\} \in A \\ k+1=s}} (E_{\{r,k+1,k+2\}} \times_m (E_{k+1,k+1} - E_{k+2,k+2}))
\end{aligned}$$

$$+ \sum_{m=1}^3 \sum_{\substack{\{r,s\} \in A \\ k+1 \notin \{r,s\}}} (E_{\{r,s,k+2\}} \times_m (E_{k+1,k+1} - E_{k+2,k+2}))$$

By the construction of  $Y$  we have that the second summand evaluates to zero as there is no pair of vertices adjacent to  $k+2$  which does not include  $k+1$ . We define  $A' := N_{\mathcal{H}}(\{k+1, k+2\})$ . Our sum thus further simplifies as

$$\begin{aligned} & \sum_{m=1}^3 \sum_{a \in A'} (E_{\{a,k+1,k+2\}} \times_m (E_{k+1,k+1} - E_{k+2,k+2})) \\ &= ((E_{(k+1,A',k+2)} + E_{(k+1,k+2,A')} - E_{(k+2,A',k+1)} - E_{(k+2,k+1,A')}) \\ & \quad + (E_{(A',k+1,k+2)} + E_{(k+2,k+1,A')} - E_{(A',k+2,k+1)} - E_{(k+1,k+2,A')}) \\ & \quad + (E_{(A',k+2,k+1)} + E_{(k+2,A',k+1)} - E_{(A',k+1,k+2)} - E_{(k+1,A',k+2)})) \\ &= 0 \end{aligned}$$

Now, we must show that  $(E_k(\vec{x}), E_k(\vec{x}), E_k(\vec{x})) \in \mathcal{D}_X$ . Unlike the previous family of derivations, we do not need to rely on strong induction. Define  $A := N_{\mathcal{H}}(\{2k+1, 2k+3\})$ . Our evaluation of the derivation is as follows:

$$\begin{aligned} & \sum_{m=1}^3 \Gamma \times_m E_k(\vec{x}) \\ &= \sum_{m=1}^3 \left( \sum_{\substack{\{r,s,t\} \in \mathcal{H} \\ 2k+2 \notin \{r,s,t\}}} E_{\{r,s,t\}} + E_{\{2k+1,2k+2,A\}} \right) \times_m \left( \sum_{i=1}^{k+1} x_i E_{2i-1} - x_i E_{2i} \right) \\ &= \sum_{m=1}^3 \left( \sum_{\substack{\{r,s,t\} \in \mathcal{H} \\ 2k+2 \notin \{r,s,t\}}} E_{\{r,s,t\}} + E_{\{2k+1,2k+2,A\}} \right) \times_m \left( \sum_{i=1}^k x_i E_{2i-1} - x_i E_{2i} \right) \\ & \quad + \sum_{m=1}^3 \left( \sum_{\substack{\{r,s,t\} \in \mathcal{H} \\ 2k+2 \notin \{r,s,t\}}} E_{\{r,s,t\}} + E_{\{2k+1,2k+2,A\}} \right) \times_m (x_{k+1} E_{2k+1} - x_{k+1} E_{2k+2}) \\ &= \sum_{m=1}^3 \Gamma \times_m E_{k-1}((x_1, x_2, \dots, x_k)) \\ & \quad + \sum_{m=1}^3 \left( \sum_{\substack{\{r,s,t\} \in \mathcal{H} \\ 2k+2 \notin \{r,s,t\}}} E_{\{r,s,t\}} + E_{\{2k+1,2k+2,A\}} \right) \times_m (x_{k+1} E_{2k+1} - x_{k+1} E_{2k+2}) \\ &= \sum_{m=1}^3 E_{\{2k+1,2k+2,A\}} \times_m (x_{k+1} E_{2k+1} - x_{k+1} E_{2k+2}) \end{aligned}$$



$$\begin{aligned}
&= \sum_{m=1}^3 (E_{\{2k+1,2k+2,A\}} \times_m (x_{k+1} E_{2k+1})) - (E_{\{2k+1,2k+2,A\}} \times_m (x_{k+1} E_{2k+2})) \\
&= x_{k+1} (E_{(2k+1,2k+2,A)} + E_{(2k+1,A,2k+2)} + E_{(2k+2,2k+1,A)} \\
&\quad + E_{(A,2k+1,2k+2)} + E_{(2k+2,A,2k+1)} + E_{(A,2k+2,2k+1)}) \\
&\quad - x_{k+1} (E_{(2k+2,2k+1,A)} + E_{(2k+2,A,2k+1)} + E_{(2k+1,2k+2,A)} \\
&\quad + E_{(A,2k+2,2k+1)} + E_{(2k+1,A,2k+2)} + E_{(A,2k+1,2k+2)}) \\
&= 0
\end{aligned}$$

□

## 2 Interpreting our Derivations

We will now aim to better understand what these theorems tell us about our hypergraphs. The renaming coming from  $Y$  is used to view our hypergraph as it is in the following figures.




Figure 1: Mountain Ranges




Figure 2: Continents

In Figure 1, we can imagine certain distinguished hyperedges, which we call *mountains*, forming a *mountain range*, where each mountain connects to an adjacent mountain at a vertex. At the top of the mountains are *continents*, which in the Figure 2 we can see represent an arbitrary conglomerate of hyperedges. Continents may intersect with each other, but the key is that the bases of the mountain range forms the given structure. What our theorem says is that if we assign weights to each vertex as in Figure 3, if we take any hyperedge in our hypergraph and add together the weights on each vertex, they sum to zero.




Figure 3: Weights on hyperedges in a mountain range

While clouds may intersect, it is relatively easy to check if this occurs. If we were to create a new hypergraph by deleting the mountain range, we can check for connected components.

**Definition 2.1.** Given a hypergraph  $\mathcal{H} = (V, E)$ , the *incidence graph* of  $\mathcal{H}$  is a bipartite graph  $\mathcal{G} = (V_V \cup V_E, E_H)$  constructed as follows:

- $V_V$  are the vertices of  $\mathcal{H}$ .
- $V_E$  are vertices corresponding to the hyperedges  $e$  of  $\mathcal{H}$ .
- There is an edge from a vertex  $v' \in V_V$  to a vertex in  $e \in V_E$  if the vertex  $v'$  in  $\mathcal{H}$  is contained in the hyperedge  $e$  in  $\mathcal{H}$ .

**Lemma 2.2.** A hypergraph  $\mathcal{H}$  is connected if and only if its incidence graph  $\mathcal{G}$  is connected.

*Proof.* If a hypergraph is connected, then between any two vertices there is a path connecting them where the path follows adjacency through hyperedges. In the incidence graph, this same path can be followed. If vertex  $v_1$  and  $v_2$  are connected via shared adjacency in hyperedge  $e_1$ , on the incidence matrix  $v_1$  and  $v_2$  are both adjacent to  $e_1$ , and so a path connects them. Therefore, any path in  $\mathcal{H}$  corresponds to a path in  $\mathcal{G}$ , and vice versa.  $\square$

By this lemma, checking if a hypergraph is connected is equivalent to checking if a bipartite graph is connected.

For the labeling of  $X$ , we have the following figure




Figure 4: Peninsulas off the Mainland

In Figure 4, we have a central continent which has coming from it various *peninsulas*. Our theorem says that if we assign weights to each vertex as in Figure 5, on any hyperedge the sum of the weights on the vertices equals to zero.

**Corollary 2.3.** With an appropriate relabeling to avoid unwanted overlaps, linear combinations of  $(E_a(\vec{x}_a), E_a(\vec{x}_a), E_a(\vec{x}_a))'$ s and  $(F_b, F_b, F_b)'$ s (with varying  $a$ 's and  $b$ 's) form a derivation which is interpretable as a combination of continents, mountain ranges, and peninsulas.

Working with the graphical interpretations of derivations assigning weights to vertices, this corollary says that we can build out our hypergraph structure by isolating mountain ranges, continents, and peninsulas all glued together, and that there exists a derivation which will identify this process. Figure 6 shows what this may look like.




Figure 5: Weights on hyperedges in a continent with peninsulas

In this example, the corresponding derivation would be  $(D, D, D)$ , where

$$D := \begin{bmatrix} X' & & & & & & \\ & -x_3 & & & & & \\ & & x_3 & & & & \\ & & & -x_3 & & & \\ & & & & -x_3 & & \\ & & & & & Y' & \\ & & & & & & Z' \\ & & & & & & & 0 \end{bmatrix},$$

$$X' := \begin{bmatrix} x_1 & & & & & & \\ & -x_1 & & & & & \\ & & x_2 & & & & \\ & & & -x_2 & & & \\ & & & & -x_4 & & \\ & & & & & \ddots & \\ & & & & & & x_{a+1} \\ & & & & & & & -x_{a+1} \end{bmatrix}$$




Figure 6: Combination of Continents, Mountain Ranges, and Peninsulas

$$Y' := \begin{bmatrix} y_1 & & & & & & \\ & -y_1 & & & & & \\ & & y_2 & & & & \\ & & & -y_2 & & & \\ & & & & -y_4 & & \\ & & & & & \ddots & \\ & & & & & & y_{b+1} \\ & & & & & & & -y_{b+1} \end{bmatrix}$$

$$Z' := \begin{bmatrix} z_1 & & & & & & \\ & -z_1 & & & & & \\ & & z_2 & & & & \\ & & & -z_2 & & & \\ & & & & -z_4 & & \\ & & & & & \ddots & \\ & & & & & & z_{c+1} \\ & & & & & & & -z_{c+1} \end{bmatrix}$$

When considering how continents and mountain ranges may be connected, we have thus far only considered trees. We must now consider loops.

**Lemma 2.4.** If a mountain range loops with  $2n$  different mountains, then (after relabeling)  $F_{2n}$  is a derivation for the corresponding tensor. If the loop is constructed with  $2n+1$  mountains, then after a relabeling  $F_{2n} + (-1)^{2n+1} 2E_{2n+2, 2n+2}$  is a derivation for the corresponding tensor.

*Proof.* With our correspondence of elements of the Cartan subalgebra of the Derivation algebra and weights on a hypergraph, we can give the proof of this lemma with an argument on the weights of hyperedges. The following two pictures show this argument: (insert tikz image here).  $\square$

Finally, the third way in which we can get a derivation is by taking the sub-hypergraph  $[\{1, 2, 3\}, \{3, 4, 5\}, \{5, 6, 7\}, \{7, 8, 9\}]$ , and give the vertex 4 a weight of 1, the vertex 5 a weight  $-1$ , and the vertex 6 a weight of 1. (this might be covered already by  $F_k$ , and my future argument sying mountain peaks connecting continents cannot have weight other than zero).

Now, we need to show that continents **MUST** have weight zero. This requiries us to further expand on our definition of a continent

**Definition 2.5.** A connected component of a hyergraph is a *continent* if there does not exist mountain ranges nor peninsulas within the connected component.

In other words, after identifying all mountain ranges and peninsulas, the hyperedges we are left with are within their respective continents.

## 2.1 Properties of continents

We will now work with the subhypergraphs which are continents.

**Lemma 2.6.** if there are too many vertices in a continent which are contained in only one hyperedge, then the adjacency tensor of the continent will be degenerate.

*Proof.* (refer back to spiky urchin).  $\square$

Our goal is to show that if we have any vertex in the continent which has weight not equal to zero, this will eventually lead to a contradiction of there not being any mountain ranges or peninsulas.

## 2.2 Generating Examples

There are two ways in which we can construct standard hypergraph structures from a graph. Both enlist the use of cones. In the first, we take an arbitrary graph  $\mathcal{G}$  with vertex set  $V$  and edge set  $E$ . We then construct a hypergraph  $\mathcal{H}$  with vertex set  $V \cup \{\infty\}$  and edge set  $E' := \{\{v_i, v_j, \infty\} \mid \{v_i, v_j\} \in E\}$ . This is equivalent to drawing our graph  $\mathcal{G}$ , including a “point at infinity,” and connecting each edge to this point at infinity.

When looking at our adjacency tensor, the evaluations are as follows

$$\Gamma(x, y, z) := \begin{cases} 1 & \{(x, y) \in E\} \wedge \{z = \infty\} \\ 0 & \text{else} \end{cases}, \quad (5)$$

where we account for symmetry in our evaluations. Once again, we will look for symmetric subalgebras of the Cartan subalgebra (diagonal matrices). For ease of our explanations, we will utilize our interpretation of weights on vertices, and that a triple of the same diagonal matrix are in the derivation algebra if and only if their corresponding weight assignments sum to zero on each hyperedge.

In this interpretation, the weight on  $\infty$  is  $a_\infty$ , and a triple of matrices is a derivation if and only if the weight assignments on the base graph  $\mathcal{G}$  has the property that the sum of weights on adjacent vertices (adjacency being defined as lying on the same hyperedge) is zero. This holds if and only if  $\mathcal{G}$  is bipartite.

*Proof.* Let  $a_\infty$  be the weight on  $\infty$ . Beginning with any pair of vertices which lie on a hyperedge with  $\infty$ , one can be assigned the weight  $x_i$  and may be given the label of 1. The other is necessarily assigned the weight of  $-x_i - a_\infty$ , and may be given the label of 2. If  $\infty$  and 1 lie on a hyperedge with a third vertex 0 which is not 2 then 0 must be given a weight of  $-x_i - a_\infty$ . Equivalently, If  $\infty$  and 2 lie on a hyperedge with a third vertex 3 which is not 1, then 3 must be given a weight of  $x_i$ . Furthermore,  $\{1, 3, \infty\}$  cannot be a hyperedge, and neither can  $\{0, 2, \infty\}$ . This construction provides us with a bipartite graph.

Likewise, if we were to begin with a bipartite 2-graph, then we can assign each side of the bipartite graph its own weight  $x_i$  and  $y_i$ , and then the point at infinity will receive a weight of  $-x_i - y_i$ .  $\square$

In this proof,  $i$  is a number of the number of connected components in our graph  $\mathcal{G}$ , which will determine how many different weight-pairs we need. Our derivations are thus of the form  $a_\infty E_{\infty,\infty} + x_i E_{11} + (-x_i - a_\infty) E_{22}$ .

In the second way of constructing a hypergraph from a graph, we instead generate a cone for each edge in the graph  $\mathcal{G}$ . We do this by assigning a distinct point at infinity  $\infty_{xy}$  for each edge  $(x, y) \in E$ . In this scenario, we are less restricted in our weight assignment. We can assign any weight to the vertices in  $\mathcal{G}$ . All we require is that if  $a_x$  and  $a_y$  are weights of adjacent vertices  $x$  and  $y$ , the weight on  $\infty_{xy}$  is  $-a_x - a_y$ . This gives us derivations with matrices of the form  $a_x E_{11} + b_y E_{22} + (-a_x - b_y) E_{33}$ .

### 3 Community Detection in a hypergraph

One way in which we can use these results is in detecting community structure. Typically, “communities” are defined as “collections of vertices in a graph which are ‘more’ connected to each other than ‘outside’ vertices.” This kind of definition leaves a very vague notion of distinct communities, which allows for a more fluid identification of communities, as well as the use of eigentheory and SVD to obtain identifications. Our use of derivations provides a stricter and more easily identifiable definition of community structure.

**Definition 3.1.** Let  $\mathcal{H}$  be a hypergraph. A *continental structure* on  $\mathcal{H}$  is an relabeling and identification of vertices so that a linear combination of  $(E_a(\vec{x}_a), E_a(\vec{x}_a), E_a(\vec{x}_a))'$ s and  $(F_b, F_b, F_b)'$ s with coefficients of 1 form a derivation which is interpretable as a combination of continents, mountain ranges, and peninsulas, where the continents do not intersect. The number of continents is  $\sum_{i=1}^j (b_i + 1)$ , where  $j$  is the number of mountain ranges, and  $b_i + 1$  is the number of mountains in the  $i^{th}$  mountain range.

### References

- [1] Peter A. Brooksbank, Joshua Maglione, and James B. Wilson. Tensor isomorphism by conjugacy of lie algebras. *Journal of Algebra*, 604:790–807, 2022.
- [2] Uriya First, Joshua Maglione, and James B. Wilson. A spectral theory for transverse tensor operators. 2022.