Observational Incompleteness: From Deterministic Dynamics to Quantum Observations

Alex Maybaum

October 2026

Status: DRAFT PRE-PRINT
Classification: Theoretical Physics / Foundations

Abstract

Observational incompleteness (OI) is the hypothesis that every physical observer is a proper part of a finite reversible deterministic system and records only the visible marginals of its dynamics. At every finite horizon, all finite stochastic laws, the visible marginals of finite reversible dynamics and the records of finite unitary systems measured in one fixed basis coincide, so stochastic, quantum-like records can arise exactly from deterministic dynamics; the known components (Egri et al.; Barandes) are stated as one machine-checked theorem. Because Markov laws lie in every class, the discriminating content is memory: every faithful realization carries the law's conditional memory in its hidden state. On one finite reversible system with a hidden prior shared by every root and drawn independently of it, the rooted visible family, with a row for every root including roots of visible probability zero, is P-indivisible somewhere in its recurrence cycle whenever some rooted transition matrix is not a permutation, which every non-deterministic visible step supplies; with a full-support visible prior this indivisibility is equivalent to conditional memory $M_k > 0$ at some order k below the recurrence time. Exact finite operational quantum mechanics is characterized by five completion conditions that OI alone does not imply. OI separates the zero-point sum from the gravitational source, and conservation of black-hole information from its accessible recovery; open are the completion conditions a physical substratum supplies, the Born exponent and the gravitational field equations.

1. Introduction

1.1 Embedded observation

An observer inside a closed system registers only part of its configuration. If the whole evolves by a bijection, hidden states compatible with the same visible state can send it to different futures, so the visible sector evolves stochastically although the whole evolves deterministically.

This is modelled by a triple (S, φ, V) [Main §1.2]: a finite total system $S = \mathcal{C}_V \times \mathcal{C}_H$ of visible configurations $\mathcal{C}_V$ and hidden configurations $\mathcal{C}_H$, a bijection φ of S, and an observer V that registers $x \in \mathcal{C}_V$ and not $h \in \mathcal{C}_H$. The bijection is the reversible representative of the hidden dynamics, motivated by observed near-unitarity, and finiteness of all of S is posited [Main §1.2, Remark (status of the early commitments)]. The finite reversible system (S, φ) is the *substratum*.

Observational incompleteness (OI) is the physical hypothesis that every physical observer is in this situation. Here incompleteness means that the hidden sector is inaccessible. It is not joint unsharpness of non-commuting observables: it is present for a single fixed readout, and no uncertainty relation constrains fixed-readout visible laws [Main §3.4].

1.2 The idea and its consequences

Marginalizing a reversible dynamics over an unobserved register can produce stochastic laws, conditional memory, and failures of stochastic divisibility, in which no stochastic map carries an earlier visible transition matrix to a later one (when the earlier matrix is invertible, the unique candidate has a negative entry). The stochastic–quantum correspondence relates these features to quantum theory [Bar23; Bar25], and at the level of finite-horizon observable laws it is exact and universal (Theorem 1, §2.2): every finite stochastic law, every fixed-basis quantum record among them, is the visible marginal of a finite reversible deterministic system. Apparently stochastic, quantum-like observations can therefore arise exactly from deterministic dynamics. Because the correspondence is universal, the discriminating content lies in where the memory of a law must sit (§2.4) and in what finite recurrence does with it (§2.5). Section 3 states how far the correspondence reaches toward operational quantum mechanics, §§4–5 give two applications, and §6 sorts the results by status. Proofs, the status ledger and the probe code are in [Main], which this paper introduces; gravitational applications are in [GR] and [Book].

Contributions. The paper is expository and claims no new theorem; each known component is attributed where it is stated (§§2.2, 2.4, 2.5). The results of §§3.2–3.5 are proved in [GR] and [Main], the Bell ceiling being Bell's theorem under the framework's hypotheses, and those of §5 in unpublished research records with exact or exhaustive checks [BHa; BHb]; the novelty of the §§3.2–3.3 and §5 results has not been assessed. Deterministic systems have been analysed with quantum methods before, for example in [tH14].

Conventions. "Machine-checked" means checked in Lean 4/Mathlib [dMU21; Mathlib20] at commit `17b830bc` by the recipe under "Code, data and formal verification" (declarations in Appendix A); "proved on paper" means a written proof without a machine check; an "exact probe" is an exhaustive computation in exact rational arithmetic at small size. A *faithful realization* of a law is a bijection φ of $\mathcal{C}_V \times \mathcal{C}_H$ with an initial law under which the visible trajectory has that law.

2. Theoretical foundation: the established results and their assumptions

2.1 Setting

Fix finite sets $\mathcal{C}_V$ (with $|\mathcal{C}_V| \geq 2$) and $\mathcal{C}_H$, a bijection φ of $\mathcal{C}_V \times \mathcal{C}_H$, and a horizon K. A law is a probability distribution on visible trajectories $(x_0, \dots, x_K) \in \mathcal{C}_V^{K+1}$. In a realization, X_t and H_t are the visible and hidden coordinates at time t, $X_{<t} = (X_0, \dots, X_{t-1})$ is the visible past and $X_{>t} = (X_{t+1}, \dots, X_K)$ the visible future. Information is measured in bits, $\|\mu - \nu\|_{\mathrm{TV}} = \frac{1}{2}\sum_k |\mu_k - \nu_k|$ is the total-variation distance, and $g_{\#}\mu$ is the pushforward of a law μ under a map g.

The visible law of a realization depends on the dynamics, the partition and the initial law (the prior), which is part of the realization datum, not selected by invariance under φ; the counting measure is a maximal-entropy default [Main §1.2, Remark (the measure as a realization datum)].

Three classes of law are compared [Main §3.4]:

- S_K: all probability distributions on $\mathcal{C}_V^{K+1}$.
- D_K: the visible trajectory laws of a bijection of $\mathcal{C}_V \times \mathcal{C}_H$, for some finite $\mathcal{C}_H$, under an initial law that may correlate the visible and hidden coordinates.
- Q_K: the *fixed-basis Born records*, the laws of finite-dimensional unitary systems read out by projective Born measurement in one fixed basis at every step, from an initial state diagonal in that basis, the record being a fixed function of each outcome.

2.2 The finite-horizon correspondence

Theorem 1 (finite-horizon stochastic–reversible–unitary equivalence [Main §3.4]). *For every finite visible alphabet and every finite horizon K,*

$$\mathsf{S}_K = \mathsf{D}_K = \mathsf{Q}_K,$$

so each class is the set of all probability laws on $\mathcal{C}_V^{K+1}$.

Theorem 1 is machine-checked: the three inclusions $\mathsf{S}_K \subseteq \mathsf{D}_K \subseteq \mathsf{Q}_K \subseteq \mathsf{S}_K$ and their compositions, with no construction entering as a hypothesis (Appendix A).

Realization. For $\mathsf{S}_K \subseteq \mathsf{D}_K$ the realization stores a sampled trajectory, with a clock, in the hidden register [Main §3.4, Theorem (finite-horizon stochastic–reversible–unitary equivalence); EGGH26, Prop. 13]. Every law also has a realization whose hidden prior is independent of x_0, by the finite-horizon process dilation and its response-table extension for arbitrary kernels (proved on paper, with exact probes) [Main §3.4, Theorem (finite-horizon process dilation) and its Extension].

Quantum representation. For $\mathsf{D}_K \subseteq \mathsf{Q}_K$ the representation is the permutation unitary $U_\varphi|x,h\rangle = |\varphi(x,h)\rangle$, read out in the configuration basis at every step [Main §3.4]; $\mathsf{Q}_K \subseteq \mathsf{S}_K$ is the Born rule.

Attribution. The trajectory-labelling realization is [EGGH26, Prop. 13], packaged as one iterated bijection; the independent-prior form is their Prop. 16, without reversibility, and its Markov one-step case is the random-mapping representation [LP17, Prop. 1.5]. A permutation dynamics is a special case of a unistochastic process, one with transition probabilities $|U_{ij}|^2$ for a unitary U [Bar25, p. 12 of arXiv v3; Bar23 §4.3; Bar26 §4]; the one-step $|\mathcal{C}_V|^2$-dimensional ancilla dilation of [Main §3.1] is that of [Bar26, Thm 3.3; Sch21, Thm 2]; and the every-step record law is [SGD19, Eq. (3)], whose consistency under omitted measurements [SGD19; Milz20] Q_K, measuring at every step, does not meet.

2.3 Scope of the correspondence

Markov laws lie in all three classes, so neither a finite reversible realization nor a fixed-basis unitary representation distinguishes a law with memory from one without [Main §3.4, Remark (what this equivalence does and does not say)].

The relation $p = |U|^2$ is part of the definition of Q_K, not an output. The unitaries supplied are permutations, with entries 0 and 1, so the recovered table is the same for $|U|^q$ at every $q > 0$, and the quadratic exponent is not selected [Main §3.4; Bar26, Thm 3.1].

A finite bijection has finite order L, so one fixed realization has an L-periodic multi-time law: the correspondence holds per horizon, and a nonperiodic law has no exact all-time realization by one fixed finite substratum [Main §3.4].

2.4 Where the memory lives

Write $M_t := I(X_{<t}; X_{t+1} \mid X_t)$ for the conditional memory at time t. In a realization, write $f_x(h) = \pi_V(\varphi(x,h))$, with π_V the projection onto $\mathcal{C}_V$, and μ_ω for the hidden posterior after a positive-probability visible history $\omega = (x_0, \dots, x_t)$ ending at $x_t = x$.

Theorem 2 (unavoidable hidden predictive memory [Main §3.4]). *In every faithful realization:*

(a) $P(X_{t+1} \mid \omega) = (f_x)_{\#}\mu_\omega$*;*

(b) for histories ω, ω' ending at x,

$$\|P(X_{t+1} \mid \omega) - P(X_{t+1} \mid \omega')\|_{\mathrm{TV}} \le \|\mu_\omega - \mu_{\omega'}\|_{\mathrm{TV}};$$

(c) $M_t \le I(X_{<t}; H_t \mid X_t) \le \log_2 |\mathcal{C}_H|$.

(b) and the first inequality of (c) are data-processing inequalities. (a)–(c) are machine-checked for a joint law of history, present and hidden state; their identification with a realization's time-t law is proved on paper [Main §3.4, Theorem (unavoidable hidden predictive memory); CT06].

A *memory witness* at t is a pair of positive-probability histories ending at the same x_t with different next-step laws; by the vanishing criterion for conditional mutual information [CT06], a law has one exactly when $M_t > 0$.

Corollary (proved on paper [Main §1.5]). $M_t > 0$ holds exactly when every faithful realization carries readout-relevant hidden memory at t, that is, a pair of posteriors with different pushforwards. The forward direction rests on Theorem 2(a) in each realization; the converse on Theorem 2(a) in one realization, which Theorem 1 supplies (`S_imp_D`); both use the standard vanishing criterion for conditional mutual information [CT06], which, like the time-t identification, is not machine-checked.

The hidden sector also has a minimum size and a minimum memory. With $I^* = \max_t I(X_{<t}; X_{>t} \mid X_t)$, a realization of the law needs $|\mathcal{C}_H| \ge 2^{I^*}$ (machine-checked in the joint-law form of Theorem 2). Over deterministic completions, not required to be bijective, the least hidden predictive memory $I(X_{<t}; H_t \mid X_t)$ is exactly $I(X_{<t}; X_{>t} \mid X_t)$, attained by the canonical predictive quotient, and equals M_t exactly when $I(X_{<t}; X_{>t+1} \mid X_t, X_{t+1}) = 0$, while bijective realizations run from time 0 carry $H(X_{<t} \mid X_t)$, Markov laws included [Main §3.4, Remark (a variational question)]; the lower bound is machine-checked for bijective realizations and proved on paper for non-injective ones. Conversely, memory has a floor: if two values of $X_{<t}$ followed by a common present state X_t, each of joint probability at least p_0, induce next-step laws at total-variation distance at least δ, then $M_t \ge p_0 \delta^2 / \ln 2$, by Pinsker's inequality (proved on paper, with exact probes) [Main §2.3, Lemma (accessible conditional memory from readback); CT06]. Observed memory thus bounds an inaccessible quantity: measurable conditional information fixes a lower bound on the hidden sector's size and the exact minimum of the predictive memory a deterministic completion must carry.

Four conditions on a hidden sector are named in [Main §1.3], as diagnostics of a realization rather than hypotheses of Theorem 1: (C1) influence; (C2) persistence of records; (C3) capacity, $\log_2 |\mathcal{C}_H| \ge I^*$; and (C4) history readback, a memory witness at some accessible order. In a faithful realization (C4) coincides with $M_t > 0$, and the Corollary places it in the hidden state; (C1) follows from (C4), and (C3) holds by the capacity bound. (C2) has two forms. (C2-structural), the persistence of a stored record until its readback, holds by reversibility in every bijective realization and constrains coarse-grained descriptions of the hidden sector; (C2-slow), a hidden coarse-grained mixing time long

compared with the accessible window, is one mechanism for it, and fast hidden dynamics that conserves the record is another [Main §1.3]. The necessity of (C2) is a theorem in the coupling-step model, in which the next visible state is a function of the current visible and hidden states: observed conditional memory $I^* > 0$ at an accessible order forces some realized hidden conditional at a coupling time to lie at total-variation distance at least $\varepsilon_{\min}(I^*, |\mathcal{C}_V|) > 0$ from the model's reference prior, with no mixing hypothesis, and under a mixing hypothesis this becomes the slow-bath form, a lower bound on the mixing time [Main §3.4, Theorem (physical memory: C2, process form)].

2.5 Recurrence and indivisibility

Fix one hidden prior shared by every root, and write Γ_t for the *rooted transition matrix* from time 0 to time t, with a row for every $x_0 \in \mathcal{C}_V$, roots of visible probability zero included. The family is *P-divisible on* $\{0, \dots, n\}$ if for all $s < t \leq n$ some stochastic $\Delta_{s,t}$ satisfies $\Gamma_t = \Gamma_s \Delta_{s,t}$. A stochastic divisor contracts total variation, so under P-divisibility $\|\delta_a \Gamma_t - \delta_b \Gamma_t\|_{\mathrm{TV}}$, with δ_a the point mass at a, cannot increase. This is the divisibility criterion of [RHP14, Thm 2.4] and the data-processing direction of [BD16, Prop. 1].

Theorem 3 (recurrence and overlap [Main §2.3]). *Let φ be a bijection of $\mathcal{C}_V \times \mathcal{C}_H$ with one hidden prior shared by every root. If two rows of Γ_s have a positive entry in a common column and $\Gamma_n = I$ for some $n > s$, the family is not P-divisible on $\{0, \dots, n\}$.*

Two such rows are at total-variation distance below 1 at time s and exactly 1 at time n, which the contraction criterion forbids; Theorem 3 is machine-checked in its rooted form, and Doukas states the same argument concurrently [Dou26, Prop. 1]. Two rows of a square stochastic matrix share a positive column exactly when it is not a permutation matrix, and [Main §2.3] also proves the theorem from the lemma that a finite square stochastic matrix with a stochastic one-sided inverse is a permutation matrix [BB12, Problem 23] (machine-checked).

Finite recurrence supplies the return, $\Gamma_L = I$ for the order L of φ:

Corollary (recurrence-cycle indivisibility). *Let φ have order L, with one hidden prior shared by every root and drawn independently of it, and let Γ_t keep a row for every root, roots of visible probability zero included. If some Γ_s with $0 < s < L$ is not a permutation, the rooted family is not P-divisible on $\{0, \dots, L\}$. Two distinct positive-probability visible histories with a common endpoint, in particular a memory witness, supply such a Γ_s, and so does every non-deterministic visible step, also when the law is Markov on every accessible window. With a full-support visible prior, this failure is equivalent to $M_k > 0$ at some order $k < L$.*

This is [Main §2.3, Theorem (history readback + finite recurrence implies indivisibility) and Scope]: the first sentence is Theorem 3 with $n = L$, since two rows of a non-permutation Γ_s share a positive column, and Theorem 3 is machine-

checked in its rooted form; the supply of a non-permutation Γ_s is proved on paper. For the equivalence, a memory witness below L gives a non-permutation Γ_s; conversely, with none the law on $\{0, \dots, L\}$ is Markov, $\Gamma_L = I$ makes every Γ_s a permutation by the stochastic-inverse lemma, and $\Gamma_s^{-1}\Gamma_t$ is a stochastic divisor, full support being needed only in this converse. Indivisibility at some return horizon is also machine-checked from the *routed* form of (C4), the realization clause in which a record of the visible history written into the hidden state is routed back into later visible conditionals within the window [Main §1.3].

Recurrence-cycle indivisibility thus follows from finite reversibility and partial access alone, Markov laws included, and does not distinguish one law from another; what an observer can discriminate is memory at an accessible order, which (C4) asserts [Main §2.3, Scope].

2.6 Three examples

A minimal model. Let $\mathcal{C}_V = \{0, 1\}$ and $\mathcal{C}_H = \{1, \dots, 6\}$ [Main §2.4]. For $h \in \{1, 2\}$, φ swaps $(0, h)$ with $(1, h)$: it flips the visible bit and keeps h, which then records the flip. At fixed x, φ swaps $h = 3$ with 4, and 5 with 6. A second application reads the record and flips the bit back. With h uniform,

$$\Gamma_1 = \begin{pmatrix} 2/3 & 1/3 \\ 1/3 & 2/3 \end{pmatrix}, \qquad \Gamma_2 = I, \qquad \Gamma_1^{-1}\Gamma_2 = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}.$$

The only candidate divisor has negative entries, so the family is P-indivisible, whereas a Markov chain with the same one-step matrix would give Γ_1^2, with diagonal entries 5/9. With x_0 uniform and independent of h, $P(x_2{=}0 \mid x_1{=}1, x_0{=}0) = 1$ while $P(x_2{=}0 \mid x_1{=}1) = 1/3$.

Memory without short-window indivisibility. Let X_0 and X_1 be independent uniform bits and $X_2 = X_0 \oplus X_1$. Every two-time transition matrix is $\frac{1}{2}J$, with J the all-ones matrix, so the family factors through $\Delta = I$ and is P-divisible on $\{0, 1, 2\}$. Yet $I(X_0; X_2 \mid X_1) = 1$ bit. Up to flipping the final bit, this is the process of [RHP14, Eq. (15)]; it appears concurrently in [Dou26, App. A].

The card table. The two-card game of [Book §1.11] uses two cards, each red (0) or black (1): a visible *hole card* x, and a face-down *burn card* b, uniform and independent of x. One shuffle step swaps the two cards, $\varphi(x, b) = (b, x)$. A *peek* reveals the hole card and overwrites the burn card, which holds the record, with a fresh card drawn uniformly and independently. Exact enumeration of the equally likely cases gives the probability that the hole card at the final reveal is the hole card as dealt:

Protocol	Probability
one shuffle	1/2
two shuffles	1

Protocol	Probability
two shuffles, with a peek between them	1/2
Markov prediction for two shuffles, Γ_1^2	1/2

Here $\Gamma_1 = \frac{1}{2}J$ and $\Gamma_2 = I$. The two rows of Γ_1 are equal, so for every Δ the two rows of $\Gamma_1\Delta$ are equal and $\Gamma_1\Delta \neq I$: the family is P-indivisible.

The table separates the deterministic information from what the player can access. The shuffle is a bijection and loses nothing; the player sees only the hole card, and the memory the player observes, certainty after two shuffles where a Markov model predicts 1/2, is carried by the face-down card. The peek row depends on the rule: if the peek only reveals or replaces the hole card, the probability is 1, so the 1/2 comes from overwriting the record, not from looking. With anticorrelated cards ($b = 1 - x$), Γ_1 is a permutation and the game is divisible.

On card inputs the game is a classical stochastic process, and no card arrangement violates a Bell inequality [Book §1.11; Main §3.3].

3. Quantum emergence: scope and open steps

3.1 Three notions kept apart

The indivisibility of §2.5 is global: it occurs somewhere in the recurrence cycle, not on every window, and conditional memory does not force short-window indivisibility (§2.6). A revival of distinguishability within a window, two roots whose rooted laws are closer at an earlier time than at a later one, does make the family P-indivisible on that window [Main §2.3].

Three notions are distinct, and no unconditional implication holds among them [Main §3.2, Remark (three notions, kept apart)]: (1) accessible memory, $M_t > 0$; (2) stochastic P-indivisibility of the visible population law; (3) quantum non-Markovianity under a chosen criterion. A family of quantum channels Φ_t is completely-positive (CP) divisible if $\Phi_t = \Psi_{t,s} \circ \Phi_s$ with each $\Psi_{t,s}$ completely positive and trace-preserving. For the family obtained by tracing out a permutation unitary from a diagonal hidden state, (2) implies CP-indivisibility [Main §3.2, Theorem (CP-indivisibility, framework scope)]. Outside the diagonal-preserving class the implication fails: with Φ_{t_1} conjugation by the Hadamard gate and $\Phi_{t_2} = \mathrm{id}$ the family is CP-divisible, while its computational-basis populations, $\frac{1}{2}J \to I$, are P-indivisible [Main §3.2, Remark (scope: P-indivisibility is not quantum non-Markovianity)]. In the card table of §2.6, embedded as a permutation unitary with the burn card in the uniform diagonal state, the one-step channel $\rho \mapsto \mathrm{tr}(\rho)\, I/2$ is entanglement-breaking (measure-and-prepare) and the two-step map is the identity, so the family is CP-indivisible [Main §3.2]; its memory is still classical in the sense of [BBS24], since the two-step identity is obtained by realizing the one-step channel as the Pauli twirl $\rho \mapsto \frac{1}{4}\sigma_o \rho \sigma_o$,

$o \in \{0, x, y, z\}$, and applying σ_o conditioned on o, a decomposition a fixed initial hidden card does not admit [Main §4.5]. Distinguishability backflow witnesses divisibility failure in one direction only [BLP09]; for bijective dynamical maps on a d-dimensional system, CP-divisibility is equivalent to monotone distinguishability with a $(d+1)$-dimensional ancilla [BJA17]. Theorem 1 states which laws admit realizations and quantum representations, not why a physical system exhibits a law with memory [Main §3.4, Remark (representation, not explanation)]; Calvo states a derivation of the textbook axioms from indivisible stochastic evolution [Cal26], whereas Theorem 1 derives no axioms.

3.2 From the correspondence to operational quantum mechanics

Operational quantum mechanics requires, beyond the fixed-basis records of Theorem 1, coherent preparations, noncommuting interventions and composites represented by one instrument theory.

Fix a nonempty finite system and an operational theory specifying which finite outcome families of linear maps are available on the system and at every ancilla level, that is, with each finite ancilla adjoined. The theory contains the identity and is closed under coarse-graining of outcomes and, at every ancilla level, under outcome-conditioned composition. It has an available preparation of a uniformly mixed ancilla, which an available operation may follow, an available spectator-independent readout of the ancilla, one that acts on every product input as the identity on the system, and ancilla discard: an available preparation followed by an available instrument, with the ancilla then discarded, is an available instrument on the system with the same outcomes [GR §3.3]. The five completion conditions are:

(i) available families send positive states to positive states, and their outcome traces sum to the input trace;
(ii) operations remain available with a one-state ancilla adjoined;
(iii) an untouched independent system can be adjoined without changing an intervention;
(iv) every finite unitary on every composite is available;
(v) attaching a fresh uniformly prepared ancilla to a composite, acting, and discarding it gives an available operation.

Theorem 4 (operational-completion characterization [GR §3.3]). *For every nonempty finite system, the available outcome families on the system and at every positive ancilla level are exactly the normalized finite Kraus instruments if and only if conditions (i)–(v) hold.*

Both directions are machine-checked (Appendix A). The sufficiency direction rests on the Stinespring dilation and on the realization of every finite instrument by a unitary interaction with an ancilla followed by ancilla readout and discard [Sti55; Kra83; Oza84]; for reconstructions of finite quantum theory from operational axioms see [Har01; CDP11; MM11].

Each condition is independent of the other four and of the observation process itself: for one fixed finite embedded-observation process satisfying (C1)–(C4), and for each condition, there is a qubit theory that realizes that process and the other four conditions, and fails that condition. In particular, a qubit theory that never creates coherence realizes the observation process and is not quantum, so OI alone does not imply quantum mechanics (both machine-checked) [GR §3.3]. Which of (iii)–(v) a physical substratum supplies is the open step toward full operational equivalence.

3.3 The evolution relating two times is not selected

Representability does not fix the unitary evolution relating two times [Main §3.4, Remark (scope: representability, not selection of the relating evolution)]. Two coherent lifts of one visible family, unitary dilations that reproduce the visible law at every time and compose exactly across times, can differ by a time-dependent right action invisible to the visible law and then have different relating evolutions (machine-checked on an exhibited pair of lifts), while a constant deformation cancels from every relating evolution (machine-checked in general).

For lifts with trivial ancilla, read in a fixed basis, two lifts have equal rooted families exactly when their one-step visible matrices are equal, and equal one-step matrices give equal trajectory laws at every horizon (machine-checked, both directions). The fixed-basis correspondence therefore acts on the quotient by visible equality, and selecting the relating evolution needs a condition beyond representability and ordinary coherence.

3.4 No fixed finite realization reproduces a qubit

A fixed finite realization with N ontic configurations generates only behaviour matrices that factor nonnegatively through N states. Simulating a qubit's prepare-and-measure statistics needs classical memory unbounded over finite experiments [GH03; Mon08], and Montina reports Hardy's result that the number of ontic states cannot be finite [Har04]. So no fixed finite realization reproduces the full prepare-and-measure theory of a qubit exactly; the factorization is proved, and the unboundedness is imported [Main §3.4, Proposition (finite-substratum operational obstruction — classical-dimension form); HKLP24]. The obstruction does not touch Theorem 1, whose realization is chosen per law.

3.5 Bell structure

[Main §3.3] proves a Bell ceiling under the framework's own hypotheses. The spatial substratum is a product of finite site factors, and its coupling graph joins site i to site j when the ith output component of φ depends on the jth input component, and d_G is its graph distance; an intervention on a region A changes, k updates later, only components within d_G-distance k of A [Main §3.3]. Let Ω be a pre-setting variable that fixes each wing's response to every setting, and

suppose that (i) the dynamics is deterministic; (ii) the setting operation acts at one step on disjoint regions A and B, each intervention reading and writing only its own region; (iii) wing A's outcome is a ±1-valued function of the state on a readout region R_A, k_A updates after the setting, with $d_G(B, R_A) > k_A$, and symmetrically $d_G(A, R_B) > k_B$; and (iv) the interventions act on one common pre-setting ensemble, $p(\Omega \mid a, b) = p(\Omega)$. If the protocol records a ±1 outcome in every run, the model is Bell-local, with deterministic local response functions, and the Clauser–Horne–Shimony–Holt combination obeys $|S_{\mathrm{CHSH}}| \leq 2$ [Bel64; CHSH69; Main §3.3, Theorem (Bell ceiling for a screened local completion)]. Reproducing the quantum values $2 < |S_{\mathrm{CHSH}}| \leq 2\sqrt{2}$ therefore requires violating at least one of these assumptions.

In the framework, determinism is part of OI and the protocol records an outcome in every run, so a completion reproducing quantum Bell correlations gives up at least one of the remaining assumptions: (ii) and (iii), which with the causal-cone property secure ontic parameter independence, or (iv), measurement independence [Main §3.3]. The framework retains measurement independence, as a posit about the initial condition, and adopts ontic parameter dependence: one wing's microstate response may depend on the remote setting, which the nearest-neighbour cubic update of [SM] alone cannot supply for wings beyond its graph cone [Main §3.3, Corollary (Bell–lattice obstruction and state-dependent escape)], while operational no-signalling is required. That a preparation-indexed coupling graph can supply it while preserving operational no-signalling and the gravitational continuum limit is the open hypothesis H-Bell [Main §3.3].

4. The cosmological constant: how OI changes the interpretation of vacuum energy

Summing zero-point energies of the quantum fields up to a Planck-scale cutoff gives a vacuum energy density some 120 orders of magnitude above the observational value, the exact power depending on the cutoff convention [Wei89, §I; §III, Eqs. (3.4)–(3.6)]. If that density sources curvature, it must be cancelled to that accuracy [Wei89, §III]; this discrepancy is here called the mismatch.

What OI establishes. In OI the zero-point sum belongs to the emergent quantum description (§§2–3), while the effective cosmological constant Λ belongs to the gravitational description of the substratum, with its classical stress-energy $T^{\mathrm{cl}}_{\mu\nu}$: the two are defined at different levels. A uniform shift $\hat{H} \to \hat{H} + C$ of the emergent Hamiltonian changes no visible transition probability at any precision, so the zero-point offset is absent from every visible statistic of the quantum description [GR §6.2]. That no non-gravitational measurement sees the absolute vacuum energy is familiar; what OI adds is the second level, at which a gravitational source can be defined independently of the emergent vacuum.

The proposed resolution. If the covariant field equations take the form [GR] targets,

$$\mathcal{G}_{\mu\nu} = 8\pi G\,(T^{\mathrm{cl}}_{\mu\nu} - \mathcal{F}[g, T^{\mathrm{cl}}]\,g_{\mu\nu}),$$

with $\mathcal{G}_{\mu\nu}$ the gravitational side, still to be derived (the Einstein tensor in general relativity), and $\mathcal{F}$ a scalar functional of the metric and the stress-energy, then the zero-point sum of the emergent description is not a source of curvature and the mismatch requires no cancellation. The vacuum term that remains, from the substratum's classical energy density and from $\mathcal{F}$, is not computed, and the observed value of Λ enters as a cosmological input.

The missing link. The quantum description alone cannot decide whether the offset sources curvature. The completions

$$\mathcal{G}_{\mu\nu} = 8\pi G\, T^{\mathrm{cl}}_{\mu\nu} \qquad \text{and} \qquad \mathcal{G}_{\mu\nu} = 8\pi G\,(T^{\mathrm{cl}}_{\mu\nu} - \rho_C\, g_{\mu\nu}),$$

with ρ_C the energy density of the offset C, share the substratum, the partition and every visible transition law [GR §6.2]. Deciding between them requires covariant field equations derived from the substratum, with their conservation and coupling to sources, which [GR] calls the G3 coupling theorem and leaves open [GR §6.2, §8.5]; the same step fixes how the energy of electroweak and QCD vacuum transitions gravitates. A derivation of these equations would turn the proposal into a theorem or refute it.

Trace-free Einstein equations [EvEMU11] and vacuum-energy sequestering [KP14] also remove matter vacuum energy as a source of curvature, and Volovik argues that the ultraviolet zero-point sum does not set the vacuum energy [Vol06]; OI locates the gravitational source in the substratum microstate.

5. Black-hole information: preservation versus accessible recovery

The black-hole information problem asks whether the radiation of an evaporating black hole carries the information of what formed it [Haw76; Pag93; Pag13]. In OI two questions separate: whether information is conserved by the global dynamics, which holds because the dynamics is a bijection, and whether it can be recovered from the records an observer can access [HP07]. Finite exact computations show that these are different properties.

The toy model [BHa]. The configuration space is a finite product of one-bit registers. A message m, drawn uniformly from a set $\mathcal{M}$, is written into designated interior registers; every other register is uniformly random and independent of m, or blank. One map is applied at every step, and at each *emission* one register passes from the interior to the radiation. The observer reads registers on a schedule fixed in advance, in the counts below the emitted radiation registers, and keeps every read; the measured quantity is P_{succ}, the probability that the best guess of m from these records is correct, and the *accessible advantage* is its excess over the blind value $1/|\mathcal{M}|$. $\mathcal{U}$ is the set of joint values of the registers left unread at the end. The layout is that of the Page-curve setup [GR App. A.7] at small size.

Results [BHa]. Conservation does not imply recovery: of the 40,320 bijections of $\{0,1\}^3$ with one message bit and two random bits, 1,376 give zero accessible

advantage with one register unread after two emissions. Nor is it necessary: 60 non-injective maps of $\{0,1\}^2$ recover the message exactly when every register is read. What bijectivity guarantees is a floor: reading every register at the end gives exact recovery, and with $\mathcal{U}$ unread $P_{\text{succ}} \geq 1/|\mathcal{U}|$, so zero advantage requires $|\mathcal{U}| \geq |\mathcal{M}|$; 32 non-injective maps of $\{0,1\}^2$ violate this floor. It is the remnant-capacity form of the fact that $|\mathcal{U}|$-valued side information multiplies a guessing probability by at most $|\mathcal{U}|$, so hiding the message completely needs at least as many hidden values as messages, as in perfect secret sharing, of which Shamir's threshold scheme is the standard example [Sha79]. The card table of §2.6 shows the same separation in its simplest form.

The Page curve. The Page-curve calculation of [GR App. A] addresses recovery in the quantum description, on $\mathcal{C}_V \times \mathcal{C}_B \times \mathcal{C}_D$, with V the exterior, B the black-hole interior, D the sector beyond the cosmological horizon, and the radiation $R \subset V$, with configurations $\mathcal{C}_R$, emitted from B [GR App. A.1, A.7]. It rests on three conditions: *H-scramble*, that the orbit ensemble equidistributes over quantum pure states on the energy shell, which fails for every bijection acting on a definite configuration with the bipartition aligned to the configuration coordinates [BHb]; *H-decouple*, that $B \cup R$ evolves uncoupled from D and from $V \setminus R$ in a pure joint state over whole cycles, which the effective-pure-state theorem [GR, Thm A.7] supplies only for times short compared with the D-sector timescale $\tau_B^D \sim 1/H$ [GR App. A.5], far shorter than the evaporation [BHa]; and constant energy on all of $\mathcal{C}_B \times \mathcal{C}_R$. Under them, for all but an exponentially small fraction of initial conditions, the full-cycle average of the radiation's entanglement entropy at each stage's bipartition equals the Page value $S_{\text{Page}}(d_B, d_R)$ up to the correction term of Theorem A.8, by measure concentration [Pag93; PSW06; HLW06; GR, Thms A.8–A.9].

6. Established results, conditional interpretations and open problems

Established mathematical results. Machine-checked: Theorem 1; the cores of Theorem 2 and of the capacity bound; Theorem 3 in its rooted form, which with the elementary fact that a non-permutation stochastic matrix has two rows sharing a positive column gives recurrence-cycle P-indivisibility of the rooted visible family, with a row for every root including roots of visible probability zero and a hidden prior shared by every root and drawn independently of it, whenever some rooted transition matrix is not a permutation; Theorem 4 and the independence of its conditions; the non-selection of the relating evolution. Proved on paper: the time-t form of Theorem 2, its Corollary and the least hidden predictive memory; that every non-deterministic visible step and every memory witness supply a non-permutation rooted transition matrix, and that with a full-support visible prior the indivisibility is equivalent to $M_k > 0$ at some order k below the recurrence time (§2.5); the Bell ceiling. The qubit obstruction combines a proved factorization with an imported unboundedness, and the separation of conservation from recovery rests on exhaustive enumeration with the

$1/|\mathcal{U}|$ floor proved. Together these results establish that stochastic, quantum-like observations can arise exactly from deterministic reversible dynamics, and they locate what is not universal: the memory a law carries, its size and position in the hidden sector, and its consequence for divisibility. The correspondence represents these observations; it does not derive their physical mechanism.

Conditional physical interpretations. If OI holds as physics (with the bijective substratum an empirically motivated representation choice, §1.1), the fixed-basis statistics of quantum theory are the visible laws of an embedded observer, and their departures from Markov behaviour are memory carried in an inaccessible sector, as the card table of §2.6 illustrates. With measurement independence retained, quantum Bell correlations then require ontic parameter dependence, subject to H-Bell (§3.5); the zero-point sum is not a gravitational source if the field equations take the targeted form (§4); and the framework's cycle-averaged Page curve holds under H-scramble, H-decouple and with all of $\mathcal{C}_B \times \mathcal{C}_R$ one energy shell (§5).

Open problems. For quantum emergence:

1. full operational equivalence: which of completion conditions (iii)–(v) a physical substratum supplies (§3.2);
2. a selection principle for the quadratic Born exponent (§2.3) and for the evolution relating two times (§3.3);
3. history readback (C4) at the cosmological partition [GR §2.2], and an experiment or memory-witness test distinguishing OI from standard quantum theory [Main §4.5];
4. the dynamical construction problems: exact gluing of per-horizon realizations for recurrent target families, approximation of a target on windows short compared with the recurrence time by one fixed substratum [Main §3.4], and the hidden-sector mixing hypothesis, motivated by eigenstate thermalization, under which observed conditional memory gives a lower bound on the hidden sector's coarse-grained mixing time, the slow-bath form of (C2) [Main §1.3, §3.4];
5. machine checks of the steps of §§2.2–2.5 proved on paper.

For the applications: a derivation of the covariant field equations with their source coupling, on which both §4 and, in [GR], the Page curve depend; H-scramble and H-decouple; the proper-shell case, which an evaporating black hole whose radiation carries away its energy occupies; the entropy along a single trajectory [BHb]; the generalized second law [GR App. A.6]; and the relation to quantum-extremal-surface and island computations [Pen20; AEMM19; AMMZ20; Book §7.6] (§5).

Code, data and formal verification

The declarations in Appendix A were checked in Lean 4.33.0 with Mathlib v4.33.0 at commit `17b830bc` (17b830bcd0ecc36e10eff5b9eeca03bdd7198d86) by building `verification/lean-mathlib` (`lake exe cache get`, `lake`

`--rehash build`) and running `python3 tools/release_gate.py`, whose `lean-axioms` step requires every `#print axioms` report to lie within `propext`, `Classical.choice` and `Quot.sound`; each declaration reports exactly these three, and the expected result and comparison recipe are in `papers/introduction_repro/`. The Lean pin is the commit at which that result was retained; the manuscripts are cited at corpus commit `55e3217b`, whose `verification/lean-mathlib` tree is identical.

The probes and the black-hole computations [BHa; BHb] are released under the MIT License, at `papers/oi_lattice_code/foundations/` (probes; also in the release archive, concept DOI 10.5281/zenodo.19060318) and `verification/programmes/black-hole-information/` (not part of the release archive) at the corpus commit.

Appendix A. Machine-checked declarations

Modules of the `OIBridge` Lean library; line numbers at commit `17b830bc`.

- **Theorem 1** (§2.2): `finite_horizon_equivalence` (Equivalence:462); directions `S_imp_D` (:422), `D_imp_Qfb` (:294), `Qfb_imp_S` (:245), compositions :466–468; supporting `permMatrix_mem_unitaryGroup` (EquivalenceChain:179).
- **Theorem 2** (§2.4): `pushforward_identity`, `distinguishability_floor`, `capacity_floor`, `unavoidable_hidden_predictive_memory` (HiddenMemory:83, :114, :170, :185), for a joint law of history, present and hidden state.
- **Capacity bound and least hidden memory** (§2.4): `c3_necessity`, `c3_necessity_via_hidden`, `card_hidden_ge_two_pow_Istar` (C3Necessity:80, :88, :117), joint-law form.
- **Stochastic-inverse lemma; Theorem 3** (§2.5): `isPermMatrix_of_stochastic_inverse` (EquivalenceChain:110); `pIndivisible_of_rooted_identity_return_of_overlap` (RecurrenceHorizon:148).
- **Recurrence-cycle indivisibility** (§2.5): `routed_forces_indivisible_somewhere` (PhysicalC4Discharge:122), indivisibility at some return horizon from routed readback; on paper for history-level (C4).
- **Relating evolution** (§3.3): `gl2_strong_gauge_moves_relative_candidate` (CoherentLiftGauge:537, an exhibited pair of lifts), `gl3_constant_gauge_preserves_relative` (:451, general); `bridge_traj`, `rooted_eq_iff_slice_eq` (both directions: rooted families equal exactly when one-step matrices are), `realData_traj_stochastic` (TrackBQfbBridge:276, :180, :286).
- **Theorem 4** (§3.2): `exactAll_iff_physical_general` (GeneralCarrier:100); ⇒ `physical_of_exactAll` (PhysicalCharacterization:301); ⇐ `exactAll_of_physical_general` (GeneralCarrier:93) via `krausSoundExt_of_validity_inert` (OperationalValidity:124), `compositeCompleteness_unconditional` (IsometryExtension:179, premise discharged by `finiteIsometryExtensionSF_discharged`, :116) and `exactAll_of_levelOne` (LevelOneSeam:196).
- **Independence; OI alone does not imply quantum mechanics** (§3.2, qubit): `five_way_minimality` (RankGapTheory:702), `oi_alone_not_qm` (GeneralCarrier:160).

References

Corpus (repository commit `55e3217b`, concept DOI 10.5281/zenodo.19060318)
[Main] A. Maybaum, "The Incompleteness of Observation: The Equivalence of Quantum Mechanics and Embedded Observation," `papers/Main.md`.

[GR] A. Maybaum, "$\hbar$, the Bekenstein-Hawking Entropy, and the Running-Vacuum Form of Dark Energy from the Cosmological Horizon," `papers/GR.md`.
[Book] A. Maybaum, *The Incompleteness of Observation*, `book/`.
[BHa] A. Maybaum, "Recovery from accessible records in finite reversible models," unpublished research record; proofs and code at `bh01_access_v1/` in the black-hole computations' directory.
[BHb] A. Maybaum, "Permutation dynamics versus quantum scrambling; Page-time fraction," unpublished research record; proofs and code at `bh02_scramble_v1/` in the black-hole computations' directory.

External

[AEMM19] A. Almheiri, N. Engelhardt, D. Marolf, and H. Maxfield, "The entropy of bulk quantum fields and the entanglement wedge of an evaporating black hole," *J. High Energy Phys.* **12**, 063 (2019); arXiv:1905.08762.
[AMMZ20] A. Almheiri, R. Mahajan, J. Maldacena, and Y. Zhao, "The Page curve of Hawking radiation from semiclassical geometry," *J. High Energy Phys.* **03**, 149 (2020); arXiv:1908.10996.
[Bar23] J. A. Barandes, "The Stochastic-Quantum Theorem," arXiv:2309.03085v2 (2023).
[Bar25] J. A. Barandes, "The Stochastic-Quantum Correspondence," *Philosophy of Physics* **3**, 8 (2025); arXiv:2302.10778v3.
[Bar26] J. A. Barandes, "The Born Representation Theorem and the Unistochastic Theorem," arXiv:2608.04354v1 (2026).
[BB12] J. C. Baez and J. Biamonte, "Quantum techniques for stochastic mechanics," arXiv:1209.3632v5 (2012).
[BBS24] C. Bäcker, K. Beyer, and W. T. Strunz, "Local disclosure of quantum memory in non-Markovian dynamics," *Phys. Rev. Lett.* **132**, 060402 (2024); arXiv:2310.01205v2.
[BD16] F. Buscemi and N. Datta, "Equivalence between divisibility and monotonic decrease of information in classical and quantum stochastic processes," *Phys. Rev. A* **93**, 012101 (2016); arXiv:1408.7062v3.
[Bel64] J. S. Bell, "On the Einstein Podolsky Rosen paradox," *Physics Physique Fizika* **1**, 195 (1964).
[BJA17] B. Bylicka, M. Johansson, and A. Acín, "Constructive method for detecting the information backflow of non-Markovian dynamics," *Phys. Rev. Lett.* **118**, 120501 (2017); arXiv:1603.04288v2.
[BLP09] H.-P. Breuer, E.-M. Laine, and J. Piilo, "Measure for the degree of non-Markovian behavior of quantum processes in open systems," *Phys. Rev. Lett.* **103**, 210401 (2009); arXiv:0908.0238v2.
[Cal26] S. Calvo, "On the stochastic-quantum correspondence," *Quantum Stud. Math. Found.* **14**, 2 (2026), doi:10.1007/s40509-026-00415-9; arXiv:2601.18720v2.
[CDP11] G. Chiribella, G. M. D'Ariano, and P. Perinotti, "Informational derivation of quantum theory," *Phys. Rev. A* **84**, 012311 (2011).
[CHSH69] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, "Proposed experiment to test local hidden-variable theories," *Phys. Rev. Lett.* **23**, 880 (1969).
[CT06] T. M. Cover and J. A. Thomas, *Elements of Information Theory*, 2nd ed. (Wiley, 2006).
[dMU21] L. de Moura and S. Ullrich, "The Lean 4 theorem prover and programming language," in *Proc. CADE 28* (Springer, 2021), p. 625.
[Dou26] J. Doukas, "On the emergence of quantum mechanics from stochastic processes," arXiv:2602.22095v3 (2026).
[EGGH26] G. Egri, M. Gömöri, B. Gyenis, and G. Hofer-Szabó, "Trajectory of probabilities, probability on trajectories, and the stochastic-quantum correspondence," arXiv:2602.23491v1 (2026).
[EvEMU11] G. F. R. Ellis, H. van Elst, J. Murugan, and J.-P. Uzan, "On the trace-free Einstein equations as a viable alternative to general relativity," *Class. Quantum Grav.* **28**, 225007 (2011); arXiv:1008.1196v2.
[GH03] E. F. Galvão and L. Hardy, "Substituting a qubit for an arbitrarily large number of

classical bits," *Phys. Rev. Lett.* **90**, 087902 (2003); arXiv:quant-ph/0110166v3.

[Har01] L. Hardy, "Quantum theory from five reasonable axioms," arXiv:quant-ph/0101012 (2001).

[Har04] L. Hardy, "Quantum ontological excess baggage," *Stud. Hist. Philos. Mod. Phys.* **35**, 267 (2004).

[Haw76] S. W. Hawking, "Breakdown of predictability in gravitational collapse," *Phys. Rev. D* **14**, 2460 (1976).

[HKLP24] T. Heinosaari, O. Kerppo, L. Leppäjärvi, and M. Plávala, "Simple information processing tasks with unbounded quantum advantage," *Phys. Rev. A* **109**, 032627 (2024); arXiv:2308.07727v4.

[HLW06] P. Hayden, D. W. Leung, and A. Winter, "Aspects of generic entanglement," *Commun. Math. Phys.* **265**, 95 (2006); arXiv:quant-ph/0407049.

[HP07] P. Hayden and J. Preskill, "Black holes as mirrors: quantum information in random subsystems," *J. High Energy Phys.* **09**, 120 (2007); arXiv:0708.4025.

[KP14] N. Kaloper and A. Padilla, "Sequestering the Standard Model vacuum energy," *Phys. Rev. Lett.* **112**, 091304 (2014); arXiv:1309.6562v2.

[Kra83] K. Kraus, *States, Effects, and Operations*, Lecture Notes in Physics **190** (Springer, 1983).

[LP17] D. A. Levin and Y. Peres, with contributions by E. L. Wilmer, *Markov Chains and Mixing Times*, 2nd ed. (American Mathematical Society, 2017).

[Mathlib20] The mathlib Community, "The Lean mathematical library," in *Proc. ACM SIGPLAN Conf. Certified Programs and Proofs* (ACM, 2020), p. 367.

[Milz20] S. Milz, D. Egloff, P. Taranto, T. Theurer, M. B. Plenio, A. Smirne, and S. F. Huelga, "When is a non-Markovian quantum process classical?," *Phys. Rev. X* **10**, 041049 (2020); arXiv:1907.05807v3.

[MM11] Ll. Masanes and M. P. Müller, "A derivation of quantum theory from physical requirements," *New J. Phys.* **13**, 063001 (2011).

[Mon08] A. Montina, "Exponential complexity and ontological theories of quantum mechanics," *Phys. Rev. A* **77**, 022104 (2008); arXiv:0711.4770v2.

[Oza84] M. Ozawa, "Quantum measuring processes of continuous observables," *J. Math. Phys.* **25**, 79 (1984).

[Pag93] D. N. Page, "Average entropy of a subsystem," *Phys. Rev. Lett.* **71**, 1291 (1993); arXiv:gr-qc/9305007v2.

[Pag13] D. N. Page, "Time dependence of Hawking radiation entropy," *J. Cosmol. Astropart. Phys.* **09**, 028 (2013); arXiv:1301.4995v3.

[Pen20] G. Penington, "Entanglement wedge reconstruction and the information paradox," *J. High Energy Phys.* **09**, 002 (2020); arXiv:1905.08255v3.

[PSW06] S. Popescu, A. J. Short, and A. Winter, "Entanglement and the foundations of statistical mechanics," *Nature Phys.* **2**, 754 (2006).

[RHP14] Á. Rivas, S. F. Huelga, and M. B. Plenio, "Quantum non-Markovianity: characterization, quantification and detection," *Rep. Prog. Phys.* **77**, 094001 (2014); arXiv:1405.0303v2.

[Sch21] H.-J. Schmidt, "Dilation of stochastic matrices by coarse graining," arXiv:2106.03513v1 (2021).

[SGD19] P. Strasberg and M. García Díaz, "Classical quantum stochastic processes," *Phys. Rev. A* **100**, 022120 (2019); arXiv:1905.03018v2.

[Sha79] A. Shamir, "How to share a secret," *Commun. ACM* **22**, 612 (1979).

[Sti55] W. F. Stinespring, "Positive functions on C-*algebras,"* Proc. Amer. Math. Soc.* **6**, 211 (1955).

[tH14] G. 't Hooft, "The cellular automaton interpretation of quantum mechanics," arXiv:1405.1548v3 (2014).

[Vol06] G. E. Volovik, "Vacuum energy: myths and reality," *Int. J. Mod. Phys. D* **15**, 1987 (2006); arXiv:gr-qc/0604062v4.

[Wei89] S. Weinberg, "The cosmological constant problem," *Rev. Mod. Phys.* **61**, 1 (1989).