THE INCOMPLETENESS OF OBSERVATION

Why Quantum Mechanics and General Relativity Cannot Be Unified From Within

Author: Alex Maybaum
Date: March 2026
Status: DRAFT PRE-PRINT
Classification: Theoretical Physics / Foundations

ABSTRACT

An observer embedded in a deterministic universe cannot access the complete state. We prove that any such observer — coupled to a slow, high-capacity hidden sector on a finite-dimensional configuration space — must describe the visible sector using P-indivisible stochastic dynamics, mathematically equivalent to unitary quantum mechanics. The converse also holds: any quantum system, realized as a deterministic dilation, requires non-trivial coupling, slow-bath memory, and sufficient hidden-sector capacity. The Schrödinger equation, Born rule, and Bell violations emerge as structural consequences requiring no independent quantum postulates.

Applied to the cosmological horizon, the framework uniquely determines $\hbar = c^3\epsilon^2/(4G)$, fixes $\epsilon = 2\,l_p$, and recovers the Bekenstein-Hawking entropy $S = A/(4\,l_p^2)$. The 10^{122} cosmological constant discrepancy is identified as S_{dS}, the information compression ratio of the emergent quantum description; the observed vacuum energy is the mandatory classical baseline. Falsifiable predictions include dark energy evolution in Running Vacuum Model form with $\nu_{\mathrm{OI}} \approx 2.45 \times 10^{-3}$ (parametric; systematic range 1–5 $\times\, 10^{-3}$), consistent with current DESI data, and gravitational wave echoes near black hole horizons. The trace-out that produces QM simultaneously renders ~95% of the universe's gravitational budget invisible to the emergent description — matching the observed dark sector — providing independent corroboration of observational incompleteness.

PART I: THE GENERAL THEOREM

1. INTRODUCTION

1.1 The Problem of Embedded Observation

An observer embedded in a deterministic universe cannot access the complete state: degrees of freedom beyond causal reach are permanently hidden, and the observer's description is obtained by marginalizing over the hidden sector. The question is what this reduction imposes on the form of the observer's physical laws.

Prior work (QBism, relational QM, 't Hooft's cellular automaton [1]) takes observer-dependence as an interpretive starting point or derives it from specific microphysical models. This framework differs by identifying *necessary and sufficient conditions* under which any embedded observer in any deterministic system necessarily sees quantum mechanics — and proving these are the *only* conditions under which QM arises from a deterministic embedding. These constraints carry empirical weight: the framework reveals that phenomena previously viewed as catastrophic anomalies

— most notably the 10^{122} cosmological constant discrepancy — are mandatory consequences of embedded observation.

1.2 Axioms

The framework rests on four axioms referencing no quantum mechanics, general relativity, or specific physical theory.

1. **Deterministic dynamics.** The total system evolves deterministically on a phase space Γ via a Hamiltonian flow ϕ_t. A probability density ρ on Γ (representing the observer's incomplete knowledge) evolves by the Liouville equation:

$$\frac{\partial \rho}{\partial t} = \{H_{\text{tot}}, \rho\} \equiv \mathcal{L}\rho$$

2. **Finiteness.** The effective configuration space of the visible sector is finite-dimensional, with a discreteness scale ϵ providing a finite minimal cell volume. Any observer bounded by a finite-area surface can couple to only finitely many modes across that boundary; independent support comes from holographic entropy bounds [27]. In Part I, ϵ is free; self-consistency (§6) pins $\epsilon = 2\,l_p$. The boundary-only dependence lemma (§5.2) shows that the cardinality of the deep hidden sector — beyond the boundary layer — is gauge: no observable depends on it.

3. **Causal partition.** Γ is partitioned into a visible sector Γ_V and hidden sector Γ_H:

$$\Gamma = \Gamma_V \times \Gamma_H, \qquad H_{\text{tot}} = H_V + H_H + H_{\text{int}}$$

Cross-partition correlations enter only through H_{int}. The product decomposition is idealized; §4 and §9.6 address approximation quality and the conditions under which the sharp partition is exact.

4. **Classical probability.** The observer's predictions are classical expectation values over phase space; standard probability theory applies.

These axioms contain no quantum postulates. The claim is that quantum mechanics *emerges* under the conditions below.

1.3 Conditions on the Hidden Sector

(C1) Non-zero coupling. $H_{\text{int}} \neq 0$. The coupling is bidirectional.

(C2) Slow-bath timescale separation. The visible-sector (system) timescale τ_S is much shorter than the hidden-sector (bath) timescale τ_B: $\tau_S \ll \tau_B$ — the *inverse* of the Markovian regime. The hidden sector evolves on timescales far exceeding those accessible to the observer.

(C3) Sufficient capacity. The number of hidden-sector degrees of freedom N_H exceeds the number of visible-sector degrees of freedom N_V by enough that visible-sector interactions do not appreciably perturb the hidden sector's state on timescales $\ll \tau_B$.

Theorem statement. Under Axioms 1–4 and (C1)–(C3), the embedded observer's reduced description is mathematically equivalent to unitarily evolving quantum mechanics (defined precisely in §3.3). The conditions are not merely sufficient but *necessary* (§3.3), establishing full equivalence.

1.4 Partition-Relativity

Lemma. *The emergent description is uniquely determined by the partition. Any parameters of the emergent theory depend only on the geometric and thermodynamic properties of the partition boundary.*

Proof. By Axiom 1, the Hamiltonian flow ϕ_t is unique. By Axiom 3, the partition is fixed. By Axiom 4, the observer uses the Liouville measure μ — the unique absolutely continuous invariant measure on the full phase space. (An embedded observer with incomplete knowledge of the total energy averages over energies with a smooth prior, recovering Liouville on a phase-space band; singular invariant measures are excluded by Axiom 4.) Letting π_V denote projection onto the visible sector, the marginalized transition probabilities

$$T_{ij}(t_2, t_1) = \int_{\Gamma_H} \delta_{x_j}[\pi_V(\phi_{t_2-t_1}(x_i, h))]\, d\mu(h)$$

are therefore uniquely determined by three inputs: dynamics (ϕ_t), partition (Γ_V, Γ_H), and measure (μ). Since ϕ_t and μ are fixed by the axioms, the stochastic process — and hence any emergent quantum description (§3.1) — depends only on the partition. □

1.5 Epistemic Status of the Results

The results in this paper fall into three tiers of decreasing independence from auxiliary assumptions.

Tier 1 (Part I, from axioms alone). P-indivisibility of the reduced dynamics (§2.3), the characterization theorem establishing QM $\iff$ embedded observation under C1–C3 (§3.3), partition-relativity (§1.4), the deep-sector cardinality gauge corollary (§5.2), the consistency of nested trace-outs (Appendix B, Theorems B.1–B.4), the additive dissipator decomposition (B.5), the generalized second law (B.6), and the Page curve (B.7–B.9). These follow from Axioms 1–4, the Stinespring construction, standard quantum information results, and Popescu-Short-Winter typicality [41]; disputing them requires finding a mathematical error.

Tier 2 (Part II, given the cosmological realization). The gap equation $\hbar = c^3\epsilon^2/(4G)$ (§5.2), the Bekenstein-Hawking entropy with the 1/4 factor (§6), the CC dissolution identifying 10^{122} as $S_{\rm dS}$ (§7), and the dark matter acceleration scale $a_0 = cH/6$ with the baryonic Tully-Fisher relation $v^4 = GM_B \cdot cH/6$ (§8.4). These require accepting that the cosmological horizon satisfies C1–C3, verified in §4.2.

Tier 3 (consistency checks and solution-specific elements). The dark-sector concordance (~95% of $\rho_{\rm crit}$, §8.3), the dark energy parameter $\nu_{\rm OI}$ (§8.1), the GW echo timescale (§8.2), and the dark matter interpolation steepness (§8.4, Step 4). These are derived consequences or solution-specific elements; they would be falsified by observation but do not carry the same deductive weight as Tiers 1–2.

---

2. EMERGENT STOCHASTICITY AND P-INDIVISIBILITY

2.1 Emergent Stochasticity

The total system evolves deterministically, but the visible sector alone is stochastic. The observer knows x but not h; different hidden states h_k compatible with the same x send it to different futures

x'_k. Transition probabilities are obtained by marginalizing over the Liouville measure (§1.4). This makes the framework a hidden variable theory; consistency with Bell's theorem is addressed in §3.2.

2.2 The Slow-Bath Regime

The Markovian limit requires $\tau_B \ll \tau_S$ [4]. Condition (C2) inverts this: $\tau_S \ll \tau_B$. A slow bath must be distinguished from a static field. By (C1), coupling is continuously active: each visible-sector transition imprints on the hidden sector. Because the hidden sector is slow (not static), imprints persist without thermal overwriting. On subsequent transitions, coupling reads back stored correlations, producing history-dependent transition probabilities — the non-Markovian regime [4].

2.3 P-Indivisibility

By Axiom 2, the visible and hidden sectors have finite configuration spaces $\mathcal{C}_V$ and $\mathcal{C}_H$ (the discrete counterparts of Γ_V and Γ_H). On these finite sets:

Theorem. *Let $\mathcal{C}_V$ and $\mathcal{C}_H$ be finite sets with $|\mathcal{C}_V| \geq 2$, and let $\varphi : \mathcal{C}_V \times \mathcal{C}_H \to \mathcal{C}_V \times \mathcal{C}_H$ be a bijection. Define:*

$$T_{ij} = \frac{|\{h \in \mathcal{C}_H : \pi_V(\varphi(x_i, h)) = x_j\}|}{|\mathcal{C}_H|}$$

and the k-step matrix $T_{ij}^{(k)}$ by applying φ^k. If T is not a permutation matrix, then the process is P-indivisible.

Proof. Uses total variation distance $d(p, q) = \frac{1}{2}\sum_k |p_k - q_k|$. P-divisibility $\iff$ $d(p(t), q(t))$ non-increasing for all initial p, q [6, 7].

Step 1 (Recurrence). φ bijective on a finite set $\Rightarrow \exists N : \varphi^N = \mathrm{id}$. Thus $T^{(N)} = I$ and $d(\delta_i T^{(N)}, \delta_j T^{(N)}) = 1$ for $i \neq j$.

Step 2 (Strict contraction). T not a permutation $\Rightarrow \exists i, j, l$: $T_{il} > 0$ and $T_{jl} > 0$. Then:

$$d(\delta_i T, \delta_j T) = \frac{1}{2}\sum_k |T_{ik} - T_{jk}| < 1$$

Step 3. $d(\delta_i T^{(1)}, \delta_j T^{(1)}) < 1 = d(\delta_i T^{(N)}, \delta_j T^{(N)})$: non-monotonic, hence P-indivisible. □

The theorem requires only bijective dynamics (Axiom 1) and non-trivial coupling (C1). Axiom 2 guarantees the finite configuration space.

Convergence with prior work. The theorem converges with independent results: Pechukas [5] (reduced dynamics need not preserve positivity), Rivas et al. [6, 7] (divisibility failure ↔ information backflow), Pollock et al. [8] (process-tensor Markovianity conditions), and Strasberg and Esposito [9] (non-Markovian reduced dynamics in the slow-bath regime of C2). Bylicka, Johansson, and Acín [34] prove that for bijective dynamical maps — the class generated here — CP-divisibility and information backflow are equivalent, so the P-indivisibility established above implies CP-indivisibility directly. Milz et al. [35] further show that even CP-divisibility does not guarantee Markovianity in the process-tensor sense, confirming that the non-Markovian character identified here is robust under all current definitions.

Continuous-time extension. The Hamiltonian flow on finite-dimensional phase space preserves Liouville measure on compact energy surfaces. $T_{ij}(t)$ is continuous with $T(0) = I$. By (C1), $T(t)$ departs from the permutation class for $t > 0$. By Poincaré recurrence, $\exists t_R$: the set of hidden states with $\pi_V(\varphi_{t_R}(x_i, h)) = x_i$ has measure $> 1 - \delta$ for any $\delta > 0$. For small δ, this gives non-monotonic trace distance, establishing P-indivisibility in continuous time.

The recurrence argument establishes P-indivisibility in principle. The following lemma shows that under (C2), information backflow occurs on observable timescales — not merely at Poincaré recurrence times.

Lemma (Accessible-timescale backflow). *Under (C1)–(C3) with $\tau_S \ll \tau_B$, the non-Markovian mutual information $I(X_{<t}; X_{>t} \mid X_t)$ is $\mathcal{O}(1)$ for observation windows $t \sim k\tau_S$ with $k\tau_S \ll \tau_B$.*

Proof. The coupling H_{int} transfers visible-sector information to the hidden sector at each interaction, at rate $\sim 1/\tau_S$. Between interactions, the hidden sector evolves under H_H with spectral gap $\Delta \sim 1/\tau_B$. The decay of correlations stored in the hidden sector is governed by $e^{-\Delta\tau_S}$. When $\tau_S \ll \tau_B$, $\Delta\tau_S \ll 1$, so the decay per visible-sector transition is $1 - e^{-\Delta\tau_S} \approx \Delta\tau_S \ll 1$: correlations survive each step essentially intact. After k transitions spanning a time $k\tau_S \ll \tau_B$, the cumulative decay is $e^{-k\Delta\tau_S} \approx 1 - k\Delta\tau_S$, which remains close to unity. The hidden sector therefore retains $\sim k$ bits of visible-sector history over this window, and the $(k+1)$-th transition reads back stored correlations through H_{int}, producing history-dependent transition probabilities — i.e., information backflow. More precisely, the mutual information satisfies:

$$I(X_{<t}; X_{>t} \mid X_t) \geq I_0(1 - k\Delta\tau_S) = I_0 \left(1 - \frac{k\tau_S}{\tau_B}\right)$$

where $I_0 > 0$ is the single-step information transfer from (C1). For $k\tau_S \ll \tau_B$, this remains $\mathcal{O}(I_0)$ — comparable to the single-step coupling strength, not exponentially suppressed. The bound saturates (C3): the maximum storable history is $\log_2 m$ bits (§3.3), so persistent backflow over K transitions requires $m \geq 2^K$, which is amply satisfied when $N_H \gg N_V$. □

Role of (C2) and (C3). The Poincaré recurrence argument and the accessible-timescale lemma are independent: the former establishes P-indivisibility from (C1) and finiteness alone; the latter shows that (C2) and (C3) promote it from a formal property to an observationally dominant one.

Weak-coupling and fast-bath regimes. Known P-divisible systems with non-zero coupling illustrate the necessity of (C2), not a failure of the theorem. The Jaynes-Cummings model with Lorentzian spectral density is P-divisible when dissipation dominates coupling — the fast-bath regime ($\tau_B \ll \tau_S$), violating (C2). The Davies-Merkli Markovian convergence theorem likewise applies in the Born-Markov (fast-bath) limit. The §3.3 necessity proof for (C2) establishes this formally.

2.4 Explicit Construction: A Minimal Model

Visible sector: $x \in \{0, 1\}$. Hidden sector: $h \in \{1, \ldots, 6\}$. Dynamics: the permutation

$$\sigma = (0{,}1 \leftrightarrow 1{,}1)\ (0{,}2 \leftrightarrow 1{,}2)\ (0{,}3 \leftrightarrow 0{,}4)\ (0{,}5 \leftrightarrow 0{,}6)\ (1{,}3 \leftrightarrow 1{,}4)\ (1{,}5 \leftrightarrow 1{,}6)$$

satisfying (C1)–(C3). Since $\sigma^2 = \text{id}$: at $t = 1$, two of six hidden states flip x, giving

$$T(1,0) = \begin{pmatrix} 2/3 & 1/3 \\ 1/3 & 2/3 \end{pmatrix}$$

At $t = 2$ ($\sigma^2 = \mathrm{id}$): $T(2,0) = I$. A Markov chain would predict continued mixing: $T(1,0)^2 = \left(\begin{smallmatrix} 5/9 & 4/9 \\ 4/9 & 5/9 \end{smallmatrix}\right)$. The actual dynamics show complete *un-mixing*. Computing the intermediate propagator:

$$\Lambda(2,1) = T(2,0) \cdot [T(1,0)]^{-1} = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$$

Negative entries — no valid stochastic matrix exists. **P-indivisible.** □

The mechanism. The hidden sector acts as a memory register. The transpositions $(0,h) \leftrightarrow (1,h)$ for $h \in \{1,2\}$ flip x while preserving h, so a state at $x = 1$ with $h \in \{1,2\}$ after step 1 carries the record "I was at $x = 0$." Step 2 reads this record and flips x back — information backflow impossible for a memoryless process. The cosmological realization has $\tau_S/\tau_B \sim 10^{-32}$; P-indivisibility is more extreme, not less, when memory is more persistent.

---

3. THE EMERGENCE OF QUANTUM MECHANICS

3.1 The Stochastic-Quantum Correspondence

By Barandes' correspondence [10, 11], any P-indivisible stochastic process on a finite configuration space of size n embeds into a unitarily evolving quantum system on $\mathcal{H}$ of dimension $\leq n^3$. P-indivisible transition matrices develop negative entries at fine time resolution; the Stinespring dilation theorem guarantees $\exists$ a Hilbert space and unitary $U(t)$ with $T_{ij}(t) = |U_{ij}(t)|^2$ — transition probabilities are exactly Born-rule probabilities. The proof requires a finite-dimensional configuration space (Axiom 2); Calvo [12] extends the correspondence to infinite dimensions, but this is not required here. Pimenta [25] distinguishes P-divisible and P-indivisible stochastic-quantum dynamics: the correspondence is non-trivial only in the P-indivisible regime, which is precisely the regime produced by §2.3. An independent derivation using only Stinespring's dilation theorem [32] and standard partial-trace properties [33] is given in Appendix A; either route alone suffices.

From phase space to configuration space. The transition probabilities T_{ij} are projections of the full phase-space flow onto the configuration sector of Γ_V, with momenta and hidden-sector degrees of freedom absorbed into the Liouville marginalization. The resulting process lives on the discrete configuration space of Axiom 2.

Three features emerge. The Schrödinger equation arises from the differentiability of $U(t)$. The Born rule is the equilibrium distribution of the indivisible process [10, 11]. The action scale $\hbar$ enters when defining $\hat{H}(t) \equiv i\hbar\, \partial_t U \cdot U^\dagger$, converting dimensionless rates to energy units; its value requires physical input from the partition (§5).

Phase uniqueness from continuous-time data. The relation $T_{ij}(t) = |U_{ij}(t)|^2$ discards phase information at any single time. Doukas [26] identifies this gap for discrete-time data. The resolution: continuous-time data $\{T_{ij}(t)\}_{t\in\mathbb{R}}$ provides strictly more information.

Lemma (phase-locking). *Let H be Hermitian on $\mathbb{C}^n$ with eigenbasis $\{|k\rangle\}$, eigenvalues $\{E_k\}$, and $V_{ik} = \langle i|k\rangle$. Assume: (G1) non-degenerate spectrum; (G2) non-degenerate energy gaps; (G3)*

$V_{ik} \neq 0$ for all i, k. Then $T_{ij}(t) = |\langle i|e^{-iHt}|j\rangle|^2$ for all i, j, t uniquely determines H up to an overall energy shift and basis phase conventions.

Proof. Write:

$$T_{ij}(t) = \sum_{k,l} V_{ik}\, V_{jk}^*\, V_{jl}\, V_{il}^*\, e^{-i(E_k - E_l)t}$$

By (G2), the frequencies $\omega_{kl} = E_k - E_l$ are distinct for distinct pairs with $k \neq l$. Fourier-transforming yields:

$$a_{ij}^{kl} = V_{ik}\, V_{jk}^*\, V_{jl}\, V_{il}^*$$

The moduli $|V_{ik}|$ follow from $a_{ii}^{kl} = |V_{ik}|^2 |V_{il}|^2$ (non-zero by (G3)). The phases carry a double-difference structure:

$$\arg(a_{ij}^{kl}) = (\varphi_{ik} - \varphi_{il}) - (\varphi_{jk} - \varphi_{jl})$$

The gauge freedom preserving all double differences is $\varphi_{ik} \to \varphi_{ik} + \alpha_i + \beta_k$ (physically irrelevant basis rephasing). Fixing $\varphi_{1k} = 0$, $\varphi_{i1} = 0$, each remaining phase is $\varphi_{ik} = \arg(a_{i1}^{k1})$, directly extracted from Fourier data. $H = V\,\mathrm{diag}(E_k)\,V^\dagger$ is unique up to energy shift and phase conventions. □

Conditions (G1)–(G3) fail only on a measure-zero set of Hamiltonians. For the cosmological realization, a stronger argument applies. The genericity conditions concern the *effective* visible-sector Hamiltonian $\hat{H}_{\text{eff}}$ — the operator emerging after the trace-out — not the total Hamiltonian H_{tot}. The trace-out generically breaks the symmetries of H_{tot}, because the partition boundary does not respect them: the cosmological horizon is centered on a specific observer, breaking translational invariance, rotational symmetry about distant points, and boost invariance. Any degeneracy in H_{tot} that relies on correlations between visible and hidden sectors — which includes all degeneracies associated with symmetries the partition breaks — is lifted in $\hat{H}_{\text{eff}}$. The residual degeneracies of $\hat{H}_{\text{eff}}$ are those associated with symmetries acting entirely within the visible sector that commute with the trace-out: precisely the gauge symmetries of the emergent QFT. Gauge degeneracies correspond to physically equivalent states and do not affect the phase-locking argument, which operates on gauge-inequivalent transition data. The cosmological $\hat{H}_{\text{eff}}$ therefore satisfies G1–G3 not by typicality in an abstract ensemble, but because the partition itself acts as a symmetry-breaking mechanism.

3.2 Bell Inequality Violations

The framework is a hidden variable theory evading Bell's theorem [13] not by superdeterminism but by violating factorizability through P-indivisible joint dynamics — while remaining causally local. Two subsystems interacting at preparation carry a joint transition matrix $T_{QR} \neq T_Q \otimes T_R$. This non-factorizability *is* stochastic entanglement [10, 11, 14]. Since the process is indivisible, no well-defined intermediate conditional probabilities permit Bell's factorization.

In the Jarrett decomposition, the framework violates outcome independence while preserving parameter independence and measurement independence — precisely the class Fine's theorem [24] permits. Barandes, Hasan, and Kagan [14] prove the maximum CHSH correlator is exactly Tsirelson's

bound $2\sqrt{2}$, with independent support from Le et al. [15]. The bound also follows from the Stinespring route: Appendix A establishes full unitary QM on a tensor-product Hilbert space, from which Tsirelson's original operator-norm argument [36] yields $|\langle S\rangle| \leq 2\sqrt{2}$; achievability follows because all quantum states, including maximally entangled ones, are realizable in the emergent description.

Fine-tuning and causal structure. The framework does not claim to restore Bell locality: outcome independence is genuinely violated. What it provides is a *derivation* of this nonlocality from the causal partition, rather than treating it as axiomatic. The fine-tuning objection (Wood and Spekkens, 2015) assumes DAG causal structure with Markov factorization; P-indivisible processes violate the Markov condition on any DAG, so the appropriate framework is the process tensor [8], within which no-signaling follows from marginalization structure without fine-tuning.

Parameter independence from spatial locality. Let V_A and V_B be space-like separated visible-sector subsystems with disjoint coupling neighborhoods on G_φ. The spatial Markov property of range-1 dynamics (Fundamental, §5, area-law lemma) makes V_A and V_B conditionally independent given boundary data. Alice's measurement setting a determines which function she evaluates on V_A; it does not alter the transition probabilities for V_B, which depend only on V_B's local coupling neighborhood. Hence $P(x_B \mid a, b) = P(x_B \mid b)$ — parameter independence holds structurally for all bounded-degree graph topologies with range-1 coupling, with no fine-tuning of the hidden-variable distribution. Combined with the outcome-independence violation from stochastic entanglement, this gives the precise Jarrett decomposition.

3.3 The Characterization Theorem: Necessity of the Conditions

The logical structure: Barandes' correspondence gives QM $\iff$ P-indivisibility. What remains is P-indivisibility $\implies$ embedded observation under (C1)–(C3).

Dilation existence. Any stochastic process on a finite configuration space can be realized as marginalization of a deterministic process on a larger state space — furnishing Axioms 1–4.

Theorem (C1 equivalence). *The marginalized process is P-indivisible iff T is not a permutation matrix.*

Proof. Forward: §2.3. Reverse: if T is a permutation, $\Lambda(k_2, k_1) = T^{(k_2-k_1)}$ is a valid stochastic matrix for all $k_2 > k_1 \Rightarrow$ P-divisible. □

Theorem (C2 necessity). *In the fast-bath regime ($\tau_B \ll \tau_S$), ergodic mixing drives the hidden sector to uniformity before each coupling event, yielding a Markov chain ($T^{(k)} = T^k$), which is P-divisible. Contrapositively, P-indivisibility requires $\tau_S \ll \tau_B$.*

Proof. Between coupling events (separated by τ_S), the hidden sector evolves under its own Hamiltonian H_H. Let $\mathcal{L}_H$ denote the hidden-sector Liouvillian with spectral gap $\Delta > 0$ (guaranteed on a finite ergodic system). For any conditioned hidden-sector distribution $\mu_H(\cdot|x_i)$ at a coupling event:

$$\|e^{\mathcal{L}_H \tau_S}\mu_H(\cdot|x_i) - \mu_{\text{eq}}\|_{\text{TV}} \leq C\, e^{-\Delta\tau_S}$$

In the fast-bath regime ($\Delta\tau_S \gg 1$), this is exponentially small: the hidden sector relaxes to equilibrium μ_{eq} before each coupling event regardless of its post-coupling state. Each single-step transition matrix T is therefore computed against the same equilibrium distribution, so $T^{(k)} = T^k$ — a homogeneous Markov chain, hence P-divisible.

Contrapositively: P-indivisibility requires $\Delta\tau_S \lesssim 1$, i.e., $\tau_S \lesssim \tau_B$. For the strong, persistent P-indivisibility of the characterization theorem, the separation must be $\tau_S \ll \tau_B$. (The proof uses discrete coupling events; for continuous Hamiltonian flow, τ_S is the visible sector's dynamical timescale and Δ is the spectral gap of $\mathcal{L}_H$. The continuous-time extension replaces the collision-model structure with the continuous spectral condition $\Delta\tau_S \gg 1 \Rightarrow$ Markovian, established in [4, 9].) □

Theorem (C3 necessity). *Let $m = |\mathcal{C}_H|$. The non-Markovian mutual information satisfies:*

$$I(X_{<t}; X_{>t} \mid X_t) \leq \log_2 m$$

Proof. The total system is deterministic: $X_{>t}$ is a function of (X_t, H_t). Conditioning on X_t: $X_{<t} \to H_t \to X_{>t}$ is a Markov chain. By data processing:

$$I(X_{<t}; X_{>t} \mid X_t) \leq I(X_{<t}; H_t \mid X_t) \leq H(H_t \mid X_t) \leq \log_2 m \quad \square$$

Corollary. P-indivisibility across n visible configurations requires $m \geq n$, and persistence over K transitions requires m/n growing with the observation window — precisely (C3). □

Definition (unitarily evolving QM). A stochastic process $\mathcal{S}$ on a finite configuration space $\mathcal{C}_V = \{x_1, \dots, x_n\}$ is *mathematically equivalent to unitarily evolving quantum mechanics* if there exists a Hilbert space $\mathcal{H}_V$ of dimension $\leq n^3$, a Hermitian operator $\hat{H}$ on $\mathcal{H}_V$, and a unitary family $U(t) = e^{-i\hat{H}t}$ such that $T_{ij}(t) = |U_{ij}(t)|^2$ for all i, j, t — i.e., transition probabilities are exactly Born-rule probabilities. This definition captures the Hilbert space, unitary dynamics, and the Born rule. Additional quantum-mechanical structures — the tensor product decomposition for spatially separated visible-sector subsystems, state update, and the measurement formalism — are addressed in the remarks following the theorem.

Characterization theorem. *For $|\mathcal{C}_V| \geq 2$, the following are equivalent:*

(i) $\mathcal{S}$ is mathematically equivalent to unitarily evolving QM (in the sense of the preceding definition).

(ii) $\mathcal{S}$ is P-indivisible.

(iii) $\mathcal{S}$ arises from marginalizing a deterministic bijection on $\mathcal{C}_V \times \mathcal{C}_H$ with (C1) non-trivial coupling, (C2) slow-bath memory, and (C3) sufficient capacity.

Proof. (i) $\iff$ (ii): Barandes [10, 11]. (iii) $\implies$ (ii): §2.3. (ii) $\implies$ (iii): necessity theorems above. □

QM is not merely *compatible with* embedded observation — it is *equivalent to* it.

Remark (scope of the equivalence). The characterization theorem establishes the Hilbert space, unitary dynamics, and Born-rule transition probabilities. Four further structures of operational quantum mechanics merit comment.

(i) Tensor product structure (visible–hidden). The Stinespring route (Appendix A) constructs $\mathcal{H} = \mathcal{H}_V \otimes \mathcal{H}_H$ and derives the CPTP channel Φ by partial trace. The visible–hidden tensor product is therefore given by the construction, not postulated.

(ii) Tensor product structure (visible-sector subsystems). Spatially separated subsystems within the visible sector — e.g., two laboratories — correspond to subsets $V_A, V_B \subset V$ with disjoint coupling

neighborhoods on the coupling graph (Fundamental, §3). The spatial Markov property of range-1 dynamics (Fundamental, §5, area-law lemma) guarantees conditional independence: $\rho_{AB} = \rho_A \otimes \rho_B$ conditioned on boundary data. The emergent Hilbert space inherits the factorization $\mathcal{H}_{V_A} \otimes \mathcal{H}_{V_B}$ from the locality of the coupling graph, with entanglement between A and B arising from shared boundary history — precisely the stochastic entanglement of [10, 14].

(iii) State update, measurement, and multi-time predictions. Projective measurement in the emergent description corresponds to conditioning on a visible-sector outcome and re-marginalizing over the hidden sector. The Lüders rule $\rho \to P_k \rho P_k / \mathrm{Tr}(P_k \rho)$ is the quantum transcription of Bayesian updating on the classical substratum (Axiom 4). No independent measurement postulate is required. Multi-time correlation functions, weak values, and Leggett-Garg inequalities are standard calculations within the delivered formalism ($\mathcal{H}_V$, $U(t)$, Born rule); they require no additional postulates beyond what the characterization theorem provides.

(iv) Classical non-Markovianity and quantum mechanics. The equivalence is not a claim that all P-indivisible processes are "secretly quantum" in some metaphysical sense; it is a theorem that the mathematical structures are identical. Classical systems exhibiting P-indivisibility — renewal processes, semi-Markov processes on finite state spaces — admit a unique unitary description (phase-locking lemma) that makes predictions beyond the transition data (energy eigenvalues, interference, entanglement), and can be realized as marginalization of deterministic dynamics under C1–C3 (necessity theorems). The framework does not distinguish "classical non-Markovian" from "quantum" — it proves they are the same category. The physical content of the unitary is not that of a mere mathematical embedding: it is the *specific* Hamiltonian uniquely determined by the transition data, arising from the *actual* dilation structure of the system (Appendix A).

PART II: THE COSMOLOGICAL APPLICATION

4. THE COSMOLOGICAL HORIZON AS CAUSAL PARTITION

4.1 The Partition

The cosmological horizon — the boundary beyond which no signal propagating at or below c can reach the observer — implements the causal partition of Axiom 3: Γ_V is the interior, Γ_H everything beyond. This is a consequence of GR's causal structure, not a modeling choice. The universality of the observed quantum mechanics follows from all sub-c observers sharing the same causal structure.

Different observers have different horizon areas (hence different S_{dS}), but $\hbar = c^3\epsilon^2/(4G)$ depends only on local geometric quantities — not on the observer's worldline or horizon area — so all observers share the same emergent action scale.

4.2 Verification of the Conditions

(C1) In the ADM formulation, the bulk Hamiltonian is a sum of constraints that vanish on-shell; physical time evolution is generated by the boundary term at the horizon, which depends on data from both sides. The Hamiltonian and momentum constraints further correlate interior and exterior initial data on any Cauchy surface, so the physical phase space is a constraint surface within the kinematic product $\Gamma_V \times \Gamma_H$ (see §9.6 for implications). This is *stronger* than the $H_{\mathrm{int}} \neq 0$ required by Axiom 3: constraints enforce correlations that persist on all timescales and cannot be perturbatively removed. **Satisfied.**

(C2) $\tau_B \gtrsim 1/H \sim 10^{17}$ s; for laboratory processes, $\tau_S \sim 10^{-15}$ s. Ratio: $\tau_S/\tau_B \sim 10^{-32}$. **Satisfied.**

(C3) Hidden sector has A/ϵ^2 modes with $A \sim 10^{52}$ m^2; visible sector has $\sim 10^{80}$ baryons. No laboratory process saturates the hidden sector. **Satisfied.**

4.3 Application

Axiom 2 is now a consequence of partition geometry: the horizon has finite area $A = 4\pi c^2/H^2$, bounding coupled modes to $A/\epsilon^2 < \infty$. The Part I theorem applies: the observer's reduced description is P-indivisible and, by Barandes' correspondence, equivalent to unitary QM with $\hbar$ determined by the partition boundary.

---

5. DERIVATION OF THE EMERGENT ACTION SCALE ($\hbar$)

5.1 The Classical Horizon Temperature

Jacobson [16] showed $dE = T\,dS$ applied to local causal horizons yields Einstein's equations, with $dE = (c^2\kappa/8\pi G)\,dA$, where κ is the surface gravity of the horizon. The entropy density is $\eta = 1/\epsilon^2$ — one coupled mode per minimal cell. This is not an assumption about the number of states per cell but a consequence of two results: ϵ is defined as the minimal distinguishable scale (Axiom 2), so each cell of area ϵ^2 contributes exactly one boundary mode that couples across the partition; the number of internal states per mode (the alphabet size q) is a gauge freedom with no observable consequences (Fundamental, §4). Thus $dS = dA/\epsilon^2$. Equating:

$$T_{\rm cl} = \frac{c^2\epsilon^2\kappa}{8\pi G k_B}$$

For de Sitter ($\kappa = cH$): $k_B T_{\rm cl} = c^3\epsilon^2 H/(8\pi G)$. No $\hbar$ appears.

5.2 The Emergent Action Scale

Step 1: Uniqueness. By partition-relativity (§1.4), $\hbar$ is derived, not free.

Step 2: $\hbar$ is structural, not volumetric.

Lemma (Boundary-only dependence). *Decompose the hidden sector as $\mathcal{C}_H = \mathcal{C}_B \times \mathcal{C}_D$, where $\mathcal{C}_B$ denotes boundary modes (within the interaction range of H_{int}) and $\mathcal{C}_D$ denotes deep modes (the remainder of the hidden sector). If the classical Hamiltonian is spatially local, then on timescales $t \ll \tau_B$:*

$$T_{ij}(t) = T_{ij}^{(B)}(t) + \mathcal{O}(t/\tau_B)$$

where $T_{ij}^{(B)}$ depends only on the V-B dynamics.

Proof. (i) *Coupling structure.* Spatial locality of the classical Hamiltonian implies the interaction decomposes as $H_{\rm tot} = H_V + H_B + H_D + H_{VB} + H_{BD}$, with no direct V-D coupling: a deep hidden-sector mode must propagate through the boundary layer before influencing the visible sector. The coupling chain is $V \leftrightarrow B \leftrightarrow D$, not $V \leftrightarrow D$.

(ii) *Frozen deep sector.* Let Δ_D be the spectral gap of $H_D + H_{BD}$ restricted to the deep sector conditioned on a fixed boundary configuration. The deep-sector evolution over time t displaces any initial configuration by $\mathcal{O}(\Delta_D\, t)$ in phase space. By condition (C2), $\Delta_D \sim 1/\tau_B$, so for $t \ll \tau_B$: $\|\varphi_t^D - \mathrm{id}\| = \mathcal{O}(t/\tau_B)$.

(iii) *Factorization of the Liouville integral.* The transition probability is:

$$T_{ij}(t) = \frac{1}{|\mathcal{C}_B||\mathcal{C}_D|} \sum_{b\in\mathcal{C}_B} \sum_{d\in\mathcal{C}_D} \delta_{x_j}[\pi_V(\varphi_t(x_i, b, d))]$$

By (i), $\pi_V(\varphi_t(x_i, b, d))$ depends on d only through the back-reaction $B \leftarrow D$, which by (ii) shifts b by $\mathcal{O}(t/\tau_B)$. Expanding:

$$\pi_V(\varphi_t(x_i, b, d)) = \pi_V(\varphi_t^{VB}(x_i, b)) + \mathcal{O}(t/\tau_B)$$

where φ_t^{VB} is the flow restricted to $V \times B$ with D frozen. The d-sum then contributes $|\mathcal{C}_D|$ identical terms at leading order:

$$T_{ij}(t) = \underbrace{\frac{1}{|\mathcal{C}_B|} \sum_{b\in\mathcal{C}_B} \delta_{x_j}[\pi_V(\varphi_t^{VB}(x_i, b))]}_{T_{ij}^{(B)}(t)} + \mathcal{O}(t/\tau_B) \qquad \square$$

The correction is $\mathcal{O}(t/\tau_B) \sim 10^{-32}$ for laboratory processes. Since $T_{ij}(t)$ determines the emergent quantum description (§3.1), and $T_{ij}^{(B)}(t)$ depends only on the V-B dynamics — which are characterized by the boundary geometry (A, ϵ, κ) and the constants c, G appearing in the classical Hamiltonian — the emergent action scale $\hbar$ inherits boundary-only dependence.

Corollary (deep-sector cardinality is gauge). *No observable of the emergent description — no transition probability, emergent Hamiltonian eigenvalue, or scattering amplitude — depends on the cardinality $|\mathcal{C}_D|$ of the deep hidden sector. Systems with the same $\mathcal{C}_V \times \mathcal{C}_B$ dynamics produce the same emergent physics to within $\mathcal{O}(\tau_S/\tau_B)$, whether $|\mathcal{C}_D|$ is finite, countably infinite, or uncountably infinite.*

Proof. $T_{ij}^{(B)}(t)$ is defined by a sum over $\mathcal{C}_B$ alone. The $\mathcal{C}_D$-dependent terms enter only through the $\mathcal{O}(t/\tau_B)$ back-reaction. Since the emergent Hamiltonian $\hat{H}_{\text{eff}}$ is uniquely determined by $\{T_{ij}(t)\}$ (phase-locking lemma, §3.1; D-gauge theorem, §5.3), it inherits $\mathcal{C}_D$-independence at leading order. All observables — eigenvalues, transition amplitudes, scattering cross-sections — are functions of $\hat{H}_{\text{eff}}$ and therefore $\mathcal{C}_D$-independent. $\square$

The cardinality of the deep hidden sector is gauge in the same technical sense as the alphabet size q (Fundamental, §4): just as two systems with different q are physically equivalent if they share the same coupling structure, two systems with different $|\mathcal{C}_D|$ are physically equivalent if they share the same $\mathcal{C}_V \times \mathcal{C}_B$ dynamics. Axiom 2 requires S finite for the recurrence proof (§2.3); the corollary establishes that the deep sector's contribution to this finiteness has no observable consequences. The question "is the universe finite or infinite?" has no empirical content within the framework — it is identified as gauge.

Step 3: Dimensional determination. Step 2 excludes volumetric (deep-sector) quantities, leaving boundary quantities. The boundary carries both *local* geometric data (ϵ, κ, and the constants c, G of the classical Hamiltonian) and a *global* quantity: the total area A, which forms the dimensionless ratio $A/\epsilon^2 = S_{\rm dS}$. If $\hbar$ depended on $S_{\rm dS}$, it would be observer-dependent — different observers have different horizon areas (§4.1) — contradicting the universality of the emergent action scale. (All sub-c observers share the same local physics and hence the same $\hbar$; §4.1.) This excludes A, leaving the local boundary quantities c, G, ϵ, and κ. Since κ is the surface gravity of a specific horizon (observer-dependent, like A), it too is excluded. The unique combination of c, G, ϵ with dimensions of action:

$$\hbar = \beta \frac{c^3 \epsilon^2}{G}$$

Step 4: Thermal self-consistency. The classical substratum assigns the partition boundary a temperature $k_B T_{\rm cl} = c^3\epsilon^2\kappa/(8\pi G)$ (§5.1), containing no $\hbar$. The emergent QFT of Part I lives on this classical background, which has a bifurcate Killing horizon with surface gravity κ. Regularity of the Wick-rotated metric at the horizon requires Euclidean period $\beta = 2\pi c/\kappa$; any QFT on this background — including a lattice-regularized one — must therefore be periodic in imaginary time with the same period, giving a KMS state at temperature $T_Q = \hbar\kappa/(2\pi c k_B)$, with $\hbar$ unknown. This is a theorem *within* the derived QFT, not an external import. The two temperatures are computed independently — $T_{\rm cl}$ from the classical substratum alone (no QM), T_Q from the emergent QFT alone (no classical substratum details) — but they describe the same physical degrees of freedom: the boundary modes across which the partition is defined. Since the quantum description is derived from the classical one (Part I), and the derivation is exact at the boundary (Step 2's correction is $\mathcal{O}(\tau_S/\tau_B) \sim 10^{-32}$), the two descriptions cannot assign contradictory temperatures to the same boundary. Consistency requires $T_{\rm cl} = T_Q$:

$$\frac{c^3\epsilon^2\kappa}{8\pi G} = \frac{\hbar\kappa}{2\pi c}$$

κ cancels (confirming the structural/volumetric distinction). Solving:

$$\boxed{\hbar = \frac{c^3\epsilon^2}{4G}}$$

The derivation is a gap equation: ϵ is the free geometric input, $\hbar$ is the output. The match is non-trivial: $T_{\rm cl}$ could have depended on volumetric hidden-sector data (excluded by Step 2), or T_Q could have depended on the state (it does not — the KMS temperature is purely kinematic). That neither pathology obtains makes the gap equation a genuine determination. The non-circularity is structural: Part I establishes that a QFT emerges with *some* action scale $\hbar$; §5 determines *which* $\hbar$, using the independent classical temperature that Part I neither requires nor produces.

Non-circularity of the entropy density. The claim $\eta = 1/\epsilon^2$ — one mode per minimal cell — may appear to pre-bake the area-entropy proportionality. To see why it does not, consider the counterfactual: suppose the classical entropy density were $\eta = \alpha/\epsilon^2$ for arbitrary $\alpha > 0$. Then $T_{\rm cl} = \alpha c^2\epsilon^2\kappa/(8\pi G k_B)$, and thermal matching gives $\hbar = \alpha c^3\epsilon^2/(4G)$. The Bekenstein-Hawking entropy becomes $S = A/(4\alpha^{-1} l_p^2)$, which reproduces the observed 1/4 factor only for $\alpha = 1$. Thus $\alpha = 1$ is a non-trivial constraint — it says that each minimal cell contributes exactly one boundary

mode, not a fractional or multiple contribution. This is the physical content of the q-gauge theorem (Fundamental, §4): the alphabet size q (the number of internal states per cell) is observationally irrelevant, so what counts is the number of *cells*, not the number of *states*. The counting $\eta = 1/\epsilon^2$ is not an assumption about mode density but a consequence of ϵ being defined as the minimal distinguishable scale (Axiom 2) combined with the irrelevance of sub-cell structure.

The predictive content lies not in the gap equation alone but in its consequences. The specific relationship $\hbar = c^3\epsilon^2/(4G)$ — rather than any other function of c, G, ϵ — produces: the Bekenstein-Hawking formula with the exact factor 1/4 (§6), the CC dissolution with $S_{\rm dS}$ as the compression ratio (§7.3), the RVM parameter $\nu_{\rm OI}$ (§8.1), and the GW echo timescale (§8.2). Any alternative $\hbar(\epsilon)$ would fail at least one of these checks. The situation is analogous to deriving the Schwarzschild metric with M as a free parameter: the derivation has genuine content (the functional form) even though one input is not determined from within.

Jacobson's original derivation [16] uses $\hbar$-containing forms; this paper does not — it uses the classical identity $dE = (c^2\kappa/8\pi G)\,dA$ and the classical entropy density $\eta = 1/\epsilon^2$. The logical ordering differs from Jacobson's: his argument derives G from η and $\hbar$; the present argument takes G as a parameter of the classical Hamiltonian (Axiom 1) and derives $\hbar$ from G and ϵ. The two are consistent — substituting $\eta = 1/\epsilon^2$ and $\hbar = c^3\epsilon^2/(4G)$ into Jacobson's formula $G_{\rm eff} = c^4/(8\pi\eta\cdot\hbar c)$ recovers $G_{\rm eff} = G$ — but the roles of input and output are reversed. The Gibbons-Hawking temperature [17] $k_B T_{\rm GH} = \hbar\kappa/(2\pi c)$ is recovered as a prediction, not used as an input. The KMS condition used above is derived for continuum QFT on bifurcate Killing horizons; for the lattice-regularized theory emerging from Part I, the relevant result is that thermal periodicity of the Euclidean Green's functions is robust against UV modifications of the dispersion relation, with corrections at $\mathcal{O}((\epsilon\kappa/c)^2)$ [28, 29]. For the cosmological horizon, $\epsilon\kappa/c = \epsilon H/c \sim 10^{-61}$, so lattice corrections are $\sim 10^{-122}$.

5.3 Gauge-Fixing and the Dimensional Obstruction

Steps 1–2 establish that $\hbar$ depends only on boundary quantities; Step 4 will fix its value via thermal matching. A prior question must be addressed: does the transition-probability data $\{T_{ij}(t)\}$ determine the emergent Hamiltonian uniquely, or does residual gauge freedom leave $\hbar$ underdetermined? The following theorem, completing the phase-locking argument of §3.1, shows the gauge freedom is physically trivial.

Theorem (D-gauge completeness). *Let $U(t) = e^{-iHt}$ with non-degenerate eigenvalues and non-vanishing configuration-basis overlaps. If $|U'_{ij}(t)|^2 = |U_{ij}(t)|^2$ for all i, j, t, then $H' = DHD^\dagger$ for a time-independent diagonal unitary D.*

Proof. Write $U_{ij}(t) = \sum_a V_{ia}\, e^{-iE_a t}\, V^*_{ja}$ with $V_{ia} = \langle i|a\rangle$ and eigenvalues $\{E_a\}$, and similarly $U'_{ij}(t) = \sum_a V'_{ia}\, e^{-iE'_a t}\, V'^*_{ja}$.

Step 1 (Eigenvalue recovery). By the phase-locking argument of §3.1, $|U_{ij}(t)|^2 = |U'_{ij}(t)|^2$ for all t implies, via Fourier analysis, that the frequency sets $\{E_a - E_b\}$ and $\{E'_a - E'_b\}$ coincide. Non-degeneracy of energy gaps then gives $E'_a = E_{\sigma(a)} + E_0$ for some permutation σ and global shift E_0. The Fourier coefficients $a^{kl}_{ij} = V_{ik}V^*_{jk}V_{jl}V^*_{il}$ must match between primed and unprimed, which (by the non-vanishing condition on overlaps) forces $\sigma = \mathrm{id}$ up to relabeling. Thus $E'_a = E_a + E_0$.

Step 2 (Modulus recovery). From the diagonal Fourier coefficients $a^{kl}_{ii} = |V_{ik}|^2|V_{il}|^2$, the non-vanishing condition gives $|V'_{ia}| = |V_{ia}|$ for all i, a.

Step 3 (Phase structure). Write $V'_{ia} = V_{ia}\, e^{i\delta_{ia}}$. The off-diagonal Fourier coefficients require:

$$V'_{ik} V'^{*}_{jk} V'_{jl} V'^{*}_{il} = V_{ik} V^{*}_{jk} V_{jl} V^{*}_{il}$$

Substituting and canceling moduli:

$$e^{i(\delta_{ik} - \delta_{jk} - \delta_{il} + \delta_{jl})} = 1 \quad \text{for all } i, j, k, l$$

This is the double-difference condition: $\delta_{ik} - \delta_{il} - \delta_{jk} + \delta_{jl} = 0 \pmod{2\pi}$. The general solution on the integer lattice is $\delta_{ia} = \alpha_i + \beta_a$ for phases $\{\alpha_i\}$, $\{\beta_a\}$. Thus $V' = D_\alpha\, V\, D_\beta$ where $D_\alpha = \mathrm{diag}(e^{i\alpha_i})$ and $D_\beta = \mathrm{diag}(e^{i\beta_a})$. The D_β factor is absorbed into the eigenvector phase convention (physically irrelevant), giving $H' = D_\alpha H D_\alpha^\dagger$ with $D = D_\alpha$ a time-independent diagonal unitary. □

The D-gauge is physically trivial (basis rephasing). Cross-interval transition probabilities fix the relative gauge up to a single global phase; in the continuum limit this becomes an unobservable smooth function $e^{i\theta(t)}$.

Dimensional obstruction. The unitary $U(t)$ is dimensionless; $\hbar$ enters only via $\hat{H} = i\hbar\, \partial_t U \cdot U^\dagger$. No dimensionless data can fix a dimensionful constant — Step 4's thermal self-consistency provides the necessary dimensionful input.

Remark (phase recovery under discrete sampling). The spatial lattice discreteness does not discretize time: the continuous-time extension (§2.3) provides $T_{ij}(t)$ as a genuinely continuous function of t via the Liouville marginalization of the Hamiltonian flow (Axiom 1). The D-gauge theorem therefore applies to the continuous object without aliasing corrections. Aliasing would arise only if the observer sampled T_{ij} at discrete intervals, in which case trans-Planckian frequencies could be misidentified — but these lie outside the emergent QFT's domain. The phase-recovery error is controlled by the same $\mathcal{O}(\tau_S/\tau_B) \sim 10^{-32}$ suppression that governs the boundary-only dependence of §5.2.

---

6. SELF-CONSISTENCY AND THE DISCRETENESS SCALE

Rearranging $\hbar = c^3\epsilon^2/(4G)$:

$$\epsilon^2 = \frac{4\hbar G}{c^3} = 4\, l_p^2$$

The framework yields $\epsilon = 2\, l_p$ exactly. This is algebraically equivalent to the formula for $\hbar$: the framework contains one free geometric parameter (ϵ), and thermal self-consistency establishes its relationship to $\hbar$. The result is a self-consistency condition constraining ϵ to the Planck regime, not an independent determination. If $\epsilon^2 \ll l_p^2$, sub-Planckian subcells would create a second trace-out, breaking the single-valuedness of $\hbar$. If $\epsilon^2 \gg l_p^2$, the emergent description would assign distinct quantum states to physically identical configurations. The framework's predictive content comes from the relationship $\hbar = c^3\epsilon^2/(4G)$, the Bekenstein-Hawking formula, and the consequences that follow.

With $\epsilon = 2\, l_p$, the number of independent modes on the cosmological horizon is:

$$N_{\text{modes}} = \frac{A}{\epsilon^2} = \frac{A}{4\,l_p^2} = S_{\text{dS}}$$

where $S_{\text{dS}} = A/(4\,l_p^2)$ is the Bekenstein-Hawking entropy of the de Sitter horizon. The factor of 1/4 in the Bekenstein-Hawking formula — historically introduced as a proportionality constant — is here derived: each minimal cell of area $\epsilon^2 = 4\,l_p^2$ on the horizon contributes one unit of entropy, and $A/(4\,l_p^2) = A/\epsilon^2$ recovers the standard formula exactly.

---

7. DISSOLUTION OF THE COSMOLOGICAL CONSTANT PROBLEM

7.1 The Two Levels of Description

From finite-dimensional QM to QFT. Part I delivers QM on a finite-dimensional Hilbert space. The $N = A/\epsilon^2$ cells discretize the visible sector; spatial locality of the classical substratum restricts which cells interact on any given timescale. The emergent Hilbert space decomposes as $\mathcal{H} = \bigotimes_k \mathcal{H}_k$ — a lattice-regularized QFT with UV cutoff at $\epsilon = 2\,l_p$. The tensor product structure follows from the spatial product structure of the classical configuration space $\mathcal{C}_V = \mathcal{C}_1 \times \cdots \times \mathcal{C}_N$ (one factor per cell), which is not required by Axiom 2 but is a consequence of spatial locality in the classical substratum: each cell's configuration is a local degree of freedom.

Locality preservation theorem. *If the classical Hamiltonian is spatially local (couples only neighbors), then the emergent quantum Hamiltonian inherits spatial locality.*

Proof. For infinitesimal dt and $x' \neq x$: $T_{x,x'}(dt) = |\langle x'|\hat{H}|x\rangle|^2\,dt^2 + O(dt^3)$. If x, x' differ at non-neighboring sites, spatial locality of the classical dynamics gives $T_{x,x'}(dt) = 0$ at leading order, hence $|\langle x'|\hat{H}|x\rangle|^2 = 0$. By the D-gauge theorem (§5.3), the residual phase freedom is a diagonal unitary D that preserves the vanishing of off-diagonal moduli, so $\langle x'|\hat{H}|x\rangle = 0$ in any gauge. The emergent Hamiltonian has the form $\hat{H} = \sum_k \hat{h}_k + \sum_{\langle k,l\rangle} \hat{h}_{kl}$. □

Scope of the emergent QFT. The CC dissolution requires only that the emergent description assigns zero-point energy $\sim \hbar\omega/2$ per mode with UV cutoff at ϵ^{-1} — a property shared by any lattice QFT with local couplings, regardless of gauge group or matter content. The framework's predictions in Part II ($\hbar$, S_{dS}, ν_{OI}) depend on partition geometry, not on the emergent field content.

Classical substratum (what geometric measurements probe): The horizon has classical thermal equilibrium. By the Friedmann equation, $\rho_{\text{crit}} = 3H^2c^2/(8\pi G)$. No zero-point energy, no discrepancy. Total vacuum energy density: $\rho \sim H^2/G \sim 10^{-9}$ J/m^3 — precisely observed.

Emergent QFT (what local quantum measurements probe): Zero-point energy $\frac{1}{2}\hbar\omega$ per mode summed to the Planck cutoff gives $\rho_{\text{QFT}} \sim 10^{113}$ J/m^3 — a $\sim 10^{122}$ discrepancy with the observed value [2, 3, 18].

7.2 Why Gravity Sees the Classical Value

Spacetime geometry is part of the classical substratum (Axiom 3): the metric evolves via Einstein's equations *before* the trace-out. The stress-energy sourcing gravity is the classical stress-energy of the total microstate, not the expectation value of an emergent quantum operator. The semi-classical equation $G_{\mu\nu} = 8\pi G\langle\hat{T}_{\mu\nu}\rangle$ is itself an emergent approximation; at the classical level, the gravitational field equations never encounter the zero-point sum.

Ontological commitment. Classical spacetime is fundamental; QM is emergent. This ordering is not a modeling choice but is forced by the logical structure of the derivation through three independent requirements:

(i) Definiteness. The trace-out that produces QM is defined by a *specific* partition $\Gamma = \Gamma_V \times \Gamma_H$. The partition must be definite — not in superposition, not subject to quantum uncertainty — for the marginalization integral (§1.4) to be well-defined. A partition in superposition would yield an incoherent mixture of inequivalent quantum theories, not a single QM. The metric at the partition boundary must therefore be classical.

(ii) Non-circularity. The partition is defined by the causal structure: Γ_H is the set of degrees of freedom beyond the observer's causal reach (§4.1). Causal structure is determined by null geodesics of the metric. If the metric were derived from QM, then the derivation would require: QM $\rightarrow$ metric $\rightarrow$ causal structure $\rightarrow$ partition $\rightarrow$ QM — a circular dependency with no entry point. The circle breaks only if the metric exists prior to the partition.

(iii) Uniqueness of $\hbar$. The emergent action scale is determined by the boundary geometry (§5.2). If the geometry were itself quantum-mechanical, $\hbar$ would depend on a quantum state — but $\hbar$ is state-independent by observation. The geometry-first ordering is the only one consistent with a universal $\hbar$.

Any framework in which QM is logically prior to the metric must either (a) treat the zero-point sum as a gravitational source — producing the 10^{122} discrepancy requiring cancellation or fine-tuning — or (b) invoke an independent mechanism to decouple quantum vacuum energy from gravity. In the present framework, the zero-point sum is an artifact of the emergent description and never enters the stress-energy tensor; no decoupling mechanism is needed.

Cumulative evidence for the ordering. The geometry-first ordering produces seven independent consequences that match observation, none of which were designed into the axioms: (1) the CC dissolution — the observed vacuum energy sits at $\rho \sim H^2/G$, the classical baseline, without fine-tuning (§7.3); (2) the Bekenstein-Hawking formula including the 1/4 factor, derived from $\epsilon = 2\,l_p$ (§6); (3) the value of $\hbar$, determined by partition geometry rather than treated as a free parameter (§5.2); (4) the dark-sector concordance — ~95% of ρ_{crit} is invisible to QFT, matching the observed dark sector (§8.3); (5) dark energy evolution with ν_{OI} in the range consistent with DESI data (§8.1); (6) a testable GW echo prediction at the discreteness scale (§8.2); (7) the MOND acceleration scale $a_0 = cH/6$ and the baryonic Tully-Fisher relation, derived from entropy displacement (§8.4). Quantum-first orderings produce the 10^{122} discrepancy, have no natural account of the dark sector, and treat $\hbar$ and the 1/4 factor as unexplained inputs. The cumulative case is not a single tiebreaker but a pattern: every quantitative output of the geometry-first ordering is consistent with observation; the central quantitative output of the quantum-first ordering is the worst prediction in physics.

The classical vacuum energy scale. The framework does not explain *why* $\rho \sim H^2/G$; this is set by initial conditions via the Friedmann equation. What it does is reclassify the CC puzzle: from quantum vacuum cancellation to classical initial conditions. The two problems are not equivalent. The QFT CC problem requires a cancellation between independent contributions — the bare cosmological constant and the zero-point energies of every field — accurate to 122 decimal places, re-tuned at each order in perturbation theory. The initial-conditions question asks why a single parameter (Λ, or equivalently H) takes its observed value; the scale $\rho \sim H^2/G$ is the natural energy density of classical cosmology, not a fine-tuned cancellation between unrelated quantities. The framework eliminates the 122-digit cancellation entirely; the remaining question — why H

takes its value — is a standard cosmological problem addressable by inflationary dynamics or other initial-conditions mechanisms.

Quantum corrections to gravity. The emergent quantum description does feed back into gravitational dynamics through state-level quantities: §8.1 derives dark energy evolution because the emergent vacuum energy depends on H through the hidden sector's volume. What the framework excludes is the zero-point sum as a gravitational source. The zero-point energy is the ground-state eigenvalue of the emergent Hamiltonian $\hat{H}_{\text{eff}}$, which acts on $\mathcal{H}_V$. But gravity is sourced by the classical Hamiltonian H_{tot}, which acts on the full phase space Γ — and $\hat{H}_{\text{eff}}$ and H_{tot} are not the same operator. The eigenvalues of $\hat{H}_{\text{eff}}$ (including the zero-point energy) are properties of the *emergent description*; the energy that H_{tot} assigns to the corresponding classical microstates is the *classical vacuum energy* $\rho \sim H^2/G$. The Equivalence Principle applies to H_{tot}: everything that contributes to H_{tot} gravitates. The zero-point sum does not contribute to H_{tot}; it appears when the classical transition probabilities are reorganized into the Hilbert space formalism (§3.1), not when the classical dynamics is evolved. The structural/volumetric distinction of §5.2 captures this: $\hbar$ (structural, a property of the correspondence) does not gravitate; the vacuum energy (volumetric, state-dependent, a property of H_{tot}) does, but at the classical scale $\rho \sim H^2/G$ rather than $\rho \sim M_{\text{Pl}}^4$.

7.3 The Structural Origin of the 10^{122} Discrepancy

$\rho_{\text{QFT}}/\rho_{\text{classical}} \sim M_{\text{Pl}}^4/(H^2/G) = S_{\text{dS}} = A/(4\,l_p^2)$ — the Bekenstein-Hawking entropy, now identified as the information compression ratio of the trace-out. The "worst prediction in physics" is the entropy of the observer's blind spot — a category error, not a fine-tuning failure. Wolpert's limits of inference [19] confirm this cannot be resolved from within: both geometric and quantum measurements are faithful to their respective descriptions, and no embedded observer can determine which is more fundamental. This is not a prediction awaiting future data — the observed vacuum energy sits at the classical geometric scale $\rho \sim H^2/G$, the value the framework expects.

---

8. PREDICTIONS AND TESTS

8.1 Dark Energy Evolution in Running Vacuum Form

$\hbar$ is H-independent (§5.2), but the emergent vacuum energy is a state-level quantity depending on the hidden sector's volume. Expanding $\Lambda_{\text{eff}}(H)$:

$$\Lambda_{\text{eff}}(H) = \Lambda_0 + \frac{3\nu_{\text{OI}}}{8\pi G}(H^2 - H_0^2) + \mathcal{O}((H^2 - H_0^2)^2)$$

This is the Running Vacuum Model [20, 21]. The S_{dS} horizon modes span frequencies $\omega_{\text{IR}} = H$ to $\omega_{\text{UV}} = c/\epsilon$. For conformal fields, the spectral density is uniform in $\ln\omega$, so the number of independent spectral channels is $\mathcal{N}(H) = \ln(c/\epsilon H)$. Each carries fraction $1/\mathcal{N}$ of the total gravitating vacuum energy, so $\rho_\Lambda \propto 1/\mathcal{N}$. Taylor-expanding:

$$\boxed{\nu_{\text{OI}} = \frac{\Omega_\Lambda}{2\,\ln(c/\epsilon H_0)}}$$

With $\epsilon = 2\,l_p$, $H_0 = 67.4$ km/s/Mpc, $\Omega_\Lambda = 0.685$: $\nu_{\text{OI}} = 2.45 \times 10^{-3}$ (parametric uncertainty

$\pm 0.03 \times 10^{-3}$ from H_0 and Ω_Λ; systematic range ~ 1–5×10^{-3} from the spectral assumption below). Independently testable ratio: $\nu_{\rm OI}/\Omega_\Lambda \approx 0.00358$.

Sensitivity to field content. The conformal assumption (uniform spectral density in $\ln \omega$) is an idealization; massive fields and QCD confinement break conformal invariance. For a spectral density $\propto \omega^\alpha$ with $\alpha \neq 0$, the effective channel count $\mathcal{N}$ is modified. Since $\nu_{\rm OI} \propto 1/\mathcal{N}$ and the conformal value is $\mathcal{N} \sim 140$, even order-unity changes to α shift $\nu_{\rm OI}$ within $\mathcal{O}(10^{-3})$: the logarithmic structure suppresses sensitivity to UV field content. The systematic range 1–5×10^{-3} spans a factor-of-two variation in $\mathcal{N}$ around the conformal value; this is the dominant uncertainty, not the parametric error. The central value 2.45×10^{-3} should be understood as the conformal benchmark, not a precision prediction.

DESI data [22, 30] report evolving dark energy at 2.8σ–4.2σ. RVM fits [31] find $\nu \sim \mathcal{O}(10^{-3})$ with $\nu = 0$ disfavored at 2.7σ–3.1σ, consistent with the prediction.

8.2 Gravitational Wave Echoes

A black hole horizon represents a regime where the causal partition differs from the cosmological one: degrees of freedom reclassify from visible to hidden sector as they cross the horizon, shifting the effective partition and the emergent description.

At proper distance $\sim \epsilon$ outside a black hole horizon r_h, an infalling mode reaches the discreteness floor and must scatter back. The predicted time delay is:

$$\Delta t_{\rm echo} \approx \frac{r_h}{c} \ln\left(\frac{r_h}{2\, l_p}\right)$$

For a $30 M_\odot$ remnant, logarithmic dependence on ϵ makes this robust (~ 0.1 ms shift within the ~ 54 ms timescale [23]). Detection would constitute direct evidence that ϵ has observable consequences at horizons other than the cosmological one — a regime where partition-relativity (§1.4) becomes empirically accessible. Joint confirmation of dark energy evolution *and* GW echoes would uniquely favor an information-theoretic origin, since standard RVMs predict $\Lambda(H)$ running but have no mechanism for discreteness-scale echoes.

8.3 The Dark-Sector Corollary

The trace-out that produces QM has an automatic gravitational consequence.

Corollary (Invisible gravitational budget). *Under Axioms 1–4 and conditions (C1)–(C3), the cosmological trace-out that produces quantum mechanics simultaneously renders ~95% of the universe's gravitational budget invisible to the emergent QFT.*

Proof. Three steps, each using only results already established.

(i) Total effective density. The horizon carries $S_{\rm dS} = A/\epsilon^2$ modes (§6). The classical temperature is $T_{\rm cl} = c^3\epsilon^2 H/(8\pi G k_B)$ (§5.1), computed entirely within the classical substratum with no reference to $\hbar$ or the emergent quantum description. The total thermal energy is $E_s = S_{\rm dS} \cdot k_B T_{\rm cl}$. By the shell theorem, distributing over $V_H = 4\pi c^3/(3H^3)$:

$$\rho_s = \frac{E_s}{V_H} = \frac{A}{\epsilon^2} \cdot \frac{c^3\epsilon^2 H}{8\pi G} \cdot \frac{3H^3}{4\pi c^3} = \frac{3c^2H^2}{8\pi G} = \rho_{\rm crit}$$

This identity, which Verlinde [37] assumes, is here derived.

Four clarifications are essential for what follows.

Classical, not emergent. The calculation above uses only pre-trace-out quantities: the number of boundary modes (A/ϵ^2), the classical temperature ($T_{\rm cl}$), and the Hubble volume (V_H). No step invokes the emergent quantum description. This is what distinguishes ρ_s from the QFT zero-point energy $\rho_{\rm QFT} \sim M_{\rm Pl}^4$: the latter is computed as $\sum \frac{1}{2}\hbar\omega$ over modes of the *emergent* quantum field theory — a quantity that exists only after the trace-out. The framework denies that $\rho_{\rm QFT}$ gravitates (§7.2), because it is an artifact of the emergent description. The boundary entropy's thermal energy, by contrast, is a classical quantity that predates the trace-out and gravitates at the classical level. That both involve $\hbar$ numerically (because $T_{\rm cl} = T_{\rm GH}$) is a consequence of the gap equation's self-consistency, not a dependence on QM. The ratio $\rho_{\rm QFT}/\rho_s \sim S_{\rm dS} \sim 10^{122}$ — the same compression ratio identified in §7.3 — confirming that the two quantities occupy different levels of the framework's hierarchy.

Physical locus. The entropy modes are physically localized on the horizon — they are the hidden-sector boundary degrees of freedom of §5.2. The uniform-medium description is a gravitational equivalence (shell theorem), not a claim about physical delocalization. No entropy occupies the bulk.

Why the equivalence is useful. When matter is present, it locally deforms the horizon geometry, redistributing the boundary entropy. The gravitational effect of this redistribution on bulk observers is most easily computed using the uniform-medium equivalence: the deformed boundary is equivalent to a non-uniform effective bulk source.

Algebraic status. The coincidence $\rho_s = \rho_{\rm crit}$ is an algebraic consequence of the Friedmann equation and the self-consistency relation $\hbar = c^3\epsilon^2/(4G)$; it is not an independent physical input.

(ii) Invisibility. The argument has two logically distinct components.

(ii-a) Construction-level invisibility (from Part I alone). The boundary modes comprising $S_{\rm dS}$ are hidden-sector degrees of freedom — the variables traced out to produce the quantum description. By the trace-out structure, no hidden-sector degree of freedom maps to an operator on $\mathcal{H}_V$. This is a theorem of Part I, independent of any gravitational content: the hidden sector is invisible to the emergent description by construction.

(ii-b) Gravitational consequence (from the ontological ordering of §7.2). The physical import — that the boundary entropy's gravitational influence appears as additional gravity with no source in $\langle \hat{T}_{\mu\nu} \rangle$ — requires the geometry-first ordering established in §7.2: the metric exists at the pre-trace-out level, and the stress-energy sourcing gravity is the classical stress-energy, not the emergent quantum expectation value. Under this ordering, the gravitational interaction between bulk matter and boundary entropy is mediated by the metric at the classical level, modifying the geometry but contributing no term to the emergent $\langle \hat{T}_{\mu\nu} \rangle$. To the embedded observer, this appears as additional gravity with no particle-sector source.

The corollary's predictions are therefore conditional on the same geometry-first ordering that dissolves the CC problem in §7.2. This is not an additional assumption: the ordering is already required by Part II's logic (the partition is defined by null geodesics, so the metric must exist prior to the partition). What the corollary provides is a second, qualitatively distinct *consequence* of that ordering.

(iii) Budget. The emergent QFT accounts for the baryonic sector (~5% of $\rho_{\rm crit}$). The remaining

~95% is boundary entropy — gravitationally active (because ρ_s is a classical quantity computed before the trace-out, unlike $\rho_{\mathrm{QFT}} \sim M_{\mathrm{Pl}}^4$ which is an artifact of the emergent description) but invisible to the emergent QFT. This matches the observed composition, in which ~95% of the gravitational content has no source in particle physics. □

Memory effects and P-indivisibility. Matter locally deforms the horizon geometry, redistributing the boundary entropy. This displacement persists because $\tau_B \gtrsim 1/H$ (condition C2): the hidden-sector modes carrying the redistributed entropy do not thermalize back to the unperturbed configuration on sub-Hubble timescales. Verlinde [37] identifies this as a "memory effect" from the non-thermalization of de Sitter states at sub-Hubble scales. In the present framework, this non-thermalization is not assumed but derived: it is P-indivisibility operating in the gravitational sector, with (C2) ensuring that entropy displacements persist without thermal overwriting.

The corollary is a consistency check for the total dark sector budget, not a derivation of its internal structure. The uniform component of the boundary entropy corresponds to dark energy (§7); the structured component — the matter-induced redistribution of boundary entropy producing local gravitational effects resembling dark matter — is derived in §8.4. The ratio $\rho_{\mathrm{QFT}}/\rho_s \sim S_{\mathrm{dS}} \sim 10^{122}$ confirms the parallel with §7.3: the information compression ratio and the gravitational occlusion fraction are two aspects of a single phenomenon.

Falsifiability. The corollary would be falsified by detection of dark matter particles accounting for the full ~27% structured dark sector, or by evidence that the dark sector's gravitational effects are unrelated to the boundary geometry.

8.4 Dark Matter from Entropy Displacement

The total dark sector budget is established by §8.3. This section derives the structured component from the entropy displacement induced by baryonic matter.

Step 1: Entropy displacement (derived). Baryonic matter M_B is coupled to the boundary entropy through H_{VB} (condition C1). The boundary is a thermal reservoir at $T_{\mathrm{dS}} = \hbar H/(2\pi c k_B)$ (derived, §5.2). For a static matter configuration in gravitational equilibrium — a quasi-static process — the Clausius relation gives the entropy displacement:

$$\Delta S = \frac{M_B c^2}{k_B T_{\mathrm{dS}}} = \frac{2\pi M_B c^3}{\hbar H}$$

This follows from the generalized second law (Theorem B.6) applied to the matter-boundary system: the boundary's entropy reduction is Q/T for a reversible process, where $Q = M_B c^2$ and $T = T_{\mathrm{dS}}$. For dynamic configurations, the Clausius inequality $\Delta S \geq Q/T$ only strengthens the displacement.

Step 2: The critical acceleration (derived). The displaced entropy redistributes over the volume between the matter and the horizon. The Jacobson mechanism converts the entropy gradient into additional radial acceleration. The crossover between Newtonian gravity and the entropic effect, accounting for the volume-to-surface ratio of spherical redistribution in de Sitter (1/6), gives:

$$a_0 = \frac{cH}{6} \approx 1.2 \times 10^{-10}\ \mathrm{m/s}^2$$

This is the MOND critical acceleration — derived, not fitted. (Milgrom's empirical value: $a_0^{\mathrm{obs}} \approx 1.2 \times 10^{-10}$ m/s².)

Step 3: The deep-MOND regime. For baryonic acceleration $\ll a_0$, the displaced entropy dominates. The Jacobson mechanism converts the entropy gradient into curvature, giving $g_{\text{total}} = \sqrt{g_B \cdot a_0}$ and rotation velocity:

$$v^2 = r \cdot g_{\text{total}} = \sqrt{GM_B \cdot cH/6} = \text{constant} \qquad \Longrightarrow \qquad v^4 = GM_B \cdot cH/6$$

This is the baryonic Tully-Fisher relation. Flat rotation curves emerge with no dark matter particles and no free parameters beyond the baryonic mass.

Step 4: Interpolation constraints. The full interpolation between the Newtonian regime ($g_B \gg a_0$) and the deep-MOND regime ($g_B \ll a_0$) is constrained by the framework but not uniquely determined. The entropy displacement defines a potential Ψ_D with $g_D = -\nabla\Psi_D$, and the framework requires:

(i) *No free parameters.* g_D depends only on g_B and $a_0 = cH/6$ — no halo-specific parameters, no fitting constants.

(ii) *Conservative.* $\nabla \times g_D = 0$, since g_D derives from the entropy potential.

(iii) *Monotonic.* $\partial g_D/\partial g_B > 0$ — more baryonic mass produces more entropy displacement, hence more dark matter effect.

(iv) *Correct limits.* $g_D \to 0$ for $g_B \gg a_0$ (Newtonian); $g_D \to \sqrt{g_B \cdot a_0}$ for $g_B \ll a_0$ (MOND).

(v) *Analytic.* The entropy redistribution is a smooth function of the coupling structure — no discontinuities or phase transitions in the interpolation.

These constraints narrow the interpolation to a one-parameter family indexed by the transition steepness — a property of the particular bijection φ, in the same category as fermion masses (Fundamental, §9.7). A representative member:

$$g_D = \frac{a_0}{2}\left(\sqrt{1 + \frac{4g_B}{a_0}} - 1\right)$$

Epistemic status. Steps 1–3 are Tier 2: the displacement formula follows from the Clausius relation applied to the boundary reservoir (§5.2, B.6), the geometric factor from spherical redistribution in de Sitter, and $a_0 = cH/6$ and $v^4 = GM_B \cdot cH/6$ are parameter-free consequences. The interpolation constraints (i)–(v) are Tier 2; the transition steepness is Tier 3 (solution-specific).

Observational predictions. (i) Dark matter "mass" within radius r is determined entirely by enclosed baryonic mass and H — no halo-specific parameters. (ii) The effect depends on $H(z)$: dark matter phenomenology at high redshift differs systematically from low-redshift observations. (iii) The Tully-Fisher relation evolves: $v^4 \propto H(z)$.

Falsifiability. The entropy displacement account would be falsified by: (a) detection of dark matter particles accounting for ~27% of ρ_{crit}; (b) high-redshift rotation curves inconsistent with the predicted $H(z)$ dependence; (c) baryonic Tully-Fisher violations exceeding baryonic mass uncertainties.

---

9. DISCUSSION

9.1 Interpretive Consequences

In the cosmological application, the hidden sector is the trans-horizon region; the degrees of freedom involved in quantum experiments — photons, electrons, slits, detectors — are all visible-sector objects. Their quantum behavior is a downstream consequence of the single cosmological trace-out: the emergent quantum mechanics, once established by the theorem, governs all visible-sector dynamics.

In the double-slit experiment, the particle traverses a single slit in the deterministic substratum. The interference pattern arises because opening or closing the second slit changes the boundary conditions of the transition matrix, altering the distribution of detection events. A which-path detector at one slit couples the trajectory to additional visible-sector degrees of freedom, changing the transition matrix and eliminating the interference terms. In Wigner's friend, the Friend has a definite outcome; Wigner's superposition reflects his epistemic deficit. The Everettian measure problem dissolves: the system evolves as a single reality; "branches" are features of the compressed description. The Born rule, often treated as an independent postulate, is derived: it is the equilibrium distribution of the indivisible stochastic process (§3.1), not an additional assumption.

9.2 Scope

The characterization theorem is a full equivalence: QM $\iff$ embedded observation under (C1)–(C3). The theorem does not identify which physical systems satisfy the conditions (empirical) nor derive $\hbar$ for specific systems (requires Part II). Universality in our universe follows from the shared causal partition defined by null geodesics.

9.3 The Status of the Discreteness Scale

ϵ does not smuggle in a quantum assumption. Axiom 2 requires finite-dimensionality, motivated by the classical fact that finite-area boundaries admit finitely many modes. The result $\epsilon = 2\,l_p$ is a *consequence* of self-consistency (§6). Holographic entropy bounds [27] provide independent, non-quantum motivation.

9.4 Vacuum Energy: Relative Effects and the Higgs Potential

Relative effects. Casimir forces and Lamb shifts are predictions *within* the emergent QFT — relative effects depending on energy differences between configurations. The framework denies that the *absolute* zero-point sum gravitates, because gravity operates at the logically prior classical level (§7.2).

The Higgs potential. The strongest CC objection is not the zero-point sum but the electroweak Higgs potential: symmetry breaking shifts $V(\phi)$ by $\sim$ (200 GeV)^4, exceeding the observed Λ by ~ 55 orders of magnitude. The framework's response follows from its ontological ordering and the structural/volumetric distinction of §5.2, applied as a precise principle:

What gravitates: Relative energy differences between configurations of the emergent fields. An electron at rest vs. no electron differs by $m_e c^2$ in the classical substratum — there is a classical microstate difference corresponding to this energy difference. Relative differences appear in the classical stress-energy tensor and gravitate. Casimir forces, Lamb shifts, and particle masses all fall in this category.

What does not gravitate: Absolute energy levels assigned to configurations by the emergent description. The Higgs vacuum energy shift $V(\phi_{\text{min}})$ is the energy the emergent QFT assigns to its ground state — the configuration obtained after symmetry breaking. But the "ground state" in the emergent description corresponds to a class of classical microstates whose energy in the classical substratum is $\rho \sim H^2/G$, not $\rho \sim (200\text{ GeV})^4$. The shift is a property of how the emergent description organizes the microstates (the correspondence), not a property of the microstates themselves (the classical dynamics).

The principle is uniform: the zero-point sum, the Higgs vacuum energy, and the QCD condensate energy are all absolute properties of the emergent ground state. The dissolution applies to each for the same reason — gravity operates at the classical level (§7.2), where none of these quantities appear.

Why QCD binding energy gravitates but the quantum vacuum does not. The proton mass is ~99% QCD binding energy — a purely quantum effect. If gravity couples to the classical substratum, how does it "know" about masses generated by emergent quantum dynamics? The answer is that QCD binding energy and vacuum zero-point energy occupy different categories in the structural/volumetric distinction of §5.2. Relative energies within the emergent QFT — energy differences between configurations — correspond to differences in the coupling structure of φ, which the Jacobson derivation (Fundamental, §5) reads as curvature. A proton vs. three free quarks corresponds to two distinct classes of classical microstates with different energies in the classical substratum; the binding energy ~ 938 MeV is a *relative* quantity that appears in H_{tot} and gravitates. The absolute baseline — the zero-point energy summed over all modes — corresponds to the total information capacity of the hidden sector (S_{dS}), which is a property of the partition geometry, not of the dynamics, and therefore does not source curvature.

9.5 Relation to Prior Work

't Hooft [1]: Differs in mechanism (epistemic trace-out vs. information loss), generality (any system satisfying axioms vs. particular rules), and Bell placement (outcome independence violation vs. superdeterminism). **QBism/relational QM:** Share observer-relative epistemic states; this framework provides a structural *why* and quantitative predictions. **Jacobson [16]:** Derives gravity from thermodynamics; this framework derives QM from causal structure — complementary, with mutual consistency as a check.

Danielson-Satishchandran-Wald (DSW) horizon decoherence [38, 39]. DSW proved within standard QFT that any Killing horizon imparts a fundamental rate of decoherence on nearby quantum superpositions: the horizon registers the long-range field of the superposed body via soft gravitons/photons, destroying coherence at a rate determined by the horizon geometry. This is precisely the framework's partition mechanism described in QFT language — the horizon is the causal boundary, information leaks through it, and the exterior state decoheres. Three specific features match: (i) the decoherence rate depends on horizon geometry (partition-relativity); (ii) decoherence vanishes for extremal black holes where the coupling to the interior is screened (C1 failure); (iii) the process is non-Markovian near the horizon, with information backflow from the interior [40]. The DSW program derives these results within the emergent QFT; the present framework explains *why* horizons produce decoherence — because the trace-out across a causal partition is the mechanism that generates quantum mechanics itself.

AdS/CFT and the geometry-entanglement connection. The Ryu-Takayanagi formula, Van Raamsdonk's disconnection argument, ER=EPR, and "it from qubit" programs all suggest entan-

glement is prior to geometry. The framework offers a uniform alternative reading: tracing out across a geometric boundary produces entanglement entropy proportional to boundary area (A/ϵ^2 modes), so each of these results is equally consistent with geometry assembling entanglement rather than the reverse. For Ryu-Takayanagi specifically, the proportionality constant must follow from the partition structure — and it does: §6 derives the 1/4 factor. In the holographic context, the boundary CFT observer is *external* to the bulk, immune to the inferential limits of embedded observers [19]; the framework does not claim to reproduce holographic results in that setting.

The point of contact is empirical. Entanglement-first programs predict spacetime breakdown at low entanglement or high energy. This framework predicts QM breakdown near $\epsilon = 2\,l_p$ while the geometric substratum persists. The geometry-first ordering produces seven independent concordances with observation (§7.2); entanglement-first orderings inherit the zero-point sum as a gravitational source and have no natural account of the dark sector.

Witten's Type II algebra program achieves finite entropy without finite-dimensional Hilbert spaces; this is compatible, since the framework's predictions depend on finite entropy (the physical content of Axiom 2), not on dimensional finiteness per se.

9.6 Assumptions, Limitations, and Falsifiability

The theorem requires Axiom 2 (independent in Part I; consequence of geometry in Part II), the stochastic-quantum bridge (§3.1 and Appendix A; established by two independent routes), and genericity conditions (non-degenerate spectrum, non-vanishing overlaps), which hold for all but a measure-zero set. The P-indivisibility proof uses finiteness via the recurrence argument ($\varphi^N = \mathrm{id}$). The deep-sector cardinality corollary (§5.2) establishes that only $\mathcal{C}_V$ and $\mathcal{C}_B$ need be finite — both guaranteed by holographic entropy bounds [27] independently of the de Sitter finite-dimensionality conjecture — while the deep sector $\mathcal{C}_D$ may be infinite. The accessible-timescale backflow lemma (§2.3) provides a second route to P-indivisibility that is independent of $|\mathcal{C}_H|$. The characterization theorem becomes an equivalence on observable timescales ($T_{\mathrm{obs}} \ll \tau_B \sim 10^{17}$ s) — physically indistinguishable from a timeless identity for all experiments within the age of the universe.

The sharp partition and constraint structure. Axiom 3's product decomposition $\Gamma = \Gamma_V \times \Gamma_H$ is the kinematic phase space. In GR, the Hamiltonian and momentum constraints restrict the physical phase space to a submanifold $\Gamma_{\mathrm{phys}} \subset \Gamma_V \times \Gamma_H$ on which interior and exterior data are correlated. P-indivisibility survives on a constraint surface because restoring P-divisibility would require decoupling the sectors entirely, violating the constraints; the constraint surface thus enforces (C1) non-perturbatively. For systems other than the cosmological horizon, the fidelity of the product approximation determines the fidelity of the emergent quantum description.

Planck-scale ordering. The classical-prior ontology may break down at l_p, where "geometric" and "quantum" lose operational distinction. If a deeper theory unifies both, this framework describes the effective macroscopic regime.

Falsifiability. The *theorem* would be falsified by a failure of both the stochastic-quantum correspondence and the Stinespring construction for the class of processes generated here — a possibility excluded by the independent proofs in §3.1 and Appendix A. The *cosmological application* would be falsified by definitive absence of dark energy evolution or GW echoes at the predicted timescale. The *dark-sector concordance* (§8.3) would be falsified by detection of dark matter particles accounting for the full structured dark sector, or by evidence that the dark sector's gravitational effects are unrelated to the cosmological boundary geometry.

Nested and time-dependent partitions. The derivation chain assumes a single partition into visible and hidden sectors. In reality, multiple causal boundaries coexist — most naturally, a black hole horizon inside the cosmological horizon — creating nested partitions with distinct geometries. Appendix B resolves this in three layers.

(i) Consistency of nested trace-outs (Appendix B, Theorems B.1–B.4). Two procedures — tracing out the trans-cosmological region and the black hole interior simultaneously (direct) or sequentially (first cosmological, then black hole) — produce the same CPTP channel, the same emergent Hamiltonian (up to D-gauge), and the same $\hbar$. The key result (Theorem B.2) is that the quantum partial trace within the emergent description agrees with the classical marginalization on the substratum: the emergent QM is self-consistent.

(ii) The generalized second law (Appendix B, Theorem B.6). The sum $S_{\text{matter}} + S_{\text{BH}} + S_{\text{dS}}$ is non-decreasing, proved from strong subadditivity and CPTP monotonicity of the emergent quantum description. Any process that shrinks the black hole (decreasing S_{BH}) transfers information to the visible sector and cosmological boundary, compensating the decrease.

(iii) The Page curve (Appendix B, Theorems B.7–B.9). For an evaporating black hole, the entanglement entropy of the radiation follows $S_{\text{rad}}(t) = \min(n_R(t), n_B(t)) \cdot \ln q$ plus exponentially small corrections, where $n_B(t)$ is the shrinking boundary mode count and $n_R(t) = n_B(0) - n_B(t)$. Unitarity is preserved (bijectivity of φ), the turnover occurs at the Page time $t_P \approx 0.646\, t_{\text{evap}}$ (structural, independent of mass and greybody factors), and the Page value is recovered for all but an exponentially negligible fraction ($\sim e^{-10^{77}}$ for $30M_\odot$) of initial conditions — proved from bijectivity (cycle decomposition on the finite energy shell) and Popescu-Short-Winter typicality [41], without assuming ergodicity. The greybody factor is solution-specific.

Laboratory tests of the characterization theorem may be possible in analogue gravity systems where the hidden sector capacity is tunable.

---

10. CONCLUSION

The general theorem (Part I). Under four axioms and conditions (C1)–(C3), the P-indivisibility of reduced dynamics is proved: any bijective dynamics on a finite system with non-trivial coupling necessarily produces P-indivisible stochastic dynamics. By the stochastic-quantum correspondence, the observer's description is necessarily quantum mechanics. Continuous-time transition data resolves all gauge freedom. The Schrödinger equation, Born rule, and Bell violations are structural consequences. The characterization theorem establishes that the conditions are necessary: QM and embedded observation under (C1)–(C3) are equivalent.

The cosmological application (Part II). The cosmological horizon provides a realization where the sharp-partition approximation is exact and all conditions hold. The theorem yields: (a) $\hbar = c^3\epsilon^2/(4G)$ with $\epsilon = 2\, l_p$ and the Bekenstein-Hawking formula as consequences; (b) dissolution of the CC problem — the 10^{122} discrepancy is S_{dS}, the compression ratio of the observer's quantum description; (c) falsifiable predictions including RVM dark energy with $\nu_{\text{OI}} \approx 2.45 \times 10^{-3}$ and gravitational wave echoes; (d) an automatic dark-sector corollary — the same trace-out renders ~95% of the universe's gravitational budget invisible to the emergent QFT; and (e) dark matter phenomenology from entropy displacement — flat rotation curves, the baryonic Tully-Fisher relation $v^4 = GM_B \cdot cH/6$, and the MOND acceleration scale $a_0 = cH/6 \approx 1.2 \times 10^{-10}$ m/s², all derived from the boundary entropy and the Jacobson mechanism (§8.4).

Nested partitions (Appendix B). When multiple causal horizons coexist, the emergent quantum description is self-consistent: the quantum partial trace within the emergent description agrees with the classical marginalization on the substratum (Theorem B.2). The generalized second law — non-decrease of $S_{\text{matter}} + S_{\text{BH}} + S_{\text{dS}}$ — follows from strong subadditivity and CPTP monotonicity (Theorem B.6). The Page curve is derived: $S_{\text{rad}}(t) = \min(n_R(t), n_B(t)) \cdot \ln q$ with turnover at $t_P \approx 0.646\, t_{\text{evap}}$, from bijectivity of φ (cycle decomposition on the finite energy shell), Popescu-Short-Winter typicality [41], and the derived Hawking temperature (Theorems B.7–B.9). Information is never lost.

The apparent conflict between quantum mechanics and general relativity, the 10^{122} vacuum energy discrepancy, and the dark sector are three faces of the same fact: the information-theoretic cost of embedded observation.

ACKNOWLEDGEMENTS

During the preparation of this work, the author used Claude Opus 4.6 (Anthropic) and Gemini 3.1 Pro (Google) to assist in drafting, refining argumentation, and verifying bibliographic details. The author reviewed and edited all content and takes full responsibility for the publication.

REFERENCES

[1] G. 't Hooft, *The Cellular Automaton Interpretation of Quantum Mechanics* (Springer, 2016). [2] J. Martin, "Everything you always wanted to know about the cosmological constant problem (but were afraid to ask)," *C. R. Phys.* **13**, 566–665 (2012). [3] S. Weinberg, "The cosmological constant problem," *Rev. Mod. Phys.* **61**, 1 (1989). [4] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, 2002). [5] P. Pechukas, "Reduced dynamics need not be completely positive," *Phys. Rev. Lett.* **73**, 1060 (1994). [6] Á. Rivas, S. F. Huelga, and M. B. Plenio, "Entanglement and non-Markovianity of quantum evolutions," *Phys. Rev. Lett.* **105**, 050403 (2010). [7] Á. Rivas, S. F. Huelga, and M. B. Plenio, "Quantum non-Markovianity: characterization, quantification and detection," *Rep. Prog. Phys.* **77**, 094001 (2014). [8] F. A. Pollock, C. Rodríguez-Rosario, T. Frauenheim, M. Paternostro, and K. Modi, "Operational Markov condition for quantum processes," *Phys. Rev. Lett.* **120**, 040405 (2018). [9] P. Strasberg and M. Esposito, "Stochastic thermodynamics in the strong coupling regime: An unambiguous approach based on coarse graining," *Phys. Rev. E* **95**, 062101 (2017). [10] J. A. Barandes, "The Stochastic-Quantum Theorem," arXiv:2309.03085 (2023). [11] J. A. Barandes, "The Stochastic-Quantum Correspondence," *Philosophy of Physics* **3**(1):8 (2025). [12] S. Calvo, "On the Stochastic-Quantum Correspondence," arXiv:2601.18720 (2026). [13] J. S. Bell, "On the Einstein Podolsky Rosen paradox," *Physics Physique Fizika* **1**, 195–200 (1964). [14] J. A. Barandes, S. Hasan, and J. Kagan, "The CHSH Game, Tsirelson's Bound, and Causal Locality," arXiv:2512.18105 (2025). [15] T. Le, F. A. Pollock, T. Paterek, M. Paternostro, and K. Modi, "Divisible quantum dynamics satisfies temporal Tsirelson's bound," *J. Phys. A* **50**, 055302 (2017). [16] T. Jacobson, "Thermodynamics of Spacetime: The Einstein Equation of State," *Phys. Rev. Lett.* **75**, 1260 (1995). [17] G. W. Gibbons and S. W. Hawking, "Cosmological event horizons, thermodynamics, and particle creation," *Phys. Rev. D* **15**, 2738 (1977). [18] S. M. Carroll, "The Cosmological Constant," *Living Rev. Relativ.* **4**, 1 (2001). [19] D. H. Wolpert, "Physical limits of inference," *Physica D* **237**, 1257–1281 (2008). [20] J. Solà Peracaula, "The cosmological constant problem and running vacuum in the

expanding universe," *Phil. Trans. R. Soc. A* **380**, 20210182 (2022). [21] J. Solà Peracaula, A. Gómez-Valent, and J. de Cruz Pérez, "Running vacuum in the Universe," *Universe* **9**, 262 (2023). [22] DESI Collaboration, "DESI 2024 VI: Cosmological Constraints from BAO," arXiv:2404.03002 (2024). [23] J. Abedi, H. Dykaar, and N. Afshordi, "Echoes from the Abyss," *Phys. Rev. D* **96**, 082004 (2017). [24] A. Fine, "Hidden Variables, Joint Probability, and the Bell Inequalities," *Phys. Rev. Lett.* **48**, 291 (1982). [25] L. S. Pimenta, "Divisible and indivisible Stochastic-Quantum dynamics," arXiv:2505.08785 (2025). [26] J. Doukas, "On the emergence of quantum mechanics from stochastic processes," arXiv:2602.22095 (2026). [27] N. Bao, S. M. Carroll, and A. Singh, "The Hilbert space of quantum gravity is locally finite-dimensional," *Int. J. Mod. Phys. D* **26**, 1743013 (2017). [28] W. G. Unruh, "Sonic analogue of black holes and the effects of high frequencies on black hole evaporation," *Phys. Rev. D* **51**, 2827 (1995). [29] K. Fredenhagen and R. Haag, "On the derivation of Hawking radiation associated with the formation of a black hole," *Commun. Math. Phys.* **127**, 273–284 (1990). [30] DESI Collaboration, "DESI DR2 Results II: Measurements of Baryon Acoustic Oscillations and Cosmological Constraints," *Phys. Rev. D* **112**, 083515 (2025); arXiv:2503.14738. [31] J. de Cruz Pérez, A. Gómez-Valent, and J. Solà Peracaula, "Dynamical Dark Energy models in light of the latest observations," arXiv:2512.20616 (2025). [32] W. F. Stinespring, "Positive functions on C-*algebras,"* Proc. Amer. Math. Soc.* **6**, 211–216 (1955). [33] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, 2000). [34] B. Bylicka, M. Johansson, and A. Acín,"Constructive method for detecting the information backflow of non-Markovian dynamics," *Phys. Rev. Lett.* **118**, 120501 (2017). [35] S. Milz, M. S. Kim, F. A. Pollock, and K. Modi, "Completely positive divisibility does not mean Markovianity," *Phys. Rev. Lett.* **123**, 040401 (2019). [36] B. S. Cirel'son, "Quantum generalizations of Bell's inequality," *Lett. Math. Phys.* **4**, 93–100 (1980). [37] E. P. Verlinde, "Emergent Gravity and the Dark Universe," *SciPost Phys.* **2**, 016 (2017); arXiv:1611.02269.

[38] D. L. Danielson, G. Satishchandran, and R. M. Wald, "Black holes decohere quantum superpositions," *Int. J. Mod. Phys. D* **31**, 2241003 (2022).

[39] D. L. Danielson, G. Satishchandran, and R. M. Wald, "Killing horizons decohere quantum superpositions," *Phys. Rev. D* **108**, 025007 (2023).

[40] Y. Li, "Non-Markovian Gravitational Decoherence and the Black Hole Information Paradox," *Int. J. Theor. Phys.* **63**, 293 (2024).

[41] S. Popescu, A. J. Short, and A. Winter, "Entanglement and the foundations of statistical mechanics," *Nature Physics* **2**, 754–758 (2006).

[42] D. N. Page, "Average entropy of a subsystem," *Phys. Rev. Lett.* **71**, 1291 (1993).

APPENDIX A: INDEPENDENT DERIVATION VIA STINESPRING DILATION

This appendix derives the emergent quantum description using only Stinespring's dilation theorem [32] and standard properties of the partial trace [33], without invoking the stochastic-quantum correspondence [10, 11]. The two routes are logically independent; either alone suffices.

A.1 Setup

The finite configuration spaces $\mathcal{C}_V = \{x_1, \dots, x_n\}$ and $\mathcal{C}_H = \{h_1, \dots, h_m\}$ (Axiom 2) embed into Hilbert spaces $\mathcal{H}_V = \mathbb{C}^n$ and $\mathcal{H}_H = \mathbb{C}^m$ via $|i\rangle \leftrightarrow x_i$ and $|k\rangle \leftrightarrow h_k$. This introduces no quantum postulates: it is the canonical identification of probability distributions on a finite set with diagonal density matrices.

A.2 Permutation Unitarity

Lemma A.1. *Any bijection $\varphi : \mathcal{C}_V \times \mathcal{C}_H \to \mathcal{C}_V \times \mathcal{C}_H$ defines a unitary U_φ on $\mathcal{H} = \mathcal{H}_V \otimes \mathcal{H}_H$.*

Proof. Define $U_\varphi|i,k\rangle = |\varphi(x_i, h_k)\rangle$. Since φ is a bijection, U_φ permutes the orthonormal basis, hence is unitary. □

For continuous-time dynamics φ_t, Stone's theorem on $\mathcal{H}$ yields $U_t = e^{-i\hat{H}t}$ for Hermitian $\hat{H}$.

A.3 The Quantum Channel

The observer's ignorance of the hidden sector (Axioms 3–4) corresponds to $\rho_H = I_m/m$. The visible-sector quantum channel is

$$\Phi(\rho_V) = \mathrm{Tr}_H\left[U_\varphi\,(\rho_V \otimes \rho_H)\,U_\varphi^\dagger\right]$$

This is CPTP by a standard result [33, Theorem 8.1], with Kraus representation $\Phi(\rho_V) = \sum_{k,l} K_{kl}\,\rho_V\,K_{kl}^\dagger$ where $K_{kl} = m^{-1/2}\langle l|U_\varphi|k\rangle_H$. The triple $(\mathcal{H}_H, U_\varphi, \rho_H)$ is the Stinespring dilation [32] of Φ.

A.4 Born Rule Recovery

Theorem A.2. *The classical transition probabilities T_{ij} (§1.4) equal the Born-rule probabilities of Φ.*

Proof. $P(j|i) = \langle j|\Phi(|i\rangle\langle i|)|j\rangle = m^{-1}\sum_{k,l}|\langle j,l|U_\varphi|i,k\rangle|^2$. Since U_φ is a permutation, $\langle j,l|U_\varphi|i,k\rangle = \delta_{(j,l),\varphi(i,k)}$. Thus $P(j|i) = m^{-1}\sum_k \delta_{j,\pi_V(\varphi(x_i,h_k))} = T_{ij}$. □

A.5 Emergent Coherence

Theorem A.3. *If T is not a permutation matrix (condition (C1)), then Φ generates genuine quantum coherence: it is not entanglement-breaking.*

Proof. If T is not a permutation, some initial state $|i\rangle$ maps to at least two distinct outputs, so $\Phi(|i\rangle\langle i|)$ has rank ≥ 2. By linearity, Φ maps some off-diagonal input $|i\rangle\langle j|$ to a non-zero operator, precluding the measure-and-prepare form of entanglement-breaking channels. □

A.6 CP-Indivisibility

Theorem A.4. *The P-indivisibility of §2.3 implies CP-indivisibility of $\{\Phi_t\}$: there exist $t_2 > t_1 > 0$ with no CPTP map Λ satisfying $\Phi_{t_2} = \Lambda \circ \Phi_{t_1}$.*

Proof. CP-divisibility restricted to diagonal inputs reduces to P-divisibility. The contrapositive gives the result. □

By the Breuer-Laine-Piilo criterion [4, 6], CP-indivisibility implies non-monotonic trace distance (information backflow) — the quantum signature of non-Markovianity.

A.7 Approximate Unitarity

On observable timescales $t \ll \tau_B$, the hidden-sector state is approximately frozen (conditions (C2)–(C3)). The channel generator decomposes as $d\Phi_t/dt|_{t=0}(\rho_V) = -i[\hat{H}_{\text{eff}}, \rho_V] + \mathcal{D}(\rho_V)$ with $\mathcal{D} \sim \mathcal{O}(\tau_S/\tau_B)$. The leading-order dynamics is the Schrödinger equation. The phase-locking lemma (§3.1) and D-gauge theorem (§5.3) then determine $\hat{H}_{\text{eff}}$ uniquely from continuous-time transition data.

A.8 Comparison of Routes

	Barandes route (§3.1)	Stinespring route (this appendix)
Input	P-indivisible process on $\mathcal{C}_V$	Bijection on $\mathcal{C}_V \times \mathcal{C}_H$ + marginalization
Bridge	Stochastic-quantum correspondence [10, 11]	Stinespring dilation [32] + partial trace [33]
Output	Unitary with $T_{ij} = \|U_{ij}\|^2$	CPTP channel with $T_{ij} = \langle j\|\Phi(\|i\rangle\langle i\|)\|j\rangle$
Born rule	Equilibrium of indivisible process	Partial trace structure (Theorem A.2)
Scope	Any P-indivisible process	Processes from marginalized bijections

The Barandes route is more general (any P-indivisible process); the Stinespring route requires only textbook results [32, 33] and delivers additional structure: the tensor product $\mathcal{H}_V \otimes \mathcal{H}_H$, genuine quantum coherence (Theorem A.3), CP-indivisibility (Theorem A.4), and approximate unitarity with $\mathcal{D} \sim \mathcal{O}(\tau_S/\tau_B) \sim 10^{-32}$ (§A.7). The two routes agree on all observables — both produce the same $T_{ij}(t)$ — establishing that the emergence of QM is robust and overdetermined. The Barandes route powers (ii) $\implies$ (i) in the characterization theorem; the Stinespring route supplies (iii) $\implies$ (i) independently.

APPENDIX B: NESTED PARTITIONS

This appendix proves the consistency of nested trace-outs, derives the generalized second law, and obtains the Page curve for an evaporating black hole. The results resolve all three layers of the nested partition problem stated in §9.6.

B.1 Setup

Let $\mathcal{C}_V$, $\mathcal{C}_B$, $\mathcal{C}_D$ be finite configuration spaces with $|\mathcal{C}_V|, |\mathcal{C}_B| \geq 2$, and let $\varphi : \mathcal{C}_V \times \mathcal{C}_B \times \mathcal{C}_D \to \mathcal{C}_V \times \mathcal{C}_B \times \mathcal{C}_D$ be a bijection. The physical picture: V is the region between a black hole horizon and the cosmological horizon, B is the black hole interior, D is the trans-cosmological region. Define $W = V \cup B$ (the cosmological interior).

Two procedures produce a quantum description on V:

Direct: Trace out $B \cup D$ in one step, marginalizing over $\mathcal{C}_B \times \mathcal{C}_D$.

Sequential: (Stage 1) Trace out D, producing emergent QM on W. (Stage 2) Within the emergent QM on W, trace out B, producing a further-reduced description on V.

B.2 Classical Associativity

Theorem B.1. *The direct and sequential procedures produce the same transition probabilities on $\mathcal{C}_V$.*

Proof. The direct transition matrix is:

$$T_{ij}^{\mathrm{dir}}(t) = \frac{1}{|\mathcal{C}_B||\mathcal{C}_D|} \sum_{b \in \mathcal{C}_B} \sum_{d \in \mathcal{C}_D} \delta_{x_j}[\pi_V(\varphi_t(x_i, b, d))]$$

The sequential procedure marginalizes over D (Stage 1) then over B (Stage 2). Composing:

$$T_{ij}^{\mathrm{seq}}(t) = \frac{1}{|\mathcal{C}_B|} \sum_b \frac{1}{|\mathcal{C}_D|} \sum_d \delta_{x_j}[\pi_V(\varphi_t(x_i, b, d))] = T_{ij}^{\mathrm{dir}}(t)$$

The sums commute (Fubini on a finite product). □

B.3 Quantum-Classical Consistency

Theorem B.2 (Quantum-classical consistency of nested trace-outs). *Let Φ_V^{dir} and Φ_V^{seq} be the CPTP channels on $\mathcal{H}_V$ produced by the direct and sequential procedures respectively. Then $\Phi_V^{dir} = \Phi_V^{seq}$.*

Proof. By the Stinespring construction (Appendix A), φ defines a unitary U_φ on $\mathcal{H} = \mathcal{H}_V \otimes \mathcal{H}_B \otimes \mathcal{H}_D$. The direct channel is $\Phi_V^{\mathrm{dir}}(\rho_V) = \mathrm{Tr}_{BD}[U_\varphi (\rho_V \otimes \rho_B \otimes \rho_D) U_\varphi^\dagger]$. The sequential channel is $\Phi_V^{\mathrm{seq}}(\rho_V) = \mathrm{Tr}_B[\mathrm{Tr}_D[U_\varphi (\rho_V \otimes \rho_B \otimes \rho_D) U_\varphi^\dagger]]$. On finite-dimensional tensor products, $\mathrm{Tr}_B \circ \mathrm{Tr}_D = \mathrm{Tr}_{BD}$, giving $\Phi_V^{\mathrm{seq}} = \Phi_V^{\mathrm{dir}}$. □

B.4 Hamiltonian and $\hbar$ Consistency

Corollary B.3 (Hamiltonian consistency). *The emergent Hamiltonians $\hat{H}_V^{dir}$ and $\hat{H}_V^{seq}$ satisfy $\hat{H}_V^{dir} = D\,\hat{H}_V^{seq}\,D^\dagger$ for a time-independent diagonal unitary D.*

Proof. Theorem B.2 gives $\Phi_V^{\mathrm{dir}} = \Phi_V^{\mathrm{seq}}$, hence $T_{ij}^{\mathrm{dir}}(t) = T_{ij}^{\mathrm{seq}}(t)$ for all t (Theorem A.2). The D-gauge theorem (§5.3) gives the result. □

Corollary B.4 ($\hbar$ universality). *Both procedures yield the same emergent action scale $\hbar = c^3\epsilon^2/(4G)$.*

Proof. $\hbar$ depends only on local boundary quantities c, G, ϵ (§5.2, Steps 2–3), not on which horizon defines the partition or how many trace-outs are performed. □

B.5 Additive Dissipators

Theorem B.5 (Additive dissipators). *Let the classical Hamiltonian be spatially local with coupling chain $V \leftrightarrow B \leftrightarrow D$ (no direct V-D coupling). Let τ_B^D be the D-sector timescale and τ_B^B the B-sector timescale. On observation timescales $t \ll \min(\tau_B^D, \tau_B^B)$, the dissipative correction to unitarity on $\mathcal{H}_V$ decomposes as:*

$$\mathcal{D}_V = \mathcal{D}_V^{(B)} + \mathcal{D}_V^{(D)} + \mathcal{R}$$

where $\|\mathcal{D}_V^{(B)}\| = \mathcal{O}(\tau_S/\tau_B^B)$, $\|\mathcal{D}_V^{(D)}\| = \mathcal{O}(\tau_S/\tau_B^D)$, and the cross term satisfies $\|\mathcal{R}\| = \mathcal{O}(\tau_S^2/(\tau_B^B \cdot \tau_B^D))$.

Proof. The proof applies the boundary-only dependence lemma (§5.2) twice, tracking the cross term at each stage.

(i) *First application: freeze D.* Apply the boundary-only dependence lemma to the partition (W, D) where $W = V \cup B$. By spatial locality, $H_{\text{tot}} = H_V + H_B + H_D + H_{VB} + H_{BD}$ with no direct V-D coupling. The D-sector is frozen on timescales $t \ll \tau_B^D$, giving:

$$T_{ij}^V(t) = \frac{1}{|\mathcal{C}_B|} \sum_{b \in \mathcal{C}_B} \delta_{x_j}[\pi_V(\varphi_t^{VB}(x_i, b))] + \mathcal{O}(t/\tau_B^D)$$

where φ_t^{VB} is the flow restricted to $V \times B$ with D frozen. The $\mathcal{O}(t/\tau_B^D)$ correction arises from D's back-reaction on B through H_{BD}.

(ii) *Dissipator from the frozen-D dynamics.* The frozen-D flow φ_t^{VB} is a bijection on $\mathcal{C}_V \times \mathcal{C}_B$ — exactly the setup of Appendix A with B as the hidden sector. The emergent dynamics on V is approximately unitary with dissipator $\mathcal{D}_V^{(B)} \sim \mathcal{O}(\tau_S/\tau_B^B)$.

(iii) *Decomposition of the D-correction.* The $\mathcal{O}(t/\tau_B^D)$ correction from step (i) has two components:

Direct: D's back-reaction shifts the boundary modes of B by $\delta b \sim \mathcal{O}(t/\tau_B^D)$. This propagates through H_{VB} to the visible sector, contributing $\mathcal{D}_V^{(D)} \sim \mathcal{O}(\tau_S/\tau_B^D)$.

Cross: The shift δb also modifies the effective B-state against which $\mathcal{D}_V^{(B)}$ is computed. The dissipator $\mathcal{D}_V^{(B)}$ depends on the B-state through the Liouville average over b. The D-induced shift changes this average by $\mathcal{O}(t/\tau_B^D)$, so:

$$\|\mathcal{R}\| = \left\| \frac{\partial \mathcal{D}_V^{(B)}}{\partial b} \right\| \cdot \|\delta b\| \leq \|\mathcal{D}_V^{(B)}\| \cdot \mathcal{O}(t/\tau_B^D) = \mathcal{O}\left(\frac{\tau_S}{\tau_B^B} \cdot \frac{\tau_S}{\tau_B^D} \right) = \mathcal{O}\left(\frac{\tau_S^2}{\tau_B^B \cdot \tau_B^D} \right)$$

The inequality uses the fact that $\mathcal{D}_V^{(B)}$ is a continuous function of the B-state distribution and the shift is first-order in t/τ_B^D. □

Bound for the cosmological + stellar black hole case. For a $30M_\odot$ black hole inside the cosmological horizon: $\tau_S \sim 10^{-15}$ s, $\tau_B^B \sim \tau_{\text{scr}} \sim \beta \log S_{\text{BH}} \sim 10^{-3}$ s, $\tau_B^D \sim 1/H \sim 10^{17}$ s. The three terms:

$$\|\mathcal{D}_V^{(B)}\| \sim 10^{-12}, \qquad \|\mathcal{D}_V^{(D)}\| \sim 10^{-32}, \qquad \|\mathcal{R}\| \sim 10^{-44}$$

The cross term is negligible by 12 orders of magnitude relative to the smaller of the two primary dissipators.

B.6 The Generalized Second Law

The Bekenstein-Hawking entropy of a horizon is the information content of the corresponding partition boundary (§6): $S_{\mathrm{BH}} = A_{\mathrm{BH}}/(4\,l_p^2)$ modes at the black hole horizon, $S_{\mathrm{dS}} = A_{\mathrm{dS}}/(4\,l_p^2)$ modes at the cosmological horizon. The matter entropy is the von Neumann entropy of the visible-sector state: $S_{\mathrm{matter}} = -\mathrm{Tr}[\rho_V \log \rho_V]$.

Theorem B.6 (Generalized second law). *For any process described by the CPTP channel Φ_V on $\mathcal{H}_V$:*

$$S_{\mathrm{matter}}(t) + S_{\mathrm{BH}}(t) + S_{\mathrm{dS}}(t) \geq S_{\mathrm{matter}}(0) + S_{\mathrm{BH}}(0) + S_{\mathrm{dS}}(0)$$

Proof. The total system (S, φ) evolves unitarily at the fundamental level, so the total fine-grained entropy $S_{\mathrm{total}} = \log|S|$ is constant. The three terms S_{matter}, S_{BH}, S_{dS} are the observer's ignorance about three sectors: V, B, D respectively.

(i) *Subadditivity.* For the tripartite system $\mathcal{H}_V \otimes \mathcal{H}_B \otimes \mathcal{H}_D$ in a pure state $|\Psi\rangle$ (the total state is deterministic, hence pure conditioned on the initial state), the strong subadditivity of von Neumann entropy [33] gives:

$$S(\rho_{VB}) + S(\rho_{BD}) \geq S(\rho_B) + S(\rho_{VBD})$$

Since $\rho_{VBD} = |\Psi\rangle\langle\Psi|$ is pure, $S(\rho_{VBD}) = 0$. Thus $S(\rho_{VB}) + S(\rho_{BD}) \geq S(\rho_B)$.

(ii) *Identification.* The observer's matter entropy is $S_{\mathrm{matter}} = S(\rho_V)$. By purification on the tripartite system: $S(\rho_V) = S(\rho_{BD})$. The black hole entropy is $S_{\mathrm{BH}} = S(\rho_B)$ restricted to the boundary modes (the number of B-modes coupled across the black hole horizon). The cosmological entropy is $S_{\mathrm{dS}} = S(\rho_D)$ restricted to the boundary modes.

(iii) *Monotonicity and compensation.* The CPTP channel Φ_V is a contraction of relative entropy [33, Theorem 11.9], so $S(\rho_V)$ is non-decreasing under the emergent dynamics. Any process that decreases S_{BH} (shrinking the black hole) transfers information from B to $V \cup D$. By the unitarity of φ, the total mutual information $I(V : B : D)$ is conserved, so the decrease in S_{BH} is compensated by an increase in $S_{\mathrm{matter}} + S_{\mathrm{dS}}$. □

B.7 The Page Curve

Setup. A black hole with initial horizon area A_0 evaporates. The partition boundary at the black hole horizon moves: modes transition from B (interior) to R (radiation, a subsystem of V) as the horizon shrinks. Define $n_B(t) = A_{\mathrm{BH}}(t)/(4\,l_p^2)$ (the number of boundary modes at the BH horizon) and $n_R(t) = n_B(0) - n_B(t)$ (the number of emitted radiation modes). Conservation: $n_B(t) + n_R(t) = n_B(0) = S_{\mathrm{BH}}(0)$.

The trans-cosmological sector D is frozen on evaporation timescales ($\tau_{\mathrm{evap}} \ll \tau_B^D$), so by Theorem B.5 the relevant dynamics is the bipartite system $B \cup R$ with D decoupled at leading order.

Theorem B.7 (Effective bipartite pure state). *On timescales $t \ll \tau_B^D$, the joint state of B and R is effectively pure: $\rho_{BR} = |\psi(t)\rangle\langle\psi(t)| + \mathcal{O}(\tau_S/\tau_B^D)$, where $|\psi(t)\rangle \in \mathcal{H}_B(t) \otimes \mathcal{H}_R(t)$.*

Proof. The total system evolves unitarily under U_φ (Appendix A). The total state $|\Psi\rangle \in \mathcal{H}_V \otimes \mathcal{H}_B \otimes \mathcal{H}_D$ is pure (deterministic evolution from a definite initial state). By the boundary-only dependence lemma, the D-sector is frozen: ρ_D is approximately constant, and $|\Psi\rangle \approx |\psi_{VB}\rangle \otimes |\chi_D\rangle + \mathcal{O}(\tau_S/\tau_B^D)$. Restricting to the $B \cup R$ subsystem (where $R \subset V$ is the radiation): $\rho_{BR} \approx |\psi\rangle\langle\psi|$ is pure at leading order. The correction is $\mathcal{O}(\tau_S/\tau_B^D) \sim 10^{-32}$. □

Theorem B.8 (Cycle typicality). *For the dynamics φ on the energy shell $\Sigma_E \subset \mathcal{H}_B \otimes \mathcal{H}_R$, the time-averaged entanglement entropy of the radiation equals the Page value [42]:*

$$\overline{S(\rho_R)} = S_{\text{Page}}(d_B, d_R) + \mathcal{O}(\epsilon)$$

for all initial conditions except a fraction $\leq 2e^{-c\, d_{\min}\, \epsilon^2}$, where $d_B = |\mathcal{C}_B(t)|$, $d_R = |\mathcal{C}_R(t)|$, $d_{\min} = \min(d_B, d_R)$, and:

$$S_{\text{Page}}(d_B, d_R) = \ln(\min(d_B, d_R)) - \frac{\min(d_B, d_R)}{2\max(d_B, d_R)}$$

Proof. The bijection φ conserves energy (the wave equation is Hamiltonian), so φ restricts to a bijection on each energy shell Σ_E. Since Σ_E is finite, this restricted bijection decomposes into disjoint cycles $C_1, \ldots, C_k$ with lengths $L_1, \ldots, L_k$ summing to $|\Sigma_E|$. On each cycle C_j, the time average of any observable f is exact:

$$\overline{f}_{C_j} = \frac{1}{L_j} \sum_{s \in C_j} f(s)$$

By Popescu-Short-Winter [41], the set of atypical states — those with $|S(\text{Tr}_B|\psi\rangle\langle\psi|) - S_{\text{Page}}| > \epsilon$ — has measure at most $\eta = e^{-c\, d_{\min}\, \epsilon^2}$ on Σ_E, where $d_{\min} = \min(d_B, d_R)$ and $c > 0$ is a universal constant. The total number of atypical states is at most $\eta|\Sigma_E|$.

Call a cycle *bad* if its time-averaged entropy deviates from S_{Page} by more than ϵ. A bad cycle must contain a disproportionate fraction of atypical states. Specifically, if more than half the states in C_j are typical (entropy within ϵ of S_{Page}), then $|\overline{S}_{C_j} - S_{\text{Page}}| \leq \epsilon + \mathcal{O}(\ln d_{\min}/L_j)$. So a bad cycle must have at least half its states atypical: $L_j \leq 2 \cdot |\{s \in C_j : s \text{ atypical}\}|$. Summing over all bad cycles:

$$\sum_{j\,\text{bad}} L_j \leq 2\eta\, |\Sigma_E|$$

The fraction of initial conditions on Σ_E that land on a bad cycle is therefore at most $2\eta = 2e^{-c\, d_{\min}\, \epsilon^2}$. For a $30M_\odot$ black hole, $d_{\min} \sim e^{S_{\text{BH}}} \sim e^{10^{77}}$, so $\eta \sim e^{-10^{77}}$. For all but an exponentially negligible fraction of initial conditions, the time-averaged entanglement entropy equals the Page value. □

Theorem B.9 (The Page curve). *The entanglement entropy of the radiation follows:*

$$S_{\text{rad}}(t) = \begin{cases} n_R(t) \cdot \ln q - \frac{q^{n_R(t)}}{2\, q^{n_B(t)}} & \text{for } n_R(t) < n_B(t) \quad \text{(early: before Page time)} \\ n_B(t) \cdot \ln q - \frac{q^{n_B(t)}}{2\, q^{n_R(t)}} & \text{for } n_R(t) > n_B(t) \quad \text{(late: after Page time)} \end{cases}$$

where q is the alphabet size (gauge — it cancels in all ratios), $n_B(t) + n_R(t) = n_B(0)$, and the Page time t_P is defined by $n_R(t_P) = n_B(t_P) = n_B(0)/2$.

Proof. Direct substitution of $d_B = q^{n_B(t)}$ and $d_R = q^{n_R(t)}$ into the Page entropy formula (Theorem B.8), using $\ln(q^n) = n \ln q$ and the conservation law $n_B + n_R = n_B(0)$. The turnover at $n_R = n_B$ follows from $\min(d_B, d_R)$ switching from d_R to d_B. □

The evaporation trajectory. The Hawking temperature (§5.2, Step 4) gives luminosity $L \sim 1/M^2$, hence $dM/dt = -\alpha/M^2$ where $\alpha = \hbar c^4 \Gamma/(15360\pi G^2)$ and Γ is the species-dependent greybody factor (solution-specific). Integrating:

$$M(t) = M_0\big(1 - t/t_{\text{evap}}\big)^{1/3}, \qquad t_{\text{evap}} = \frac{M_0^3}{3\alpha}$$

The boundary mode count $n_B(t) = 4\pi G M(t)^2/(\hbar c) = n_B(0) \cdot (1 - t/t_{\text{evap}})^{2/3}$ and $n_R(t) = n_B(0) - n_B(t)$. The Page time t_P satisfies $n_B(t_P) = n_B(0)/2$:

$$t_P = t_{\text{evap}}\big(1 - 2^{-3/2}\big) \approx 0.646\, t_{\text{evap}}$$

The entropy of the radiation at the Page time is $S_{\text{rad}}(t_P) = n_B(0) \ln q/2 = S_{\text{BH}}(0) \ln q/2$ — half the initial black hole entropy, as expected.