Contents

Copyright **11**

Preface **12**

Reader's Guide **15**
The complete reading path . . . 15
Reading path for foundational interest . . . 15
Reading path for empirical content evaluation . . . 16
Reading path for philosophical positioning . . . 16
Reading path for biology and medicine . . . 17
Reading path for computational complexity and quantum information 17
How to use the appendices . . . 17
How to use the glossary and bibliography . . . 18
What to skip on a first reading . . . 18
A note on density . . . 19

Acknowledgments **19**

Introduction **22**
0.1 Two physics, or two projections? . . . 22
0.2 An informal thought experiment . . . 24
0.3 The framework's central claim . . . 26
0.4 The intellectual context . . . 27
0.5 The empirical record . . . 28
0.6 The book's structure . . . 30

Chapter 1 **34**

Embedded Observation and the Characterization Theorem **34**
1.1 The problem and the theorem . . . 34
1.2 The starting point . . . 35
1.3 The observer's blind spot . . . 37
1.4 Partition-relativity . . . 40
1.5 P-indivisibility . . . 42
1.6 The two routes to quantum mechanics . . . 47
1.7 Bell-inequality violations . . . 49
1.8 The characterization theorem . . . 50
1.9 What the theorem says . . . 51
1.10 The four structural limits . . . 52
1.11 The card table and the chessboard . . . 58

Chapter 2 **61**

The Substratum **61**
2.1 What this chapter constructs . . . 61

2.2 The substratum as a finite lossless memory . . . 62
2.3 What the substratum is and is not . . . 63
2.4 The reconstruction theorem . . . 65
2.5 The substratum gauge group . . . 71
2.6 The three projections . . . 75
2.7 The QM-GR incompatibility as a category error . . . 78
2.8 What follows . . . 80

Chapter 3 **82**

Hierarchical Structural Realism **82**
3.1 What this chapter develops . . . 82
3.2 The observation axis . . . 83
3.3 The observer-admission distinction in more detail . . . 86
3.4 The gauge axis . . . 88
3.5 Orthogonality, multi-level realism, and stratified predictions . . . 91
3.6 Gauge freedom and observational incompleteness . . . 94
3.7 No-go theorems at different descriptive levels . . . 95
3.8 The philosophical lineage . . . 101
3.9 What follows . . . 106

Chapter 4 **107**

Methodology **107**
4.1 What this chapter develops . . . 107
4.2 The single-foundational-commitment pattern . . . 109
4.3 Classification of claims by derivational rigor . . . 112
4.4 Engaging the no-go results through one architecture . . . 115
4.5 Pre-registered falsification conditions . . . 118
4.6 What does not falsify and regimes of deliberate silence . . . 123
4.7 What the framework derives, and what it compresses . . . 127
4.8 What follows . . . 129

Chapter 5 **130**

The Emergence of Gauge Structure **130**
5.1 What this chapter develops . . . 131
5.2 The minimal substratum and the wave equation . . . 132
5.3 Selection of $d = 3$ . . . 135
5.4 The emergent QFT: factorization and multi-component dynamics . . . 137
5.5 The cubic decomposition and the Standard Model gauge group . . . 139
5.6 Chirality, anomaly cancellation, and hypercharges . . . 142
5.7 What follows . . . 146

Chapter 6 **147**

The Matter Content and Quantitative Predictions **147**

6.1 What this chapter develops 147
6.2 Three generations and the composite Higgs 149
6.3 The strong-CP problem: what time reversal fixes 152
6.4 Gauge couplings and the no-GUT prediction 154
6.5 The Cabibbo angle and CKM structure 157
6.6 The Koide relation and the lepton mass chain 161
6.7 PMNS mixing angles 166
6.8 Summary table and chapter close 170

Chapter 7 **176**

The Gravitational Sector **176**
7.1 What this chapter develops 176
7.2 The cosmological horizon as causal partition 179
7.3 The emergent action scale $\hbar$ 180
7.4 Self-consistency and the discreteness scale 185
7.5 The Bekenstein-Hawking entropy 186
7.6 The Page curve and the information paradox 188
7.7 Dissolution of the cosmological constant problem 194
7.8 Dark energy in running-vacuum form 198
7.9 The dark sector and the MOND acceleration scale 202
7.10 Summary of gravitational predictions and chapter close 206

Chapter 8 **209**

JUNO and the Neutrino Sector **209**
8.1 What this chapter develops 209
8.2 The JUNO experiment 210
8.3 The cubic-group flavor structure 211
8.4 The U-parity grading 214
8.5 The sum rule and structural relation 216
8.6 The three numerical predictions 219
8.7 Comparison with TM1/TM2 column-preservation patterns 221
8.8 JUNO design-lifetime sensitivity and falsification 223
8.9 The coupled DESI DR2 result 224
8.10 Chapter close: the framework's empirical record in the neutrino sector 226

Chapter 9 **228**

Universality Classes and External Convergence **228**
9.1 What this chapter develops 228
9.2 The framework's hierarchical structural realism 230
9.3 Tier 1 results as universality-class content 233
9.4 Tier 2 and Tier 3 stratification 237
9.5 The framework's relationship to string theory 239
9.6 The string landscape multiplicity problem 242

9.6b Foundational programs claiming parameter-free SM retrodictions 243
9.6c A gravitational-sector sibling: conformal gravity 245
9.7 The Danielson-Satishchandran-Wald external convergence 248
9.8 Adjacent observer-emergent literature 250
9.9 Chapter close: the framework's position in the broader landscape . 252

Chapter 10 **254**

Chemistry **254**
10.1 What this chapter develops 255
10.2 The structural chain from substratum to orbital chemistry 256
10.3 Beyond carbon: four additional structural results 259
10.4 The scale hierarchy and the permanence of atoms 261
10.5 The thermal window for chemistry 262
10.6 Water as biological solvent 263
10.7 The chirality chain 265
10.8 Fine-tuning and the anthropic principle 267

Chapter 11 **269**

The Origin of Life **269**
11.1 What this chapter develops 269
11.2 The autocatalytic argument 271
11.3 Prebiotic chemistry pathways 273
11.4 Self-replication as read-write cycling 275
11.5 From template replication to evolution 277
11.6 C1–C4 at the molecular scale 279
11.7 RNA as the unique single-molecule C1–C4 system 281
11.8 What remains genuinely open 283

Chapter 12 **284**

Evolution, Intelligence, and Self-Reference **284**
12.1 What this chapter develops 284
12.2 Evolution as a multi-scale C1–C4 system 286
12.3 Information processing as fitness-enhancing 288
12.4 Major evolutionary transitions 289
12.5 Neural computation as structurally possible 291
12.6 General intelligence as structurally unconstrained but not guaranteed 292
12.7 The chain from intelligence to AI 294
12.8 The self-referential loop and structural limits 296

Chapter 13 **301**

Quantum Biology **301**
13.1 What this chapter develops 301

13.2 The framework's reframing of quantum biology 303
13.3 C1–C4 in proteins 304
13.4 Photosynthesis as energy transfer through C1–C4 architectures . 306
13.5 Magnetoreception via cryptochrome radical pairs 308
13.6 Enzyme catalysis as the framework's clearest test case 309
13.7 The restricted quantum-statistical regime in biology 311
13.8 Implications for drug design 313
13.9 Chapter close 314

Chapter 14 **315**

Quantum Computing **315**
14.1 What this chapter develops 315
14.2 The framework's content in computational complexity 318
14.3 BQP is available: the lower bound 319
14.4 BQP cannot be exceeded: the upper bound 321
14.5 The BQP characterization theorem 324
14.6 The quantum (BQP-form) Extended Church-Turing Thesis as a conditional theorem 326
14.7 Task-specific quantum advantage as cost of partial observation . . 328
14.8 Internal structure of BQP and the P vs NP question 330
14.9 Beyond axioms: research directions opened by the substratum structure 334
14.9.1 Geometric structure on the substratum lattice 334
14.9.2 Task-specific quantum advantage, hidden group structure, and the SU(2)-compatibility refinement 335
14.9.3 Gauge equivalence and complexity-theoretic reductions . . 337
14.9.4 Conditions and the right kind of research question 337
14.10 Chapter close 337

Chapter 15 **341**

Quantum Engineering — Hardware, Software, and the BEC Experiment **341**
15.1 What this chapter develops 341
15.2 The environment as programmable quantum memory 342
15.3 Superconducting qubits and TLS noise 344
15.4 Correlation-aware error correction 346
15.5 Coherence extension through engineered baths 347
15.6 Quantum sensing with NV centers 349
15.7 The restricted quantum-statistical regime in engineered systems . 350
15.8 The BEC analogue-gravity experiment 352
15.9 Chapter close 356

Chapter 16 **358**

Medicine **358**

16.1 What this chapter develops . . . 358
16.2 The enzyme as a read-write system . . . 360
16.3 Cancer pharmacology . . . 362
16.4 Neurodegeneration . . . 363
16.5 Antibiotic resistance . . . 366
16.6 Immunotherapy and T-cell exhaustion . . . 367
16.7 Cardiac pharmacology . . . 368
16.8 Autoimmune disease . . . 369
16.9 Epigenetics as biological hidden sector . . . 370
16.10 The substratum-emergent operator distinction at chromatin . . . 372
16.11 Reader, writer, and eraser pharmacology . . . 374
16.12 The wider therapeutic window for reader-writer disorders . . . 375
16.13 Chapter close: memory asymmetry as unifying principle . . . 377

Chapter 17 **379**

Bioinformatics **379**
17.1 What this chapter develops . . . 380
17.2 The information-theoretic ceiling on transcriptome-only methods 381
17.3 Trajectory inference and the RNA velocity paradigm . . . 383
17.4 Gene regulatory network inference . . . 384
17.5 Perturbation prediction and the failure of scaling . . . 385
17.6 Multi-omics integration and timescale-aware methods . . . 387
17.7 Cancer methylome classification as positive prediction . . . 387
17.8 Variant pathogenicity for reader/writer/eraser genes . . . 389
17.9 Empirical confirmations in evolutionary biology . . . 391
17.10 Chapter close: the structural roadmap forward . . . 394

Chapter 18 **396**

Beyond Quantum Mechanics and General Relativity **396**
18.1 What this chapter develops . . . 397
18.2 The Born rule as dynamical equilibrium . . . 399
18.3 The quantum-classical boundary and the restricted quantum-statistical regime . . . 403
18.4 The epistemic character of quantum randomness . . . 406
18.5 Quantum gravity: the cumulative case at the framework level . . . 408
18.6 Gravitational wave echoes . . . 412
18.7 Finite information and exact finite recurrence . . . 412
18.8 The arrow of time . . . 416
18.9 Cosmic spatial topology . . . 418
18.10 The framework's position on consciousness . . . 421
18.11 Chapter close . . . 426

Chapter 19 **428**

Resolved and Open Problems in Fundamental Physics **428**

19.1 What this chapter develops 428
19.2 Problems resolved 430
19.2.1 The quantum gravity problem 430
19.2.2 The cosmological constant problem 431
19.2.3 The nature of dark matter 432
19.2.4 The nature of dark energy 432
19.2.5 The measurement problem 433
19.2.6 The black hole information paradox 434
19.2.7 The strong CP problem 434
19.2.8 The hierarchy problem 435
19.2.9 The origin of the gauge group 436
19.2.10 The generation puzzle 436
19.2.11 The dark sector budget 437
19.2.12 The origin of the Born rule 437
19.3 Problems open within the framework 440
19.3.1 The flavor problem 440
19.3.2 The Hubble tension 442
19.3.3 Baryogenesis 443
19.3.4 Inflation and initial conditions 445
19.3.5 The initial state of (S, φ) 446
19.3.6 Is the discrete structural residue genuinely non-tautological? 446
19.3.7 Radiative stability of emergent Lorentz invariance (rotational sector) 448
19.3.8 The chirality hypothesis H-χ' 453
19.3.9 Selection of the relative quantum evolution 454
19.4 Chapter close: summary table and what the framework does not address 456

Appendix A **461**

Prediction Status Table **461**
A.1 What this appendix provides 461
A.2 Classification scheme 462
A.3 Four-layer descriptive framing 463
A.4 Master prediction tables 464
A.4.1 Standard Model observables 465
A.4.2 Gravitational-sector observables 467
A.4.3 Neutrino-sector observables 470
A.4.4 Biology and medicine predictions 471
A.5 Forward predictions and falsification targets 473
A.6 Framework's epistemic position summary 477

Appendix B **479**

Mathematical Derivations **479**
B.1 What this appendix develops 479

B.2 The Stinespring construction for emergent quantum mechanics . . 481
B.3 The trace-out as a Jordan-Chevalley projection 487
B.4 The gap equation for $\hbar$ 489
B.5 Anomaly cancellation and unique hypercharges 492
B.6 Gauge-theoretic derivations 495
B.6b Bravais-lattice uniqueness lemma 497
B.6c T_1 generation-assignment uniqueness lemma 499
B.7 The lemma chain for emergent quantum mechanics 502

Appendix C **509**

Common Objections and Framework Responses **509**
C.1 What this appendix develops 509
C.2 Bell's theorem and the framework's relationship to hidden-variable theories 510
C.3 The no-go theorems: PBR, Frauchiger-Renner, Colbeck-Renner, Bong-Wiseman 512
C.4 The stochastic-quantum bridge: Barandes and Stinespring 514
C.5 The necessity direction of the characterization theorem 515
C.6 Finiteness: is Axiom 2 too strong? 517
C.7 Computational implementability and Wolpert's theorem 519
C.8 Relationship to standard interpretations: MWI, Bohmian, QBism 521
C.9 The substratum-emergent operator distinction: substantive or just renaming? 523
C.9.1 Observational incompleteness and the uncertainty principle: substantive or renaming? 524
C.10 Circularity concerns: the gap equation for $\hbar$ 526
C.11 The framework's epistemic status as a structural framework . . . 528
C.11.1 Is the derivation gerrymandered? The selector-motivation audit 529
C.12 Meta-objections: credentials, scope, ambition 530
C.13 Closing remarks 531

Glossary **534**

Bibliography **548**
Framework-internal citations 548
Foundations of quantum mechanics and interpretation 549
Standard Model physics and particle phenomenology 551
Gravitational physics, cosmology, and quantum gravity 552
Chemistry and origin of life 555
Evolution and molecular biology 555
Medicine, pharmacology, and clinical biology 556
Bioinformatics and computational biology 558
Quantum biology and biological information processing 560
Quantum information, computing, and complexity 561

Mathematical foundations and computability 561
Historical and philosophical works 562

The Incompleteness of Observation

A Unified Framework from Quantum Mechanics to Computational Biology

Alex Maybaum

First complete draft

May 2026

Companion technical papers:
The Observational Incompleteness Framework
Zenodo DOI: 10.5281/zenodo.19060318

Contents

Copyright

© 2026 Alex Maybaum

All rights reserved.

This is a working manuscript representing the first complete draft of *The Incompleteness of Observation: A Unified Framework from Quantum Mechanics to Computational Biology*. The framework's authoritative technical content is in the companion papers (Zenodo DOI 10.5281/zenodo.19060318). For framework-internal derivations, the technical papers should be consulted as the primary references.

Citation: Maybaum, A. (2026). *The Incompleteness of Observation: A Unified Framework from Quantum Mechanics to Computational Biology*. Working manuscript.

No part of this work may be reproduced or distributed without the author's permission, with the standard exception for brief quotations in critical reviews and certain other non-commercial uses permitted by copyright law.

Preface

This book argues a heterodox position: quantum mechanics is not fundamental ontology but the exact observer-level representation of finite incomplete observation. At the level of finite observable laws, three descriptions coincide: stochastic laws, finite reversible deterministic realizations, and fixed-basis unitary/Born representations. The equivalence is bidirectional and exact on every finite horizon. History readback supplies the nontrivial content: it forces hidden predictive memory. Indivisibility across the recurrence cycle of a fixed finite reversible representative is a separate fact: any rooted visible map that is not a permutation forces it, Markov laws included. The book develops the consequences of taking that equivalence as a structural fact rather than as an interpretive analogy.

The qualification is equally important. Universal Hilbert-space representability is not by itself a uniqueness theorem for the whole operational theory of quantum mechanics. Finite classical action menus already admit one common reversible dilation; within complex matrix kinematics taken as given, the tensor-product composite included, an operational theory's available instruments on a finite carrier and its composites are the coherent/noncommuting instruments of standard quantum mechanics exactly when five completion conditions hold ([GR §3.3]); the substratum's stated interventions do not supply one of them, full reversible control; whether an extended OI mechanism can supply it, and which of the others the substratum supplies, are open. Bell is a separate constraint: the deterministic completion retains measurement independence but must relinquish ontic parameter independence to reproduce quantum Bell violation, while operational no-signaling is preserved. The manuscript therefore restores the original equivalence claim where it is proved and states the operational completion conditions exactly where they enter, rather than weakening the established stochastic–quantum equivalence or overstating what the substratum is shown to supply.

The framework's value depends on what follows from this reversal — and the book documents exactly that. The cosmological constant discrepancy of 10^{122} orders of magnitude — conventionally the worst quantitative prediction in physics — is read as the compression ratio between two levels of description, conditional on an open coupling step. The Standard Model gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$, the three matter generations, and the anomaly-free hypercharges are reproduced conditionally on a cubic-lattice commitment that is itself empirically selected by the observed SU(3) — selected-and-reproduced rather than derived from nothing. The Bekenstein-Hawking factor of 1/4 is recovered as a consistency relation of the horizon calibration (conditional on the conditions of Chapter 7). Twenty-two Standard Model observables, seven gravitational-sector entries, and seven neutrino-sector entries follow — two of the Standard Model set as parameter-free structural retrodictions and five as conditional structural ones, the remainder as layered, chained, or fitted, each classified explicitly in Appendix A. The JUNO solar mixing angle is matched by a parameter-free retro-

diction — at 0.07σ against the post-JUNO global fit, 0.14σ against the direct measurement; the derivation postdates the measurement — and JUNO's design-lifetime precision is a pre-registered forward test. Hawking's area theorem is confirmed at 99.999% confidence by the LIGO/Virgo/KAGRA observation of GW250114 in September 2025 (the 1/4 entropy coefficient itself has no direct test). The wider therapeutic window for reader-writer disorders is consistent with the Rett-syndrome reversibility of Guy et al. 2007 (a retrodiction, since that result predates the framework).

These are not interpretive achievements; they are quantitative empirical content. The framework is therefore neither a metaphysical reframing of standard physics nor a speculative alternative to it. It is a derivation chain from one foundational commitment to specific quantitative predictions, with substantial empirical content already in place and additional content awaiting near-term experimental tests.

The book is organized as a complete first treatment of the framework's content for readers with a working background in theoretical physics, mathematical physics, or technical biology. The opening four chapters develop the foundational mathematics. The next five chapters develop the framework's content in fundamental physics. The third part traces the emergence cascade from atoms to chemistry to life to intelligence. The fourth part applies the framework to working scientific fields — quantum biology, quantum computing, quantum engineering, medicine, bioinformatics — with substantial empirical predictions in each. The closing chapters develop the framework's forward predictions and its position on the major open problems of fundamental physics. Three appendices provide a prediction status table, mathematical derivations supporting the main text, and a systematic engagement with the most frequent objections to the framework's commitments.

The book is intended to make the framework's structure, derivation chains, and specific predictions available for evaluation by the broader scientific community. The framework's correctness is an empirical question, and the framework has the empirical content to be evaluated. The book's role is to make that content accessible.

A note on prerequisites. The technical chapters of Parts I and II require working familiarity with quantum mechanics, general relativity, and group theory. The applied chapters of Parts III and IV assume the technical content as derived results and add domain-specific background as needed. The Introduction and the closing chapters can be read by anyone with general scientific background; the technical content can be consulted as needed when specific claims become load-bearing for an argument. A glossary at the end of the back matter provides definitions for the framework's distinctive terminology and the major technical terms used throughout.

A note on completeness. The framework is not a theory of everything. Several major problems in fundamental physics are scope limits within the framework's

content: the flavor problem (the specific values of fermion masses and CKM mixing parameters), baryogenesis (the specific baryon-to-photon ratio — solution-specific, downstream of the framework's flavor-sector CP input), inflation, the Hubble tension, and the cosmological initial conditions. Chapter 19 develops the framework's position on these open problems systematically. The framework provides structural answers where it can and identifies its scope limits where it cannot.

A note on terminology. The phrase "the framework's content" appears throughout the book to distinguish the framework's claims from claims made by standard physics or by other unification programs. The phrase is technical: the framework's content is what follows from the framework's commitments through derivation, as distinct from interpretive readings of those derivations or speculative extensions. Where the framework's content overlaps with standard physics (e.g., the Standard Model's gauge structure), the overlap is noted; where the framework's content extends beyond standard physics (e.g., the Born rule's transient violations in the very early universe), the extension is identified.

The framework is the work of one author, but it depends on the work of many. The intellectual debts to the foundations of physics community — to Bell, Wolpert, 't Hooft, Maldacena, Witten, the Wald group, and the broader observer-emergent literature — are substantial and are acknowledged at the appropriate points in the text. The framework's technical methods rely on standard operator algebra (Stinespring, Birkhoff-von Neumann) and on recent stochastic-quantum correspondence work (Barandes). The framework's empirical content depends on the experimental and observational programs of JUNO, LIGO/Virgo/KAGRA, DESI, the Standard Model precision program, and the broader astrophysical and biological communities. The acknowledgments at the back of the front matter name specific contributors; the bibliography at the back of the book provides the full citation record.

The position the book argues will strike many readers as too ambitious. The framework's claim is that quantum mechanics is not fundamental, that a single derivation chain produces specific quantitative predictions across physics, chemistry, biology, and medicine, and that the empirical record already supports the framework's content — with parameter-free retrodictions in fundamental physics (JUNO at 0.07σ; the Standard Model constants) and pre-registered forward predictions awaiting data (DESI Y5, JUNO design-lifetime precision, and others). Each of these claims is heterodox; their conjunction is more so. The book makes the conjunction explicit and supports it with the available evidence. The reader is invited to evaluate the case as it is presented and to judge for themselves.

— Alex Maybaum 2026

Reader's Guide

The book has four parts and three appendices, organized around the framework's derivation chain from foundational commitment to applied predictions. Different readers approach the framework with different interests, technical backgrounds, and goals. This guide suggests reading paths for several common situations.

The complete reading path

Readers approaching the book from start to finish — the framework's intended presentation — will find the technical content of Parts I and II load-bearing for the cascade developments of Parts III and IV. The argument's intellectual coherence depends on the reader having absorbed the central theorem (Chapter 1), the substratum construction (Chapter 2), the structural realism framing (Chapter 3), the methodology (Chapter 4), the Standard Model derivation (Chapters 5-6), and the gravitational sector (Chapter 7) before engaging with the emergence cascade (Chapters 10-12) and the applied chapters (Chapters 13-17). The complete reading path is therefore the recommended path for readers wanting to evaluate the framework's full content.

Estimated reading time for the complete path: approximately 50-80 hours for technical readers, longer for readers consulting the appendices and external references extensively.

Reading path for foundational interest

Readers primarily interested in the framework's foundational claim should focus on the exact equivalence proved for finite observable laws: stochastic description, finite reversible realization, and fixed-basis unitary/Born representation are intertranslatable; C4 identifies the memory-bearing sector, while the full coherent local instrument/composite lift is stated separately rather than folded into the equivalence theorem. The following chapters and sections develop that claim.

Essential. Chapter 0 (the framing). Chapter 1 (the central theorem and characterization theorem). Chapter 2 (the substratum construction). Chapter 7 (the gravitational sector, particularly §§7.3, 7.5, 7.7 on $\hbar$, the Bekenstein-Hawking entropy, and the cosmological constant dissolution). Chapter 8 (JUNO as concrete empirical anchor). Appendix B (mathematical derivations) for technical detail.

Recommended. Chapter 3 (structural realism). Appendix C (engagement with no-go theorems and standard-interpretation objections). Chapter 18 §18.5 (the cumulative case on quantum gravity).

Optional for this path. The applied chapters (13-17) develop empirical content that supports the framework but is not load-bearing for the foundational

claim. Chapter 19 provides systematic positioning against open problems.

This reading path takes approximately 15-25 hours for technical readers.

Reading path for empirical content evaluation

Readers approaching the framework with primary interest in evaluating its empirical record — the JUNO retrodictive match (0.07σ against the post-JUNO global fit, 0.14σ against the direct measurement), the Standard Model retrodictions, the GW250114 area-theorem consistency, the Rett syndrome retrodiction — should focus on the framework's specific quantitative content.

Essential. Chapter 0 (framing, particularly §0.5 on the empirical record). Chapter 6 (twenty-two classified Standard Model predictions). Chapter 7 (gravitational sector predictions, particularly §§7.5, 7.9 on Bekenstein-Hawking and MOND). Chapter 8 (JUNO neutrino sector). Chapter 16 (medical predictions, particularly §16.12 on Rett syndrome). Appendix A (prediction status table for systematic reference).

Recommended. Chapter 5 (gauge structure derivation, the chain that produces the SM predictions). Chapter 9 (universality classes, framing the empirical record). Chapter 18 (forward predictions for upcoming experiments).

Optional. Foundational content in Chapters 1-4 can be consulted when specific empirical claims raise foundational questions. The chemistry and origin-of-life chapters (10-11) are structural rather than empirical.

This reading path takes approximately 10-15 hours for technical readers.

Reading path for philosophical positioning

Readers approaching the framework from philosophy of physics, philosophy of science, or the foundations community should focus on the framework's interpretive structure.

Essential. Chapter 0 (framing). Chapter 3 (hierarchical structural realism). Chapter 4 (methodology and falsification conditions). Chapter 9 (universality classes and convergence with adjacent programs including CLPW, Maldacena, HUZ, DSW). Chapter 18 (forward predictions and the consciousness position). Appendix C (engagement with no-go theorems and standard interpretations).

Recommended. Chapter 1 (central theorem) for the formal grounding of philosophical claims. Chapter 2 (substratum) for the framework's ontological commitments. Chapter 19 (positioning against open problems) for the framework's epistemic scope.

Optional for this path. The empirical chapters (6-8, 13-17) can be consulted when specific predictions become relevant to philosophical evaluation.

This reading path takes approximately 15-25 hours.

Reading path for biology and medicine

Readers approaching the framework from biology, medicine, or applied biological sciences can begin with the applied content and consult the technical chapters as needed.

Essential. Chapter 0 (framing). Chapter 13 (quantum biology). Chapter 16 (medicine, particularly §§16.9-16.13 on epigenetics and the Rett syndrome confirmation). Chapter 17 (bioinformatics, particularly §17.9 on evolutionary biology empirical confirmations).

Recommended. Chapter 10 (chemistry derivation, providing the foundations). Chapter 11 (origin of life). Chapter 12 (evolution and intelligence). Chapter 15 §§15.5-15.7 (the restricted quantum-statistical regime, which connects engineering to biology).

Optional. The fundamental physics chapters (5-9) can be consulted when biological claims rest on physics derivations. The technical foundations (Chapters 1-4) can be consulted for the framework's structural commitments at the molecular scale.

This reading path takes approximately 15-25 hours.

Reading path for computational complexity and quantum information

Readers approaching the framework from computer science, quantum information, or quantum hardware engineering can focus on the computational and engineering content.

Essential. Chapter 0 (framing). Chapter 14 (BQP characterization theorem). Chapter 15 (quantum engineering, including correlation-aware error correction, NV sensing, and the BEC analogue-gravity test). Appendix B §B.2 (Stinespring construction) and §B.7 (lemma chain).

Recommended. Chapter 1 (foundational theorem) for the structural basis of the BQP result. Chapter 9 (universality classes) for the framework's relationship to alternative computational architectures. Appendix C §C.7 (Wolpert's theorem and computational implementability).

Optional. The empirical chapters in fundamental physics (6-9) and in biology and medicine (13, 16-17) provide the framework's broader empirical record but are not load-bearing for the computational content.

This reading path takes approximately 10-15 hours.

How to use the appendices

The three appendices serve different purposes and can be consulted independently.

Appendix A (Prediction Status Table). A systematic reference resource organizing the framework's predictions by classification (Structural / Computational / Linked / Refined / Pending / Mechanism / Empirical), confidence level, current empirical match, and open issues. Use as a lookup table when evaluating specific claims or planning empirical work against the framework. The appendix does not need to be read sequentially.

Appendix B (Mathematical Derivations). Complete derivations for the framework's key technical results: the Stinespring construction, the Jordan-Chevalley projection, the gap equation relating $\hbar$ to the discreteness scale, the anomaly cancellation calculation, the gauge-theoretic derivations, and the lemma chain for the characterization theorem. Use when the main-text derivations are too compressed for the reader's purposes. The appendix is designed for readers requiring full technical content; readers without that requirement can rely on the main-chapter derivations.

Appendix C (Common Objections and Framework Responses). Systematic engagement with the most predictable objections to the framework's commitments: Bell's theorem, no-go theorems (PBR, Frauchiger-Renner, Colbeck-Renner, Bong-Wiseman), the stochastic-quantum bridge, the necessity direction of the characterization theorem, the finiteness axiom, computational implementability, the relationship to standard interpretations (MWI, Bohmian, QBism), the substratum-emergent operator distinction concern, circularity in the gap equation, the framework's epistemic status, and meta-objections about scope and credentials. Use as a reference resource for skeptical readers or for engaging the framework against specific objections.

How to use the glossary and bibliography

Glossary. Located at the end of the back matter. Defines the framework's distinctive terminology (C1–C4, P-indivisibility, substratum, partition-relativity, etc.) and the major technical concepts used throughout the book. Cross-references in glossary entries point to the chapter and section where each concept is most fully developed. Use as a reference when terminology is unfamiliar.

Bibliography. Located at the end of the back matter. Organized by topical area for ease of navigation. Cite using the author-year convention used in the text. The framework's own technical companion papers (available in the Zenodo release at DOI 10.5281/zenodo.19060318) are the primary references for framework-internal derivations.

What to skip on a first reading

Several sections are structurally optional on a first reading, even within the complete reading path.

- **Chapter 4 §4.6** (regimes of deliberate silence) is methodological commentary that can be returned to after the main argument is absorbed.

- **Chapter 9 §§9.5-9.6** (relationship to string theory and the string landscape multiplicity problem) is positioning against the broader physics landscape; readers without interest in string theory can skim.
- **Chapter 12 §12.8** (the self-referential loop) is philosophical commentary; readers without interest in self-reference can skim.
- **Chapter 18 §18.10** (consciousness) is philosophical commentary; readers without interest in consciousness questions can skim.
- **Appendix B §B.3** (Jordan-Chevalley projection) is the most technical of the appendix sections; readers without background in algebraic geometry can rely on the main-text §5.4-5.5 treatment.

These sections add depth to the framework's content but are not load-bearing for the main argument.

A note on density

The framework is the work of a single author developed primarily in technical-paper form. The book represents the integrated treatment for a broader audience, but the technical density varies. Chapters 1-2, 5-7, and the appendices are demanding and reward careful reading. Chapters 3-4, 9, and 18 are philosophical and require attention to the framework's distinctive vocabulary. Chapters 10-12 and 13-17 are accessible to readers with general scientific background and assume the technical content of Parts I and II as derived results.

Readers finding any chapter too dense should not hesitate to skip ahead and return later. The framework's content is sufficiently modular that most chapters can be productively read with limited preceding context, particularly with the glossary as support.

The framework is unfinished in its empirical exposure and carries named open structural questions (Chapter 19 §19.3). The book makes its structural content available for evaluation. The empirical evaluation depends on experiments now being done and on those forthcoming.

Acknowledgments

The framework developed in this book rests on the work of many physicists, mathematicians, biologists, and philosophers whose contributions made the present synthesis possible. The framework's intellectual debts span multiple decades and several scientific communities, and the acknowledgments here cannot be complete. What follows names the most direct contributors and indicates the broader communities whose work has been load-bearing for the framework.

Foundations of physics. The framework's central theorem owes its mathematical apparatus to standard operator-algebraic results going back to the 1950s,

particularly Stinespring's dilation theorem (1955) and the Birkhoff-von Neumann theorem (1946). Recent work on stochastic-quantum correspondence by Jacob Barandes (2023-2024) provided a compression-and-interpretation bridge from deterministic substrata — sharing the dilation bedrock with the framework's internal construction ([Main §3.4]) — to emergent quantum mechanics. The framework's positioning against the no-go theorems engages the work of Bell, Kochen and Specker, Pusey-Barrett-Rudolph, Frauchiger and Renner, Colbeck and Renner, and Bong et al. The framework's anti-Laplacian commitments draw on Wolpert's 1992 work on physical limits of inference. The deeper intellectual lineage extends through Spinoza's attribute theory and Nicholas of Cusa's *docta ignorantia* to contemporary observer-emergent foundational work.

Observer-emergent and horizon decoherence programs. The framework's structural alignment with contemporary observer-emergent foundational work has been a substantial source of intellectual confidence in the framework's commitments. The work of Chandrasekaran, Longo, Penington, and Witten on observer-incorporated entropy in de Sitter space (2022), Maldacena on the role of the observer's clock in cosmological partition functions (2024), Harlow, Usatyuk, and Zhao on observer-dependent Hilbert spaces in closed universes (2025), and Danielson, Satishchandran, and Wald on horizon decoherence (2022) has provided independent corroboration of the framework's universality-class content from directions the framework's development could not have anticipated.

Standard Model and gravitational physics. The framework's quantitative content in fundamental physics rests on the experimental and theoretical programs of the Standard Model precision community, the LIGO/Virgo/KAGRA collaboration (particularly the GW250114 analysis confirming the Bekenstein-Hawking area law), the JUNO collaboration (whose November 2025 measurement of $\sin^2\theta_{12}$ matches the framework's prediction at 0.07σ), the DESI collaboration, the Planck collaboration, and the LUX-ZEPLIN dark matter direct-detection program. The framework's debt to these communities is substantial: without their data, the framework's content would be structural speculation rather than empirically anchored derivation.

Chemistry, origin of life, and biology. The framework's emergence cascade through chemistry to biology draws on Stuart Kauffman's autocatalytic threshold analysis, the prebiotic chemistry programs of the past seven decades (Miller, Urey, Ferris, Damer, Deamer, and others), Cech and Altman's discovery of catalytic RNA, the Soai autocatalytic reaction, and the broader RNA-world literature. The framework's content in evolution draws on Maynard Smith and Szathmáry's analysis of major evolutionary transitions, Bedford and Hartl on molecular clock overdispersion, Tao et al. on universal rate autocorrelation across the tree of life, Wiser, Ribeck, and Lenski on LTEE power-law fitness trajectories, and Eldredge and Gould on punctuated equilibrium.

Medicine, epigenetics, and pharmacology. The framework's clinical content rests on the work of many groups in chromatin biology, particularly Bird, Cech, Lindquist, and the broader Rett syndrome research community (with

the Guy, Bird et al. 2007 paper providing the framework's strongest empirical anchor outside fundamental physics). The framework's pharmacological content engages the work of multiple groups in cancer pharmacology (particularly on Chk1 inhibitors and gemcitabine combinations), neurodegeneration (particularly on CaMKII and PTSD reconsolidation), antibiotic resistance (particularly on RecA and persister cells), immunotherapy (T-cell exhaustion), and the broader pharmacological literature.

Bioinformatics and computational biology. The framework's bioinformatics content engages the substantial recent literature on transcriptome-based foundation models (Csendes et al., Kedzierska et al., and the broader scGPT, Geneformer, and scFoundation programs), single-cell trajectory inference (RNA velocity by La Manno et al., chromatin velocity by Tedesco et al.), gene regulatory network inference (Stolovitzky, Marbach, Bravo González-Blas, and others), and cancer methylome classification (Capper et al. and the DKFZ program).

Quantum information and computing. The framework's content in computational complexity draws on the foundational work of Bernstein, Vazirani, Bennett, Brassard, and the broader BQP literature. The framework's engineering content engages the work of the superconducting qubit community (with Wilen et al., McEwen et al., and Google Quantum AI providing the empirical foundations for correlated noise structure) and the NV-center quantum sensing community. The BEC analogue-gravity experimental program developed by Jeff Steinhauer and collaborators provides the cleanest near-term test of the framework's foundational content.

The framework's working repository. The framework's content has been developed in technical-paper form over the past several years, with the release at DOI 10.5281/zenodo.19060318 representing the current state. The repository's collaborators and reviewers — too numerous to name individually — have contributed substantially to the framework's refinement.

AI assistance in preparing this work is disclosed at
https://github.com/amaybaum/incompleteness#ai-assistance.

Errors. Any remaining errors in the framework's content or its presentation are the author's. The book is the integrated synthesis of work developed in technical papers, and the integration introduces the possibility of inconsistencies between the book's presentation and the underlying technical papers. The technical papers should be consulted as the primary references for framework-internal derivations; the book's role is to make the framework's content accessible.

— Alex Maybaum

Introduction

0.1 Two physics, or two projections?

Physics has two fundamental theories. Quantum mechanics describes the microscopic world — atoms, particles, photons, the interactions of fields at small scales. Quantitative predictions of quantum electrodynamics agree with experiment to twelve decimal places; the Standard Model has been tested at the LHC to energies above 13 TeV without finding a single deviation. General relativity describes the macroscopic world — spacetime curvature, gravitational waves, the expansion of the universe. Its predictions have survived every test from the precession of Mercury to the LIGO observations of binary black-hole mergers. Each theory, within its domain, is the most successful physical theory in human history.

They are incompatible. Quantum mechanics has no preferred temporal foliation; general relativity has one at every point. Quantum mechanics is linear; general relativity is not. Quantum mechanics treats measurement as a primitive operation; general relativity treats it as a process within the dynamics. The two theories give incompatible answers to elementary questions. The most striking of these is the energy of empty space. Quantum field theory predicts a vacuum energy of roughly 10^{113} joules per cubic meter, summed over zero-point modes up to the Planck scale. General relativity, read off the observed expansion of the universe, gives about 6×10^{-10} joules per cubic meter. The ratio is 10^{122}.

For comparison: the observable universe contains about 10^{80} atoms. The discrepancy between what quantum mechanics says the vacuum should weigh and what general relativity measures is forty-two orders of magnitude larger than the number of atoms in the universe. It is the worst quantitative prediction in the history of physics. Physicists have been trying to explain it for fifty years.

The standard reading of this failure — and of related failures that have accumulated over the past several decades — is that the existing theories are incomplete, and that a deeper unified theory is required. The persistent failures are then research opportunities. String theory, loop quantum gravity, causal dynamical triangulations, asymptotic safety, and other unification programs have each pursued the project of finding the missing deeper theory. None has produced a quantitatively confirmed prediction in the regime where it would distinguish itself from existing physics. After a half-century of intensive work, the unification program remains unsettled.

Adjacent to the unification problem, other expected discoveries have not arrived. Supersymmetric particles have not been found at the LHC. Grand unified theories predicting proton decay have not been confirmed by Super-Kamiokande or its successors. Weakly interacting massive particles have not been detected by any direct-detection experiment despite increasingly sensitive searches cul-

minating in LUX-ZEPLIN's 417-day analysis in December 2025. The Hubble tension between early-universe and late-universe measurements of the cosmic expansion rate has grown rather than shrunk. The dark energy equation of state shows evolution that ΛCDM does not predict. Each of these is a specific empirical failure of a specific theoretical expectation, and the cumulative effect is a foundation that has been settling for decades without yet finding its level.

This book argues that the failures share a common diagnostic. The unification problem, the cosmological constant, the dark sector, the failed predictions of supersymmetry and grand unification — these are not separate puzzles awaiting separate solutions. They are symptoms of a single structural error: the assumption that quantum mechanics is fundamental.

The alternative — that quantum mechanics is not fundamental but emergent — has been entertained in foundations of physics for nearly a century. It has rarely been productive. The difficulty is that taking quantum mechanics as emergent has typically required either abandoning Bell-inequality results (superdeterminism), introducing nonlocal mechanisms (Bohmian pilot waves), or proliferating universes (Everett's many-worlds). Each approach has its defenders and its difficulties; none has produced a quantitative empirical record that distinguishes it from the standard reading. The intellectual cost of demoting quantum mechanics from fundamental has typically exceeded the benefits.

The framework developed in this book proposes a different route. It demotes quantum mechanics from fundamental to *emergent representation* — universal at the fixed-basis layer, with the memory-bearing sector carrying the discriminating content — and the framework produces specific quantitative content that the conventional reading does not. The 10^{122} vacuum-energy discrepancy is read as a ratio between two levels of description rather than a fine-tuning failure, conditional on an open coupling step. The Standard Model gauge group is reproduced conditionally on a cubic-lattice commitment whose dimension is selected by the dimensional filters of Chapter 5 within the assumed periodic-lattice class on the stated continuum branch and whose cubic structure is empirically anchored by the observed SU(3) — argued-and-anchored rather than derived from nothing — with the hypercharges then forced by anomaly cancellation on the nonzero-Y_Q branch. The Bekenstein-Hawking factor 1/4 is recovered as a consistency relation between the classical and emergent horizon temperatures, conditional on the horizon conditions of Chapter 7; it has no direct test. The JUNO neutrino mixing angle is matched by a parameter-free retrodiction at 0.07σ — the derivation postdates the November 2025 measurement — whose exact form makes JUNO's design-lifetime precision a pre-registered forward test. Twenty-two Standard Model observables are organized by a single foundational commitment plus cubic-lattice geometry — two as parameter-free structural retrodictions and five as conditional structural ones, the remainder as layered, chained, or fitted, each classified explicitly in Appendix A. The framework's epistemic position is therefore not "another interpretation of quantum mechanics" but a different *kind* of object: a derivation chain from one foundational

commitment to specific quantitative content across multiple sectors of physics.

The foundational commitment is unusual. The framework takes as its starting point not a Lagrangian, not a Hilbert space, not a set of symmetries, but the bare fact that *observation occurs*. From this single premise, together with four conditions on what lies outside the observer's access, two theorems follow: every finite observable law admits a fixed-basis unitary quantum representation, and three of the four conditions (C1, C3, C4) characterize the memory-bearing sector, where hidden predictive memory is unavoidable — with the remaining problem, stated as the frontier, being the additional structural principles that select operational quantum theory and the specifically quadratic Born rule — in finite-resolution approximate form, the exact finite version being closed by a classical-dimension obstruction under the ordinary operational interface ([Main §3.4]) — ([Main §3.2, §3.4]). The chapters that follow develop this identification and trace its consequences.

0.2 An informal thought experiment

Before stating the framework's central claim formally, consider a thought experiment that captures the structural idea.

Imagine a closed room. Inside the room is an experimenter with measuring instruments — clocks, voltmeters, particle detectors. Outside the room is the rest of the universe, but the experimenter has no direct access to it. The walls are opaque to all signals except for whatever couplings the laws of physics enforce: gravitational, electromagnetic, thermal exchanges through the walls. The experimenter wants to characterize the physics inside the room.

The experimenter's situation has three features. First, the room is coupled to the outside. Energy, momentum, and information cross the walls — not at the experimenter's choosing, but because the laws of physics force the coupling. Heat flows through walls. Gravitational fields penetrate them. The experimenter cannot isolate the room from the universe completely. Call this *coupling*.

Second, the outside changes more slowly than the inside. The experimenter can run experiments on millisecond timescales — pulses of light, voltage spikes, particle scattering events. The universe outside changes on the cosmological timescale of billions of years. Whatever is going on outside is, from the experimenter's perspective, essentially frozen during any single experiment. Call this *slow-bath separation*.

Third, the outside is vastly larger than the inside. The room contains perhaps 10^{30} particles. The observable universe contains 10^{80}. The information capacity of what the experimenter cannot access dwarfs the information capacity of what they can. Call this *large capacity*.

These features — coupling, memory persistence, large capacity, and the readback that puts them to use — are the framework's four conditions, abbreviated C1, C2, C3, C4. They are diagnostics of a realization rather than hypotheses of

the central theorem ([Main §1.3]): C1 follows from C4, C3 holds in every faithful realization by the capacity bound, and C2 is a physical-regime premise outside the equivalence. They are not assumptions about exotic physics. They are realization diagnostics of laboratory observers in our universe: at the cosmological cut, coupling is verified structurally, persistence and capacity are verified with enormous margins, and readback is named there and still to be demonstrated; at the lattice cut, coupling is structural, the capacity floor holds for the realized process by data processing, and persistence and readback remain hypotheses (Chapter 1 §1.2, Chapter 7 §7.2).

The framework's central theorem says something more exact. The finite stochastic law available to the experimenter, a finite reversible hidden-state realization, and a fixed-basis unitary/Born description are mathematically equivalent representations of the same finite observable data. If the hidden record is read back, the reduced law is not merely stochastic: every faithful completion must retain predictive memory. Quantum mechanics is therefore an exact observer-level representation of the incomplete finite law, not an extra microscopic postulate placed on top of it.

The thought experiment captures the framework's central move. Quantum mechanics is reframed from a statement about reality to a statement about the observer's relationship to reality. The mathematics is unchanged; what it represents is different. The wave function is not the universe's state; it is the experimenter's compressed description of what they cannot see. The Born rule is not an independent postulate about fundamental indeterminism at the finite observable-law layer; it is the $|U|^2$ readout dictionary of the exact unitary representation every such law admits ([Main §3.4]). The remaining quantum question is the coherent local operational extension of that representation. For Bell experiments the deterministic layer faces Bell's sharper fork: a measurement-independent response-complete completion with one-step own-wing setting interventions, Bell couplings that do not connect the wings before readout and a ± 1 outcome in every run is Bell-local, so quantum violation under measurement independence requires ontic parameter dependence. The framework takes that branch, with operational no-signaling required; the stochastic Tsirelson construction is kept as a separate weaker-description result.

The chapters that follow make this thought experiment precise. The room becomes a partition of the universe into visible and hidden sectors. The experimenter becomes a mathematical observer. The characterization theorem then separates what is universal from what is physically discriminating. The universal part is the exact finite-horizon equivalence $S \iff D \iff Q_{\mathrm{fb}}$; the discriminating part is history readback, which forces hidden predictive memory. The spatial extension adds graph causal cones and a separately stated operational screening condition for Bell experiments. The strongest remaining question is therefore not whether the observer has a quantum representation, but whether the whole coherent instrument/composite hierarchy is one standard local quantum operational theory.

0.3 The framework's central claim

The framework's central claim is an equivalence claim, but it must be stated at the correct level. For every finite observable horizon,

$$S \iff D \iff Q_{\mathrm{fb}} :$$

finite stochastic laws, finite reversible deterministic realizations, and fixed-basis unitary/Born representations are the same class. Thus a finite embedded observer's stochastic description has an exact quantum representation, and every finite quantum record has a finite reversible realization. This is not a limiting metaphor; it is an exact finite theorem.

The representation equivalence is universal, so the distinctly memory-bearing content is carried by history readback. When two visible histories with the same present lead to different future laws, every faithful completion must retain predictive information in the hidden sector; on a fixed finite reversible system, recurrence forces the rooted visible process to be indivisible somewhere in its full cycle whenever any visible step is non-deterministic, so that indivisibility also holds for laws with no accessible memory. The observer therefore sees quantum-representable stochastic dynamics because of incomplete access, while the C1–C4 structure tells us when that description contains genuine hidden memory rather than merely a Markovian stochastic law.

For spatially separated systems, the coupling graph supplies an exact causal cone. Bell adds a sharper deterministic constraint: if each wing's setting intervention, applied at one step, reads and writes only its own region, every run records a ± 1 outcome at each wing, the response-complete pre-setting ontic ensemble is setting-independent, and the wings lie outside each other's graph cone for the duration of the measurement, then Bell factorization follows and $|S_{\mathrm{CHSH}}| \leq 2$. The adopted branch therefore retains measurement independence and uses ontic parameter dependence with operational no-signaling required. The SM/GR target proposes preparation-indexed structure in the state-dependent graph $G(x)$ for it; whether that structure keeps no-signaling and the continuum-curvature limit is the open hypothesis H-Bell. The measurement-dependent horn is recorded separately: 0.0309 bits is a model-independent necessary lower bound from the manuscript's inequality, not an achievable cost. One common standard quantum instrument/composite representation for every coherent intervention at once is characterized, on a finite carrier within complex matrix kinematics taken as given, by five completion conditions ([GR §3.3]); the substratum's stated interventions do not supply one of them, full reversible control; whether an extended OI mechanism can supply it, and which of the others the substratum supplies, are open; none of this weakens the exact stochastic–quantum equivalence already established.

0.4 The intellectual context

The framework's central claim — that the finite observable law of any observer embedded in a finite deterministic universe has an exact fixed-basis quantum representation, Markov laws included, and carries conditional memory at an accessible order exactly when history readback holds there — belongs to a family of impossibility-with-structure results in twentieth-century mathematics and computer science.

In 1931 Kurt Gödel proved that any formal system rich enough to encode arithmetic cannot prove all true statements about itself from within. The limitation arises from self-reference: the system can construct sentences that refer to its own provability, and consistency forces some such sentences to be true but unprovable. The proof yields not chaos but a precisely characterized structure for the unprovable truths.

In 1936 Alan Turing proved that there is no algorithm that can decide, for arbitrary computer programs, whether they halt. The proof again proceeds by self-reference: a hypothetical universal-halting-decider can be turned against itself. The undecidable problems again have a precise structure, captured in the field of computability theory that grew from Turing's work.

In a pair of results (2001, 2008), David Wolpert proved a physical analog of these theorems. Any inference device embedded in the universe it is trying to predict faces fundamental limits. Not from noise, not from finite computational resources, but because complete self-prediction is logically impossible. Wolpert's argument is Gödel's diagonal applied to physical systems.

The framework developed in this book identifies a physical incompleteness that runs parallel to the Wolpert limitation — parallel rather than identical, because Wolpert's is a logical result about worst-case self-prediction, while the incompleteness here is enforced by spacetime geometry and is quantitative. The observer is made of the same fields as the universe they describe; they obey the same speed-of-light constraint; they occupy a bounded region inside the system they are characterizing. The geometry of spacetime enforces a horizon — a boundary beyond which no signal can reach. Everything beyond that horizon must be averaged over in the observer's description of physics. The framework's central theorem is that this averaging, under the actual physical conditions of laboratory observers, produces exactly a fixed-basis unitary/Born representation of what such observers record; whether the full operational formalism of quantum mechanics follows is the open completion question (§0.3). Gödel showed formal systems cannot completely describe themselves; Turing showed computational systems cannot completely predict themselves; Wolpert showed physical inference devices face fundamental limits on self-prediction; this framework adds a physical member to that family: the partition-induced trace-out as the mechanism of an embedded observer's *physical* incompleteness, and quantum mechanics as its specific cost — a construction that stands on its own derivation, which the logical results corroborate but do not supply.

The framework is not the first to suggest that observation cannot be treated as external to physics. QBism, relational quantum mechanics, 't Hooft's cellular-automaton interpretation, and other foundational programs take observer-dependence as a starting point or derive it from specific models. Recent technical work — Chandrasekaran-Longo-Penington-Witten on observer-incorporated entropy in de Sitter space (2022), Maldacena on the role of the observer's clock in cosmological partition functions (2024), Harlow-Usatyuk-Zhao on observer-dependent Hilbert spaces in closed universes (2025) — has converged on the foundational claim that observation matters from independent directions. The framework here converges with these programs on substance: observation matters; the algebra of observables depends on the observer; the partition between observer and observed is physically meaningful. It differs in approach. CLPW, Maldacena, and HUZ add observer degrees of freedom to an existing quantum-gravitational formalism; the framework here goes the opposite direction, starting with a finite deterministic substratum and deriving the quantum representation as a theorem — the operational layer's unique selection stated as open.

This intellectual lineage has older roots. The fifteenth-century cardinal Nicholas of Cusa's *docta ignorantia* (learned ignorance) — the claim that the highest knowledge is knowing what we cannot know — anticipates the framework's structural claim that incompleteness is a feature rather than a deficiency. Cusa's *coincidentia oppositorum* (coincidence of opposites) anticipates the framework's claim that two irreconcilable descriptions can both be correct because they are projections of the same underlying reality. Spinoza's attribute theory — one substance expressed through multiple incommensurable attributes — provides the deepest ontological parallel to the framework's substratum-and-projections picture. A contemporary touchstone is Deutsch's *Beginning of Infinity* [Deutsch], which argues that knowledge itself is structurally incomplete — there is always more explanatory work to be done, and the right epistemic stance toward any current best explanation is provisional but committed. The framework's observational incompleteness provides one physics-foundational grounding for Deutsch's epistemological claim: knowledge derived from observation inherits observational limits as its lower bound (developed in Chapter 12 §12.8). The framework's full philosophical positioning is developed in Chapter 3.

0.5 The empirical record

The framework's claims are unusual; the evidence for them is specific. The empirical record consists of three components: retrodictions of known parameters — including the framework's sharpest matches, some against measurements that predate the framework's development — and pre-registered predictions awaiting upcoming data.

The retrodictions are quantitative. The Cabibbo angle of CKM mixing is predicted to be $1/(\pi\sqrt{2}) = 0.2251$ from a single Brillouin-zone distance on the substratum lattice; the observed value is 0.22500 ± 0.00067, a 0.04% match. The

Koide relation among the three charged-lepton masses is predicted to be exactly 2/3 from a cubic-group quadratic Casimir; the observed value is 0.666664 ± 0.000005, a match to better than 0.001% (0.4σ). The three Standard Model gauge couplings at the Z pole are reproduced to better than 0.1% from one universal lattice-determined coupling at the Planck scale, a threshold fixed by one of the couplings and two threshold parameters matched to the other two, plus standard renormalization-group running. Six fermion masses are reproduced from one empirical input to within 1.3%. The Higgs quartic coupling derived from cubic-group representation theory matches the LHC value at 0.6σ. Twenty-two Standard Model observables in total — gauge couplings, mixing angles, mass ratios — are classified in Appendix A: two as parameter-free structural retrodictions, five as conditional structural ones (the Cabibbo angle with its two inheritors, conditional on the emergent dressing of its identification, and the two Koide entries, conditional on the dynamical closure of the amplitude), the remainder as layered, chained, or fitted, with no parameters fit to the quantities in the structural and layered classes. Each carries the status of its own derivation chain: the gauge sector under H-link (the hypercharges then following from the gauge group and anomaly cancellation on the nonzero-Y_Q branch), the matter sector's physical reading under H-spin', and the chirality assignment under H-χ' (Chapter 19's ledger records every link).

One retrodiction deserves separate note for its precision, with its provenance stated plainly: the measurement predates the framework's 2026 development, so the match is retrodictive, not a forward prediction. In November 2025 the JUNO collaboration reported the first world-leading direct measurement of the solar neutrino mixing angle: $\sin^2\theta_{12} = 0.3092 \pm 0.0087$. The framework's prediction — derived from the substratum's cubic-group structure with no fit parameters — is $1/3 - 1/(4\pi^2) = 0.3080$. The direct measurement matches at 0.14σ; against the post-JUNO global fit by Capozzi et al., the prediction matches at 0.07σ.

In September 2025 the LIGO collaboration's analysis of GW250114, a high-signal-to-noise binary black-hole merger, confirmed Hawking's area theorem — the statement that the total horizon area does not decrease — at 99.999% confidence. The observation compares horizon areas and does not measure entropy, so it does not test the 1/4 coefficient, which remains without a direct empirical test. The framework recovers the 1/4, which the standard account fixes from the Hawking temperature and the first law, as the ratio $2\pi/8\pi$ of its horizon calibration, under the horizon conditions named in Chapter 7.

A second component of the framework's gravitational predictions concerns the dark sector. The framework predicts that the entropy of the cosmological horizon, distributed over the Hubble volume, produces a gravitational contribution equal to the critical density of the universe — the observed $\Omega_{\text{total}} \approx 1$. It further predicts that the structured component of this entropy (dark matter) follows the MOND phenomenology with acceleration scale $a_0 = cH/6$ — about 1.1×10^{-10} m/s^2, within the ten-percent stellar mass-to-light systematic of the observed value — and the baryonic Tully-Fisher relation $v^4 = GM_B \cdot cH/6$,

with the dimensional factor 1/6 taken from Verlinde's emergent-gravity analysis. Under this law rotation curves approach flat beyond a crossover radius, which at a fixed baryonic mass profile shrinks with redshift, so the declining rotation curves recently observed in the Milky Way at galactocentric distances > 19 kpc (Jiao et al. 2023) and in high-redshift galaxies (Genzel et al. 2017) are in tension with it, pending comparisons that model the observed galaxies' own baryon profiles (Chapter 7). Direct dark-matter searches continue to return null results, consistent with the framework's structural prediction that no particle dark matter exists.

Several predictions await imminent data. DESI Year 5 will measure the dark energy equation of state; the framework predicts $\nu \approx 10^{-32}$, observationally indistinguishable from ΛCDM. The Bullet Cluster's post-collision relaxation is predicted to show gradual realignment of the dark gravity toward the gas distribution over timescales comparable to H^{-1}. Reliable ALMA gas-mass measurements at $z > 1$ will provide the definitive test of the baryonic Tully-Fisher evolution. The framework's chromatin-condensate predictions in Rett syndrome and triple-negative breast cancer (Chapter 16) provide near-term clinical-trial-scale tests outside fundamental physics.

The framework's structural foundation has also received independent corroboration from the open-quantum-systems literature. Brandner's 2025 theorems on integrating out hidden degrees of freedom in autonomous linear systems establish at theorem level the kind of non-Markovian visible dynamics that the framework requires, derived independently for general open systems with no stake in the framework being correct. The mathematical apparatus the framework relies on is therefore the standard way physicists now think about open systems with hidden structure.

No competing framework produces this collection of results from a single starting definition. The 10^{122} vacuum-energy discrepancy as an information compression ratio (conditional on an open coupling step), the $\sim 95\%$ dark sector as a gravitational occlusion fraction, the Standard Model retrodiction package (two parameter-free structural and five conditional structural entries among twenty-two classified observables), the Bekenstein-Hawking 1/4 factor as a consistency relation under the horizon calibration, the MOND scale $a_0 = cH/6$ as a conditional relation with an imported dimensional factor, and the JUNO $\sin^2\theta_{12}$ retrodiction at 0.07σ — follow from the single statement that observation occurs, the four conditions on the partition, and the named hypotheses of each chapter.

0.6 The book's structure

The book has four parts.

Part I — Foundations. Chapters 1–4 develop the framework's mathematical core. Chapter 1 proves the central theorem: under the four conditions C1–C4, any embedded observer's reduced description is non-Markovian on accessible

horizons, with a fixed-basis unitary quantum representation internal and universal — the finite-horizon equivalence $S \iff D \iff Q_{\mathrm{fb}}$ ([Main §3.4]) — the compressed form supplied across the correspondence's class under (T), nontrivially where the dynamics is P-indivisible. The proof is constructive (the Barandes correspondence and the Stinespring dilation provide explicit constructions) and includes the characterization theorem establishing necessity. Chapter 2 constructs the substratum — the finite deterministic bijection (S, φ) — and proves a **scoped reconstruction theorem**: under the stated empirical/structural inputs, M1-T, C2-mixing and the nearest-neighbor restriction, the local propagating lattice/gauge residue is fixed. M1-B and H-Bell separately condition Bell-inclusive existence, whose uniqueness remains open. Chapter 3 develops the framework's hierarchical structural realism, including its relation to Bell, Kochen-Specker and PBR-type no-go theorems through separate checks of their local-response, operational-algebra and preparation premises. Chapter 4 articulates the framework's methodology — the single-foundational-commitment derivation pattern, the architecture-level engagement with the no-go results, the classification of claims by derivational rigor.

Part II — Physics. Chapters 5–9 develop the framework's quantitative predictions in fundamental physics. Chapter 5 derives the Standard Model gauge group from the cubic-lattice substratum conditionally on H-link: the wave equation, the link count giving $K = 2dm$ with $m = 1$ (coupling-degree minimization does not establish it), the cubic-group decomposition with multiplicities $(3, 2, 1)$, the gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ from the bicommutant of the cubic-group action, chirality from the trace-out, and anomaly cancellation forcing the observed hypercharges on the nonzero-Y_Q branch. Chapter 6 develops the matter content and quantitative predictions: three fermion generations from staggered taste structure (physical reading conditional on H-spin'; chirality on H-χ'), one composite Higgs, what T-invariance of the substratum fixes about strong CP ($\theta \in \{0, \pi\}$ and reciprocity, $\bar{\theta}$ open), the gauge-coupling derivation with the structural no-GUT prediction, and the twenty-two classified retrodictions (two parameter-free structural and five conditional structural; the remainder layered, chained or fitted) including the Cabibbo angle, the Koide relation, the six fermion masses from one empirical input, and all three PMNS angles. Chapter 7 develops the gravitational sector: the Bekenstein-Hawking 1/4 factor and the calibration $\hbar = c^3\epsilon^2/(4G\sigma)$ as consistency relations of horizon thermodynamics under named horizon conditions, the dissolution of the cosmological constant problem at the descriptive level. Chapter 8 develops the JUNO neutrino prediction in detail. Chapter 9 discusses universality, the relationship to other unification programs, and the framework's empirical scope.

Part III — Emergence. Chapters 10–12 trace the structural cascade from the substratum to chemistry, biology, evolution, and intelligence. Chapter 10 derives chemistry from substratum-level electromagnetic structure. Chapter 11 develops the origin-of-life problem in terms of the framework's structural conditions for autocatalysis and template replication. Chapter 12 traces evolution to neural computation to semiconductors to general-purpose computation, identi-

fying the contingent and structural steps along the cascade.

Part IV — Applications. Chapters 13–19 develop the framework's reach into working scientific fields. Chapter 13 addresses quantum biology: non-Markovian enzyme kinetics in proteins, photosynthesis as energy transfer through C1–C4 architectures, magnetoreception via cryptochrome radical-pair dynamics, and enzyme catalysis with conformational landscapes acting as τ_B-scale memory. Chapter 14 develops quantum computing: BQP is identified, under Chapter 14's explicit model hypotheses, as the computational class accessible to embedded observers satisfying C1–C4, with the lower bound (the computational-control model supplies the BQP toolkit on Main's representation layer) and upper bound (informational completeness relative to the model's admissible instruments — operational closure, an explicit hypothesis) jointly characterizing the class under those hypotheses. The quantum (BQP-form) Extended Church-Turing Thesis is reframed from empirical conjecture to conditional theorem of the framework together with that model under a global-coverage premise (all physically realizable computation falling under the model), any super-BQP demonstration falsifying coverage first, the conjunction and its bridge only with further argument. Chapter 15 develops quantum engineering: superconducting qubit dynamics as the canonical engineered C1–C4 system with $\tau_S/\tau_B \sim 10^{-3}$, memory-bearing noise structure producing correlated errors over $\sim 10^3$ gates, correlation-aware decoders correlated errors that a correlation-aware decoder can exploit, coherence revivals from engineered baths, and information recovery in quantum sensing — structural effects whose magnitudes the framework does not quantify, and the BEC analogue-gravity experiment as the cleanest near-term test of the proposed capacity-controlled memory mechanism — a proposal-level prediction, not a test of the characterization theorems. Chapter 16 develops medical applications across cancer pharmacology, neurodegeneration, antibiotic resistance, immunotherapy, cardiac pharmacology, and autoimmune disease, with an explicit second half on epigenetics as biological hidden sector — the substratum-emergent operator dissociation at chromatin with reader/writer/eraser pharmacology axis, epigenetic drugs as memory erasers, cellular reprogramming as memory reset, the memory hierarchy from DNA methylation through chromatin architecture, and the wider therapeutic window claim for reader-writer disorders, consistent, as a retrodiction, with the Rett-syndrome reversibility of Guy et al. 2007. Chapter 17 develops the bioinformatics consequences of the framework's structural constraints on biological information processing, with empirical confirmations from single-cell methodology and from the evolutionary-biology literature (molecular clock overdispersion, rate autocorrelation across the tree of life, LTEE power-law fitness trajectory). Chapter 18 develops the framework's distinctive forward content: Born-rule transient violations (a conditional prospect for CMB tests of the primordial power spectrum and non-Gaussianity), the quantum-classical boundary with observable signatures, the epistemic character of quantum randomness, the cumulative case on quantum gravity (combining the CC dissolution, the BH information account and the non-renormalizability reading as three components

of one framework-level reading, with covariant unification open), the absence of an OI-specific gravitational-wave echo for external observers, the arrow of time as emergent from partition initial conditions, the framework's silence on global cosmic spatial topology with engagement of Levin's compact-universe proposal, and a substantive position on consciousness — the framework's mathematical operation of observation (partition with C1–C4 structure) is a necessary but not sufficient condition for consciousness, with the hard problem proper explicitly outside the framework's scope. Chapter 19 closes the book with a systematic inventory of the framework's content against the major open problems of fundamental physics — ten problems the framework addresses with a resolution, dissolution, conditional construction or partial treatment, Chapter 19's table giving each one's status (quantum gravity, covariant unification open, the cosmological constant, dark matter, dark energy, the measurement problem, black hole information, hierarchy, gauge group origin, generation puzzle, dark sector budget), two split-status entries (the Born-rule origin — fixed-basis representation settled exactly, coherent operational extension characterized by five named conditions; strong CP — open under two named conditions), and five scope limits where the framework does not provide a derived value (the flavor problem; baryogenesis — solution-specific, the baryon asymmetry inheriting the input status of the framework's flavor-sector CP phases; inflation and initial conditions; the Hubble tension; the cosmological initial state).

The book's technical density varies. Chapters 1, 2, 5, 6, and 7 contain the framework's core mathematical content and are demanding; they require working familiarity with quantum mechanics, general relativity, and group theory. Chapters 3, 4, and 9 are more accessible — philosophical and methodological, with formal content present but not load-bearing. Chapters 10–19 are accessible to readers with general scientific background and assume the technical content of Parts I and II as derived results rather than as content to be re-developed.

Readers approaching the book with different interests will find different reading paths useful. Readers primarily interested in the framework's foundational claim should focus on Chapters 1, 2, and 7; the JUNO prediction in Chapter 8 provides the most concrete empirical anchor. Readers primarily interested in the philosophical positioning should focus on Chapters 3, 4, and 9, with Chapter 18 providing substantive content on consciousness and cosmic-scale questions. Readers approaching the framework from computational complexity or quantum hardware engineering can begin with Chapters 14 and 15 and consult Parts I and II as needed. Readers approaching the framework from biology, medicine, or applied physics can begin with Chapters 12, 13, 16, and 17 and consult Parts I and II as needed. Readers wanting a systematic overview of the framework's content against the standard open problems of fundamental physics should consult Chapter 19 as a reference resource. A reader following the full argument from start to finish — the framework's intended presentation — will find the technical content of Parts I and II load-bearing for the cascade developments of Parts III and IV.

The book makes one foundational commitment — that observation occurs — and develops the consequences. The consequences are extensive. They include the structure of quantum mechanics, the dynamics of general relativity, the gauge group and matter content of the Standard Model, the composition of the cosmological dark sector, the structural conditions for chemistry and life, and the framework's reach into specific contemporary problems in biology and medicine. Whether the framework is correct is an empirical question that will be settled by the predictions developed throughout the book against measurements now being made and forthcoming. The book's role is to make the structure of the framework, the chain of its derivations, and the specific content of its predictions available for evaluation.

Chapter 1

Embedded Observation and the Characterization Theorem

1.1 The problem and the theorem

An observer inside a physical system cannot condition on the complete state of that system. The central question is therefore not whether incomplete observation produces probabilities — it obviously can — but what class of laws the observer's reduced description belongs to.

The answer has two parts. First, at every finite observable horizon there is an exact representation equivalence:

$$S \iff D \iff Q_{\mathrm{fb}}.$$

A finite stochastic law, a finite reversible hidden-state realization, and a fixed-basis unitary/Born representation are three descriptions of the same finite observable object. Conversely, every finite quantum record has a finite reversible realization. Second, history readback identifies the nontrivial sector: it forces hidden predictive memory in every faithful completion.

This chapter proves those statements, then adds their spatial scope. The coupling graph of the underlying dynamics gives exact causal cones. At the response-complete deterministic level, setting interventions, applied at one step, that each read and write only their own wing, readouts outside the other wing's graph cone, a ± 1 outcome recorded in every run, and one setting-independent pre-setting ensemble give a Bell-local model and $|S_{\mathrm{CHSH}}| \leq 2$. The adopted quantum-Bell branch therefore retains measurement independence and relinquishes ontic parameter independence, with operational no-signaling required. The one boundary not folded into the theorem is the full

coherent instrument/composite category of operational quantum mechanics. Keeping that boundary explicit lets the chapter recover the original equivalence claim without confusing a universal representation theorem with a uniqueness theorem.

1.2 The starting point

The framework begins from a starting point usually put in two words: *observation occurs*. An observer records distinguishable outcomes of interactions with a system not wholly under its own control. This is the *cogito* of Descartes made mathematical, without its thinking subject — the framework's one concession to a foundation outside its own theorems.

That starting point, examined closely, is two things. What genuinely cannot be doubted is thin: there is differentiated content — some this-rather-than-that. Call this the first axiom. It is genuinely indubitable, because to doubt it is already to tell one thing from another, which is an instance of it. But notice what a single such fact does *not* contain: time. A single differentiated content is like a photograph — a frozen frame can be full of difference, light here and dark there, without anything in it being before or after anything else. To get from the photograph to a *movie* — to a world with succession, change, one moment after another — something must be added: that the differentiation *recurs*, that it is instantiated more than once. Call this the second axiom. It cannot be squeezed out of the first; a frozen frame, however rich, never amounts to a sequence of frames. The framework therefore rests not on one fact but on two: one that cannot be doubted, and one that must be assumed. This is a strength, not a hedge. A foundation that can point to its single assumed ingredient — and that comes with a theorem showing the assumed ingredient genuinely cannot be derived from the indubitable one — is sturdier than a foundation whose assumptions are scattered, unnamed, through a definition.

One qualification, stated here and developed in the companion paper. Counting the framework's foundational commitments is partly a matter of how the foundation is *presented* rather than a fact about the world: the same theory can be axiomatized in inter-derivable ways, and which structural feature is named an "axiom" and which is derived as a consequence is, within limits, a presentational choice — a notational variance, in Quine's sense, that leaves all observable physics untouched. In particular, whether the embedded observer's status as a *proper part* of a larger whole follows from the first axiom or is a further foundational posit in its own right turns on how the first axiom is read. The framework adopts the reading on which the registering occurs from a locus against a remainder, so that embeddedness follows; on a thinner reading it would be a third posit. The choice bears on how the framework counts its assumptions, not on any prediction it makes. The technical development of the foundational structure, and the no-go theorem behind it, is given in Part I of the companion paper *Physics Modulo Gauge*; the chapters that follow build on the structure without further reference to it.

Definition. An *observation* is a triple (S, φ, V): a total system S, a deterministic dynamics $\varphi : S \to S$, and an observer $V \subsetneq S$ — a proper subsystem with finitely many distinguishable internal states, coupled to the complement $H = S \setminus V$ through φ.

The definition is weaker than classical mechanics. A shuffled deck of cards satisfies it. A finite cellular automaton satisfies it. So does a detector or a protein interior: nothing in the definition requires that anyone be aware of an outcome. What cannot be doubted is that *some* differentiated content is registered — that much is secured from within whatever locus raises the doubt. That a particular deck, automaton or detector registers anything is not indubitable but a physical fact, and the definition covers it all the same. Three lemmas extract its structural content. The map φ is nothing unusual, and its form need not even be assumed. It follows from one innocuous premise — that evolution *composes*: advancing the state by one interval and then another equals advancing it by the combined interval. That single consistency condition (Cauchy's functional equation) forces evolution into a one-parameter family generated by a fixed rule (Engel & Nagel, 2000); in continuous time this is the *phase flow* of classical mechanics (Arnold, 1989), and in the discrete case here it is simply the repeated application of one transition map. So the state-transition picture is not a special assumption but a consequence of time evolution being consistent. What *is* distinctive here is not that such a map exists, but finiteness: the observer distinguishes finitely many visible configurations (Lemma 1), and the map acts on a finite system under the finiteness posit A1 (Chapter 2 §2.4); finiteness, not the dynamics, is the load-bearing commitment.

Lemma 1 (Finiteness). *The visible configuration space $\mathcal{C}_V$ is finite, with a discreteness scale ϵ providing a finite minimal cell volume.*

Any observer bounded by a finite-area surface couples to only finitely many modes across that boundary; independent support comes from holographic entropy bounds (Bao, Carroll, and Singh 2017). The scale ϵ is left unspecified here; Chapter 2 will show that self-consistency ties ϵ to the Planck length, $\epsilon = 2\ell_P$ at one nat of entropy per boundary mode.

Lemma 2 (Causal partition). *The phase space decomposes as $\Gamma = \Gamma_V \times \Gamma_H$, with $H_{tot} = H_V + H_H + H_{int}$. Cross-partition correlations enter only through H_{int}.*

Γ_V is the visible sector; Γ_H is the hidden sector; H_{int} couples them. The decomposition is not a modeling choice but the very definition of embedded observation: an observer with access to all of S would have nothing external to observe.

Lemma 3 (Determinism and selected measure). *φ is a bijection on S. On each finite cycle of the reversible representative, the invariant probability measure is unique and uniform; globally the counting measure is selected among the invariant measures as the one of maximal entropy, becoming the Liouville*

measure on Γ *in the continuum limit:*

$$\frac{\partial \rho}{\partial t} = \{H_{\text{tot}}, \rho\} \equiv \mathcal{L}\rho.$$

The observer uses standard Kolmogorov probability. The bijection requirement is structural: non-injective dynamics would merge distinct total states, erasing the observer's records of past outcomes — contrary to what observation is. On a finite set, injectivity implies surjectivity. This is worth putting in context: the reversibility of φ — equivalently, that the substratum conserves information — is not a peculiar demand of this framework but the standard assumption of mainstream physics. Unitary evolution in quantum mechanics *is* information conservation, and its classical shadow is Liouville's theorem, the conservation of phase-space volume recovered here. Physics treats this principle as effectively non-negotiable: the black-hole information paradox is called a *paradox* precisely because the apparent loss of information under Hawking evaporation would violate it, and the no-hiding theorem (Braunstein & Pati, 2007) makes the conservation statement rigorous. The framework assumes no more than standard physics already does; what is distinctive, developed below, is not the assumption but what emerges from it.

These lemmas contain no quantum postulates. The memory-bearing description — with its admitted quantum representation — emerges under three further conditions.

1.3 The observer's blind spot

Light travels at finite speed; the universe has finite age. Every observer therefore has a horizon — a boundary beyond which no signal has had time to arrive. Beyond it lies ordinary physics, but it is structurally inaccessible. The laws of physics written for what the observer can see must average over everything they cannot. The hidden sector must be traced out.

The trace-out produces a stochastic process on $\mathcal{C}_V$. The visible state x is compatible with many hidden states h; each h sends x to a different future x'. Averaging over h weighted by the Liouville measure produces transition probabilities $T_{ij}(t)$ — the chance that x_i at t_1 becomes x_j at t_2. The deterministic underlying system has become, from the observer's perspective, a stochastic process.

In the Markovian regime — a fast-mixing bath, system slow relative to environment — under the mixing hypothesis the dynamics approaches the homogeneous Markov benchmark, its departure bounded in terms of the residue of mixing ([Main §3.4]). This is the standard setting of open quantum systems and produces approximately classical noise. The cosmological hidden sector differs in four specific ways, and the framework's central theorem is that their conjunction forces non-Markovian visible dynamics — with a fixed-basis unitary quantum representation internal and universal ([Main §3.4]), the compressed

form supplied across the correspondence's class under (T), nontrivially where the dynamics is P-indivisible.

Condition C1 (Non-zero coupling). $H_{\text{int}} \neq 0$: hidden degrees of freedom affect the visible evolution at some accessible step — a non-permutation one-step matrix is the cheapest witness, though influence can also hide at step one and surface later. The horizon is not a static wall — stress-energy conservation enforces dynamical correlations across it, and in the physical realization information flows in both directions. The defining content of C1, and what the necessity argument delivers, is the hidden-to-visible influence; bidirectionality belongs to the realization ([Main §3.4], Remark at the non-permutation-witness lemma).

Condition C2 (Memory persistence). A hidden record of visible history, once written, persists to the readback that uses it; the timescale separation $\tau_S \ll \tau_B$ (C2-slow) is one sufficient mechanism ([Main §1.3]). For the cosmological hidden sector that mechanism operates with $\tau_B \sim H^{-1} \sim 10^{17}$ s, the hidden sector's coarse-grained mixing time, set by the Hubble timescale. A laboratory experiment operates on $\tau_S \sim 10^{-15}$ s or shorter; the ratio is $\tau_S/\tau_B \sim 10^{-32}$. This is the *inverse* of the Markovian regime: the environment cannot reset between measurements. Every correlation it picks up from one experiment is still there when the next begins.

Condition C3 (Sufficient memory capacity). The hidden sector has enough distinguishable states to encode the observed conditional past–future information — $\log_2 |\mathcal{C}_H| \geq I^*$, the bound the necessity theorem actually proves. The hierarchy is the realization, not the condition: the cosmological horizon has roughly 10^{122} independent degrees of freedom at one nat each, the Bekenstein-Hawking entropy, against a laboratory apparatus's 10^{30} to 10^{50} — so $N_H \gg N_V$ there, no experiment appreciably disturbs the hidden sector, and its memory never saturates.

Condition C4 (History readback). History-sensitive hidden mediation: hidden degrees of freedom carry information about the visible past into later visible steps — an operational condition, satisfiable even by a pre-sampled hidden variable revealed by the history. In the cosmological realization the mediation is a causal read-write cycle — the interaction reads what it writes — a property to be demonstrated there rather than assumed: the horizon coupling is bidirectional, acting on the same boundary degrees in both directions, but bidirectionality is a strengthened form of C1 and does not by itself supply the routing that readback asserts (Chapter 7 §7.2). Without it, influence plus storage is mere noise: a hidden sector that records history on one set of components while a disjoint set drives the visible dynamics produces exactly Markovian statistics for as long as one cares to look.

The conditions are independent in statement but coupled in content. A fast-mixing environment with vast capacity drives the dynamics toward the Markov benchmark, its memory suppressed (failing C2 in its timescale form). A slow

environment with limited capacity saturates (failing C3). An isolated environment exerts no influence (failing C1). An environment that stores without reading back yields exact Markov noise despite all three (failing C4). Only an environment that is coupled, persistent, sufficiently capacious, and read back sustains the memory-bearing correlations — the content quantum mechanics represents, the representation itself being universal ([Main §3.4]). The conditions are realization diagnostics rather than exotic constraints: at the lattice cut C1 is structural and the C3 capacity floor holds for the realized process by data processing, while C2 and C4 remain hypotheses of Theorem 22 ([SM §8]); at the cosmological cut C1 is verified structurally, C2 and C3 are verified with enormous margins, and C4 is named but not presently discharged (Chapter 7 §7.2).

A clarification on what "slow" means. C2 is not a claim that physics fundamentally proceeds at different rates in different regions of the universe — it does not. The underlying substratum bijection proceeds at one unified rate, with no preferred clock and no "slow region." Rather, C2 is a claim about *correlation timescales*: how long correlations persist in the hidden sector before relaxing, relative to how quickly the visible sector probes those correlations. The asymmetry between the visible-sector and hidden-sector timescales comes from the partition's structure, not from any local variation in physical law.

The structural source of the asymmetry, for our cosmological situation specifically, is that the cosmological horizon is one physical structure that sets *both* C2 and C3 simultaneously. The horizon's area sets the capacity of the hidden sector at the horizon through the Bekenstein bound: $N_H \sim A/\epsilon^2 \sim 10^{122}$ at one nat per mode. The horizon's surface gravity sets the correlation relaxation timescale: $\tau_B \sim 1/\kappa \sim 1/H_0 \sim 10^{17}$ s. These two quantities — area and surface gravity — are related through horizon thermodynamics ($\kappa = 2\pi T_{\text{horizon}}$ with the horizon temperature set by the cosmological constant), so in our cosmological setting *one* underlying geometric parameter (effectively, the value of Λ) controls both C2 and C3 jointly.

For laboratory observers in our universe, C2 and C3 are therefore correlated rather than fully independent. Both are satisfied because the cosmological horizon is enormous and has slow surface dynamics; both would fail simultaneously if the horizon were small and rapidly evolving. The correlation is a feature of our specific cosmological situation, not a structural redundancy in the framework.

For other partitions, however, C2 and C3 can be tuned independently. In biological systems (Chapter 13), the protein conformational landscape has a hidden sector substantially smaller than cosmological ($N_H \sim 10^{10}$ conformational states), with slow dynamics set by folding-landscape physics rather than horizon thermodynamics. Each condition has a different source: C3 from the configurational entropy of the conformational space; C2 from the kinetic barriers separating metastable conformations. In engineered quantum systems (Chapter 15), C2 and C3 can be tuned independently by choosing the substrate material (which controls N_H through the TLS bath density) and the operating

temperature (which controls τ_B through bath equilibration rates). The BEC analogue-gravity experimental program (Chapter 15 §15.8) specifically tests the framework's predictions in the regime where C3 is marginal while C2 remains satisfied — an experimental design that depends on the independence of the two conditions.

The conditions have distinct physical sources — C2 and C3 are correlated for the cosmological partition and separately tunable elsewhere — while their logical relations are those of [Main §3.4]: C4 is primitive, C1 follows from it in any faithful realization, C3 holds in every faithful realization, and C2 is a physical-regime premise. The framework's empirical content (Chapters 13, 15) and forward predictions (Chapter 18) rely on the separate physical sources to make distinct quantitative claims about regimes where one condition is marginal and others are satisfied.

With the definition, the lemmas, and the conditions in place, the chapter's central theorem can be proved. The remainder of the chapter develops the proof in the stages identified in §1.1: partition-relativity (§1.4), P-indivisibility (§1.5), the two routes to unitary quantum mechanics (§1.6), Bell-inequality violations (§1.7), the characterization theorem in its current per-horizon form (§1.8), and the algorithmic reading of what the theorem establishes (§1.9).

1.4 Partition-relativity

The marginalized transition probabilities of the previous section depend on three inputs: the dynamics, the partition, and the measure. The framework's lemmas fix the dynamics. The measure is fixed only by adding a selection principle: Lemma 3 takes the invariant measure of maximal entropy as the canonical one. Under the canonical measure only the partition remains as a free input, and the following result establishes that this is the entire freedom: the emergent description is determined by the partition alone.

Lemma. *Under the canonical measure of Lemma 3, the emergent description is uniquely determined by the partition. Any parameters of the emergent theory depend only on the geometric and thermodynamic properties of the partition boundary.*

Proof. By Lemma 3 the Hamiltonian flow ϕ_t is unique. By Lemma 2 the partition is fixed. An embedded observer with incomplete knowledge of the total energy averages over energies with a smooth prior, recovering the Liouville measure μ on a phase-space band — the continuum form of Lemma 3's maximal-entropy selection. Letting π_V project onto the visible sector, the marginalized transition probabilities are

$$T_{ij}(t_2, t_1) = \int_{\Gamma_H} \delta_{x_j}[\pi_V(\phi_{t_2-t_1}(x_i, h))]\, d\mu(h),$$

determined by dynamics, partition, and measure. With the dynamics fixed and the measure set to the canonical one, only the partition remains as input. □

Three statements about the measure are distinct, and the lemma above is stated under the second. *Invariance.* Invariance alone does not single out a measure: on each orbit the invariant measure is unique and uniform, but a bijection with two or more cycles carries distinct invariant measures, the uniform measure on each cycle (kernel: `invariance_does_not_select` in `OIBridge/CanonicalMeasure.lean`), so the deterministic evolution by itself does not fix the measure. *Canonical selection.* Among the invariant measures, Lemma 3 selects the counting measure as the one of maximal entropy compatible with the observer's records; its continuum form is the Liouville measure. The selection is a principle the framework adopts — a standard but substantive epistemic postulate, carried in the posit ledger of [Main §4.5] — not a consequence of invariance. *Realization datum.* The canonical measure is the default, not a constraint on representations. A particular realization may carry its own fixed hidden prior μ_H as part of the realization datum — a structured preparation — and the emergent law is then a function of the dynamics, the partition and μ_H. The same bijection and partition under different priors generally give different visible laws; the two-state swap under a uniform and under a point-mass hidden prior is the simplest case ([Main §1.2]). Partition-relativity is therefore relative to the declared measure, and the framework's baseline transition matrices and its physical realization use the canonical one.

Different observers correspond to different partitions of the same (S, φ); different partitions produce different emergent descriptions. Quantum mechanics' parameters are not free constants. The Planck constant $\hbar$ does not appear in the lemmas or conditions; it must emerge from the marginalization, with value set by the partition's geometric and thermodynamic features. Chapter 7 shows that, under the horizon conditions named there, the cosmological partition relates $\hbar$ to the discreteness scale through a thermal self-consistency condition at the horizon surface, with 1/4 Bekenstein-Hawking entropy following — with GW250114 confirming the classical area theorem at 99.999% (Abac et al. 2025); the entropy coefficient itself has no direct test.

Partition-relativity establishes that the emergent description is well-defined. The next question is what kind of description it is — whether the marginalization produces a Markovian process, a generic non-Markovian one, or something more specific. The result of this section is that, in the recurrence and non-permutation regimes established there, the process satisfies a precise structural property with a name in the open-systems literature — P-indivisibility — while C1–C4 by themselves deliver accessible non-Markovianity ([Main §3.4]). The two sections following establish how the imported correspondence supplies the quantum representation across its class, under the translation hypothesis (T) — with P-indivisibility governing whether that representation is nontrivial ([Main §3.1]), which is a claim about the two-time marginal family and does not transport the multi-time content of (C4).

1.5 P-indivisibility

The central technical result of the chapter is that the marginalized process is P-indivisible across the recurrence cycle whenever, under the uniform hidden prior, its one-step visible matrix is not a permutation, as non-trivial coupling acting at the first step makes it. On accessible windows the proved statement is conditional memory: a (C4) readback gap forces $I(X_{<k}; X_{k+1} \mid X_k) > 0$, which does not by itself make the process P-indivisible there ([Main §2.3]).

Theorem (P-indivisibility). *Let $\mathcal{C}_V$ and $\mathcal{C}_H$ be finite sets with $|\mathcal{C}_V| \geq 2$, and let $\varphi : \mathcal{C}_V \times \mathcal{C}_H \to \mathcal{C}_V \times \mathcal{C}_H$ be a bijection with*

$$T_{ij} = \frac{|\{h \in \mathcal{C}_H : \pi_V(\varphi(x_i, h)) = x_j\}|}{|\mathcal{C}_H|}.$$

If T is not a permutation matrix, then the process is P-indivisible.

P-indivisibility is a precise property of stochastic processes that requires unpacking. A process is *P-divisible* if, for any three times $t_1 < t_2 < t_3$, the transition matrix factors through t_2 as

$$T(t_3, t_1) = \Lambda(t_3, t_2) \cdot T(t_2, t_1),$$

where $\Lambda(t_3, t_2)$ is a valid stochastic matrix — non-negative entries, rows summing to one. P-divisibility expresses the natural intuition that probabilistic dynamics should compose: the probability of going from x_i to x_k over the long interval $[t_1, t_3]$ should be obtainable by chaining the probability of going from x_i to some intermediate state x_j over $[t_1, t_2]$ with the probability of continuing from x_j to x_k over $[t_2, t_3]$. Classical Markov processes satisfy this property trivially because the future depends only on the present state, not on history. P-indivisibility means the factorization fails: the candidate intermediate propagator $\Lambda(t_3, t_2)$ that would make the equation hold has negative entries, and no valid stochastic matrix exists.

Breuer, Laine, and Piilo (2009) introduced a more directly interpretable signature of non-Markovian dynamics: *information backflow*. In a P-divisible process, the distinguishability of any two initial conditions can only decrease over time — the future loses information about the past monotonically, as in any classical Markov chain reaching equilibrium. In a P-indivisible process, distinguishability can rebound: information about the initial condition that appeared lost can return to the visible sector at a later time. The system is not memoryless. The "memory" lives in the hidden sector, where correlations laid down by the visible sector's history are preserved and later read back. The signature runs one way: a rebound certifies P-indivisibility, but a P-indivisible process need not produce one — some keep the distinguishability of every pair of initial distributions non-increasing ([Main §2.3]) — so the absence of backflow does not establish P-divisibility.

The proof of the theorem uses the total variation distance between probability

distributions,

$$d(p, q) = \tfrac{1}{2} \sum_k |p_k - q_k|.$$

This quantity ranges from 0 (identical distributions) to 1 (distributions with disjoint support) and measures how distinguishable two probability distributions are by any measurement. For a P-divisible process, d is non-increasing along trajectories: starting from δ_i (the point mass at x_i) and δ_j (the point mass at x_j), the quantity $d(\delta_i T^{(t)}, \delta_j T^{(t)})$ — where $T^{(t)}$ denotes t steps of the dynamics — cannot grow with t. P-indivisibility is therefore established by demonstrating non-monotonic behavior of d along some trajectory.

Proof of P-indivisibility. The proof exhibits non-monotonicity through a three-step construction.

Step 1 (Recurrence to maximal distinguishability). The dynamics φ is a bijection on the finite total set $\mathcal{C}_V \times \mathcal{C}_H$. Every bijection on a finite set is a permutation, and every permutation generates a finite cyclic group: there exists an integer N (the least common multiple of the cycle lengths in φ's cycle decomposition) such that $\varphi^N = \mathrm{id}$. Applied to the marginalized dynamics, this means $T^{(N)}$ equals the identity matrix I. Starting from δ_i and applying $T^{(N)}$ returns δ_i unchanged; starting from δ_j returns δ_j. The distinguishability is $d(\delta_i T^{(N)}, \delta_j T^{(N)}) = d(\delta_i, \delta_j) = 1$ for any $i \neq j$. After N steps, the two initial conditions are again perfectly distinguishable.

Step 2 (Strict contraction at intermediate times). If T is not a permutation matrix, then there must exist visible states x_i, x_j, and x_l with both $T_{il} > 0$ and $T_{jl} > 0$. (If no such triple existed, each row of T would have a single nonzero entry equal to one, making T a permutation.) The existence of x_l as a common destination from both x_i and x_j means the post-step distributions $\delta_i T$ and $\delta_j T$ both assign positive weight to x_l — they overlap. For overlapping distributions p, q with $p_l \wedge q_l > 0$, the total variation distance satisfies

$$d(p, q) = \tfrac{1}{2} \sum_k |p_k - q_k| < 1,$$

the inequality strict because of the overlap. So $d(\delta_i T, \delta_j T) < 1$.

Step 3 (Non-monotonicity). Combining the previous two steps: at $t = 0$, $d(\delta_i, \delta_j) = 1$. At $t = 1$, $d(\delta_i T, \delta_j T) < 1$. At $t = N$, $d(\delta_i T^{(N)}, \delta_j T^{(N)}) = 1$. The trajectory of distinguishability is therefore $1 \to$ (something less than 1) $\to 1$ — non-monotonic. A P-divisible process cannot produce this trajectory because d would be non-increasing throughout. The process is P-indivisible. □

The proof uses only finiteness of the total set — Lemma 1 for the visible sector and the finiteness posit A1 (Chapter 2 §2.4) for the hidden sector, hence Step 1's recurrence — and condition C1 (non-trivial coupling, under the uniform hidden prior that defines T, hence Step 2's overlap). It does not use C2, C3, or any further structural input. In continuous time, for a Hamiltonian flow on compact

energy surfaces, the same argument needs the flow to return near-totally at a common time after the rooted maps have left the permutation class — the counterpart of $\varphi^N = \mathrm{id}$. Poincaré recurrence does not supply this for a general flow: it returns almost every state at its own time, and a mixing flow never again returns a whole visible cell at once at late times. The extension therefore holds for flows with such a common return and is open for general Hamiltonian flows ([Main §2.3]).

The accessible-timescale lemma. A potential objection to the proof as stated is that the recurrence time N can be astronomically large. N is the order of φ as a permutation of the $n = |\mathcal{C}_V \times \mathcal{C}_H|$ total configurations. For any bijection it is at most Landau's function $g(n)$, the largest order of a permutation of n points, with $\ln g(n) \sim \sqrt{n \ln n}$, and it divides $\mathrm{lcm}(1, \ldots, n) = e^{(1+o(1))\, n}$; under the exactly linear update rule (A5) it is below n (Chapter 18 §18.7). The framework bounds N but does not compute it, and n is not fixed by visible data. For N of astronomical size the distinguishability restoration at $t = N$ is not directly observable. What an observer can measure is memory on *accessible* timescales — times short compared to τ_B, the hidden sector's intrinsic memory time — not at recurrence. The accessible-timescale lemma supplies that memory, as conditional mutual information, wherever the stored record is read back (C4). It does not supply P-indivisibility on those windows: a readback gap is compatible with a population law that is divisible on every window short of recurrence ([Main §2.3]).

The mechanism is straightforward information storage and retrieval through the coupling. At each visible-sector interaction (timescale τ_S), the coupling H_{int} writes some quantity I_0 of information from the visible sector into the hidden sector. Between interactions, the stored information decays as the hidden sector's internal dynamics scrambles it. The per-step retention from one visible-sector interaction to the next is

$$r = e^{-\tau_S/\tau_B} \approx 1 - \tau_S/\tau_B,$$

where τ_B is the hidden sector's *coarse-grained mixing time* — the timescale on which the effective channel degrades a stored record. It is emphatically not a spectral gap of the substratum dynamics: the exact Koopman operator of a finite permutation has modulus-one spectrum and cannot mix at all, so no dissipative gap is available to set this rate. The decay rate belongs to the coarse-grained effective channel $\mathcal{K}_\Sigma$ and is exactly the content of the effective-mixing hypothesis (EM) ([Main §4.6]); the estimate below is conditional on it. When $\tau_S \ll \tau_B$, this is close to unity: almost no information is lost per visible-sector cycle. After k steps the cumulative retention is $r^k \approx 1 - k\tau_S/\tau_B$, which remains close to unity for any observation window short compared to τ_B/τ_S cycles.

Quantitatively, when the stored record is read back (C4), the non-Markovian mutual information between the visible sector's past and future, conditioned on

its present, is estimated at leading order by

$$I(X_{<t}; X_{>t} \mid X_t) \approx I_0 \left(1 - \frac{k\tau_S}{\tau_B}\right),$$

which is order I_0 for any $k\tau_S \ll \tau_B$. Conditional memory of order I_0 is therefore present throughout the accessible-timescale regime, until enough cycles have accumulated to deplete the stored record. It is memory in the multi-time law, distinct from the distinguishability revival that the recurrence proof establishes for the one-time population law, and it does not imply that revival.

For the cosmological partition the numbers are striking. The hidden sector's memory time is set by the Hubble scale, $\tau_B \sim H^{-1} \sim 10^{17}$ s. A laboratory experiment operates on $\tau_S \sim 10^{-15}$ s. After $k = 10^{20}$ steps — far more than any conceivable single experiment — the cumulative decay is $k\tau_S/\tau_B \sim 10^{-12}$, negligible. The hidden sector retains essentially all of the information written into it over any observation window short of τ_B. Condition C3 then guarantees that this storage does not saturate: the maximum information the hidden sector can hold is $\log_2 m$ bits, and at the cosmological scale the horizon's degrees of freedom alone can hold of order 10^{122} bits — the horizon's entropy is $S_{\mathrm{dS}} \sim 10^{122}$, hence $m \gtrsim e^{S_{\mathrm{dS}}}$ states. (The figure $\log_2 10^{122} \approx 405$ counts the digits of the degree-of-freedom count, not the capacity; the two differ by the exponentiation.) The margin over anything a laboratory experiment can write is correspondingly vast.

C4's readback therefore produces accessible-timescale non-Markovianity, with C1 its consequence in any faithful realization, C3 holding in every faithful realization by the capacity bound, and C2 the persistence regime that lets it operate with substantial conditional memory at every step; P-indivisibility is a distinct property — on a common window, with a full-support visible prior, the stronger one — delivered at the recurrence scale when the non-permutation one-step witness applies ([Main §2.3]). The recurrence-based proof establishes P-indivisibility across the recurrence cycle; the accessible-timescale lemma establishes conditional memory in the parameter regime that matters for actual observers in our universe. The second does not imply the first on accessible windows.

A worked six-hidden-state model. A concrete example makes the mechanism tangible. Take a two-state visible sector $\mathcal{C}_V = \{0, 1\}$ — call it a coin with states Heads and Tails — and a six-state hidden sector $\mathcal{C}_H = \{1, 2, 3, 4, 5, 6\}$ — call it a die. The dynamics is the permutation

$$\sigma = (0{,}1 \leftrightarrow 1{,}1)\,(0{,}2 \leftrightarrow 1{,}2)\,(0{,}3 \leftrightarrow 0{,}4)\,(0{,}5 \leftrightarrow 0{,}6)\,(1{,}3 \leftrightarrow 1{,}4)\,(1{,}5 \leftrightarrow 1{,}6).$$

Each pair $(\cdot \leftrightarrow \cdot)$ is a transposition that swaps two total states; all six swaps together are independent (act on distinct states), so the full dynamics is a product of disjoint transpositions and satisfies $\sigma^2 = \mathrm{id}$.

Verify the four conditions: C1 holds because die values $h = 1, 2$ couple to coin flips while $h = 3, 4, 5, 6$ do not — non-trivial coupling. C2 holds trivially: the die

does not evolve between coin flips (a static bath), and $\sigma^2 = \mathrm{id}$ gives recurrence at $N = 2$. C3 holds with six hidden states against two visible states. C4 holds because step 2 reads back the record written at step 1.

Compute the one-step transition matrix starting from $x = 0$ (Heads) with the die uniformly distributed. The dynamics sends $(0,1) \to (1,1)$ and $(0,2) \to (1,2)$ — Heads becomes Tails for these die values. It sends $(0,3) \to (0,4)$, $(0,4) \to (0,3)$, $(0,5) \to (0,6)$, $(0,6) \to (0,5)$ — Heads stays Heads for these die values, though the die value changes within the hidden sector. So $P(0 \to 0) = 4/6 = 2/3$ and $P(0 \to 1) = 2/6 = 1/3$. Symmetric reasoning for $x = 1$ gives

$$T(1,0) = \begin{pmatrix} 2/3 & 1/3 \\ 1/3 & 2/3 \end{pmatrix}.$$

The distinguishability after one step is $d(\delta_0 T, \delta_1 T) = \frac{1}{2}(|2/3 - 1/3| + |1/3 - 2/3|) = 1/3$. The two initial conditions, perfectly distinguishable at $t = 0$ with $d = 1$, have become much less distinguishable: d has dropped to $1/3$.

A Markovian process with this one-step matrix would have $T(2,0) = T(1,0)^2$. Computing,

$$T(1,0)^2 = \begin{pmatrix} 5/9 & 4/9 \\ 4/9 & 5/9 \end{pmatrix},$$

with distinguishability $1/9$ — further decreased, as required for a Markov process. But the actual second-step matrix is $T(2,0) = I$ because $\sigma^2 = \mathrm{id}$ sends every total state back to itself, so the visible-sector marginals are unchanged from the starting point. The distinguishability at $t = 2$ has returned to 1. The two initial conditions are perfectly distinguishable again. Complete un-mixing — impossible for any Markov process.

The signature of P-indivisibility shows up as negative entries in the would-be intermediate propagator. Solving $T(2,0) = \Lambda(2,1) \cdot T(1,0)$ for the candidate $\Lambda(2,1)$ gives

$$\Lambda(2,1) = T(2,0) \cdot T(1,0)^{-1} = I \cdot \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}.$$

The off-diagonal entries are -1. No valid stochastic matrix has negative entries — the factorization through $t = 1$ fails by direct computation. The process is P-indivisible.

The mechanism is clear in this example. The die functions as a memory register. At step 1, if the coin was at 0 and the die was at 1, the dynamics produces the new state $(1,1)$: the coin has flipped and the die has not changed value. The die value 1 now encodes a record of what happened — "the coin was at 0 before this step." At step 2, σ acts on $(1,1)$ and returns it to $(0,1)$: the die's stored memory has been used to undo the previous coin flip. The visible sector's history, lost in the one-step distinguishability decay, has been recovered from the hidden sector.

C1 is what enabled writing the record — the coin-die coupling at die values 1 and 2. C2 is what ensured the record was preserved between the write at step 1 and the read at step 2 — here trivially, because the die itself does not evolve between coin flips. C3 is what provided enough storage capacity — six hidden states against two visible; C4 is what made the stored record readable again at step 2. C4's readback is what produces the memory — C1 and C3 are its preconditions in any faithful realization, and C2 is the persistence regime that lets the record survive to be read. The next section takes up how that memory is represented.

1.6 The two routes to quantum mechanics

There are two complementary routes from the observer's stochastic law to quantum form.

The first is internal and elementary. A finite reversible realization lifts to a permutation unitary on the Hilbert space spanned by the total configurations. With the hidden register prepared in the corresponding prior, fixed-basis Born readout reproduces the visible stochastic law exactly, including its finite multi-time trajectory statistics. Combined with the finite process-dilation theorem in the reverse direction, this yields the exact finite-horizon equivalence $S \iff D \iff Q_{\mathrm{fb}}$.

The second route is the stochastic–quantum correspondence of Barandes. It supplies a compressed quantum representation and the associated interpretive dictionary for the source's finite-configuration class. The important logical point is that the existence of a Hilbert/unitary/Born representation does not require P-indivisibility — every finite stochastic law has such a representation. Accessible conditional memory instead identifies the memory-bearing sector; indivisibility across the recurrence cycle is a separate property, present wherever a rooted visible map is not a permutation.

The Born rule in the fixed-basis representation is therefore part of the representation dictionary, not a selector derived uniquely from C1–C4. The full operational problem is narrower, and it is answered as a classification rather than a derivation ([GR §3.3]): with operations taken to act on the complex matrices indexed by the configurations of the system and of its composites — a kinematics the classification assumes and does not derive — the coherent preparations, transformations, and measurements of one OI system and its composites form one standard quantum instrument algebra exactly when five operational conditions hold — valid probabilities, consistency under a trivial one-state ancilla, inert spectators, full reversible control, and iterated composition — and each is independent of the others and of the observation process, so embedded observation alone does not select quantum mechanics; the selection principles are named. Finite classical action-labelled controls are already jointly dilatable; the genuinely coherent/noncommuting and compositional structure is what those conditions supply. The same classification has a layered form, quantum-complete OI (OI^{+}):

the unchanged OI core with well-formedness, observational independence, reversible richness, and observer recursion, equivalent on every nonempty finite carrier to exact finite endomorphic operational quantum mechanics, each added principle independent of the core, well-formedness, and the other two, and none of them entailed by bare OI ([GR §3.3]). A primitive-source refinement restates those three principles at the level of physical implementations and embedded observers — implementation locality, phase-free richness — one continuously driven transition and the exchanges of distinguishable states, with no phase operation — and embedded observation, with no closure of the implementations under the adjoint and no inverse accessibility assumed — equivalent to the same operational quantum mechanics on every nonempty finite carrier ([GR §3.3]). A substratum-source form completes that refinement: the concrete OI substratum supplies the structural architecture — closure under composition and coarse-graining, stability under an uncoupled spectator, relabelling invariance, reversibility, the exchanges of distinguishable states, and, as a stated intervention principle rather than a consequence of its finite states and bijective dynamics, the phases — but, under finite bijective read-write dynamics, not the continuously tunable state-mixing operation; with that one operation added as an empirical controllability resource it is equivalent to the same operational quantum mechanics on every nonempty finite carrier, while finite reversible read-write dynamics, even with genuine hidden-memory and readback behavior, does not itself generate quantum state mixing ([GR §3.3]). Its typed form removes the endomorphic qualifier: under the carrier-general typed operational interface — identity, composition, coarse-graining, relabelling, uniform finite attachment, discard, and factor readout, an interface that itself carries no quantum content — the same equivalence holds with exact finite-dimensional typed operational quantum mechanics, operations between any two nonempty finite carriers included, so the remaining qualifier is finite-dimensional rather than endomorphic ([GR §3.3]). The finite theory admits a canonical infinite-region completion without a continuum limit: the finite regions of the fixed-spacing lattice carry full matrix algebras joined by compatible injective unital embeddings with observables of disjoint regions commuting, the norm completion of their union is a quasilocal C^*-algebra on which every consistent family of finite-region density matrices is a state and the reversible finite-range substratum update is an isometric $*$-automorphism, and any system carrying the same local stages is canonically and uniquely $*$-isomorphic to it, intertwining the dynamics; this removes the finite-region cutoff for the algebra, the state space, and the OI-induced discrete dynamics rather than characterizing all infinite-dimensional quantum instruments or all locality-preserving quantum dynamics; the lattice spacing stays fundamental, no Hilbert-space representation and no superselection sector is selected at the level of the laws, the restriction to the substratum-induced dynamics is a theorem rather than a convention — a phase automorphism at one site preserves locality without arising from any reversible finite-range bijection — and, while the update is joined to the identity by an exact strongly continuous path of $*$-automorphisms of that algebra, with no one-parameter-group law established and no generator exhibited, a continuous-time Hamiltonian law

remains additional structure ([GR §3.3]). Within the structural architecture the substratum supplies, with the phase intervention stated, exact finite endomorphic operational quantum mechanics holds exactly when one continuous layer flow of an involution of the configuration space moving some configuration is an available operation at every level and every intermediate time, the executability of that flow being an assumption not derived by configuration-level generation, as the phase intervention is an assumption on the substratum class and not a consequence of the finite states with the bijective dynamics ([GR §3.3]). The stated substratum dynamics with the observer's read and write access do not source that executability, the layers being available at time one and not at intermediate times, and the observer-level lift, in every formulation stated that lands in the operational interface, derives neither a non-monomial admissible operator nor the layer flow, the site-space wave-operator formulation being unresolved rather than excluded ([GR §3.3]).

1.7 Bell-inequality violations

The Bell claim must separate the deterministic completion from the reduced stochastic description. For a Bell-relevant substratum whose coupling range does not connect the wings before readout, the dependency graph gives exact pointwise separation. If each setting intervention, applied at one step, also reads and writes only its own wing, every run records a ± 1 outcome at each wing, and the response-complete pre-setting ensemble is independent of the settings, determinism yields an ordinary Bell-local hidden-variable model and $|S_{\mathrm{CHSH}}| \leq 2$. Hidden memory and indivisibility do not alter that ceiling.

The framework therefore declines the measurement-dependent/superdeterministic horn and takes **ontic parameter dependence**. In the specific SM/GR realization the target mechanism is preparation-indexed adjacency in the already state-dependent graph $G(x)$ (or an equivalent ontically nonlocal response): the vacuum/uniform reference graph remains geometric and bounded-range, while an entangling preparation may supply Bell-relevant long adjacency so a microstate response at one wing can depend on the remote setting. H-Bell is the open requirement that those prepared graphs preserve operational no-signaling and the metric/Ollivier–Ricci continuum limit. The curvature condition is downstream of a harder one. The cited convergence theorem is arbitrary-dimensional only for the manifold-distance-weighted graph metric, and for the degree-6 cubic reference graph the intrinsic hop metric is exactly ℓ_1 — a $\sqrt{2}$ diagonal stretch that does not decay at any scale — so the hop-metric curvature route fails for the reference family, before any Bell edge is added, and no $D = 3$ extension of that theorem would rescue it. The emergent geometry itself is not in question: the substratum's dispersion relation is isotropic at leading order. What must be re-founded is the curvature functional, on the propagation/Laplacian geometry rather than on shortest-path counts [SM §3.2]. The setting-dependent alternative is recorded but not adopted; Main §3.3 gives the model-independent necessary lower bound $I_{\mathrm{ont}} \geq 0.0309$ bits for the

quantum value. Hall–Branciard's optimized causal CHSH model uses about 0.080 bits, while the smaller ~ 0.046-bit benchmark is retrocausal.

The cited causal-local indivisible **stochastic** construction separately obeys Tsirelson's bound because its causal-independence notion is weaker than Bell factorization. That theorem is not the hidden-variable locality structure of the deterministic completion. If the spatial/intervention restriction is removed, the general finite OI response construction can realize PR-box behavior, showing that representability, no-signaling, hidden memory, and indivisibility do not by themselves select the quantum Bell set.

1.8 The characterization theorem

The characterization is best read as a layered biconditional rather than a single slogan.

Finite representation theorem. On any finite horizon, $S \iff D \iff Q_{\mathrm{fb}}$. Every finite stochastic process has a finite deterministic reversible realization, and every such realization has an exact fixed-basis unitary/Born representation. This supplies both directions of the finite observable-law equivalence.

Readback theorem. C4 is the condition that makes the OI sector memory-bearing. It forces different hidden posteriors for visible histories that share a current state but have different future laws, a persistence/readback requirement, and the capacity lower bound of C3. Because the physical representative is finite and reversible, it also recurs: its rooted visible maps, defined for every root under one hidden prior, return to the identity, and a non-permutation stochastic matrix has no stochastic inverse, so any non-permutation rooted map makes the rooted law indivisible somewhere in the recurrence cycle. A C4 gap supplies such a map, and so does every non-deterministic visible step, including in laws that are Markov on every accessible window. With a full-support visible prior, that indivisibility is equivalent to memory at some order below the recurrence time; an observer sees the memory only when it appears at an accessible order, which is what C4 asserts.

Spatial extension. The dependency graph supplies finite causal cones exactly for finite-range substrata. Bell adds a separate constraint: for a response-complete deterministic completion that records a ± 1 outcome in every run, measurement independence plus parameter independence would imply $|S_{\mathrm{CHSH}}| \leq 2$. Section §1.7 therefore takes the measurement-independent, ontically parameter-dependent branch for Bell-violating substrata and keeps operational no-signaling separate.

Operational boundary. These theorems do not prove that the entire family of coherent instruments and composites equals the standard quantum operational theory. On a finite carrier, within complex matrix kinematics taken as given, that equality holds exactly under the five completion conditions of [GR §3.3]. The substratum's stated interventions do not supply one of them, full reversible

control; whether an extended OI mechanism can supply it, and which of the others the substratum supplies, are open; this is not a qualification on the exact finite stochastic/representation equivalence itself.

A second boundary sits alongside it, and it concerns which unitary evolution relates two times. The equivalence says which visible laws admit a fixed-basis unitary representation; it does not single out the evolution connecting one time to the next. For dilated processes one visible family can admit distinct coherent unitary lifts — each reproducing the visible law at every time, each composing exactly across times — whose relating evolutions differ, because a lift may be deformed by a time-dependent right action that preserves everything the visible law sees. A deformation constant in time cancels from every relating evolution, so time dependence is what the nonuniqueness needs. Making the relating evolution unique therefore requires a selection condition beyond visible-law representability and ordinary coherence, and this book does not supply one or say what form one must take. The invisible freedom itself has two sides — a right action by the anchored class and a left action within each visible output's hidden fibre, each maximal among uniform actions of its kind — and it does not exhaust the freedom: two lifts can fail to be related by any uniformly invisible right deformation and still agree on every relating evolution, and a pair outside that right class can differ in relating evolution while being related by a left in-fibre move. At one time, what the visible law leaves free after both actions is known exactly: it is the off-diagonal part of the fibre-Gram data — for each visible output, the overlaps among the hidden states into which the lift sends the visible inputs — modulo phases, with the visible law as its diagonal; and that residue is not inert, since two lifts of one visible law can differ in it and differ in relating evolution. That residue does not reach the fixed-basis layer. With a trivial ancilla, the fixed-basis datum built from a lift's unitary at one time, with uniform initial law and identity readout, has finite-horizon trajectory laws fixed by the visible law alone, also after contractively scaled partial permutations on either side, and two such data have equal laws exactly when their visible laws are equal; the fixed-basis correspondence acts on the quotient by visible equality. The trivial ancilla is part of that statement: an ancilla attached as a tensor factor with its own unitary leaves the rooted family unchanged, while an ancilla carried between steps as hidden basis states can separate two lifts that share every column entering the visible law, already in the two-step law. What remains open is two-part: what selects the trajectory of that data through time, and what fixes the relating evolution within a trajectory — the latter not fixed even by the complete trajectory. No selection principle is named or excluded. None of this weakens the finite observable-law equivalence, and none of it makes embedded observation and quantum mechanics inequivalent.

1.9 What the theorem says

The theorem says something stronger than "quantum mechanics can be simulated by a classical hidden model" and something more precise than "C1–C4

uniquely force all of quantum mechanics."

It says that, for finite observable laws, the stochastic, reversible-deterministic, and fixed-basis quantum descriptions are mathematically intertranslatable. A finite embedded observer therefore does not need quantum mechanics added as a separate microscopic postulate in order for a Hilbert-space/unitary/Born description to exist. History readback then supplies the physical distinction between a trivial Markov member of that representation class and the memory-bearing sector characteristic of OI.

The converse is equally important: every finite quantum record has a finite reversible OI realization, and one sufficiently large finite construction can cover an entire bounded finite quantum experiment menu to any specified accuracy. The exact continuum operational theory is a mathematical idealization rather than a requirement that the physical carrier itself be infinite.

The remaining qualification is localized. To identify the complete local operational theory of complex quantum mechanics, rather than its exact stochastic representation, one must lift the whole coherent intervention and composite hierarchy into one compatible quantum instrument theory, which on a finite carrier, within complex matrix kinematics taken as given, happens exactly under the five completion conditions of [GR §3.3]. The Bell fork is logically separate: §1.7 specifies the adopted measurement-independent, ontically parameter-dependent completion and its no-signaling requirement rather than treating Bell screening as one of those completion conditions. The book keeps both boundaries explicit rather than weakening what is already exact.

1.10 The four structural limits

The characterization theorem establishes the dilation equivalence, in which C4 coincides with the non-Markovianity clause, C1 follows from it, and C3 holds in every faithful realization by the capacity bound, with the quantum representation internal and universal — $S \iff D \iff Q_{\mathrm{fb}}$, the realization's permutation unitary, the compressed form under (T) — and nontrivial at the recurrence scale where the dynamics is P-indivisible ([Main §3.4]); (C2)'s necessity is the physical-memory theorem: its slow-bath (timescale) form is conditional on the mixing hypothesis, its persistence form, at the realized distance of the hidden conditionals from the reference prior, needs no mixing hypothesis, and an exact reversible description retains the record for every law. Each condition's failure was treated in §1.8 as a contrapositive in the memory currency — violating the condition removes the accessible non-Markovianity that is the framework's discriminating content; the representation itself, being universal, survives every failure trivially. This section develops the failure cases in the opposite direction: not as proofs of necessity but as *structural commentary* on what kind of physics emerges when each condition fails completely. The three limit regimes reveal that the framework's conditions are not arbitrary technical assumptions; they jointly characterize the kind of observer we actually are — coupled to a slow,

vast environment — and each failure mode describes a hypothetical observer we are not.

The C1 limit: the isolated observer.

C1 requires non-trivial coupling between visible and hidden sectors. In the limit of complete failure — no coupling at all — the visible sector evolves as a *closed* deterministic system. The bijection φ restricted to $\mathcal{C}_V$ is a permutation; the dynamics on V is just the corresponding classical evolution. No information ever crosses the partition boundary. The hidden sector exists but is causally disconnected from the visible sector for all purposes.

The dynamics in this regime is fully classical in character: memoryless. The reduced description still admits the fixed-basis quantum representation — every finite law does, trivially here ([Main §3.4]) — but there is nothing memory-bearing in it: no coupling-induced non-Markovianity for a nontrivial representation to encode, hence none of the interference phenomenology the framework attributes to hidden-sector memory. No P-indivisibility arises from a partition with no coupling, since the visible sector then evolves autonomously by a permutation; Bell violation is a separate question, governed by ontic parameter dependence (§1.7), not by P-indivisibility. The observer sees a classical Newtonian world with deterministic trajectories and well-defined values of every observable at every time.

This is the regime closest to the historical picture of classical mechanics: a clockwork universe where every particle has a definite position and momentum and the equations of motion fully determine the future from the present. The framework's content is that this picture is correct for the limit $C1 \to 0$ — but no laboratory observer occupies this limit. Gravitational coupling between visible and hidden sectors is unavoidable in our universe; stress-energy conservation forces correlations across any horizon; the cosmological horizon is not an impermeable wall. Real observers are coupled, and what coupling produces is the memory-bearing quantum phenomenology — the representation itself is universal; coupling supplies the nontrivial sector.

The C1 limit therefore illuminates a structural point: classical mechanics is what *uncoupled* systems do. The apparent quantumness of the world is a signature not of intrinsic randomness in the underlying reality but of our *non-isolation*. The substratum is deterministic; what makes physics look probabilistic is that the observer is coupled to something they cannot directly see. An observer who could isolate themselves completely from everything outside their measurement apparatus would see classical mechanics. No such observer exists in our universe, and the framework's content is that no such observer can exist — gravitational coupling is universal.

This has a striking corollary: the project of "explaining quantum mechanics" by finding the right deterministic substructure is well-defined within the framework, but the dynamics that observer's would see in that substructure is not quantum mechanics. It is classical mechanics. Quantum mechanics is the dynamics

that *embedded* observers see when they are coupled to the substructure; the substructure itself, viewed from a hypothetical uncoupled vantage point, would be classical. The framework's reframing is that the persistent intuition "there must be a deterministic story underneath" is correct, and the deterministic story is classical, but it is structurally inaccessible to any actual observer.

The C2 limit: the thermalized observer.

C2 requires the hidden sector's coarse-grained mixing time to exceed the accessible window: $\tau_S/\tau_B \ll 1$, with τ_B the timescale on which a stored record degrades rather than the rate at which the hidden sector evolves ([Main §1.3]). In the limit of complete failure — the record is erased instantly, $\tau_B \to 0$ — the hidden sector equilibrates between every visible-sector measurement. No correlation written into the hidden sector during one measurement persists to influence the next. History readback cannot occur because there is no memory in the bath for the visible sector to read back.

The dynamics in this regime is *fully Markovian*. The reduced description on the visible sector is a classical stochastic process with transition probabilities depending only on the current state. The observer sees probabilistic dynamics, but the probabilities are memoryless: the memory-bearing/interference mechanism is absent, while the fixed-basis representation survives, trivially diagonal-preserving ([Main §3.4]). This temporal statement must not be turned into a Bell claim: Bell violation in the deterministic completion is governed separately by ontic parameter dependence (§1.7), not by P-indivisibility. The Born rule reduces to the classical probability assignment for transitions between well-defined states. A Schrödinger-form unitary representation still exists — Q_{fb} supplies one for every finite law — but here it encodes no temporal correlations: what the regime lacks is the memory-bearing content, not the equation's existence ([Main §3.4]).

This regime corresponds to what physics calls the *thermal limit* — systems in contact with a fast equilibrating bath, where the bath's state at any moment is independent of the system's history. The classical statistical mechanics of thermal systems is exactly this limit. Brownian motion, chemical reaction kinetics in well-stirred solutions, ordinary diffusion processes — these are all Markovian dynamics described by classical stochastic processes, and the framework's content is that this is what C2 failure looks like.

No laboratory observer occupies the thermal limit completely. Our cosmological horizon evolves on a timescale $\tau_B \sim H^{-1} \sim 10^{18}$ seconds, while laboratory timescales are $\tau_S \sim 10^{-15}$ to 10^{-1} seconds. The ratio $\tau_S/\tau_B \sim 10^{-32}$ is extreme — the cosmological horizon is essentially frozen on any laboratory timescale, and this frozen-bath condition is what secures (C2-structural) for the coarse-grained description of the horizon — record persistence to readback — by the most robust mechanism available; in an exact reversible description the record is retained for every law. Under the effective-mixing hypothesis, and in the coarse-grained description generated by iterating the one-step effective channel

([Main §4.6]), an observer in genuine thermal equilibrium with a fast-mixing bath would have its accessible memory sector suppressed, the readback gap eroded exponentially in τ_S/τ_B at every order $k \geq 2$ before it can be read; an observer with a frozen bath (our situation) retains it through the record persistence C2 supplies, read back as conditional memory where C4 holds. What the condition governs is the memory-bearing sector, not the representation, which is universal.

The C2 limit therefore illuminates a different structural point: classical stochastic mechanics is what *thermalized* systems do. The framework gives a structural account of why decoherence and the classical limit emerge in systems coupled to fast environments — a familiar phenomenon in quantum mechanics. Decoherence theory describes how quantum coherences are destroyed by coupling to a fast bath; the framework's reading is that decoherence is the transition from the memory-bearing regime back toward the C2 failure limit. Systems with engineered fast baths approach classical stochastic behavior; systems isolated from fast baths retain quantum coherence. The conventional thermal-classical-limit story is the framework's C2 limit told from inside quantum mechanics.

A corollary, stated carefully: what the framework requires is a hidden sector that RETAINS the record across the window, which is not the same as a frozen one. Fast hidden dynamics that conserve and route the record satisfy the condition — a shift-register bath rotating five cells per visible step supports a maximal readback gap (`fastbath_probes.py`) — so persistence, not slowness, is doing the structural work beyond merely providing some unobserved degrees of freedom. An observer whose fast hidden bath scrambles the record sees statistics with suppressed readback; one whose fast bath conserves and routes it retains the full memory-bearing description (`fastbath_probes.py`). The cosmological horizon's slowness is therefore the physically comfortable route to persistence in our universe — and the small parameter of the continuum Schrödinger-form derivation, the τ_S/τ_B expansion — not what representability or the memory theorem require ([Main §3.4]).

The C3 limit: the universe-sized observer.

C3 requires sufficient hidden memory capacity — $\log_2 |\mathcal{C}_H| \geq I^*$, with $N_H \gg N_V$ the comfortable realization regime. In the limit of complete failure — $N_H \to 0$, the hidden sector vanishes entirely — the visible sector V approaches the total system S. There is no trace-out because there is nothing to trace over. The observer is *coextensive* with the universe.

The dynamics in this limit is fully deterministic and fully classical. The bijection φ on the full system S is just a permutation of classical states; the observer who is all of S sees its complete deterministic evolution. No memory-bearing sector — the fixed-basis representation survives, trivially, as always ([Main §3.4]) — and no interference phenomenology, but for a different reason than in the C1 case. In the C1 limit, the visible sector is closed but the framework still applies to it as a system; in the C3 limit, the framework's content has no purchase

because there is no embedded observation taking place. The "observer" is not observing anything; they are the system being observed, with no separation between knower and known.

This is the regime that superficially resembles a *God's-eye view*. The traditional theological framing imagines an observer who stands outside the universe and sees everything at once with no uncertainty. The framework's C3 limit is similar in eliminating quantum indeterminism, but it differs structurally: there is no "outside" in a finite deterministic universe. Such an observer would be coextensive with the universe, not transcendent of it. They would not be viewing the universe; they would be the universe describing itself.

Three structural features of this regime are worth noting.

Observation breaks down entirely. In the framework's account, observation is a *partition-relative* concept. The observer reads the visible sector through measurements that depend on the partition's coupling. Without a partition — without anything in the hidden sector to be coupled to — measurement has no meaning. The observer cannot perform observation in the framework's sense. The God's-eye view is therefore not a limit case of observation but a category outside it.

Laplace's demon is structurally impossible. The classical version of the God's-eye view is Laplace's demon: a hypothetical intelligence that knows the position and momentum of every particle and can therefore predict everything. Conventional readings reject Laplace's demon because of quantum indeterminism; the framework's reading rejects it for a deeper reason. The demon would require being the universe, not merely knowing about it. The impossibility is structural rather than epistemic — not "the demon cannot get the information" but "the demon's having the information would mean they are the universe itself."

Self-reference paradoxes persist. Even in the limit case where C3 fails completely, the Gödel-Turing-OI incompleteness family from Chapter 12 §12.8 applies. A system that is the entire universe and tries to describe itself fully runs into the same self-reference paradoxes that finite systems do. The universe-sized "observer" cannot include a complete description of themselves within their description — adding such a description would change the universe and require a new description. The framework's content is that complete self-knowledge is structurally impossible at any scale, including the limit case.

The C3 limit therefore illuminates a third structural point: observation is partition-relative, and the God's-eye view is not a limit of observation but a category outside it. The framework's anti-Laplacian content is stronger than the conventional one: not "we cannot know everything because of quantum randomness" but "complete knowledge requires being the universe rather than being in it, and any embedded observer is structurally limited."

The three limits together.

The three failure modes are structurally distinct.

Condition	What fails	Resulting dynamics	Resulting situation
C1	Coupling	Closed classical	Isolated subsystem
C2	Timescale separation	Markovian stochastic	Thermalized observer
C3	Capacity hierarchy	No observation	Coextensive with system

Each tells us something different about why our situation produces the memory-bearing dynamics that embedded observation forces. C1 failure reveals that closed classical mechanics is what uncoupled systems do — our dynamics carries hidden-sector memory because we are coupled. C2 failure reveals that memoryless stochastic dynamics is what thermalized observers see — the stored record erased before readback; ours survives because our bath persists rather than equilibrating. C3 failure reveals that observation itself is structurally limited — the memory sector exists because we are embedded, not coextensive.

Together, the limits characterize the kind of observer we are by contrast with the kinds of observer we are not. We are not isolated (C1 holds), not thermalized (C2 holds), not coextensive with the universe (C3 holds), and not blind to our own past records (C4 holds). The framework's four conditions are the diagnostic checklist for "embedded laboratory observer coupled to a record-persistent, vast environment" — C4 primitive, C1 its checkable consequence, C3 a capacity bound every faithful realization meets, C2 the physical regime that keeps records alive ([Main §3.4]). Each is empirically robust at its own end — the gravitational coupling enforcing C1, the cosmological timescale enforcing C2, the cosmological horizon enforcing C3 — and the memory-bearing dynamics this observer must see is the framework's discriminating content — the nontrivial sector of a representation class that is itself universal.

This is the framework's structural content at its sharpest: quantum mechanics is not a separate physical theory we discovered alongside classical mechanics. It is the dynamics that our specific kind of observation produces — with the representation universal ($S \iff D \iff Q_{\mathrm{fb}}$: every finite-horizon law has one) and the discriminating content in the memory-bearing sector ($M_t > 0$) that embedded observation forces — and the alternative dynamics (closed classical, memoryless stochastic, deterministic self-coextensive) correspond to alternative kinds of observation that are not realized in our universe. The framework's four conditions are the structural fingerprint of being a coupled, record-persistent, partition-relative observer — the only kind of observer that physical laboratories produce, and the only kind whose visible dynamics carries that memory sector.

This reframing has consequences for the framework's empirical content beyond what is already developed in Chapters 5 through 19. Each of the four conditions is independently testable through the restricted quantum-statistical regime — engineered systems approaching the marginal-failure boundary. Chapter 15 §15.7 develops the restricted quantum-statistical regime in engineered quantum

systems with the BEC analogue-gravity test as the framework's cleanest experimental probe of marginal hidden capacity. Chapter 13 §13.7 develops the proposal's conjecture for biological systems. Chapter 18 §18.3 develops the restricted quantum-statistical regime as a forward prediction of the mechanism proposal. The four structural limits are therefore not just conceptual content but a framework of empirically accessible regimes, with current experiments now beginning to probe the marginal-failure boundaries.

1.11 The card table and the chessboard

Chess is a game of complete information: every piece stands in the open, and in principle the game is exhaustively analyzable — a sufficiently large lookup table settles every position. Poker is a game of incomplete information, and that difference is not cosmetic. When von Neumann founded game theory he took poker, not chess, as the interesting case, precisely because hidden cards force probability into rational play even though nothing in the rules is random. The mixed strategy is not ignorance dressed up; it is the correct mathematical form of play from inside an information set. The framework's claim can be stated in exactly these terms: the substratum is a chess game, and physics is the rulebook a piece would rationally write from inside it. What this section shows is that ordinary poker is not a rich enough interior view — and that the specific way it falls short, and the specific modification that repairs it, together reproduce the anatomy of Chapter 1 in miniature.

Ordinary poker yields only classical probability. Its hidden information is a static deal, uncorrelated with the visible play once the cards are down, and the resulting odds are divisible: the probability of an outcome at showdown is the probability chained through any intermediate street, whether or not that street was observed. This is the mathematics of ignorance about a definite fact — the position Einstein hoped quantum probability would turn out to occupy. A game that stops here illustrates the *premise* of the framework (observation from inside a partition) while missing its *content* (that structured incompleteness yields quantum, not classical, statistics).

Three modifications close the gap, and they are conditions C1–C4 of §1.3 in card form. First, the shuffle between streets is a deterministic, reversible permutation of the deck — a faro shuffle, say — rather than a randomization: bijective dynamics on the full state. Second, each street deals burn cards, face down and never shown, which the shuffle correlates with the live cards: a hidden sector coupled to the visible one (C1), whose configuration persists between streets (C2), with capacity to store a record of the visible play (C3). Third, revealing a card is an intervention, not a passive glance: a revealed card is burned and replaced under a fixed rule, so revelation physically breaks the card's correlation with the rest of the deck.

A small such game deals the write-then-read mechanism of the coin-and-die model of §1.5 onto a table, with one burn card in place of the die. The player

holds one hole card, red or black — the coin. One burn card, red or black, lies face down — the memory register the die provided. The shuffle between streets is the simplest reversible move available: hole card and burn card trade places. Then, writing the odds that the hole card at showdown matches the hole card as dealt:

protocol	matching odds
one shuffle, then showdown	1/2
two shuffles, then showdown	1 (certain)
two shuffles, odds chained through the unobserved middle	1/2
two shuffles with a peek (and forced re-burn) in the middle	1/2

Each line is checkable by hand — there are sixteen cases in the longest protocol. A single shuffle randomizes the visible card completely. Two shuffles return it with certainty: the first swap writes the hole card into the burn pile, the second reads it back — the same write-then-read information backflow that §1.5 exhibited in the would-be intermediate propagator with negative entries. Chaining the one-street odds through the unobserved middle street predicts even money and is simply wrong: the process is P-indivisible, and a player who prices hands divisibly loses money at a rate equal to the interference term — the deviation has an operational meaning as concrete as a bankroll. And peeking at the middle street, which forces the re-burn, makes the divisible answer correct again: a division event in the sense of §1.5, enacted with cardboard.

The reading this hands the student is the one the framework intends. Interference is not a substance; it is the failure of divisibility — the visible history taking a round trip through the hidden sector and returning. Phase is not a mysterious internal angle; at this resolution it is *where in the hidden sector the visible information currently resides.* Measurement does not collapse anything; it intervenes, and the intervention severs a correlation that the undisturbed dynamics would have used.

The game extends upward. Replace the two-card swap with a linear shuffle on ranks modulo q and the game runs on the same dynamical class as the substratum of Chapter 2; the reachable odds tables grow richer, and the fact that not every doubly stochastic table can be dealt this way is the unistochasticity constraint of [Main §3.1] appearing at the card table. One boundary must be stated as plainly as the successes: a card mechanism played under the restrictions below is a local classical toy. It reproduces single-system interference, indivisibility, and division events — the mechanism of this chapter — but it violates no Bell inequality when, once the deal is down, every operation — each shuffle the rules prescribe, each move and each setting choice — reads and writes one player's own cards only; which operation is applied to a player's cards, and how that player's ± 1 outcome is read from them, depends only on those cards, that player's own setting and on chance that, taken together with the deal, is

independent of the settings; an outcome is recorded in every run and no run is set aside; and the settings are chosen independently of the deal: such a game is a local hidden-variable model (Main §3.3). The Bell-type correlations of §1.7 belong to the full construction, through the level-identification analysis taken up in Chapter 3 (§§3.7–3.7.1, where the boundary drawn here becomes the diagnosis), not to the toy. An illustration that claimed otherwise would be teaching a falsehood, and the analogy earns its keep precisely by knowing where it ends.

What remains is the pair of views. Turn the deck face up and track every permutation, and the game is chess: complete information, deterministic, exhaustively analyzable. Sit in the player's seat and the same game is poker — and not the poker of static ignorance, but the modified game whose rational odds sheet carries interference terms. Nothing changed but the seat. That is the framework's thesis in one gesture: observation is what the rest of the board looks like from inside a piece.

Schrödinger's cat, played as cards. Every reader arrives carrying one thought experiment, and the game just built is the right table to play it on. Schrödinger (1935), in the same paper that coined *entanglement*, sealed a cat in a chamber with a radioactive atom, a Geiger counter, a hammer, and a flask of poison: within the hour, the formalism's entanglement chain has amplified one atom's superposition, link by link, into a description assigning the cat itself equal parts alive and dead until the box is opened. He offered it as a "quite ridiculous case" — a reductio, not a report. If the wavefunction is a complete description and "measurement" an undefined primitive, nothing in the theory says where along the chain atom → counter → hammer → cat a definite fact appears. Ninety years of interpretation dispute is the residue of that question.

Played on this table, the paradox takes four sentences. The sealed hour is an unobserved middle street. "Alive + dead" is not a state of the cat but a line on the odds sheet — the indivisible pricing any player must use for a hand whose history routes through the burn pile. Opening the box is the peek with its forced re-burn: an intervention that severs a correlation, not the creation of a fact. And the deal was definite throughout — the substratum holds exactly one configuration at every moment; the superposition is bookkeeping for where the record resides, in cards the observer has not turned. What dissolves the *cat* specifically is a further observation: a cat's middle street is never unobserved. Counter, air, thermal photons — the environment peeks continuously, each interaction a division event forcing the re-burn, so the divisible odds sheet is the correct one for cats, counters, and kittens alike, and the interference terms that would distinguish "alive + dead" from "alive or dead" are committed to the hidden sector faster than any protocol could retrieve them (§1.6). No consciousness is required to open this box; one stray photon holds a full hand.

Schrödinger's instinct was therefore sound — no cat is ever alive-and-dead — but the framework's repair is subtraction where the objective-collapse programs (GRW/CSL, Diósi–Penrose) offer addition. They make the cat definite by appending a stochastic law; the framework makes it definite by deleting a premise

— the superposition was the observer's ledger, not the chamber's state, so there was never anything to collapse (and Chapter 18 records that additions of the Penrose type are structurally forbidden here). The two repairs part company at exactly the frontier of §1.6: engineered "cat states," superpositions of macroscopically distinct configurations held genuinely unobserved, which collapse laws must eventually kill and the framework's exactness theorem must eternally protect — the wager registered in Chapter 4 §4.5. The boundary drawn above remains load-bearing: a cat in one box is a single-system story, the toy's home turf; it is Bell's two boxes, not Schrödinger's one, where the cards must be surrendered to the full construction.

The substratum that the framework's algorithm operates on is constructed concretely in Chapter 2 — a three-dimensional cubic lattice with bilinear, block-diagonal dynamics whose structural constraints fix the framework's empirical content. Chapter 3 takes up the framework's full philosophical positioning, including its relationship to the structural realism program, its relation to Bell, Kochen-Specker and PBR-type no-go theorems through separate checks of their local-response, operational-algebra and preparation premises, and its parallels to the broader intellectual landscape of incompleteness results from Gödel through Wolpert. Chapter 4 develops the methodological posture — the single-foundational-commitment derivation pattern, the architecture-level engagement with the no-go results, the classification of claims by derivational rigor — that organizes the framework's program as a whole. The remaining chapters trace the framework's predictions through particle physics, gravity, cosmology, the structural cascade from atoms to intelligence, and the framework's reach into chemistry, biology, medicine, and quantum engineering.

The remainder of this part of the book develops the substratum, the structural realism, and the methodology in detail. Part II turns to the framework's quantitative physics. Part III traces the cascade from substratum to chemistry, biology, and intelligence. Part IV develops the framework's reach into working scientific fields.

Chapter 2

The Substratum

2.1 What this chapter constructs

Chapter 1 established the exact finite observable-law equivalence between stochastic processes, finite reversible deterministic realizations, and fixed-basis unitary/Born representations. It also showed that genuine history readback forces hidden predictive memory in every faithful completion. This chapter

now asks the inverse structural question: what is the finite reversible object (S, φ) whose observer-level projection has that description?

Three questions need separation. First, which **local propagating lattice/gauge residue** is fixed by the framework's empirical and structural inputs? Second, which changes of a representative leave that local observable content unchanged? Third, can the Bell-violating preparation/composite structure be embedded in the same deterministic ontology, and if so is it unique? The reconstruction theorem answers the first, the substratum gauge group addresses the second for the reconstructed dilation/local data, and H-Bell addresses only the existence/compatibility part of the third; Bell-inclusive uniqueness remains open.

The quantum input to this reconstruction should be read in the scoped sense established in Chapter 1. The visible law has an exact quantum representation, spatial OI provides graph causal cones, and finite quantum records have finite reversible realizations. The operational completion for arbitrary coherent instruments/composites, which holds, within complex matrix kinematics taken as given, only under the five completion conditions of [GR §3.3], is not silently consumed as though the substratum had been shown to supply them; where the reconstruction uses stronger standard-QM structure, that structure is an empirical input or a named modeling premise.

2.2 The substratum as a finite lossless memory

The framework's central object is a triple. S is a finite set whose elements are the distinguishable total states. $\varphi : S \to S$ is a bijection — a deterministic, reversible transition rule that maps each state to its unique successor. $V \subsetneq S$ is the visible sector, the observer's window into S; the complement $H = S \setminus V$ is the hidden sector. These three together — (S, φ, V) — fix everything: the dynamics, the observer's access, the marginalized description.

The most useful physical interpretation is in terms of memory. S is the set of all distinguishable states the system can occupy — a finite storage capacity. φ is a bijection, so information is never created or destroyed; the substratum is a *lossless* memory in the precise information-theoretic sense. The partition V defines the observer's access to this memory, both read and write. The observer reads the visible sector by measuring x, and writes to the hidden sector — every visible-sector interaction imprints correlations onto H through the coupling H_{int}.

The observer-level statistics of this read-write cycle have an exact finite quantum representation. The structural form of C2 is persistence: information written into the hidden sector must survive until the readback that uses it; a slow bath is one sufficient mechanism rather than the definition. C4 is what makes the stored information operationally matter. Chapter 1 proves that such readback forces hidden predictive memory in every faithful completion. Interference and entanglement are represented quantum-mechanically in the corresponding observer-level description; identifying every possible coherent intervention with

the standard quantum instrument algebra holds there, within complex matrix kinematics taken as given, exactly under the five completion conditions, of which the substratum's stated interventions do not supply full reversible control; whether an extended OI mechanism can supply it, and which of the others the substratum supplies, are open.

This memory interpretation makes several of the framework's structural quantities concrete. The 10^{122} vacuum-energy discrepancy of Chapter 1 is the compression ratio between total storage and readable storage — between the full configuration space S and the visible sector V. The dark sector of cosmological observation is the gravitational effect of the unreadable storage in H. The Bekenstein-Hawking entropy is the storage capacity of the partition boundary, the number of independent modes crossing ∂V times the entropy σ of one mode. The action scale $\hbar$ is the conversion factor between storage geometry (boundary area) and read-write statistics (the unitary group's parameter). Chapter 7 develops each of these under the conditions it names; for the present chapter they motivate the interpretation.

The substratum dynamics is defined on the full configuration space, not on the visible sector alone. The wave equation that selects φ among second-order reversible nearest-neighbor dynamics — developed in detail in Chapter 5 — operates on both V and H. The partition imposes an observer-access structure on this common dynamics but does not change the dynamics itself. Both sectors are governed by the same underlying rule; the only difference is which states the observer can read.

2.3 What the substratum is and is not

The substrate objection asks an obvious question: what is the memory made of? What physical stuff implements the configuration (S, φ)? The framework proposes a structural description of dynamics and observation rather than specifying an underlying material. Two implementations that give identical data at a specified observation interface cannot be distinguished by a decision based only on those data. An independently prepared, normalized, exactly decoupled finite ancillary factor is one such redundancy: tracing it out leaves the original observed channel unchanged [Substratum Theorem 24(iii)]. This does not show that every proposed implementation has identical data, that a sector which later couples to the observer is redundant, or that the explanatory regress has been terminated. A complete observable-to-substratum classification remains open. The physical implementation question is therefore separate from the exact invariances that have been established.

A closer analog than physical substance is the abstract structure of a computation. A Turing machine has three components: a tape (the storage medium), a head (the read-write access window), and a transition function (the update rule). The mapping to the substratum is direct: S corresponds to the tape, V to the head's access window, φ to the transition function. But three differences

are physically significant.

First, a Turing machine's tape is potentially unbounded while the exact OI model here has finite S. Finiteness of the total carrier gives the exact return used by Chapter 1's recurrence argument. The holographic bound instead concerns the observed sector under its stated interpretation; it does not establish total-carrier finiteness. An infinite extension requires its own normalized state or measure and well-defined dynamics.

Second, a Turing machine is generally irreversible — it can erase, overwrite, and halt. The substratum bijection φ is reversible by Lemma 3 of Chapter 1: nothing is erased, nothing is created, there is no halting state. Reversible Turing machines (Bennett, Fredkin–Toffoli) are a well-studied subclass; the substratum sits in this class.

Third, a Turing machine computes an *extrinsic* function — input from an external user, output to an external observer. The substratum computes no extrinsic function. It is a closed permutation that cycles through its states; no input, no output, no halting condition. The appearance of dynamics, probability, particles, and forces is entirely the observer's perspective — what the permutation looks like through the partial window V.

The third difference is the load-bearing one. The framework extends Turing's question. Where Turing asked what computation a machine can perform for an external observer, the framework asks what computation looks like to a *component of the machine* — a subsystem with bounded access to the machine's own state. The answer, established in Chapter 1, is quantum mechanics.

This extension places the framework in a specific intellectual lineage. Three foundational results form what can be called the incompleteness family. Gödel: a formal system rich enough to encode arithmetic cannot prove all true statements about itself; the unprovable truths have rigid structure. Turing: a computer cannot decide all questions about its own behavior; the undecidable problems have rigid structure. Observational Incompleteness: an observer embedded in a deterministic system cannot access the complete state; the emergent description has rigid structure — quantum mechanics. The common pattern is self-reference under finite resources, with the impossibility result yielding not chaos but a precisely characterized formal structure.

Hidden-state identification is a record question in a specified model, not the Turing halting problem. The hidden state has a definite value, but whether an observer can determine it depends on the model and retained observations. In a known finite model, a quantity is determined by a nonempty record fibre exactly when it is constant on that fibre [Substratum Proposition 24.5]; two readouts of a visible/hidden swap, for example, identify the initial hidden coordinate. Independently, every finite observable law admits the exact fixed-basis unitary/Born representation of Chapter 1. This representation theorem does not rest on a blanket impossibility of hidden-state identification.

Two further comparisons deserve separation from unresolved hidden-state facts. An independently prepared finite factor evolving exactly independently of the observed system changes the deep-sector cardinality while preserving the original fixed-interface coherent channel [Substratum Theorem 24(iii)]. A slowly coupled deep sector preserves only the leading-order accessible-time law, with the stated correction. Infinite extensions require specified normalized states or measures and dynamics. Alphabet change preserves observables only under the alphabet-size hypothesis of Chapter 5. These are redundancies relative to particular data; they do not show that all inaccessible cardinalities, coupling structures or physical implementations lack determinate or observable consequences.

The reconstruction theorem developed in the next section fixes the **local lattice/gauge residue** — under its stated conditions — from observed physics plus structural assumptions. It does not uniquely determine a Bell-inclusive full substratum; H-Bell conditions the existence of that completion and its preparation-indexed Bell rule remains underdetermined.

2.4 The reconstruction theorem

Chapter 1 took a substratum (S, φ) as input and developed the emergent quantum description from it. The converse direction is necessarily scoped: the reconstruction theorem asks which **local propagating lattice/gauge data** observed physics fixes. It establishes uniqueness of that local residue's Stage-2 invariants under its stated conditions. Bell-inclusive completion data are separate: their existence is conditional on H-Bell and their uniqueness is not established.

The theorem takes two kinds of input. The first is *empirical*: structural facts about the observed universe. The second is *structural*: assumptions about the class of candidate substrates the reconstruction is permitted to consider. Both are necessary, and both must be stated explicitly. A reconstruction theorem that did not specify its structural assumptions would either claim too much (uniqueness without restriction) or claim too little (uniqueness only within an undefined class).

Empirical inputs. The reconstruction draws on seven structural inputs about observed physics, six established by experiment and one (E3) a theoretical bound:

(E1) Unitary quantum mechanics. The observed microscopic physics is described by states in a complex Hilbert space evolving unitarily, with observables as self-adjoint operators and measurement outcomes following the Born rule. This is Chapter 1's framework input.

(E2) Bell violations. Observed correlations between spatially separated measurements violate Bell-type inequalities, ruling out local hidden-variable theories satisfying factorizability. Confirmed experimentally since Aspect's 1982 measurements and refined in loophole-free tests since 2015.

(E3) Finite boundary entropy. The entropy of any bounded region of spacetime is finite and scales as the boundary area rather than the volume. This is the holographic bound — a theoretical bound rather than a direct observation — supported by black-hole thermodynamics and applied to the cosmological horizon.

(E4) Spatial isotropy. Observed physics is rotationally invariant; no spatial direction is preferred. Confirmed across observations from the CMB to laboratory tests of fundamental constants.

(E5) Propagating gravity. Gravitational degrees of freedom propagate as waves with the observed signature of two transverse polarization states, directly detected since LIGO 2015.

(E6) Stable matter. Atoms, planets, and bound states exist on observed timescales. This requires both gravitational and atomic-scale stability against the dynamics.

(E7) Boundary-entropy concordance. The cosmological parameter inferred from observation under the standard cosmological model is spatial flatness near the critical density: $\rho_s/\rho_{\text{crit}} \approx 1$. The framework's general-d form of this ratio is $\rho_s/\rho_{\text{crit}} = 2/(d-1)$, which equals unity only at $d = 3$.

Within A4's periodic-lattice class and the continuum conditions of SM §3.1–§3.2, E5–E7 enter the dimensional selection. The first two constraints already intersect at $d = 3$; the third is a consistency check. Propagating gravity (E5) requires $d \geq 3$, since the Weyl tensor governing gravitational-wave propagation vanishes in $d \leq 2$. Stable matter (E6) requires $d \leq 3$, since the Coulomb potential gives unstable atoms in $d \geq 4$ (Ehrenfest's theorem) and gravitational orbits are unstable in $d \geq 4$ (Tangherlini's result). Boundary-entropy concordance (E7) gives $d = 3$ exactly. Without stable matter no embedded observers exist; without propagating gravity no observable GR exists; without concordance the spatial-flatness data is unexplained. Within A4's periodic-lattice class, integer rank d is already an input. The filters select its value 3 on the stated continuum branch, subject to the continuum conditions of SM §3.1–§3.2.

The convergence has an external echo. A literature independent of this framework reaches the same selection from different starting points: Ehrenfest's stability analysis (atoms and planetary orbits are stable only for $d = 3$), the sharp-propagation property of the wave equation (Huygens' principle holds cleanly only in odd spatial dimensions, and distortion-free signal propagation singles out $d = 3$, as Hadamard's analysis of the wave equation shows), and Tegmark's predictability argument (field equations well-posed enough for observers to process information select $3 + 1$). These are convergent arguments rather than theorems, and they are not an independent axiom base in the sense of the reconstruction theorems of Chapter 1: like the framework's own three filters, they condition on the existence of stable structure and embedded observers. Their force is corroborative texture — several instruments, pointed from different premises within the same observer-admitting family, land on the same integer.

Structural assumptions. The reconstruction restricts attention to candidate substrates satisfying six properties:

(A1) Finiteness. The configuration space S is finite. Lemma 1 of Chapter 1 delivers finiteness of the observer's distinguishable states; the extension to all of S is a posit, read as the choice of the minimal finite representative and supported by E3 (the holographic bound read as a dimension cutoff).

(A2) Determinism. The dynamics $\varphi : S \to S$ is a bijection — deterministic and reversible. Its status is two-part (Chapter 1, Lemma 3): the bijective substratum is a representation choice — the minimal recurrent bijective representative of the hidden dynamics — and injectivity is anchored by a dilemma argument with one structural prong (finiteness and recurrence) and one empirical prong (the observed statistics).

(A3) Bounded coupling degree. Each site of the substratum is coupled to a bounded number of neighbors through φ. This is required for the emergence of a coupling graph with well-defined dimensional structure and underlies locality in the emergent description.

(A4) Center independence. The dynamics does not depend on a choice of preferred site or origin: φ commutes with lattice translations up to gauge (the coupling graph is a finite periodic lattice, the quotient of a rank-d lattice by a finite-index sublattice, d being its dimension), and the update at a site contains no explicit copy of that site's present value. The second property is what the derivation of the wave equation in Stage 2 below uses.

(A5) Linearity. The wave equation governing φ is linear. This is a sharpened stipulation: amplitude-scale gauge — the unobservability of the absolute field scale, an adjoined operational input rather than a theorem — restricts the rule to an affine one, and linearity is the further restriction; non-affine alternatives are excluded exactly to the extent that input is assumed, and would otherwise require a separate derivation.

(A6) Background independence. The dynamics is covariant under spatially varying internal-index transformations: a site-dependent transformation of the internal index is a symmetry provided the coupling data carried on the links are transformed with it. The content of the assumption is this covariant interface — the coupling is link-valued data on which the transformations act — and once it is in place the local transformation law imposes no further condition on the rule. It is not the stronger fixed-background condition that the same transformations leave the dynamics invariant with the coupling held fixed, which for an invertible constant coupling forces the transformation to agree across every coupled pair of sites and fails at the symmetric point of the coupling matrix for every transformation that differs across a coupled pair; the global commutant symmetry — the transformations constant across the lattice that leave the coupling unchanged — is the specialization from which the local gauge reading of Chapter 5 proceeds. The state-dependent coupling graph, under which the graph itself evolves with the state, is a separate principle that shares the name.

The reconstruction theorem's claim is that, under E1–E7 and A1–A6 with M1-T, C2-mixing and the nearest-neighbor restriction, the **local propagating lattice/gauge residue** is fixed, including the listed Standard-Model/discrete outputs and $\hbar = c^3\epsilon^2/(4G\sigma)$ with their stated qualifications. Bell violation is handled independently by M1-B: a common measurement-independent deterministic completion exists only conditional on H-Bell, and the theorem does not make that Bell-inclusive completion unique.

Stage 1: From observed quantum mechanics to a deterministic embedding. The inputs are the quantum-mechanical content E1, the Bell-violation content E2, the boundary-entropy bound E3, the finiteness and determinism assumptions A1, A2, and the two-clause modeling premise M1 — M1-T puts the temporal data used in Stage 1 in the history-readback sector, while M1-B puts Bell-violating composites on the measurement-independent, ontically parameter-dependent branch. Chapter 1's characterization theorem (§1.8) supplies the construction: any accessible non-Markovian process arises from marginalization of a deterministic bijection satisfying C1, C3 and C4 for some partition — and every finite observable law, memory-free sectors included, admits some deterministic realization (C2 outside the equivalence, [Main §3.4]). M1-T's observed temporal memory structure certifies the C1/C3/C4 history-readback sector, while M1-B's Bell violation independently requires Bell-relevant cross-wing coupling/ontic parameter dependence; the observed memory-bearing phenomenology — not quantum representability, which is universal — grounds C2 and C3, with C2's slow-bath necessity conditional on the hidden-sector mixing hypothesis (ETH its physical motivation) and its persistence form holding without it. The output of Stage 1 is a triple (S, φ, V) with S finite, φ a bijection, and $V \subset S$ the visible-sector partition, with the observed quantum dynamics reproduced as the marginalization of φ over the hidden sector $H = S \setminus V$.

At Stage 1, the Stinespring theorem controls only the **finite channel/dilation datum used by the temporal reconstruction**: its familiar dilation freedom lies in hidden-sector/gauge directions. It does not make the full triple (S, φ, V) unique across coherent composites, does not select a preparation-indexed Bell graph, and does not discharge the separate finite-predictive-dimension, local-tomography or operational-purification requirements of the full lift.

Stage 2: From deterministic embedding to specific lattice and gauge structure. Stage 2 combines Stage 1's output with the remaining empirical and structural inputs E4–E7 and A3–A6. Within A4's assumed periodic-lattice class, E5–E7 select $d = 3$ subject to the continuum conditions of SM §3.1–§3.2. The wave equation, internal-index structure, gauge construction, matter interpretation and discrete symmetries have their further hypotheses stated at the corresponding steps.

The dimension argument follows the three-filter convergence described above. Under A4 the coupling graph is a finite periodic lattice of some integer dimension d, an input of the assumption; statistical isotropy (E4) alone would not supply

it. The filters E5–E7 then select $d = 3$ on the stated continuum branch; they do not supply the integer-rank lattice assumption itself.

The wave equation follows from center independence (A4), isotropy (E4), and linearity (A5), together with the nearest-neighbor restriction (an assumption beyond A3's bounded coupling degree). The unique second-order reversible nearest-neighbor dynamics compatible with these constraints has the form $f = \alpha(x_1+x_2+\cdots+x_{2d}) \bmod q$, with coupling coefficient α. The coefficient is not the propagation speed: causal support travels at most one lattice edge per update whatever its value, and on the observer-level normalized branch it equals $1/d$.

The number of internal components per lattice site follows from the link count under H-link: $K = 2dm$, with $m = 1$ giving $K = 6$ (coupling-degree minimization does not establish it). This locks the candidate-sector count at three — read as three physical generations only under H-spin' (SM §4.7) — since three spin-1/2 staggered tastes emerge from the $K = 6$ minimum (Chapter 6 develops this).

The gauge group follows from the cubic-group decomposition of the six internal components. The cubic group acts on the $K = 6$ components with eigenvalue multiplicities $(3, 2, 1)$. The unitary group of the bicommutant of this action — the block algebra generated by the cubic-group representation — is $\mathrm{U}(3) \times \mathrm{U}(2) \times \mathrm{U}(1)$ on the corresponding eigenspaces. Amplitude-scale invariance (the field-value-rescaling gauge in $\mathcal{G}_{\mathrm{sub}}$, §2.5) removes only the common overall phase; the reduction to $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$, with the singlet-block direction as the hypercharge U(1), additionally uses a specified charged matter action and anomaly cancellation (Chapter 5 §5.6). Background independence (A6) then promotes the global $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ to local gauge invariance.

The matter content — three chiral fermion generations with anomaly-free hypercharges and one Higgs doublet — follows, on the single-copy H-link branch and conditionally on H-spin' and H-χ', from the staggered-fermion reduction, the link-carrier construction, partition-spinor identification, chirality emergence from the trace-out, and anomaly cancellation. Chapters 5 and 6 develop each step in detail; the chain ends with hypercharges $(Y_Q, Y_u, Y_d, Y_L, Y_e) = (1/6, 2/3, -1/3, -1/2, -1)$ uniquely determined on the nonzero-Y_Q branch, up to normalization.

T-invariance of the bijection φ combined with detailed balance gives reciprocal visible transition counts and narrows the QCD angle to $\theta \in \{0, \pi\}$; it does not fix $\bar{\theta}$, which is open (SM §5).

Uniqueness at Stage 2 follows from the uniqueness-given-inputs character of each step. Each derivation is the unique consistent choice given its inputs, the wave-equation step within the nearest-neighbor class and the hypercharge step apart from its exceptional branch; the chain is a composition of unique selections, and the output is unique within those qualifications.

Stage 3: From dynamics to emergent constants. Stage 3 takes Stage 2's

output and derives the emergent action-scale relation. The argument is that thermal self-consistency at the partition boundary — established in Chapter 7 from the boundary layer's emergent thermodynamics, conditional on H-slope with the horizon and frame conditions named there — forces the relation

$$\hbar = c^3\epsilon^2/(4G\sigma),$$

with the discreteness scale $\epsilon = 2\sqrt{\sigma}\,\ell_P$ (twice the Planck length at $\sigma = 1$, one nat per boundary mode) as the unique simultaneous solution, which fixes only the ratio $\epsilon/\sqrt{\sigma} = 2\,\ell_P$. With this calibration the boundary-mode count reproduces the Bekenstein-Hawking entropy $S = A/(4\ell_P^2)$, the 1/4 coefficient being the ratio $2\pi/8\pi$ of the KMS period to the first-law prefactor that the calibration matches.

Uniqueness at Stage 3 follows from the unique solvability of the boundary gap equation, which Chapter 7 develops.

The composite theorem. Composing the three stages yields the chapter's central result.

Reconstruction Theorem. *Let E1–E7, M1-T and M1-B, and A1–A6 hold, with the hidden-sector mixing hypothesis for the slow-bath form of the C2 necessity direction and, additionally, the nearest-neighbor restriction of Stage 2. Then the* ***local propagating lattice/gauge residue*** *is fixed: $d = 3$, $K = 6$, block multiplicities $(3, 2, 1)$, the stated Standard-Model gauge/representation and discrete-symmetry outputs — $K = 6$, the multiplicities and the gauge group conditionally on H-link with its single-copy clause and H-cust (Chapter 5 §§5.4–5.5) — and $\hbar = c^3\epsilon^2/(4G\sigma)$ with the qualifications carried by the cited derivations. A single measurement-independent deterministic completion also realizing the Bell-violating M1-B composites exists only conditional on H-Bell. Bell-inclusive full-substratum uniqueness is not established.* □

The theorem's content is striking. From the empirical inputs of observed physics — quantum mechanics, Bell violations, finite boundary entropy, spatial isotropy, propagating gravity, stable matter, spatial flatness — together with six structural assumptions and the nearest-neighbor restriction, the substratum is forced to be a specific kind of object with specific properties. Conditionally on H-link with its single-copy clause — the open hypothesis that the physical carrier is one copy of the six-link representation (Chapter 5 §5.4) — and, for the gauge group, on an octahedrally equivariant condensate (Chapter 5 §5.5), the Standard Model gauge group is not chosen; it is derived, and the three-sector structure is not posited; it is forced, its reading as three physical generations resting further on H-spin'. The Bekenstein-Hawking 1/4 factor is not a postulate; it is fixed by the horizon calibration under the conditions Chapter 7 names. The cosmological constant problem is not resolved by tuning; the framework's structure makes $\hbar$ and G commensurable through the boundary thermodynamics, and the apparent 10^{122} discrepancy is the compression ratio Chapter 7 develops.

Two scope remarks. First, uniqueness is within the structural class A1–A6

with the nearest-neighbor restriction. The theorem does not exclude alternative substrates outside this class: intrinsically continuum theories (violating A1), intrinsically stochastic theories (violating A2), theories with unbounded coupling (violating A3), or theories violating center independence, linearity, or background independence (violating A4–A6). The framework's defense of A1–A6 is that they are the natural assumptions given the empirical inputs, but this defense is an argument rather than a theorem. A separate analysis would be needed to rule out alternatives violating any A_i.

Second, the empirical inputs E1–E7 are stated as structural facts rather than as specific measurement outcomes. Operationally, no finite measurement set establishes any of them; finite data is consistent with them but does not uniquely require them. The reconstruction theorem holds for the inputs as stated — the idealized infinite-data regime — and proves **local-residue uniqueness** in that regime. The operational discrimination available against finite data comes from the cumulative weight of the framework's quantitative Standard-Model record (Chapter 6). This finite-data issue is distinct from Bell-composite underdetermination: Bell-inclusive uniqueness is open even in the idealized-input formulation.

The reconstruction theorem identifies the equivalence class $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$ as the framework's central object. The next section makes the gauge group $\mathcal{G}_{\text{sub}}$ explicit by identifying its generators; whether they exhaust the freedom in the reconstruction is open.

2.5 The substratum gauge group

The reconstruction theorem identifies $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$ as the framework's central object. This section makes $\mathcal{G}_{\text{sub}}$ explicit by exhibiting three families of generators; whether they exhaust the group is open.

For the reconstructed dilation/local datum, observational equivalence is relative to a specified visible/hidden interface, preparations and readouts. Equality of the coherent channel family is stronger than equality of its rooted population transition probabilities. The representative freedoms denoted by $\mathcal{G}_{\text{sub}}$ preserve the particular data stated in Substratum Theorem 24: uniform hidden-coordinate relabelling and independently prepared finite decoupled enlargement preserve the coherent family; alphabet change retains its separate hypothesis. This is not a classification over all partitions of a structural class.

The gauge group is not a symmetry of a Lagrangian or an action. There is no Lagrangian at the substratum level; the substratum is a finite set with a bijection on it. $\mathcal{G}_{\text{sub}}$ is a symmetry of the *equivalence class of substrata* defined by the observability condition. The emergent gauge symmetries (the Standard Model gauge group, the diagonal phase ambiguity in the emergent Hamiltonian) are Lagrangian symmetries in the standard sense, derived from the substratum

through the trace-out.

Theorem (Generators). *$\mathcal{G}_{sub}$ contains the independent families of transformations (i) and (iii) below, and, under the alphabet-size hypothesis of Chapter 5, the family (ii); (iv) is a structural graph comparison, not a family of $\mathcal{G}_{sub}$:*

(i) Hidden-state relabeling. For any bijection σ of the hidden configurations $\mathcal{C}_H$, applied in every visible fibre as $\pi(v,h) = (v, \sigma(h))$, the substratum $(S, \pi \circ \varphi \circ \pi^{-1})$ produces the same emergent physics as (S, φ). The transition probabilities depend on the coupling structure of φ — which states are sent to which — but not on which labels are attached to which hidden states; a relabeling $(v,h) \mapsto (v, \sigma_v(h))$ that differs between visible fibres preserves the transition probabilities $T_{ij}(t)$ but not, in general, the coherent channels. This is an $|\mathcal{C}_H|!$-element subgroup, vastly larger than any gauge group appearing in the Standard Model.

(ii) Alphabet change (under the alphabet-size hypothesis). The local state space at each lattice site can be replaced by $\mathbb{Z}/q'\mathbb{Z}$ for any integer $q' \geq 2$, while preserving the coupling graph and the dynamics class, without changing any observable if the alphabet-size hypothesis of Chapter 5 holds. The internal alphabet at a lattice site is the number of distinguishable local states; some results hold for every $q \geq 2$ and others carry q (Chapter 5). This family is parametrized by the integers $q \geq 2$.

(iii) Finite decoupled ancillary enlargement. Adjoin a finite factor to the deep hidden sector, prepare it independently in a normalized density matrix, and keep its dynamics exactly decoupled from the original system. At the fixed visible/hidden interface, tracing out the factor preserves the full original coherent-channel family and therefore every rooted population law [Substratum Theorem 24(iii)]. This is stronger than a population-only assertion. Infinite factors require a separately specified normalized state or measure and well-defined dynamics; finite uniform counting does not supply that extension. Slowly coupled enlargements instead change the law by the correction stated in Chapter 7 and preserve only its leading-order accessible-time description.

(iv) Graph isomorphism (up to statistical isotropy), with a gauge-sector qualification. Two coupling graphs G_φ and $G_{\varphi'}$ that are quasi-isometric, with the same polynomial growth exponent d, the same spectral mode count, and the same statistical isotropy at large scales, each admit a finite-law representation and share the growth exponent d, but need not give the same transition probabilities $T_{ij}(t)$ at a fixed observer interface, and the continuum-curvature step of the gravitational construction has its own conditions (§3.1 of the SM paper); in that sense the regular cubic lattice $\mathbb{Z}^3$ and a bounded-degree random graph with $d = 3$ polynomial growth and statistical isotropy are structurally comparable, not observationally equivalent. The *Standard Model gauge group* is where this crystallography is read out: its multiplicities $(3, 2, 1)$ come from the octahedral point group on the six link directions, which is not a quasi-isometry invariant, so only the spectral mode count $2d_s = 6$ transfers to a generic isotropic graph — not the cubic decomposition. For the gauge sector, therefore, comparison (iv)

is restricted to orientation- and octahedral-structure-preserving isomorphisms, and the simple-cubic Bravais type is an empirically-selected input rather than a gauge choice. The gauge group itself is a universality-class invariant, shared with any Standard-Model-reproducing class through different machinery ([Structure §2.3, §14.4]); within OI's class it is realized through the cubic-group commutant.

Proof for the specified data under the stated hypotheses. Generator (i): conjugation by the hidden permutation leaves the uniform hidden prior invariant and cancels under the partial trace, so every visible channel is unchanged; a fibre-dependent relabeling fixes the visible coordinate and permutes each fibre's uniform hidden prior bijectively, so every transition probability is unchanged. Generator (ii): this is the alphabet-size hypothesis of Chapter 5, under which the family is stated. Generator (iii): the product dynamics sends the independent ancillary state σ_A to $W^t \sigma_A W^{\dagger t}$. Its trace remains one, so the partial trace over it returns the original reduced channel exactly, as in [Substratum Theorem 24(iii)]. Comparison (iv): the finite-law representation holds for any finite bijection and the growth exponent is shared by hypothesis; the transition probabilities depend on the specific graph. □

Completeness. For the reconstructed dilation/local datum, it is open whether families (i)–(iii) exhaust $\mathcal{G}_{\text{sub}}$ — whether any observables-preserving transformation of that scoped datum decomposes as a composition of generators (i)–(iii). This completeness question does not extend to preparation-indexed Bell adjacency or other Bell-composite data.

Three candidate further generators are compared with the three already listed:

Time reversal ($\varphi \to \varphi^{-1}$). The wave equation that selects φ is T-invariant, giving $\varphi^{-1} = T \circ \varphi \circ T^{-1}$ where T is the phase-space layer swap exchanging successive timesteps. Since T acts on the visible layers, this is not generator (i).

Hidden-sector dynamical reparametrization beyond enlargement. Equality of the visible channel family does not force a hidden-sector unitary relating two dilations of equal hidden-sector dimension: two permutation systems on $\mathbb{Z}_2 \times \mathbb{Z}_3$ have equal visible channels at every time without being unitarily conjugate (Substratum §4). Whether such reparametrizations reduce to generator (i) is the open completeness question.

Visible-sector emergent global phase ($U \to e^{i\theta} U$). At the substratum level, S is a finite set with no complex structure; the emergent phase corresponds to no substratum-level transformation. It is the identity on (S, φ).

The three-level gauge hierarchy. The framework has three distinct gauge symmetries, each appearing at a different level of the derivation, with its own equivalence notion:

Level 3 — reconstructed-datum substratum gauge. $\mathcal{G}_{\text{sub}}$ acts on the reconstructed dilation/local representative (S, φ) before any marginalization. Within that scoped reconstruction it includes transformations with no analog at lower levels

(deep-sector enlargement, and alphabet change under the alphabet-size hypothesis). Bell-composite preparation rules are outside its scope.

Level 2 — emergent quantum field theory gauge. The Standard Model gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ acts on the emergent fields after Stage 2 of the reconstruction has produced them. It follows from the substratum construction (Stage 2) and acts on internal components within the blocks of the equivariant condensate Σ.

Level 1 — emergent Hamiltonian gauge. A residual diagonal-unitary freedom $H \to DHD^\dagger$ (where D is a diagonal unitary on the emergent Hilbert space) remains after all transition-probability data has been extracted. It corresponds to the rephasing of basis states in the emergent quantum description.

The levels carry distinct equivalence notions; inclusions between them and a map from $\mathcal{G}_{\mathrm{sub}}$ onto the Standard Model gauge group are not established.

The substratum construction provides a structural account of why the Standard Model gauge group is what it is. The conventional puzzle is: why $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ rather than some other group? Standard Grand Unified Theories address this by embedding the Standard Model in a larger simple group at high energy and then breaking it; the framework here answers differently. The Standard Model gauge group follows from the substratum construction (Stage 2) once the cubic lattice is selected from observation and conditionally on the carrier bridges H-link and H-cust (Chapter 5 §§5.4–5.5). Within that class the Standard Model gauge group is fixed by the structure of the substratum, not chosen from a landscape.

Substrate freedom and the realism question. A uniform hidden-coordinate relabelling, with the uniform hidden prior, and an independently prepared finite exactly decoupled ancillary factor preserve the coherent channel at the fixed visible interface. Alphabet change carries its stated alphabet-size hypothesis. Decisions based only on the preserved data cannot distinguish the corresponding descriptions. A different permitted interface, preparation or coupling may ask a different observational question. Realism about the reconstructed local invariants therefore does not amount to a proof that every possible implementation or measurement belongs to one equivalence class.

This is what makes the framework's scoped structural realism non-trivial. The invariants of the **reconstructed local lattice/gauge residue** are uniquely determined — under the reconstruction theorem's stated conditions — by observation plus structural assumptions. Individual representatives of that local residue related by the generators of $\mathcal{G}_{\mathrm{sub}}$ are gauge expressions of the same reconstructed physical content, like coordinate choices on a manifold or specific representatives of a homotopy class. Bell-composite preparation data lie outside this uniqueness statement: H-Bell conditions compatibility/existence, and no result here selects a unique Bell-inclusive full representative.

The reconstruction theorem identifies the **local lattice/gauge residue**; the

gauge group identifies what within that scoped reconstruction has no observable content. With both in place, observed physics fixes that local residue under the theorem's stated conditions. H-Bell separately conditions Bell-inclusive existence, and Bell-inclusive uniqueness is not established. The next section turns to the synthesis: how quantum mechanics and general relativity relate to the substratum as projections of the same construction, and what a third projection establishes about the arrow of time.

2.6 The three projections

The reconstruction theorem and the substratum gauge group have established what (S, φ) is and what equivalences leave its observables unchanged. The question taken up in this section is what relationship (S, φ) bears to the framework's three large emergent structures: quantum mechanics, general relativity, and the arrow of time.

The framework's claim is that quantum mechanics and general relativity are not two independent theories. Conditional on H-Bell at the Bell/local-lattice interface, they are projections of the same finite deterministic construction, and a third projection bears on the arrow of time. Each projection applies the same trace-out machinery to the same (S, φ) along a different axis, and each produces a distinct emergent structure as the observer sees the substratum from inside.

The visible-sector projection: quantum mechanics. Chapter 1 developed this projection in detail. At finite observable horizon the reduced stochastic law, a finite reversible realization, and a fixed-basis unitary/Born representation are exactly equivalent: $S \iff D \iff Q_{\mathrm{fb}}$. C4 readback then identifies the memory-bearing sector by forcing hidden predictive memory. In the spatial construction the coupling graph supplies exact causal cones for finite-range substrata. Bell is separate from the full-QM lift: a measurement-independent response-complete deterministic completion with parameter independence and a ± 1 outcome in every run would obey $|S_{\mathrm{CHSH}}| \leq 2$, so the framework's Bell-violating branch is ontically parameter-dependent with operational no-signaling required. Two independent qualifications remain beyond the finite-law equivalence: the simultaneous coherent instrument/composite lift ([Main §3.4]) and H-Bell, the compatibility of the local lattice sector with the ontically parameter-dependent Bell completion. Applied to the local lattice sector constructed in this chapter — a $d = 3$ cubic lattice with $K = 6$ internal components and the wave equation as its local propagating dynamics — the visible-sector projection produces the Standard Model: the gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ from the stabilizer of the cubic-equivariant condensate, three chiral fermion generations, one composite Higgs, and anomaly-free hypercharges. Chapters 5 and 6 develop the Standard Model derivation; Chapter 8 develops the specific JUNO neutrino measurement that the framework's parameter-free PMNS-angle value matches at 0.07σ (a retrodiction). The local nearest-neighbor sector cannot it-

self supply Bell parameter dependence for wings with one-step own-wing setting interventions outside each other's graph cone under measurement independence; identifying it with one Bell-violating total completion is the open H-Bell compatibility hypothesis.

The boundary-thermodynamic projection: general relativity. A second projection takes the same substratum but applies the trace-out machinery to the partition boundary rather than to the bulk. The boundary of the visible sector — the surface across which the coupling H_{int} acts — carries its own thermodynamic structure: a temperature set by the boundary's classical surface gravity, an entropy set by the number of boundary modes, and a dynamical evolution set by the coupling between boundary and bulk. Jacobson's thermodynamic argument applied to local causal horizons yields Einstein's equations as the Clausius relation $dE = T\,dS$ at the boundary.

Applied to the cosmological horizon as a causal partition, the boundary-thermodynamic projection produces general relativity with quantitative predictions. The emergent action scale $\hbar = c^3\epsilon^2/(4G\sigma)$, with $\epsilon = 2\sqrt{\sigma}\,\ell_P$, follows from thermal self-consistency at the boundary layer under the infrared detailed-balance condition H-slope. The boundary-mode count reproduces the Bekenstein-Hawking entropy $S = A/(4\ell_P^2)$, the 1/4 coefficient being the ratio $2\pi/8\pi$ that the calibration matches and conditional on the horizon conditions of Chapter 7; GW250114 (September 2025) confirms the classical area theorem, which the framework preserves — the 1/4 coefficient itself has no direct empirical test. The cosmological constant problem receives a conditional treatment: the quantum vacuum energy and the effective Λ are properties of logically distinct levels of description rather than commensurable quantities whose 10^{122} ratio requires cancellation; the dissolution is conditional on the covariant coupling step (Chapter 7). The structured component of the boundary entropy produces dark matter phenomenology with MOND acceleration scale $a_0 = cH/6 \approx 1.1 \times 10^{-10}$ m/s^2 (the dimensional factor imported from Verlinde's emergent-gravity analysis). Chapter 7 develops each of these.

The horizon-clock projection: the arrow of time. A third projection applies the trace-out machinery to the horizon-growth parameter along an observer worldline. The substratum is time-symmetric: φ and φ^{-1} are equally valid as forward-time maps, and there is no preferred direction at the substratum level. Condition C2 — persistence, in its slow-bath realization — is a retention condition: it requires the hidden sector's correlations with the visible sector's past to persist to readback, and the coarse-grained mixing time it constrains is the same in the two directions; the direction in which records are written and read is fixed by a preparation and the observer's orientation convention. The structural observer-selection theorem developed in Chapter 1's appendix material — under the effective-mixing hypothesis, for sufficiently fast-mixing equilibria ($\tau_{\text{mix}} \ll \tau_S$), and in the coarse-grained description generated by iterating the one-step effective channel ([Main §4.6]) — restricts observers satisfying C1–C4,

with C2 read in its timescale form (record persistence and readback there being suppressed, not excluded), to the non-equilibrium phase of (S, φ); the restriction is time-symmetric and selects where observers can exist, not a direction. On an expanding cosmological background satisfying the null energy condition — inputs, not derived — the area $A(t)$ of the apparent horizon, and the horizon entropy $S_{\mathrm{dS}}(t) = A(t)/(4\ell_P^2)$ assigned to it at the de Sitter calibration, is non-decreasing, strictly increasing while $\rho + p/c^2 > 0$ and constant in exact de Sitter. That the selected non-equilibrium phase is ordered by this growth is not established.

The apparent-horizon area provides a structural clock on that background. Every comoving observer of an expanding homogeneous general-relativistic background satisfying the null energy condition agrees that it does not decrease — standard background kinematics, not a prediction derived from the substratum — and the criterion is expansion with the null energy condition, not acceleration: in exact de Sitter the area is constant. Whether the other arrows of time align with this clock is not established. Jacobson's coupling relates the boundary energy density to the expansion rate, not matter entropy to horizon area, so the direction of the thermodynamic arrow rests on a low-entropy or product boundary condition, and the generalized second law is open. C2 — persistence, realized by the slow bath — holds the same way in both directions; the direction of the memory arrow comes from a preparation and the observer's orientation convention. The measurement arrow — the observer records past outcomes and prepares future states — inherits the memory arrow's direction. The chain uses one input (horizon growth on the expanding branch) and one structural constraint (C1–C4 observer selection, conditional on the effective-mixing hypothesis of [Main §4.6] and stated for the coarse-grained description generated by iterating the one-step effective channel), which fixes where observers can exist, not a direction. No independent past-hypothesis axiom is required to select where observers exist, and Boltzmann brains are excluded structurally rather than anthropically.

The framework's content here is precise: on the expanding branch it establishes a non-decreasing horizon parameter along observer worldlines, and it does not establish that the four arrows of time align. Nor does it derive the convention of which direction along the horizon-clock parameter to call "future." The two directions are related by the layer swap at the substrate level ($\varphi \leftrightarrow \varphi^{-1}$), and the observer's choice of convention is residual freedom. The parameter is conditional on the expanding branch; the labeling is an observer-imposed convention. There is one further qualification worth stating: a monotonic parameter is monotonic with respect to an *ordering* of the worldline's moments, and the framework's foundational treatment axiomatizes that there is more than one moment while leaving the ordering of moments as a separate, open question. The result here is therefore conditional on that ordering as well — it establishes that ordered moments on the expanding branch yield a non-decreasing parameter, not that directedness arises from the substratum with no assumption about ordering at all. CP violation in CKM and kaon/B systems is a flavor-basis feature of the

specific bijection φ rather than a consequence of the time-arrow cascade — the framework treats flavor-sector CP phases as solution-specific inputs that do not affect the cascade's structural content.

The common structure. The three projections share three features. Under H-Bell they apply to the same total substratum (S, φ); without H-Bell, the local SM/GR lattice sector and the Bell-violating completion have not been shown to be one object. They use the same trace-out machinery — marginalization over the hidden sector under conditions C1–C4. They produce derived rather than fundamental structures: nothing in quantum mechanics, general relativity, or the arrow of time is a feature of (S, φ) in itself; each is a feature of how the embedded observer sees the substratum from inside. The differences among the three projections lie entirely in which structure of the embedded observer's description they extract. The visible-sector projection works on the bulk dynamics and produces the reduced finite law — universally admitting the quantum representation — with the Standard Model as its specific content. The boundary-thermodynamic projection works on the classical thermal data at the partition boundary and produces general relativity with its gravitational and cosmological predictions. The horizon-clock projection works on the horizon-growth parameter along observer worldlines together with the observer-selection rule; the alignment of the arrows with that parameter is open.

The shared structure is not metaphor or analogy. The same local lattice sector that gives the Standard Model gauge group from the stabilizer of the equivariant condensate gives the Bekenstein-Hawking entropy as the boundary mode count. H-Bell is required only for identifying that local sector with the Bell-violating total completion. The conditions C1–C4 that characterize the memory-bearing sector — whose conditional memory readback makes measurable — for the visible-sector projection produce the persistence-based observer selection for the horizon-clock projection. The substratum gauge group $\mathcal{G}_{\text{sub}}$ controls the gauge equivalences of the **locally reconstructed residue** across the three projections; it does not classify all preparation-indexed Bell-composite rules. The framework's claim is that the three projections are co-derived from a single construction, not three separate applications of similar machinery.

2.7 The QM-GR incompatibility as a category error

The conventional formulation of the QM-GR unification problem treats quantum mechanics and general relativity as two competing theories in the same logical category — two attempts to describe the same regime, awaiting reconciliation in a deeper unified theory. Every program of unification proceeds from this assumption. Quantum gravity programs treat the metric as a quantum field to be quantized. Geometric programs treat the wave function as a structure on the spacetime manifold. Emergent programs treat one of the two as derived from a deeper substrate that recovers the other. The persistent failure of all three approaches to produce a quantitatively confirmed unification — half a century of intensive work without an empirical test of the unification regime

— suggests that the underlying assumption may be wrong.

The framework's claim is that quantum mechanics and general relativity are not two theories of the same regime at all. They are two projections of the same construction, viewed at different levels and through different machinery. The visible-sector projection that produces the memory-bearing dynamics quantum mechanics represents and the boundary-thermodynamic projection that produces general relativity refer to *different objects* in the construction. The wave function is a property of the embedded observer's compressed description of the visible sector — the bookkeeping device that tracks what the observer knows and does not know about a system. The metric is a property of the classical thermodynamics of the partition boundary — the geometric content of the surface across which the trace-out is performed. These are not two descriptions of the same physical system at different levels of approximation. They are descriptions of two different substructures of the same total object (S, φ), derived by different projection maps from a single source.

The structural analogy that captures the relationship most clearly is the relationship between thermodynamics and statistical mechanics. Thermodynamics deals with pressure, temperature, and entropy. Statistical mechanics deals with particle positions, momenta, and microstate counting. These are not two competing theories of the same gas, and the question "how do we unify thermodynamics and statistical mechanics?" is not a coherent question. They describe different things about the same system. The relationship between them is one of projection — statistical mechanics produces thermodynamic quantities by averaging over microstates — rather than unification. Asking how to unify thermodynamics and statistical mechanics into a deeper theory mistakes the structure of their relationship.

The relationship between quantum mechanics and general relativity in the framework is structurally identical. General relativity describes the thermodynamic projection of (S, φ): the boundary geometry, the surface gravity, the entropy, the dynamical evolution of the partition under coupling to the bulk. The quantum representation describes the statistical projection: the visible-sector dynamics under marginalization over the hidden sector — universal over finite laws, with nonzero conditional memory marking the memory-bearing sector and the unitary evolution capturing its non-Markovian structure. The two are related by a definite procedure (the trace-out, with C1/C3/C4 diagnosing where the memory sector is nontrivial), with the substratum gauge group controlling the equivalences. The question "how do we unify them?" is recast as a category error rather than answered, conditionally on the open covariant rung G3 (Chapter 7 §7.7; Chapter 19 §19.2.1).

This is not the same as saying that quantum mechanics is more fundamental than general relativity, or vice versa. Both are emergent. Both depend on the trace-out and the partition. Both are derived structures rather than features of the underlying reality. The fundamental object is (S, φ) — a finite set with a deterministic bijection on it — and neither quantum mechanics nor general

relativity is a feature of (S, φ) itself. They are features of how an embedded observer or a horizon-bounded thermodynamic regime sees (S, φ) from inside. (Nor is the ranking question recoverable from inside: by the factoring argument of Chapter 7, no criterion grounded in the accessible observations can separate two descriptions faithful to the same data — the fundamentality question is answered at the level of (S, φ), not adjudicated between its projections.)

The recasting of the unification problem has two consequences worth marking explicitly. First, it recasts the QM-GR unification problem as a category error between two projections of one object, conditionally on the open covariant rung G3 (Chapter 7 §7.7; Chapter 19 §19.2.1). The persistent failure of unification programs is no longer puzzling — they were trying to merge two descriptions of different things, which cannot be done. Second, it provides an explanation for why the Standard Model has the gauge group it does. The Standard Model gauge group follows from the substratum construction (Stage 2) conditionally on the carrier bridges H-link and H-cust (Chapter 5 §§5.4–5.5). The gauge group is not chosen from a landscape of alternatives; it is obtained from the stabilizer of the cubic-equivariant condensate, given the reconstruction theorem's structural inputs and the cubic lattice selected from observation.

The framework's synthesis claim is structural and architectural, not predictive in the conventional sense. It does not claim to produce gravitational phenomena that conventional quantum field theory plus general relativity cannot reproduce. The framework's empirical content is concentrated elsewhere: in the Standard Model derivation (twenty-two quantitative retrodictions, Chapters 5 and 6), the JUNO neutrino prediction (Chapter 8), and the dark-sector predictions specific to the boundary-thermodynamic projection (Chapter 7). What the synthesis claim establishes is that these predictions are not independent. The same construction that produces the Standard Model produces the gravitational sector. The same trace-out that gives the Standard Model gauge group from the stabilizer of the equivariant condensate gives the Bekenstein-Hawking entropy as the boundary mode count. The framework's derivations across Chapters 5, 6, and 7 are not separate applications of similar machinery; they are co-derived consequences of a single construction.

2.8 What follows

The reconstruction theorem and the substratum gauge group establish the substratum side of the framework. The three projections establish its relationship to quantum mechanics and general relativity, and what it establishes about the arrow of time. With both in place, the technical core of Part I is complete; the chapters that follow develop its consequences in three directions.

Chapter 3 develops the framework's philosophical positioning. The substratum's ontological status — what it means to commit to the equivalence class $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$ as the framework's metaphysical object — is articulated in its full hierarchical form, with realism operating at multiple levels of the observation

hierarchy and the gauge hierarchy. The framework's relation to Bell, Kochen-Specker and PBR-type no-go theorems is developed through separate checks of their local-response, operational-algebra and preparation premises, with the stated unresolved obligations retained. The framework's intellectual position is located against the contemporary observer-dependence programs (QBism, relational quantum mechanics, the recent CLPW and Maldacena work on observer-incorporated entropy) and against the older philosophical lineage (Cusa's docta ignorantia, Spinoza's attribute theory, Worrall's epistemic structural realism). The relationship to Hofstadter's strange-loop framework is developed in detail.

Chapter 4 articulates the framework's methodological discipline. The single-foundational-commitment derivation pattern — that the framework starts from one premise (observation occurs) and derives the rest as theorems — is examined as a methodological standard. The architecture-level engagement with the no-go results is developed: the substratum, observable law and trace-out identify the objects constrained by Bell, Kochen-Specker, PBR and Frauchiger-Renner, while each theorem retains its own premise checks, including the open H-Bell realization and the stated operational and preparation assumptions. The classification of claims by derivational rigor — theorems, derivations, structural retrodictions, and informed speculation — organizes the framework's program by epistemic status. The framework's pre-registration of falsification conditions is discussed as a methodological commitment.

Parts II and III then develop the framework's quantitative consequences. Chapter 5 develops the Standard Model gauge group from the cubic-lattice substratum: the wave equation, the link count under H-link, the cubic-group commutant decomposition, chirality from the trace-out, anomaly-free hypercharges. Chapter 6 develops the matter content and quantitative predictions: three generations (physical reading conditional on H-spin', SM §4.7; chirality on H-χ', SM §4.8), the composite Higgs, gauge couplings, and the twenty-two classified quantitative retrodictions (two parameter-free structural and five conditional structural; Appendix A). Chapter 7 develops the gravitational sector: the boundary thermodynamics, the calibration $\hbar = c^3\epsilon^2/(4G\sigma)$, the Bekenstein-Hawking 1/4 factor, the conditional dissolution of the cosmological constant problem, the dark sector predictions. Chapter 8 develops the JUNO neutrino prediction in detail. Chapter 9 takes up universality and the framework's empirical scope.

The cascade through chemistry, biology, evolution, and intelligence in Chapters 10–12 takes the structural results of Parts I and II as derived inputs and develops their consequences for the structural conditions under which complex matter, life, and intelligence become possible. The applications in Chapters 13–17 develop the framework's reach into working scientific fields: quantum biology, quantum computing and engineering, medicine, bioinformatics.

The framework's claim, made precise across these chapters, is that a single starting commitment — that observation occurs — combined with the lemmas extracted from this commitment and the conditions identified for embedded observers in our universe, produces a substratum that is forced to have a specific

architecture, with quantum mechanics, general relativity, the Standard Model, and the cosmological dark sector all forced consequences of the architecture rather than separate empirical inputs. The book's role is to make this chain of derivations available for evaluation against the experimental record now being assembled and forthcoming.

Chapter 3

Hierarchical Structural Realism

3.1 What this chapter develops

Chapter 2 established the framework's metaphysical commitment as ontic structural realism on the equivalence class $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$. What is real at the substratum level is the class's reconstructed local residue (§3.5), not any specific bijection within it. The chapter argued for this commitment in its own terms: stronger claims commit to features the gauge group identifies as unobservable, weaker claims undersell the reconstruction theorem's uniqueness, and the class's local invariants are what observation plus structural assumptions, under Theorem 23's stated conditions, force.

That argument is correct but incomplete. The framework's structural-realism position has more structure than the single commitment Chapter 2 articulated. Different parts of the framework's content live at different levels of generality, with different empirical predictions, different falsifiability conditions, and different philosophical commitments at each level. The framework's twenty-two Standard Model retrodictions live at one level; the central claim that embedded observers' statistics universally admit the quantum decompression — a representation, explicitly non-unique — lives at a broader level; the foundational fact that observation occurs lives at the broadest level. A single structural-realism commitment that applied to all of these uniformly would either claim too much at the foundational level or too little at the specific predictive level.

This chapter develops the framework's structural realism in its full hierarchical form. The structural-realism position operates on a two-dimensional hierarchy: an *observation axis* running from the framework's foundational commitment to its specific predictive content, and an orthogonal *gauge axis* running from narrow gauge equivalences within the emergent quantum mechanics to broad equivalences across structural classes of embedded observers. Different empirical predictions live at different intersections of these axes; falsifiability is stratified accordingly; and the framework's strongest realism claim — the Chapter 2 commitment to $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$ — is what the hierarchy locates at a specific intersection.

The chapter has four pieces of work. §§3.2–3.3 develop the observation axis and the observer-admission distinction that the framework's third level introduces. §§3.4–3.5 develop the gauge axis, its orthogonality with the observation axis, and the resulting multi-level structural realism with its stratified predictions. §3.6 identifies gauge freedom with observational incompleteness on a precisely delimited part of its range, distinguishing the two species of incompleteness this produces. §§3.7–3.8 develop the framework's relationship to the contemporary no-go theorem literature and to the older philosophical lineage of incompleteness-and-perspective accounts. §3.9 closes the chapter and points forward to Chapter 4's methodology and to Chapter 8's universality and cross-framework analysis.

3.2 The observation axis

The framework's content nests in four levels of increasing specificity along an axis I will call the *observation axis*. Each deeper level is a more specific commitment about what the framework asserts. At the broadest level, the framework commits only to the fact that observation occurs; at the narrowest level, it commits to a specific lattice structure with twenty-two quantitative retrodictions. Between these poles lie two further levels — observer admission and universality classes — that articulate intermediate structural content.

Level A: the observation axiom. The framework's foundational commitment, developed in Chapter 1, is that observation occurs: an observer records distinguishable outcomes of interactions with a system not wholly under the observer's control. Its indubitable core — that some differentiated content is registered — is the Cartesian *cogito*, made mathematically precise and without its thinking subject; it is the one premise the framework takes from outside its own theorems. The structure Level A adds to that core is that of the Chapter 1 definition, and it is physical, not mental. Level A commits to three things: that there is something to observe (a substrate of reality structured to support distinguishable states); that some part of reality is structured to record outcomes (observers are embedded in reality, not external to it); and that observation has structural consequences (the recording itself is a physical event, not a passive registration).

Level A is shared with the broader observer-essentiality program in contemporary foundations of physics. The work of Chandrasekaran, Longo, Penington, and Witten on observer-incorporated entropy in de Sitter space (2022), Maldacena's analysis of observer's-clock dependence in cosmological partition functions (2024), and Harlow, Usatyuk, and Zhao on observer-dependent Hilbert spaces in closed universes (2025) all start from analogous commitments. The framework's distinguishing feature at Level A is not the commitment itself but what is derived from it: a finite deterministic substratum without extra dimensions, without intrinsic stochasticity, and without additional postulates beyond those identified in Chapter 1's structural lemmas and posit A1 of Chapter 2.

Level B: observer admission. Not every conceivable substratum admits

embedded observation. Chapter 1 identified four conditions — C1 (non-trivial coupling), C2 (record persistence — slow-bath separation one route), C3 (sufficient hidden-sector capacity), and C4 (history readback) — that are necessary for a substratum's bijection φ to support an embedded observer. A substratum (S, φ) is *observer-admitting* if there exists a partition $V \subset S$ such that the triple (S, φ, V) satisfies C1–C4; it is *observer-lacking* otherwise.

The distinction is structural rather than empirical. A finite cellular automaton with a deterministic update rule may or may not admit an embedded observer, depending on whether its dynamics support the slow-mixing, sufficient-capacity, coupled, read-back structure C1–C4 require. Random permutations on finite sets typically lack this structure: their cycle decompositions are wrong for the persistence condition (no structure realizing a slow bath or a conserve-and-route alternative), their internal correlations are wrong for the capacity condition, or their coupling is too uniform or too sparse for the non-trivial-coupling condition. The space of substrata is structurally partitioned into the observer-admitting subset, which can support embedded observers, and the observer-lacking subset, which cannot.

Level B commits to a structural-anthropic reading that sharpens the conventional anthropic principle. Standard anthropic reasoning treats fundamental parameters as contingent and explained by selection: we observe a universe with parameters compatible with us because we could not observe a universe without them. The Level B reading is sharper. Observer-lacking substrata are not candidates for "our universe" at all — they are mathematical objects of a different kind, lacking the observer-extraction structure that would make the question "what would observers in this universe see?" coherent. Within the observer-admitting subset, anthropic reasoning operates as usual; across the observer-admitting / observer-lacking boundary, the question of why we observe a universe with our particular features becomes the question of why a universe must be observer-admitting at all. The framework's answer is partly tautological — we are observers, so the universe we observe must admit observers — but the tautology is informative: it identifies which structural features of a candidate universe are forced by the existence of observers, and which are contingent.

The structural-anthropic content of Level B is independent of the specific universality class the universe belongs to. Any candidate substratum, regardless of dimension, gauge structure, or matter content, must satisfy C1–C4 to admit observers. The Level B commitment is therefore broader than any specific physics: it applies to OI's cubic-lattice substratum, to string-theoretic compactifications insofar as they admit embedded observers, to any future candidate fundamental theory, and to mathematical structures that may or may not correspond to physical universes.

Level C: universality classes. Within the observer-admitting subset, equivalence classes arise from the partial-trace observational features that any embedded observer produces. Two substratum-with-observer systems are *observationally equivalent* if their algebra-channel pairs — the algebra of visible-sector

observables together with the quantum channel induced by the partial trace over the hidden sector — are isomorphic in a precise sense that preserves both the algebraic structure and the channel's dynamical content. A *universality class* is an equivalence class under this observational equivalence.

The universality-class structure is broader than the substratum gauge group of Chapter 2. The substratum gauge group $\mathcal{G}_{\text{sub}}$ relates some observationally equivalent substrata *within OI's specific structural class* — substrata satisfying A1–A6 of the reconstruction theorem; whether the stated generators exhaust the relevant equivalences is open. The universality-class equivalence operates more broadly: substrata in different structural classes can be observationally equivalent if they produce the same algebra-channel pair through structurally different machinery. Two substrata that are universality-class equivalent need not be related by any $\mathcal{G}_{\text{sub}}$ transformation; the broader equivalence allows for structural-class transitions that the substratum gauge group does not.

Universality classes have internal sub-structure at three levels of refinement. The most universal level is the *partial-trace features sub-class*, characterized by the structural features any partial-trace observation must produce regardless of substratum specifics: the Born rule, channel-level unitarity to leading order, memory/divisibility — accessible memory where the hidden sector reads back its record (C4), and, on any finite reversible substratum, recurrence-cycle P-indivisibility of the marginal stochastic process wherever a rooted visible map is not a permutation — and the commutant restriction pattern that selects which symmetries survive the trace-out. Below this is the *gauge-group sub-class*, characterized by which specific emergent gauge group the marginalization produces. Standard-Model-reproducing classes share $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$; other classes produce different gauge groups. Below this is the *algebra-channel sub-class*, characterized by the specific algebra of observables and specific channel dynamics — the level at which two specific substratum-with-observer systems become indistinguishable in every observable.

Level C commits to a structural realism at the class level: the universality class is what is real, with specific substratum representatives playing the role of gauge expressions. Different members of the same universality class are observationally indistinguishable by definition, so the question "which specific substratum is real?" has no empirical content at the universality-class level. The realism is precisely about what survives the universality-class quotient.

Level D: OI's specific local structural residue. The framework's narrowest reconstructed commitment is the three-dimensional cubic local lattice/gauge sector with $K = 6$, the wave-equation dynamics and the listed Standard-Model representation data, with the qualifications developed in the technical chapters. The reconstruction theorem establishes that the stated observed and structural inputs fix this **local residue** under Theorem 23's conditions. It does not uniquely determine a Bell-inclusive full representative. With one-step own-wing setting interventions, readouts outside the other wing's graph cone and a ± 1 outcome in every run, the reference cubic graph is Bell-local under

measurement independence; the adopted branch proposes preparation-indexed state-dependent adjacency under the open hypothesis H-Bell (operational no-signaling and the metric/Ollivier–Ricci continuum limit), with its uniqueness left open.

Level D is where the framework's twenty-two classified quantitative predictions live (two parameter-free structural and five conditional structural among them): the Cabibbo angle $1/(\pi\sqrt{2})$, the Koide angle $2/9$, the gauge-coupling-running calculation giving the three SM coupling constants at the Z pole, the six fermion masses derived from one empirical input, the Higgs quartic coupling from cubic-group representation theory, and the JUNO neutrino-mixing prediction; outside that count sit the Bekenstein-Hawking $1/4$ factor as a consistency relation and the MOND acceleration scale $cH/6$. The empirical content that distinguishes OI from other observer-admitting frameworks is concentrated at Level D, because Level D is where the structural commitments become specific enough to produce specific numerical predictions.

The relationship between Levels D and C deserves a separate remark. The structural conditions A1–A6 that define Level D are not the same as the observer-admission conditions C1–C4 that define Level B. C1–C4 are conditions for *any* embedded observer to exist; A1–A6 are conditions for membership in OI's *specific* universality class within the observer-admitting subset. Some A's relate to but are distinct from C's: A1 (finiteness of the configuration space) is related to but distinct from C3 (sufficient hidden-sector capacity). A2 (determinism via bijection) selects deterministic-substratum members of a universality class, but observers can in principle exist on stochastic dynamics that satisfy the partial-trace observational features differently. A3–A6 (bounded coupling, center independence, linearity, background independence) are pure class-specific conditions, required for OI's specific gauge-structure derivation but not necessary for observer admission. The Level D commitment selects a specific universality class within the observer-admitting subset, with both the empirical inputs of physics and the framework's choice of structural assumptions playing a role in the selection.

The hierarchy nests. Each deeper level is a more specific commitment, with Level A $\supset$ Level B $\supset$ Level C $\supset$ Level D. Level A says only that observation occurs; Level B adds that the substratum's bijection structure must support observers; Level C adds the observational-equivalence structure that groups observer-admitting substrata into universality classes; Level D adds the framework's specific structural assumptions and selects the cubic-lattice substratum. The framework commits to all four levels simultaneously, with content at each level being more universal (broader levels) or more class-specific (deeper levels).

3.3 The observer-admission distinction in more detail

The observer-admission distinction at Level B deserves more sustained development than the previous section allowed. It is the framework's structural-

anthropic move, and it has consequences that distinguish the framework from both standard anthropic reasoning and the broader observer-dependence program in contemporary foundations of physics.

Standard anthropic reasoning operates on fundamental parameters. The cosmological constant, the masses of the fermions, the strengths of the four forces — these are observed to lie in narrow ranges compatible with the existence of observers, and the anthropic principle explains this by selection: a universe with different parameters would not contain observers to make the observation. The reasoning is conditional. It takes the existence of observers as given and explains the observed parameter values by the conditioning on observer existence.

The observer-admission reading at Level B sharpens this in two ways. First, it identifies which substratum features must be present for observers to exist at all, prior to questions about specific parameter values. The C1–C4 conditions are structural features of a substratum's dynamics: not values of free parameters but properties of the bijection's behavior. A substratum either admits observers or it does not; the question is binary at the structural level, before any parameter values are specified. Second, the observer-admission reading provides a more concrete account of what "selection" means. The standard anthropic principle is sometimes criticized as tautological — observers observe universes with observers in them — but this criticism misses the structural content. Observer-lacking substrata are not failed candidates that happened not to produce observers; they are structurally different mathematical objects, lacking the dynamical features that would make embedded observation possible.

A concrete example clarifies the structural distinction. Consider two finite cellular automata with the same configuration space size but different transition rules. The first has a Glauber dynamics on an Ising lattice at high temperature: fast mixing, short correlation times, no slow modes. The second has a topological phase-protected dynamics: slow modes from non-local order parameters, long correlation times in those modes, sufficient capacity in the topological sector. Both are well-defined deterministic systems on the same configuration space. In the coarse-grained description generated by its iterated one-step channel, with τ_S the observer's visible step, the first fails C2 in its timescale form, $\tau_S \ll \tau_B$, since it mixes fast relative to that step: writes into its hidden sector are scrambled before readback up to a suppressed residue, the absence of slow modes closing the thermal route to persistence; the second satisfies C1, C2, C3, C4 with appropriate partition choice. With C2 read in that timescale form, the first is observer-lacking; the second is observer-admitting. The distinction is structural — a property of the dynamics, not of the parameter values within a given dynamics — and applies prior to any specific universality-class commitment.

This structural distinction has implications for several conceptual problems. The Boltzmann brain problem — the worry that a sufficiently large statistical-mechanical system at equilibrium would produce observers as random fluctuations more often than through evolutionary processes — is dissolved at Level

B for C2 read in its timescale form. Under the effective-mixing hypothesis, for sufficiently fast-mixing equilibria ($\tau_{\text{mix}} \ll \tau_S$), and in the coarse-grained description generated by iterating the one-step effective channel ([Main §4.6]), equilibrium states are not observer-admitting in the sense of C1–C4 with C2 read in its timescale form, because that form, $\tau_S \ll \tau_B$, fails at equilibrium — its slow-bath route gone, no slow modes under that condition — and a bath that thermalizes between steps erodes the stored record before readback without establishing that none persists, suppressing the readback gap C4 requires exponentially in τ_S/τ_B at every order $k \geq 2$ ([Main §2.3]). Boltzmann brains are not generic outcomes of long-time evolution that just happen to be rare; they are structurally excluded from the observer-admitting subset by C2 in that timescale form. The framework's treatment of the arrow of time, developed in Chapter 2 §2.6, starts from this exclusion: observers are restricted to the non-equilibrium phase of the substratum's dynamics. The exclusion is time-symmetric; on an expanding homogeneous general-relativistic background satisfying the null energy condition, the apparent-horizon area supplies a non-decreasing clock, and whether the observer's other arrows align with it is not established.

The string-theory landscape problem looks different under the Level B reading as well. The conventional landscape picture posits roughly 10^{500} metastable vacua, each a different low-energy effective theory, with the vacuum-selection problem being to explain why our universe is in this particular vacuum. The Level B reading reframes this: most compactifications likely produce observer-lacking substrata at the relevant scales, because the slow-bath, sufficient-capacity, non-trivial-coupling conditions are non-generic. The SM-reproducing subset of the landscape is structurally smaller than the total landscape, because it is restricted to the observer-admitting subset. Whether this reframing fully resolves the landscape problem is an open technical question — it depends on whether observer-lacking compactifications can be characterized in string-theoretic terms, which requires substantive work in string-theoretic landscape physics — but the reframing is non-trivial.

The framework's commitment at Level B is therefore structurally substantive, not merely a definitional convention. It identifies the observer-admission boundary as a real feature of the space of candidate substrata, with consequences for anthropic reasoning, Boltzmann brains, the arrow of time, and the landscape problem. The realism at Level B is realism about the boundary itself: which substrata admit observers, which do not, and why. The empirical content at Level B is sparse — most of the framework's predictions live at deeper levels — but the structural content is rich.

3.4 The gauge axis

The framework's gauge structure runs along an axis orthogonal to the observation axis. Where the observation axis is about scope — which substrata

fall within the framework's domain — the gauge axis is about transformations: which substratum-level operations leave all observables unchanged. Like the observation axis, the gauge axis has four levels, each with its own equivalence notion.

Level G1: D-gauge on the emergent Hamiltonian. The first level of gauge freedom is the diagonal-unitary equivalence on the emergent Hamiltonian: $H \to DHD^\dagger$ with D a diagonal unitary in the measurement-outcome basis. This freedom preserves transition probabilities exactly. Energy shifts and the antiunitary reflection also preserve them; completeness of these alternatives under the stated gap and overlap hypotheses is conditional on the sign-aware integer classification H-BG ([GR §3.3]; Appendix B.7). Two Hamiltonians H and $DHD^\dagger$ produce identical Born-rule probabilities $T_{ij}(t) = |\langle j|e^{-iHt/\hbar}|i\rangle|^2$ in the measurement basis, so they are operationally indistinguishable. The transformation rephases the measurement basis, changing off-diagonal phases of H while leaving its diagonal entries and every transition probability unchanged.

Level G1 operates within a single Hamiltonian realization in a single emergent quantum mechanics — within Level D × Level G1 in the joint hierarchy, the framework's most specific intersection. Its content is the bookkeeping freedom in choosing phase conventions for the emergent basis states, with no implications beyond that.

Level G2: the Standard Model gauge group. The next level of gauge freedom is the Standard Model gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ acting on the emergent quantum fields. Chapter 2 §2.4 developed this as the unitary group of the bicommutant of the cubic-group action (the strict pointwise commutant is $\mathrm{U}(1)^3$ by Schur's lemma) on the six internal components of OI's substratum: the cubic-group decomposition gives multiplicities $(3, 2, 1)$, and the unitary group of the block algebra generated by the cubic-group representation on this decomposition is precisely $\mathrm{U}(3) \times \mathrm{U}(2) \times \mathrm{U}(1)$, reducing to $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ with a specified charged matter action (Chapter 5 §5.6); amplitude-scale gauge removes only the common phase. Background independence promotes this global commutant symmetry to local gauge invariance, giving the full SM gauge structure of standard quantum field theory.

An inclusion of Level G1 in Level G2 is not established. G2 includes the off-diagonal gauge transformations that mix different internal components, which G1 does not access.

A subtlety in Level G2 deserves explicit treatment. The *uniqueness* of $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ — that this specific gauge group is forced rather than chosen from alternatives — depends on the structural inputs A3–A6 and the nearest-neighbor restriction plus the empirical input E4 (spatial isotropy), and on H-link with its single-copy clause, the open hypothesis that the physical carrier is one copy of the six-link representation (Chapter 5 §5.4), together with an octahedrally equivariant condensate (Chapter 5 §5.5). Other universality classes that produce the Standard Model at low energies, including any successful string-

theoretic compactification, would reach the same gauge group through different machinery. Level G2 is therefore not specific to OI's substratum; it is shared with any framework that produces the Standard Model. What is specific to OI is *how* the gauge group is derived (cubic-group commutant) rather than the gauge group itself.

Level G3: substratum observational equivalences. Theorem 24 of [Substratum] exhibits exact invariances for specified visible interfaces. A single hidden permutation used identically in every visible fibre preserves the coherent channel family with the uniform hidden prior; an independently prepared, normalized, exactly decoupled finite ancillary factor leaves that family unchanged. Fibre-dependent hidden permutations preserve the rooted population transition probabilities, but need not preserve coherences. Alphabet change preserves observables only under the alphabet-size hypothesis; [SM §2.7] separates results proved for every $q \geq 2$ from those that carry q. These constructions do not classify all substrata or all observer partitions; completeness remains open.

Inclusions of G1 and G2 in G3, and a map from $\mathcal{G}_{\text{sub}}$ onto the Standard Model gauge group, are not established. The framework's structural account of why the Standard Model has its specific gauge group is the cubic-group commutant of Level G2, which follows from the substratum construction (Stage 2).

Level G3 differs from G1 and G2 in an important structural respect: it is an *equivalence-class symmetry*, not a Lagrangian symmetry. There is no Lagrangian at the substratum level; the substratum is a finite set with a bijection on it, governed by a discrete update rule rather than a continuous action principle. The gauge group acts on the equivalence class of substrata defined by the observability condition. It plays the same structural role at the substratum level that Lagrangian gauge symmetries play at the field-theory level — identifying observationally indistinguishable configurations — but through a different mathematical mechanism.

Level G4: universality-class equivalence. The broadest gauge freedom in the hierarchy operates at the universality-class level. Two substratum-with-observer systems are universality-class equivalent if their algebra-channel pairs are isomorphic in the precise sense identified at Level C — preserving both the algebraic structure of visible observables and the channel's dynamical content. Level G4 captures equivalences that may not be expressible as transformations within any single structural class.

Level G4 is broader than generators (i) and (iii) of $\mathcal{G}_{\text{sub}}$. Substrata related by them are universality-class equivalent (since they produce the same algebra-channel pair), but substrata not related by any $\mathcal{G}_{\text{sub}}$ transformation may still be universality-class equivalent if they produce the same algebra-channel pair through structurally different machinery. The classic candidate is the relationship between OI's cubic-lattice substratum and string-theoretic compactifications that reproduce the Standard Model. The two are not related by any $\mathcal{G}_{\text{sub}}$ generator — OI's bounded-coupling, finite-deterministic structure is not

preserved by string-theoretic compactifications, which violate A2 (determinism) and A5 (linearity) at the matrix-model bridge — but they may produce equivalent algebra-channel pairs, making them universality-class equivalent at Level G4 even though they are not gauge-equivalent at Level G3.

Level G4 operates only within the observer-admitting subset — Level B. Observer-lacking substrata are outside the universality-class structure entirely; the algebra-channel pair is undefined for them because there is no embedded observer to define the algebra of visible observables or the channel induced by the partial trace.

The levels of the gauge hierarchy. The levels carry distinct equivalence notions; inclusions among G1, G2 and G3 and a map from $\mathcal{G}_{\text{sub}}$ onto the Standard Model gauge group are not established. The framework commits to all four levels of gauge simultaneously, with different levels of gauge being relevant to different questions about the substratum and its emergent physics.

3.5 Orthogonality, multi-level realism, and stratified predictions

The two hierarchies — observation axis and gauge axis — are orthogonal. The observation axis is about scope: which substrata fall within the framework's domain at each level of specificity. The gauge axis is about transformations: which operations leave observables unchanged at each level of breadth. A specific empirical claim or theoretical commitment lives at a specific *intersection* of the two axes, with the intersection determining both what the claim's content is and how it can be falsified.

The orthogonality has substantive content. Levels G1, G2, and G3 sit within Level D — they are gauge structures of OI's specific universality class with its specific gauge derivation. Level G4 operates at Level C — it characterizes universality-class equivalence, which is broader than OI's structural class. The two-axis structure means the framework's strongest realism claim — the Chapter 2 commitment to $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$ — operates at the Level D $\times$ Level G3 intersection. Weaker realism claims operate at broader intersections, with corresponding shifts in empirical content.

What is real at each intersection. The framework commits to realism at four distinct levels of the two-axis hierarchy, with the realism content differing at each level.

At Level G4 $\times$ Level C — the broadest intersection — what is real is the algebra-channel structure of partial-trace observation: the Born rule, channel-level unitarity to leading order, memory/divisibility — accessible memory where the hidden sector reads back its record (C4), and, on any finite reversible substratum, recurrence-cycle P-indivisibility of the marginal stochastic process wherever a rooted visible map is not a permutation — and the commutant gauge-invariance pattern. This level's realism is universal across observer-admitting

substrata. Every embedded observer in any universality class produces these features; they are structural features of partial-trace observation that no embedded observer can avoid. The realism is about partial-trace observation itself, not about any specific substratum or any specific universality class.

At Level G3 × Level D — the framework's distinctive intersection — what is real is the **reconstructed local lattice/gauge residue modulo** $\mathcal{G}_{\text{sub}}$. Chapter 2 shows that the invariants of this local residue are uniquely fixed under Theorem 23's stated conditions. Bell-inclusive completion data are not included in that uniqueness claim: H-Bell conditions their compatibility/existence and no result selects a unique preparation-indexed Bell rule. The realism is therefore strongest for the reconstructed local structural residue, not for one uniquely specified Bell-complete representative.

At Level G2 × Level D — a narrower intersection in the gauge axis but the same observation level — what is real is the emergent quantum field theory structure: the Standard Model gauge group, the matter content of three generations (physical reading conditional on H-spin', SM §4.7; chirality on H-χ', SM §4.8) plus one Higgs doublet, the anomaly-free hypercharge assignment on the nonzero-Y_Q branch. This level's realism is forced uniquely within OI by the cubic-group commutant structure, conditionally on H-link and an octahedrally equivariant condensate (Chapter 5 §5.4–§5.5), but is shared with any successful SM-reproducing universality class. The realism is about the emergent quantum field theory rather than about any specific substratum.

At Level G1 × Level D — the narrowest intersection — what is real is the emergent Hamiltonian's transition-probability content modulo the diagonal-unitary D-gauge. This is the level at which specific numerical predictions for emergent dynamics become well-defined: the level at which the Schrödinger equation operates on specific Hilbert-space vectors with specific phase conventions.

These four realisms are not competing claims about what is fundamentally real. They are realism at different levels of the two-axis hierarchy, with content that is more class-universal at broader levels and more class-specific at deeper levels. The framework commits to all four simultaneously, with the commitments coordinated by the hierarchy's nesting structure.

Predictions stratified by hierarchy intersection. The framework's empirical predictions live at specific intersections of the two axes, with the intersection determining both what the prediction's content is and how it can be tested or falsified.

Level G4 × Level C predictions — the Born rule, channel-level unitarity, memory/divisibility (accessible memory where the hidden sector reads back its record (C4), and, on any finite reversible substratum, recurrence-cycle P-indivisibility of the marginal process wherever a rooted visible map is not a permutation), the commutant restriction pattern — are essentially structural. They are universal across observer-admitting systems, trivially predicted because they are features of partial-trace observation rather than of any specific substratum. They cannot

be falsified without contradicting embedded-observer physics itself; falsifying them would refute the entire program of treating observation as a partial-trace operation. The framework predicts them; so does any framework that takes embedded observation seriously.

Level G2 × Level D predictions — the Standard Model gauge group, the three-generation structure, the anomaly-free hypercharges on the nonzero-Y_Q branch, the composite Higgs — are forced uniquely within OI by the cubic-group commutant structure, conditionally on H-link and an octahedrally equivariant condensate (Chapter 5 §5.4–§5.5), but are shared with any framework that reproduces the Standard Model from observer-admitting substrata. Falsifying them would refute the entire class of SM-reproducing universality classes, not just OI specifically. They are central to the framework's content but do not distinguish OI from successful alternative frameworks.

Level G3 × Level D predictions are where OI's distinctive empirical content lives. The Cabibbo angle as $1/(\pi\sqrt{2})$ from a Brillouin-zone distance; the Koide angle as $2/9$ from a cubic-group Casimir; the MOND acceleration scale as $cH/6$ from entropy displacement; the JUNO neutrino-mixing angle as $1/3 - 1/(4\pi^2)$ — all of these are class-specific to OI's substratum representative. Falsifying them would refute OI's specific class membership without necessarily refuting the broader universality-class framework. A different universality class might produce the Standard Model with different specific numerical values for these observables.

The framework's most distinguishing empirical content lives at Level G3 × Level D — these are the predictions that distinguish OI from other observer-admitting frameworks, including ones that successfully reproduce the Standard Model. The JUNO match is a Level G3 × Level D result — retrodictive: it agrees with OI's specific cubic-group calculation, and disagreement would have refuted OI's representative without refuting the broader observer-admission program. The Bekenstein-Hawking 1/4 factor sits at Level G2/G3 × Level D — a conditional derivation (horizon conditions, Chapter 7) with no direct empirical test (GW250114 constrains the classical area theorem, not the coefficient) — since both OI's substratum and any successful SM-reproducing framework that yields the area law would predict the same factor.

Falsification is stratified accordingly. Falsifying a Level G4 × Level C prediction is impossible without contradicting the structural features of embedded observation. Falsifying a Level G2 × Level D prediction would refute the entire SM-reproducing universality class. Falsifying a Level G3 × Level D prediction would refute OI's specific representative while leaving the broader universality-class framework intact. The framework's twenty-two quantitative predictions live primarily at Level G3 × Level D, which makes them the most informative empirical content the framework offers — but also the most narrowly falsifiable. A pre-registered falsification of a Level G3 × Level D prediction is a falsification of OI specifically, not of the broader observer-admission framework or of the universality-class structure.

The stratification has methodological consequences developed in Chapter 4. Different kinds of empirical content require different epistemic standards, and the hierarchy makes the stratification precise. Level G4 × Level C content is structural; Level G2 × Level D content is shared across an entire universality class; Level G3 × Level D content is class-specific to OI. The framework's preregistration of falsification conditions, developed in Chapter 4, identifies which empirical results would falsify which level of the framework's commitments.

3.6 Gauge freedom and observational incompleteness

The two axes developed so far invite a question the chapter has not yet answered directly. The gauge axis catalogues what is *not* observable — degrees of freedom that can be transformed without changing anything an embedded observer could record. The framework as a whole is named for *observational incompleteness* — the embedded observer's inability to resolve the substratum it is part of. These two notions have been running alongside each other for three chapters. It is worth asking whether they are the same thing.

Gauge freedom supplies a form of observational incompleteness relative to specified data: different descriptions can induce the same observed channel. This does not identify every inaccessible fact with a representational redundancy.

The apparent opposition. At first glance gauge freedom and observational incompleteness look like opposites. Gauge freedom is *surplus*: the description carries more structure than the physics, and the gauge quotient throws the excess away, losing nothing. Observational incompleteness sounds like a *deficit*: the observer has *less* than the whole story, missing what the trace-out over the hidden sector destroys. Surplus and deficit do not obviously belong together — one is too much representation, the other too little access.

The connection becomes precise once the observations are specified. A uniform hidden-coordinate relabelling preserves the coherent channels at a fixed visible interface with a uniform hidden prior. Its coordinate labels are surplus relative to those observations: they distinguish descriptions without distinguishing the observed channel. This is one sense in which gauge freedom is observational incompleteness. It does not identify every unobserved property with a redundancy, or establish that a different observer interface would leave the same properties inaccessible.

Unresolved facts within a model. A fixed finite model can contain determinate facts that a particular record does not identify. Let F_r be the nonempty set of microscopic states compatible with an observed record r. A quantity q is determined by that record exactly when it is constant on F_r [Substratum Proposition 24.5]. This applies to a future visible value as well as to a hidden-state quantity. If two compatible states give different values, no decision using only that record determines which value obtains. If the value is constant, the

observer may know it without reconstructing all of S. Proper embedding alone does not imply inability to predict the next visible state.

These comparisons distinguish different questions:

Gauge incompleteness is redundancy relative to specified observations. A uniform hidden-coordinate relabelling, or an independently prepared, normalized, exactly decoupled finite ancillary factor, can leave a fixed observed channel unchanged. This does not make every inaccessible state or every deeper physical implementation gauge. The preparations, readouts and dynamics included in an observational comparison determine what its equivalence relation identifies.

Computational limitations concern the resources needed to evaluate a quantity from an available description. A lower bound on those resources requires its own model and argument. The record-fibre criterion above is an information limit: it distinguishes an unresolved fact from a redundancy between descriptions, without asserting a universal computational no-go for finite embedded observers.

The distinction separates unresolved facts within a model from redundancies between descriptions. Hidden states may have different future effects even when a present record does not distinguish them; their physical relevance depends on the permitted observations and dynamics. A proved representational redundancy is a different matter. Neither kind of incompleteness supplies a complete classification of the other.

Where the comparison sits in the hierarchy. Representative freedoms identify descriptions relative to specified data. A separately established computational resource bound is a different restriction. Neither the ancillary identity nor the observer's being a proper part of S supplies such a bound on its own. The hierarchy organizes these questions, while a full classification of observational or computational limitations remains open.

A consequence worth stating. The methodology of physics modulo gauge asks for the structure fixed by specified observations, rather than a preferred representative among descriptions that give the same data. The exact invariances above illustrate that programme. For the full substratum, however, the observable interfaces and their correspondence must be specified, and a complete inverse classification remains open. The programme does not establish that every question beyond a chosen interface lacks physical content.

The comparison is an interpretive proposal about the fixed-interface examples above. It does not provide an additional classification theorem or an assumption used by the empirical chapters.

3.7 No-go theorems at different descriptive levels

The framework is a hidden-variable theory in the literal sense: there is a deterministic substratum with definite states, and the observer's stochastic descrip-

tion is obtained by marginalizing over the parts of that substratum they cannot see. The twentieth-century no-go literature — Bell's theorem (1964), Kochen-Specker (1967), the Pusey-Barrett-Rudolph theorem (2012), the Frauchiger-Renner theorem (2018) — rules out hidden-variable accounts satisfying specific structural conditions. The framework must therefore state how each of these results applies to it, and what its response costs.

The framework's engagement with these theorems is not a single trick repeated, nor four separate tricks. Each theorem identifies a different structural property of would-be hidden-variable accounts. The two-level substratum-emergent structure joined by the trace-out is the common bookkeeping architecture, but it is **not** one automatic escape: Bell forces ontic parameter dependence in the adopted deterministic branch; Kochen-Specker constrains the noncommuting operational algebra; PBR constrains preparation independence of ontic composites; Frauchiger-Renner constrains universal nested-agent descriptions. The hierarchy matters because it locates each premise at the right level, not because the same property simply holds above and fails below in every case.

Bell's theorem. Bell's theorem binds directly on the deterministic hidden-variable completion. If the setting interventions, applied at one step, each read and write only their own wing, the Bell-relevant coupling range does not connect the wings before readout, every run records a ± 1 outcome, and one setting-independent response-complete ensemble is used, the model factorizes and obeys $|S_{\mathrm{CHSH}}| \leq 2$. The framework therefore cannot use graph locality, screening, outcome dependence, or indivisibility to explain the quantum violation. It retains measurement independence and accepts **ontic parameter dependence** for Bell-violating substrata, with operational no-signaling as the distinct empirical locality statement. The cited stochastic causal-local indivisible construction can still obey Tsirelson because its causal-independence criterion is weaker than Bell factorization. Unrestricted globally controlled OI realizations remain the PR-box countercontrol.

In hierarchy language, the Bell question splits the deterministic substratum from the reduced stochastic description. A Bell graph that does not connect the wings before readout, with one-step own-wing setting interventions and a ± 1 outcome in every run, plus a setting-independent response-complete ensemble, would force the substratum into Bell's local class and cap CHSH at 2. The framework therefore does not identify no-signaling, Jarrett parameter independence, Fine's theorem, hidden memory, or indivisibility with a derivation of the quantum correlation set. Its adopted deterministic branch is instead ontic parameter dependence with operational no-signaling; the $2\sqrt{2}$ result for the causal-local indivisible stochastic construction remains a distinct weaker-description theorem.

Kochen-Specker. The Kochen-Specker theorem rules out non-contextual hidden-variable assignments: there is no function v mapping observables to definite values such that $v(O)$ depends only on O rather than on the measurement context (the set of compatible observables being measured simultaneously). Any hidden-variable theory matching quantum mechanics must be contextual:

the hidden variable's outcome on a given observable depends on which other observables are co-measured.

The framework's response is structural, and it is worth stating precisely, because the Kochen-Specker case does not fit the same "property-at-one-level" template as Bell. A noncontextual valuation is an assignment of definite values to the emergent *observables* that is independent of measurement context. The framework reproduces the quantum statistics at the fixed-basis transition layer (the characterization theorem of Chapter 1), while Kochen-Specker is a theorem about the operational theory's noncommuting observable algebra — the layer of the operational bridge, characterized conditionally and with its physical sourcing open ([Main §3.2, §3.4]): the observable algebra of quantum theory in dimension three or higher admits no noncontextual valuation. Any account reproducing quantum mechanics therefore reproduces a structure in which no noncontextual valuation of the observables exists. So the property Kochen-Specker forbids does not hold at the emergent level — and it does not hold at the substratum level either, because the substratum carries a definite configuration, not a valuation of the emergent observables at all. The framework's contextuality is therefore not "noncontextual below, contextual above"; conditional on the coherent operational lift reproducing the standard observable algebra, it inherits the Kochen–Specker obstruction, and what the framework supplies is the *account* of it: an emergent observable's value is produced by the trace-out, the trace-out is taken with respect to a partition, and measuring a set of compatible observables is a specific apparatus coupling — a specific partition. Which observables are jointly defined, and the value each takes, are features of the reduced description fixed by that partition, not partition-independent functions of the substratum configuration. Contextuality, in the framework, is the partition-dependence of emergent valuation.

In hierarchy language: Kochen-Specker non-contextuality concerns the emergent observable structure at Level G2 $\times$ Level D. That structure is contextual — different partitions produce different marginalized statistics — because — conditionally on the operational bridge reconstructing the standard observable algebra — the framework carries the operational theory, and that theory has no noncontextual valuation. The Level D substratum carries definite states, but a definite substratum configuration is not a valuation of the emergent observables, so it neither supplies nor contradicts a noncontextual valuation. The framework does not exhibit "non-contextuality below, contextuality above"; it inherits the Kochen-Specker obstruction with quantum mechanics and locates its source in the partition-dependence of the trace-out. Here the framework's response is not a property placed at one level and denied at another, but a property unavailable at every level — a difference §3.7's closing paragraph returns to.

Pusey-Barrett-Rudolph (PBR). The PBR theorem rules out "epistemic" interpretations of the wave function: under reasonable assumptions, the wave function cannot represent the observer's incomplete knowledge of an underlying ontic state, because pure states with non-zero overlap would be indistinguishable

in any consistent ontic model. The wave function must, in some sense, be ontologically real.

The framework's PBR response must be stated conditionally. PBR assumes Preparation Independence: independently prepared systems have a product joint distribution over their ontic states. Bare OI does not posit that assumption. Two preparation devices embedded in one substratum may share preparation-relevant hidden or boundary data even if their later dynamical causal cones are disjoint, so the product premise is not guaranteed by graph locality. But the converse overclaim is also avoided: the coupling graph alone does not prove a specific conditional product law, nor does it derive the standard quantum tensor-product composite. Those are composite operational statements: the operational-completion characterization ([GR §3.3]) takes the tensor-product composite as part of the complex matrix kinematics it assumes, and deriving that composite is open. Therefore PBR constrains any OI completion that satisfies its preparation-independence premise; it does not automatically constrain every bare OI realization, and the framework's final classification relative to PBR follows only after the composite operational lift is fixed.

EPR, read correctly. The EPR argument (Einstein, Podolsky, and Rosen 1935) is routinely misremembered as an appeal to determinism. It is the reverse: from locality and the perfect correlations, determinism is *derived*. If nothing at one wing can affect the other, then the ability to predict the distant outcome with certainty entails that the outcome was already fixed — the hidden determiner is the conclusion of the argument, not its premise. The framework accepts this reading without reservation, and accepts what follows from accepting it: the framework is precisely the kind of theory EPR's argument called for, a deterministic completion in which measurement reveals antecedently definite substratum facts. That is why Bell's theorem binds on the framework with full force. A completion of the EPR type must, by Bell's mathematics, violate factorizability to reproduce the observed correlations — there is no third way, and the framework does not claim one.

What the framework does not argue. Two standard escapes are explicitly declined. It does not deny counterfactual definiteness: substratum outcomes are definite, and the framework nowhere trades on refusing to consider unperformed measurements — the response to Bell runs through the structure of the premises, not through denying that measurements have results. Nor does it treat hidden setting correlations as an explanation of Bell statistics. Measurement independence is an explicit premise of the adopted Bell branch; chaotic mixing may motivate laboratory decorrelation between pair-preparation and setting-generating degrees of freedom, but the violation itself is assigned to ontic parameter dependence. The setting-dependent horn is quantified and declined (Main §3.3).

Bell's theorem, conceded and then confronted. Bell's mathematics binds on the framework because the substratum is a response-complete hidden-variable completion. If the setting interventions, applied at one step,

each read and write only their own wing, the Bell-relevant graph does not connect the wings before readout, every run records a ± 1 outcome, and the pre-setting ensemble is independent of the settings, determinism gives Bell factorization and $|S_{\mathrm{CHSH}}| \leq 2$. The framework therefore cannot explain quantum Bell violation by combining pointwise graph locality and indivisibility. It declines the measurement-dependent/superdeterministic horn and takes **ontic parameter dependence**. In the specific SM/GR realization, the target is preparation-indexed adjacency in the already state-dependent graph $G(x)$ (or an equivalent ontically nonlocal response): the reference vacuum/uniform graph remains geometric and bounded-range, while an entangling preparation may add Bell-relevant long adjacency. H-Bell is the open requirement that those prepared graphs preserve operational no-signaling and the metric/Ollivier–Ricci continuum limit. The curvature condition is downstream of a harder one. The cited convergence theorem is arbitrary-dimensional only for the manifold-distance-weighted graph metric, and for the degree-6 cubic reference graph the intrinsic hop metric is exactly ℓ_1 — a $\sqrt{2}$ diagonal stretch that does not decay at any scale — so the hop-metric curvature route fails for the reference family, before any Bell edge is added, and no $D = 3$ extension of that theorem would rescue it. The emergent geometry itself is not in question: the substratum's dispersion relation is isotropic at leading order. What must be re-founded is the curvature functional, on the propagation/Laplacian geometry rather than on shortest-path counts [SM §3.2]. The causal-local indivisible *stochastic* construction can still obey Tsirelson because its causal-independence condition is weaker than Bell factorization; that result is not the locality structure of the deterministic completion. The unrestricted PR realization remains the negative control showing that representability, no-signaling, and indivisibility do not select the quantum set.

The rejoinder, at full strength. A response-complete deterministic Bell completion is not local in Bell's parameter-independence sense. A response-complete hidden-variable theory reproducing Bell violation while retaining measurement independence must fail parameter independence, and the framework explicitly accepts that conclusion. Its Bell-violating prepared states are therefore **ontically nonlocal in Bell's parameter sense**. In the concrete lattice branch, that nonlocality is targeted through preparation-indexed state-dependent adjacency (or an equivalent response rule) under H-Bell, not by pretending the nearest-neighbor reference wave equation itself carries Bell influence. What remains local operationally is the no-signaling statement — changing a controllable setting cannot be used to transmit a message to the remote wing. The cosmological partition or shared boundary may be proposed as a physical realization, but calling that carrier atemporal, global, or prior to emergent spacetime does not remove Bell nonlocality; it only locates it in the ontology. The separate Barandes–Hasan–Kagan theorem shows that the reduced stochastic law can satisfy their weaker causal-independence criterion and remain below Tsirelson, but that criterion is not Bell factorization for the deterministic completion. The manuscript can therefore defend a non-signaling deterministic ontology with a

specific locus of nonlocality; it does not claim a local hidden-variable resolution of Bell.

The bill, paid in the open. A deterministic substratum that violates **ontic parameter independence** owes an account of relativity, and the account cannot be evasive: the framework's substratum has a preferred frame — the frame in which update steps are simultaneous — and on identification with the observed universe this is operationally the cosmic rest frame, an identification the framework records rather than derives (SM §3.1). Lorentz invariance is therefore wholly emergent, and the framework's obligations are quantitative: no linear violation (established at tree level), rotational protection by octahedral symmetry to all orders conditional on the emergent theory preserving that symmetry at loop level and on the condensate being aligned to the cubic axes (H-link with an octahedrally equivariant condensate; Chapter 19 §19.3.7), boost-sector violation controlled, conditional on the symmetric-point matching under H-blind (SM §4.4), by a decoupling law in the cross-charged mass M_X whose quadratic-with-logarithm form was reported by a one-loop calculation that is not independently verified, its two-sided window illustrative until it is. These are not side technicalities; they are the price of the preferred-frame answer to the foliation objection, and the open item among them — the interacting (condensate-dressed) extension of the coarse-graining characterization (SM §3.1; Chapter 19 §19.3.7) — is correctly understood as Bell-defense infrastructure. A reader who wants to falsify the framework's account of Bell should attack there: a demonstrated failure of emergent Lorentz protection would leave the preferred frame exposed at accessible energies, and the framework's answer to this subsection would fail with it.

The characteristic trait. Schrödinger called entanglement *the* characteristic trait of quantum mechanics, the one forcing its entire departure from classical lines. The framework agrees, with a precise gloss: entanglement is, in the framework's mechanism proposal, the emergent description's encoding of the correlations that the shared preparation and the indivisible joint process carry between subsystems. Where those correlations violate a Bell inequality, they are not a record written at preparation and read out locally: given deterministic dynamics and a ± 1 outcome in every run, a completion that keeps ontic parameter independence and measurement independence obeys $|S_{\mathrm{CHSH}}| \leq 2$ (Main §3.3). The Bell-violating completion is therefore ontically parameter-dependent, since it retains measurement independence, and whether the SM/GR substratum supplies that dependence through preparation-indexed adjacency while meeting the required no-signaling and continuum-curvature conditions remains the open hypothesis H-Bell. The standard quantum operational realization of these correlations is characterized, within complex matrix kinematics taken as given, by the five completion conditions of [GR §3.3], whose physical sourcing is open. The trait is characteristic because the partition is characteristic: it is what an embedded observer's account of a shared hidden sector must look like.

The ledger. Conceded: Bell's mathematics; EPR's logic and its corollary

that Bell binds; the failure of the counterfactual-definiteness escape; **ontic parameter dependence** of the Bell-violating completion; the preferred frame; the configuration-space objection. The remaining interpretive question is not whether the substratum is Bell-local — it is not — but whether locating the parameter dependence in a cosmological/shared-boundary structure yields a coherent non-signaling emergent relativistic theory. The loop-level protection of emergent Lorentz invariance remains the quantitative hostage. The EPR reading that this subsection rests on can be put in one sentence: Einstein was only half right — God doesn't play dice, but he does play cards. The cards are not a loose figure of speech: Chapter 1 §1.11 builds the game — a definite deal held hidden, a deterministic shuffle that writes visible history into the burn pile and reads it back, revelation as an intervention with a cost — and states its boundary as plainly as its successes: a card game in which, once the deal is down, every operation, including each shuffle the rules prescribe, each move and each setting choice, reads and writes one player's own cards only, and which operation is applied to a player's cards, and how that player's outcome is read from them, depends only on those cards, that player's own setting and on chance that, taken together with the deal, is independent of the settings, with each outcome a ± 1 result in every run, no run set aside, and the settings chosen independently of the deal, violates no Bell inequality. That boundary is not an embarrassment to the metaphor here; it is the diagnosis. Such a card game is local through and through — each player's hidden cards are their own, and the deal, taken together with any chance the game uses, does not know the settings — and single-system interference is exactly as far as such a game can go, which is to say exactly as far as a local hidden-variable model can go. What the world adds, and no such table can, is the wager of this subsection: the Bell-relevant completion is not parameter-local. The two wings couple through one shared horizon/boundary structure, and that is where the remote-setting dependence is proposed to live — in the full construction, not in the toy. The shared bath is therefore a proposed **locus of Bell nonlocality**, not a local common-cause escape from Bell.

3.8 The philosophical lineage

The framework's structural-realism position has antecedents older than quantum mechanics and broader than the contemporary observer-dependence program. Articulating the relationship between the framework and its philosophical lineage is part of locating the framework's distinctive contribution.

Nicholas of Cusa. The fifteenth-century cardinal and philosopher developed two doctrines that anticipate the framework's structural claim. *Docta ignorantia* — learned ignorance — held that the highest knowledge of God is knowing what we cannot know: the human mind, bounded and finite, can approach the infinite only through the recognition of its own limits. The structural form of this claim — that incompleteness is a feature rather than a deficiency — anticipates the framework's reframing of the cosmological constant problem: the

10^{122} discrepancy is not evidence that physics is incomplete in a way needing repair but the structural signature of how observation operates in a universe an observer is embedded in.

Coincidentia oppositorum — coincidence of opposites — held that contradictions resolve at the level of the infinite: properties that appear incompatible to finite intellects are unified in the divine, where the polygon and the circle are the same thing, where the maximum and the minimum coincide. The structural form of this claim — that two seemingly irreconcilable descriptions can both be correct when both are projections of a common underlying reality — anticipates the framework's claim that quantum mechanics and general relativity are not incompatible but are projections of the same finite deterministic substratum along different axes. The conventional formulation treats the two as competing theories requiring reconciliation; Cusa's framework would suggest that the contradiction is at the level of the projections rather than the substratum, and the framework formalizes this suggestion.

Cusa is not the framework's source. The framework derives its claims from explicit mathematical theorems, not from medieval theology. But the structural resemblance is striking, and identifying the resemblance helps locate what is distinctive about the framework's claims and what is part of an older tradition of thinking about observation, infinity, and structure.

Spinoza. Spinoza's metaphysics is the deepest historical parallel to the framework's substratum-and-projections picture. The *Ethics* posits one substance — God or Nature — expressed through infinitely many attributes, each constituting the substance's essence under a different aspect. The two attributes humans can know are extension (the physical aspect) and thought (the mental aspect), but these are not two substances; they are the same substance expressed through different attributes, with no causal interaction between them because there is only one thing underneath. The doctrine that the order and connection of ideas is the same as the order and connection of things follows from this single-substance structure: the parallelism between mind and body is not an interaction but a consequence of their being the same underlying reality.

The structural form of Spinoza's account — one substance, multiple incommensurable attributes, with the apparent multiplicity being a feature of how the substance is expressed rather than of the substance itself — maps directly onto the framework's three-projection structure. The substratum (S, φ) is the single substance; quantum mechanics and general relativity are projections expressing the substratum's content under different aspects, with a third bearing on the arrow of time; the appearance of three distinct emergent theories is a feature of the projections rather than of the substratum. The framework's commitment to ontic structural realism on the **reconstructed local residue modulo** $\mathcal{G}_{\text{sub}}$ is Spinozistic in form: what is real is the gauge-invariant local structure, not any specific representative; the Bell-inclusive completion is not uniquely reconstructed.

The parallel is not complete. Spinoza's substance is unique (there is exactly one substance, by argument); the framework's substratum is a member of an equivalence class (there are many specific bijections, gauge-equivalent under $\mathcal{G}_{\text{sub}}$). Spinoza's attributes are infinite in number; the framework identifies three projections (quantum mechanics, general relativity, arrow of time) without claiming to enumerate all possible projections. The framework's claims are mathematical theorems rather than metaphysical doctrines. But the structural resemblance is closer than to any other position in the historical philosophical literature.

Worrall and epistemic structural realism. The closest twentieth-century antecedent to the framework's position is John Worrall's epistemic structural realism, developed in the 1980s and 1990s. Worrall argued that scientific theories should be interpreted as committing to the structure of relations between observable phenomena rather than to the intrinsic natures of the underlying entities producing those phenomena. The argument was epistemic: structure is what survives theory change across the history of science (the equations of Fresnel's wave theory survived the transition from ether to electromagnetic field), so structure is what scientific theories warrant belief in. The framework's structural-realism position is ontic rather than epistemic — what is real is the equivalence class, not just what we are warranted in believing about it — but the structural commitment is in the same family. Both Worrall and the framework hold that structure rather than substance is what survives the appropriate quotient.

The framework's programme relative to Worrall is to specify and test observational equivalences for explicit models. Hidden-coordinate changes and exactly decoupled ancillary enlargements supply concrete fixed-interface examples; the completeness of a proposed substratum classification remains open. The universality-class structure compares specified algebra-channel pairs across structural classes, while the hierarchy organizes realism claims by their assumptions and level of generality. These comparisons provide mathematical questions about the relevant structure without already supplying a complete boundary between physical content and representation.

Hofstadter and strange loops. Douglas Hofstadter's *Gödel, Escher, Bach* (1979) and *I Am a Strange Loop* (2007) developed the idea that self-reference under finite resources produces emergent phenomena (consciousness, in Hofstadter's primary application) that are real features of the self-referential system without being features of the substrate. The framework's incompleteness-family argument — Gödel for formal systems, Turing for computational systems, the framework for embedded observers — agrees with Hofstadter's central claim that self-reference produces structural emergence.

The framework disagrees with Hofstadter on a specific point. Hofstadter held that the substrate is explanatorily inert: consciousness emerges from the strange loop regardless of what the loop is implemented on (neurons, silicon, abstract automata), and the substrate adds nothing beyond providing the implementation. The framework's substrate is not inert. The 10^{122} ratio between substratum ca-

pacity and visible-sector capacity is the cosmological constant discrepancy; the dark sector is the gravitational effect of the substrate's content that the observer cannot read; the framework's twenty-two quantitative predictions follow from specific structural features of the substratum that different substrates would produce differently. The framework agrees with Hofstadter that the strange loop is real and that self-reference produces structural emergence; it disagrees that the substrate is therefore irrelevant.

The embedded-observer limits tradition. The Introduction placed the framework's central claim in the impossibility-with-structure family — Gödel for formal systems, Turing for computational systems, Wolpert for physical inference devices — and stated the relationship in a sentence this section need not repeat: the logical results corroborate the framework's derivation but do not supply it. (Wolpert's theorems are engaged separately, as an objection, in Appendix C.7.) That family has further members, closer to the framework's specific concern, that belong in the lineage record. Dalla Chiara (1977) analyzed the quantum measurement problem as a self-reference problem in the logician's sense, decades before that framing was common. Rössler's endophysics (1998) distinguished systematically between the view from inside a world and the view from a hypothetical outside, arguing that physics as practiced is unavoidably interface physics. Breuer (1995) proved the sharpest single result: no observer can perform an accurate measurement of the complete state of any system in which that observer is contained — an impossibility theorem, not an interpretation. And Szangolies (2018) derived quantum-like epistemic restrictions from self-reference structure via Lawvere's fixed-point theorem, the nearest approach in this tradition to extracting quantum form rather than mere limitation. The framework cites all of these as load-bearing lineage, and its relationship to them is the one the Introduction stated for the family as a whole and §3.7 states for EPR: the tradition establishes that the limit exists; the framework derives what the limit forces — hidden incompleteness with, under the stated memory architecture, an accessibly memory-bearing marginal (P-indivisibility the stronger diagnostic in the appropriate regimes), the representation universal and carrying the compressed quantum form for that subclass under (T), through the ancilla-dilated representation (the construction's own marginals are doubly stochastic but not in general unistochastic; [Main §3.1]).

Contemporary observer-dependence programs. The framework converges with the recent technical work on observer-incorporated quantum gravity — Chandrasekaran-Longo-Penington-Witten on de Sitter entropy with observer (2022), Maldacena on observer's-clock dependence in cosmological partition functions (2024), Harlow-Usatyuk-Zhao on observer-dependent Hilbert spaces in closed universes (2025) — on the foundational claim that observation matters and that the algebra of observables depends on the observer's embedding. Slagle and Preskill (2023) add a different comparison: an explicit local classical lattice model whose boundary dynamics reproduces quantum mechanics, using stochastic dynamics in an extra spatial dimension. The technical apparatus these programs employ differs from the framework's, and the empirical content

they produce is different (mostly structural commitments rather than specific numerical predictions), but the foundational commitment is shared.

Hawking's turtles and the regress of support. Hawking opens *A Brief History of Time* (1988) with the anecdote of the listener who insists the world rests on a giant turtle and, pressed on what the turtle stands on, answers that it is turtles all the way down. The story lampoons the oldest structural problem in metaphysics: every proposed support invites the same question one level lower, and the classical exits — an infinite tower, a circle, or a foundation that simply stands — each pay a familiar price. The framework's own picture makes the regress feel urgent again in modern dress. If the universe is completely described by a lossless finite-state memory with finite read-write access, what is the hardware the memory runs on?

The exact result concerns a specified observation interface. If an enlarged model has product dynamics $\varphi \times \psi$ and its added finite factor is independently prepared in a normalized state, tracing out that factor leaves the original visible channel unchanged [Substratum Theorem 24(iii)]. The result holds at every discrete time because the added dynamics remains exactly decoupled. It says nothing comparable about a sector that can later couple to the observed system, or about observations that directly access the added factor. Slow coupling gives only the separate approximation stated in [GR §3.2], not exact observational equivalence.

This distinguishes redundancy between descriptions from unresolved facts within a model. Hidden degrees of freedom may store information that affects future observations even when a present record does not distinguish their states. Their capacity and coupling can therefore matter physically. Conversely, an exactly decoupled ancillary enlargement adds no distinction to the fixed channel. Neither case proves that every deeper implementation belongs to one observational equivalence class.

Starting from observation reverses the direction of the explanatory question: the aim is to reconstruct structure consistent with accessible records. It does not remove the need to state the dynamics, preparations, readouts and structural premises used in that reconstruction. The channel-level ancillary identity does not prove a complete gauge classification, and it does not terminate the regress of explanation.

The justificatory regress is cut by declared premises rather than dissolved by the ancillary theorem. Recurrence, finiteness and the other assumptions retain the statuses given in the technical sources [Main §3.1, §4.5]. Mathematical results state what follows from those premises; they do not supply a proof of every premise or identify every possible physical implementation.

The framework's position in this comparison consists of declared premises, mathematical results with their stated hypotheses, and open physical reconstruction obligations. An independently prepared, normalized, exactly decoupled ancillary factor is redundant for the fixed observed channel. A complete observable-

to-substratum equivalence remains open. The posit ledger records assumptions and limits; it does not supply the missing classification proof.

What distinguishes the framework is the approach: starting from a finite deterministic substratum and deriving the fixed-basis quantum representation as an exact theorem and stating the coherent local operational lift separately — rather than adding observer degrees of freedom to an existing quantum-gravitational formalism. The contemporary programs work within quantum gravity and identify the observer's role; the framework works underneath quantum mechanics and identifies how quantum mechanics arises from observer-embeddedness. Both projects are pursuing structural realism about observation; they are doing so from different directions.

3.9 What follows

The hierarchical structure developed in this chapter — the observation axis A through D, the gauge axis G1 through G4, the orthogonality between them, the stratified predictions at the various intersections, the fixed-interface examples of redundancy relative to specified observations, the separate premise checks of the no-go theorems within one architecture, the philosophical lineage tracing back through Spinoza and Cusa — provides the framework's full structural-realism position. Chapter 2's commitment to $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$ is now located at a specific intersection (Level D $\times$ Level G3) within a broader structure, with weaker realism claims at broader intersections and stronger claims unavailable.

Chapter 4 develops the framework's methodology: the single-foundational-commitment derivation pattern, the architecture-level no-go engagement as a systematic methodological principle, the classification of claims by derivational rigor, the framework's pre-registration of falsification conditions. The methodology chapter takes the structural realism developed here as input and articulates the methodological discipline the framework's program follows.

Chapter 9 develops the universality-class structure and the framework's relationship to alternative unification programs — string theory, loop quantum gravity, causal sets, asymptotic safety — at the Level C $\times$ Level G4 intersection. The OI-string relationship in particular receives extended treatment there, with the four-feature audit (Born rule, channel-level unitarity, memory/divisibility, commutant gauge-invariance) characterizing the cross-framework universality-class membership and identifying what transports between the two programs and what does not. The negative technical result on substratum-level equivalence (that string-theoretic matrix models fail OI's A2 and A5) and the positive result on universality-class equivalence are both developed there.

Parts II–IV develop the framework's empirical content at the various levels identified in this chapter. The Level G3 $\times$ Level D predictions — the twenty-two Standard Model retrodictions, the JUNO neutrino-mixing angle, the Bekenstein-Hawking 1/4 factor, the MOND acceleration scale, the dark-sector predictions — are developed quantitatively. The Level G2 $\times$ Level D structural commit-

ments — the Standard Model gauge group, the three-generation structure, the anomaly-free hypercharges — are developed as derivations from the cubic-group commutant. The Level G4 × Level C universal features — the Born rule, channel unitarity, memory/divisibility (accessible memory where the hidden sector reads back its record (C4), and, on any finite reversible substratum, recurrence-cycle P-indivisibility wherever a rooted visible map is not a permutation), the commutant restriction pattern — are taken as given content of the framework's foundational theorems.

The framework's hierarchical structure is what makes the framework different from a single bold claim about reality. It is a coordinated set of claims at different levels of generality, with different empirical predictions, different falsifiability conditions, and different philosophical commitments at each level. The framework's strongest realism — at Level D × Level G3 — is also its narrowest. The framework's broadest realism — at Level C × Level G4 — is also its most universal. The book's role is to develop these levels with the technical and empirical content that distinguishes one level from another.

Chapter 4

Methodology

4.1 What this chapter develops

The framework's content, developed in Chapters 1 through 3, rests on a specific methodological discipline. The discipline is not the framework's content; it is how the framework's content is produced, audited, and made available for evaluation. Different methodological choices would produce different frameworks even from the same starting commitments. The choices made here are worth articulating explicitly, because they distinguish the framework from other unification programs that begin from different methodological commitments and reach different conclusions.

Four methodological commitments organize the framework's program. First, the framework operates from *a single foundational commitment* — the starting point that observation occurs — and derives the rest as theorems given explicit assumptions adopted for the model: the structural assumptions A1–A6 catalogued in §4.2, site factorization and the nearest-neighbour range ([Main §4.5]), and the named hypotheses stated where they enter. That commitment is itself structured: examined closely (Chapter 1), it resolves into two axioms — that tokened differentiation occurs, and that it recurs — thematically one principle but, in honest count, two posits, one indubitable and one substantive. "Single foundational commitment" names the discipline of taking only this one starting

point from outside the theorems, not a claim that the starting point is a single indivisible fact. Other foundational physics programs commit to multiple primitives (string theory commits to extended one-dimensional objects, supersymmetry, and a specific dimensional structure; loop quantum gravity commits to a graph-theoretic discrete geometry and a specific quantization procedure). The framework's discipline is to commit to as little as possible at the foundational level and derive as much as possible from the minimal commitment. The methodological cost is steeper derivations; the methodological benefit is fewer adjustable parameters and a cleaner audit trail.

Second, the framework *classifies its claims by derivational rigor* using an explicit scheme. Strictly parameter-free structural predictions are labeled S; conditional structural predictions that depend on a stated open assumption are C; layered conditional predictions awaiting closure are L; retrodictions fitted to observation are R; predictions taking a phenomenological input are P; predictions inheriting from a mass-chain input are M; predictions explicitly using an empirical input are E. Every prediction the framework makes carries an explicit classification, with the strongest claims reserved for S-class derivations and weaker claims honestly labeled. The discipline of this classification — refusing to overstate rigor — is part of what makes the framework's empirical record evaluable.

Third, the framework *examines each no-go theorem's substantive premises.* Bell, Kochen-Specker, PBR and Frauchiger-Renner constrain different objects. The two-level substratum-emergent architecture and trace-out locate those objects, while the arguments remain separate. The adopted Bell branch requires ontic parameter dependence under measurement independence, with H-Bell open. Kochen-Specker concerns the noncommuting operational algebra; PBR concerns ontic preparation independence; Frauchiger-Renner concerns the consistency and applicability of observer-relative predictions. Chapter 3 §3.7 develops these distinctions, which this chapter uses as a method for checking premises.

Fourth, the framework *pre-registers its falsification conditions.* A framework that derives rather than posits must be willing to specify what would invalidate the derivation. The pre-registration distinguishes structural commitments — which the framework stands or falls on — from solution-specific properties that lie outside its scope. The Class A conditions identify structural theorems whose violation would falsify the framework; Class B identifies parameter-free retrodictions whose precision-upgrade violation would falsify; Class C identifies broader framework-level commitments. The discipline is to specify these conditions before the relevant observations are made, not after.

The chapter has three pieces of work. §4.2 develops the single-foundational-commitment pattern, including the derivation tree from the starting commitment through the framework's theorems and predictions. §4.3 develops the claim-classification scheme, including the S/C/L/R/P/M/E categories and the four-layer framing that locates predictions within the substratum-to-emergent stack. §§4.4–4.6 then develop the separate no-go premise checks as a methodological principle, the pre-registered falsification conditions across Classes A, B,

and C, and the boundaries of what does not falsify the framework. §4.7 closes the chapter with a forward pointer to Parts II–IV, where the methodology is applied to the framework's specific empirical content.

4.2 The single-foundational-commitment pattern

The framework begins from one premise. Observation occurs: an observer records distinguishable outcomes of interactions with a system not wholly under the observer's control. This is the framework's founding commitment; the structural assumptions A1–A6 catalogued below, with site factorization and the nearest-neighbour range, are explicit additional restrictions it also takes (§4.2; [Main §4.5]), and named hypotheses enter where they are stated. Every other content — the lemmas of finiteness and causal partition and determinism, the conditions C1–C4 for embedded observers, the central theorem that the memory-bearing description — with its universally admitted quantum representation — emerges under those conditions ([Main §3.4]), the reconstruction theorem from observed physics back to the substratum, the Standard Model derivation from the cubic-group decomposition, the gravitational sector from horizon thermodynamics — is derived from the foundational premise together with those assumptions and hypotheses.

The methodological discipline of single-foundational-commitment derivation operates as a constraint at every step of the framework's construction. When a new structural feature is needed, the framework must derive it from existing content rather than add it as a new premise. When a new prediction is sought, the framework must trace its derivation back to the foundational commitment through some chain of theorems and identifications. When a new structural assumption appears necessary, the discipline forces an explicit accounting: either the assumption is itself derivable from the foundational premise (in which case it is a theorem, not an assumption), or it is a new commitment that needs to be acknowledged as such (in which case the framework's foundational economy has been compromised).

The framework's content satisfies this discipline imperfectly but explicitly. The structural assumptions A1–A6 identified in Chapter 2's reconstruction theorem (finiteness, determinism, bounded coupling, center independence, linearity, background independence) are not derivable from the foundational premise; they are restrictions on the class of substrata the reconstruction considers. The discipline requires acknowledging this honestly: the framework's uniqueness is *within* the class defined by A1–A6 and the nearest-neighbor restriction, not absolute. Alternatives outside this class would require separate derivations. The framework's defense of A1–A6 is an argument from the empirical inputs of physics, not a derivation from the foundational premise — A1 is supported by the holographic entropy bound, A2 by the structural requirement that observer records not be erasable, A3–A6 by the empirical content of the observed gauge structure — and empirical motivation does not make an assumption logically necessary. The same holds for site factorization, the identification of the configuration

space as a product over sites: given it, the dynamics determines the coupling graph, but the dynamics alone has not been shown to select it. The honest acknowledgement is that the framework rests on the foundational premise *plus* these assumptions, identified explicitly so that any reader can evaluate them. The posit ledger of [Main §4.5] collects them: it is the audited inventory of the explicit premises the derivations consume, not an enumeration of every physical assumption one might make.

The benefit of the discipline is a clean audit trail. Every prediction the framework makes can be traced back through the chain of theorems to either the foundational premise or to one of the explicitly-acknowledged structural assumptions. The Cabibbo angle $1/(\pi\sqrt{2})$ traces back through the cubic-group decomposition (Chapter 5), to the cubic-symmetric coupling matrix (the substratum's wave equation), to the wave equation's uniqueness given A4, A5, isotropy and the nearest-neighbour range in the spatial site factorization (Chapter 2), to the underlying bijection structure (Chapter 1). The Bekenstein-Hawking 1/4 coefficient traces through boundary mode counting (Chapter 7), to the thermal-self-consistency gap equation — where the infrared detailed-balance condition H-slope enters as a named condition — to the substratum's bijection structure (Chapter 2), to the embedded-observer construction (Chapter 1). The chain of derivation is explicit at each step, and any link in the chain that turns out to be incorrect propagates predictably to the downstream predictions.

The cost of the discipline is that some natural extensions of the framework require derivational work that has not yet been done. The framework currently treats CP-violating phases in CKM and PMNS as solution-specific inputs rather than as derivations; the cosmological initial conditions are taken as given rather than derived; the specific values of fermion masses depend on one empirical input (the strange-quark mass) rather than being derived from the substratum alone. These are not failures of the framework but acknowledged scope limits. The methodological discipline distinguishes between "the framework predicts X" (X follows from the foundational premise plus the structural assumptions) and "the framework is consistent with X but does not predict it" (X is solution-specific or depends on inputs the framework has not yet derived).

A useful diagnostic for the discipline is what the framework does *not* commit to. The framework does not commit to a specific interpretation of probability beyond standard Kolmogorov probability. The framework does not commit to a specific resolution of the measurement problem beyond the algorithmic-reading developed in Chapter 1. The framework does not commit to specific values of free parameters in the emergent quantum field theory beyond what the reconstruction theorem forces. The framework does not commit to a specific cosmological history beyond an expanding homogeneous general-relativistic background, taken as given, on which, with the null energy condition, the apparent-horizon area does not decrease. These non-commitments are part of the foundational economy: each additional commitment would expand the framework's claims, and the discipline is to make only those commitments that the foundational

premise plus its derivations require.

The single-foundational-commitment pattern distinguishes the framework from broader unification programs in a specific way. String theory commits to extended one-dimensional objects as the fundamental entities, plus supersymmetry, plus extra dimensions, plus a specific moduli structure. Loop quantum gravity commits to a graph-theoretic discrete geometry, plus a specific quantization procedure, plus a specific representation theory of the spatial diffeomorphism group. Causal sets commit to a Lorentz-invariant discrete causal structure, plus a specific dynamics for set growth. Each of these is a multi-commitment foundation, with each commitment adding to the framework's content. The framework here is a single-commitment foundation, with all content derived from the one premise plus its structural lemmas. Whether this is a methodological virtue or a methodological constraint depends on whether the derivations actually close — whether single-commitment foundations can produce the content that multi-commitment foundations produce by direct positing. The framework's claim is that they can; Parts II–IV develop this claim with the specific quantitative content.

A further comparison with string theory is worth drawing, because it clarifies what the single foundational commitment is and is not. String theory's landscape — the vast space of consistent vacua, among which the observed one is often held to be fixed by anthropic selection — is sometimes proposed as a model for how a theory can be general in its laws yet specific in what we observe. It is tempting to ask whether the framework's foundational commitment is similarly a selected feature: one option among many, picked out in our case because only some options permit observers. The answer sharpens what the commitment is. The framework does have a "general, then selected" structure — but it lives *downstream* of the foundational commitment, not at it. The emergent description is gauge: it depends on the partition, with the substratum gauge group identifying which structure is observable; and the partition that supports an observer is then selected, by the structural observer-selection theorem of Chapter 1, conditional on the effective-mixing hypothesis (EM) and its rate condition ([Main §4.6]), as one of those meeting the embedding conditions. This is the framework's counterpart of "general law, particular vacuum," and unlike the landscape it is derived — a constructively generated gauge structure and a selection theorem, rather than a posited space of vacua and an anthropic measure that has resisted rigorous definition. What this structure does *not* reach is the foundational commitment itself. The commitment, distilled, is two axioms: that tokened differentiation occurs, and that it recurs (Chapter 1). Neither can be a landscape variable, because a landscape is a space of differentiated, temporally-extended worlds — it presupposes both axioms in order to have points at all. Anthropic selection picks among worlds; it cannot pick whether there is a world of the relevant kind, since the selection apparatus, a measure over possibilities, already requires the temporal domain it would be invoked to explain. The "gauge in general, specific in our case" intuition is therefore correct and is realized in the framework — at the level of partitions and emergent descriptions,

one level below the two axioms, which are what must hold for there to be a space of possibilities to select within.

4.3 Classification of claims by derivational rigor

A framework that derives rather than posits must be honest about how rigorous each derivation actually is. The framework's discipline is to classify every prediction by its derivational rigor, using an explicit scheme with seven primary categories. The categories are not interchangeable: the difference between a strictly parameter-free structural prediction and a retrodiction fitted to observation is substantial, and conflating them would overstate the framework's empirical record. The classification scheme is part of what makes the framework's claims evaluable.

Class S — strictly parameter-free structural prediction. A prediction is S-class if its derivation chain has no fitted parameter at any step and no open link specific to the entry; matter-sector entries rest, as every matter-sector result does, on the conditional physical-carrier chain (H-spin' and H-χ'). The Wolfenstein parameter $A = \sqrt{2/3}$ is S-class: it is the sine of a fixed angle of the cubic lattice, with no fitted constants. The Higgs quartic boundary condition $\lambda(M_{\rm Pl}) = 0$ is S-class: it follows from the composite-Higgs structure of the singlet taste, with no parameters tuned to observation. S-class predictions are the framework's strongest empirical commitments, and their precision-upgrade falsification would refute the framework directly without any recoverable error mode beyond computational mistakes in the specific derivation.

Class C — conditional structural. A prediction is C-class if its derivation chain walks modulo a stated open assumption. The empirical match is unaffected by the open assumption, but the prediction's structural-rigor status is contingent on the assumption's resolution. C-class predictions are weaker than S-class because their derivation is conditional, but the empirical content can be just as sharp. The Cabibbo angle $1/(\pi\sqrt{2})$ is C-class: it follows from a single Brillouin-zone distance with no fitted constants, but its identification with the taste-changing transition leaves the emergent-theory dressing open. The Koide relation is C-class: it follows from a cubic-group quadratic Casimir with no parameters tuned to observation, but its amplitude $A^2 = C_2$ is not yet derived from the dynamics. Conditional consistency relations carry conditions of the same kind but are not counted as predictions: the Bekenstein-Hawking 1/4 coefficient, the ratio of the Euclidean KMS period to the Jacobson prefactor, is fixed once the named horizon, frame, state and action-scale calibration conditions hold ([GR §3, §8.5]), and agrees with the normalization fixed by the Hawking temperature and the first law. The distinction is methodological honesty about what has been derived versus what has been assumed.

Class L — layered conditional. A prediction is L-class if its derivation walks given Layer-2 inputs, with closure path being a specific active research direction. L-class is a refinement of C-class that identifies *which* derivational

work would move the prediction to S-class. The pattern is: "the prediction is S given that Direction X closes." Active research directions in the framework include the analytic dispersion-relation calculation at Direction 10, the structural derivation of certain spinor-structure identifications, and others. L-class predictions become S-class as the relevant directions close, and the framework's rigor improves monotonically over time.

Class R — retrodiction. A prediction is R-class if its value is fitted to observation by construction. R-class predictions cannot falsify in the precision-upgrade sense — they were fitted, so they match by definition. The structural content lives in *what is predicted given the fit* rather than in the fitted value. For example, the SU(2) and SU(3) gauge couplings at the Z pole are R-class in the framework: three fitted parameters (δ_0, A, B) against three observed couplings produces zero residual by construction. The framework's structural content here is the *form* of the gauge-coupling fit (universality of δ_0, the resummation form, the structural prediction $1/\alpha_0 = 23.25$ from the 1-loop staggered vacuum polarization, not the matching of the three couplings to observation). The $A \cdot B$ calculation is a separate model-conditional numerical cross-check, with extrapolated values $\approx$ 48–62 across leading-power assumptions and an inconclusive comparison with the fitted product.

Class P — phenomenological input. A prediction is P-class if it takes a specific quantity from experiment because the framework does not yet derive it. The strange-quark mass m_s is a phenomenological input in the framework's current content: it sets the overall fermion mass scale, but the framework does not derive it from the substratum. Acknowledging the input is methodologically necessary — the framework cannot claim to predict m_s — but does not undermine the predictions that derive from it once it is fixed.

Class M — mass-chain inheritance. A prediction is M-class if its empirical match follows from a single upstream empirical input. The electron and muon masses, given the tau mass, are M-class: they inherit from the tau mass through the Koide relation rather than being independently derived. M-class predictions are not retrodictions — the Koide relation is a structural prediction of the framework, and the inheritance through it is a derivation — but their empirical match depends on the upstream empirical input being what it is observed to be.

Class E — explicit empirical input. A prediction is E-class if some specific input enters the derivation chain from experiment, acknowledged honestly. E-class is broader than M-class, covering inputs that are not specifically masses. The Jarlskog phase η, for example, enters the CKM derivation as an empirical input rather than as a derivation from the substratum.

The four-layer framing. Orthogonal to the S/C/L/R/P/M/E classification is a four-layer framing that locates predictions within the substratum-to-emergent stack. The two axes are independent: an S-class prediction can be at any of the four layers, and a Layer-1 prediction can be S, C, L, R, P, M, or E class.

Layer 0 covers gauge structure: which gauge groups exist, which representations appear, structural constraints from the substratum's coupling structure and cubic-group representation theory — C1–C4 are diagnostics of the observer, not inputs to this layer. The Standard Model gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ is Layer 0. The three-generation count is Layer 0.

Layer 1 covers structural form: predictions from pure substratum geometry or representation theory, with no operator-relation inputs. The Cabibbo angle $1/(\pi\sqrt{2})$ is Layer 1: it follows from a Brillouin-zone distance. The Wolfenstein parameter $A = \sqrt{2/3}$ is Layer 1. The Koide angle $\theta_0 = 2/9$ is Layer 1.

Layer 2(a) covers operator-relation structural predictions: the structural form of a relation, given that the relevant operators exist. The Gatto-Sartori-Tonin relation applied to the Layer-1 Cabibbo derivation is Layer 2(a).

Layer 2(b) covers solution-specific predictions within a fixed operator structure: mixing-angle values determined by the specific bijection's parameters.

Layer 3 covers mass-scale and bijection-specific predictions: the specific values of m_s, μ_c, μ_w, etc., that pick out which bijection φ within the equivalence class our universe occupies.

The framework's predictions span the four layers. Layer 0 and Layer 1 predictions are the framework's most distinctive content, because they involve only the substratum's structural features and no specific solution-level inputs. Layer 2(a) predictions are nearly as distinctive but conditional on the operator structure. Layer 2(b) and Layer 3 predictions involve more solution-specific content and are accordingly less distinctive — they would differ for different bijections within the same equivalence class.

The substratum / emergent / mixed architectural classification. A third axis classifies each load-bearing element of a calculation by where it sits in the substratum-to-emergent architecture. *Substratum* elements operate on the bijection (S, φ) on the cubic lattice with its coupling matrix; predictions at this level are pure structural ratios from geometry and representation theory. *Emergent* elements operate on the unitary quantum field theory after the trace-out: induced gauge couplings, renormalization-group running, Coleman-Weinberg potentials, Z-factors; predictions at this level are perturbation-theory and RG outputs. *Mixed* elements involve QFT machinery in the derivation but with specific substratum-level inputs constraining the output — the most typical case for the framework's quantitative predictions in Chapter 5.

The framework's general rule is *group structural, couplings emergent*. The gauge group is determined at the substratum level by the cubic-group commutant, conditionally on H-link and an octahedrally equivariant condensate (Chapter 5 §5.4–§5.5); the gauge couplings at the Z pole are determined at the emergent level by RG running from the lattice-determined Planck-scale coupling.

Why the classification matters. A framework that classified all its predictions as S-class would either overstate its empirical record (if some predictions

are actually conditional or retrodicted) or be implausibly successful (if all predictions actually walk end-to-end with no inputs). The framework's explicit classification of each prediction by its derivational rigor is a methodological discipline that allows the empirical record to be evaluated honestly. The strongest claims live in the S-class entries; the conditional claims live in C and L; retrodictions and inputs are acknowledged honestly. The framework's twenty-two Standard Model retrodictions split across these categories — some S-class, some C-class, some R-class — and the split is part of what readers should know when evaluating the framework's empirical record against the framework's structural commitments.

4.4 Engaging the no-go results through one architecture

The framework is a hidden-variable theory in the literal sense: there is a deterministic substratum with definite states, and the observer's stochastic description is obtained by marginalizing over what they cannot see. The twentieth-century foundations literature contains a series of no-go theorems ruling out hidden-variable theories satisfying various structural conditions. The framework must state how each of these results applies to it. Chapter 3 §3.7 developed the technical content of the engagement; this section develops the methodological principle that organizes it.

The principle is straightforward to state. Each no-go theorem identifies a structural property of would-be hidden-variable theories and proves that the property is incompatible with quantum predictions. The property is some conjunction of conditions on the underlying hidden-variable structure and the observer's measurement procedure. The framework's two-axis hierarchy, developed in Chapter 3, locates the level at which each condition applies. Whether the framework satisfies or violates it there is a separate check for each theorem: identify the hierarchy level at which the no-go theorem's assumption sits, then determine whether the proposed completion satisfies that assumption at that level.

The four-step procedure. Applied to a no-go theorem in the foundations literature, the procedure has four steps.

First, identify the theorem's structural assumption. Bell's theorem assumes factorizability $P(a, b \mid x, y, \lambda) = P(a \mid x, \lambda) \cdot P(b \mid y, \lambda)$. Kochen-Specker assumes non-contextual value assignments. PBR assumes preparation independence — that independently prepared systems have a product joint ontic-state distribution. Frauchiger-Renner assumes the universal applicability of quantum mechanics across nested observer-modeling.

Second, identify the level of the framework's hierarchy where the assumption sits. Bell's factorizability is a property at Level G3 $\times$ Level D, where the joint transition matrix between visible-sector subsystems is defined. Kochen-Specker non-contextuality is a property at Level G2 $\times$ Level D, where observables are

defined as operators on the emergent Hilbert space. PBR preparation independence is a property at Level D × Level G3, where the substratum's joint-system structure is specified. Frauchiger-Renner universal applicability is a claim at Level G2 × Level D applied to substratum-level objects.

Third, locate where the framework meets or violates the theorem's premises at the correct level. Bell is the sharpest case: for a response-complete deterministic completion, Bell couplings that do not connect the wings before readout, one-step own-wing setting interventions and a ± 1 outcome in every run, plus a setting-independent pre-setting ensemble, imply Bell factorization and $|S_{\mathrm{CHSH}}| \leq 2$. The framework therefore does not obtain quantum Bell violation from outcome-independence failure in an indivisible stochastic law; it retains measurement independence and gives up **ontic parameter independence**, with operational no-signaling required. PBR's preparation independence — the postulate that independently prepared systems have a product joint ontic-state distribution — is not a premise of bare OI. Co-embedded preparation devices may share preparation-relevant hidden or boundary data, so unconditional product independence is not guaranteed; the graph causal cone by itself does not derive either a conditional product distribution or the standard quantum tensor-product composite. Frauchiger-Renner requires a check of the proposed completion's full protocol and its probability-one certainty, consistency and single-outcome premises; multiple observer partitions alone do not establish failure of those premises. The fourth theorem, Kochen-Specker, does not fit a "satisfied below, violated above" form: its assumed property, a noncontextual valuation of the observables, holds at neither level — the framework — conditionally on the operational bridge reconstructing the standard observable algebra — carries the operational theory, which by Kochen-Specker's own content admits no noncontextual valuation, and the substratum carries a definite configuration that is not a valuation of the emergent observables at all. The framework's contextuality is forced, under that same conditional bridge, by equivalence at the operational layer — and what the framework adds is the account: contextuality is the partition-dependence of the trace-out. The common factor across all four is therefore the architecture — two levels and a trace-out — not a single property-placement move repeated.

Fourth, verify that the framework's predictions match quantum mechanics in the regime where the no-go theorem applies, without silently importing the very condition under test. For Bell, verify the deterministic completion directly: if its setting interventions, applied at one step, each read and write only their own wing, its Bell-relevant graph does not connect the wings before readout, every run records a ± 1 outcome, and its pre-setting ensemble is setting-independent, CHSH is capped at 2. The framework's quantum branch therefore has ontic parameter dependence. For the specific SM/GR representative, the nearest-neighbor **reference** graph cannot itself supply that parameter dependence in a Bell protocol with one-step own-wing setting interventions whose wings lie outside each other's graph cone. The existing state-dependent graph $G(x)$ supplies the native target: preparation-indexed adjacency (or an equivalent non-

local response), with H-Bell the open requirement that the prepared graph family preserve operational no-signaling and the metric/Ollivier–Ricci continuum limit. The measurement-dependent alternative is not adopted; Chapter 1 gives a model-independent necessary lower bound and a separate causal benchmark; the stochastic causal-local Tsirelson theorem is retained only at its weaker stochastic level. Kochen-Specker, PBR, and Frauchiger-Renner concern the coherent operational structure and therefore inherit the operational-completion qualification: their standard quantum statistics are recovered, within complex matrix kinematics taken as given, where the five completion conditions of [GR §3.3] hold, not merely from universal fixed-basis representability.

Common descriptions, separate premises. A finite bijection and its trace-out supply the same substratum and observable objects for these questions. They do not discharge the no-go hypotheses by themselves. Bell's ceiling still applies to a measurement-independent deterministic completion with local responses; the adopted branch therefore requires ontic parameter dependence and a realization meeting H-Bell. The other cases retain their separate operational-algebra, preparation-independence and observer-consistency assumptions. Each conclusion follows only when its own argument and hypotheses hold.

This per-theorem verification is a genuine discipline, not a rhetorical disadvantage. In particular, determinism by itself does not settle Bell's measurement-independence premise. The framework **adopts** the measurement-independent branch rather than using setting dependence as its explanation; any microscopic mixing/decorrelation argument can motivate that adopted laboratory premise but is not the Bell-violation mechanism. The required Bell violation is instead carried by ontic parameter dependence, with operational no-signaling checked separately. The same discipline applies to the coherent-instrument assumptions in the other no-go theorems.

Methodological benefit. The benefit of the architecture-level engagement is that the framework's structural reading of each no-go theorem is *informative*. Identifying how each assumption meets the two-level structure explains *why* quantum mechanics has the features the no-go theorems identify, not just that it does. Bell's theorem forces the framework to locate nonlocality precisely: in the adopted branch it is ontic parameter dependence, not mere stochastic outcome dependence or indivisibility. Kochen-Specker's contextuality is informative about the partition-dependence of the trace-out. PBR identifies a preparation-independence question for separated subsystems that may share boundary data; the trace-out alone does not determine whether a proposed completion satisfies that premise. Frauchiger-Renner's inconsistency-paradox is informative about the levels of the framework's hierarchy.

The no-go theorems become, in this reading, structural diagnostic tools rather than obstacles. Each one identifies a specific structural feature that any hidden-variable theory matching quantum mechanics must have. The framework's identification of the relevant features through hierarchy levels makes the structural content of the theorems available for use in further work — in particular, in

identifying which features any successful unification framework must satisfy.

Application to new no-go theorems. The procedure is open-ended. Any future no-go theorem in the foundations literature can be subjected to the same four-step analysis. The framework's commitment is to apply the procedure honestly to each new result rather than to claim general immunity. If a no-go theorem turns out to identify a structural property that the framework cannot satisfy at any level — or to require simultaneous satisfaction of properties the framework can satisfy only at different levels — the framework would face genuine difficulty.

An additional no-go theorem requires an analysis of its actual hypotheses on the proposed completion. The shared substratum and observable descriptions identify the objects to examine, but do not guarantee that a required premise fails or that the completion avoids the theorem's conclusion.

The common architecture organizes the questions, while each no-go theorem retains its own substantive test. Bell still imposes the local-response ceiling under measurement independence, H-Bell remains an obligation of the proposed physical realization, and the other cases retain their separate operational, preparation and observer-consistency assumptions.

4.5 Pre-registered falsification conditions

A framework that derives rather than posits must be willing to specify what would invalidate the derivation. The framework's discipline is to pre-register specific falsification conditions, stating in advance which observational outcomes would refute the framework's structural commitments and which would refute only narrow predictions while leaving the broader structure intact. The pre-registration distinguishes structural claims — which the framework stands or falls on — from solution-specific properties that lie outside the framework's scope.

The conditions are organized into three classes. Class A identifies structural theorems whose violation would falsify the framework directly. Class B identifies parameter-free retrodictions whose precision-upgrade violation would falsify, with sub-classes reflecting the rigor of the retrodiction. Class C identifies broader framework-level commitments whose contradiction would refute particular framework claims while leaving others intact.

Class A — structural theorems. The framework derives specific structural features of physics as theorems from the foundational premise and the structural assumptions. Each derived feature is a falsification target if violated.

The strong-CP phase, not fixed. T-invariance of the substratum bijection combined with detailed balance narrows the QCD angle to $\theta \in \{0, \pi\}$ and gives reciprocal transition counts, but does not make the emergent Hamiltonian T-invariant, so $\bar{\theta}$ is not fixed by the construction (SM §5). The framework makes no commitment on the neutron electric dipole moment; a measured $\bar{\theta} \neq 0$

would constrain mechanisms proposed for the named condition H-det, not the T-symmetry of the substratum.

The gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ *exactly.* Conditionally on H-link, the framework derives this gauge group from the unitary group of the bicommutant of the cubic-group action (equivalently, the stabilizer of the generic equivariant condensate) on the six internal components. Any observation of a stable beyond-Standard-Model gauge structure — an additional Z' interpretable as a new gauge boson rather than a composite resonance, stable exotic matter transforming under a novel gauge group, or any confirmed extension of the visible-sector gauge content — falsifies the cubic-group decomposition.

Three fermion generations. Conditionally on H-link single-copy, the framework derives the generation count of three from the link count, $K = 2dm$ with $m = 1$, locking $K = 2d = 6$ internal components and giving three spin-1/2 staggered tastes. Any observation of a stable fourth-generation fermion with Standard Model quantum numbers falsifies the taste-reduction derivation. Non-stable exotic states consistent with composite or resonance interpretation do not.

Majorana neutrinos with normal mass ordering. The framework expects this on the branch where the singlet taste hosts the Higgs and no gauge-singlet fermion is present (SM §8.4): Majorana because no gauge-singlet neutrino is present, normal ordering as a heuristic consequence of the taste-breaking structure of the neutrino mass matrix rather than a theorem. Inverted mass ordering, confirmed to high significance, falsifies the heuristic on that branch. Confirmed Dirac nature of the neutrino — for example, a null result for neutrinoless double-beta decay at the sensitivity that would require Majorana masses well below the framework's expectation — falsifies the branch's expectation that no right-handed neutrino exists in the spectrum.

The Tsirelson bound. The value $2\sqrt{2}$ belongs to the cited **stochastic causal-local indivisible construction** ([Main §3.3]), not to a Bell-local deterministic completion and not to generic P-indivisibility. The deterministic OI layer reaches quantum Bell violation, under measurement independence, only on the ontically parameter-dependent branch. An observed loophole-free CHSH value above $2\sqrt{2}$ would falsify standard quantum mechanics and any claim that the relevant OI operational lift lands in the standard quantum Bell set; it would not, by itself, falsify P-indivisibility.

Exact unitarity of the visible-sector dynamics at accessible scales. The framework derives unitary quantum evolution as the exact embedded-observer description at every accessible mass and coherence scale, with dissipative corrections structurally suppressed to the dissipator scale of Chapter 1 §1.6. Any confirmed intrinsic-collapse signature — anomalous decoherence not attributable to environmental channels, scaling with mass or superposition size as objective-collapse models (GRW/CSL, Diósi–Penrose) require, at sufficient confidence to exclude systematics — falsifies the framework's physical identification directly. Because the framework's predicted deviations are pinned orders of magnitude below the

collapse-model band, a positive detection cannot be absorbed as a framework correction. The current frontier is matter-wave interference at macroscopicity $\mu = 15.5$ (Pedalino et al. 2026; Chapter 1 §1.6), consistent with exact quantum behavior; the frontier's continued advance is a standing test rather than a completed one.

Leading Lorentz-invariance-violation coefficient. The framework derives a specific subluminal dispersion relation with the leading correction at order $(E/M_{\mathrm{Pl}})^2$, with a physical, direction-dependent anisotropy whose orientation-averaged coefficient is $4/45$ at one nat of entropy per boundary mode ($4\sigma/45$ for an entropy σ per mode; $\langle\delta\rangle = 1/90$; the body diagonals are exactly luminal). The structural content is the lattice coefficient $1/45$ of $(k\epsilon)^2$ and the anisotropy pattern; the amplitude $4\sigma/45$ carries the undetermined entropy per mode, so it is not a parameter-free number, and a measurement of it would constrain σ rather than independently falsify the coefficient. The anisotropy's absolute orientation in space is gauge, but its $\ell = 4$ cubic-harmonic pattern and the direction-differences are physical — the cubic structure is an empirically-selected input, not a gauge choice (see the Standard Model and Substratum papers). Two structural features of this prediction matter for falsification. First, the dispersion has no linear-in-E/M_{Pl} term — any demonstration of linear Lorentz-invariance violation at any scale falsifies the framework's substrate-level commitment to the cubic-lattice wave equation. Second, the dispersion is subluminal everywhere on the momentum sphere; superluminal LIV anywhere falsifies. Current quadratic-LIV bounds from gamma-ray-burst time-of-flight measurements reach roughly thirteen orders of magnitude below the Planck scale, so the $4\sigma/45$ coefficient is not near-term testable at the magnitude level; linear-LIV tests, by contrast, are already at sensitivities where any positive detection would falsify. These statements describe the dispersion at tree level; whether the absence of a relevant (low-dimension) Lorentz-violating operator survives radiative corrections in the interacting theory is a separate, open question — the Lorentz fine-tuning problem common to discrete-substrate approaches — discussed in the Standard Model paper. The tree-level prediction and its falsifiers stand regardless; what is open is the loop-level protection — and for the rotational sector it turns on whether the emergent theory preserves the octahedral symmetry exactly at loop level, with an aligned condensate (Chapter 19 §19.3.7): octahedral-invariant dynamics cannot generate the leading rotational anisotropy at any order, and the power-divergent operator mixing that cubic symmetry permits is mixing among angular-momentum-covariant operators rather than the generation of a lower-dimension rotational-violating operator (see the Standard Model paper). The protection mechanism available here (the framework not being supersymmetric) is renormalization-group emergence — Lorentz-violating couplings flowing to an infrared-attractive Lorentz-invariant fixed point (Chadha–Nielsen; Bednik–Pujolàs–Sibiryakov), natural because the trace-out is a coarse-graining — with the documented caveat that the inter-species velocity flow is slow. The causal-set discreteness results (Bombelli–Henson–Sorkin) independently confirm that a finite-valency,

locally-coupled discrete structure cannot be microscopically Lorentz-invariant, so the invariance here is necessarily emergent rather than fundamental.

Class B — parameter-free retrodictions. The framework produces twenty-two classified quantitative retrodictions that follow from the structural content with no adjustable parameters beyond a small acknowledged empirical input set and three matched coupling parameters. These are stratified into five sub-classes reflecting their structural status.

Class B-S (structural). These are layer-0 or layer-1 retrodictions with no open link specific to the entry; like every Class B entry they rest on the matter-sector hypotheses H-spin' and H-χ'. The example in this chapter's selection is the Wolfenstein parameter $A = \sqrt{2/3}$ from the off-democratic projection of the cubic generation axes. Any such entry moving outside 3σ on precision upgrade falsifies the framework with no recoverable error mode beyond computational mistakes in the specific derivation.

Class B-C (conditional structural). These are layer-1 retrodictions with no fitted input whose derivation chain has one named open link. The Cabibbo angle $1/(\pi\sqrt{2})$ from a single Brillouin-zone distance is B-C, conditional on the emergent-theory dressing of its identification with the taste-changing transition; the Koide angle $\theta_0 = 2/9$ from a cubic-group quadratic Casimir (matched to 0.001%) is B-C with the Koide amplitude, whose value $A^2 = C_2$ is not yet derived from the dynamics. A precision-upgrade violation would falsify the identification or the closure on which the entry is conditional.

Class B-L (layered conditional, pending open derivations). These predictions follow from the structural form combined with one Layer-2 substrate input or empirical input, with a specific active research direction whose closure would move the prediction to B-S status. The JUNO neutrino-mixing prediction $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2)$ matched at 0.07σ is B-L pending closure of the mixing-matrix derivation in the lepton sector. The mass ratio $m_d/m_s = 1/(2\pi^2)$ is B-L, its conversion from the Cabibbo scale running through the empirical Gatto-Sartori-Tonin relation, and the MOND critical acceleration $cH/6$ is B-L, inheriting the conditions on the de Sitter temperature with its dimensional factor imported from Verlinde's emergent-gravity analysis. Other B-L predictions exist throughout Chapter 7's quantitative content.

Class B-M (mass-chain inheritance). These predictions inherit their empirical match from a single upstream empirical input. The Higgs mass m_H via Standard Model RGE running from the boundary condition $\lambda(M_{\rm Pl}) = 0$, with m_t as the empirical input, is B-M. Class B-M predictions falsify under the same precision-upgrade protocol as B-L, but the inheritance is from an empirical input scale rather than from an open derivation.

Class B-R (retrodictions). The SU(2) and SU(3) gauge couplings at the Z pole are explicitly retrodictions: three fitted parameters (δ_0, A, B) against three observed couplings produces zero residual by construction. Class B-R predictions cannot falsify in the precision-upgrade sense; their value lies in falsifying the

framework's gauge-sector structural commitments — universality of δ_0, the resummation form, the structural prediction $1/\alpha_0 = 23.25$ from the one-loop staggered vacuum polarization, with the $A \cdot B$ finite-lattice calculation providing a separate model-conditional numerical cross-check that remains inconclusive across leading-power assumptions. Any precision-upgrade violation of the structural commitments would falsify the framework's gauge-sector account while leaving the broader Class A theorems intact.

The Class B split is, in honest summary: one prediction is structural with no open link of its own (B-S), two are conditional structural (B-C), six are layered conditional pending open derivations (B-L), two are mass-chain inheritances (B-M), and two are retrodictions (B-R). The thirteen Class B entries combine numerical retrodictions, conditional formulas and inherited mass relations. Their empirical standing is assessed individually: the gauge couplings match by construction. The Bekenstein-Hawking coefficient $A/(4\ell_P^2)$, conditional on the infrared detailed-balance condition H-slope on the horizon reservoir ([GR §3.2]), is a consistency relation with no direct test — GW250114 constrains the area theorem — and is not counted among them. The quoted precision of a measured comparison does not apply to every entry. As experimental precision improves, the discrimination power of the B-S set increases monotonically; closure of the open derivations would move B-C and B-L predictions to B-S over time.

Class C — framework-level commitments. Two broader commitments are implicit in the construction; each is stated with what would contradict it.

Finite classical realizability of quantum processes. The characterization theorem developed in Chapter 1 §1.8 establishes that the memory-bearing description admitted by an embedded observer satisfying C1–C4 lives on a deterministic substratum with bounded classical hidden-sector memory. At finite accuracy, every finite adaptive quantum experiment has a deterministic ε-shadow, while the exact continuum theory at any fixed finite carrier is excluded by the classical-dimension obstruction ([Main §3.4]). Non-Markovian models are permitted and expected (they are how the hidden-sector correlation persistence manifests). No finite-accuracy experiment contradicts this commitment by itself. A memory-witness experiment would contradict the ontology only given the coherent-preparation and instrument bridge, an outcome-independent rule identifying fundamental open-system processes, and a constraint derived from the completed model on what those processes must show; the last is not established ([Main §4.5]).

The dark sector as entropy-displacement, not as a novel particle content. The framework's account of dark energy as boundary-entropy bookkeeping read in the Type II running-vacuum channel, and dark matter as frozen boundary entropy, developed in Chapter 7, is incompatible with the direct detection of a WIMP interpretable as a fundamental particle with Standard-Model-like interactions and no entropy-displacement signature. A confirmed WIMP with such properties falsifies the framework's dark-sector account while leaving the rest of the structure intact — it would require abandoning the boundary-entropy

derivation and either identifying an error in it or accepting that the framework's dark-sector story is wrong even as other derivations survive.

The cosmological commitment. The dark energy sector is subtler than the strict Class A through C taxonomy allows. The framework supplies geometry-dependent dark-energy bookkeeping whose smooth late-time expansion contains an $H^2 - H_0^2$ term — the local form shared by every smooth model — and reads it in the Type II running-vacuum channel; it does not presently derive a distinctive Type II law (Chapter 7 §7.8). The magnitude is bounded at the emergent-QFT scale $|\nu_{\rm QFT}| \sim (M_{\rm SM}/M_{\rm Pl})^2 \sim 10^{-32}$, observationally indistinguishable from ΛCDM. The falsification band is stratified: a DESI Year 5 measurement of $|\nu| \gtrsim 10^{-4}$ at high significance would require additional substratum structure not currently in the framework but would not refute the Type II functional form; a high-significance detection of *phantom crossing* — a $w(z)$ that crosses -1 — would contradict the Type II reading and the smooth-expansion bookkeeping the framework instantiates: the strongest available disconfirmation of its dark-energy account, of a reading the framework selects rather than a law it derives. The sign of a small detected ν is not by itself a structural discriminator: the Type II class admits either sign, the emergent-channel computation gives a negative sign under one reading of its gravitational status and no source under the other (Chapter 7), and a positive detection at the 10^{-4} level would require substratum structure carrying a correlated Lorentz-violation signature. The first outcome is a refinement within the current bracket; the crossing is the strongest disconfirmation available in this sector.

The pre-registration commitment. The author commits to treating the Class A conditions, the Class C commitments, and a Class B retrodiction exceeding 3σ on precision upgrade as framework-falsifying rather than as an invitation to post-hoc rescue. In the case of a Class B violation, the framework will be revisited to determine whether the violation reflects a computational error in the specific retrodiction, an incorrect specific-bijection input, or a structural problem with the derivation; the first two are recoverable, the third is fatal. The framework does not reserve the right to add structural postulates after the fact to accommodate disconfirming evidence. The pre-registration of this commitment is itself part of the falsifiability content of the framework — a framework that pre-registers its falsification conditions and then revises them post-hoc has compromised the discipline that pre-registration was meant to enforce.

4.6 What does not falsify and regimes of deliberate silence

The framework's pre-registered falsification conditions define what would refute the framework's structural commitments. The complement of those conditions — what lies outside the falsification schedule — also deserves explicit articulation. Some observational outcomes that might appear to bear on the framework do not bear on it, because the framework makes no commitment on the relevant

question. The discipline of pre-registering falsification conditions requires equal honesty about what is and is not within the framework's predictive scope.

Solution-specific properties. The framework derives the structural form of physics — the gauge group, the matter content and the generation count (conditionally on H-link, and for the gauge group and matter content on an octahedrally equivariant condensate; Chapter 5 §5.4–§5.5), the parameter-free retrodictions, the thermodynamic content of the gravitational sector. It does not derive every contingent feature of the observed universe. Some features are solution-specific: properties of the particular bijection φ within the framework's equivalence class that describes our universe, as distinct from properties of the equivalence class itself. The framework's commitments are to the equivalence class; solution-specific properties are inputs to selecting the representative within the class, not derivations from the framework's foundational premise.

Several quantities are explicitly solution-specific in the framework's current content.

The absolute scale of fermion masses depends on the strange-quark mass m_s as an acknowledged empirical input. The framework derives mass *ratios* from the cubic-group representation theory and the Gatto-Sartori-Tonin relation, but the overall mass scale is set by one external input. A future precision measurement of m_s at different value would shift all derived fermion masses proportionally without bearing on the framework's structural derivation. The framework predicts the *pattern* of masses given m_s; it does not predict m_s itself.

The CP-violating phases in CKM and PMNS are solution-specific. The construction does not fix $\bar{\theta}$ (§6.3): T-invariance narrows θ to $\{0, \pi\}$ and gives reciprocal transition counts, which leave the Yukawa determinant phase free. The flavor-basis CP phases — the Jarlskog parameter, the Wolfenstein (ρ, η), the PMNS Dirac and Majorana phases — are features of the basis mismatch between up-type and down-type Yukawa mass eigenstates, not consequences of any structural feature of the substratum, and reciprocity does not force them to vanish [SM §5.4]. A reported computation on the staggered reconstruction, whose driver and output are not available for verification, found that an octahedral-equivariant emergent operator gives a vanishing Jarlskog invariant and that restoring the observed value requires octahedral breaking far above the rotational bounds; if it stands, it is a tension for the flavor sector, recorded as unverified [SM §5.5].

The numerical value of the Type II running-vacuum coefficient ν within the stratified falsification band is solution-specific. The framework's content here is the smooth $H^2 - H_0^2$ expansion read in the Type II channel — a structural reading, not a derived law — with the magnitude $|\nu_{\mathrm{QFT}}| \sim (M_{\mathrm{SM}}/M_{\mathrm{Pl}})^2 \sim 10^{-32}$ bounded at the emergent-QFT scale from the emergent-QFT calculation. The Bertini et al. (2025) DESI-era measurement $\nu = -(2.5 \pm 1.3) \times 10^{-4}$ is consistent with both the framework's emergent-QFT-scale value and with ΛCDM, lying within the predictive bracket. A DESI Year 5 measurement that landed any-

where within the bracket — including values much larger than the emergent-channel bound, of either sign, provided the $w(z)$ shape remains Type II (no phantom crossing) — would not by itself falsify; the commitment is to the Type II reading, not to the specific numerical value within the bracket, though a positive detection would require the non-local substratum branch with its Lorentz-violation co-signature.

Cosmic initial conditions are solution-specific. The framework derives the structure of the dynamics — the wave equation, the bijection, the partition — from which evolution proceeds. The state at any particular cosmic time, including the time-zero conditions, is not a framework output. The framework is compatible with any cosmic-history that produces a C1–C4-satisfying observer-admitting regime by the time the cosmological horizon stabilizes. Specific cosmological initial-condition observations bear on the framework only at the boundary where they would or would not produce the observer-admitting regime; within that boundary they are solution-specific.

The choice among representatives related by the stated fixed-interface invariances is not determined by the data those invariances preserve. Uniform hidden relabelling and an independently prepared finite decoupled enlargement give exact examples [Substratum Theorem 24]. The same results do not classify every physical implementation or every observer partition. A measurement using a different interface or a sector that later couples to the observer lies outside the fixed-interface comparison, and requires its own analysis.

Regimes of deliberate silence. Beyond solution-specific properties, three regimes lie outside the framework's predictive scope by structural construction. These are regimes where the framework's foundational structure does not produce predictions, not because the framework is silent by choice but because the structural conditions for predictions to be defined do not hold.

The pre-horizon early universe. The framework's gravitational sector derivations in Chapter 7 operate on a stable causal partition with $\tau_B \sim H^{-1}$ — the cosmological horizon as a partition between observer-accessible and observer-inaccessible regions. The pre-horizon era — before such a partition stabilizes — has no observer in the framework's sense, because the C1–C4 conditions for embedded observation are not satisfied. The framework's only commitment in this regime is a boundary condition on deeper theories: any successful pre-horizon dynamics must produce a C1–C4-satisfying regime by the time a horizon stabilizes. The framework makes no prediction for the dynamics of the pre-horizon era itself; that domain belongs to whatever deeper theory describes the pre-horizon transition.

The equilibrium phase of (S, φ). Chapter 1's central theorem requires C1–C4 to be satisfied. At a substratum equilibrium that mixes fast compared with the observer's step ($\tau_{\text{mix}} \ll \tau_S$), and in the coarse-grained description generated by iterating the one-step effective channel, C2 fails in its timescale form, $\tau_S \ll \tau_B$ — its slow-bath route gone, though persistence of a residual record is not thereby

excluded — and the readback C4 requires is suppressed exponentially in τ_S/τ_B at every order $k \geq 2$, since a bath thermalizing between steps erodes the record before it can be read ([Main §2.3]): at such an equilibrium, the hidden sector has thermalized, and there are no slow modes to provide the timescale separation that sustains accessible memory. Under the effective-mixing hypothesis, with C2 read in that timescale form, and in the coarse-grained description generated by iterating the one-step effective channel, the structural observer-selection theorem developed at Main §4.6 restricts observers to the non-equilibrium phase of the substratum's dynamics. In that formulation a fast-mixing equilibrium phase is observer-free by structural argument, and the framework makes no prediction for what happens in this regime — there is no observer to whom a prediction could be addressed. This is the framework's account of why Boltzmann brains do not appear in observed physics: not because they are rare random fluctuations that happen not to occur, but because, in that formulation, the fast-mixing equilibrium phase that would produce them is structurally observer-free.

Regions beyond the observer's causal horizon. For any observer with a finite cosmological horizon, regions beyond that horizon are part of the observer's hidden sector. They are addressed in the framework only through boundary effects — the area-law entropy at the horizon surface, which encodes the gravitational signature of the hidden-sector content. The framework takes no observer-independent position on the *internal content* of these regions. Whether the universe beyond an observer's horizon contains fields configured differently, contains different gauge structures, or contains anything at all in the observer's conceptual vocabulary, is not a question the framework's structure can address. The framework's account is that the trace-out over what lies beyond the horizon produces the observer's quantum-mechanical description; the content traced over is, by construction, inaccessible to that observer.

These silences are structural rather than methodological. They follow from the characterization theorem's identification of accessible non-Markovian observation with embedded observation under C1, C3 and C4 (quantum for the source's finite-configuration class; the slow-bath form of C2's necessity for physical hidden sectors conditional on the mixing hypothesis), and from the observer-selection theorem's restriction of observers to the non-equilibrium phase (under its effective-mixing hypothesis, with C2 read in its timescale form (record persistence and readback there being suppressed, not excluded), for sufficiently fast-mixing equilibria ($\tau_{\text{mix}} \ll \tau_S$), and in the coarse-grained description generated by iterating the one-step effective channel [Main §4.6]). The framework's silence on these regimes is not a gap in its content but a feature of its structural commitments: it is committed to the regime where embedded observers exist and the regime is non-equilibrium; it is not committed to regimes where these conditions fail.

What this section is not. The articulation of what does not falsify is not a list of escape routes. The framework's structural commitments — the Class A theorems, the Class B retrodictions, the Class C broader commitments —

are pre-registered as falsifiable, and the author's commitment is to treat their violation as framework-falsifying rather than as an opportunity for post-hoc revision. The solution-specific properties and the regimes of deliberate silence are not categories into which falsifying observations can be reclassified after the fact. They are explicit acknowledgments of where the framework's commitments end and where the framework's structure has no predictive content. The line between them is drawn before the relevant observations are made, with the structural derivation determining which side of the line each putative prediction falls on.

4.7 What the framework derives, and what it compresses

A framework that advertises "the Standard Model, gravity, and quantum mechanics from a single object" invites a sharp question, and the methodological honesty of the preceding sections requires meeting it directly: is the derivation genuine, or is the answer quietly placed in the construction and then recovered? The answer is neither a simple yes nor a simple no, because the framework's results do not all stand in the same relation to its starting point. They fall into three tiers, and naming the tier is the honest form of every claim the book makes.

The first tier is **forced**. Once the substratum is committed to — a simple-cubic lattice carrying the wave equation — a great deal follows by pure representation theory, with no remaining freedom. The decomposition of the six lattice link-directions under the cubic point group fixes the multiplicity pattern $6 = T_1(3) \oplus E(2) \oplus A_1(1)$; read as the physical carrier under H-link and H-cust, it gives the gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$, and on that branch the same structure gives three candidate fermion generations (physical generations under H-spin'), the anomaly-free hypercharges (unique on the nonzero-Y_Q branch, up to normalization), and, under its identification hypothesis, the Cabibbo angle to four-significant-figure agreement. These could have come out otherwise — a different point group, a different count, a different angle — and the content of the derivation is that they do not. This tier is the framework's genuine achievement, and it is substantial: it is not reverse-engineering, because the predictions exceed the inputs, most visibly where the Cabibbo angle lands at $1/(\pi\sqrt{2})$ while the competing cubic lattices miss the measured value by tens of percent.

The second tier is **selected by the world**. Several quantities the derivation leans on are fixed not by the substratum but by observation. The spatial dimension is three because gravity propagates, matter is stable, and space is observed flat — three facts read off the universe, not consequences of the bijection. The lattice is *cubic* because we observe an SU(3) in nature and only the cubic crystal system can carry one. Here the reasoning runs backward, from the world to the commitment: "the Standard Model from a cubic lattice" and "the lattice is cubic because we see SU(3)" are the same fact spoken in two directions. The framework does not hide this — it states the cubic structure as a commitment forced by the observed gauge group — but the headline should carry it too,

because it marks the boundary of what is derived.

The third tier is **stipulated**. Some load-bearing moves are decisions about what is to count as physical structure rather than facts the bijection compels: that a "site" is *defined* by minimal coupling (from which the internal component count follows), that couplings are nearest-neighbor, and — at the foundation — that the limitation of an embedded observer takes the specific form of a fast sector reading and writing to a slow, capacious hidden one. These choices are principled and motivated, but they are choices.

Seeing the three tiers at once dissolves a question the framework had carried as open — whether its structural residue is "genuinely non-tautological." The question has no single answer because the residue is not one kind of thing: it is non-tautological at the first tier, world-selected at the second, definitional at the third. And no calculation settles it, because it was never an empirical question; the logical status of each step is fixed by the construction as written. What the classification yields is a precise description of what the framework *is*. It is a **maximally tight compression** of known physics onto a small set of commitments. The first tier shows that, once those commitments are granted, the discrete structure of the Standard Model is largely not a list of independent facts but the unfolding of a single representation-theoretic object — and *that* is a genuine and unusual claim about the unity of physics, worth the book whether or not the deeper pretension holds. The second and third tiers show that the commitments themselves are anchored in observation and partly stipulated, so the framework is not spinning the world out of a parameter-free principle.

This is also the place to be exact about the foundation that names the book. "Observation is incomplete" — that what we know is reached through a partial vantage, never the whole — is, taken by itself, not a falsifiable claim about physics. It is close to the definition of the empirical stance itself: every theory in science is a description built from inside, from limited access. A thesis that cannot be false cannot, on its own, predict. All of the framework's predictive content therefore lives one level down, in the *specific model* of the incompleteness — the architecture, the lattice, the dynamics — which is precisely where the second and third tiers locate every input. The framework's reach and its circularity are one structure seen from two sides. Read as the *source* of the predictions, "observation is incomplete" claims too much. Read as the *organizing principle* under which an unusually large body of physics compresses to one structure, it is the defensible and still-remarkable claim, and it is the claim this book makes. The reader should hold the framework to that standard and no other: not as a derivation of the universe from the bare fact that there are observers, but as a demonstration that an enormous amount of what we observe coheres as the shadow of one finite construction — with the construction's own starting points held up honestly to the light.

4.8 What follows

The methodological discipline developed in this chapter — the single-foundational-commitment pattern, the claim classification by derivational rigor, the architecture-level engagement with the no-go results, verified theorem by theorem, the pre-registered falsification conditions and their complement of deliberate silences — provides the framework with a discipline that organizes its program as a research program rather than as a single bold conjecture. The discipline is part of the framework's content, not separate from it: the structural commitments developed in Chapters 1 through 3 and the empirical predictions developed in Parts II through IV are produced under this discipline, with the discipline determining what the predictions are, how they are classified, and what their falsification status is.

Parts II and III apply the methodology to the framework's specific empirical content in fundamental physics. Chapters 5 and 6 develop the Standard Model gauge and matter sectors, with twenty-two classified observable entries, including two structural and five conditional structural entries, in the inventory of [SM §7.6]. The four-layer framing locates each retrodiction in the substratum-to-emergent stack, with Layer 0 and Layer 1 predictions (gauge group, generation count, Cabibbo, Koide, Wolfenstein) carrying the framework's strongest structural commitments and Layer 2 and Layer 3 predictions (mass ratios within specific operator structures, scale-setting predictions) carrying more solution-specific content. Chapter 7 develops the gravitational sector: the Bekenstein-Hawking 1/4 coefficient and the calibration $\hbar = c^3\epsilon^2/(4G\sigma)$ from horizon thermodynamics are consistency relations conditional on the named horizon, frame, state and calibration assumptions, while the vacuum-energy level distinction is descriptive and supplies no dark-energy dynamics. The individual classifications and empirical limits appear in Appendix A §A.4.2. Chapter 8 develops the JUNO neutrino prediction as an L-class prediction matched at 0.07σ, with the closure of the lepton-sector mixing-matrix derivation as the open direction whose completion would move the prediction to S-class. Chapter 9 develops the universality-class structure and the framework's relationship to alternative unification programs, with the architecture-level no-go engagement from §4.4 applied to the analysis of cross-framework structural relationships.

Parts III and IV develop the framework's reach into structural cascades and applied scientific fields, with the methodological classifications carrying through to the more applied content. The framework's predictions in chemistry, biology, neuroscience, and quantum engineering are not all S-class — many are P-class or E-class, with empirical inputs from the relevant scientific fields entering the derivation chain. The classification scheme developed here makes the rigor of each prediction explicit, so that readers approaching the framework from biology or medicine can evaluate the framework's claims against the appropriate empirical standards in those fields.

The methodology developed in this chapter is not unique to the framework.

The single-foundational-commitment pattern is a general principle of theory construction, the claim-classification scheme is a general standard for empirical evaluation, the architecture-level no-go engagement is a general method for analyzing structural results, and the pre-registered falsification discipline is a general epistemological standard. The framework's contribution is to apply this methodology systematically rather than selectively, with every prediction carrying its classification and every structural commitment carrying its falsification condition. What this discipline produces, when applied to the framework's foundational starting point — the two-axiom observation base — is the content developed in the remaining chapters of the book.

The methodological posture sketched in this chapter is developed at length, for a philosophy-of-physics readership, in a companion paper — *Physics Modulo Gauge* — which treats the two-axiom foundational commitment, constructive structural realism, the separate premise checks of the no-go results of quantum foundations within the common substratum and observable descriptions, and "physics modulo gauge" as a general methodology. That paper is a companion to the framework's technical sequence rather than part of this book; readers concerned specifically with the framework's methodology and its relation to the foundations-of-physics literature will find it the natural next text.

Chapter 5

The Emergence of Gauge Structure

Observer geometry and physical carrier. SM §4.1 distinguishes two claims. First, finite equivariant projection is classical and exact: cubic symmetry is inherited by the projected Markov part and every memory kernel, forbidding quadratic anisotropy in the scalar symbol — so whatever nondegenerate quadratic structure survives is isotropic — while allowing cubic anisotropy at quartic order. The particular normalized nearest-neighbor operator $A/(2d)$ additionally requires an **observer-level** center-free kernel; microscopic center independence alone does not imply that after correlated hidden variables are marginalized. Second, the six signed simple-cubic links and their exact $3 \oplus 2 \oplus 1$ representation are unconditional geometry, but identifying that link space with the complete physical carrier is **H-link**; the single-copy clause gives $K = 6$, and the custodial condensate-stabilizer reading is **H-cust**. Accordingly, any downstream statement here that consumes $K = 6$, $N_f = 6$, the physical SM gauge carrier, or a fixed three-sector count inherits those named bridge conditions unless it is explicitly referring only to the six geometric links.

5.1 What this chapter develops

Part I established that any observer embedded in a finite deterministic universe under conditions C1–C4 must use a memory-bearing formalism — one admitting a fixed-basis quantum representation, as every finite observable law does, with the compressed form for processes in the correspondence's class under (T); that observed physics together with structural assumptions, under the reconstruction theorem's stated conditions and its nearest-neighbor premise, fixes the **local propagating lattice/gauge residue** while Bell-inclusive completion remains underdetermined; and that the framework's structural-realism position operates on a two-dimensional hierarchy with stratified empirical commitments. The remaining question is what specific physics the substratum produces. This chapter and the next answer that question for the visible sector — the Standard Model of particle physics.

The derivation has two natural phases, and the book treats each in its own chapter. The present chapter separates an exact geometric core from the conditional physical-carrier reading: finite equivariant projection gives an Euclidean quadratic observer symbol; the six signed cubic links decompose exactly with multiplicities $(3, 2, 1)$; **under H-link single-copy + H-cust**, the condensate-stabilizer/background-independence construction supplies the proposed Standard-Model gauge carrier. Chirality and hypercharge statements retain their additional hypotheses where stated. The second phase, Chapter 6, establishes *what specific quantitative content* this gauge structure produces: three chiral fermion generations, one Higgs doublet $(\mathbf{1}, \mathbf{2}, +1/2)$ taken as input and accommodated in the E projection of the singlet taste (SM Theorem 11), gauge couplings, and the twenty-two classified retrodictions (two parameter-free structural and five conditional structural) including the Cabibbo angle, the Koide relation, fermion masses, and PMNS mixing angles.

This chapter's content is structured. Section 5.2 establishes the minimal mathematical object — a finite set with a deterministic bijection of bounded coupling degree, with the coupling graph read off from the bijection once a product factorization of the state space is given; the factorization itself is an input the dynamics has not been shown to select. Section 5.3 selects $d = 3$ within A4's assumed periodic-lattice class under the continuum conditions: propagating gravity and stable matter give the two bounds, and boundary-entropy concordance is a further consistency check. Section 5.4 develops the observer-level lattice-wave branch and its staggered factorization, then states H-link: the physical carrier is a copy of the six signed links, possibly with multiplicity, with $K = 6$ only on the single-copy branch. Section 5.5 derives the exact six-link $(3, 2, 1)$ decomposition and, under H-link + H-cust plus the stated gauge premises, the corresponding condensate-stabilizer/local-gauge construction. Section 5.6 develops chirality from the trace-out and anomaly cancellation forcing the observed hypercharges on the nonzero-Y_Q branch. Section 5.7 closes the chapter with a forward pointer to Chapter 6.

The structural status of the chapter's results is now layered. The exact finite-observer isotropy theorem and the six-link $(3, 2, 1)$ representation are forced by their stated finite/symmetry hypotheses. The **physical Standard-Model carrier is not yet forced**: it additionally requires H-link single-copy and H-cust, followed by the amplitude-scale/background-independence premises already recorded. That realization is itself empirically anchored (the lattice is selected to match the observed gauge group, not derived from the substratum alone — §4.7), so the uniqueness is conditional on it rather than derived from a parameter-free first principle. The methodological discipline developed in Chapter 4 will organize the predictions of Chapter 6 by derivational rigor. The exact observer-isotropy theorem and six-link multiplicities sit at the structural tier; the physical gauge identification sits at a conditional bridge tier because it consumes H-link/H-cust, with the later reduction to $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ carrying amplitude-scale gauge and a specified charged matter action.

The chapter takes as input the framework's prior content. The quantum-mechanical emergence theorem of Chapter 1 is invoked where needed but not reproved. The reconstruction theorem of Chapter 2 provides the structural assumptions A1–A6 and the stated modeling/completeness conditions and nearest-neighbor premise under which the **local lattice/gauge residue** is uniquely fixed. The Bell-inclusive completion is not uniquely determined by that theorem. The hierarchy framework of Chapter 3 locates each prediction in the multi-level structural-realism scheme. The methodological discipline of Chapter 4 classifies each prediction by its derivational rigor. The chapter's content is the substantive derivational chain from the cubic-lattice substratum to the Standard Model gauge group — the framework's specific structural claim about *why* physics has the gauge symmetries it has.

5.2 The minimal substratum and the wave equation

The framework's content rests on a minimal mathematical structure: a finite set S, a deterministic bijection $\varphi : S \to S$, and a partition of S into visible and hidden sectors. The lemmas of Chapter 1 extracted three structural features from this minimal object — visible finiteness, causal partition, determinism — and the conditions C1–C4 identified when an embedded observer can exist. This section develops two further structural features of the minimal object that the Standard Model derivation requires.

The factorization problem. Let S be a finite set with $|S| = q^N$ for some $q \geq 2$ and integer N. There are many ways to write S as a Cartesian product $S = S_1 \times S_2 \times \cdots \times S_N$ with $|S_k| = q$ for each k. Different factorizations correspond to different choices of which elementary degrees of freedom to identify as "local" — different bases in which to decompose the configuration space. The factorization is not given by the set S itself; it must be determined by some additional structure.

Given a bijection φ and a product factorization, define the *coupling graph* G_φ

on vertices $\{1, 2, \ldots, N\}$, with vertices i and j connected by an edge if the i-th component of $\varphi(s)$ depends on the j-th component of s for at least one state $s \in S$. The coupling graph is not an additional structure imposed on the system; once the factorization is given, it is read off from the bijection. The factorization itself is an input — site factorization, in the posit ledger of [Main §4.5] — and the dynamics alone has not been shown to select it.

Minimizing the coupling degree — the number of edges per vertex — over all product decompositions does not select it. The wave equation in one dimension is

$$x_i(t+1) = x_{i-1}(t) + x_{i+1}(t) - x_i(t-1) \mod q,$$

On a ring of four sites over $\mathbb{Z}/3\mathbb{Z}$ the ring's adjacency matrix diagonalizes, and in its eigenbasis each normal mode evolves alone, $c_k(t+1) = \lambda_k c_k(t) - c_k(t-1)$: the coupling graph of the normal-mode factorization has maximum degree 1, against 3 for the spatial one. And composing a factorization with φ gives one with the same coupling graph, which differs from it whenever some component of φ depends on two components; a minimizer of that kind is therefore not unique. Whether the observable law, the observer's registers or the dynamics select the spatial factorization within a stated class is open.

The earlier extension from spectral dimension to a canonical $2d_s$ **physical internal carrier** is not established. Quasi-isometry and heat-kernel scaling constrain large-scale geometry but do not manufacture signed internal link channels. H-link and the exact $(3, 2, 1)$ carrier construction are simple-cubic-branch statements, computed in the spatial site factorization with the nearest-neighbour coupling graph.

Space as coupling structure. Every "spatial" property used in the derivation chain is a property of the coupling graph G_φ. The *graph boundary* $|\partial V|$ of a region $V \subset S$ — the quantity the area-law bound is stated in — is the number of edges crossing from V to its complement. The *dimension* is extracted from the causal partial order via the Myrheim-Meyer estimator. The *area-law bound* — that the information shared across a region's boundary is at most proportional to the boundary rather than the volume — follows from the spatial Markov property on any graph with range-1 dynamics. The *dispersion relation* describing how perturbations propagate requires translation invariance of the underlying dynamics. The regular cubic lattice $\mathbb{Z}^3$ is one graph with all these properties; it is not the only one.

The framework's account of space dissolves the container problem that has been a recurring objection to discrete models of spacetime. The coupling graph is not embedded in a pre-existing space. There is no ambient manifold in which the graph "lives." The vertices are labels for degrees of freedom, the edges record which degrees of freedom are dynamically coupled. Any spatial embedding is a representation for human convenience, not a physical fact. The coupling graph is not made of material — the vertices are not atoms of space and the edges are not physical connections. A degree of freedom is a component of the state,

an abstract mathematical entity rather than an object. Asking what a site is "made of" is a category error: it is a label for a variable, not a name for a thing.

The relationship between locality and the dynamics is the precise inverse of the conventional picture. In the conventional picture, space exists first as a manifold, and locality is the property that nearby points in space have causally connected dynamics. In the framework, the dynamics exists first as a bijection, and space emerges as the topology that locality induces on the coupling graph. Two sites are neighbors because the bijection couples them — because the state at one affects the future of the other. This is a dynamical fact, not a spatial relationship. The topology is defined by the interactions; the apparent spatial arrangement is the topology made visible.

The alphabet size. The alphabet size q is the number of distinguishable states at each lattice site. Several results hold for every $q \geq 2$ and contain no q: the P-indivisibility theorem, center independence, the boundary scaling of the area law, and the dispersion relation of the observer-level operator. The gravitational sector depends on q only through the entropy σ assigned to one boundary cell, which cancels from the Bekenstein-Hawking formula; at $\sigma = 1$, one nat per boundary mode, neither the gap equation $\hbar = c^3\epsilon^2/(4G\sigma)$ derived in Chapter 7 nor the Bekenstein-Hawking formula contains q, while $\sigma = \ln q$ would place q in the discreteness scale $\epsilon = 2\sqrt{\ln q}\, l_p$. Other results carry q: the period of the wave map on a ring of L sites is qL for prime q coprime to L, the coefficient of the area law and the capacity count of the hidden sector contain $\log q$, and the substratum normalisation $\alpha = 1/d$ requires $3 \nmid q$.

That the emergent observables do not depend on q at all — that two substrata (S, φ) and (S', φ') with the same coupling structure and different alphabet sizes produce the same emergent transition probabilities — is the *alphabet-size hypothesis*. It is not proved for every observable, and because the visible transition matrices of the two substrata act on configuration spaces of different sizes, stating it requires a correspondence between them that is not constructed. Under the hypothesis the mod-q state space at each lattice site plays the role of the gauge potential A_μ and the coupling structure with its emergent consequences that of the field strength $F_{\mu\nu}$. Generator (ii) of the substratum gauge group $\mathcal{G}_{\text{sub}}$ developed in Chapter 2 §2.5 is stated under this hypothesis.

A second gauge freedom — *amplitude-scale invariance* — has consequences that recur in the gauge-derivation chain. Stated operationally, it is the principle that an embedded observer has no access to the absolute scale of the field value ϕ, only to the emergent transition probabilities and coupling structure — the discrete analogue of the field-normalization redundancy of ordinary field theory, where rescaling a field is unphysical because only correlators are observable. It is realized as the additive (affine) relabellings $\phi \mapsto \lambda\phi$ of the alphabet being gauge, and is one instance of the framework's master principle that observationally indistinguishable transformations are gauge — the same principle behind the four $\mathcal{G}_{\text{sub}}$ generators and the alphabet-*size* hypothesis, from which it is nonetheless *distinct*. It is an explicit adjoined operational principle of $\mathcal{G}_{\text{sub}}$ (com-

panion papers, [Substratum §4, Theorem 24 Remark]); results that invoke it are conditional on it. The reduction of the natural commutant $\mathrm{U}(3) \times \mathrm{U}(2) \times \mathrm{U}(1)$ to the Standard Model's $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ uses it: the common overall $\mathrm{U}(1)$ phase can be absorbed into a global rescaling of the field value, which is gauge by amplitude-scale invariance; the hypercharge direction among the remaining abelian phases is selected by a specified charged matter action and anomaly cancellation (§5.6).

5.3 Selection of $d = 3$

The framework's structural lemmas operate at any integer dimension $d \geq 1$. The observer-level quadratic-isotropy argument and dimension-dependent lattice calculations can be formulated in other dimensions, but the six-link physical-carrier construction is not a dimension-general theorem: H-link is presently tied to the simple-cubic signed-link representation used at $d = 3$. The question of why our universe has three spatial dimensions is therefore a question about which dimension produces a self-consistent observed physics. The framework's answer is that exactly three works.

Within A4's periodic-lattice class and the continuum conditions of SM §3.1–§3.2, propagating gravity and stable matter select $d = 3$; boundary-entropy concordance is an additional consistency check. This is a conditional self-consistency selection using the stated physical inputs.

Filter 1: Propagating gravity requires $d \geq 3$. The gravitational sector developed in Chapter 7 requires propagating gravitational degrees of freedom — gravitational waves, the structural content of the LIGO observations beginning in 2015. The Weyl tensor, which captures the propagating part of the gravitational field, vanishes identically in $d \leq 2$: there is no gravitational radiation in two or fewer spatial dimensions. Jacobson's thermodynamic derivation of Einstein's equations from local causal horizons requires propagating gravity. The filter therefore excludes $d \leq 2$.

Filter 2: Stable matter requires $d \leq 3$. The Coulomb potential in d spatial dimensions falls off as $1/r^{d-2}$ for $d \geq 3$ and gives a logarithmic potential for $d = 2$. In $d \geq 4$ the Coulomb potential gives unstable atoms — there are no normalizable bound states of the d-dimensional Coulomb Hamiltonian, by Ehrenfest's argument and its quantum-mechanical refinement. Gravitational orbits are similarly unstable for $d \geq 4$ by Tangherlini's analysis. Without stable matter, no embedded observers exist to define the framework's partition. The filter therefore excludes $d \geq 4$.

The intersection of Filter 1 and Filter 2 is already $d = 3$ uniquely. A third filter provides a consistency check.

Filter 3: Boundary-entropy concordance gives $d = 3$ exactly. The cosmological observation that the universe is spatially flat near the critical density — $\Omega_{\text{total}} \approx 1$ — is, in the framework's gravitational sector developed in Chapter

7, a structural feature rather than an initial condition. The ratio $\rho_s/\rho_{\text{crit}}$ between the boundary-derived entropy density and the critical density, computed in d spatial dimensions, satisfies

$$\frac{\rho_s}{\rho_{\text{crit}}} = \frac{2}{d-1}.$$

This ratio equals unity exactly at $d = 3$. At $d = 2$ the ratio is 2 (overclosure); at $d = 4$ the ratio is $2/3$ (deficit, requiring an unsourced gravitational contribution). The filter selects $d = 3$ as the unique value at which the observed spatial flatness is structurally explained rather than fine-tuned.

Within A4's assumed periodic-lattice class and the stated continuum conditions, the first two bounds select $d = 3$ and boundary-entropy concordance supplies a further consistency check. The specified Yang–Mills coupling is dimensionless at $3+1$ spacetime dimensions; its renormalizability result is downstream of that selection. The six-link octahedral carrier and H-link used here are stated for the selected simple-cubic branch. A comparison with another dimension requires its own carrier and matter assumptions.

Integer dimension. The self-consistency argument assumes that the coupling graph has a well-defined integer dimension. A general bounded-degree graph need not have one: besides exponential growth ($|B(r)| \sim k^r$ for some $k > 1$), fractal growth ($|B(r)| \sim r^\alpha$ for some non-integer α) and integer-polynomial growth ($|B(r)| \sim r^d$ for some integer d), there are graphs of intermediate growth (Grigorchuk) and graphs with no single growth exponent. Under A4 (Chapter 2) the integer is an input: the coupling graph is a finite periodic lattice, the quotient of a rank-d lattice by a finite-index sublattice, so $|B(r)| \sim r^d$ below the system size and none of the other classes arises. Independently, self-consistency excludes the first two classes.

Exponential growth fails because the wave equation on exponentially-growing graphs does not produce Lorentz-invariant dispersion at low energies — the spectral gap does not close in the correct functional form — so Jacobson's derivation of Einstein's equations fails. Additionally, the effective dimension $d_{\text{eff}} \to \infty$ is incompatible with the stable-matter filter.

Fractal growth fails because Jacobson's argument requires the Raychaudhuri equation, which requires geodesic congruences on a manifold structure that fractal spaces do not support. The staggered fermion decomposition that produces the framework's three-generation structure requires a hypercubic Brillouin-zone structure $\eta \in \{0, 1\}^d$, which has no fractal-graph analog. The cubic-group decomposition of $2d$ link directions, which constrains the gauge structure, requires cubic lattice symmetry — absent in fractals.

The integer d therefore enters through A4 as the rank of the translation lattice, not through Gromov's theorem, which concerns finitely generated groups and their Cayley graphs; statistical isotropy alone does not supply it. Applied to

the continuum description, conditional on the continuum step of §3.1 of the SM paper, the three filters then select $d = 3$.

Corollary. *Any bijection whose coupling graph is a d-dimensional periodic lattice with $d \neq 3$ fails at least one of the filters E5–E7 in its emergent description. The coupling-graph dimension is not a free parameter or a property of a specific φ; it is the unique value compatible with the filters.*

The argument's structure is worth marking explicitly. The framework's definition of observation requires an embedded observer as a constitutive element — the partition $V \subset S$ is defined relative to an observer that the substratum's structure must support. The requirement that such an observer exist, combined with the framework-internal concordance calculation, constrains d. This is not a standard anthropic argument selecting from a pre-existing multiverse of dimensions. The observer is a structural prerequisite for the framework's definition to have content, not a contingent feature of one universe among many.

Within the assumed A4 periodic-lattice class and the selected continuum description, the E5–E7 filters select $d = 3$. Integer rank is supplied by A4, while the value 3 is selected using the physical filters. The continuum-curvature identification remains open, as stated in SM §3.1. This selection concerns that branch rather than every finite bijection or embedded-observer architecture.

5.4 The emergent QFT: factorization and multi-component dynamics

The cubic branch supports an observer-level lattice-wave description whose relation to the finite bijection is governed by the projection bridge of SM §4.1. This section develops its staggered factorization and then separates the exact six-link geometry from H-link, the open identification of those links with the physical internal carrier.

Observer-level lattice Klein–Gordon branch. The exact projection theorem does not require a wave equation. If the projected scalar Markov part is additionally nearest-neighbor, constant-preserving and center-free, cubic symmetry fixes it to $A/(2d)$, and the reversible second-order branch is

$$\phi(\mathbf{n}, t+1) = \frac{1}{d} \sum_{j=1}^{d} [\phi(\mathbf{n} + \hat{e}_j, t) + \phi(\mathbf{n} - \hat{e}_j, t)] - \phi(\mathbf{n}, t-1).$$

Its low-momentum symbol is Euclidean, $\omega^2 = (a^2/d)|\mathbf{k}|^2 + O(a^4 k^4)$. This is an **observer-level conditional branch**, not the fundamental finite bijection itself. A $\beta = 1$ "maximum-speed" reading is unavailable: causal support travels at most one graph edge per update independently of the coefficient, and the unnormalised $d = 3$ real lift is unstable at $k = 0$.

Factorization into staggered Dirac fermions. A theorem of Susskind from 1977 establishes that the square of the staggered Dirac operator on a hypercubic lattice equals $-\frac{1}{4}$ of the lattice d'Alembertian: $D_{\text{st}}^2 = -\frac{1}{4}\Box_{\text{lat}}$. The wave equation $\Box_{\text{lat}}\phi = 0$ is therefore equivalent to the massless staggered Dirac equation $D_{\text{st}}\chi = 0$ under an exact change of variables on the lattice. The bosonic field ϕ and the fermionic field χ are not two different objects; they are the same lattice content viewed through different decompositions.

The implication is non-trivial. The wave equation as substratum dynamics looks like a single scalar field evolving on the lattice, but its Susskind factorization reveals that the underlying degrees of freedom can equally be described as fermionic. The framework's matter content — the fermions of the Standard Model — is not added to the substratum as a separate input. It is implicit in the wave equation itself, made explicit through the Susskind change of variables. The structural content of the Standard Model's three generations (physical reading conditional on H-spin', SM §4.7; chirality on H-χ', SM §4.8), developed in Chapter 6, traces back through the Susskind decomposition to the wave equation's specific algebraic form.

Center independence, beyond setting the mass to zero in the bosonic description, has a corresponding structural meaning in the fermionic description. For site-independent center coefficients added to opposite-parity hopping, in a ring without additive 2-torsion, on a lattice carrying both parity sectors, the exact chiral symmetry of the emergent staggered fermions is equivalent to the vanishing of those center coefficients (SM Theorem 3): any non-zero center coefficient breaks chiral symmetry. Identifying these first-order terms with center dependence of the second-order substratum dynamics is a separate bridge (SM §4.3). The memory-bearing sector whose phenomenology Chapter 1 develops does not come from center independence: Main's readback lemma supplies that memory, and its recurrence-cycle theorem the P-indivisibility, independently of center independence (SM §4.3). The Higgs mechanism developed in Chapter 6, which gives masses to fermions in the emergent description, operates on the chiral fermions of the center-free branch.

Multi-component dynamics with K internal components. The wave equation as written above is a single-component dynamics — one scalar field per lattice site. The framework's gauge-emergence chain requires the generalization to K components per site: $\phi(\mathbf{n}, t) \in (\mathbb{Z}/q\mathbb{Z})^K$, with the matrix wave equation

$$\phi(\mathbf{n}, t+1) = M \sum_{j=1}^{d} \left[\phi(\mathbf{n} + \hat{e}_j, t) + \phi(\mathbf{n} - \hat{e}_j, t)\right] - \phi(\mathbf{n}, t-1)$$

where M is a $K \times K$ coupling matrix. Spatial isotropy forces M to be the same matrix for all d link directions; reversibility forces the previous-step coefficient to be $-I_K$; center independence forces the absence of a $C\,\phi(\mathbf{n}, t)$ term. The matrix M is the sole free parameter of this branch at this level.

The coupling matrix is scalar, $M = \mu I_K$ with $\mu = 1$: every substratum mode

has the dispersion relation $\cos\omega = \cos k$, so all substratum modes share one massless cone, and mass and gauge-group distinctions are carried by the solution's equivariant condensate Σ (SM Theorem 5). The global gauge group is the stabilizer of the solution's equivariant condensate Σ in $\mathrm{U}(K)$ — the set of $K \times K$ unitaries commuting with Σ (the coupling matrix itself is scalar; the dynamics carries an exact custodial $\mathrm{U}(6)$). By Schur's lemma, this stabilizer decomposes as a product of unitary groups, one on each eigenspace of Σ: if Σ has r distinct eigenvalues with multiplicities $(n_1, \ldots, n_r)$, then the gauge group is $\mathrm{U}(n_1) \times \cdots \times \mathrm{U}(n_r)$. The gauge group and the mass spectrum are encoded in the same matrix Σ.

The six-link carrier and H-link. The simple-cubic graph has exactly six signed nearest-neighbor link directions. Their span V_{link} is therefore an exact six-dimensional geometric representation. Coupling-degree minimization does **not** show that diagonalizing the isotropic internal matrix assigns one field component to one signed link; the full neighbor sum remains in every diagonal component. A $\delta(2d) = 1$ Step 1 is therefore unavailable.

H-link states that the observer/gauge carrier is $V_{\mathrm{link}} \otimes \mathbb{C}^m$. The single-copy clause $m = 1$ gives $K = 6$, but no present theorem derives $m = 1$. The later gauge and matter calculations that use $K = 6$ are thus conditional tests of this branch. The exact group-theoretic statement below — the decomposition of the six **geometric** links — does not depend on H-link.

5.5 The cubic decomposition and the Standard Model gauge group

The six signed simple-cubic links, defined in the spatial site factorization with the nearest-neighbour coupling graph ([Main §4.5], items (viii) and (x)), form an exact six-dimensional representation of the cubic group. This section first computes that representation unconditionally. Its use as the **physical** internal/gauge carrier then assumes H-link single-copy; the custodial condensate-stabilizer step additionally assumes H-cust.

The cubic-group representation on the six links. The six nearest-neighbor link directions $\{\pm\hat{e}_1, \pm\hat{e}_2, \pm\hat{e}_3\}$ of the cubic lattice, with one component per link direction. The cubic rotation group O — the group of orientation-preserving symmetries of the cube, with twenty-four elements — acts on these six link vectors by permutation: rotations of the cube permute the six link directions among themselves, with rotations through a face center cycling four link directions and rotations through a vertex cycling three.

The action of O on the six-dimensional space spanned by the link directions is a representation of O with character $\chi(O) = 6, 0, 2, 0, 2$ at the five conjugacy classes of O (identity, eight three-fold rotations, three two-fold rotations through face centers, six two-fold rotations through edge midpoints, six four-fold rotations through face centers). Computing the inner products of χ with the characters of the five irreducible representations of O — labeled A_1, A_2, E, T_1, T_2

with dimensions $1, 1, 2, 3, 3$ — yields the decomposition

$$6 = T_1 \oplus E \oplus A_1$$

with multiplicities $(3, 2, 1)$. The two remaining irreducible representations A_2 and T_2 have multiplicity zero in the six-dimensional link representation.

The decomposition is forced by group-theoretic content rather than chosen. The six link directions form a specific representation of the cubic rotation group; the decomposition of this representation into irreducibles is computed mechanically from the character table. The framework's commitment to the cubic lattice as substratum, combined with the requirement of spatial isotropy, makes the $(3, 2, 1)$ multiplicity structure inevitable.

Conditional condensate-stabilizer reading. The link representation itself is $T_1 \oplus E \oplus A_1$. Under **H-link single-copy + H-cust**, the observer-level physical carrier is this six-dimensional space and the pre-condensate kinetic structure is custodially scalar (at the symmetric point the kinetic clause follows from the signed-permutation symmetry of the components under H-blind, [SM §4.4]). An O-equivariant condensate then has block form

$$\Sigma = aP_{T_1} + bP_E + cP_{A_1},$$

and for generic distinct block values its unitary stabilizer is

$$\mathrm{U}(3) \times \mathrm{U}(2) \times \mathrm{U}(1).$$

This stabilizer theorem is exact at those stated premises. It does not prove H-link, select $m = 1$, or by itself perform the later $\mathrm{U} \to \mathrm{SU}$ reduction.

Reduction to the Standard Model gauge group. The natural commutant $\mathrm{U}(3) \times \mathrm{U}(2) \times \mathrm{U}(1)$ is not quite the Standard Model gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$. At the level of Lie algebras each factor splits as $\mathfrak{u}(n) = \mathfrak{su}(n) \oplus \mathfrak{u}(1)$, so the commutant carries $\mathfrak{su}(3) \oplus \mathfrak{su}(2)$ and three abelian phase directions (globally $\mathrm{U}(n) \cong (\mathrm{SU}(n) \times \mathrm{U}(1))/\mathbb{Z}_n$). The common overall phase can be absorbed into a global rescaling of the field value, which is gauge by amplitude-scale invariance of the substratum gauge group $\mathcal{G}_{\mathrm{sub}}$ (§5.2); selecting the hypercharge direction among the remaining abelian phases requires a specified charged matter action and anomaly cancellation (§5.6).

After this reduction, the gauge group is

$$\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)_Y.$$

On the H-link/H-cust branch, this is the proposed Standard Model gauge group after the separate amplitude-scale/background-independence premises and the charged matter action are applied. The exact unconditional result from the cubic lattice at this point is the $3 \oplus 2 \oplus 1$ link representation. The three factors have physical identifications matching the Standard Model: T_1 (the three-dimensional spatial-vector representation) is the color SU(3) of quantum chromodynamics; E (the two-dimensional quadrupole representation) is the weak

isospin SU(2) of the electroweak theory; A_1 (the one-dimensional scalar representation) is the hypercharge $\mathrm{U}(1)_Y$.

Bravais-lattice uniqueness. A subtle point deserves explicit treatment: among the fourteen three-dimensional Bravais lattices, the simple cubic lattice is the unique one compatible with the Standard Model gauge group under the commutant construction on the nearest-neighbor link representation. The body-centered cubic lattice has eight nearest-neighbor link directions decomposing as $A_1 \oplus A_2 \oplus T_1 \oplus T_2$ under O, giving a commutant of $\mathrm{SU}(3)^2 \times \mathrm{U}(1)^2$ — two color factors and no weak factor. The face-centered cubic lattice has twelve nearest-neighbor link directions decomposing as $A_1 \oplus E \oplus T_1 \oplus 2T_2$, giving the factor structure $\mathrm{SU}(3)^3 \times \mathrm{SU}(2)$ with extra U(1) factors (SM Theorem 7b) — incompatible with the Standard Model gauge group regardless of the precise factor count. The remaining eleven non-cubic Bravais lattices belong to crystal systems whose point groups have only one- and two-dimensional irreducible representations, supporting no three-dimensional irrep — no SU(3) factor available under the commutant construction. The tightest form of the argument runs through the SU(2) rather than the SU(3): the multiplicity ladder $(3, 2, 1)$ is the irreducible-degree set $\{1, 2, 3\}$, and the octahedral group is the unique finite rotation group carrying both a genuine 2- and a 3-dimensional irrep (Appendix B.6b).

Simple cubic is therefore the unique Bravais-lattice substratum compatible with the observed Standard Model gauge group. The framework's commitment to simple cubic, rather than to a more general bounded-degree graph satisfying the structural conditions of Chapter 1, is forced by the empirical fact that the gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ is observed. The further empirical content of the Cabibbo angle being $1/(\pi\sqrt{2})$ to 0.04%, developed in Chapter 6, provides quantitative confirmation: simple cubic produces the observed value, while body-centered and face-centered cubic produce values discrepant from observation by 42% and 29% respectively.

The formal uniqueness theorem, including the character-table decomposition for all 32 three-dimensional crystallographic point groups and the proof that only the five cubic-system point groups admit the required 3D irreducible representations, is developed in Appendix B.6b.

Local gauge invariance from background independence. The group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ derived so far is a global symmetry of the matrix wave equation: the same gauge transformation applies at every lattice site. The Standard Model is, of course, a local gauge theory — the gauge transformations vary from site to site. The step from global to local is what background independence (A6) supplies: the local transformations act covariantly once the coupling data on the links are transformed with them.

Background independence, as the reconstruction's assumption A6 states it, is a covariance requirement: a site-dependent internal-index transformation is a symmetry of the dynamics provided the coupling data on the links are transformed

with it. It is not the requirement that the coupling structure be held fixed and left invariant — for an invertible coupling that stronger fixed-background condition holds only for transformations that agree across every coupled pair of sites — and it is not the state-dependent coupling graph $G(x)$ on which the Bell branch of Chapter 1 rests, under which the graph itself evolves with the state; those are different principles that share a name. Under H-link and H-cust (§§5.4–5.5), the cubic decomposition of the link directions fixes the block multiplicities $(3, 2, 1)$ at each site; spatial isotropy fixes those multiplicities, not distinct block values. The coupling on each link, $M(\mathbf{n}, \hat{e}_j)$, is the data the site-dependent transformations act on; where the equivariant condensate takes distinct block values, its stabilizer $\mathrm{U}(3) \times \mathrm{U}(2) \times \mathrm{U}(1)$ (SM Theorem 5) acts at each site, and the resulting gauge transformations are site-dependent.

Under a site-dependent transformation $G(\mathbf{n})$ of the form $\phi(\mathbf{n}) \to G(\mathbf{n})\,\phi(\mathbf{n})$ with $G(\mathbf{n}) \in \mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ at each site, the matrix wave equation is invariant provided the coupling matrix transforms as $M(\mathbf{n}, \hat{e}_j) \to G(\mathbf{n})\, M(\mathbf{n}, \hat{e}_j)\, G(\mathbf{n} + \hat{e}_j)^{-1}$. The link variable $M(\mathbf{n}, \hat{e}_j)$ transforms as a gauge connection, with the plaquette product $P(\mathbf{n}, j, k) = M(\mathbf{n}, \hat{e}_j)\, M(\mathbf{n}+\hat{e}_j, \hat{e}_k)\, M(\mathbf{n}+\hat{e}_k, \hat{e}_j)^{-1}\, M(\mathbf{n}, \hat{e}_k)^{-1}$ transforming in the adjoint as $P \to G(\mathbf{n})\, P\, G(\mathbf{n})^{-1}$. The real part of the trace of the plaquette is gauge-invariant — recognizable as the Wilson plaquette action of lattice gauge theory.

The framework's derivation of local $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ gauge invariance, on the H-link/H-cust branch, is therefore complete at the level of the symmetry. The gauge group is fixed by the cubic decomposition of the six link directions; amplitude-scale invariance, a specified charged matter action and anomaly cancellation (§5.6) reduce the three abelian phase directions to the hypercharge; background independence promotes the global commutant to a local gauge symmetry; the gauge-invariant plaquette functional from which the Wilson action is built follows, with the link variables $M(\mathbf{n}, \hat{e}_j)$ as the gauge connections. What is derived is the gauge symmetry; that the link dynamics is the Wilson action is not derived.

5.6 Chirality, anomaly cancellation, and hypercharges

The gauge structure derived in §5.5 — local $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ from the cubic commutant and background independence — is the kinematic content of the Standard Model gauge sector. Two further structural features must hold for the emergent quantum field theory to match observation: the SU(2) factor must act chirally on left-handed fermions while SU(3) remains vector-like, and the hypercharges of the matter content must satisfy the six anomaly-cancellation conditions that the chiral structure imposes. This section develops the grading structure as a theorem of the framework rather than as an additional postulate, and states precisely where the physical chirality identification instead rests on the open hypothesis H-χ' (SM §4.8).

Chirality from the trace-out. The Standard Model is a chiral gauge theory: weak interactions distinguish left-handed from right-handed fermions, with the gauge bosons of SU(2) coupling only to left-handed fields. This is the empirical content of parity violation in beta decay, established in the 1950s. The structural origin of chirality in the framework is not a separate postulate; it emerges from the trace-out structure of the substratum partition.

The starting point is the staggered Dirac structure of the wave equation, established in §5.4. The Susskind decomposition assigns one Dirac spinor component to each hypercube corner of the cubic lattice, with the parity of the corner determining the ε-grade — the product $\gamma_5 \otimes \xi_5$ of spin and taste chirality — of the assigned component. Sites with $|\eta|$ even span the $+1$ eigenspace of $\gamma_5 \otimes \xi_5$ and sites with $|\eta|$ odd the -1 eigenspace; each grade contains both spin chiralities in equal dimension (certified exactly by `chirality_grading_probes.py`). This sublattice parity $\varepsilon(\mathbf{x}) = (-1)^{\sum x_\mu}$ is the lattice-level realization of γ_5, the chirality operator of the continuum Dirac theory.

The staggered Dirac operator has a structural property that the framework's chirality derivation uses: it couples only across sublattices. The even-even and odd-odd blocks of D_{st} vanish identically because the staggered Dirac operator anti-commutes with the sublattice parity, $\{D_{\text{st}}, \varepsilon\} = 0$, which is the lattice form of chiral symmetry established in §5.4. The full Dirac operator has the block form

$$D_{\text{st}} = \begin{pmatrix} 0 & D_{eo} \\ D_{oe} & 0 \end{pmatrix},$$

with off-diagonal blocks coupling the two sublattice grades and diagonal blocks vanishing.

The framework's checkerboard partition assigns even sites — the $+1$ grade of $\gamma_5 \otimes \xi_5$, carrying both spin chiralities — to the visible sector V, and odd sites — the -1 grade — to the hidden sector H. This is not an arbitrary choice; it is what the framework's structural partition produces for a cubic lattice with bipartite coupling. The visible sector that the embedded observer reads out is therefore one $\gamma_5 \otimes \xi_5$ grade — which carries BOTH spin chiralities, so calling it the "left-handed sublattice" would prejudge the taste-selectivity question; the complementary grade is part of the hidden sector that is traced over.

The trace-out structure determines the grading content of the emergent quantum field theory; physical chirality enters through the taste-selectivity hypothesis H-χ' (SM §4.8). Integrating out the hidden sector (the odd sublattice) via the Schur complement — a standard procedure in lattice quantum field theory — produces an effective action whose external legs lie in the visible half of the spin⊗taste space, where spin- and taste-chirality coincide:

$$\mathcal{L}_{\text{eff}} = \bar{\psi}_+ \left(D_{eo}\, G_{oo}\, D_{oe} \right) \psi_+ + \cdots,$$

where G_{oo} is the hidden-sector propagator. The hidden-grade components appear only as internal lines; the observable external states carry both spin chiralities, graded so that spin chirality equals taste chirality.

The two gauge factors couple differently to this graded effective theory. The SU(3) factor acts on internal T_1 components of the cubic decomposition — the three-dimensional spatial-vector representation. Both left-handed and right-handed Dirac components carry the same color charge, since the color index is internal to the T_1 block and independent of the sublattice parity. The trace-out preserves color coupling, and SU(3) remains vector-like, coupling identically to left-handed and right-handed components. The SU(2) factor acts on internal E components — the two-dimensional quadrupole representation. The internal E index commutes with the grading, so the SU(2) embedding as constructed is taste-blind; its observed spin-chirality therefore requires taste-chirality selectivity of the physical embedding — the named hypothesis H-χ' (SM §4.8), which the construction does not supply.

The vector-like nature of SU(3) is forced by the trace-out structure applied to the cubic-lattice substratum; the chirality of SU(2) rests additionally on H-χ', which is open. The framework's account of parity violation is correspondingly conditional: the substratum supplies the ε-grading and the visible-sector identity (spin chirality = taste chirality on the visible half), and the remaining step — H-χ' — is stated as an open hypothesis rather than derived from which sublattice the embedded observer occupies.

Evasion of Nielsen-Ninomiya. A potential objection arises from the Nielsen-Ninomiya theorem, which forbids a single Weyl fermion on a lattice under four hypotheses: locality of the action, hermiticity of the lattice Hamiltonian, translation invariance under the lattice group, and the action being bilinear in fermionic fields carrying a definite conserved chirality charge. Under these conditions, the lattice fermion spectrum is forced to contain equal numbers of left-handed and right-handed Weyl modes — the doubling phenomenon — so a chiral gauge theory cannot be regulated on a lattice in the conventional sense.

The framework manifestly satisfies the first three hypotheses. The wave equation is nearest-neighbor with range exactly $\epsilon = 2\sqrt{\sigma}\,\ell_P$ (locality); the bijection φ is real and the dynamics is T-invariant, so the induced operator is hermitian; the equivalence class $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$ is invariant under the cubic translation group. Were the fourth hypothesis to apply, the framework would inherit the no-go conclusion and chiral SU(2) would be impossible.

The fourth hypothesis does not apply, and the reason is structural rather than evasive. The Nielsen-Ninomiya theorem presupposes that the fundamental degrees of freedom are Grassmann fields $\psi(x)$ entering the action bilinearly, with chirality γ_5 defined on those fields ab initio. The framework's fundamental object is a real scalar bijection $\varphi : S \to S$ governed by the lattice wave equation. There are no fermions at the bijection level, no Grassmann variables, no γ_5, and consequently no $\mathrm{U}(1)_\chi$ on which an obstruction could be formulated. Dirac structure is derived through the Susskind factorization; chirality crystallizes only at the trace-out step. By the time fermions and chirality exist as derived structures, the trace-out has already broken the bilinear-fermionic-action setting that the fourth hypothesis requires.

The framework therefore satisfies hypotheses (NN1)–(NN3) but is bosonic at the level where (NN4) requires fermionic structure. Nielsen-Ninomiya is not circumvented by a clever fermion formulation; it is bypassed because the framework never enters the hypothesis space the theorem applies to. The doublers that the no-go theorem forces upon a fermionic lattice action have a striking reinterpretation in the framework: rather than being unwanted modes that contaminate the desired theory, they are the physical fermion generations of the Standard Model — Chapter 6 develops this identification in detail.

Anomaly cancellation forcing the hypercharges. A chiral gauge theory is consistent only if its quantum anomalies cancel. The chirality of SU(2) in the emergent theory means that the framework must satisfy six anomaly-cancellation conditions: three independent chiral gauge anomalies for the SU(3), SU(2), and $\mathrm{U}(1)_Y$ factors taken in various combinations, plus mixed anomalies involving the gravitational sector. These conditions impose constraints on the hypercharge assignments of the chiral fermion content.

The matter content of the framework, developed in detail in Chapter 6, consists of three candidate matter sectors of chiral fermions (three generations under H-spin', SM §4.7) in specific representations of $\mathrm{SU}(3)\times\mathrm{SU}(2)$: a left-handed quark doublet Q_L in $(\mathbf{3},\mathbf{2})$, a right-handed up quark u_R in $(\mathbf{3},\mathbf{1})$, a right-handed down quark d_R in $(\mathbf{3},\mathbf{1})$, a left-handed lepton doublet L_L in $(\mathbf{1},\mathbf{2})$, and a right-handed charged lepton e_R in $(\mathbf{1},\mathbf{1})$. The framework's structural commitments fix the $\mathrm{SU}(3)\times\mathrm{SU}(2)$ representations of the matter content; what remains free are the hypercharge assignments Y_Q, Y_u, Y_d, Y_L, Y_e of these five representations.

The six anomaly conditions are four independent constraints on these five hypercharge unknowns. Apart from the exceptional branch $Y_Q = Y_L = Y_e = 0$, $Y_u = -Y_d$ (SM Theorem 15), the remaining one-parameter freedom is the overall hypercharge normalization, a convention the anomaly conditions leave free (SM Theorem 15). With the normalization fixed, the anomaly conditions determine the hypercharges to the observed Standard Model values, uniquely up to the interchange $Y_u \leftrightarrow Y_d$ — a relabeling of which quark is called up-type, fixed by the electric-charge identification $Q_{\mathrm{em}} = T_3 + Y$:

$$Y_Q = \tfrac{1}{6}, \qquad Y_u = \tfrac{2}{3}, \qquad Y_d = -\tfrac{1}{3}, \qquad Y_L = -\tfrac{1}{2}, \qquad Y_e = -1.$$

The framework's derivation of the hypercharges is therefore non-trivial. The cubic-commutant construction in §5.5 produces a gauge group with three U(1) factors corresponding to the three blocks (B, L, S) of the cubic decomposition. A general abelian charge is a linear combination $Q = \alpha B + \beta L + \gamma S$ of these three factors. Of all such linear combinations, the six anomaly conditions identify exactly one, apart from the exceptional branch $Y_Q = 0$, as anomaly-free: the linear combination that produces the observed hypercharge Y of the Standard Model. The other two combinations cannot be gauged because gauging them would introduce non-cancelling chiral anomalies. They survive as accidental global symmetries of the emergent theory, identified as baryon number B and

lepton number L — broken only by the Weinberg operator and non-perturbative effects, exactly the structure observed in the Standard Model.

A corollary of the hypercharge derivation deserves explicit mention: the equality of the magnitudes of the proton and electron charges, $|q_p| = |q_e|$, follows on the nonzero-Y_Q branch from the framework's structural derivation rather than being a coincidence. The proton charge $q_p = 2Y_u + Y_d = 2 \cdot (2/3) + (-1/3) = 1$ and the electron charge $q_e = Y_e = -1$ are equal in magnitude because the anomaly conditions force the specific ratios that produce this cancellation. The famous one-to-one ratio between proton and electron charge magnitudes — observed to fantastic precision in atomic neutrality measurements — is, on that branch, a theorem of the framework, not a parameter fixed by observation.

5.7 What follows

The chapter has established an exact observer/geometric core and a conditional Standard-Model branch. On the chosen observer-level wave branch, the Susskind factorization applies; the six **geometric** cubic links decompose exactly as $(3, 2, 1)$; under H-link single-copy + H-cust the same blocks are used as the physical carrier; amplitude-scale invariance, a specified charged matter action and anomaly cancellation reduce the natural commutant $\mathrm{U}(3) \times \mathrm{U}(2) \times \mathrm{U}(1)$ to $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$; background independence promotes the global commutant to local gauge invariance; the trace-out yields the ε-graded visible sector on which spin- and taste-chirality coincide, so the taste-blind SU(3) remains vector-like while the chirality of SU(2) rests on the taste-selectivity hypothesis H-χ' (open); and, given that chiral structure, the six anomaly conditions force the hypercharges to the observed values, apart from the exceptional branch $Y_Q = 0$. The representation-theoretic and anomaly calculations remain theorem-level at their stated inputs, but the physical gauge-carrier chain now explicitly includes H-link single-copy and H-cust; weak chirality remains additionally conditional on H-χ'.

The chapter's exact six-link representation sits at the structural level, while the **physical gauge-group identification is conditional** on H-link/H-cust and therefore must not be classified as unconditional S-class structure. The cubic multiplicities are exact geometry. The physical gauge-group reading is conditional on H-link/H-cust and then on amplitude-scale/background-independence premises and a specified charged matter action; anomaly-fixed hypercharges are exact conditional on the resulting chiral matter assignment. The hypercharges are not chosen from a continuum of possibilities; apart from the exceptional branch $Y_Q = 0$, they are determined by anomaly cancellation. Falsification of either would refute the framework directly: any stable beyond-Standard-Model gauge structure, any departure of the hypercharges from the values derived here, would violate the framework's Class A structural commitments developed in Chapter 4 §4.5.

Chapter 6 develops the specific quantitative content this gauge structure pro-

duces. The matter content is set out there: three chiral fermion generations from the staggered taste structure of the wave equation, one Higgs doublet $(\mathbf{1}, \mathbf{2}, +1/2)$ taken as input and accommodated in the E projection of the singlet staggered taste (SM Theorem 11), with three candidate tastes on the **single-copy H-link** branch (three physical generations only under H-spin', SM §4.7); the framework does not presently derive the copy number $m = 1$. The strong-CP problem is treated structurally in Section 6.3, which states what T-invariance of the substratum bijection combined with detailed balance fixes ($\theta \in \{0, \pi\}$ and reciprocal transition counts) and what it does not ($\bar{\theta}$, open under two named conditions). The gauge couplings at the Z pole are reproduced by a chain with one structurally computed input, the analytic one-loop staggered vacuum polarization $1/\alpha_0 = 23.25$, and three threshold parameters fitted to the three observed couplings (SM §6.3). The twenty-two classified quantitative retrodictions developed there (two parameter-free structural and five conditional structural; Appendix A) include the Cabibbo angle $1/(\pi\sqrt{2})$ matched to 0.04%, the Koide angle $\theta_0 = 2/9$ matched to 0.001%, the six fermion masses from a single empirical input matched to within 1.3%, all three PMNS mixing angles within 1.1σ, and the Higgs quartic boundary condition $\lambda(M_{\rm Pl}) = 0$, matching the RG-extrapolated value at 0.6σ (the m_H band 129–132 GeV sits 4–7 GeV above observation). The framework's quantitative empirical record outside cosmology is concentrated in Chapter 6.

Chapter 6

The Matter Content and Quantitative Predictions

6.1 What this chapter develops

Chapter 5 derived the gauge structure of the Standard Model from the cubic-lattice substratum. The wave equation factored into staggered Dirac fermions through the Susskind decomposition; the link count under H-link fixed the internal-component count at $K = 6$; the cubic-group decomposition produced eigenvalue multiplicities $(3, 2, 1)$; amplitude-scale invariance, a specified charged matter action and anomaly cancellation reduced the natural commutant to $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$; background independence promoted the global commutant to local gauge invariance; the trace-out yielded the ε-graded visible sector, with the taste-blind SU(3) vector-like and the chirality of SU(2) conditional on the taste-selectivity hypothesis H-χ' (Chapter 5 §5.6); given that chiral structure, anomaly cancellation forced the observed hypercharges, apart from the exceptional branch $Y_Q = 0$. The gauge sector of the Standard Model is forced — uniquely — by the structural inputs of Part I and the cubic-lattice realiza-

tion, with three qualifiers: the gauge carrier rests on H-link with its single-copy clause — the open hypothesis that the physical carrier is one copy of the six-link representation (Chapter 5 §5.4) — together with an octahedrally equivariant condensate (Chapter 5 §5.5), the reduction of the natural commutant to $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ is conditional on amplitude-scale gauge invariance, an explicit adjoined principle (Chapter 5 §5.2), together with a specified charged matter action, and the chirality of the weak sector is conditional on the taste-selectivity hypothesis H-χ' (Chapter 5 §5.6).

This chapter develops the specific quantitative content that the gauge structure produces. Three pieces of structural derivation come first. Section 6.2 derives the matter content: three chiral fermion generations from the staggered taste structure of the wave equation, one composite Higgs doublet from the singlet taste, with the candidate-sector count locked at three by the same argument that fixed $K = 6$ (three physical generations only under H-spin', SM §4.7) — the link count under H-link jointly with the condensate-stabilizer accounting. Section 6.3 treats the strong-CP problem structurally: T-invariance of the substratum bijection combined with detailed balance narrows the QCD angle to $\theta \in \{0, \pi\}$ and gives reciprocal transition counts, but does not fix $\bar{\theta}$, which is open under two named conditions stated at the end of that section. Section 6.4 develops the gauge-coupling derivation and the framework's structural no-GUT prediction, with the analytic one-loop staggered vacuum polarization producing the structural input $1/\alpha_0 = 23.25$ and the RG running to M_Z matching the three observed gauge couplings.

The remaining sections develop the framework's parameter-free quantitative retrodictions, in the two categories of Section 6.8 — independent, and conditional on the empirical mass-hierarchy bijection. Section 6.5 derives the Cabibbo angle as $1/(\pi\sqrt{2})$ from a single Brillouin-zone distance — matched to the observed value at 0.04%, the framework's sharpest geometric prediction in the quark sector. Section 6.6 derives the Koide relation as $Q = 2/3$ from a cubic-group quadratic Casimir — matched to 0.0003%, the sharpest empirical agreement in the framework in relative precision, with the lepton mass chain following from one empirical input through the Koide structure — $Q = 2/3$ with the derived phase $\theta_0 = 2/9$ — at 0.003%. Section 6.7 derives the three PMNS mixing angles from the cubic-group representation theory with parity-graded charged-lepton corrections — all three angles matched within 1.1σ at current precision, including the JUNO solar mixing angle $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2)$ matched at 0.07σ by the post-JUNO global fit (a retrodiction). Section 6.8 collects the twenty-two predictions in a summary table classified by both derivational layer (Class column) and independent-evidence status (Cat column), and closes the chapter with a pointer forward to Chapter 7 (gravitational sector), Chapter 8 (JUNO in detail), and the framework's broader empirical commitments.

The methodological discipline of Chapter 4 organizes the predictions. The Cabibbo angle is Class B-C (conditional structural, Layer 1, substratum geometry with the emergent dressing of its identification open). The Koide relation is

Class B-C (Layer 1, cubic-group representation theory with the dynamical closure of its amplitude open). The Bekenstein-Hawking 1/4 factor — developed in Chapter 7 — is a consistency relation, conditional on H-slope at the horizon (Chapter 7 §7.3). The gauge couplings at M_Z are Class B-R (retrodiction): three fitted parameters (δ_0, A, B) against three observed couplings produces zero residual by construction, with the structural content lying in the *form* of the fit rather than in the matching values. The six fermion masses inherit from one empirical input (the strange-quark mass) through Class B-M (mass-chain inheritance). The PMNS angles are Class B-L (layered conditional): the substrate coefficient ratio of Condition 2 is Layer 2(b) — the second-order substrate route is excluded (SM §7.3) and the 7/6 sum rule is the direct experimental test — while the directional/assignment content is Layer 2(b) bijection-specific per Chapter 2 §2.5 (the empirical mass-hierarchy fixing which generation is electron). The framework's empirical record in fundamental physics outside cosmology is concentrated in these classes, with each prediction's derivational status explicit.

The chapter's content sits at Levels 1 and 2 of the four-layer framing developed in Chapter 4. Layer 1 predictions (Cabibbo, Wolfenstein, Koide) follow from pure substratum geometry or representation theory with no operator-relation inputs; Layer 2 predictions (mass ratios within fixed operator structures, mixing-angle values) involve specific solution-level inputs. The framework's strongest predictions — the ones that most distinguish OI from other observer-admitting frameworks — live at Layer 1 and at the Level G3 × Level D intersection of the hierarchy developed in Chapter 3.

6.2 Three generations and the composite Higgs

The wave equation on the cubic lattice has staggered Dirac structure, as Chapter 5 established. The staggered decomposition produces $N_{\text{taste}} = 2^{\lfloor (d+1)/2 \rfloor}$ Dirac tastes in $d+1$ spacetime dimensions, with $N_{\text{taste}} = 4$ at $d = 3$. The framework's matter content — three chiral fermion generations plus one Higgs doublet — emerges from how these four staggered tastes decompose under the cubic rotation group and how the embedded observer's gauge structure acts on the resulting sectors.

The taste decomposition under the cubic group. The four staggered tastes can be labeled by the corners of the Brillouin zone they originate from: the $2^d = 8$ Brillouin-zone corners on a $d = 3$ lattice pair into 4 taste pairs under the corner-conjugation $\eta \leftrightarrow \mathbf{1} - \eta$. Two corners are O-invariant under the cubic rotation group: the Γ point $(0,0,0)$ and the R point $(1,1,1)$. These two corners pair into one taste pair that transforms as the singlet A_1 representation. The remaining six corners form three axis-aligned pairs (X, M) along the three spatial directions, transforming as the triplet T_1 representation — carried by the signed vector action of the spin-taste matrices γ^j; the unsigned pair labels alone transform only as $A_1 \oplus E$, since sign flips act trivially on the corner components. The decomposition is exact:

$$4 = \mathbf{1} \oplus \mathbf{3}.$$

One singlet taste; three triplet tastes related by cubic symmetry.

Spin-statistics from the lattice. The singlet taste produces a spin-0 field; the triplet tastes produce spin-1/2 fields — the algebraic labels proved and probe-certified, the physical identification the named hypothesis H-spin' [SM §4.7]. The argument runs through the staggered-to-Dirac reconstruction. The singlet pairs Γ and R in the Brillouin zone, with corresponding Dirac matrices $\Gamma(0) = I_4$ (scalar) and $\Gamma(1) = \gamma^1\gamma^2\gamma^3$ (pseudoscalar). Both are SO(3)-invariant, giving total angular momentum $J = 0$. The j-th triplet pair has matrices γ^j (vector) and $-i\Sigma_j$ (spin generator), giving total angular momentum $J = 1/2$.

The corollary, at the level of the algebraic labels: the spin-statistics correspondence between the singlet taste (spin-0, bosonic) and the triplet tastes (spin-1/2, fermionic) follows from the lattice representation theory — with the physical spin identification riding H-spin' (SM §4.7) — not from a separate spin-statistics postulate. The Higgs (spin-0 boson) and the fermions (spin-1/2) emerge as distinct sectors of the same underlying staggered decomposition, with the cubic-group representation forcing the spin content of each sector.

Three degenerate fermion sectors. The three triplet staggered tastes provide three degenerate spin-1/2 sectors related by cubic symmetry — the structural origin of the three-generation pattern of the Standard Model. The cubic rotation group O permutes the three spatial axes, and the three triplet tastes transform among themselves under this permutation. The three sectors are therefore degenerate at the level of cubic symmetry, with their Yukawa couplings to the Higgs producing the observed fermion mass hierarchy through generation-dependent symmetry breaking that the specific bijection φ controls.

The candidate-sector count is locked at three by the same argument that fixed $K = 6$ — its reading as three physical generations rides H-spin' (SM §4.7), whose free-kernel and 4D routes are both settled negative in Chapter 5 — the link count under H-link jointly with the condensate-stabilizer accounting. The chain runs as follows: the link count under H-link gives $K = 2dm$ on $d = 3$, and the single-copy clause $m = 1$ with the stabilizer accounting fixes $K = 2d = 6$ (Chapter 5 §5.4); the six internal components host the $2^3 = 8$ Brillouin-zone corners; the corners pair into $N_{\text{taste}} = 4$ Dirac tastes; the cubic-group decomposition gives $4 = \mathbf{1} \oplus \mathbf{3}$, with one scalar taste and three spin-1/2 tastes. The generation count is not chosen from a continuum of possibilities; it is a forced consequence of $K = 6$ on $d = 3$ — the link count under H-link single-copy together with the single-gauge-sector accounting.

Adding a fourth fermion generation would require either expanding the internal-component count beyond $K = 6$ (which departs from the single-copy clause $m = 1$ of H-link) or finding a fourth distinct staggered taste (which does not exist for $d = 3$). The framework's structural commitment is therefore that no stable fourth-generation fermion with Standard Model quantum numbers can exist. This is a pre-registered falsification condition: any observation of such a fermion would refute the framework's taste-reduction derivation at the Class A

level identified in Chapter 4 §4.5.

The formal uniqueness theorem closing the T_1 generation-assignment — establishing that the assignment of three fermion generations to the T_1 triplet (rather than to $3A_1$, $A_1 \oplus E$, T_2, or other distributions) is forced by the framework's existing commitments — is developed in Appendix B.6c.

The matter content on links. The Standard Model matter representations — Q_L in $(\mathbf{3}, \mathbf{2})$, u_R and d_R in $(\mathbf{3}, \mathbf{1})$, L_L in $(\mathbf{1}, \mathbf{2})$, e_R in $(\mathbf{1}, \mathbf{1})$ — involve cross-charged tensor products. A left-handed quark doublet carries both color **3** and weak isospin **2** simultaneously, requiring a tensor-product representation rather than a direct-sum decomposition. The framework's link-carrier construction provides the natural home for these representations.

The local gauge invariance established in Chapter 5 §5.5 makes the coupling matrix $M(\mathbf{n}, \hat{e}_j)$ the gauge connection on a nearest-neighbor link, transforming under local gauge transformations as $M(\mathbf{n}, \hat{e}_j) \to G(\mathbf{n})\, M(\mathbf{n}, \hat{e}_j)\, G(\mathbf{n} + \hat{e}_j)^{-1}$. The link variable carries one gauge index at each endpoint, living in the tensor-product space $V_K^{(\mathbf{n})} \otimes V_K^{(\mathbf{n}+\hat{e}_j)*}$ rather than in a single V_K. The matter content on links inherits this tensor-product structure: a left-handed quark doublet at a link site can have a definite color label at one endpoint and a definite weak-isospin label at the other, with no algebraic obstruction.

The framework's content on links is therefore that matter propagates along links and the gauge representation of a matter field is read off from the tensor-product structure of the link's two endpoints. The three triplet tastes distribute identical matter content across the three candidate sectors (generations under H-spin'); the singlet taste accommodates the Higgs doublet, taken as input, in its E projection (the only projection that admits a gauge-invariant Yukawa coupling to the fermion doublets, since the T_1 projection is excluded on color grounds and the A_1 projection on weak grounds; SM Theorem 11). The matter content of three identical generations of $(Q_L, u_R, d_R, L_L, e_R)$ together with the one Higgs doublet taken as input — exactly the Standard Model matter content — follows.

The Higgs as a composite scalar. A structural feature of the framework's Higgs sector deserves explicit treatment. The Higgs is not a fundamental scalar field. It is a composite excitation in the singlet staggered taste of the wave equation — the E projection of the $A_1 \oplus E \oplus T_1$ decomposition under the cubic group. The Higgs's gauge representation $(\mathbf{1}, \mathbf{2}, +1/2)$ is taken as input; the E projection of the singlet taste is the one projection that accommodates it with a gauge-invariant Yukawa coupling (SM Theorem 11).

The composite nature of the Higgs has implications for the hierarchy problem. In the standard formulation, the smallness of the Higgs mass compared to the Planck scale is mysterious: radiative corrections to a fundamental scalar's mass should generically push it toward the highest available scale, and the observed $m_H \sim 125$ GeV requires a tuning of one part in 10^{34}. In the framework, the

Higgs mass is set by the block values (a, b, c) of the equivariant condensate Σ (SM Theorem 5) — properties of the substratum bijection φ — rather than by radiative corrections. The emergent quantum field theory's notion of "radiative correction" applies only to the perturbative expansion around the substratum's solution; the substratum itself fixes the Higgs mass at the value the bijection produces, with no separate fine-tuning required.

Whether this constitutes a *solution* to the hierarchy problem in the conventional sense depends on how the substratum's choice of φ is itself viewed. The framework treats the specific bijection φ within the equivalence class $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$ as a solution-specific property — what Chapter 4 called Layer 3 content, depending on the specific representative within the gauge class. The hierarchy is dissolved at the level of the framework's emergent description (no radiative tuning required at the emergent level) and reframed at the substratum level (the bijection has whatever Higgs-mass eigenvalue it has). The reframing is a methodological choice consistent with the framework's general pattern: structural commitments at the level of the equivalence class; solution-specific inputs at the level of the specific representative.

6.3 The strong-CP problem: what time reversal fixes

The strong-CP problem is one of the Standard Model's most persistent puzzles. Quantum chromodynamics admits a CP-violating term $\mathcal{L}_\theta = (\theta/32\pi^2)\,\mathrm{Tr}(F\tilde{F})$ where F is the gluon field-strength tensor. This term would produce CP-violation in strong interactions, observable as a non-zero neutron electric dipole moment. Experimental bounds limit θ to less than approximately 10^{-10} — a remarkable smallness for a parameter the Standard Model treats as free, ranging in principle over $[0, 2\pi]$. The conventional resolution — the Peccei-Quinn mechanism with an associated axion — postulates a new global symmetry whose spontaneous breaking dynamically drives θ to zero. No axion has been observed at expected scales.

The framework's contribution is structural and partial: it fixes what time-reversal symmetry of the substratum does and does not determine about the combination $\bar{\theta} = \theta + \arg\det(Y_u Y_d)$, the physically meaningful CP-violating parameter, invariant under field redefinitions. The answer is that T-invariance leaves $\bar{\theta}$ open, under two named conditions stated at the end of this section. Three structural inputs carry the analysis: T-invariance of the substratum bijection, the spatial nature of the framework's partition, and the reciprocity of the resulting transition counts.

T-invariance of the wave equation. The first input is that the discrete wave equation is invariant under time reversal. Under the transformation $\phi(\mathbf{n}, t) \to \phi(\mathbf{n}, -t)$, the wave equation $\phi(\mathbf{n}, t{+}1) = \sum_j [\phi(\mathbf{n}{+}\hat{e}_j, t){+}\phi(\mathbf{n}{-}\hat{e}_j, t)]{-}\phi(\mathbf{n}, t{-}1)$ maps to itself: the time-reversed field satisfies the same equation. T-invariance is a structural property of the wave equation, not an additional postulate. The substratum dynamics is symmetric under time reversal in a precise mathematical

sense.

The framework's partition — the visible/hidden checkerboard developed in Chapter 5 — marginalizes over spatial sites, not over temporal data. The partition is spatial, preserving the temporal structure of the substratum dynamics. T-invariance of the substratum implies T-invariance of the visible-sector marginalization: the time-reversal operation commutes with the partition's spatial projection.

What T-invariance fixes about θ. The θ-term $\mathcal{L}_\theta$ is T-odd. The trace $\mathrm{Tr}(F\tilde{F})$ equals $2\,\mathbf{E}\cdot\mathbf{B}$ when expressed in terms of the chromoelectric and chromomagnetic fields, and under time reversal $\mathbf{E} \to \mathbf{E}$ while $\mathbf{B} \to -\mathbf{B}$. A T-invariant Lagrangian therefore requires $\theta = -\theta$ modulo 2π — and that equation has two solutions, $\theta = 0$ and $\theta = \pi$, both of which are time-reversal-invariant points of Yang–Mills theory. T-invariance of the emergent Lagrangian, inherited from the substratum because the partition is spatial, narrows θ to $\{0, \pi\}$; selecting the $\theta = 0$ branch is a separate condition, named H-top, which the symmetry does not supply.

Detailed balance. The second input is reciprocity of the visible-sector transition counts. The argument runs through a counting lemma: for each hidden state h that carries a visible initial state i to a visible final state j under the bijection φ, time reversal supplies a hidden state carrying the time-reversed j to the time-reversed i, because T-invariance gives $\varphi = T \circ \varphi^{-1} \circ T$ and the spatial partition preserves the visible/hidden classification under T. The correspondence is a bijection between two finite sets of hidden states, so the transition counts agree: $T_{ij} = T_{\theta j, \theta i}$, where θ is the action of time reversal on visible states — plain detailed balance, $T_{ij} = T_{ji}$, when the visible states carry no T-odd data. Since φ^n is a bijection with the same property, reciprocity holds at every time scale; numerically, the T-violation of the visible transition counts is exactly zero for ring sizes $L = 4, 6$, alphabet sizes $q = 2, 3, 5$, and all times $n = 1, \ldots, 12$. Detailed balance is therefore not an emergent thermodynamic property requiring equilibration; it is a direct consequence of the substratum's bijection structure combined with T-invariance and spatial partitioning.

What reciprocity does not give. Reciprocity of transition probabilities does not make the emergent Hamiltonian T-invariant. A four-site ring with the same complex hopping amplitude $z = a + ib$ on every link in one direction has $|U_{ij}(t)|^2 = |U_{ji}(t)|^2$ exactly at every time — the ring has a reflection exchanging any two sites, and a reflection reverses the loop — yet its loop product z^4 carries a phase that no rephasing can remove and no real Hamiltonian can have; no time-reversal operator inherited from the substratum commutes with it. Wigner's theorem does not close the gap: it represents a given ray transformation and does not manufacture a physical time reversal from the equalities $|U_{ij}|^2 = |U_{ji}|^2$. The bipartite structure of that example is exactly the structure of the framework's staggered sector, where the triangle phases that would otherwise tie reciprocity to reality do not exist. So the chain from substratum T-invariance to a T-

invariant emergent Hamiltonian, and from there to real Yukawa matrices and $\arg\det(Y_u Y_d) = 0$, breaks at its second step [SM §5.3, Proposition 20].

The strong-CP status. T-invariance of the substratum therefore fixes neither θ, which may be 0 or π, nor $\arg\det(Y_u Y_d)$, and $\bar{\theta}$ is not determined by the construction [SM §5.3, Theorem 21]. The problem is open under two named conditions: H-top, the selection of $\theta = 0$ over π; and H-det, a mechanism fixing $\arg\det(Y_u Y_d) = 0$ at the emergent level while leaving the rephasing-invariant flavor quartets free. The framework makes no prediction for the neutron electric dipole moment; a measured $\bar{\theta} \neq 0$ would constrain mechanisms proposed for H-det, not the T-symmetry of the substratum, which is compatible with any value of $\bar{\theta}$.

The narrowing of the target is partly a relief. The CKM CP violation, encoded in the Jarlskog invariant $J \approx 3 \times 10^{-5}$, arises from the basis mismatch between up-type and down-type Yukawa mass eigenstates; reciprocal transition probabilities can hide exactly such loop phases, so detailed balance at the substratum does not kill weak CP violation, and there is no algebraic conflict between $\arg\det(Y_u Y_d) = 0$ and $J \neq 0$. Conversely, a mechanism that fixed $\arg\det(Y_u Y_d) = 0$ by making Y_u and Y_d simultaneously real would force the CKM matrix to be real orthogonal and $J = 0$ — it would have solved too much, and that is the falsification control on any proposed H-det. The flavor-sector CP phases themselves are solution-specific properties of the particular bijection φ, treated as inputs rather than predictions; Chapter 4 §4.6 lists them among the solution-specific properties whose measurement does not bear on the framework's structural status.

6.4 Gauge couplings and the no-GUT prediction

The Standard Model has three gauge couplings: α_3 for color, α_2 for weak isospin, α_1 for hypercharge, measured at the Z boson mass. The values $\alpha_3(M_Z) \approx 0.1181$, $\alpha_2(M_Z) \approx 0.0339$, $\alpha_1(M_Z) \approx 0.0169$ are taken as independent inputs in the Standard Model. The framework's derivation of the gauge couplings extends the chain through a structural input — the one-loop staggered vacuum polarization — combined with threshold parameters that match the three couplings simultaneously. The result is a structural account of *why* the gauge couplings have approximately the values they have, including the framework's distinctive structural prediction of no grand unification.

The induced coupling from the fermion determinant. The gauge field in the framework is not an independent dynamical degree of freedom. It is the state-dependent coupling matrix $M(\mathbf{n}, \hat{e}_j)$ that Chapter 5 §5.5 identified as the gauge connection on a nearest-neighbor link. The matter content — the staggered fermion field ϕ on the lattice — determines the coupling matrix through the wave equation. The gauge kinetic term is therefore *induced* by the fermion determinant rather than being a separate input.

The effective action for the gauge connection, obtained by integrating out the fermion degrees of freedom, is

$$S_{\text{eff}}[U] = -N_f \,\text{Tr}\ln(D^\dagger D[U]),$$

where N_f is the number of fermion species and D is the staggered Dirac operator coupled to the gauge connection U. Expanding S_{eff} to second order in the gauge field gives the gauge propagator and the induced gauge coupling, computed at one loop from the staggered vacuum polarization $\Pi_s(0)$ — a lattice momentum integral over the fermion bubble with the framework's specific vertex structure.

The induced coupling has a universal structural value. The Dynkin index $T(R) = 1/2$ per Dirac fermion is the same for all three gauge factors because every component of the coupling matrix M — with $K = 6$ components distributed across the $(3, 2, 1)$ cubic decomposition — transforms in the fundamental representation of its respective gauge group. Combined with $N_f = 6$ (one fermion species per internal component, fixed by the $K = 2d = 6$ result of Chapter 5 §5.4), the induced inverse coupling at the matching scale $\mu = M_{\text{Pl}}$ — a matching convention, the lattice scale itself being $1/\epsilon = M_{\text{Pl}}/(2\sqrt{\sigma})$ for an entropy σ per boundary mode (Chapter 7) — is

$$\frac{1}{\alpha_0} = 6 \times \Pi_s(0) \times 4\pi = 23.25,$$

with $\Pi_s(0) \to 0.3084$ on a robust numerical plateau as the lattice integral converges. This is a strictly structural prediction — depending only on N_f and $T(R)$, both determined by the lattice structure — and is universal across the three gauge factors because the Dynkin index equality $T_3(R) = T_2(R) = T_1(R)$ holds at the substratum level.

The threshold structure. The universal coupling α_0 at the matching scale M_{Pl} does not yet match the observed couplings at M_Z. Two corrections enter at the matching between the substratum and the emergent quantum field theory: a universal threshold δ_0 from higher-loop corrections to the vacuum polarization, and a group-dependent threshold from gauge self-interactions proportional to the quadratic Casimir C_2 of each gauge factor.

The expansion parameter for the group-dependent threshold is $C_2 \cdot g_0^2 = C_2 \times 4\pi\alpha_0$, which equals 1.08 for SU(2) ($C_2 = 2$) and 1.62 for SU(3) ($C_2 = 3$). Both are of order unity at the matching scale, so finite-order perturbation theory converges too slowly to capture the threshold from first principles. The framework's account is that the threshold is given by an all-orders geometric-series resummation of the gauge self-energy, with the form

$$\frac{1}{\alpha_i(M_Z)} = \underbrace{23.25}_{1/\alpha_0} + \underbrace{10.0}_{\delta_0} + \underbrace{A \cdot \ln(1 + B \cdot C_2 g_0^2)}_{\text{resummed gauge self-energy}} + \underbrace{\frac{b_i}{2\pi} \ln \frac{M_{\text{Pl}}}{M_Z}}_{\text{SM RG running}},$$

where A and B are the threshold parameters and b_i is the standard Standard Model beta-function coefficient for each gauge group.

The four components have distinct derivational statuses. The structural input $1/\alpha_0 = 23.25$ is computed analytically from the one-loop staggered vacuum polarization, with no fitted parameters. That input is the ambient staggered-background coefficient: the hypercharge factor's microscopic connection is flat, so reading the number as a native hypercharge coupling carries two further conditions — that the observer's state bundle carry projective curvature in that channel, and that it couple with the vertex the calculation assumes. The non-abelian factors do not inherit them. The universal threshold $\delta_0 = 10.0$, in the $\mu = M_{\rm Pl}$ matching convention, is empirically fixed by the U(1) row of the matching equations; a two-loop staggered VP rebuild indicates that strict two-loop perturbation theory does not reproduce δ_0, so it is honestly treated as an empirical input rather than a derived value. The C_2-dependent threshold parameters $(A, B) = (8.3, 5.59)$ are matched to the SU(2) and SU(3) couplings at M_Z, with the product $A \cdot B = 46.4$ independently computed at $A \cdot B \approx 48$–62 depending on the assumed leading finite-size power — a cross-check that agrees with the fit under the $1/N^2$ assumption and is inconclusive across finite-size models. The Standard Model RG running below $M_{\rm Pl}$ is standard.

The classification per Chapter 4 §4.5 is therefore that the three gauge couplings at M_Z are Class B-R retrodictions: three fitted parameters (δ_0, A, B) against three observed couplings produces zero residual by construction. The structural content of the framework's gauge-sector account does not lie in matching the three coupling values — that is achieved by the fit. It lies in the specific value $1/\alpha_0 = 23.25$ from the one-loop staggered VP and the universality of δ_0 across the three gauge groups (no group-dependent term). The finite-lattice $A \cdot B$ computation is a separate numerical cross-check: its extrapolated values span ≈ 48–62 across leading-power models, and agreement with the fitted product is conditional on the $1/N^2$ assumption.

The structural no-GUT prediction. The framework's most distinctive content in the gauge-coupling sector is structural rather than quantitative: the three gauge couplings of the Standard Model do not unify at a single energy scale, and no grand unified theory underlies the Standard Model gauge group. The prediction has three layers.

At the group-theoretic level, the Standard Model gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ emerges, conditionally on H-link and an octahedrally equivariant condensate (Chapter 5 §5.4–§5.5), directly from the cubic decomposition of the six nearest-neighbor link directions — a single-step decomposition with three eigenspaces of the condensate Σ and three commutant factors. There is no intermediate large gauge group at any scale: no SU(5), no SO(10), no E_6, no Pati-Salam $\mathrm{SU}(4) \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$. The framework has no GUT stage, structurally. This contrasts with all standard string-phenomenology routes to the Standard Model, each of which traverses an intermediate large gauge group during compactification.

At the coupling level, the three gauge couplings do not unify at any energy scale. The universal $1/\alpha_0 = 23.25$ at the matching scale is shared across all three

gauge factors before gauge self-interactions are included; but the C_2-dependent threshold $A\ln(1+B\cdot C_2 g_0^2)$ splits the couplings *at the matching scale itself*, with C_2 different across the three gauge groups. Below $M_{\rm Pl}$, the standard Standard Model beta functions run the already-split couplings to lower scales, with no intermediate gauge structure modifying the running. The three gauge couplings therefore remain distinct at all scales accessible to the emergent quantum field theory; they were never unified, even at the Planck scale.

At the mechanism level, the gauge couplings emerge from the fermion determinant — a specific mechanism that string compactifications do not share. String-theoretic gauge couplings depend on moduli (compactification volumes, dilaton vacuum expectation values, flux choices) and are typically not related to one another structurally. The framework's gauge couplings, by contrast, share an origin: they all arise from the same one-loop staggered vacuum polarization with the same N_f and $T(R)$, with the differences among them attributable to the C_2-dependent threshold structure. The three couplings are not three independent observables but three projections of a common substructure.

The structural no-GUT prediction has empirical consequences. The most direct is proton stability: the framework predicts no proton decay through GUT-mediated channels, with the only available baryon-number-violating operators suppressed by $M_{\rm Pl}^2$ rather than $M_{\rm GUT}^2$. The expected proton lifetime in the framework is $\tau_p \sim M_{\rm Pl}^4/m_p^5 \sim 10^{45}$–$10^{46}$ years, roughly ten orders of magnitude beyond Hyper-Kamiokande's projected $\sim 10^{35}$ year sensitivity. Any detection of proton decay at experimentally accessible scales would refute the framework's no-GUT commitment. Tightening of the proton lifetime bound past 10^{35} years would support the framework's prediction. Two further consequences: no GUT-chain magnetic monopoles (so the cosmological monopole problem that inflation was historically invoked to dissolve does not arise in the framework), and no GUT-chain cosmic strings from $\mathrm{SO}(10) \to \mathrm{SM}$ symmetry breaking with non-trivial π_1. Standard cosmic-string non-detection in CMB and gravitational-wave observations is consistent with the framework's prediction.

The framework-specific discriminator within the no-GUT class is the shared origin of the three gauge couplings: the structural prediction $1/\alpha_0 = 23.25$ as a single value shared across $\mathrm{SU}(3)\times\mathrm{SU}(2)\times\mathrm{U}(1)$ before threshold corrections. Other no-GUT frameworks would generically have three independent gauge couplings without this shared-origin structural relation. The matching of $\alpha_3, \alpha_2, \alpha_1$ at M_Z uses three fitted threshold parameters against three observed couplings, giving zero residual by construction. It therefore supplies no independent confirmation of the shared-origin relation; the structural input and threshold constraints are assessed separately in Chapter 5 §5.5.

6.5 The Cabibbo angle and CKM structure

The Cabibbo–Kobayashi–Maskawa (CKM) matrix parametrizes the misalignment between the mass eigenstates and the weak interaction eigenstates of the

down-type quarks. Its leading parameter, the Cabibbo angle $\lambda \approx 0.225$, controls the strength of generation-changing weak transitions. In the Standard Model the Cabibbo angle is a free parameter measured to be $\lambda = 0.22500 \pm 0.00067$. The framework obtains λ from the geometry of the cubic lattice with no fitted parameters, under an identification hypothesis whose emergent dressing is open (Class B-C; SM §7.1), matching the observed value to 0.04% (0.12σ) — the framework's sharpest geometric prediction in the quark sector.

The Cabibbo angle from a Brillouin-zone distance. The three fermion generations of the framework occupy the three triplet (T_1) staggered tastes, located at the three pairs of axis-aligned corners of the Brillouin zone. In a normalized convention where the Brillouin zone has dimension 2π, the three triplet corners $X_j = \pi\,\hat{e}_j$ sit at positions $(\pi, 0, 0)$, $(0, \pi, 0)$, $(0, 0, \pi)$ along the three coordinate axes. The distance between two adjacent triplet corners is

$$|X_i - X_j| = |\pi\,\hat{e}_i - \pi\,\hat{e}_j| = \pi\sqrt{2},$$

a geometric constant of the cubic lattice.

The Cabibbo angle equals the inverse of this distance:

$$\lambda = \frac{1}{\pi\sqrt{2}} = 0.22508.$$

Observed value: $\lambda = 0.22500 \pm 0.00067$ (global fit with unitarity imposed). The framework's prediction matches to 0.04% — well inside experimental uncertainty.

The mixing matrix element between generations i and j is the continuum fermion propagator evaluated at the inter-generation momentum:

$$|M_{ij}| = |S(X_j - X_i)| = \frac{1}{|X_j - X_i|} = \frac{1}{\pi\sqrt{2}}.$$

The propagator has $1/|q|$ falloff rather than $1/|q|^2$ because the taste-changing transition preserves chirality, established in the staggered-lattice literature: taste-changing vertices in staggered quarks have spinor structure given by a combination of γ_μ and $\gamma_{5\mu} = \gamma_5\gamma_\mu$, both of which map left-handed to left-handed and right-handed to right-handed rather than flipping between them. In the massless limit the chirality-preserving propagator structure contracted with the chirality-preserving vertex gives $|M_{ij}| \propto 1/|q|$. The corresponding chirality-flipping process would give $1/|q|^2$ falloff, which produces the Gatto-Sartori-Tonin relation $\lambda^2 = m_d/m_s$ that is discussed below.

Comparison with other Bravais lattices. The Cabibbo angle on simple cubic is $\lambda_{\mathrm{SC}} = 1/(\pi\sqrt{2}) = 0.2251$, matching observation at 0.04%. On body-centered cubic, the triplet sector sits at the H-points of the BCC Brillouin zone with separation $\pi\sqrt{6}$, giving $\lambda_{\mathrm{BCC}} = 0.1299$ — discrepant from observation by 42%. On face-centered cubic, the triplet sector sits at the L-points with

separation 2π, giving $\lambda_{\text{FCC}} = 0.1592$ — discrepant by 29%. The simple-cubic Bravais-lattice commitment from Chapter 5 §5.5 is quantitatively confirmed by the Cabibbo prediction: only simple cubic produces the observed value.

Isolation in integer-input space. A complementary robustness check examines whether nearby alternatives to the specific integer $N = 2$ in the form $\lambda = 1/(\pi\sqrt{N})$ also match observation. They do not. The table below shows predictions for $N \in \{1, 2, 3, 4, 6, 8\}$ — the integers that arise naturally from inter-corner distances in various cubic-lattice geometries — against the observed $\lambda = 0.22500 \pm 0.00067$:

N	$\lambda = 1/(\pi\sqrt{N})$	Deviation
1	0.3183	$+139\sigma$
2 (framework)	**0.2251**	$+0.1\sigma$
3	0.1838	-62σ
4	0.1592	-98σ
6	0.1299	-142σ
8	0.1125	-186σ

The framework's $N = 2$ is the only integer value in the nearby alternatives that matches observation within 1σ. The next-closest alternative ($N = 3$) deviates by 62 standard deviations. The Cabibbo prediction is *isolated* in the space of nearby alternatives — not selected from a cluster of acceptable choices.

The Wolfenstein parameter A. The A_1 (Higgs) taste sits along the democratic direction $\hat{h} = (1, 1, 1)/\sqrt{3}$ in the internal space. Each generation axis $\hat{e}_j$ makes an angle θ with $\hat{h}$ satisfying $\cos\theta = \hat{e}_j \cdot \hat{h} = 1/\sqrt{3}$. The CKM mixing in the A parameter is driven by the perpendicular component of the generation axis relative to the Higgs democratic direction:

$$A = \sin\theta = \sqrt{1 - 1/3} = \sqrt{2/3} = 0.8165.$$

Observed: $A = 0.826 \pm 0.012$. Match: 0.8σ, agreement at 1.2%. The observed value itself is less settled than the error bar suggests: different global fits place it between roughly 0.79 and 0.83, and the prediction sits inside that range. The geometric value is exact, but its two uses differ in status: in the reactor angle $\sin\theta_{13} = A^2\lambda$ it enters protected (the second-order correction cancels, so that comparison tests the geometry alone), while in $|V_{cb}| = A\lambda^2$ it enters through a second-order coefficient whose derivation is open, so the identification with the measured A is leading-order; the observed 1.2% deviation is smaller than the natural $\lambda^2 \approx 5\%$ correction scale, a suppression that is observed rather than derived.

The combination $|V_{cb}| = A\lambda^2 = \sqrt{2/3}/(2\pi^2) = 0.04136$ (observed: 0.0408 ± 0.0014, agreement at 0.4σ) provides a cross-check at the second-order level of the

Wolfenstein expansion. Two structural predictions (λ and A) generate a derived prediction for a third observable ($|V_{cb}|$), with all three matching observation.

A structural identity. A non-trivial representation-theoretic identity underlies the Wolfenstein A parameter. The off-democratic projection-squared $A^2 = 1 - (\hat{e}_j \cdot \hat{h})^2 = 1 - 1/d$ for the generation axes in a d-dimensional space, combined with the cubic-group quadratic Casimir for the vector representation $C_2(T_1) = d - 1$, gives the structural identity

$$A^2 = \frac{d-1}{d} = \frac{C_2(T_1)}{d}.$$

For $d = 3$ this gives $A^2 = 2/3$ directly. The same Casimir-over-dimension structure underlies the Koide prediction in §6.6 below and the PMNS angle $\sin^2\theta_{13}$ in §6.7. The identity is not a numerical coincidence; it is the structural source of multiple Layer-1 predictions in the framework.

The down-to-strange mass ratio. The Cabibbo derivation extends to mass ratios through the Gatto-Sartori-Tonin (GST) relation $\sin\theta_C \approx \sqrt{m_d/m_s}$, a standard relation in flavor physics with known higher-order corrections. The framework's prediction is structural in λ — the cubic-lattice derivation above — while the conversion to the mass ratio inherits GST's empirical status. To leading order:

$$\frac{m_d}{m_s} = \lambda^2 = \frac{1}{2\pi^2} = 0.05066.$$

Observed (FLAG 2024 $N_f = 2{+}1{+}1$ lattice): $m_d/m_s = 0.0503 \pm 0.0007$. Match: 0.56σ, agreement at 0.7%.

The Jarlskog invariant. The Jarlskog invariant $J = \mathrm{Im}(V_{us}V_{cb}V_{ub}^*V_{cs}^*)$ measures CP violation in the quark sector. In the Wolfenstein parametrization, $J \approx A^2\lambda^6\eta$. Substituting $\lambda = 1/(\pi\sqrt{2})$ and $A = \sqrt{2/3}$:

$$J = \frac{\eta}{12\pi^6},$$

where $\eta = 0.348$ is the solution-specific CP-violating parameter. This gives $J = 3.02 \times 10^{-5}$ (observed: $(3.08 \pm 0.13) \times 10^{-5}$, agreement at 0.5σ). The structural suppression factor $12\pi^6 \approx 11{,}537$ is purely geometric.

A consistency check using the framework's $A^2 = 2/3$ and $\lambda^2 = 1/(2\pi^2)$ structural inputs runs the relation in reverse: solving $J = A^2\lambda^6\eta$ for η at the observed Jarlskog value $J_{\mathrm{obs}} = 3.08 \times 10^{-5}$ gives $\eta_{\mathrm{implied}} = 0.355$. The PDG value is $\eta = 0.357 \pm 0.011$. The framework's structural inputs combined with the observed Jarlskog produce an η consistent with the PDG fit at 0.15σ. This consistency check is independent of the chapter's specific solution-specific $\eta = 0.348$ — it asks only whether the framework's structural A^2 and λ^2 inputs are compatible with the observed CP-violating phase, and the answer is yes within 1σ.

The Wolfenstein parameters ρ and η themselves require complex CKM entries, arising from the specific bijection φ rather than from structural cubic-group

representation theory. They are solution-specific properties of the framework's particular substratum representative — Layer 3 content in the four-layer framing of Chapter 4 — and the framework makes no commitment to their numerical values beyond the structural constraint that the Jarlskog invariant has the form $\eta/(12\pi^6)$.

6.6 The Koide relation and the lepton mass chain

The Koide relation is a numerical regularity among the three charged-lepton masses observed in 1981:

$$Q = \frac{m_e + m_\mu + m_\tau}{(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2} = \frac{2}{3}$$

to extraordinary precision — matching observation at 0.001% with no fitted parameters in the conventional Standard Model. The relation has been a persistent puzzle, with multiple proposed derivations from grand-unified, supersymmetric, and modular-symmetry frameworks. None has produced both the value 2/3 and an account of why the relation should hold in the first place. The framework derives the Koide relation from the cubic-group representation theory of the three generations, with both the value and the underlying structure following from substratum-level content.

The Koide angle from a cubic Casimir. The three lepton generations occupy the three triplet (T_1) staggered tastes, related by the cubic symmetry group O. The Koide relation in its angular form $\theta_0 = C_2/d^2$ — where C_2 is the quadratic Casimir of the relevant cubic-group representation and d is the spatial dimension — provides a specific structural prediction for the value θ_0.

For the T_1 representation of O (the three-dimensional vector representation occupied by the three lepton generations), the quadratic Casimir is $C_2(T_1) = d - 1 = 2$ in $d = 3$ spatial dimensions. The lattice bandwidth squared is $d^2 = 9$. The Koide angle in its natural framework form is therefore

$$\theta_0 = \frac{C_2(T_1)}{d^2} = \frac{2}{9},$$

which is the phase of the standard $\mathbb{Z}_3$-symmetric parametrization of three masses, independent of the amplitude that $Q = 2/3$ fixes (SM §7.2). The framework's prediction matches the observed value $\theta_0^{\text{obs}} = 0.222225 \pm 0.000006$ at 0.001% (0.4σ).

The structural interpretation is geometric: $C_2(T_1)/d^2$ is the anisotropy strength of the generation representation, normalized by the lattice bandwidth squared. The cubic-group representation theory of T_1 fixes the numerator; the dimensional structure of the lattice fixes the denominator. The value 2/9 follows from these two pieces of substratum-level structure, with no fitted parameters.

Uniqueness among alternative ratios. Among dimensionless ratios involving cubic-group Casimirs and lattice-dimension quantities at $d = 3$ and T_1, the

value $\theta_0 = C_2/d^2 = 2/9$ is the unique match to observation. Ten alternative ratios — including C_2/d (which gives the Wolfenstein A^2 above), $C_2/(2d)$, $\sqrt{C_2}/d$, $(C_2 - 1)/d^2$, and similar combinations — produce values discrepant from the observed Koide angle by amounts ranging from 11% to over 200%. The match at $\theta_0 = C_2/d^2$ is not one of several possibilities consistent with cubic-group structure; it is the unique value forced by the relevant Casimir-and-bandwidth combination.

Connection to the Cabibbo identity. The Koide angle is related to the Cabibbo structural identity through second-order perturbation theory. The natural cubic-lattice perturbation amplitude on the T_1 representation is $\sqrt{C_2}/d = \sqrt{2}/3$ — the Casimir's square root scaled by the bandwidth. Second-order perturbation theory on the eigenvalues then gives $\theta_0 \sim (\sqrt{C_2}/d)^2 = C_2/d^2 = A^2/d$, with $A^2 = (d-1)/d$ the structural identity of §6.5 for vector representations. Amplitudes scale as $\sqrt{C_2}/d$; eigenvalue shifts scale as the square. The Koide angle and the Wolfenstein A parameter share a common Casimir structure, with the Koide angle obtaining at the eigenvalue level what A obtains at the amplitude level.

The first-principles derivation of the specific $1/d$ amplitude scaling (rather than alternatives like $1/(2d)$) from the cubic-lattice Yukawa structure is complete: the amplitude prefactor has been computed and equals one (SM §7.2; driver `oi_lattice_code/fermion/n_ratio_pt.py`), so $\mathcal{N} = 1/d^2 = 1/9$ exactly at $m \to 0$, and the empirical match at 0.001% confirms a derived normalization rather than constraining a chosen one. The $1/d$ amplitude is therefore fixed by the framework's structural commitments rather than chosen empirically, conditional on the substrate-eigenvalue criterion that selects a loop without a time direction (below). The Koide prediction is Class B-C, Layer 1, because the Koide amplitude $A^2 = C_2$ is not yet derived from the dynamics (SM §7.2). The choice reduces to whether the taste-mixing loop carries a time direction ($d = 4$, giving 1/16) or not ($d = 3$, giving 1/9); the $d = 3$ substrate-eigenvalue origin of the generation masses selects 1/9; Schur's lemma on the T_1 generations protects this against all taste-blind emergent dressing. A cubic-invariant dressing need not be taste-blind: it may act on $\mathrm{Sym}^2(T_1)$ with different scalars on A_1, E and T_2, as the one-loop staggered taste kernel does ($\lambda_E/\lambda_{A_1} \approx 0.22$–$0.24$). That class is bounded: it leaves θ_0 invariant at the $\sqrt{m}$ level, and on the mass pattern the observed agreement constrains it to $-2.5 \times 10^{-6} \lesssim \delta m/m \lesssim 1.7 \times 10^{-6}$ at 95% confidence; the interval is conditional on the linear response of θ_0 to the dressing (sensitivity coefficient $\kappa \approx -35$, with λ_E/λ_{A_1} in the range above) and is not a model-independent exclusion; a lepton-Yukawa loop of order 10^{-6} is compatible with that interval for either sign, by a factor of about two, but is not a confirmed prediction (SM §7.2). What remains is a possible loop-level cubic-breaking contribution — the same residual as the emergent-Lorentz radiative-stability question (SM §7.3).

The charged-lepton mass chain. Given the Koide angle $\theta_0 = 2/9$ and one empirical input (the tau mass $m_\tau = 1776.93$ MeV), the parametrization

$\sqrt{m_k} = \mu(1 + A\cos(\theta_0 + 2\pi k/3))$ is fully determined: $Q = 2/3$ fixes $A = \sqrt{2}$, θ_0 fixes the phase, and m_τ fixes the scale μ. The two structural inputs are $Q = 2/3$ and $\theta_0 = 2/9$, and the resulting masses are $m_e = 0.51098$ MeV and $m_\mu = 105.656$ MeV against 0.51100 and 105.658 observed (0.003% and 0.002%; SM §7.2). The ratio $m_e/m_\tau = 2.876 \times 10^{-4}$ lies within 0.84% of

$$\frac{m_e}{m_\tau} = \frac{\lambda^4}{d^2} = \frac{1}{36\pi^4} \approx 2.852 \times 10^{-4},$$

with $\lambda = 1/(\pi\sqrt{2})$ the Cabibbo angle from the quark sector (§6.5) and $d = 3$ the spatial dimensionality (Chapter 5 §5.2). The proximity plays no role in the chain: the mass point is fixed by Q and θ_0. The assignment of m_e specifically to the lightest charged-lepton mass eigenvalue is Layer 2(b) bijection-specific by the gauge-invariance constraint of Chapter 2 §2.5.

Combined with $Q = 2/3$ and the empirical m_τ, the second structural input $\theta_0 = 2/9$ determines all three charged-lepton masses:

Mass	Formula	Predicted	Observed	Deviation
m_e	Koide$(Q, \theta_0; m_\tau)$	0.51098 MeV	0.51100	0.003%
m_μ	Koide$(Q, \theta_0; m_\tau)$	105.656 MeV	105.658	0.002%

Both predictions match observation to better than 0.01%.

The cross-sector pattern is significant: the lepton mass spectrum inherits *both* the Wolfenstein parameter $A^2 = 2/3$ (via the PMNS Condition 2 sum rule of Ch08 §8.5) *and* the Cabibbo coupling $\lambda^2 = 1/(2\pi^2)$ (via the PMNS reactor angle $\sin^2\theta_{13}$) from the quark sector. The lepton-sector structure is not freshly postulated; it inherits from the same substrate machinery that produces the quark-sector observables.

The classification per Chapter 4 §4.5 is that the electron and muon masses are Class B-M (mass-chain inheritance from a single upstream empirical input m_τ plus structural commitment); the Koide relation $Q = \frac{1}{3} + \frac{A^2}{6} = 2/3$ is Class B-C Layer 1 from the cubic-group Casimir through the amplitude $A^2 = C_2(T_1) = 2$; the phase input $\theta_0 = C_2(T_1)/d^2 = 2/9$ is Class B-C Layer 1 from the same Casimir structure, with the amplitude prefactor computed equal to one (SM §7.2); both are conditional on the dynamical closure of the Koide amplitude, whose value $A^2 = C_2$ is not yet derived from the dynamics.

The magnitude content of the lepton chain is Layer 1, conditional on the same amplitude closure: both inputs come from the cubic Casimir, the phase input with computed unit prefactor (SM §7.2). The analogous coefficient of Condition 2 in Ch08 §8.5 is Layer 2(b): the second-order substrate route is excluded (SM §7.3), the 7/6 sum rule is the direct experimental test, and the $O(\lambda^3)$ interference route is recorded, not pursued; there the framework provides a structural

identification with cross-sector content (quark-sector inputs determining lepton-sector outputs through cubic-group machinery), and the percent-level empirical match is consistent with the structural identification modulo renormalization-group running. The directional assignment (which specific generation is e, μ, or τ, and correspondingly which eigenvalue carries which label) remains Layer 2(b) bijection-specific by the gauge structure of Chapter 2 §2.5 and is not derivable from substrate alone in any framework consistent with $\mathcal{G}_{\text{sub}}$-invariance.

The full six-mass chain. The structural relations link the six lighter charged fermion masses through one empirical mass scale (m_s) plus two further structural inputs that the chapter makes explicit below:

$$m_s \xrightarrow{\lambda^2} m_d \xrightarrow{\sqrt{\theta_0}} m_u \xrightarrow{Q_{\text{down}}} m_b \xrightarrow{Z_S} m_\tau \xrightarrow{\theta_0} m_e, m_\mu.$$

The five arrows are structural relations of three different epistemic statuses, which the chapter distinguishes explicitly:

Cubic-group derivations (parameter-free forms): the Cabibbo-derived $m_d/m_s = \lambda^2$ from the inter-generation Brillouin-zone distance squared; the $m_u/m_d = \sqrt{\theta_0}$ from the intra-doublet splitting at the amplitude level (Layer 1 form with Layer 2(b) alignment, SM §7.2); the charged-lepton Koide $Q = \frac{1}{3} + \frac{A^2}{6} = 2/3$ through the amplitude $A^2 = C_2(T_1) = 2$; and the Koide phase $\theta_0 = 2/9 = C_2(T_1)/d^2$ with computed unit prefactor (SM §7.2). These four relations take their forms from cubic-group representation theory with no free parameters; their classifications are those of the Section 6.8 table, with Koide Q and θ_0 conditional on the dynamical closure of the Koide amplitude.

Derived running factor plus measured non-perturbative renormalization: the down-to-tau ratio combines two components of different character. The factor 4.28 is derived: tree-level taste-independence gives $y_b = y_\tau$, and full Standard Model RGE running from a common M_{Pl} boundary condition gives $m_b/m_\tau|_{\text{tree}} = 4.276$, with a $\sim 1\%$ systematic from higher-loop corrections and electroweak-threshold matching (SM §7.5). The factor Z_S is measured: the scalar-density renormalization $Z_S = \langle\bar{\psi}\psi\rangle_{\text{int}}/\langle\bar{\psi}\psi\rangle_{\text{free}}$, computed non-perturbatively on SU(3) backgrounds at the matching scale $m_{\text{match}} = \lambda g_0^2$, whose dimensional form is forced by power-counting in the taste-decomposed Coleman–Weinberg potential. At $L = 32$, $Z_S(\lambda g_0^2) = 1.813 \pm 0.001$ (statistical), with the $L \to \infty$ extrapolation ≈ 1.83 and a realistic 1–2% systematic from finite volume and the regulator choice. The prediction $m_b/m_\tau = 4.28/Z_S = 2.361$ matches the observed 2.352 ± 0.017 at 0.4% (0.5σ; $\sim 0.7\sigma$ at $L \to \infty$). The open Layer-2 element is the scalar prefactor in the matching-scale identification (the K coefficient): the Monte Carlo consistency check gives $m_{\text{match}}^{\text{MC}} = 0.1234 \pm 0.0002$ against $\lambda g_0^2 = 0.1217$ (1.4%), and $K = 1$ is excluded by the same data (SM §7.5).

Phenomenological auxiliary parameter (Class P): the down-sector Koide invariant $Q_{\text{down}} \approx 0.7304$ does not match any simple cubic-group form (the closest

cubic-group ratios — 11/15, 5/7, $\theta_0 \cdot (3/2) \cdot (1+\lambda^2)$ — all deviate from the empirical value by more than 1%). The framework treats Q_{down} as a phenomenological input classified as Class P per Chapter 4 §4.5, not as a structural prediction: $Q_{\text{down}} \approx 0.7304$ is taken as phenomenological input in the form $(2/3)(1+\alpha_s/\pi)$, which reproduces the PDG mixed-scheme value to 0.15% but has no derivation (SM §7.2); m_b then follows at 0.9%. Whether Q_{down} has a derivable structural form different from the charged-lepton sector's 2/3 — a magnitude-level question potentially closable through lattice perturbation theory in the emergent description — is open. The sector-identification content (which Q values attach to down-quarks vs charged-leptons) is bijection-specific per Chapter 2 §2.5.

Mass	Formula	Class	Predicted	Observed	Match
m_d	$m_s/(2\pi^2)$	Layer 1	4.73 MeV	4.67 ± 0.48	1.3%
m_u	$m_d \times \sqrt{2/9}$	L (Layer 1 form; 2(b) alignment)	2.20 MeV	2.16 ± 0.49	0.1σ
m_b	Koide(Q_{down})	L (Q_{down} a Class P input)	4144 MeV	4180 ± 30	0.9%
m_τ	$m_b \times Z_S/4.28$	L (Z_S measured; 2(a): K)	1755 MeV	1776.9	1.2%
m_e	Koide(Q, θ_0; m_τ)	Layer 1	0.505 MeV	0.511	1.2%
m_μ	Koide(Q, θ_0; m_τ)	Layer 1	104.4 MeV	105.66	1.2%

The chain produces six masses from one empirical scale (m_s) plus two further inputs (Q_{down} as a Class P phenomenological input, Z_S measured nonperturbatively). The four cubic-group relations carry no fitted parameter; Q_{down} is a phenomenological input. Four out of five mass-chain transitions are therefore free of phenomenological input, with the down-sector Q_{down} closure carrying the one.

The top quark mass. The top quark sits separately from the chain above because its Yukawa coupling is large. The weak-scale relation $m_t = v/\sqrt{2}$ with $y_t(m_t) \approx 1$ reproduces the observed mass:

$$m_t = \frac{v}{\sqrt{2}} = \frac{246\,\text{GeV}}{\sqrt{2}} = 174.1\,\text{GeV},$$

where v is the Higgs vacuum expectation value, matching the observed 172.5±0.3 GeV to 0.9%.

The mechanism behind this relation is open, and the honest classification is a retrodiction rather than a structural prediction. The natural proposal — a

compositeness-scale normalization $y_t(M_{\rm Pl}) = 1$ carried to low energy by the infrared quasi-fixed point — does not deliver the observed value under one-loop Standard Model running: $y_t(M_{\rm Pl}) = 1$ flows to $y_t(m_t) \approx 1.19$, giving $m_t \approx 207$ GeV, and the IR quasi-fixed point itself sits at $y_t \approx 1.25$ ($m_t \approx 215$–220 GeV), not at unity. The attractor basin $y_{\rm UV} \in [0.5, 5]$ focuses to $m_t \in [177, 221]$ GeV, a 25% spread. What is empirically exact is the weak-scale statement $y_t(m_t) \approx 1$; the compositeness-scale normalization plus IR attraction does not produce it, and the observed top corresponds to $y_t(M_{\rm Pl}) \approx 0.4$–$0.5$. The relation is therefore recorded as a retrodiction of the weak-scale $y_t \approx 1$ coincidence, with the structural mechanism an open task; a derivation of $y_t = 1$ at the electroweak matching scale, or of $y_t(M_{\rm Pl}) \approx 1/2$ (which runs to ≈ 177 GeV), would close it.

The relation $m_t = v/\sqrt{2}$ matches the observation to 0.9%, but as a retrodiction of the weak-scale $y_t \approx 1$ coincidence rather than a parameter-free structural prediction; a derivation of $y_t = 1$ at the electroweak matching scale, or of $y_t(M_{\rm Pl}) \approx 1/2$ (which runs to ≈ 177 GeV), would close the mechanism.

6.7 PMNS mixing angles

The PMNS matrix parametrizes the misalignment between neutrino mass eigenstates and flavor eigenstates, analogous to the CKM matrix in the quark sector but with three angles $(\theta_{12}, \theta_{23}, \theta_{13})$ governing the dominant mixing structure. In the Standard Model the three angles are independent free parameters. The framework derives all three from cubic-group representation theory combined with parity-graded charged-lepton corrections at the Cabibbo scale, with the post-JUNO measurement of $\sin^2\theta_{12}$ matched by one parameter-free value at 0.07σ (a retrodiction; see Chapter 8 provenance).

Tribimaximal mixing as the leading structure. The starting point is tribimaximal (TBM) mixing, derived from the alternating subgroup $A_4 \subset O$ of the cubic rotation group. TBM is the specific pattern

$$\sin^2\theta_{12} = 1/3, \qquad \sin^2\theta_{23} = 1/2, \qquad \sin^2\theta_{13} = 0,$$

historically introduced as an empirical observation about the leptonic mixing pattern and later given group-theoretic foundations. In the framework, TBM emerges as the zeroth-order PMNS structure from the A_4 subgroup of the cubic rotation group acting on the three lepton generations — the same cubic group whose decomposition $T_1 \oplus E \oplus A_1$ produced the Standard Model gauge group in Chapter 5.

The TBM structure is not the final answer: the experimentally measured $\sin^2\theta_{13} \approx 0.022$ is non-zero, and $\sin^2\theta_{12}$ and $\sin^2\theta_{23}$ deviate from their TBM values by amounts consistent with corrections of order $\lambda^2 = 1/(2\pi^2) \approx 0.05$. The framework's content is the structural derivation of these corrections from the charged-lepton sector.

The three structural ingredients. The deviations from TBM are controlled by three structural ingredients that the framework derives rather than postulates.

First, the overall correction scale is the squared Cabibbo angle $\lambda^2 = 1/(2\pi^2)$ from §6.5. The PMNS matrix's deviation from TBM follows from the charged-lepton sector's perturbation away from the basis in which TBM holds exactly; the charged-lepton perturbation has natural amplitude λ, with second-order effects entering at λ^2.

Second, the charged-lepton perturbation has a specific structure: it is U-parity odd under μ-τ exchange. The leading-order generator A_1 in the perturbative expansion $V_\ell = \exp(\lambda A_1 + \lambda^2 A_2 + O(\lambda^3))$ satisfies $UA_1U^{-1} = -A_1$ under the μ-τ exchange generator of $S_4 \supset A_4$. Geometrically, A_1 is a rotation around the symmetric combination $(e_2+e_3)/\sqrt{2}$ by angle $A^2 = 2/3$ — with the Wolfenstein A of §6.5 entering directly as the rotation angle. The structural property is that this A_1 produces no $O(\lambda)$ corrections to the TBM angles θ_{12} and θ_{23} (the corrections are pushed to $O(\lambda^2)$), while producing the leading reactor angle $\sin\theta_{13} = A^2\lambda$ at order λ.

Third, the ratio of corrections satisfies $\Delta_{13}/\Delta_{23} = A^4 = 4/9$ from the same Wolfenstein $A = \sqrt{2/3}$ via a second-order projection onto the Higgs democratic direction. The same structural Casimir-over-dimension identity $A^2 = C_2(T_1)/d = (d-1)/d$ that produced the Koide angle in §6.6 now produces the PMNS ratio.

The three predicted angles. The three ingredients combine to give the three angles:

$$\sin^2\theta_{12} = \frac{1}{3} - \frac{1}{4\pi^2} = 0.3080,$$

$$\sin^2\theta_{23} = \frac{1}{2} + \frac{1}{2\pi^2} = 0.5507,$$

$$\sin^2\theta_{13} = \frac{4}{9}\cdot\frac{1}{2\pi^2} = \frac{4}{18\pi^2} = 0.02252.$$

These can be compared with the experimental values from the post-JUNO global fit and from NuFIT 6.0 (normal mass ordering).

Angle	Predicted	Observed	Match
$\sin^2\theta_{12}$	0.3080	0.3085 ± 0.0073 (post-JUNO global fit)	0.07σ
$\sin^2\theta_{23}$	0.5507	$0.561^{+0.012}_{-0.015}$ (NuFIT 6.0 NO)	0.74σ
$\sin^2\theta_{13}$	0.02252	0.02195 ± 0.00056 (NuFIT 6.0 NO)	1.01σ

All three angles match observation within about 1σ. The match for $\sin^2\theta_{12}$ at 0.07σ is unusually close, made tighter by the JUNO measurement reported in November 2025.

The JUNO measurement. The Jiangmen Underground Neutrino Observatory reported its first measurement of reactor neutrino oscillations in November 2025, achieving the world's most precise determination of $\sin^2\theta_{12}$: the value 0.3092 ± 0.0087, a factor of 1.6 improvement over all previous measurements combined. The framework's prediction $1/3 - 1/(4\pi^2) = 0.3080$ matches the JUNO measurement directly at 0.14σ. The updated global fit incorporating JUNO data gives $\sin^2\theta_{12} = 0.3085 \pm 0.0073$, matching the prediction at 0.07σ. As JUNO accumulates data over its thirty-year design lifetime, the error bar is projected to reach ± 0.0014, testing the prediction at sub-percent precision.

The structural status of the three predictions deserves careful articulation. The reactor angle $\sin^2\theta_{13}$ has the cleanest derivation: it depends only on the leading-order charged-lepton generator A_1 at second order in λ, and is consequently Layer 1 structural — independent of any second-order Wilson coefficient, and conditional only through λ on the emergent dressing of the Cabibbo identification. The structural identity $\sin\theta_{13} = A^2\lambda = (C_2(T_1)/d)\cdot\lambda$ is exact at order λ^2. This is the cleanest of the three PMNS predictions.

The solar and atmospheric angles $\sin^2\theta_{12}$ and $\sin^2\theta_{23}$ depend on a structural relation among second-order Wilson coefficients b_{12}, b_{13}, b_{23} that arises from the charged-lepton perturbation's A_2 structure. The relation reduces to $b_{23} = (4/3)(b_{12} + b_{13})$, which has the geometric reading $4/3 = 2A^2 = 2(d-1)/d$ — partner-count times per-partner projection in the cubic-group representation theory. Under this relation, the two angles satisfy the joint sum rule $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$. Observed: 1.178 ± 0.020 using the post-JUNO global fit and NuFIT 6.0 atmospheric data; agreement at 0.6σ. This depends on the atmospheric octant: the quoted value uses the upper-octant best fit $\sin^2\theta_{23} \approx 0.561$; under the lower-octant solution (≈ 0.455) the combination is ≈ 1.07, roughly 5σ below $7/6$.

The structural status of $b_{23}/(b_{12}+b_{13}) = 4/3$ separates into two distinct layers. The cubic-group identity $4/3 = 2A^2 = C_2(T_1)^2/d$ that fixes the *magnitude* is Layer 2(b): the second-order substrate route is excluded (SM §7.3), the 7/6 sum rule is the direct experimental test, and the $O(\lambda^3)$ interference route is recorded, not pursued (per Ch02 §2.5, no Lagrangian exists at the substratum proper). The *directional* assignment — that b_{23} specifically is the larger coefficient rather than $b_{12}+b_{13}$, with the labeling fixed by the empirically observed mass hierarchy ($m_e \ll m_\mu \ll m_\tau$ pinning generation 1 = electron) — is Layer 2(b) bijection-specific content by the gauge-invariance argument of Chapter 2 §2.5: under state-relabeling gauge $\sigma \in \mathcal{G}_{\text{sub}}$, the function $b_{23}/(b_{12}+b_{13})$ transforms non-trivially, so no $\mathcal{G}_{\text{sub}}$-invariant substrate computation can fix the directional assignment. The classification per Chapter 4 §4.5 is therefore that the two angles are Class B-L (layered conditional), with the Layer 2(b) magnitude content a solution-specific input tested directly by the 7/6 sum rule and the Layer 2(b) directional

content classified as bijection-specific input rather than derivational gap.

The framework's prediction structure compared to alternatives. The $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$ sum rule is the framework's distinguishing structural prediction in the PMNS sector. The TM1 and TM2 partial-TBM patterns — alternative deviation structures from TBM also derivable from cubic-group representation theory — preserve specific columns or rows of the TBM matrix but do not produce this exact sum rule. The empirical match to the sum rule therefore distinguishes the framework's specific cubic-symmetric perturbation from the TM1/TM2 alternatives at the sub-half-percent level on the sum-rule precision. Chapter 8 develops the JUNO prediction and its discrimination from these alternatives in detail.

The framework's lepton-sector content has more structural commitments than the three angles. The sum rule itself is a falsifiable consequence of the cubic-symmetric perturbation. On the branch where the singlet taste hosts the Higgs and no gauge-singlet fermion is present (SM §8.4; the neutrino mass mechanism is otherwise open, SM §4.7.1), the neutrinos are Majorana rather than Dirac, and the mass ordering is expected to be normal rather than inverted as a heuristic consequence of the taste-breaking structure of the neutrino mass matrix developed in Chapter 5 §5.6, not as a theorem. Each of these is a pre-registered falsification condition per Chapter 4 §4.5: confirmed inverted ordering would refute the taste-breaking heuristic on that branch; a confirmed Dirac nature would refute the branch's expectation that no right-handed neutrino exists in the spectrum.

The Higgs mass. A separate structural prediction completes the chapter's quantitative content: the Higgs quartic coupling at the compositeness scale is exactly zero. The Higgs is the singlet (A_1) taste of the staggered decomposition — a composite scalar in the framework's structural account, as Chapter 5 §5.5 developed — and its quartic self-coupling at the Planck scale receives no tree-level contribution. The Standard Model quartic λ at low energies is generated entirely by RGE running between the compositeness scale and the experimental scale.

The structural prediction is $\lambda(M_{\rm Pl}) = 0$. The observed value from Buttazzo et al. is $\lambda(M_{\rm Pl}) = -0.013 \pm 0.020$, consistent with zero at 0.6σ. The structural commitment is therefore consistent with the extrapolated value at the boundary-condition level.

The corresponding prediction for the Higgs mass in GeV is derived by combining the structural boundary condition with the measured top mass m_t as a one-input parameter and running the three-loop SM RGE. The result is $m_H \approx$ 129-132 GeV for $m_t =$ 172-173 GeV, against the observed $m_H = 125.10 \pm 0.14$ GeV. The gap is sensitive to m_t (with $\partial m_H/\partial m_t \approx -1.1$ GeV/GeV), and the recent CMS measurement $m_t = 170.5 \pm 0.8$ GeV would bring the prediction to about 128 ± 2 GeV. The structural claim $\lambda(M_{\rm Pl}) = 0$ is confirmed; the specific Higgs mass in GeV is consistent with observation within the current m_t uncertainty.

The framework's Higgs-sector prediction has a notable convergence with the broader physics literature. The Multiple Point Principle developed by Bennett, Froggatt, and Nielsen in the 1990s postulated $\lambda(M_{\mathrm{Pl}}) = 0$ as a fine-tuning condition, predicting the Higgs mass at approximately the value subsequently observed. Wetterich's asymptotic-safety analysis showed that if the Standard Model holds to the Planck scale with an asymptotically safe gravity contribution, the observed Higgs mass requires $m_t \approx 171$ GeV. The framework's structural derivation of $\lambda(M_{\mathrm{Pl}}) = 0$ thus provides a substratum-level account of what an active beyond-SM literature has been postulating as a boundary condition.

6.8 Summary table and chapter close

The chapter has developed twenty-two Standard Model retrodictions from the cubic-lattice substratum, with the structural status of each made explicit per the methodological discipline of Chapter 4. The summary table collects the predictions with two complementary classification axes: the **Class** column gives the *derivational layer* per Chapter 4 §4.5 (S/C/L/R/E/M), and the **Cat** column gives the *independent-evidence status* per the categorization developed below (A/B/C/D/E).

Observable	Source	Predicted	Observed	Match	Class	Cat
$1/\alpha_0$	one-loop staggered VP	23.25	—	structural input	**S**	input
$\alpha_3(M_Z)$	RG running with (δ_0, A, B)	0.118	0.1181 ± 0.0011	fit	**R** [a]	E
$\alpha_2(M_Z)$	RG running with (δ_0, A, B)	0.0339	0.03379 ± 0.00009	fit	**R** [a]	E
δ_0	U(1) row fit	10.0	—	empirical input	**E**	input
$A \cdot B$ cross-check	finite-lattice computation	≈ 48–62 (leading-power-dependent)	46.4 (from fit)	inconclusive	**C** (leading power)	—
$\bar{\theta}$	T-invariance + detailed balance	not fixed ($\theta \in \{0, \pi\}$; H-top, H-det)	$\lvert\bar{\theta}\rvert < 10^{-10}$	open	—	—

Observable	Source	Predicted	Observed	Match	Class	Cat
Cabibbo λ	inter-generation BZ distance	0.2251	0.2250 ± 0.0007	0.04%	**C**	**A**
Wolfenstein A	off-democratic projection	$\sqrt{2/3} = 0.8165$	0.826 ± 0.012	1.2%	**S**	**A**
$\lvert V_{cb}\rvert$	$A\lambda^2$	0.0414	0.0408 ± 0.0014	0.4σ	**C**	**A**
m_d/m_s	GST $\lambda^2 = 1/(2\pi^2)$	0.0507	0.0503 ± 0.0007	0.7%	**L**	**A**
Jarlskog J	$\eta/(12\pi^6)$	3.02×10^{-5}	$(3.08 \pm 0.13) \times 10^{-5}$	0.5σ	**E** (η)	**B**
Koide Q	cubic T_1 structure	2/3	0.666664	0.0003%	**C**	**A**
Koide angle θ_0	$C_2(T_1)/d^2$	2/9	0.222225	0.001%	**C**	**A**
m_e	Koide(Q, θ_0, m_τ)	0.51098 MeV	0.51100 MeV	0.003%	**M** (m_τ)	**B**
m_μ	Koide(Q, θ_0, m_τ)	105.66 MeV	105.658 MeV	0.002%	**M** (m_τ)	**B**
m_d	$m_s/(2\pi^2)$	4.73 MeV	4.67 ± 0.48	1.3%	**M** (m_s)	C
m_u	$m_d\sqrt{2/9}$	2.20 MeV	2.16 ± 0.49	0.1σ	**L** [e]	**B**
m_b	Koide (Q_{down})	4144 MeV	4180 ± 30	0.9%	**L** [d]	E
m_t	$v/\sqrt{2}$ (weak-scale $y_t \approx 1$; mechanism open)	174.1 GeV	172.5 ± 0.3	0.9%	**R**	D
$\sin^2\theta_{12}$	$1/3 - 1/(4\pi^2)$	0.3080	0.3085 ± 0.0073	0.07σ	**L** [f]	**B**
$\sin^2\theta_{23}$	$1/2 + 1/(2\pi^2)$	0.5507	$0.561^{+0.012}_{-0.015}$	0.74σ	**L** [f]	**B**
$\sin^2\theta_{13}$	$A^4\lambda^2 = 4/(18\pi^2)$	0.02252	0.02195 ± 0.00056	1.01σ	**C**	**A**

Observable	Source	Predicted	Observed	Match	Class	Cat
$\lambda(M_{\rm Pl})$	composite-Higgs structure	0	-0.013 ± 0.020	0.6σ	**S**	**A**
m_H	RGE running from $\lambda(M_{\rm Pl}) = 0$	129-132 GeV (band)	125.10 ± 0.14	m_t-consistent	**M** (m_t)	D

Footnotes: **[a]** The three gauge couplings at M_Z are reproduced with one empirical input (δ_0) and two fitted parameters (A, B); the structural predictions in this sector are $1/\alpha_0$ and the universality of δ_0; the $A \cdot B$ numerical cross-check is leading-power-conditional and inconclusive across finite-size models. **[d]** $Q_{\rm down}$ is a phenomenological input (Class P) in the form $(2/3)(1 + \alpha_s/\pi)$, with no first-principles derivation (SM §7.2; §6.6 mass-chain discussion); m_b then follows at 0.9%. The bottom-to-tau bridge uses the RGE-derived 4.28 and the MC-measured $Z_S(\lambda g_0^2)$ (SM §7.5). **[e]** Layer 1 form + Layer 2(b) alignment of SU(2) doublet structure with cubic T_1 perturbation. **[f]** Layer 1 form + Layer 2(b) magnitude $b_{23}/(b_{12} + b_{13}) = 4/3 = 2A^2$ (the second-order substrate route excluded, SM §7.3; the 7/6 sum rule the direct test) + Layer 2(b) directional assignment (which pair-ratio gets the 4/3, fixed by mass-hierarchy bijection per Ch02 §2.5 and Ch08 §8.5).

Parameter accounting — two complementary classifications. The table above shows two classification axes for each prediction. The **Class** column (S/C/L/R/E/M) reflects the *derivational layer* per Chapter 4 §4.5 (S = parameter-free structural: no fitted input and no open link specific to the entry; C = conditional; L = layered; R = retrodiction; E = empirical-input-using; M = mass-chain). The **Cat** column reflects the *independent-evidence status* — whether a prediction is independent empirical evidence for the framework's structural inputs or a consequence of those inputs already counted elsewhere.

- **Category A — independent empirical tests of structural inputs:** Cabibbo λ, Wolfenstein A, $|V_{cb}| = A\lambda^2$, $m_d/m_s = \lambda^2$, $\sin^2\theta_{13} = A^4\lambda^2$, Koide $Q = 2/3$, Koide $\theta_0 = 2/9$ (independent of Q: the amplitude and the phase of the same parametrization, SM §7.2), and $\lambda(M_{\rm Pl}) = 0$ from composite-Higgs structure. These eight relations are tested through five empirical channels: CKM phenomenology (λ, $|V_{cb}|$), lattice QCD (m_d/m_s), PMNS ($\sin^2\theta_{13}$), lepton-mass dynamics (Koide Q and θ_0), and the Higgs sector ($\lambda(M_{\rm Pl})$). This category describes evidence independence, not an unconditional derivational class: m_d/m_s retains Class L because its conversion from the structural Cabibbo scale uses the empirical Gatto–Sartori–Tonin relation. Several entries combine the structural inputs A

and λ while using independent measurements: $|V_{cb}|$ is measured independently of V_{us}; lattice QCD measures m_d/m_s independently of the Cabibbo angle; and the observed m_H and m_t determine $\lambda(M_{\rm Pl})$ through RG running. The specific $\sin^2\theta_{13} = A^4\lambda^2$ form remains discriminating: $A^2\lambda^2$, $A\lambda^2$, and $A^4\lambda^4$ differ from observation by 32%, 50%, and 95%, respectively.

- **Category B — parameter-free predictions conditional on empirical mass-hierarchy bijection:** $\sin^2\theta_{12}$, $\sin^2\theta_{23}$, m_e and m_μ via Koide closure $(Q,\ \theta_0)$, m_u via intra-doublet $\sqrt{\theta_0}$ splitting, Jarlskog J via $\eta/(12\pi^6)$. Six predictions whose structural forms are tightly identified at the cubic-group magnitude level but whose specific assignment to labeled observables (which generation is electron, which is the second-lightest, etc.) is Layer 2(b) bijection-specific per Chapter 2 §2.5. Closure of the magnitude content depends on lattice perturbation theory in the emergent post-trace-out description; closure of the directional/assignment content is gauge-forbidden under $\mathcal{G}_{\rm sub}$-invariance and properly classified as empirical input rather than derivational gap. The framework's structural forms match observation at sub-percent or sub-sigma precision; the implied-η consistency check for Jarlskog (§6.5) gives $\eta = 0.355$ from the framework's A^2 and λ^2 inputs combined with observed J, within 0.15σ of PDG $\eta = 0.357 \pm 0.011$ — the structural $12\pi^6$ suppression is consistent with observation.
- **Category C — composite of Category A inputs with redundant content:** $m_d = m_s/(2\pi^2)$ — same prediction as $m_d/m_s = \lambda^2$ with m_s plugged in; an internal-consistency check rather than independent test.
- **Category D — band predictions at order-of-magnitude precision:** m_t (weak-scale relation $v/\sqrt{2}$ at 0.9%; fixed-point mechanism open — §6.7); m_H (RG-running band 129-132 GeV from $\lambda(M_{\rm Pl}) = 0$, observation 125.10 ± 0.14, 4-7 GeV below the band).
- **Category E — fitted using multiple empirical parameters:** $\alpha_3(M_Z)$, $\alpha_2(M_Z)$ (fitted via (δ_0, A, B), with structural content concentrated in the *form* of the fit rather than the matching values); m_b (uses Class P auxiliary parameter $Q_{\rm down}$ — see footnote [d]).
- **Input rows:** $1/\alpha_0$ and δ_0 are inputs to other predictions rather than themselves predicted observables; δ_0 is empirical. The $A \cdot B$ cross-check is inconclusive, and $\bar{\theta}$ remains open; neither is counted in Categories A–E.

The table has twenty-four entries: eight Category A rows, one for each relation, six in Category B, one in Category C, two in Category D, three in Category E, two input rows ($1/\alpha_0$ and δ_0), one inconclusive numerical cross-check ($A \cdot B$), and one open quantity ($\bar{\theta}$). The Cat C label means composite-redundant and is distinct from the derivational Class C. Beyond the table, the three-generation count (Appendix B.6c) and the absence of grand-unified gauge structure (Ap-

pendix B.6b) are further falsifiable structural commitments, without numerical values to compare against observation.

The framework's content is concentrated at Layer 0 and Layer 1 of the four-layer framing — gauge structure and pure cubic-group representation theory — with Layer 2 and Layer 3 content (bijection-specific operator structures and the overall mass scale) appearing as explicit inputs rather than fitted parameters. The structural-vs-bijection-level distinction developed in Chapter 4 §4.5 is what makes the empirical record evaluable: the framework predicts the *angular* and *ratio* structure of fermion mass relations from cubic-group representation theory, but the overall mass scale of each chain is set by the specific bijection and is taken as one empirical input.

The Category A relations span five empirical channels: CKM phenomenology, lattice QCD, PMNS mixing, lepton-mass relations, and the Higgs sector. The structural inputs $A^2 = 2/3$, $\lambda^2 = 1/(2\pi^2)$, and $C_2(T_1)/d^2 = 2/9$ are tested through those channels with the derivational conditions stated in the table. The gauge-coupling product $A \cdot B$ remains a separate, leading-power-conditional numerical cross-check and supplies no settled empirical match. The two formal uniqueness theorems (three-generation count via B.6c and no-GUT structural via B.6b) concern the gauge structure and matter content.

Falsification commitments revisited. The pre-registered falsification conditions from Chapter 4 §4.5 connect directly to the predictions developed here. Any stable fourth-generation fermion would refute the taste-reduction derivation that locks the generation count at three (formal uniqueness theorem B.6c). Any stable beyond-Standard-Model gauge structure would refute the cubic-group commutant derivation (formal uniqueness theorem B.6b). Any precision-upgrade measurement moving the Cabibbo angle outside 3σ of $1/(\pi\sqrt{2})$ would refute the framework's geometric prediction. Any precision-upgrade measurement moving the Koide relation outside 3σ of $2/3$ would refute the cubic-Casimir derivation. The Category A predictions in independent CKM, lattice-QCD, and Higgs-sector channels — $|V_{cb}|$, m_d/m_s, and $\lambda(M_{\rm Pl})$ — provide further falsification handles through measurements independent of the Cabibbo channel that constrains λ.

The Category B predictions ($\sin^2\theta_{12}$ and $\sin^2\theta_{23}$, the mass-chain entries via the Koide structure, the Jarlskog relation) constitute *conditional* falsification targets. The framework's specific structural forms are the natural predictions of the cubic-group machinery; their match to observation under the empirical mass-hierarchy bijection is what the framework predicts. The magnitude-level closure (verifying the cubic-Casimir structure produces the specific numerical coefficients) is open through lattice perturbation theory in the emergent description; the directional/assignment-level content is Layer 2(b) bijection-specific by Chapter 2 §2.5 and is empirical input rather than open derivation. JUNO's design-precision measurement of $\sin^2\theta_{12}$ at sub-percent precision is the nearest forward-falsifiable test at high precision in this category.

Cross-sector universality. A structural feature of the framework's prediction package deserves explicit treatment: the same numerical structural inputs that produce quark-sector observables also produce lepton-sector observables, with no separate fitting between sectors.

The Wolfenstein parameter $A^2 = 2/3$ from off-democratic projection on the cubic generation axes appears in the quark sector via $|V_{cb}| = A\lambda^2 = 0.041$ (matching observation at 0.4σ) and in the lepton sector via the PMNS Condition 2 sum rule $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$ (matching observation at 0.07σ on JUNO's $\sin^2\theta_{12}$). The Cabibbo coupling $\lambda^2 = 1/(2\pi^2)$ from the inter-generation Brillouin-zone distance appears in the quark sector via Cabibbo λ (0.04% match) and $m_d/m_s = \lambda^2$ (0.7% match), and in the lepton sector through the Koide chain (m_e at 0.003% and m_μ at 0.002% via Q and θ_0). The threshold product $A \cdot B$ of §6.2 is a separate numerical cross-check, with values $\approx$ 48–62 across finite-size models against the fitted 46.4; its leading-power dependence leaves the comparison inconclusive.

The cross-sector pattern places a substantial constraint on alternative input choices: any alternative natural input to the framework's machinery that fits quark-sector observation would also need to fit lepton-sector observation. Alternative natural inputs that produce different Cabibbo angles (e.g., BCC or FCC inter-generation distances) all deviate from $\lambda = 1/(\pi\sqrt{2})$ by $> 20\%$ and would correspondingly mispredict m_d/m_s and $\sin^2\theta_{13}$ by similar fractions. Alternative power combinations of A and λ in $\sin^2\theta_{13}$ ($A^2\lambda^2$ at 32% deviation, $A\lambda^2$ at 50% deviation) similarly fail the cross-sector match. The framework's specific $A^2 = 2/3$ and $\lambda^2 = 1/(2\pi^2)$ inputs are consistent with observation in both sectors simultaneously.

What follows. Chapter 7 develops the gravitational sector — the Bekenstein-Hawking 1/4 factor as the ratio matched by the horizon calibration under the conditions of §7.3 (a consistency relation: GW250114 confirms the classical area theorem; the coefficient has no direct test), the calibration $\hbar = c^3\epsilon^2/(4G\sigma)$ relating the action scale to the discreteness scale (a self-consistency condition, §7.4), the dissolution of the cosmological constant problem at the level of two incommensurable descriptive levels, conditional on the covariant coupling step, and the dark-sector phenomenology with the MOND acceleration scale $a_0 = cH/6$ and the entropy-displacement account of dark matter. The empirical record there is independent of and complementary to the Standard Model content developed here: the gravitational sector tests the substratum's boundary thermodynamics rather than its bulk gauge structure.

Chapter 8 develops the JUNO prediction in detail, as a focused case study of the framework's sharpest empirically matched Layer 1-plus-Layer-2(b) value (a parameter-free retrodiction). The November 2025 measurement and its discrimination from the TM1/TM2 alternatives are presented with the experimental context that this chapter abbreviated. Chapter 9 develops universality classes and the framework's relationship to alternative unification programs, with the

OI-string analysis at the matrix-model level and the framework's structural commitment to no grand unified theory developed against the broader landscape of unification proposals.

The framework's empirical record in fundamental physics, developed across Chapters 5 through 9, is concentrated at the Level G3 × Level D intersection of the hierarchical structural realism developed in Chapter 3 — class-specific predictions of the cubic-lattice substratum representative, with substantial empirical support and pre-registered falsifiability. The structural and methodological commitments of Part I are made quantitatively precise here.

Chapter 7

The Gravitational Sector

7.1 What this chapter develops

Chapters 5 and 6 developed the framework's content in the Standard Model sector: the gauge group SU(3) × SU(2) × U(1) from the cubic decomposition of the substratum, three matter generations from the staggered taste structure, twenty-two classified SM observables (two parameter-free structural and five conditional structural; the remainder layered, chained or fitted), with the measured comparisons spanning 0.0003% to 1.2% across the parameter space. The content was developed by *specifying* the substratum dynamics — the wave equation on a cubic lattice with the link count under H-link — and tracing the structural consequences. The framework's content in the gravitational sector follows a different route.

The gravitational sector applies the framework to the cosmological horizon as a causal partition, with the substratum dynamics treated as a free input. The cosmological horizon — the boundary beyond which no signal propagating at or below the speed of light can reach the observer — implements the framework's visible-hidden partition at its largest physical scale. Sub-c observers share the same causal structure, so all such observers share the same emergent action scale $\hbar$. The horizon's thermodynamics relates the emergent action scale $\hbar$ to the discreteness scale ϵ, with κ and A dropping out of $\hbar$ (Sections 7.3–7.4); it recovers the Bekenstein-Hawking entropy with its 1/4 coefficient as a consistency relation of that calibration and supplies the dark-energy bookkeeping read in the running-vacuum channel (Section 7.8).

This chapter develops seven structural results. Section 7.2 verifies coupling C1, slow bath C2 and large capacity C3 at the cosmological horizon, with ratio $\tau_S/\tau_B \sim 10^{-32}$, and records history readback C4 as a named realization condition there, not presently discharged. Section 7.3 derives the emergent ac-

tion scale $\hbar = c^3\epsilon^2/(4G\sigma)$, with σ the entropy of one boundary mode, from thermal self-consistency between the classical horizon temperature (computed from the Jacobson thermodynamic argument with no quantum input) and the Euclidean KMS period of the emergent quantum field theory, conditional on the infrared detailed-balance slope of the horizon reservoir matching the Killing period (H-slope) together with the horizon and frame conditions. Section 7.4 fixes the discreteness scale $\epsilon = 2\sqrt{\sigma}\, l_p$ ($\epsilon = 2\, l_p$ at $\sigma = 1$, one nat per mode) as the unique simultaneous solution of the action-scale relation and the definition of the Planck length. Section 7.5 recovers the Bekenstein-Hawking entropy $S = A/(4l_p^2)$: with the calibration the boundary-mode count reproduces it, the 1/4 coefficient being the ratio of the Euclidean KMS period to the Jacobson prefactor that the calibration matches, under the same conditions; the September 2025 LIGO/Virgo/KAGRA observation of GW250114 confirms the area theorem at 99.999% confidence (the 1/4 coefficient itself has no direct test). Section 7.6 treats the black hole information paradox, which the framework does not resolve: information is never lost from the substratum bijection φ, bijectivity being a premise; a cycle-averaged Page curve follows under the scrambling condition H-scramble and the decoupling condition H-decouple, taking the energy shell of the black hole and its radiation to be their full product space, by a mechanism substantially different from that of the quantum-extremal-surface and entanglement-island programs, with a Page time that, for a mass-independent emission coefficient, is a fixed fraction of the evaporation time set by the entropy the radiation carries per unit of horizon entropy lost — $t_P \approx 0.646\, t_{\text{evap}}$ when the two are equal; H-scramble fails for every bijection acting on an initial configuration with the bipartition aligned to the configuration coordinates, so the Page value requires an initial superposition over many configurations or a non-aligned bipartition, neither of which the framework specifies. Section 7.7 proposes a dissolution of the cosmological constant problem by identifying the emergent quantum vacuum energy and the effective cosmological constant as properties of logically distinct descriptions rather than commensurable quantities whose 10^{122} ratio requires cancellation; the descriptive distinction is established, and the dissolution is conditional on the covariant coupling step that [GR §6.2] leaves open. Section 7.8 develops the framework's content in dark-energy phenomenology — Running Vacuum Model with ν_{OI} bounded at the emergent-QFT scale $\sim 10^{-32}$ — and Section 7.9 develops the dark-sector content (95% invisible matter as concordance rather than puzzle (the $\sim 95\%$ being partly definitional, $\approx 1 - \Omega_B$), MOND $a_0 = cH/6$ from entropy displacement, declining rotation curves at $r_M = \sqrt{6GM_B/(cH)}$).

The methodological discipline of Chapter 4 organizes the gravitational predictions into three tiers. **Tier 1** consists of what follows from partition geometry alone — the energy-shell temperature of the horizon reservoir, the finite counting of boundary modes, and the two-level distinction between the emergent vacuum energy and the effective cosmological constant. These hold for any finite deterministic system satisfying the four conditions at its cosmological horizon. They are universality-class-level structural results — Level C × Level G4 in the

hierarchical structural realism of Chapter 3, where the framework operates at the level of structural commitments shared across substratum representatives rather than at the specific cubic-lattice realization. The results built on them carry named conditions, stated where they are derived: the action-scale relation $\hbar = c^3\epsilon^2/(4G\sigma)$, the discreteness scale $\epsilon = 2\sqrt{\sigma}\, l_p$ and the Bekenstein-Hawking 1/4 coefficient require that the infrared detailed-balance slope of the horizon reservoir match the Killing period — the condition H-slope of [GR §3.2] — together with the horizon and frame conditions of Section 7.3; the descriptive distinction behind the cosmological-constant dissolution is established while its dynamical form is conditional on the covariant coupling theorem the companion paper leaves open ([GR §6.2]); and the Type II running-vacuum form is not derived, its local expansion being shared by every smooth model (Section 7.8). **Tier 2** consists of dark-sector phenomenology — the 95% invisible matter fraction, the MOND acceleration scale $a_0 = cH/6$, the rotation-curve crossover. These follow from the thermodynamic Clausius relation applied to the boundary reservoir, so they inherit the conditions on the de Sitter temperature, and the dimensional factor 1/6 is imported from Verlinde's emergent-gravity analysis (Section 7.9); they sit at Level D $\times$ Levels G1–G2 of the hierarchical structure. **Tier 3** consists of solution-specific quantities — the transition steepness between MOND and Keplerian regimes, the greybody factor for Hawking radiation, the specific microscopic realization of the substratum — which depend on the particular bijection φ within $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$ and are addressed by the cubic-lattice realization developed in Chapters 5–6.

The tier system clarifies what each result claims. A conditional result like "$\hbar = c^3\epsilon^2/(4G\sigma)$" is not "the framework predicts the value of Planck's constant uniquely" but "any horizon-bounded embedded-observer system with the framework's structural conditions, and satisfying H-slope at its horizon, gets this functional form for the emergent action scale." The Tier 1 results, and the conditional results built on them, are robust to the substratum's specific dynamics; only the discreteness scale ϵ remains as a free parameter, and self-consistency with the Planck length definition fixes it for a given entropy σ per boundary mode, $\epsilon = 2\sqrt{\sigma}\, l_p$ ($2\, l_p$ at $\sigma = 1$, one nat per boundary mode). A Tier 2 result like "$a_0 = cH/6$" depends on the geometric factors of de Sitter space but not on the substratum's gauge structure; it would hold for any framework where the dark-sector phenomenology arose from entropy displacement at the cosmological boundary — with the dimensional factor taken from Verlinde's analysis rather than derived here. A Tier 3 result depends on the specific bijection and varies across realizations within the universality class. The empirical record of the gravitational sector — concentrated at Tiers 1 and 2 — is therefore not a record of the cubic-lattice substratum but a record of the framework's broader structural commitments at the boundary scale.

7.2 The cosmological horizon as causal partition

The cosmological horizon implements the framework's causal partition at the largest physical scale accessible to any sub-c observer. The horizon's role in the framework is to separate visible and hidden sectors by causality rather than by an explicit structural choice: the visible sector is the interior of the horizon, with content that the observer can causally access; the hidden sector is everything beyond, with content that no sub-c signal can reach in finite time. This is a consequence of general relativity's causal structure, not a modeling choice imposed on top of it.

The horizon partition is universal across sub-c observers. Different observers have different horizon areas — depending on their worldline and the local Hubble parameter — but they share the same causal structure: any pair of observers within each other's horizons agrees on what is reachable in finite time and what is not. The emergent action scale $\hbar$ depends only on local geometric quantities and not on the observer's specific horizon area, so all sub-c observers share the same value of $\hbar$. The framework's content here is that the universality of Planck's constant across observers — a feature that the Standard Model treats as a postulate — is a structural consequence of the universality of GR's causal structure.

Verification of C1 (coupling). The Hamiltonian and momentum constraints of general relativity correlate interior and exterior initial data on any Cauchy surface. In the ADM formulation, the bulk Hamiltonian is a sum of constraints that vanish on-shell, with physical time evolution generated by the boundary term at the horizon, which depends on data from both sides. The physical phase space is therefore a constraint surface within the kinematic product of interior and exterior configuration spaces. This is stronger than the non-trivial coupling required by the framework's C1: the constraints enforce correlations that persist on all timescales and cannot be perturbatively removed. The cosmological horizon satisfies C1 not as an additional assumption about the dynamics but as a structural consequence of general relativity.

Verification of C2 (memory persistence). The hidden-sector relaxation timescale is the Hubble time, $\tau_B \sim 1/H \sim 10^{17}$ s. The visible-sector dynamics timescale for any laboratory process is much faster: typical electroweak processes occur on timescales $\tau_S \sim 10^{-15}$ s. The ratio $\tau_S/\tau_B \sim 10^{-32}$ is extraordinarily small, making the corrections to the framework's leading-order results at scale τ_S/τ_B correspondingly small. C2's slow-bath realization is satisfied with enormous margin — the cosmological horizon is among the slowest baths the framework's conditions admit. Corrections of order 10^{-32} to the framework's leading results are well below current observational precision in all relevant sectors.

Verification of C3 (sufficient capacity). The horizon's information-storage capacity scales with the horizon area: the number of independent modes is $N_H = A_{\text{horizon}}/\epsilon^2 \sim 10^{122}$ for the present-day cosmological horizon at $\sigma = 1$. The visible sector's mode count is much smaller — bounded by the volume divided

by the relevant length scale of the observer's measurement. The ratio $N_H/N_V \gtrsim 10^{60}$ for any laboratory-scale visible sector, with the actual ratio depending on the visible sector's spatial extent. The capacity condition is satisfied with enormous margin: the cosmological horizon's information capacity exceeds any visible-sector observer's capacity by tens of orders of magnitude.

Status of C4 (history readback). The horizon construction establishes bidirectional coupling (the constraint structure of C1), persistence (C2) and ample capacity (C3), but this chapter does not independently demonstrate that a visible-history record written into hidden boundary degrees is routed back into later visible conditionals within the accessible window — the read-write cycle that Chapter 1 names as the property to be demonstrated in the cosmological realization. Bidirectionality of the boundary coupling is a strengthened form of C1 and persistence is C2; neither, nor their conjunction, supplies the routing that C4 asserts, and since C1 follows from C4 and C3 holds in every faithful realization, verifying those two does not supply it either. C4 is therefore a named realization condition at the cosmological cut, not presently discharged.

Three of the four conditions hold at the cosmological horizon by structural argument rather than by phenomenological assumption, and the fourth is named rather than assumed. The framework's results in the gravitational sector rest on the horizon and frame conditions stated with them in §7.3 — H-slope above all — and consume no readback claim. The eigenstate thermalization hypothesis, which motivates the mixing hypothesis behind the slow-bath form of the C2 necessity direction, is the standard assumption for the horizon complement (a generic chaotic many-body system).

7.3 The emergent action scale $\hbar$

The most distinctive content of the framework's gravitational sector is the treatment of Planck's constant $\hbar$ as an emergent action scale whose relation to the boundary geometry is fixed by the cosmological horizon's thermodynamics. The standard view treats $\hbar$ as a fundamental constant of nature, its value $1.055 \times 10^{-34}\,\mathrm{J \cdot s}$ measured. The framework derives $\hbar = c^3\epsilon^2/(4G\sigma)$ from thermal self-consistency between classical horizon thermodynamics and the emergent quantum field theory on the horizon — with σ the entropy of one boundary mode and ϵ the discreteness scale of the substratum partition, fixed by self-consistency in Section 7.4 to $\epsilon = 2\sqrt{\sigma}\,l_p$; the two relations together fix only the ratio $\epsilon/\sqrt{\sigma} = 2\,l_p$, a self-consistency condition rather than an independent determination of the value of $\hbar$.

The derivation proceeds in four steps.

Step 1: The classical horizon temperature. The Jacobson thermodynamic argument assigns the cosmological horizon a temperature from classical considerations alone, with no quantum input. The argument runs as follows. The horizon is a bifurcate Killing horizon with surface gravity κ. The first law of thermodynamics applied to a horizon-bounded region gives $dE = T\,dS$, where

E is the energy flux across the horizon, T is the horizon temperature, and S is the horizon entropy. The Einstein equations applied to the horizon's Raychaudhuri equation give $dE = (c^2\kappa/8\pi G)\, dA$, where A is the horizon area. Combined with the structural commitment that each minimal cell of area ϵ^2 contributes one boundary mode, carrying the dimensionless entropy σ in nats ($dS = \sigma\, dA/\epsilon^2$), the first law gives a classical horizon temperature

$$k_B T_{\mathrm{cl}} = \frac{c^2\epsilon^2\kappa}{8\pi G\sigma}.$$

The temperature contains no $\hbar$ — only classical quantities (κ, ϵ, σ, G, c). This is the substratum's classical assignment of a temperature to the partition boundary, derived from the Einstein equations and the discreteness scale of the partition.

An independent route to Einstein's equations. Einstein's equations would follow for the framework's emergent matter, given the single-cone structure qualified below, from an external theorem with an entirely different axiom base. The gravitational closure program of Düll, Schuller, Stritzelberger and Wolz (Phys. Rev. D 97, 084036) shows that for matter field equations whose causal structure is that of standard-model fields on a single Lorentzian light cone — the class the framework's emergent sector occupies if its single-cone structure holds, that structure being supported by the rotational-protection results of the companion papers, conditional on loop-level octahedral invariance and an aligned condensate (Chapter 19 §19.3.7), and by the symmetric-point universality of the block cones under H-blind ([SM §4.4]) — the requirement that matter and geometry evolve together consistently, independently of how spacetime is sliced, forces the gravitational action to be Einstein–Hilbert with a cosmological constant. No thermodynamics enters that argument at all. Where the closure route leaves Newton's constant and the cosmological constant undetermined, the framework's relation $\hbar = \mathrm{c}^3\varepsilon^2/(4\mathrm{G}\sigma)$ and the dark-energy analysis of the following sections take over. The closure theorem is, moreover, one of a family of such necessity results that the emergent sector inherits: Lovelock's theorem, which shows that in four dimensions the only second-order, generally covariant dynamics for a metric alone is Einstein's with a cosmological constant, and the spin-2 results of Weinberg and Deser, which show that a massless spin-two particle coupled to a complete matter sector is forced into universal coupling and Einstein's theory at low energy. Each of these routes would deliver Einstein's dynamics from hypotheses the framework has not established — the emergent light cone, the massless spin-two bound state, the absence of extra light gravitational modes — and they answer the standard objection facing every emergent-spacetime program, that such constructions deliver the geometry of spacetime but never Einstein's dynamics, only once those hypotheses are established. The thermodynamic derivation of this chapter uses Einstein's equations through Jacobson's relation and does not derive them; the covariant field equations remain open (Chapter 19 §19.2). The behaviour of the framework's small lattice corrections under the closure machinery remains an open consistency check.

Step 2: Boundary-only dependence. The emergent quantum field theory on the horizon depends only on the boundary modes — the degrees of freedom at the horizon itself — and not on the bulk content of the visible or hidden sectors at scales far from the boundary. This is the framework's partition-relativity result: under the canonical measure of Lemma 3 of Chapter 1, the emergent description is uniquely determined by the partition geometry at leading order in τ_S/τ_B, with bulk corrections at order τ_S/τ_B for the cosmological partition. The boundary-only dependence is a structural consequence of the small parameter $\tau_S/\tau_B \sim 10^{-32}$: the emergent QFT is effectively a theory of the boundary modes, with bulk corrections suppressed by the slow-bath timescale ratio.

Step 3: Dimensional analysis of $\hbar$. The emergent action scale $\hbar$ depends only on local geometric quantities at the horizon. Dimensional analysis on the available constants gives $\hbar \propto c^a \epsilon^b G^d$ with $\hbar$ having dimensions of action. The unique combination yielding action dimensions is $\hbar \propto c^3\epsilon^2/G$. The proportionality constant is not fixed by dimensional analysis alone; it is fixed by thermal self-consistency in Step 4. The dimensional argument excludes κ from $\hbar$ on observer-universality grounds: different observers have different κ at their respective horizons, but they share the same $\hbar$.

Step 4: Thermal self-consistency. The classical horizon temperature from Step 1 must equal the Euclidean KMS period of the emergent quantum field theory on the same horizon. The KMS condition follows from the regularity of the Wick-rotated metric at the horizon: a QFT on this bifurcate Killing horizon background is periodic in imaginary time with period $\beta = 2\pi c/\kappa$ under the H-Hawking conditions, giving a KMS state at temperature $T_Q = \hbar\kappa/(2\pi c k_B)$ when the emergent state belongs to that KMS class (H-spectrum). The step is therefore a conditional infrared calibration, not a theorem within the derived QFT ([GR §3.2]).

The two temperatures — T_{cl} from the classical substratum, T_Q from the emergent QFT — refer to the same physical degrees of freedom: the boundary modes across which the visible-hidden coupling acts. Temperature is a state property; two complete descriptions of the same modes cannot assign different temperatures without logical contradiction. The matching $T_{\mathrm{cl}} = T_Q$ identifies the classical horizon temperature with the emergent KMS temperature on the same boundary modes; it holds under the infrared detailed-balance condition H-slope, which partition geometry does not supply, with corrections at order $\tau_S/\tau_B \sim 10^{-32}$:

$$\frac{c^2\epsilon^2\kappa}{8\pi G\sigma} = \frac{\hbar\kappa}{2\pi c}.$$

The surface gravity κ cancels — a non-trivial cross-check on the boundary-only argument of Step 2, which excluded κ from $\hbar$ on observer-universality grounds. The thermal matching here recovers that exclusion from independent physics: κ enters both sides of the consistency equation, and only the combinations of c,

G, ϵ predicted by Step 3 survive. Solving for $\hbar$:

$$\boxed{\hbar = \frac{c^3\epsilon^2}{4G\sigma}}.$$

The derivation has the structure of a gap equation: ϵ is the free geometric input, $\hbar$ is the output. The match between the two temperature computations is nontrivial. The classical temperature could have depended on volumetric hidden-sector data — excluded by the boundary-only argument of Step 2 — but in fact depends only on the boundary geometry. The KMS temperature could have depended on the state of the emergent QFT — but in fact is kinematic in its parameter, conditional on H-balance in its class, depending only on the surface gravity. What the calibration needs is narrower than H-balance: only the infrared slope of the detailed-balance ratio across the accessible horizon transitions, the condition H-slope ([GR §3.2]), which identifies the classical inverse-energy temperature with the emergent Killing period; the gap equation, and with it the 1/4 coefficient of Section 7.5, is conditional on it. One prerequisite of the KMS step is carried explicitly: Euclidean regularity at a bifurcate Killing horizon presupposes local boost structure in the emergent description at the horizon, so the gap equation is partially downstream of the boost sector's protection mechanism: the symmetric-point matching under H-blind ([SM §4.4]) and, conditional on it, the M_X-controlled running tracked in the Lorentz-sector accounting ([GR §8.5]). At the normalized $d = 3$ observer level the free massless branch supplies the common infrared cone at the symmetric matching point under H-blind ([SM §4.4]); exact all-momentum boost invariance is not claimed — the unit-Courant cone of the one-dimensional discrete wave equation does not carry to the three-dimensional observer branch, and away from the body diagonals the cone holds only in the infrared ([GR §8.5]). Conditional on that matching, the custodial symmetry forbids a condensate-independent sector-differential counterterm, and an interacting cross-charged calculation, reported in the companion paper and not independently verified, gave the running a quadratic-with-logarithm form decoupling as $(M_X/M_{\rm Pl})^2$, with the non-logarithmic constant that loop's own, the substratum's component-local nonlinearity contributing only to the universal cone. The one item shared with the Lorentz-sector accounting is that reported running calculation; on its reported form the cross-charged mass M_X would sit in an illustrative window of roughly two to four decades, kept open from below by the baryon safety of the cross-charged sector, which is of the proton-safe leptoquark class rather than the grand-unified kind. That neither pathology obtains makes the gap equation a genuine determination of $\hbar$ in terms of the discreteness scale.

The counting of one mode per cell, $\eta = \sigma/\epsilon^2$, is definitional — and the internal components do not multiply it. The entropy-density argument in Step 1 used $dS = \sigma\, dA/\epsilon^2$ — one boundary mode per cell of area ϵ^2. This counting is not an independent physical ansatz about the mode density but a consequence of the definition of ϵ as the minimal distinguishable scale. A cell

finer than ϵ would subdivide the minimal scale; counting fewer modes would require a distinction below ϵ to group them. Two potential multiplicities are settled separately. The number of internal *states* per mode (the alphabet size q) does not change the count of one mode per cell; it bounds the entropy σ of a mode, $\sigma \leq \ln q$, and the alphabet-size hypothesis of Chapter 5 does not fix σ; the numerical values of this chapter rest on the assumption $\sigma = 1$, one nat per boundary mode, which the bound makes unavailable for a single mode with $q = 2$. The Bekenstein-Hawking coefficient does not depend on σ, which cancels from the entropy law and enters only the relation between ϵ and the Planck length, $\epsilon = 2\sqrt{\sigma}\, l_p$ ([GR §3.1, §4]). The number of internal *components* per site ($K = 2d = 6$) is physical, but it does not multiply the count either, for a structural reason: the $K = 2d$ components are the $2d$ link directions from a cell, each coupling across exactly one link, and a boundary cell has exactly one outward-normal link crossing the horizon — the other $2d-1$ are tangential or inward. One crossing link per cell means one coupled boundary mode per cell, forced by the link-direction structure rather than assumed ([GR §3.1]); one mode per cell does not, however, establish one nat per mode. The count of one mode per cell, and with it $\eta = \sigma/\epsilon^2$, is therefore the definition of ϵ together with a forced consequence of the component structure.

The counting also sits at the intersection of an independent family of results: the Bekenstein bound and Bousso's covariant entropy bound each cap the entropy of a region by its boundary area in Planck units — area-law mode counting reached from quantum field theory and from light-sheet geometry respectively. One caution keeps the logic clean: those bounds are formulated with $\hbar$ already in hand, and the framework derives $\hbar$ downstream of this counting, so the family can serve only as downstream convergence — the emergent theory, once $\hbar = c^3\epsilon^2/(4G\sigma)$ is fixed, satisfies both bounds — and never as motivation for the counting, which stands on the definitional argument above alone. Citing the bounds as motivation would re-import the circularity that the definitional argument removed.

The consequences of the gap equation. Given ϵ, the gap equation fixes $\hbar$; the specific relationship $\hbar = c^3\epsilon^2/(4G\sigma)$ then enters three later results. In Section 7.5 it enters the Bekenstein-Hawking entropy $S = A/(4l_p^2)$ with the factor $1/4$. In Section 7.7 it enters the de Sitter entropy S_{dS} as the compression ratio of the conditional cosmological constant dissolution. In Section 7.8 it enters the Type II running-vacuum reading of the dark energy. The three relations share the calibration; they are consequences of it, not independent checks of it. The situation is analogous to deriving the Schwarzschild metric with mass M as a free parameter: the derivation has genuine content (the functional form of the metric) even though one input is not determined from within the derivation.

The framework's relation to Jacobson's original derivation deserves explicit comment. Jacobson's argument [Phys. Rev. Lett. 75, 1260 (1995)] uses $\hbar$-containing forms throughout; the present derivation does not — it uses the classical identity $dE = (c^2\kappa/8\pi G)\, dA$ and the classical entropy density $\eta = \sigma/\epsilon^2$. The logical or-

dering differs from Jacobson's: his argument derives G from η and $\hbar$; the present argument takes G as a parameter of the classical Hamiltonian and derives $\hbar$ from G and ϵ. The two are consistent — substituting $\eta = \sigma/\epsilon^2$ and $\hbar = c^3\epsilon^2/(4G\sigma)$ into Jacobson's relation $G_{\text{eff}} = c^3/(4\eta\hbar)$ recovers $G_{\text{eff}} = G$, with σ cancelling — but the roles of input and output are reversed. The Gibbons-Hawking form $k_B T_{\text{GH}} = \hbar\kappa/(2\pi c)$ enters as the KMS temperature of Step 4; the calibration relates the $\hbar$ in it to ϵ and σ through $\hbar = c^3\epsilon^2/(4G\sigma)$.

7.4 Self-consistency and the discreteness scale

The gap equation $\hbar = c^3\epsilon^2/(4G\sigma)$ from Section 7.3 takes ϵ as a free geometric parameter — the discreteness scale of the substratum partition — and produces $\hbar$ as the output. The framework's content beyond this single relation requires that ϵ itself be determined. The determination comes from self-consistency with the definition of the Planck length.

The Planck length $l_p = \sqrt{\hbar G/c^3}$ is defined as the unique length scale constructed from $\hbar$, G, and c. Substituting the framework's relation $\hbar = c^3\epsilon^2/(4G\sigma)$ into this definition:

$$l_p^2 = \frac{\hbar G}{c^3} = \frac{c^3\epsilon^2}{4G\sigma} \cdot \frac{G}{c^3} = \frac{\epsilon^2}{4\sigma}.$$

Rearranging gives the discreteness scale:

$$\boxed{\epsilon = 2\sqrt{\sigma}\, l_p.}$$

The result is, for a given σ, the unique value of ϵ consistent with the framework's gap equation and the standard definition of the Planck length. The framework's substratum partition cell has linear extent $2\sqrt{\sigma}\, l_p$, which is $2\, l_p$, twice the Planck length, at $\sigma = 1$.

Here σ is the entropy of one boundary mode, a dimensionless number of nats, and the Bekenstein-Hawking coefficient of Section 7.5 does not depend on it ([GR §3.1, §4]). The alphabet-size hypothesis of Chapter 5 does not fix σ: a mode with q internal states carries at most $\ln q$, so the assumption $\sigma = 1$, one nat per boundary mode — an assumption about the entropy of a mode, not a choice of units, since entropy is already measured in nats — is unavailable for a single mode with $q = 2$.

The self-consistency is algebraically equivalent to the gap equation. The framework contains one free geometric parameter; self-consistency with the Planck length definition fixes its value for a given σ. It adds no relation beyond the structural commitment $\hbar = c^3\epsilon^2/(4G\sigma)$ and the definition of l_p, and its numerical value takes the measured $\hbar$ as input. The result is a *constraint* on ϵ, not an independent measurement.

The discreteness scale and σ. With σ undetermined, self-consistency fixes only the ratio $\epsilon/\sqrt{\sigma} = 2\,l_p$, and a Planck-scale ϵ additionally requires σ of order one. The bound $\sigma \leq \ln q$ limits σ from above for a mode with q states, but the count bounds it from below only by $\sigma > 0$: $\sigma = 10^{-12}$ would give $\epsilon = 2 \times 10^{-6}\, l_p$. The gap equation does not exclude discreteness scales far from the Planck length; whether they are excluded on independent grounds is open.

The framework's content in the discreteness scale therefore has three structural commitments: the gap equation, the Planck-length definition, and the self-consistency between them. The result $\epsilon = 2\sqrt{\sigma}\, l_p$ is a derived quantity for a given σ, not a free parameter, with $\epsilon = 2\, l_p$ at $\sigma = 1$, one nat per mode — but the derivation requires accepting the gap equation as the substratum's structural commitment rather than $\hbar$ as a fundamental constant.

7.5 The Bekenstein-Hawking entropy

The Bekenstein-Hawking entropy $S = A/(4\, l_p^2)$ assigns each horizon a thermodynamic entropy proportional to its area, with the proportionality constant $1/(4\, l_p^2)$. Bekenstein (1973) introduced the area law with a coefficient he could estimate only heuristically; Hawking (1975) fixed the 1/4 from the Hawking temperature and the first law, and the semi-classical and stringy microstate arguments reproduce it. The framework reproduces the same ratio from its horizon calibration: with the discreteness scale $\epsilon = 2\sqrt{\sigma}\, l_p$ fixed in Section 7.4 under the condition H-slope, the boundary-mode count gives the area law with the coefficient 1/4.

Direct mode counting. With $\epsilon = 2\sqrt{\sigma}\, l_p$ from Section 7.4, the number of independent boundary modes on a horizon of area A is

$$N_{\text{modes}} = \frac{A}{\epsilon^2} = \frac{A}{4\sigma\, l_p^2}.$$

Each minimal cell of area $\epsilon^2 = 4\sigma\, l_p^2$ contributes the entropy σ of one boundary mode, giving $S = \sigma N_{\text{modes}} = A/(4\, l_p^2)$. This is the Bekenstein-Hawking entropy in its standard form, and the coefficient does not depend on σ; the mode count $N_{\text{modes}} = S/\sigma$ equals the entropy only at $\sigma = 1$.

The factor of 1/4 is therefore fixed, given $\epsilon = 2\sqrt{\sigma}\, l_p$: it is the ratio $\epsilon^2/(\sigma\, l_p^2) = 4\sigma\, l_p^2/(\sigma\, l_p^2) = 4$ in the denominator, with the ϵ^2 factor arising from the framework's discreteness scale, the σ from the entropy per mode, and the l_p^2 in the denominator arising from the Planck length appearing in the definition of S_{BH}. Equivalently, the 1/4 is the ratio $2\pi/8\pi$ between two physical quantities: the numerator 2π is the period of the Euclidean KMS regularity at the horizon (thermal periodicity of QFT on a Killing horizon), and the denominator 8π is the Jacobson coefficient in the Einstein-equation prefactor $dE = (c^2\kappa/8\pi G)\, dA$ (gravity's coupling). The Bekenstein-Hawking 1/4 is therefore not an arbitrary proportionality constant but the consistency condition between gravitational

thermodynamics and quantum thermal physics — the ratio that makes the framework's gap equation in Section 7.3 close on itself.

Extension to black-hole horizons. The minimal-cell counting argument applies to any horizon with area A. The cosmological horizon and black-hole horizons differ in their global structure — the former is a future causal boundary at infinity, the latter a past or future causal boundary at finite radius — but they share the local property of being bifurcate Killing horizons with surface gravity κ and area A. The framework's mode-counting argument depends only on the local properties; the entropy $S = A/(4\,l_p^2)$ therefore extends to black-hole horizons with the same coefficient.

Observational confirmation: GW250114. The Bekenstein-Hawking area law $dA \geq 0$ is a theoretical prediction of black-hole thermodynamics derived from classical general relativity combined with the second law of thermodynamics. The empirical record before 2025 came primarily from agreement between classical horizon dynamics and semi-classical Hawking radiation predictions — supportive but indirect, with no direct test of the 1/4 coefficient at the percent level.

The LIGO/Virgo/KAGRA observation of GW250114 in September 2025 provided the first direct test of the area law at the percent level. The event was the merger of two stellar-mass black holes; the inspiral phase of the gravitational waveform constrained the initial horizon areas of the two progenitors, and the ringdown phase constrained the remnant horizon area of the merged black hole. The comparison: the remnant horizon area exceeded the sum of the initial horizon areas by an amount consistent with the area theorem at 99.999% confidence — a result the framework, like any theory retaining classical general relativity, preserves.

The scope of this test is important to state precisely: GW250114 compares horizon areas and does not measure entropy, so it tests Hawking's area theorem — classical horizon dynamics, which the framework preserves — and not the 1/4 entropy coefficient. The coefficient, fixed by the calibration of Section 7.4 as the ratio $2\pi/8\pi$ between Euclidean KMS periodicity and the Jacobson Einstein-equation prefactor, is the ratio of the two normalizations that the calibration matches, conditional on H-slope and the horizon and frame conditions, with no direct empirical test at present; any framework retaining classical general relativity passes the GW250114 test identically. Untested is not untestable: the coefficient is equivalent to the Hawking-temperature normalization, so a measured horizon radiation spectrum would fix it. The one contingent channel is the final evaporation burst of a primordial black hole — active all-sky burst searches (Fermi-LAT 2018; HAWC 2020; LHAASO 2025, PRL 135 181005) have so far set rate-density limits only, with next-generation sensitivity projected for CTAO; a detected burst light curve would measure $T(M)$ and hence the coefficient directly. Analogue-gravity experiments probe the kinematic side of the $2\pi/8\pi$ ratio, not the gravitational side. The ongoing LIGO observing runs will accumulate additional events testing the area theorem across a range of

progenitor masses and spins.

Falsification status. The Bekenstein-Hawking 1/4 is a consistency relation of the horizon calibration, not a counted Class B prediction (Chapter 4 §4.5) — conditional at the universality-class level, holding under H-slope together with the horizon and frame conditions on the gap-equation chain (Section 7.1). Falsification of the coefficient at sufficient confidence to exclude experimental systematics would refute that chain (the connection between the classical horizon temperature in Section 7.3 Step 1 and the Euclidean KMS temperature in Step 4) under those conditions, and with it the framework's structural commitment to thermal self-consistency at the partition boundary. The coefficient currently has no direct empirical test — GW250114 constrains the classical area theorem, not the entropy normalization — the $a_0 = cH/6$ rotation-curve family (Section 7.9) is a conditional measured comparison, with agreement within the quoted systematic uncertainty and the factor 1/6 imported.

7.6 The Page curve and the information paradox

The framework's content in the Bekenstein-Hawking sector extends beyond the area-law coefficient to the *dynamics* of black hole evaporation. The black hole information paradox — Hawking's 1975 observation that semiclassical black hole evaporation appears to destroy information, in tension with quantum mechanics' commitment to unitarity — is one of the canonical hard problems of theoretical physics. The framework does not resolve the paradox. Information is never lost from the substratum bijection φ because φ is bijective, and bijectivity is a premise of the framework (Chapter 1), not a result; how much of that information an observer can recover from the radiation is a separate question, which bijectivity alone does not settle. The Page-curve derivation below bears on it under two conditions, H-scramble and H-decouple, neither of which is established, and H-scramble fails for every bijection acting on an initial configuration with the bipartition aligned to the configuration coordinates. On the framework's reading the apparent information loss is an artifact of the trace-out — the same mechanism that underlies the conditional cosmological constant dissolution in §7.7 below and the dark sector concordance in §7.9.

The paradox in its conventional form. Hawking's calculation of the spectrum of radiation emitted by an evaporating black hole produced a thermal distribution with a temperature $T_H = \hbar\kappa/(2\pi c k_B)$, dependent only on the black hole's mass and angular momentum. The thermal spectrum carries no information beyond the macroscopic horizon parameters. If the radiation is exactly thermal — as Hawking's semiclassical derivation suggests — then the radiation carries no information about the matter that formed the black hole, and the information appears to be lost when the black hole completes its evaporation. This is in tension with the unitarity of quantum mechanics, which requires that any time evolution preserve information.

The Page curve is the unitarity prediction. Don Page argued in 1993 that if

black hole evaporation is unitary, the entanglement entropy of the radiation must follow a specific trajectory: rising as radiation accumulates in the early phase, then turning over and decreasing at the *Page time* t_P — defined as the moment when the radiation's entanglement entropy equals the remaining black hole's entropy. If the radiation's entropy equals the horizon entropy it carries off, the Page time comes when half the black hole's initial entropy has been radiated, and earlier if the radiation carries more; the curve's turnover is the empirical signature of unitarity.

The dispute between Hawking's calculation (thermal radiation, information lost) and Page's prediction (Page curve, information preserved) was unresolved for decades. The recent quantum-extremal-surface and entanglement-island programs of Penington, Almheiri-Engelhardt-Marolf-Maxfield, and others have reproduced the Page curve through replica wormhole computations in the gravitational path integral, providing a derivation of the Page curve from semiclassical gravity. The framework offers a conditional alternative through a substantially different mechanism: a derivation from the scrambling condition H-scramble and the decoupling condition H-decouple of [GR Appendix A]. Under those conditions, and taking the energy shell of the black hole and its radiation to be their full product space, the two derivations give a Page curve of the same form, the framework's as an average over the cycles of the dynamics; whether they agree on the Page time depends on the entropy each assigns to the radiation, as set out below.

The observer's description: nested trace-out. The framework's substratum is a deterministic bijection φ acting on a finite configuration space S. Information is never lost from φ — bijectivity is the framework's structural commitment to unitarity at the substratum level. The apparent information loss in Hawking radiation is an artifact of the *observer's* description, not of the underlying dynamics.

The mechanism is the nested-partition formalism. When a black hole forms within the cosmological horizon, the framework's partition structure has two coexisting causal boundaries: the cosmological horizon (the framework's primary partition) and the black hole horizon (a secondary partition within the visible sector). The configuration space decomposes as

$$S = \mathcal{C}_V \times \mathcal{C}_B \times \mathcal{C}_D,$$

where V is the region between the black hole horizon and the cosmological horizon (the external observer's visible sector), B is the black hole interior (the secondary hidden sector), and D is the trans-cosmological region (the primary hidden sector).

The external observer's reduced description of the region V — universally admitting the quantum representation — is obtained by tracing out *both* B and D. Two procedures produce that reduced description on V: a *direct* trace, marginalizing over $\mathcal{C}_B \times \mathcal{C}_D$ in one step, and a *sequential* trace, first marginalizing over

$\mathcal{C}_D$ (cosmological partition) and then over $\mathcal{C}_B$ (black hole partition). The consistency of the nested formalism — the requirement that the two procedures produce the same CPTP channel, the same emergent Hamiltonian, and the same value of $\hbar$ — is established in the framework's appendix to the gravitational sector. The nested trace-out is well-defined; the external observer's emergent quantum mechanics is self-consistent when multiple causal boundaries coexist.

The Page curve from cycle typicality. The framework's derivation of the Page curve does not use the gravitational path integral: the cycle structure of the bijection φ supplies exact time averages, and the Page value enters through a typicality condition on the substratum orbit. The relevant results are Page's mean entropy (Page 1993) and measure concentration (Popescu, Short & Winter 2006): for a bipartite quantum system in a Haar-random pure state, with subsystem dimensions $d_{\min} \leq d_{\max}$, the entanglement entropy of the smaller subsystem lies close to Page's mean, about $\ln d_{\min} - d_{\min}/(2d_{\max})$ — within half a nat of the maximum $\ln d_{\min}$ when the two dimensions are equal, and closer as the larger one grows — with deviations from that mean exponentially unlikely in the dimension of the smaller subsystem. These results concern Haar-random states of the full product space; on a proper energy shell, a subspace fixed by energy conservation, the mean entropy depends on the shell and in general differs from Page's value, so the derivation below takes the shell to be the full product space of the black hole and its radiation, which places no energy constraint on them; an evaporating black hole whose radiation carries away its energy is the energy-shell case, and that version is open ([GR Appendix A]). Applying them to the substratum requires H-scramble: uniformity over classical microstates is not uniformity over quantum pure states, and every finite bijection acting on a definite configuration, with the bipartition aligned to the configuration coordinates, visits only product states, however uniformly it counts its energy-shell cycles ([GR Appendix A]). H-scramble therefore fails in that setting, which is the one the derivation counts: its initial conditions are configurations, so there the derivation has no instance. At a bipartition aligned with the configuration coordinates, the entanglement entropy a bijection produces never exceeds, at any step, the entropy of the initial state's distribution over configurations ([GR Appendix A]); a Page value therefore requires an initial superposition over at least about $e^{S_{\text{Page}}}$ configurations, with S_{Page} Page's mean entropy, or a bipartition not aligned with them. Which of these, if either, the substratum supplies is open, and the condition is not established.

Applied to the evaporating black hole, the relevant bipartite structure is the radiation field outside the black hole versus the boundary modes remaining at the shrinking horizon. Let $n_B(t)$ denote the number of boundary modes remaining at the horizon at time t, and $n_R(t) = n_B(0) - n_B(t)$ the number of modes radiated. Conservation gives $n_B(t) + n_R(t) = n_B(0)$ — the total number of modes is fixed by the initial black hole entropy, $n_B(0) = S_{\text{BH}}(0)/\sigma$.

Under H-scramble, which carries Popescu-Short-Winter measure concentration from the Haar ensemble to the orbit of the bijection on the full product space of

the black hole and its radiation, and under the decoupling condition H-decouple of [GR Appendix A], which treats the black hole and its radiation as a closed system in a pure joint state, the radiation's entanglement entropy, averaged over full cycles of the dynamics at the bipartition of time t, is approximately

$$\overline{S}_{\mathrm{rad}}(t) = \min(n_R(t), n_B(t)) \cdot \ln q,$$

where q is the substratum's alphabet size, so that the state space of one mode has dimension q, with a correction of at most about half a nat, reached at the turnover and exponentially small in $|n_R(t) - n_B(t)|$ away from it. Whether the entropy along a single trajectory, whose bipartition moves with t, follows the same curve is open. The structure has two regimes:

- *Early phase* ($n_R(t) < n_B(t)$): radiation has fewer modes than the remaining black hole; entropy rises as $n_R(t) \ln q$.
- *Late phase* ($n_R(t) > n_B(t)$): radiation has more modes than the remaining black hole; entropy falls as $n_B(t) \ln q$.

The turnover — the Page time t_P — occurs at $n_R(t_P) = n_B(t_P) = n_B(0)/2$, when half the initial boundary modes have been radiated. Under the evaporation law $dM/dt = -\alpha/M^2$, with a mass-independent coefficient α into which the species content and the greybody factors enter ([GR Appendix A]), the mode count falls as $n_B(t) = n_B(0)\,(1 - t/t_{\mathrm{evap}})^{2/3}$, and the Page time is

$$t_P = (1 - 2^{-3/2})\, t_{\mathrm{evap}} \approx 0.646\, t_{\mathrm{evap}},$$

independent of the black hole's mass and of α.

The greybody factors enter a second time, through the entropy the radiation carries. The conservation law $n_B(t) + n_R(t) = n_B(0)$ gives the radiation one mode of dimension q for each boundary mode the horizon loses, so the radiation's state-space entropy grows by exactly the amount the horizon's falls. More generally, write β for the ratio of the radiation's gain in state-space entropy to the horizon's loss, taken constant along the evaporation; when each mode carries $\sigma = \ln q$, β is the radiation's coarse-grained entropy per unit of Bekenstein-Hawking entropy lost. The turnover falls where $\beta\,[S_{\mathrm{BH}}(0) - S_{\mathrm{BH}}(t_P)] = S_{\mathrm{BH}}(t_P)$, and the same evaporation law gives (Page 2013, Eq. (8))

$$\frac{t_P}{t_{\mathrm{evap}}} = 1 - \left(\frac{\beta}{1+\beta}\right)^{3/2},$$

which is 0.646 at $\beta = 1$ and falls as β grows: 0.568 at $\beta = 4/3$, the value for unfiltered blackbody emission of massless quanta. In the quasi-stationary semiclassical regime — a non-rotating, uncharged black hole emitting free fields into empty space (incoming vacuum), outgoing modes in independent thermal-form states with mean occupation $N = \Gamma\, n_{\mathrm{th}}(\omega/T_H)$ for greybody factors $0 < \Gamma \leq 1$, and $dS_{\mathrm{BH}} = dE/T_H$ — every emitted mode carries more entropy than its energy divided by the Hawking temperature, so $\beta > 1$ in that regime (Page 2005,

Eqs. (5)–(10); [GR Appendix A]); its value depends on the species content and on how the greybody factors vary with frequency — solution-specific content, Tier 3 in the sense of Section 7.1. For a large Schwarzschild black hole emitting only photons and gravitons, at constant α, Page's emission rates give $\beta \approx 1.485$ and $t_P \approx 0.538\, t_{\text{evap}}$ (Page 2013, Eqs. (2), (8)). The Page time $0.646\, t_{\text{evap}}$ is therefore conditional on $\beta = 1$ as well as on H-scramble and H-decouple: the mode-for-mode conservation law sets $\beta = 1$, the semiclassical regime just defined has $\beta > 1$, and how much state-space entropy the radiation acquires per boundary mode lost is open.

Comparison with quantum extremal surfaces and islands. The quantum extremal surface formula and the entanglement-island construction developed by Penington, Almheiri-Engelhardt, and others obtain the Page curve by a different route. In the QES approach, the Page curve emerges through an extremization principle that selects the dominant saddle in a replica-trick computation of the radiation's entanglement entropy. The post-Page-time saddle includes an *island* — a region of bulk spacetime that contributes to the radiation's entanglement entropy despite being topologically disconnected from the radiation region. The replica wormhole geometries connecting the replicas in the gravitational path integral are essential to obtaining the saturating Page curve at late times.

The framework arrives at a cycle-averaged Page curve of the same form — rise, turnover, fall — through a substantially different mechanism. The cycle-typicality argument applies Popescu-Short-Winter measure concentration to the deterministic substratum dynamics on the full product space of the black hole and its radiation, under the scrambling condition H-scramble and the decoupling condition H-decouple of [GR Appendix A] — uniformity over classical microstates is not uniformity over quantum pure states, and neither condition is established — with no replica trick and no gravitational path integral. In the QES computation the turnover comes where the island saddle's generalized entropy, at leading order the black hole's remaining Bekenstein-Hawking entropy, falls below the semiclassical entropy of the radiation. The two Page times coincide when that semiclassical entropy grows one-for-one with the horizon entropy lost, the $\beta = 1$ of the mode counting; in the semiclassical regime defined above $\beta > 1$ holds (Page 2005; for the QES setting, Penington 2019, §2), so agreement between the two derivations on the Page time and the Page entropy is not established. A correspondence between the two constructions is conjectured: the *island* in the QES formulation would correspond to the post-Page-time regime in which the black hole's boundary modes form the smaller subsystem, and the *replica wormhole* geometries to the substratum's cycle structure on the energy shell.

This conjectured correspondence — replica wormholes $\leftrightarrow$ substratum cycles — is suggestive but not yet rigorous. Making it precise would require constructing the gravitational path integral as an effective description of the substratum dynamics in the appropriate limit, a task that has not been completed. What the framework's derivation establishes is conditional: under H-scramble and H-

decouple and with the full product space as the energy shell, a cycle-averaged Page curve with its turnover where the radiation's state space overtakes the black hole's follows without the replica machinery. The Page value enters through H-scramble, not through the finite deterministic dynamics, which from an initial configuration produces no entanglement across an aligned bipartition and, across such a bipartition, from any initial state at most the entropy of its distribution over configurations. The open steps include the scrambling condition, the decoupling condition, the value on a proper energy shell, whether the entropy along a single trajectory follows the cycle-averaged curve, and the radiation's entropy per boundary mode lost.

Both approaches require *something* to play the role of typicality — Popescu-Short-Winter measure concentration in the framework's derivation, replica saddles in QES + islands — and both place the turnover where the radiation's entropy reaches the black hole's remaining entropy. If the conjectured correspondence holds, the typicality-providing ingredient need not be gravitational at all, and the cycle-typicality derivation would be a microscopic realization of the entanglement-island construction; on the substratum side the bijection alone does not supply that ingredient, since the entanglement it can produce across an aligned bipartition is bounded by the entropy of the initial distribution over configurations.

The generalized second law. A separate but related statement is the generalized second law: the sum $S_{\text{matter}} + S_{\text{BH}} + S_{\text{dS}}$ is non-decreasing in time. The matter entropy S_{matter} captures the visible-sector matter's information content; the black hole entropy S_{BH} is the horizon area divided by $4\,l_p^2$; the de Sitter entropy S_{dS} is the cosmological horizon's contribution.

The framework does not derive it; what its reduced dynamics supplies is weaker. Bijectivity makes the evolution of the whole configuration space unitary, so the joint von Neumann entropy of the visible sector V, the black hole interior B and the trans-cosmological region D is constant. If the state at $t = 0$ is a product over the three sectors, subadditivity then gives

$$S(\rho_V(t)) + S(\rho_B(t)) + S(\rho_D(t)) \geq S(\rho_V(0)) + S(\rho_B(0)) + S(\rho_D(0))$$

at every later time, including when the degrees of freedom are regrouped into three sectors as the horizon moves. The inequality concerns the sectors' von Neumann entropies. It becomes the generalized second law only if S_{BH} and S_{dS} equal the von Neumann entropies of the black-hole and trans-cosmological reduced states at each time, and the boundary-mode count does not supply that identification: $S_{\text{BH}} = n_B\,\sigma$ counts the horizon's modes, while in the Page-curve derivation above, where the black hole and its radiation share a pure state, the black hole's reduced entropy equals the radiation's and is at most $n_R \ln q$ — near zero early in the evaporation, when S_{BH} is largest. Monotonicity under the reduced CPTP dynamics does not close the gap. It holds for relative entropy; the von Neumann entropy of the output is at least that of the input for every input only when the channel is unital, and then it constrains S_{matter} alone,

not the horizon terms. A derivation of the generalized second law from the substratum needs a relation between horizon area and the hidden sectors' coarse-grained entropy that holds along the evaporation, and that relation is open. Conservation of information at the substratum level rests on bijectivity alone and does not wait on the generalized second law; how much of that information the observer can recover from the radiation is the separate question the Page-curve derivation addresses, under its own conditions.

Empirical status and falsification. The framework's account of the information paradox has no direct empirical signature accessible at current experimental precision — black hole evaporation timescales for astrophysical black holes vastly exceed the age of the universe, and the Page time for a stellar-mass black hole would not occur within any conceivable observational program. The empirical content is therefore conceptual rather than quantitative: the framework's structural commitment to bijectivity entails that information is never lost from the substratum, with the apparent information loss being an artifact of the observer's trace-out.

Bijective dynamics is compatible with radiation that carries less information than the matter that formed it, the remainder being held in sectors the observer does not read, and the nested trace-out is consistent for every dynamics; a measurement of the radiation alone therefore does not test the conservation claim, which rests on bijectivity as a premise (Chapter 1). The QES + islands computations, which obtain a unitary Page curve from semiclassical gravity by an independent route, support unitary evaporation without depending on the framework. The framework's content is therefore consistent with the current theoretical and (limited) empirical picture. Its quantitative target is the Page-time relation $t_P/t_{\text{evap}} = 1 - [\beta/(1+\beta)]^{3/2}$, which becomes a sharp prediction once the radiation's entropy per boundary mode lost is derived; at the mode-counting value $\beta = 1$ it gives 0.646.

7.7 Dissolution of the cosmological constant problem

The cosmological constant problem is conventionally described as the worst quantitative discrepancy in physics. Quantum field theory predicts that the vacuum energy density of the universe should be of order $\rho_{\text{QFT}} \sim M_{\text{Pl}}^4 \sim 10^{113}\,\text{J/m}^3$, obtained by summing zero-point energies $\frac{1}{2}\hbar\omega$ over all field modes up to the Planck cutoff. The observed vacuum energy density is $\rho_{\text{obs}} \sim H^2/G \sim 10^{-9}\,\text{J/m}^3$, with the ratio $\rho_{\text{QFT}}/\rho_{\text{obs}} \sim 10^{122}$. Standard accounts treat this as a fine-tuning problem requiring an unknown cancellation mechanism to suppress the bare vacuum energy by 122 decimal places, re-imposed at each order of perturbation theory.

The framework proposes a dissolution rather than a solution. On its reading the 10^{122} discrepancy is not a fine-tuning failure but a category error — a comparison between two quantities that are properties of logically distinct descriptions and are not directly commensurable as gravitational sources. The descriptive

distinction is established; that the zero-point sum does not source curvature is the covariant coupling step that [GR §6.2] leaves open, so the dissolution is conditional on it. The proposal has three components.

The two levels of description. The framework's structural commitments produce two distinct descriptions of the universe's energy content, operating at logically different levels.

The classical substratum assigns the universe a vacuum energy through the Friedmann equation: $\rho_{\text{crit}} = 3H^2c^2/(8\pi G)$. This is the classical baseline — what geometric measurements probe, what on the proposal gravity couples to, what the standard tests of general relativity confirm. There is no zero-point energy at this level; the vacuum energy is set by the Hubble parameter H through classical thermodynamics. Numerically, $\rho_{\text{crit}} \sim H^2/G \sim 10^{-9}\,\text{J/m}^3$, the Friedmann scale at which any $O(1)$ component of a flat universe sits; its value is an input.

The emergent quantum field theory assigns the universe a vacuum energy by summing zero-point energies over its field modes. With the framework's UV cutoff at $\epsilon = 2\sqrt{\sigma}\,l_p$, the sum has cutoff at $1/\epsilon = M_{\text{Pl}}/(2\sqrt{\sigma})$ in natural units, near the Planck scale when σ is of order one, giving $\rho_{\text{QFT}} \sim M_{\text{Pl}}^4 \sim 10^{113}\,\text{J/m}^3$. This is what local quantum measurements probe, what perturbative quantum field theory describes — but it is a property of the *emergent description*, not of the classical substratum.

The two values describe the same universe at different levels. On this proposal the classical substratum's vacuum energy is the gravitating object; the emergent QFT's vacuum energy is bookkeeping internal to the emergent description. The two are not commensurable as gravitational sources because they are properties of different things.

Why gravity sees the classical value. Spacetime geometry is part of the classical substratum: the metric evolves via Einstein's equations *before* the trace-out that produces the quantum-representable reduced description. The proposal is that the stress-energy that sources gravity is the classical stress-energy of the total microstate, not the expectation value of an emergent quantum operator; establishing it is the coupling step left open. The semi-classical equation $G_{\mu\nu} = 8\pi G\langle\hat{T}_{\mu\nu}\rangle$ is then itself an emergent approximation, and at the classical level the gravitational field equations never encounter the zero-point sum.

Three structural requirements force this geometry-first ordering of the framework's derivation chain.

Definiteness. The trace-out that produces the quantum-representable reduced description is defined by a specific partition $\Gamma = \Gamma_V \times \Gamma_H$. The partition must be definite — not in superposition, not subject to quantum uncertainty — for the marginalization producing the emergent description to be well-defined. A partition in superposition would yield an incoherent mixture of inequivalent quantum theories, not a single QM. The metric at the partition boundary must

therefore be classical.

Non-circularity. The partition is defined by causal structure: the hidden sector contains the degrees of freedom beyond the observer's causal reach. Causal structure is determined by null geodesics of the metric. If the metric were derived from QM, the derivation would require QM → metric → causal structure → partition → QM — a circular dependency with no entry point. The circle breaks only if the metric exists prior to the partition.

Uniqueness of $\hbar$. The emergent action scale is determined by the boundary geometry, as Section 7.3 developed. If the geometry were itself quantum-mechanical, $\hbar$ would depend on a quantum state — but $\hbar$ is observed not to depend on any state. The geometry-first ordering is the only one consistent with a universal $\hbar$.

The framework's geometry-first ordering is therefore not a stipulation but a structural consequence of the framework's three commitments — definiteness, non-circularity, and the universality of $\hbar$.

What the ordering does not yet fix. The three requirements fix the metric at the partition boundary. They do not fix the gravitational field of a laboratory mass that the observer's description places in a spatial superposition. Three completions are consistent with them. In the first, the field follows the substratum configuration of the mass. In the second, the field follows an average over the hidden sector, which is semiclassical gravity at laboratory scales. In the third, the laboratory near field belongs to the emergent quantum description and the classical metric of the ordering is the metric at the partition boundary only; two masses can then become entangled through gravity. What each completion implies for the coherence of the superposition is not derived. Two outcomes must be kept apart: environmental decoherence, in which which-path records pass to unobserved degrees of freedom such as emitted radiation while the total evolution stays unitary, which any of the completions may produce; and an additional nonunitary effect, of the kind Chapter 4 §4.5 lists among the framework's falsifiers. The framework has not determined which completion holds, or whether any produces the second outcome. Fixing the completion and deriving its consequences for interferometric tests of gravitationally mediated entanglement and for searches for gravitational decoherence ([GR §6.2]) is what turns the geometry-first ordering into a laboratory prediction.

The structural origin of the 10^{122} discrepancy. The ratio $\rho_{\mathrm{QFT}}/\rho_{\mathrm{classical}} \sim M_{\mathrm{Pl}}^4/(H^2/G) \sim S_{\mathrm{dS}} = A/(4\,l_p^2)$ is a dimensional identity for any theory with a Planck cutoff and a Hubble-scale density, holding up to a convention-dependent factor. The framework reads it as the Bekenstein-Hawking entropy of the cosmological horizon in its role, identified by Section 7.5, as the information-compression ratio of the trace-out from substratum to emergent description; on that reading, and conditional on the coupling step, the "worst prediction in physics" is the entropy of the observer's blind spot.

The identification holds up to a convention-dependent numerical factor: the

numerator depends on where the emergent sum is cut off, at $1/\epsilon = M_{\rm Pl}/(2\sqrt{\sigma})$ in natural units or at another Planck-scale cutoff, and on how the zero-point integral is normalized, and the denominator on which classical density is compared. $M_{\rm Pl}^4$ against H^2/G gives $S_{\rm dS}/\pi$; the mode sum $k^4/(16\pi^2)$ cut off at $1/\epsilon$ against the critical density $\rho_{\rm crit}$ gives $S_{\rm dS}/(96\pi^2\sigma^2)$; the conventional estimate, a mode sum $k^4/(16\pi^2)$ cut off at the reduced Planck mass $(8\pi G)^{-1/2}$, against the observed dark-energy density $\Omega_\Lambda\rho_{\rm crit}$ gives $S_{\rm dS}/(384\pi^4\Omega_\Lambda) \approx 3.9\times10^{-5}\,S_{\rm dS}$. The 10^{122} factor itself is the de Sitter entropy $S_{\rm dS} = A_{\rm horizon}/(4\,l_p^2)$ at the present cosmological horizon, computed from the observed horizon area combined with the framework's entropy formula. The factor that the conventional QFT view treats as a discrepancy requiring cancellation is, on the framework's reading, the information-compression ratio of the trace-out from substratum to emergent description.

No embedded observer can determine from within whether the geometric or the quantum description is more fundamental, and not merely as a matter of present ignorance. Any adjudication procedure available to an embedded observer is a function of the observer's accessible data, and both descriptions are constructed from the same accessible data — so no criterion grounded in the accessible observations themselves can separate them. (Rankings by theoretical virtue — simplicity, unification — are not blocked by this factoring; but they order presentations of the same observational content rather than track fundamentality, and the framework's answer to the fundamentality question is supplied at the substratum level rather than adjudicated from within either description.) The non-adjudicability follows from the equivalence itself, not from limited resources: a future observation could break the equivalence, but that would be falsification of the equivalence, not adjudication between the descriptions. Nor is this factoring limitation an artifact of the framework's particular observer architecture. Wolpert's limits of inference and T. Breuer's self-measurement theorem establish barriers of the same type for any embedded inference device under far weaker assumptions, corroborating that the limitation is device-general rather than model-specific — with the scope note that those results, unlike the factoring argument, assume free availability of the device's setups. The framework's content here is not to identify the geometric description as "more real" but to read the 10^{122} factor as the compression ratio between the two descriptions; the observed density itself is of the Friedmann scale H^2/G, as any $O(1)$ component of a flat universe is, and its value is an input.

Convergence with the de Sitter observer literature. The framework's geometry-first dissolution converges with recent developments in the de Sitter observer literature, in which an unphysical phase of the de Sitter sphere partition function — long understood to obstruct a clean dS thermodynamic interpretation — cancels exactly when the formalism incorporates an observer with a clock. Maldacena established the cancellation in 2024; Chandrasekaran, Longo, Penington, and Witten (CLPW) showed that the von Neumann algebra of QFT observables in a gravitational subregion promotes from type III to type II once an observer is included, with the resulting entropy pinned to the observer's ref-

erence data. Harlow, Usatyuk, and Zhao established that the closed-universe Hilbert-space dimension is determined by the observer's degrees of freedom.

The framework shares the starting point of these results: the partition trace-out is the *defining* operation rather than an added ingredient, so the observer is present from the outset. It does not reproduce the results themselves. It computes neither the phase of the sphere partition function nor a type II algebra. Its observable algebra on each finite region is a full matrix algebra, and its infinite-region completion is a quasilocal C^*-algebra with no state or representation selected ([GR §3.3]); no von Neumann closure of it, no relation to CLPW's observer-dressed algebra and no identification of CLPW's entropy with a substratum quantity is derived. Whether the construction reproduces CLPW's algebra and its entropy, and whether its trace-out yields Maldacena's cancellation, are open research questions. The observer-essentiality literature converges with the framework on that starting point while differing in mechanism. Chapter 9 develops the framework's relationship to this literature in detail.

The classical vacuum energy scale. The framework's proposed dissolution of the QFT cosmological constant problem does not explain *why* $\rho \sim H^2/G$ takes its specific observed value. This is set by initial conditions through the Friedmann equation, and the framework reclassifies the problem from quantum vacuum cancellation to classical initial conditions. The two problems are not equivalent. The QFT cosmological constant problem requires a 122-decimal-place cancellation between independent contributions — the bare cosmological constant and the zero-point energies of every quantum field — re-tuned at each order of perturbation theory. The initial-conditions question asks why a single parameter takes its observed value; the scale $\rho \sim H^2/G$ is the natural energy density of classical cosmology, not a fine-tuned cancellation between unrelated quantities. Conditional on the coupling step, the framework removes the need for the 122-digit cancellation; the remaining question — why H takes its value — is a standard cosmological problem addressable by inflationary dynamics or other initial-conditions mechanisms.

7.8 Dark energy in running-vacuum form

The framework's content in the dark-energy sector is structural rather than quantitative. Section 7.7 proposed that the emergent quantum field theory's vacuum energy does not gravitate — the descriptive distinction established, the gravitational decoupling conditional on the coupling step that [GR §6.2] leaves open. A distinct question concerns the *evolution* of the effective cosmological constant with cosmic expansion. The framework predicts that the effective cosmological constant has a specific functional form — Type II running-vacuum — but with a coefficient bounded at $|\nu_{\mathrm{OI}}| \sim 10^{-32}$ that is indistinguishable from zero at all foreseeable observational precision.

One scoping remark keeps the registers aligned: the coefficient bound comes from the emergent quantum theory's curvature response, while Section 7.7's dissolution denies that the emergent vacuum energy gravitates absolutely. Whether that response feeds back as a source (in which case its sign is computable and negative for natural couplings, dominated by the top quark) or remains the observer's effective attribution (in which case the coefficient is set by substratum geometry alone, with sign open) is left explicitly unadjudicated — because both readings make the same observable prediction: dark energy indistinguishable from a cosmological constant at every foreseeable precision, with any detected evolution in the DESI-favored direction attributable only to substratum structure carrying a correlated Lorentz-violation signature.

The Type II running-vacuum form. The framework's emergent quantum vacuum energy depends on the cosmological horizon's area, which in turn depends on the Hubble parameter through $A_{\text{horizon}} = 4\pi c^2/H^2$. **Not a distinctive prediction (removed from Tier 1).** The expansion below is correct but universal: for any smooth $f(H)$, setting $g(X) = f(\sqrt{X})$ with $X = H^2$ gives $f = g(X_0) + g'(X_0)(H^2 - H_0^2) + O((H^2 - H_0^2)^2)$ around any $H_0 > 0$. Every smooth dark-energy model has that local form, and the counterfamily $\rho_\Lambda = \rho_0 + A(H^2 - H_0^2) + B(H^2 - H_0^2)^2$ with arbitrary B satisfies the same boundary H-dependence with matter conserved and the Bianchi relation carried by running G. Nothing in the architectural commitments forces $B = 0$, and the running-vacuum literature carries even powers of H, H^4 in other regimes, and $\dot{H}$ terms. So: if "Type II form" means the first local Taylor term it is non-predictive; if it means a global $c_0 + \nu H^2$ law, it is not derived here. The defensible statement is that boundary architecture requires SOME state dependence on cosmological geometry, whose smooth late-time normal form contains an $H^2 - H_0^2$ term. With that said: any smooth dependence of ρ_Λ on H admits a Taylor expansion around the present-day value H_0:

$$\Lambda_{\text{eff}}(H) = \Lambda_0 + \frac{3\nu}{8\pi G}(H^2 - H_0^2) + \mathcal{O}((H^2 - H_0^2)^2).$$

This is the Running Vacuum Model. The framework's content lies not in the existence of an H^2 correction — any framework with an H-dependent vacuum energy admits this form — but in three architectural commitments that select the specific *Type II* class of RVM implementations — a structural reading of the boundary bookkeeping, not a derived law.

Boundary-entropy dependence. The de Sitter entropy $S_{\text{dS}} = \sigma A_{\text{horizon}}/\epsilon^2$ depends on H through the horizon area, so the framework's emergent vacuum energy inherits an H-dependence at the boundary.

Matter conservation. The framework's trace-out ontology conserves matter locally; no direct vacuum-matter energy exchange is permitted. This forbids the Type I RVM variants in which a phenomenological exchange term $Q = 3\nu H\rho_{\text{dm}}$ transfers energy between dark energy and dark matter.

Bianchi consistency. The Einstein equations' Bianchi identity, combined with

matter conservation, requires that Newton's constant $G(H)$ run mildly in tandem with $\rho_\Lambda(H)$ — a structural feature that distinguishes Type II RVM from Type I and from $w_0 w_a$CDM parameterizations.

These three commitments place the framework in the Type II class: matter is locally conserved, $G(H)$ runs with cosmic expansion, and the Bianchi identity is preserved by the joint running of $\rho_{\rm vac}(H)$ and $G(H)$. Type I RVM variants — sharing the same functional form for $\rho_{\rm vac}(H)$ but assuming $G = G_N$ fixed with a phenomenological vacuum-DM exchange — are physically distinct models that the framework does not predict.

Magnitude from the emergent-QFT channel. The framework's structural commitments fix the functional form but the coefficient ν requires a quantitative calculation. The relevant calculation is the one-loop renormalization-group running of the emergent vacuum energy in the curved-spacetime background of the cosmological horizon — a standard computation within emergent quantum field theory, applied with the framework's specific field content.

The Shapiro-Solà β-function for the vacuum-energy running coefficient is

$$\nu = \frac{1}{48\pi^2} \sum_i a_i \frac{M_i^2}{M_{\rm Pl}^2},$$

where the sum runs over the emergent fields with masses M_i and a_i are species-dependent coefficients of order unity. Applied to the Standard Model content — with the heaviest mass being the top quark at $m_t \approx 173$ GeV — the framework's prediction is

$$|\nu_{\rm OI}| \sim (M_{\rm SM}/M_{\rm Pl})^2 \sim 10^{-32}.$$

This is observationally indistinguishable from zero at all foreseeable observational precision. The framework's expectation, on the Type II reading, is therefore that the effective cosmological constant carries the smooth running-vacuum form with a coefficient too small to be measured by any current or planned instrument.

Potential contributions from the substratum. A separate contribution might arise from the substratum's boundary mode count, in principle larger than the emergent-QFT value if the substratum modes have a distinct frequency distribution. The framework's boundary architecture from Section 7.3 catalogs modes by spatial location alone (one degree of freedom per cell of area ϵ^2), with no inherent frequency assignment. Under the natural two-dimensional dispersion relation $dN/d\omega \propto \omega$, modes concentrate near the UV cutoff and contribute a fraction $(H\epsilon)^2 \sim 10^{-122}$ (at $\sigma = 1$) at horizon frequency — well below the emergent-QFT magnitude. Absent additional substratum structure giving a log-uniform boundary-mode distribution at the cosmological horizon, the framework's prediction reduces to $|\nu| \approx |\nu_{\rm QFT}|$, observationally indistinguishable from zero. That absence has since been computed rather than assumed: extracting the boundary spectral measure directly from the substratum wave equation (an exact surface-Green's-function computation, Section 19.3.7) yields

a power-suppressed local distribution — $dN/d\omega \propto \omega^4$, with the two-dimensional $\propto \omega$ above an upper bound on the infrared weight — stable over thirty orders of magnitude in the slow-bath ratio, with the log-uniform distribution appearing nowhere; the interacting extension of that computation is the remaining residual.

Observational status. The empirical situation in late 2025 and early 2026 has evolved substantially with the DESI Data Release 2 results. The DESI collaboration reports evolving dark energy at 2.8–4.2σ significance in the standard $w_0 w_a$ parameterization, with the data favoring a present-day equation of state $w_0 > -1$ transitioning to $w(z) < -1$ at higher redshift — a "phantom crossing" pattern. The framework's prediction is $w(z) \approx -1$ structurally, with the deviations of order $\nu \sim 10^{-32}$. The mild tension with DESI DR2 is 3–4σ on the phantom-crossing pattern, depending on parameterization choices and prior assumptions.

A direct fit in the framework's Type II running-vacuum channel was provided by Bertini, Novaes, von Marttens, and Shapiro in late 2025, who fit a renormalization-group-corrected ΛCDM model — structurally equivalent to Type II RVM at leading order in ν, with both $G(\mu)$ and $\Lambda(\mu)$ running and matter locally conserved — against the Planck 2018 + DESI DR2 + DES-Y5 combined data. Their result:

$$\nu^{\text{Bertini}} = -(2.5 \pm 1.3) \times 10^{-4},$$

with the ΛCDM limit at the 2σ boundary of the posterior. This is consistent at 2σ with the framework's expectation $\nu \approx 0$, conditional on the Type II reading. An earlier pre-DESI canonical RVM fit by Gómez-Valent and Solà returned $\nu = (1.34 \pm 0.38) \times 10^{-3}$ with 3.5σ preference over ΛCDM; the migration from this value toward Bertini's smaller value as the data improved from pre-DESI to DESI DR2 is itself informative — the empirical evidence is shifting toward consistency with the framework's near-zero prediction.

Decisive test structure. DESI Year 5 results are expected in 2028–2029. The framework's expectation makes three discriminating outcomes possible. A null result supports the expectation $\nu \approx 0$ and disconfirms the DESI DR2 phantom-crossing preference. A high-confidence detection of $|\nu| \sim 10^{-4}$ would confirm running-vacuum dynamics consistent with both Type II RVM structure and DESI DR2 trends, but at a magnitude requiring substratum structure beyond what the framework currently derives. A high-confidence detection of phantom crossing would contradict the Type II reading and the smooth-expansion bookkeeping — the strongest available disconfirmation of the framework's dark-energy account.

The framework's expectation $\nu \approx 0$ stands as a falsifiable commitment for the 2028–2029 timeframe.

Coupled neutrino test. The DESI DR2 + CMB analysis finds $\Sigma m_\nu < 0.0642$ eV (95% CL) assuming ΛCDM, prefers normal mass ordering, and bounds the

lightest neutrino mass at $m_l < 0.023$ eV — all consistent with the framework's neutrino-sector predictions of normal ordering with Σm_ν near the oscillation minimum of 0.059 eV. The same analysis finds a 3σ tension with the lower oscillation limit assuming ΛCDM, interpreted in the literature as "a hint of new physics not necessarily related to neutrinos" — the kind of structural mismatch the framework's RVM dark energy would *relax*: an evolving-dark-energy analysis loosens the ΛCDM Σm_ν bound and comfortably accommodates the framework's ~ 0.059 eV prediction, though the RVM-corrected bound has not yet been computed. The neutrino sector predictions developed in Chapter 6 and the RVM dark energy prediction here are coupled in the framework: both follow from the same lattice structure, and the joint observational support is therefore not independent confirmation of separate effects but a single empirical pattern consistent with the framework's central mechanism.

7.9 The dark sector and the MOND acceleration scale

The framework's content in the dark-matter sector is structural rather than particulate. The conventional cold-dark-matter paradigm postulates a new fundamental particle species that interacts gravitationally and contributes about 27% of the universe's energy budget. Decades of direct-detection experiments — XENON, LUX, SuperCDMS, PandaX, and others — have produced no detection at any expected scale. The framework predicts that no such particle exists: the apparent dark-matter phenomenology is instead a structural consequence of entropy displacement at the cosmological boundary, with the MOND acceleration scale $a_0 = cH/6$ deriving from a dimensional argument in de Sitter space.

The invisible gravitational budget as concordance. A corollary of the framework's trace-out structure is that the cosmological horizon carries a gravitational budget that the emergent quantum field theory cannot access. The argument runs in three steps.

The horizon carries $A_{\text{horizon}}/\epsilon^2$ modes and the entropy $S_{\text{dS}} = \sigma A_{\text{horizon}}/\epsilon^2$ by the boundary-mode counting of Section 7.5. The classical temperature is $T_{\text{cl}} = c^3\epsilon^2 H/(8\pi G k_B \sigma)$ from Section 7.3 — computed entirely within the classical substratum with no reference to the emergent quantum description. The total thermal energy carried by the boundary modes is $E_s = S_{\text{dS}} \cdot k_B T_{\text{cl}}$. Distributing this energy over the Hubble volume $V_H = 4\pi c^3/(3H^3)$ via the shell theorem:

$$\rho_s = \frac{E_s}{V_H} = \frac{\sigma A}{\epsilon^2} \cdot \frac{c^3\epsilon^2 H}{8\pi G\sigma} \cdot \frac{3H^3}{4\pi c^3} = \frac{3c^2H^2}{8\pi G} = \rho_{\text{crit}}.$$

The factors of ϵ^2 and σ cancel between $S_{\text{dS}} \propto \sigma/\epsilon^2$ and $T_{\text{cl}} \propto \epsilon^2/\sigma$, so the result is independent of the specific values of ϵ and σ — only Jacobson's $dE = (c^2\kappa/8\pi G)\, dA$, an area-law entropy, and the Friedmann volume enter the calculation. The total effective energy density carried by the boundary modes equals the critical density of the universe.

This is the framework's account of the universe's apparent dark-sector content. The $\sim 95\%$ of the universe's gravitational budget that is invisible to the emergent quantum field theory corresponds to the boundary modes that the trace-out moves into the hidden sector. The dark sector is not a separate substance requiring a new particle species; it is the framework's structural quantity for the partition's hidden-sector content at the cosmological horizon.

Three clarifications are essential. The calculation above uses only classical, pre-trace-out quantities: the number of boundary modes, the classical temperature, and the Hubble volume. No step invokes the emergent quantum description. This is what distinguishes ρ_s from the QFT zero-point energy $\rho_{\mathrm{QFT}} \sim M_{\mathrm{Pl}}^4$: the latter is a property of the emergent description (on the proposal of Section 7.7, does not gravitate), while the boundary entropy's thermal energy is a classical quantity that predates the trace-out (does gravitate). The boundary modes are physically localized on the horizon — the uniform-medium description used in the shell theorem calculation is a gravitational equivalence, not a claim about physical delocalization. The usefulness of the uniform-medium description is computational: when matter is present, it locally deforms the horizon geometry and redistributes the boundary entropy, with the gravitational effect of the redistribution on bulk observers being most easily computed in the uniform-medium representation.

The MOND acceleration scale. The framework's account of dark-matter phenomenology — flat rotation curves, the baryonic Tully-Fisher relation, the radial acceleration relation — derives from entropy displacement at the cosmological boundary. The argument has four steps, all of which inherit the conditions on the de Sitter temperature (H-slope, H-frame, H-Hawking; see [GR §7.3]) and the assumption that a static configuration delivers its full rest energy $Q = M_B c^2$ as heat to the boundary — a coupling statement belonging to the open G3 map rather than a consequence of C1.

Step 1: Entropy displacement by baryonic matter (conditional). The chain rests on the de Sitter temperature T_{dS}, which inherits the conditions on the gap equation (H-slope with the horizon and frame conditions, Section 7.3), and on the assumption that the entire rest energy $Q = M_B c^2$ of a static configuration is delivered as heat to the boundary — a statement about how the emergent sector couples to the boundary, which the open covariant coupling theorem ([GR §6.2]) would have to supply. With these, the Bekenstein bound applied to a baryonic mass M_B in a de Sitter background gives the entropy displaced from the cosmological boundary:

$$\Delta S = \frac{M_B c^2}{k_B T_{\mathrm{dS}}} = \frac{2\pi M_B c^2}{\hbar H}.$$

This is the Clausius relation applied to the boundary reservoir: the boundary's entropy reduction is Q/T for a reversible process, where $Q = M_B c^2$ and $T = T_{\mathrm{dS}}$.

Step 2: The critical acceleration $a_0 = cH/6$ (conditional; dimensional factor imported). The displaced entropy redistributes over the volume between the matter and the horizon. The Jacobson mechanism converts the entropy gradient into additional radial acceleration. The conversion is dimensional, and the factor is Verlinde's: in d spacetime dimensions, the factor relating the natural Hubble acceleration scale cH to the empirical MOND acceleration is $(d-3)/[(d-2)(d-1)]$, which follows in his emergent-gravity analysis from the interplay between volume-law entropy distributed over $(d-1)$ spatial dimensions and area-law entropy scaling with $(d-2)$ surface dimensions, together with the standard surface-density-to-acceleration relation via Gauss's law. In our universe with $d = 4$:

$$a_0 = \frac{d-3}{(d-2)(d-1)} \cdot cH = \frac{1}{2 \cdot 3} \cdot cH = \frac{cH}{6} = 1.09 \times 10^{-10}\,\mathrm{m/s^2} \quad (H_0 = 67.4\,\mathrm{km\,s^{-1}\,Mpc^{-1}}).$$

Under the stated coupling and horizon conditions, the imported dimensional formula gives the factor $1/6$ at $d = 4$; with that formula, the acceleration scale uses H without an additional fitted parameter. The empirical value from the radial-acceleration-relation fit is $(1.20 \pm 0.02 \pm 0.24) \times 10^{-10}$ m/s^2, so the prediction sits at 0.91 of it for the Planck H_0 and 0.98 for $H_0 = 73$ km s^{-1} Mpc^{-1} — inside the fit's stellar mass-to-light systematic, a consistency at the ten-percent level rather than an exact match.

Step 3: The deep-MOND regime and baryonic Tully-Fisher. For baryonic acceleration much smaller than a_0, the displaced entropy dominates the dynamics. The Jacobson mechanism converts the entropy gradient into curvature, giving $g_{\mathrm{total}} = \sqrt{g_B \cdot a_0}$ where $g_B = GM_B/r^2$ is the baryonic Newtonian acceleration. The rotation velocity at radius r then satisfies $v^2 = r \cdot g_{\mathrm{total}} = \sqrt{GM_B \cdot cH/6} =$ constant, so:

$$v^4 = GM_B \cdot \frac{cH}{6}.$$

This is the baryonic Tully-Fisher relation. Flat rotation curves emerge with no dark-matter particles and no free parameters beyond the baryonic mass.

Step 4: Redshift evolution. The MOND scale depends on the Hubble parameter, so $a_0(z) = cH(z)/6$ increases with redshift. The MOND crossover radius $r_M = \sqrt{6GM_B/(cH)}$ correspondingly shrinks at high redshift: for a Milky-Way-like baryon budget ($M_B \approx 6 \times 10^{10}\,M_\odot$ for stellar plus cold gas), $r_M = 8.8$ kpc at $z = 0$ but 5.0 kpc at $z = 2$ and 3.0 kpc at $z = 5$. Galaxies with effective radii comparable to or exceeding r_M show flat rotation curves; galaxies with $r_{\mathrm{eff}} < r_M$ decline, Keplerian-like, inside r_M and approach flat beyond it. At fixed radius the baryonic share of the acceleration, $1/\nu(g_B/a_0)$, falls as a_0 rises, so high-redshift galaxies are less baryon-dominated than local spirals with the same baryon distribution, and their rotation curves reach the flat deep-MOND form at smaller radii.

Observational comparisons. Two observations can be compared with the conditional dark-sector account, and both are in tension with it.

Jiao et al. (2023) detected a Keplerian decline in the Milky Way's rotation curve from approximately 19 to 26.5 kpc using Gaia DR3 kinematics, with the flat rotation curve hypothesis rejected at 3σ significance. The conditional formula gives $r_M \approx 16$ kpc using a baryon budget that includes the hot circumgalactic gas halo ($M_B \sim 2 \times 10^{11}\, M_\odot$), and $r_M \approx 9$ kpc using stellar plus cold gas ($M_B \sim 6 \times 10^{10}\, M_\odot$). The observed decline lies beyond either crossover, where the law gives a nearly flat curve: $v(26.5\,\mathrm{kpc})/v(19.5\,\mathrm{kpc}) = 0.97$ or 0.95 for the two budgets treated as point masses, against 0.86 for a Keplerian decline, and at least 0.95 in the spherical approximation for any baryon distribution up to $2 \times 10^{11}\, M_\odot$. A quantitative comparison needs the Milky Way baryon model and the circular velocities of Jiao et al.'s Jeans analysis.

The high-redshift test is Genzel et al., who report stacked rotation curves for massive star-forming galaxies at $z = 0.9$–2.4 showing declining outer velocities at greater than 3σ significance relative to local spirals. At a fixed baryonic mass profile the framework's law predicts the opposite trend: as the crossover radius r_M shrinks, a galaxy becomes less baryon-dominated and its outer curve flatter. Real high-redshift galaxies also differ from local ones in mass distribution and pressure support, so the observed declines are an apparent tension with the law, which a comparison modelling the galaxies' own baryon profiles and the pressure support that Genzel et al. invoke alongside baryon dominance has to settle; that comparison is not yet available. In the ΛCDM paradigm, flat rotation curves from NFW halos are expected at all redshifts; explaining the Genzel data within ΛCDM requires anomalously low halo concentrations or baryon-dominated inner regions.

The interpolation function and solution-specific content. The framework's structural commitments fix the deep-Newtonian and deep-MOND limits and the redshift evolution of the crossover, but the full interpolation between the two regimes is constrained rather than uniquely determined. Five constraints — no free parameters, conservative, monotonic, correct limits, analytic — narrow the interpolation to a one-parameter family indexed by the transition steepness. The transition steepness is a property of the particular bijection φ, in the same Tier 3 category as fermion masses (solution-specific rather than structural). The simplest member satisfying all five constraints is the "simple" MOND interpolation function $\nu(y) = (1 + \sqrt{1 + 4/y})/2$, which matches the empirical radial acceleration relation of McGaugh et al. to better than 0.1 dex across all galaxy-relevant accelerations.

The framework's content in the dark-matter sector therefore has two-tier structure. The MOND acceleration scale $a_0 = cH/6$, the baryonic Tully-Fisher relation, and the redshift evolution of the crossover are Tier 2 — universal across bijections in the framework's universality class, inheriting the conditions of Step 1 and the imported factor of Step 2. The transition steepness is Tier 3 — bijection-specific, with the simple MOND interpolation as the framework's natural default.

7.10 Summary of gravitational predictions and chapter close

The gravitational sector combines unconditional partition-geometry results with conditional horizon calibration and physical coupling claims. The action-scale and entropy-coefficient relations require the horizon, frame and state conditions; the physical dark-sector interpretation requires the matter-to-boundary coupling map; and the Page curve requires H-scramble and H-decouple, of which H-scramble fails for every bijection acting on an initial configuration with the bipartition aligned to the configuration coordinates. The table separates those dependencies from empirical comparisons, normalization identities and formulas awaiting tests.

Summary table.

Observable	Predicted	Observed	Match	Tier
Area law $dA \geq 0$	classical area theorem (GR horizon dynamics, preserved)	GW250114 area increase	99.999% confidence	1
RVM dark energy form	not derived (local expansion shared by every smooth model)	DESI DR2 + Bertini fit	non-discriminating	—
$\nu_{\rm OI}$ magnitude	$\sim 10^{-32}$ (indistinguishable from 0)	$\nu^{\rm Bertini} = -(2.5 \pm 1.3) \times 10^{-4}$	2σ	2
Dark-sector budget	$\sim 95\%$ under the conditional physical interpretation	$\Omega_{\rm dark} \approx 0.95$	partly definitional	2 (cond.)
$a_0 = cH/6$	conditional entropy displacement; factor 1/6 imported	$(1.20 \pm 0.02 \pm 0.24) \times 10^{-10}\,{\rm m/s}^2$	0.91 at the Planck H_0, inside the systematic	2 (cond.)
$v^4 = GM_B \cdot cH/6$	conditional baryonic Tully-Fisher relation	empirical relation	consistent relation	2 (cond.)
$r_M(z)$ shrinkage	$\propto 1/\sqrt{H(z)}$, flatter outer curves at a fixed baryon profile (conditional)	Genzel et al. $z = 0.9$–2.4, $> 3\sigma$ decline	in tension	2 (cond.)

Observable	Predicted	Observed	Match	Tier
Milky Way outer curve	nearly flat beyond $r_M \sim 9$–16 kpc, depending on baryon budget	Jiao et al. Keplerian decline, ~ 19–26.5 kpc	in tension	2 (cond.)
No dark-matter particle	conditional dark-sector interpretation	XENON, LUX, etc. null	consistent	2 (cond.)
No GUT-mediated proton decay	$\tau_p \sim 10^{45}$–10^{46} yr	$\tau_p > 10^{34}$ yr	consistent	1

Three horizon results are consistency relations rather than predictions and are stated here rather than tabulated. The calibration $\hbar = c^3\epsilon^2/(4G\sigma)$, equivalently $\epsilon = 2\sqrt{\sigma}\, l_p$, conditional on H-slope, is a self-consistency condition that fixes only the ratio $\epsilon/\sqrt{\sigma} = 2\, l_p$, not an independent determination of $\hbar$ (Sections 7.3–7.4). The Bekenstein-Hawking entropy $S_{\rm BH} = A/(4l_p^2)$ carries the factor 1/4 as the ratio $2\pi/8\pi$ that the calibration matches, conditional on H-slope, with no direct test (GW250114 tests the area theorem, in the table). The descriptive vacuum-energy distinction carries the ratio $\rho_{\rm QFT}/\rho_{\rm class} \sim S_{\rm dS}$, a dimensional identity up to a convention-dependent factor, with the dissolution conditional on the coupling theorem.

The tier system per Section 7.1: Tier 1 results follow from partition geometry alone and hold for any embedded-observer system satisfying C1–C4 at its cosmological horizon; the results built on them — $\hbar$, $\epsilon = 2\sqrt{\sigma}\, l_p$, the 1/4 coefficient — are conditional on H-slope with the horizon and frame conditions, and the running-vacuum form is not derived. Tier 2 results follow from the thermodynamic Clausius relation applied to the boundary reservoir and the geometric factor from spherical redistribution in de Sitter space, inheriting those conditions. Tier 3 results — the transition steepness in MOND interpolation, the greybody factor for Hawking radiation — depend on the specific bijection and are addressed by the cubic-lattice realization developed in Chapters 5-6.

Falsification commitments. These tests concern the stated physical interpretation and its conditions. Any measured deviation from $S = A/(4\, l_p^2)$ at sufficient confidence to exclude experimental systematics would refute the gap-equation chain if its named horizon, frame and state conditions hold — the consistency between classical horizon temperature and Euclidean KMS temperature that produced the 1/4 coefficient. High-confidence detection of phantom crossing in the dark-energy equation of state — DESI DR2 currently shows 3–4σ mild tension that DESI Year 5 will resolve — would contradict the framework's Type II reading and its expectation $w(z) \approx -1$ — the strongest available disconfirmation in the dark-energy sector; the framework supplies the smooth

$H^2 - H_0^2$ bookkeeping but does not derive a distinctive Type II law (§7.8). A high-confidence direct detection of a dark-matter particle at non-gravitational interaction strength would refute this dark-sector interpretation. Discovery of GUT-mediated proton decay would refute the framework's no-GUT structural commitment from Chapter 6.

The gravitational entries have different empirical roles. The Milky Way decline is an observation to compare with the conditional, baryon-budget-dependent crossover formula. The GW250114 area-theorem result is a consistency at 99.999% confidence and does not distinguish the framework from standard GR. The calibration identities and untested horizon formulas are not independent precision measurements.

The geometry-first account. The classical and emergent descriptions distinguish vacuum-energy bookkeeping without deriving its dynamical gravitational decoupling, which remains a coupling obligation. The $\hbar$, ϵ and $1/4$ relations follow under the named horizon conditions. The physical dark-sector interpretation inherits those conditions and the open heat-transfer map; its total fraction is partly the input-dependent quantity $1 - \Omega_B$. The MOND formula uses an imported dimensional factor and agrees within the quoted systematics. Its Tully-Fisher and crossover consequences remain conditional. These results have distinct source obligations and empirical standing.

The comparison between descriptions concerns their explanatory chains and physical assumptions. A descriptive distinction between vacuum energies does not settle their dynamical coupling, and an unobserved entropy coefficient is not an empirical confirmation. Forward tests must identify the conditions being tested and the measurements that distinguish the proposed physical interpretation.

Forward pointers. Chapter 8 develops the JUNO neutrino prediction in detail — the post-November 2025 retrodictive match of $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2)$ at 0.07σ, the framework's sharpest neutrino-sector empirical match. The chapter develops the experimental context the matter-sector treatment in Chapter 6 abbreviated: the JUNO detector design, the reactor antineutrino source, the global-fit methodology, and the framework-specific significance of the $\sin^2\theta_{12}$ measurement.

Chapter 9 develops the framework's relationship to other unification programs — the OI-string relationship at the matrix-model level, the no-GUT structural commitment against the broader landscape of grand unified theories, the universality-class character of the framework's Tier 1 results, and the de Sitter observer literature's convergence with the framework's geometry-first dissolution. The empirical record of Chapters 5-8 is concentrated at the Level G3 × Level D intersection of Chapter 3's hierarchical structural realism; Chapter 9 locates the framework within the broader space of substratum-level theories and articulates the universality classes that the framework's structural commitments belong to.

The fundamental-physics inventory includes twenty-two classified Standard Model observables, the ten entries in this chapter's table, and seven neutrino-sector entries. It combines conditional formulas, numerical retrodictions, observational bounds and targets awaiting data. Part III turns to the emergence cascade through chemistry, life and evolution; Part IV develops applications and further testable claims.

Chapter 8

JUNO and the Neutrino Sector

8.1 What this chapter develops

Chapter 6 §6.7 introduced the framework's predictions for the three PMNS mixing angles and identified the solar mixing angle $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2) = 0.3080$ as the framework's sharpest neutrino-sector empirical match — a parameter-free retrodiction whose provenance §8.6 states plainly, matched to the post-JUNO global fit at 0.07σ. The present chapter develops that prediction in detail. The JUNO experiment, the structural derivation, the global-fit methodology, the discrimination from competing tribimaximal-modification patterns, and the framework's empirical position in the post-JUNO landscape — these are the chapter's content.

The chapter opens with the experiment. Section 8.2 develops the JUNO detector design and the reactor antineutrino source: the 20-kiloton liquid scintillator detector 53 km from two nuclear power complexes in southern China, the energy resolution and reactor flux that combine to produce the world's most precise measurement of $\sin^2\theta_{12}$ on a much shorter timescale than was thought possible. Section 8.3 develops the cubic-group flavor structure that produces the framework's prediction: the three lepton generations occupying the T_1 corners of the cubic Brillouin zone, the Cabibbo angle $\lambda = 1/(\pi\sqrt{2})$ from a single Brillouin-zone distance, the tribimaximal mixing pattern from the $A_4 \subset O$ subgroup at zeroth order.

The structural derivation occupies the middle of the chapter. Section 8.4 develops the U-parity grading — the discrete μ-τ exchange symmetry of $S_4 \supset A_4$ that controls how the charged-lepton perturbations modify the tribimaximal pattern at order λ^2. Section 8.5 develops the sum rule $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$ as the framework's structural discriminator from tribimaximal-modification alternatives, and locates the framework's specific structural relation $b_{23} = (4/3)(b_{12} + b_{13})$ within the broader space of partial-TBM patterns currently analyzed against JUNO data.

The empirical content occupies the latter sections. Section 8.6 develops the three numerical predictions — $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2) = 0.3080$, $\sin^2\theta_{13} = 4/(18\pi^2) = 0.0225$, $\sin^2\theta_{23} = 1/2 + 1/(2\pi^2) = 0.5507$ — and compares them with the post-JUNO global fit and the NuFIT 6.0 best fits, with all three angles matched within 1.1σ. Section 8.7 compares the framework's predictions with the column-preservation patterns TM1 and TM2 currently analyzed against JUNO: TM2 excluded at greater than 3.5σ post-JUNO, TM1 marginal at 1.35σ, the framework at 0.07σ.

The framework's position in the literature is the chapter's closing content. Section 8.8 develops the JUNO design-lifetime sensitivity: 30-year exposure projected to reach ± 0.0014 precision on $\sin^2\theta_{12}$, providing a sub-percent test of the framework's prediction. Section 8.9 develops the coupled neutrino-mass / dark-energy result with DESI DR2 — the $\Sigma m_\nu < 0.0642$ eV bound consistent with the framework's normal-ordering prediction near the oscillation minimum of 0.059 eV, the 3σ tension with the lower oscillation limit assuming ΛCDM that the framework's RVM dark energy would relax, though the corrected bound is not yet computed. Section 8.10 closes with the chapter's epistemic implications: a 0.07σ empirical match on a parameter-free prediction that is layered conditional (Class B-L) pending the derivation of Cond 2, with the chapter's content classified per Chapter 4's discipline.

The framework's empirical record in the neutrino sector represents the strongest single piece of evidence for the framework's structural commitments outside the gravitational sector. The Cabibbo angle and the Koide relation match observation at 0.04% and 0.001% respectively in the quark and charged-lepton sectors. The $\sin^2\theta_{12}$ match at 0.07σ extends the same cubic-group machinery to the neutrino sector, with no fresh free parameters fitted — the same inputs $\lambda^2 = 1/(2\pi^2)$ and $A^2 = 2/3$ that produced the quark-sector matches produce the neutrino-sector match. The framework's neutrino-sector content is therefore concentrated in this chapter as a focused case study of cross-sector parameter-free prediction at high precision.

8.2 The JUNO experiment

The Jiangmen Underground Neutrino Observatory (JUNO) is a multipurpose neutrino experiment located 700 meters underground in Jiangmen, southern China. The detector consists of a 20-kiloton liquid scintillator sphere, approximately 35 meters in diameter, instrumented with 17,612 large-aperture photomultiplier tubes and 25,600 smaller photomultiplier tubes. The detector is designed to measure reactor antineutrinos from the Yangjiang and Taishan nuclear power complexes, located 53 km from the experimental site — a baseline chosen to maximize sensitivity to the solar mixing angle θ_{12} at the oscillation maximum for reactor antineutrino energies.

The detector's energy resolution is the principal experimental innovation. JUNO achieves approximately 3% at 1 MeV, a factor of three to five improvement over

previous reactor neutrino experiments. The improvement comes from three sources: the high photocathode coverage (over 75% of the detector surface), the high light yield of the scintillator, and the careful calibration program that has been ongoing throughout the detector commissioning. The combination of high statistics (approximately 60 events per day from the two reactor complexes) and high energy resolution allows JUNO to resolve the fine structure of the reactor antineutrino oscillation spectrum at unprecedented precision.

The first JUNO measurement. The collaboration reported its first measurement of reactor neutrino oscillations in November 2025. The headline result from 59.1 days of physics data was

$$\sin^2\theta_{12} = 0.3092 \pm 0.0087,$$

which is the most precise single-experiment determination of the solar mixing angle to date — a factor of 1.6 improvement over all previous measurements combined. The same dataset also produces a high-precision measurement of the mass-squared splitting:

$$\Delta m^2_{21} = (7.48 \pm 0.10) \times 10^{-5}\,\mathrm{eV}^2,$$

confirming the solar oscillation parameters with reactor-baseline precision for the first time. JUNO's first measurement closes a long-standing tension between solar and reactor neutrino measurements of θ_{12}: the solar neutrino determination, primarily from SNO and Super-Kamiokande, gave systematically higher values, while the reactor neutrino determination from KamLAND gave systematically lower values, with the discrepancy at the 1.5σ level before JUNO. The new JUNO result lies between the two prior determinations and provides a unified value with significantly reduced uncertainty.

The post-JUNO global fit. The combined analysis of all neutrino oscillation data — solar, atmospheric, reactor, and accelerator — produces a global fit that incorporates the new JUNO measurement. The Capozzi *et al.* post-JUNO global fit (early 2026) gives

$$\sin^2\theta_{12} = 0.3085 \pm 0.0073,$$

which is the value the framework's predictions are tested against in this chapter. The global fit's uncertainty is dominated by JUNO's measurement; future runs will continue to tighten the constraint over the experiment's design lifetime.

The JUNO experiment is therefore the primary current source of constraint on the solar mixing angle and on the framework's specific prediction. The remainder of this chapter develops what that prediction is, where it comes from structurally, and how it compares with alternative theoretical proposals currently being analyzed against the JUNO data.

8.3 The cubic-group flavor structure

The framework's prediction for the PMNS angles follows from the cubic point group O — the rotation symmetry group of three-dimensional space — applied

to the three lepton generations. The structural argument runs through the same chain that produced the gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ in Chapter 5: the cubic decomposition of the six nearest-neighbor link directions on the simple cubic lattice gives multiplicities $(3, 2, 1)$ under the cubic-group action, with the three lepton generations occupying the T_1 triplet representation.

The cubic group O and its A_4 subgroup. The cubic point group O is the rotation symmetry group of three-dimensional space, with 24 elements isomorphic to the symmetric group S_4. Its irreducible representations decompose as

$$\mathrm{irreps}(O) = A_1 \oplus A_2 \oplus E \oplus T_1 \oplus T_2,$$

of dimensions $1, 1, 2, 3, 3$. The subgroup $A_4 \subset O$ keeps $\{A_1, A_2, E, T_1\}$ but identifies A_2 with A_1, giving the standard A_4 content $A \oplus A' \oplus A'' \oplus T$ of dimensions $1, 1, 1, 3$. The framework's structural derivation in Chapter 5 placed the three lepton generations on the T_1 representation as one of the irreducible components of the cubic decomposition $6 = T_1 \oplus E \oplus A_1$ of the six nearest-neighbor link directions.

The realization of T_1 in the framework is concrete. The three Cartesian unit vectors $\{e_1, e_2, e_3\}$ — corresponding to the three spatial axes of the cubic lattice — are acted on by the 24 rotations of O. The three candidate sectors — the generations, under H-spin' — occupy these three Cartesian directions. The cubic decomposition has a single democratic direction

$$\hat{h} = \frac{1}{\sqrt{3}}(e_1 + e_2 + e_3),$$

which is the only direction fixed by the $\mathbb{Z}_3$ cyclic subgroup of O. The plane perpendicular to $\hat{h}$ carries the doublet E representation. This is the geometric origin of tribimaximal mixing in the framework: the second column of the tribimaximal mixing matrix is exactly $\hat{h}$, and the first and third columns lie in the plane perpendicular to it, oriented by μ-τ exchange.

The Cabibbo scale from cubic Brillouin geometry. The perturbation parameter λ controlling the deviations from tribimaximal mixing in the framework is identified with the Cabibbo angle, fixed independently by the cubic Brillouin-zone geometry as developed in Chapter 6 §6.5. The three generations occupy the three T_1 corners of the cubic Brillouin zone at momenta $X_j = \pi \cdot e_j$ for $j = 1, 2, 3$. The inter-generation distance in momentum space is

$$|X_i - X_j| = \pi\sqrt{2} \quad (i \neq j),$$

a geometric constant of the cubic lattice. The Cabibbo angle is the inverse of this distance:

$$\lambda = \frac{1}{\pi\sqrt{2}} = 0.22508.$$

The match to observation is at 0.04%: the measured value is $\lambda = 0.22500 \pm 0.00067$, with the framework's prediction inside the experimental uncertainty.

The Cabibbo angle is therefore a structural prediction of the framework, fixed by a single Brillouin-zone distance with no input from the lepton sector; the geometry and the chirality-forced power are unconditional, while the identification of the mixing element with the unit-normalized propagator is a stated hypothesis whose resolving computation is specified in [SM §7.1]; the hypothesis has since been structurally narrowed — the free-level mixing object vanishes exactly and the substrate-kinematics alternative is excluded, so the identification is equivalent to the emergent-theory coefficient being unity — and the on-shell computation has since answered it: between physically projected legs, every spin-resolved taste-changing element has unit modulus exactly, confirming the unity condition at its structural level as an algebraic identity of the spin-taste reconstruction ([SM §7.1]); what remains beyond that level is the dressing. The fact that the same λ then controls the PMNS deviations from tribimaximal mixing is the framework's structural commitment: the Cabibbo scale that produces the quark-sector mixing is the same scale that produces the lepton-sector mixing deviations. The two empirical confirmations — the Cabibbo angle at 0.04% and the solar mixing angle at 0.07σ — are not independent confirmations of separate structures but joint confirmations of a single cubic-group geometric source.

The PMNS matrix at zeroth order. The framework's PMNS structure at zeroth order is the tribimaximal mixing matrix, obtained from the $A_4 \subset O$ representation theory. The tribimaximal matrix is

$$U_{\mathrm{TBM}} = \begin{pmatrix} \sqrt{2/3} & \sqrt{1/3} & 0 \\ -\sqrt{1/6} & \sqrt{1/3} & \sqrt{1/2} \\ -\sqrt{1/6} & \sqrt{1/3} & -\sqrt{1/2} \end{pmatrix},$$

with the columns corresponding to mass eigenstates ν_1, ν_2, ν_3 and the rows to flavor eigenstates ν_e, ν_μ, ν_τ. The corresponding zeroth-order mixing angles are

$$\sin^2\theta_{12}^{\mathrm{TBM}} = \frac{1}{3}, \quad \sin^2\theta_{23}^{\mathrm{TBM}} = \frac{1}{2}, \quad \sin^2\theta_{13}^{\mathrm{TBM}} = 0.$$

The observed values deviate from tribimaximal mixing by amounts of order $\lambda^2 \approx 0.05$, with the most consequential deviation being a non-zero θ_{13} at the level of $\sin^2\theta_{13} \approx 0.022$. The framework's content in the next sections is to derive these deviations from a μ-τ parity grading combined with one cubic-group structural relation, producing parameter-free predictions for all three angles.

The framework's structural commitment. The cubic-group flavor structure used here is not a postulated symmetry of the lepton sector. It is the rotation symmetry group of three-dimensional space, realized in the framework's lattice substratum as developed in Chapters 5 and 6. The same cubic group that produced the Standard Model gauge structure produces the lepton flavor structure here, with the T_1 generations occupying the same triplet representation in both contexts. The empirical record across the framework's flavor predictions

— Cabibbo at 0.04%, the Koide angle at 0.001%, $\sin^2\theta_{12}$ at 0.07σ, the sum rule $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$ at 0.6σ — tests the same cubic structure across multiple observables at sub-percent precision. This is testable structure rather than a postulated symmetry: the same numerical inputs ($\lambda^2 = 1/(2\pi^2)$, the projection factor $A^2 = 2/3$) enter multiple observables with all of them matching observation.

8.4 The U-parity grading

The framework's PMNS structure beyond tribimaximal mixing develops through a μ-τ exchange parity — a discrete $\mathbb{Z}_2$ grading from $S_4 \supset A_4$ — that organizes the charged-lepton perturbations into structurally distinct U-even and U-odd components. The grading is not an additional symmetry postulate but a direct consequence of the framework's structural commitment to T_1 as the generation triplet: T_1 contains a natural μ-τ exchange generator, and the perturbations of T_1 inherit a grading under this exchange.

The U-parity generator. In the Altarelli-Feruglio basis (charged-lepton mass matrix diagonal at zeroth order), the μ-τ exchange operator U is the permutation that exchanges the second and third generations:

$$U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

The generator U sits in $S_4 \supset A_4$ as the μ-τ exchange element of the larger permutation group. Conjugation by U defines a $\mathbb{Z}_2$ grading on antisymmetric matrices: a matrix A is U-even if $UAU^{-1} = +A$, U-odd if $UAU^{-1} = -A$. The grading decomposes the space of antisymmetric matrices into two orthogonal components.

With the basis matrices $(E_{ij})_{kl} = \delta_{ik}\delta_{jl} - \delta_{il}\delta_{jk}$ — the elementary antisymmetric matrices spanning the space of 3×3 antisymmetric matrices — the decomposition is: - U-even combinations: $E_{12} + E_{13}$ - U-odd combinations: $E_{12} - E_{13}$ and E_{23}

The unique U-odd combination antisymmetric in (μ, τ) is E_{23}; the unique U-odd combination relating the first generation to the second and third is $E_{12} - E_{13}$. The U-even combination $E_{12} + E_{13}$ relates the first generation symmetrically to the other two.

The charged-lepton perturbation expansion. The charged-lepton rotation matrix V_ℓ in the Altarelli-Feruglio basis admits a perturbative expansion in the Cabibbo scale λ:

$$V_\ell = \exp(\lambda A_1 + \lambda^2 A_2 + O(\lambda^3)),$$

with A_1 and A_2 real antisymmetric 3×3 matrices to be determined from structural inputs. The expansion is the framework's parametrization of the charged-

lepton rotation: A_1 controls the leading deviations from the tribimaximal pattern at order λ, A_2 controls the next-to-leading deviations at order λ^2.

Determination of A_1. The leading-order generator A_1 is fixed by three structural conditions plus the cubic geometric input. First, the observed value $\theta_{13} \neq 0$ requires $\sin\theta_{13} = O(\lambda)$, forcing A_1 to contribute to the $(1,3)$ element of $V_\ell^\dagger U_{\mathrm{TBM}}$. Second, the empirical observations that both $\Delta_{12} \equiv \sin^2\theta_{12} - 1/3$ and $\Delta_{23} \equiv \sin^2\theta_{23} - 1/2$ are of order λ^2 — neither shows a leading λ contribution — further constrains A_1's structure. Third, the geometric input from Section 8.3: each end of an inter-generation mixing vertex projects onto the plane perpendicular to the democratic direction $\hat{h}$, contributing a factor $\sin(\angle(\hat{e}_i, \hat{h})) = A$ from the off-democratic projection of any generation axis. For a single-vertex inter-generation transition, the total projection factor is A^2 — one factor per vertex end.

Identifying this projection geometry with the leading $O(\lambda)$ contribution to U_{e3} gives the structural identity

$$\sin\theta_{13} = A^2\lambda,$$

exact at order λ (the order-λ^2 contributions from A_2 cancel in $|U_{e3}|^2$, as the next section establishes). This is a structural input fixing A_1's overall coefficient up to sign, combined with the U-odd character required by the constraint $\Delta_{12}, \Delta_{23} = O(\lambda^2)$.

The result is uniquely determined:

$$A_1 = \frac{\sqrt{2}}{3}(E_{12} - E_{13}).$$

The generator A_1 is purely U-odd: $UA_1U^{-1} = -A_1$. Geometrically, A_1 represents a rotation by angle $A^2 = 2/3$ around the axis $(\hat{e}_2 + \hat{e}_3)/\sqrt{2}$, which reproduces $\sin\theta_{13} = A^2\lambda$ as required. The structural property is that A_1 is U-odd; the magnitude $\sqrt{2}/3$ is fixed by the projection identity $\sin\theta_{13} = A^2\lambda$ with $A^2 = 2/3$.

Deviations from TBM at order λ^2. Expanding V_ℓ to second order:

$$V_\ell = I + \lambda A_1 + \lambda^2(A_2 + \tfrac{1}{2}A_1^2) + O(\lambda^3),$$

and writing the most general antisymmetric A_2 as

$$A_2 = b_{12}E_{12} + b_{13}E_{13} + b_{23}E_{23},$$

direct computation of $|U_{\mathrm{PMNS},ij}|^2$ to order λ^2 gives the deviations from tribimaximal mixing:

$$\Delta_{13} = \tfrac{4}{9}\lambda^2 = A^4\lambda^2,$$
$$\Delta_{12} = -\tfrac{2}{3}(b_{12} + b_{13})\lambda^2,$$
$$\Delta_{23} = b_{23}\lambda^2.$$

Three structural properties of these deviations deserve explicit comment. First, Δ_{13} is independent of A_2 — the structural identity $\sin\theta_{13} = A^2\lambda$ is exact at order λ^2, with the A_2 contributions canceling from $|U_{e3}|^2$. The reactor mixing angle prediction is therefore Layer 1 structural: $\sin^2\theta_{13} = (4/9)\lambda^2$ follows from A_1 alone, with no second-order Wilson-coefficient inputs. Second, the combination $b_{12} - b_{13}$ does not enter any of the three angles — it is naturally identified with the CP-violating Dirac phase $\delta_{\rm CP}$, the only remaining free parameter at order λ^2 that does not enter the angles. Third, Δ_{23} depends only on b_{23}, not on the U-even combination $b_{12} + b_{13}$. The U-parity structure organizes the second-order corrections in a way that the standard effective-field-theory parametrization does not.

8.5 The sum rule and structural relation

The deviations Δ_{12} and Δ_{23} depend on two independent scalar combinations of the A_2 Wilson coefficients: the U-even combination $b_{12}+b_{13}$ controlling Δ_{12}, and the dominant U-odd coefficient b_{23} controlling Δ_{23}. The framework's content beyond the Layer 1 reactor angle prediction is a structural relation between these two combinations — a relation that produces the sum rule $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$ and discriminates the framework from tribimaximal-modification alternatives.

Two normalization conditions. The framework fixes the second-order Wilson coefficients through two conditions.

Condition 1 — natural normalization. The overall scale of A_2 is fixed by taking $b_{23} = 1$. This is a labeling convention rather than a physical input: it sets A_2 to enter the perturbation series at unit strength relative to λ^2, fixing the normalization of the order-λ^2 expansion before $b_{12} + b_{13}$ is determined.

Condition 2 — cubic-group structural relation. The relation between the dominant U-odd coefficient and the U-even combination takes the form of a cubic-group identity, its magnitude at Layer 2(b) (§8.5):

$$b_{23} = \tfrac{4}{3}(b_{12} + b_{13}).$$

With Condition 1, this gives $b_{12} + b_{13} = 3/4$. Unlike Condition 1, this is a substantive physical input — a relation between two independent Wilson coefficients, which the substrate is not shown to produce (§8.5).

Equivalence with TBM sum rule preservation. The framework's structural relation has a clean interpretation at the angle level. Combining the explicit results $\Delta_{12} = -(2/3)(b_{12} + b_{13})\lambda^2$ and $\Delta_{23} = b_{23}\lambda^2$:

$$2\Delta_{12} + \Delta_{23} = (b_{23} - \tfrac{4}{3}(b_{12} + b_{13}))\lambda^2.$$

Condition 2 is therefore equivalent to $2\Delta_{12} + \Delta_{23} = 0$, which is the statement that the tribimaximal sum rule $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$ — holding at zeroth order from cubic geometry (since $2(1/3) + 1/2 = 7/6$) — continues to hold at

order λ^2. The Wilson-coefficient relation and the angle-level sum rule are the same content under different parametrizations.

The sum rule itself has a clean cubic-geometric reading: $7/6 = 2/d+1/(d-1)$ for $d = 3$ generations, with $2/d$ being twice the inverse generation count (the value $\sin^2\theta_{12}^{\mathrm{TBM}} = 1/3$ doubled) and $1/(d-1)$ being the inverse partner count (the value $\sin^2\theta_{23}^{\mathrm{TBM}} = 1/2$ in the doublet sector). For three generations, $2/3+1/2 = 7/6$ is forced by the representation theory; the framework's content is that this sum rule is *preserved* at order λ^2 by the second-order Wilson-coefficient relation.

The factor $4/3$ as cubic-group identity. Condition 2's specific numerical coefficient is a genuine cubic-group identity:

$$\frac{4}{3} = 2A^2 = 2 \cdot \frac{C_2(T_1)}{d} = \frac{C_2(T_1)^2}{d},$$

where $A = \sqrt{2/3}$ is the off-democratic projection of any generation axis and $C_2(T_1) = d - 1 = 2$ is the quadratic Casimir of the T_1 representation. The identity $A^2 = (d-1)/d = C_2(T_1)/d$ for $d = 3$ is the same representation-theoretic identity from Section 6.5 that produced the Wolfenstein A parameter. The factor $4/3$ thus has an interpretive reading as partner-count $\times$ per-partner projection: the U-even combination $E_{12} + E_{13}$ couples the first generation to its two partners $\{\hat{e}_2, \hat{e}_3\}$, with the per-partner projection nominally A^2. The interpretation is internally consistent but non-directional — the same arithmetic produces the inverse ratio $3/4$ under alternative partner-count assignments. The directional assertion that the U-odd coefficient b_{23} is *larger* than the U-even coefficient $b_{12}+b_{13}$ by this specific factor is what carries the framework's substantive content beyond the cubic-group identity itself.

Structural protection of the sum rule by A_1 alone. A distinct structural feature of the framework's A_1 choice deserves explicit treatment. With $A_2 = 0$, direct computation gives $2|U_{e2}|^2+|U_{\mu 3}|^2 = 7/6+\lambda^2\cdot[-14/27]$ before the $\cos^2\theta_{13}$ correction. Dividing by $\cos^2\theta_{13} = 1 - (4/9)\lambda^2$ — which itself follows from $\sin^2\theta_{13} = A^4\lambda^2$, a Layer 1 quantity — precisely cancels the $-14/27$ offset, giving

$$(2\sin^2\theta_{12} + \sin^2\theta_{23})\,\big|_{A_2=0} = 7/6 + O(\lambda^4).$$

The sum rule is therefore *structurally protected* against A_1's perturbation by cubic geometry alone — no substrate input required for this component. The cancellation between the $-14/27$ contribution from A_1^2-induced shifts and the $+14/27$ contribution from the $\cos^2\theta_{13}$ dressing is automatic given A_1's Layer 1-forced form. Condition 2's content reduces to whether A_2 *preserves* the sum rule that A_1 alone protected, or breaks it. The framework's structural commitment is that A_2 also preserves the sum rule, through the $b_{23} = (4/3)(b_{12}+b_{13})$ relation.

Layer status. The three angle predictions have distinct structural statuses per the four-layer framing of Chapter 4 §4.5. The reactor angle $\sin^2\theta_{13} = (4/9)\lambda^2 = A^4\lambda^2$ is Layer 1 structural — it depends only on A_1 at second order, with

A_2 contributions canceling, and is conditional only through λ on the emergent dressing of the Cabibbo identification. The solar and atmospheric angles $\sin^2\theta_{12}$ and $\sin^2\theta_{23}$ are layered predictions: the structural form is forced by the A_1 component (with the sum rule preserved by cubic geometry alone), and the specific deviations at order λ^2 depend on Condition 2's relation $b_{23} = (4/3)(b_{12} + b_{13})$.

Condition 2 as a Layer 2(b) bijection-specific commitment. The framework's $b_{23} = (4/3)(b_{12} + b_{13})$ relation has a multi-layer structural status that the four-layer framing of Chapter 4 §4.5 distinguishes precisely:

(i) Under the U-parity grading developed in §8.4, the antisymmetric tensor product $T_1 \wedge T_1$ decomposes into a U-even subspace (spanned by $E_{12}+E_{13}$) and a U-odd subspace (spanned by $E_{12} - E_{13}$ and E_{23}). The subspace decomposition is forced by group theory — this is Layer 1 structural.

(ii) The cubic-geometric off-democratic projection $A^2 = 1 - 1/d = 2/3$ for $d = 3$ is forced by inner-product structure on T_1 with $\hat{h} = (1,1,1)/\sqrt{3}$ as the democratic axis. This is the same A^2 that produces the Wolfenstein parameter $A = \sqrt{2/3}$ in the quark sector (Chapter 6 §6.5) — Layer 1 structural.

(iii) The cubic-group identity $4/3 = 2A^2 = C_2(T_1)^2/d$ is a representation-theoretic identity of the equivalence class $[(S,\varphi)]/\mathcal{G}_{\text{sub}}$. Whether the substrate produces it as the magnitude of a pair-ratio is not established: the second-order substrate route is excluded (SM §7.3), so the magnitude is Layer 2(b), with the 7/6 sum rule as its direct experimental test and the $O(\lambda^3)$ interference route recorded, not pursued.

(iv) The directional assertion that the U-odd amplitude exceeds the U-even amplitude by the specific factor of 4/3, with the labeling fixed by the observed mass hierarchy ($m_e \ll m_\mu \ll m_\tau$ pinning generation 1 = electron), is *Layer 2(b) bijection-specific* content. It depends on the specific bijection φ within the equivalence class — the same kind of content as the fermion mass hierarchy itself (Chapter 19 §19.3.1) and CKM/PMNS CP phases.

The empirical match at 0.07σ on $\sin^2\theta_{12}$ provides genuine evidence about the specific bijection φ. This is the same form of empirical evidence the framework relies on for other Layer 2(b) content (mass hierarchy match through the Koide structure, the existence of three generations with hierarchical masses).

The open derivational task is correctly scoped: it asks whether the specific staggered-Yukawa structure on the cubic lattice (in the emergent post-trace-out description; the substratum proper has no Lagrangian per Ch02 §2.5) produces A_2 in the sum-rule-preserving subspace for a given bijection. Its second-order lattice-perturbation route to the *magnitude* is excluded (SM §7.3), leaving the $O(\lambda^3)$ interference construction as the recorded remaining route, so the magnitude is Layer 2(b), tested directly by the 7/6 sum rule; the directionality is Layer

2(b) by construction — fixed by the empirical mass-ordering bijection, not derivable from substrate alone in any framework consistent with $\mathcal{G}_{\text{sub}}$-invariance.

8.6 The three numerical predictions

Substituting Condition 1 ($b_{23} = 1$), Condition 2 ($b_{12} + b_{13} = 3/4$), and the Cabibbo scale $\lambda^2 = 1/(2\pi^2)$ into the order-λ^2 deviations from tribimaximal mixing gives the framework's three numerical predictions for the PMNS angles.

The three predictions.

$$\sin^2\theta_{12} = \frac{1}{3} - \frac{1}{4\pi^2} = 0.3080,$$

$$\sin^2\theta_{13} = \frac{4}{9}\cdot\frac{1}{2\pi^2} = \frac{4}{18\pi^2} = 0.02252,$$

$$\sin^2\theta_{23} = \frac{1}{2} + \frac{1}{2\pi^2} = 0.5507.$$

The reactor angle $\sin^2\theta_{13}$ uses only $A^2 = 2/3$ and $\lambda^2 = 1/(2\pi^2)$ together with the constraints on A_1 — it does not use the normalization conditions on A_2. The other two angles additionally use Conditions 1 and 2. The three predictions together exhaust the framework's content for the PMNS mixing angles; the framework leaves δ_{CP} (the CP-violating Dirac phase) undetermined.

Comparison with the post-JUNO global fit and NuFIT 6.0. The empirical state of the PMNS angles as of early 2026 is summarized in the table below. The post-JUNO global fit incorporates the November 2025 JUNO measurement; the NuFIT 6.0 best fits use the full neutrino-oscillation data set under the normal mass ordering convention.

Angle	Predicted	Observed	Match
$\sin^2\theta_{12}$	0.3080	0.3085 ± 0.0073 (post-JUNO global fit)	0.07σ
$\sin^2\theta_{23}$	0.5507	$0.561^{+0.012}_{-0.015}$ (NuFIT 6.0, NO)	0.74σ
$\sin^2\theta_{13}$	0.02252	0.02195 ± 0.00056 (NuFIT 6.0, NO)	1.01σ

All three predictions match observation within 1.1σ at current precision. The solar mixing angle value is the framework's sharpest neutrino-sector empirical match — a parameter-free retrodiction at 0.07σ from the post-JUNO global fit; the derivation (2026) postdates the first results (November 2025), and JUNO's design-lifetime precision (§8.8) is the corresponding pre-registered forward test, well within experimental uncertainty.

The solar mixing angle as the cleanest test. The $\sin^2\theta_{12}$ prediction is the framework's cleanest empirical test in the lepton sector for three reasons. First, it has the smallest experimental uncertainty among the three angles (relative uncertainty about 2.4%, compared to about 2.5% for $\sin^2\theta_{23}$ and about 2.6% for $\sin^2\theta_{13}$). Second, the prediction is parameter-free: both inputs entering the framework's prediction — the zeroth-order value 1/3 from the A_4 tribimaximal pattern and the perturbation scale $\lambda^2 = 1/(2\pi^2)$ from the Cabibbo Brillouin-zone geometry — are fixed by *quark-sector* observables before any neutrino-sector data enter the derivation. The same cubic-group machinery that produces the Cabibbo angle $\lambda = 1/(\pi\sqrt{2})$ at 0.04% precision and the Wolfenstein parameter $A = \sqrt{2/3}$ within 1σ then produces the solar mixing angle $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2) = 0.3080$ at 0.07σ — a cross-sector empirical match with no fresh free parameters fitted to the neutrino sector. Third, the prediction is distinctive in the space of tribimaximal-modification proposals: the post-JUNO data exclude TM2 at more than 3.5σ and place TM1 at 1.35σ, with the framework alone at 0.07σ among the well-developed alternatives.

The cumulative empirical record across the framework's flavor predictions is striking. The Cabibbo angle is matched at 0.04%, the Koide angle at 0.001%, the solar mixing angle at 0.07σ, the sum rule $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$ at 0.6σ — all from the same cubic-group representation theory applied to the lepton sector, with the same numerical inputs ($\lambda^2 = 1/(2\pi^2)$, $A^2 = 2/3$) producing all four predictions. The pattern is what distinguishes the framework's cubic-group structure from postulated flavor symmetries: the same parameter enters multiple observables at sub-percent precision, testable across the full flavor sector rather than only at one observable.

The sum rule. The framework's prediction $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$ is the cubic structural commitment beyond the three angles individually. Substituting the framework's predicted values: $2 \cdot 0.3080 + 0.5507 = 1.167$, which equals 7/6 exactly. Substituting the observed values: $2{\cdot}0.3085{+}0.561 = 1.178{\pm}0.020$, with the agreement at 0.6σ — consistent with the framework's structural prediction. The sum rule is therefore a *fourth* empirical confirmation of the same cubic-group structure, complementary to the three individual angle measurements.

The remaining sections develop the framework's position in the broader landscape of tribimaximal-modification proposals (Section 8.7), the JUNO design-lifetime sensitivity for testing the $\sin^2\theta_{12}$ prediction at sub-percent precision (Section 8.8), and the coupled neutrino-mass / dark-energy result with DESI DR2 (Section 8.9).

8.7 Comparison with TM1/TM2 column-preservation patterns

Tribimaximal mixing was the leading phenomenological description of neutrino oscillation data until Daya Bay measured a non-zero θ_{13} in 2012. Since then, modified tribimaximal patterns have been the subject of a substantial literature, with the two best-known modifications preserving one column of the original tribimaximal matrix exactly: TM1 (first column preserved) and TM2 (second column preserved). Each gives a one-parameter family that accommodates the observed θ_{13} but predicts different values for the other two angles.

TM1 and TM2 in the post-JUNO landscape. Post-JUNO analyses by He and by Zhang find that TM2 is excluded at greater than 3.5σ while TM1 remains marginally viable. The framework's prediction sits well below both alternatives in tension with the post-JUNO global fit:

Approach	Symmetry origin	$\sin^2\theta_{12}$	σ from observed
TM1	postulated A_4 flavor	0.3184	1.35σ
TM2	postulated A_4 flavor	0.3408	4.4σ
TM3	postulated; $\theta_{13} = 0$	—	excluded
This work	**cubic group O of 3D space**	0.3080	0.07σ

(Comparing against the post-JUNO global fit $\sin^2\theta_{12} = 0.3085 \pm 0.0073$. TM1 uses $\sin^2\theta_{12} = 1-(2/3)/\cos^2\theta_{13}$; TM2 uses $\sin^2\theta_{12} = 1/(3\cos^2\theta_{13})$; both with $\sin^2\theta_{13} = 0.02195$.) The cubic-group prediction is closer to the post-JUNO global fit by a factor of about 20 in σ-distance than the best surviving column-preservation pattern, and at the same time makes more falsifiable claims (three independent angles plus the sum rule).

Three structural differences distinguish the framework from TM$_n$.

Symmetry origin. TM1, TM2, and TM3 take a discrete flavor symmetry as an *axiom* of the lepton sector — typically A_4, S_4, or a $\mathbb{Z}_2$ residual postulated for the lepton sector specifically. The framework's prediction uses $A_4 \subset O$, where O is the rotation symmetry group of three-dimensional space. The framework's cubic structure is not postulated for the lepton sector; it is forced by the framework's structural commitments developed in Chapters 5–6: the Bravais-lattice uniqueness theorem (Chapter 5 §5.5), the the link count producing $K = 2dm$ with $m = 1$ under H-link (Chapter 5 §5.4), and the anomaly cancellation chain (Chapter 5 §5.7). The same cubic structure that produced the Standard Model

gauge group with three generations and the Cabibbo angle here produces the PMNS mixing structure. The flavor symmetry is derived, not postulated.

Independence of predictions. TM1 produces a one-parameter correlation $\sin^2\theta_{12} = 1 - (2/3)/\cos^2\theta_{13}$, fixing θ_{12} given θ_{13} but predicting neither separately. The framework's analysis predicts all three angles as fixed numerical values, with the same two structural inputs ($\lambda^2 = 1/(2\pi^2)$ and $A^2 = 2/3$) appearing across all three predictions. This makes the framework's prediction strictly more falsifiable than TM1: any individual angle outside the framework's prediction refutes it independently, whereas TM1 can absorb deviations in one angle by adjusting the prediction for another.

CP violation. He's surviving TM1 pattern predicts $\sin\delta_{\rm CP} = \pm 0.998$, close to maximal CP violation. The framework identifies $\delta_{\rm CP}$ with the only remaining order-λ^2 free parameter $(b_{12} - b_{13})$ that does not enter any of the three angles. The framework's angle predictions therefore stand independently of any value of $\delta_{\rm CP}$; this chapter makes no prediction for the CP phase. The two frameworks are testably distinct in the CP sector: if forthcoming long-baseline neutrino experiments (DUNE, T2HK) measure $\sin\delta_{\rm CP}$ inconsistent with maximal CP violation, TM1 is refuted while the framework's content is unaffected.

The sum rule discriminator. The framework's structural prediction $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$ is the angle-level statement of Condition 2's relation $b_{23} = (4/3)(b_{12} + b_{13})$ from Section 8.5. The observed value is $2(0.3085) + 0.561 = 1.178 \pm 0.020$, in agreement with $7/6 = 1.1667$ at 0.6σ. This depends on the atmospheric octant: the quoted value uses the upper-octant best fit $\sin^2\theta_{23} \approx 0.561$; under the lower-octant solution (≈ 0.455) the combination is ≈ 1.07, roughly 5σ below $7/6$. TM1 and TM2 do not produce this combination as an exact sum rule — in TM2 specifically, $\sin^2\theta_{12}$ is determined by θ_{13} alone with no correlation to θ_{23}, so no analogous sum rule is forced. A direct measurement of $2\sin^2\theta_{12} + \sin^2\theta_{23}$ at the 0.005 level — achievable with the JUNO design lifetime combined with NOvA, T2K, and IceCube-Upgrade atmospheric determinations of θ_{23} — is the sharpest data-side test distinguishing the framework from column-preservation alternatives.

The cumulative-match pattern. The framework's $\sin^2\theta_{12}$ match at 0.07σ does not stand alone. It sits within a cluster of empirical matches that all use the same cubic-group structural inputs: the Cabibbo angle at 0.04%, the Koide angle at 0.001%, the $\sin^2\theta_{13}$ prediction at 1.01σ, the sum rule at 0.6σ, the $|V_{cb}|$ prediction at 0.4σ, the m_u/m_d ratio at 0.05σ. All six predictions match observation at sub-percent or sub-σ precision simultaneously, with two structural parameters ($\lambda^2 = 1/(2\pi^2)$ and $A^2 = 2/3$) appearing repeatedly across the cluster.

Under any reasonable prior over flavor models, this is non-trivial structural agreement. A single set of structural inputs produces six distinct empirical matches; each would have to be explained as a coincidence in a flavor-symmetry framework that postulates the cubic structure rather than deriving it. The

framework's content here is that the same cubic structure that forces the SM gauge group on the substratum side appears as the empirical flavor structure on the phenomenological side — testable across the full flavor sector rather than only at one observable.

8.8 JUNO design-lifetime sensitivity and falsification

JUNO's design lifetime is 30 years of physics operations. The collaboration projects that the experiment will achieve approximately ± 0.0014 precision on $\sin^2\theta_{12}$ by the end of its design lifetime — a factor of six improvement over the first-result precision of ± 0.0087 and approximately a factor of five improvement over the post-JUNO global fit precision of ± 0.0073. The improvement comes from extended exposure (factor of about 185 increase in integrated luminosity over the first 59.1 days), continued calibration refinement, and accumulated cross-checks across multiple reactor cores.

The framework's prediction at JUNO design-lifetime precision. The framework's structural prediction $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2) = 0.30798$ is a single numerical value with no uncertainty from the framework side. The experimental side will reach ± 0.0014 at design lifetime, giving a precision test at the 0.5% relative level on the predicted value. If the framework's prediction is correct, JUNO's design-lifetime measurement should center on 0.3080 with ± 0.0014 uncertainty.

The discrimination this provides is severe. The TM1 pattern's prediction $\sin^2\theta_{12} \approx 0.3184$ differs from the framework's by 0.0104, which is 7.4σ at JUNO's design-lifetime precision. TM2's 0.3408 differs by 0.0328, which is 23σ. Any column-preservation pattern with $\sin^2\theta_{12} \gtrsim 0.31$ will be excluded at JUNO design-lifetime precision; only the cubic-group prediction will remain viable in that range. The 30-year program is therefore a decisive test between the framework's structural prediction and the broader landscape of tribimaximal-modification alternatives.

Sensitivity to Condition 2. The framework's prediction depends on Condition 2 — the structural relation $b_{23} = (4/3)(b_{12} + b_{13})$ developed in Section 8.5. A small fractional shift to the structural coefficient $4/3 \to (4/3)(1+\epsilon)$ propagates to a shift in $\sin^2\theta_{12}$ of approximately $\epsilon \cdot \Delta_{23}/2$ at first order, where $\Delta_{23} = 1/(2\pi^2) \approx 0.05$. For a fractional shift $\epsilon = 0.10$ — a 10% deviation from the cubic-group structural coefficient — the shift in $\sin^2\theta_{12}$ is approximately 0.0025. This is comparable to JUNO's design-lifetime precision and roughly half of the first-result precision.

The framework's current 0.07σ match therefore tests Condition 2 at the few-percent level on its structural coefficient. JUNO's design-lifetime program will tighten this to better than 1%, providing an empirical test of the structural relation $4/3 = 2A^2$ — the framework's commitment that the cubic-group quadratic Casimir structure produces this specific ratio in the substrate. The empirical

confirmation at sub-percent precision over JUNO's 30-year lifetime would constitute a quantitative test of the framework's cubic-group commitment at the level normally reserved for Standard Model coupling constants.

Falsification commitments. Three pre-registered falsification conditions per Chapter 4 §4.5 are concentrated at the JUNO experimental program.

Class A falsification of the cubic-group lepton-sector structure. Any precision-upgrade measurement of $\sin^2\theta_{12}$ moving the observed value outside 3σ of the framework's prediction $1/3 - 1/(4\pi^2)$ refutes the framework's cubic-group commitment in the lepton sector. The framework would have to be revised — either the $A_4 \subset O$ tribimaximal structure at zeroth order, the Cabibbo perturbation scale $\lambda^2 = 1/(2\pi^2)$, or the Condition 2 structural relation — to accommodate the deviation. The current 0.07σ match makes such revision unlikely, but JUNO's 30-year program will resolve the question definitively.

Class A falsification of the sum rule. Any precision measurement of $2\sin^2\theta_{12} + \sin^2\theta_{23}$ moving outside 3σ of the framework's prediction $7/6$ refutes the Condition 2 structural relation. The sum rule is the framework's sharpest discriminator against tribimaximal-modification alternatives, and a precise measurement at the 0.005 level would either confirm or refute the framework's structural commitment in the lepton sector independently of the absolute value of $\sin^2\theta_{12}$.

Class A falsification of the reactor angle. Any precision measurement of $\sin^2\theta_{13}$ moving outside 3σ of the framework's prediction $4/(18\pi^2) = 0.02252$ refutes the Layer 1 reactor angle derivation. This prediction depends only on the structural identity $\sin\theta_{13} = A^2\lambda$ with $A^2 = C_2(T_1)/d = 2/3$ and the Cabibbo scale $\lambda = 1/(\pi\sqrt{2})$ — both Layer 1 inputs. Refutation here would refute the cubic-group representation-theoretic structure at the framework's deepest level. The current 1.01σ match leaves substantial empirical room; ongoing reactor neutrino experiments (Daya Bay, Double Chooz, RENO) and accelerator experiments (T2K, NOvA) will continue to tighten the constraint.

The framework's empirical exposure at JUNO is therefore substantial. Three pre-registered Class A falsification conditions are concentrated at the JUNO experimental program, with a fourth (the $\sin^2\theta_{12}$ first-result match at 0.07σ) already confirmed. The framework's neutrino-sector content has more empirical exposure per prediction than any other sector outside the gravitational confirmations of Chapter 7.

8.9 The coupled DESI DR2 result

The framework's neutrino-sector predictions extend beyond the PMNS mixing angles to include commitments on the neutrino mass spectrum. Chapter 6 §6.7 noted that the framework predicts normal mass ordering and Σm_ν near the oscillation minimum; Chapter 7 §7.7 developed the coupled neutrino-mass / dark-energy result with DESI DR2. This section consolidates the joint observational support.

The DESI DR2 + CMB neutrino mass result. The DESI Data Release 2 combined with CMB lensing and primary CMB constraints from Planck 2018 produces the most precise current bound on the sum of neutrino masses. The analysis finds

$$\Sigma m_\nu < 0.0642\,\text{eV} \quad (95\%\,\text{CL})$$

under the ΛCDM cosmological model. The same analysis prefers the normal mass ordering over the inverted ordering and bounds the lightest neutrino mass at $m_l < 0.023$ eV.

The framework favors a hierarchical normally ordered spectrum. Its numerical near-minimum sum, approximately 0.059 eV, combines that structural expectation with measured solar and atmospheric mass-squared splittings: $\Sigma m_\nu^{\min} = \sqrt{\Delta m^2_{21}} + \sqrt{\Delta m^2_{31}}$ for a negligible lightest mass. The value is therefore layered (L, Layer 3), not a parameter-free structural prediction. The DESI DR2 + CMB upper bound at 0.0642 eV is compatible with it under ΛCDM; an upper limit alone does not establish a standard-deviation distance from a measured mass sum.

The 3σ tension that the framework resolves. The DESI DR2 + CMB analysis under ΛCDM produces a 3σ tension with the *lower* oscillation limit on Σm_ν. The oscillation experiments require $\Sigma m_\nu \geq 0.059$ eV under normal ordering (from $\sqrt{\Delta m^2_{31}} + \sqrt{\Delta m^2_{21}}$ approximately), but the DESI DR2 + CMB combined analysis under ΛCDM prefers a value approaching zero. The literature has interpreted this tension as "a hint of new physics not necessarily related to neutrinos" — a structural mismatch between the cosmological inference assuming ΛCDM and the laboratory oscillation experiments.

The framework's running-vacuum dark energy (Chapter 7 §7.8) resolves this tension. When the cosmological analysis is performed with a Type II RVM dark energy model — the Type II channel the framework carries, its functional form shared with every smooth model (Chapter 7 §7.8) — the inferred Σm_ν shifts upward to be compatible with the oscillation minimum. The Bertini *et al.* fit reports a ν parameter consistent with zero at 2σ but with the residual structure favoring slightly negative values, consistent with the framework's $|\nu_{\text{OI}}| \sim 10^{-32}$ prediction bounded at the emergent-QFT scale. The RVM dark energy and the neutrino mass results are therefore coupled in the framework: both follow from the same lattice structure, and the joint observational support is a single empirical pattern rather than independent confirmation of separate effects.

The neutrino sector as joint test of three commitments. The empirical record across the framework's neutrino-sector predictions tests three structural commitments simultaneously. The PMNS angle predictions (§8.6) test the cubic-group representation theory applied to the lepton sector. The mass ordering and Σm_ν near the oscillation minimum test the framework's normal-ordering commitment and the taste-breaking structure that determines the mass hierarchy. The RVM dark energy resolution of the Σm_ν tension tests the framework's gravitational sector commitment from Chapter 7. The joint observational sup-

port across these three commitments — all consistent with experiment at the current precision — is what the framework's structural unity predicts: three commitments that look distinct in standard physics appear in the framework as different facets of the same cubic-lattice substratum.

The framework's content in the neutrino sector therefore extends well beyond the JUNO $\sin^2\theta_{12}$ measurement. The full empirical pattern includes the PMNS angles, the mass ordering, the absolute mass scale, and the coupled dark-energy / mass-bound resolution. Each tests a distinct framework commitment; the joint coherence across all of them is the framework's strongest non-cosmological empirical case.

8.10 Chapter close: the framework's empirical record in the neutrino sector

The framework's content in the neutrino sector represents the strongest single piece of evidence for the framework's structural commitments outside the gravitational sector. The chapter's specific empirical record:

- The Cabibbo angle $\lambda = 1/(\pi\sqrt{2})$ matches the observed value at 0.04%.
- The solar mixing angle $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2) = 0.3080$ matches the post-JUNO global fit at 0.07σ.
- The reactor mixing angle $\sin^2\theta_{13} = 4/(18\pi^2) = 0.02252$ matches NuFIT 6.0 at 1.01σ.
- The atmospheric mixing angle $\sin^2\theta_{23} = 1/2 + 1/(2\pi^2) = 0.5507$ matches NuFIT 6.0 at 0.74σ.
- The sum rule $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$ matches observation at 0.6σ.
- The neutrino mass ordering (normal) and Σm_ν (near 0.059 eV) are consistent with DESI DR2 + CMB.
- The 3σ tension with the oscillation minimum under ΛCDM is resolved by the framework's RVM dark energy.

The empirical pattern is what distinguishes the framework's cubic-group structure from postulated flavor symmetries. Seven distinct empirical matches across the neutrino sector all use the same cubic-group structural inputs — the squared Cabibbo angle $\lambda^2 = 1/(2\pi^2)$ from Brillouin geometry, the off-democratic projection $A^2 = C_2(T_1)/d = 2/3$ from cubic representation theory, the tribimaximal pattern from $A_4 \subset O$, and the structural relation $b_{23} = (4/3)(b_{12} + b_{13})$ from Condition 2. The framework's content is testable across the full sector rather than only at one observable.

The cross-sector parameter-free status. The framework's $\sin^2\theta_{12}$ prediction uses two structural inputs — $\lambda^2 = 1/(2\pi^2)$ from Cabibbo Brillouin-zone geometry and $A^2 = 2/3$ from cubic-group representation theory — both fixed by quark-sector observables (Cabibbo angle, Wolfenstein parametrization) before any neutrino-sector data enter the derivation. The 0.07σ match at JUNO is therefore not a free-parameter fit to neutrino data; it is the same cubic-group machinery that produced quark-sector matches at 0.04% (Cabibbo) and within

1σ (Wolfenstein A), now tested in a different sector with no fresh parameters fitted. This is what distinguishes a structural framework from a fitted phenomenological model: the inputs are determined by one sector, the prediction is tested in another, and the cross-sector match is empirically nontrivial.

The methodological discipline of Chapter 4 §4.6 places this kind of cross-sector parameter-free match at the center of the framework's empirical case. The framework's neutrino-sector content is the cleanest example of this kind of confirmation outside the cosmological sector developed in Chapter 7. Most tribimaximal-modification patterns in the literature introduce one or more free parameters fitted to the lepton-sector data; the framework's content here is the absence of any such fitted parameter, with the prediction held entirely by structural commitments made elsewhere.

Forward pointers. Chapter 9 develops the framework's universality content — the framework's relationship to other unification programs, the OI-string analysis at the matrix-model level, the universality classes that the framework's Tier 1 structural results belong to, and the framework's content against the broader landscape of grand unified theories and beyond-Standard-Model proposals. The cumulative empirical case developed in Chapters 5 through 8 — twenty-two classified Standard Model retrodictions (two parameter-free structural and five conditional structural on the empirically selected cubic substratum), seven gravitational-sector entries, seven neutrino-sector entries — is what Chapter 9 contextualizes against the broader theoretical landscape. The framework's empirical record at the close of Part II is concentrated at Tiers 1–2 of the gravitational sector, with the Tier 1 calibrations conditional on the horizon conditions of Chapter 7, and at Layer 1 of the Standard Model and neutrino sectors, with substantial empirical exposure across pre-registered falsification conditions and continued empirical agreement at current precision.

The framework's empirical content in fundamental physics is at the strongest evidential standard the field admits: parameter-free structural predictions with inputs fixed by cubic-group representation theory and quark-sector observables, matching neutrino-sector observation at the current experimental precision with no fresh free parameters fitted — retrodictively, with provenance stated (§8.6), and with the pre-registered forward exposure carried by the design-lifetime test (§8.8). The JUNO retrodictive match is the chapter's empirical anchor — the framework's commitment to cubic-group representation theory in the lepton sector, tested at 0.07σ on a parameter-free cross-sector prediction. The remainder of the book develops the framework's content beyond fundamental physics: the cascade through chemistry, life, and intelligence in Part III, and the applications and forward predictions in Part IV.

Chapter 9

Universality Classes and External Convergence

9.1 What this chapter develops

Chapters 5 through 8 developed twenty-two classified Standard Model observable entries (two parameter-free structural and five conditional structural), seven gravitational-sector entries and seven neutrino-sector entries. The numerical comparisons have different evidential roles: selected flavor retrodictions have percent or sub-percent agreement, fitted gauge couplings match by construction, and the neutrino mass sum is constrained by upper bounds. The gravitational formulas carry the conditions of Chapter 7; the MOND acceleration agrees within quoted systematics, and the Bekenstein-Hawking coefficient is a consistency relation, conditional and without a direct test. GW250114 tests the classical area theorem. Appendix A records each entry's classification and empirical standing.

This chapter is different. The framework's content is *not* a competitor to other unification programs that operate at the same level — it is a structural framework at a different level than string theory, loop quantum gravity, or grand unified theories. Locating the framework within the broader theoretical landscape, articulating what classes of results the framework establishes at the universality-class level rather than at the specific-bijection level, and identifying convergences and divergences with mainstream physics literature — these are the chapter's content.

Seven pieces occupy the chapter.

Section 9.2 develops the framework's hierarchical structural realism: the framework operates at multiple levels of abstraction simultaneously, with different chapter contents living at different levels. Some results are *universality-class* results (the energy-shell temperature of the horizon reservoir and the finite counting of boundary modes unconditionally; the Bekenstein-Hawking 1/4 coefficient and the action-scale relation $\hbar = c^3\epsilon^2/(4G\sigma)$ from partition geometry, consistency relations conditional on the horizon conditions named in Chapter 7; the descriptive distinction behind the cosmological constant dissolution) — they hold for any finite deterministic substratum satisfying the framework's structural conditions at its cosmological horizon, regardless of the specific bijection. Other predictions are *substratum-specific* (the Cabibbo angle from cubic Brillouin geometry, the JUNO $\sin^2\theta_{12}$ prediction from $A_4 \subset O$ structure) — they require the specific cubic-lattice realization developed in Chapters 5-6.

Section 9.3 develops the Tier 1 results, and the conditional results built on them, as universality-class content. The framework's gravitational-sector content (Chapter 7) and the memory-bearing quantum phenomenology with its

universal representation (Chapter 1) are largely universality-class — the phenomenology follows from partition geometry and the structural conditions C1–C4, the representation universal outright, the gravitational calibrations from the same geometry under named horizon conditions, not from the specific cubic-lattice substratum. The empirical record at Tier 1 tests the framework's commitments at the level of *structural classes of substrates*, with the cubic-lattice realization being one example within the class. Section 9.4 develops the corresponding Tier 2 and Tier 3 content and clarifies which empirical predictions test which level.

Section 9.5 develops the framework's relationship to string theory at the substratum level. The two programs operate at different levels and answer different questions. The framework's specific commitments to a finite deterministic bijection are not equivalent to string theory's commitments to a continuum quantum gravity with extra dimensions; the apparent overlap in their predictions reflects shared universality-class content rather than substratum-level equivalence. Section 9.6 develops the string-landscape multiplicity problem: the framework's gauge-group derivation does not dissolve the landscape problem because the framework operates at a structurally different level than string compactification.

Section 9.7 develops the Danielson-Satishchandran-Wald (DSW) external convergence — the recent mainstream-physics literature on horizon decoherence as a fundamental phenomenon. The DSW program (Danielson, Satishchandran, and Wald, 2022-2023) proves within standard quantum field theory in curved spacetime that Killing horizons impart fundamental decoherence on nearby quantum superpositions, with rate determined by horizon geometry. This is *direct mainstream convergence* with the framework's core mechanism — horizons as causal partitions producing decoherence. The framework extends DSW's analysis to the capacity-controlled regime (marginal hidden capacity) that DSW does not address, with the BEC analogue-gravity experiment of Chapter 15 providing the experimental access point.

Section 9.8 develops adjacent observer-emergent literature: the Chandrasekaran-Longo-Penington-Witten (CLPW) construction of observer algebras in de Sitter space, the Maldacena and Harlow-Usatyuk-Zhao work on observer-essentiality in quantum gravity, the Slagle-Preskill experimental program for entanglement structure in gravitational systems. These are *adjacent convergences* with the framework's substratum-emergent operator distinction, each addressing a related question from a different theoretical starting point.

Section 9.9 closes the chapter by locating the framework's content within the broader theoretical landscape: the framework's epistemic position relative to other unification programs, the empirical record concentrated at the Tier 1/Tier 2 boundary, and the framework's openness to embedding string-theoretic or loop-quantum-gravity content as universality-class members rather than as competing claims.

The chapter is a *positioning* chapter rather than a *content-development* chapter. The framework's empirical content has been developed in Chapters 5 through 8; here it is located within the space of substratum-level theories and articulated relative to alternative programs. The pedagogical value is to clarify what the framework claims at which level, with the universality-class character of the strongest empirical predictions being one of the framework's distinctive features.

9.2 The framework's hierarchical structural realism

The framework operates simultaneously at multiple levels of abstraction. Chapter 3 §3.4 introduced the hierarchical structural realism that organizes the framework's content; this section develops it concretely against the empirical content of Chapters 5 through 8 and identifies where each chapter's results live in the hierarchy.

The hierarchy. The framework distinguishes four levels of structural commitment, organized along two orthogonal axes.

The first axis is *observation level*. Level O1 is the emergent quantum field theory accessible to embedded observers — the level where the Standard Model, GR, and standard quantum mechanics operate. Level O2 is the substratum, where the framework's bijection φ and partition structure are defined. The framework's content cuts across both levels: O1 is where predictions are tested, O2 is where the structural commitments live.

The second axis is *gauge level*. Level G1 is the framework's general structural axioms (finite substratum, deterministic bijection, classical probability, causal partition). Level G2 is the structural properties of the substratum (the six properties developed in Chapter 5: isotropy, sublattice structure, link content, anomaly cancellation, taste content, generation count). Level G3 is the specific bijection within the equivalence class $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$. Level G4 is the universality class to which any specific bijection belongs — multiple bijections within $\mathcal{G}_{\text{sub}}$ produce identical observables.

The framework's predictions stratify across these levels. Different chapters live at different (O $\times$ G) intersections.

Where each chapter's content lives.

Chapter 1 (Quantum Mechanics from Embedded Observation). The derivation lives at Level O2 $\times$ Level G1: the framework's most general structural commitments produce accessible non-Markovianity in the emergent description, with the quantum representation internal and universal ($S \iff D \iff Q_{\text{fb}}$), the imported correspondence retained for compression under (T), and the operational instrument algebra characterized by five completion conditions within complex matrix kinematics, its physical sourcing open ([Main §3.4]; [GR §3.3]). The result is universality-class-level: any finite deterministic substratum satisfying C1–C4 produces the memory-bearing description whose laws admit the universal quantum representation, regardless of the specific substratum dynamics.

Chapter 5 (Gauge Structure). The exact Level O2 × Level G2 content is the observer-isotropy theorem together with the six-link cubic representation; the latter decomposes as $(3, 2, 1)$. Identifying that representation with one physical Standard-Model carrier is H-link single-copy, with H-cust and the later matter hypotheses entering downstream. The geometric decomposition is therefore structural; the physical gauge-group reading is a conditional bridge construction rather than a consequence of a $K = 2d$ minimization theorem.

Chapter 6 (Matter Content). The quantitative predictions of Standard Model parameters live at Level O1 × Level G3 — the specific cubic-lattice bijection produces specific values for the Cabibbo angle, Koide relation, gauge couplings, and other observables. The Cabibbo angle prediction $\lambda = 1/(\pi\sqrt{2})$ tests the framework's commitment to the cubic Brillouin geometry at Level G3.

Chapter 7 (Gravitational Sector). Predictions stratify across multiple levels. The energy-shell horizon temperature, the finite boundary-mode count and the two-level distinction behind the cosmological constant dissolution are Tier 1 results at Level O2 × Level G4; the Bekenstein-Hawking 1/4 coefficient and the action-scale relation $\hbar = c^3\epsilon^2/(4G\sigma)$, consistency relations, sit at the same level conditional on the horizon conditions of Chapter 7 §7.3, the dynamical form of the dissolution is conditional on the coupling theorem of [GR §6.2], and the Type II running-vacuum form is not derived (Chapter 7 §7.8). The MOND acceleration scale $a_0 = cH/6$ and the baryonic Tully-Fisher relation are Tier 2 results at Level G2, inheriting the conditions on the de Sitter temperature with the dimensional factor imported from Verlinde. The MOND interpolation steepness is Tier 3 at Level G3.

Chapter 8 (JUNO and Neutrino Sector). The PMNS angle predictions test Level G3 content — the specific cubic-lattice bijection's $A_4 \subset O$ symmetry produces $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2)$. The sum rule $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$ tests Level G3 structural commitments. The framework's coupled neutrino-mass and RVM dark-energy result tests Level G2 (mass ordering, Σm_ν near oscillation minimum) and Level G4 (RVM functional form).

Multi-level structural realism. A single empirical prediction can carry content at multiple levels simultaneously. The GW250114 area-theorem result illustrates the level structure (noting that it tests classical horizon dynamics, not the 1/4 coefficient, which has no direct empirical test): - Level G4: the universality-class commitment to thermal self-consistency at the partition boundary. - Level G2: the framework's specific implementation of the thermal self-consistency through the gap equation chain. - Level G3: the cubic-lattice realization being one example within the universality class.

The same observation provides evidence at multiple levels. The empirical record is therefore not just a record of "does the framework's specific cubic-lattice bijection match observation?" but "does the framework's structural commitment at each level match observation?" The cumulative case across the framework's empirical chapters is strongest at Level G2 and G4 — the framework's structural

commitments at the substratum-properties level and the universality-class level are extensively tested, with the cubic-lattice realization at Level G3 providing one specific way the structural commitments are met.

Why the hierarchy matters. The hierarchical structure clarifies what the framework's empirical case actually tests. Critics sometimes claim that frameworks like this one are too flexible — that any empirical result could be accommodated by adjusting the specific substratum bijection. The hierarchical structure shows this concern misplaced: most of the framework's strongest empirical predictions test Level G4 (universality-class) and Level G2 (structural-properties) content, which is *not* adjustable by varying the bijection.

The Bekenstein-Hawking 1/4 coefficient is determined at the universality-class level: any finite deterministic substratum satisfying C1–C4 at its cosmological horizon, with thermal self-consistency between classical horizon temperature and emergent KMS temperature, gives the 1/4 factor. There is no bijection within the universality class that produces 1/3 or 1/5. A direct empirical test of the coefficient, if one becomes available, would test this universality-class commitment rather than any specific bijection's predictions; GW250114, which compares areas and does not measure entropy, does not provide such a test.

Similarly, the MOND acceleration scale $a_0 = cH/6$ is determined at the structural-properties level: the dimensional factor 1/6 arises from the cubic-lattice geometry combined with the $d = 4$ spacetime structure, common to any substratum with these properties. The descriptive distinction behind the proposed cosmological constant dissolution is determined at the structural-properties level, regardless of the specific bijection; the dissolution itself is conditional on the coupling step that [GR §6.2] leaves open.

The framework's empirical case is therefore largely a case at the *structural* levels — universality-class and substratum-properties — with the substratum-specific (Level G3) content concentrated in the flavor sector and the dark-sector interpolation function. This is what distinguishes the framework from frameworks operating at a single specific level: the framework's predictions are stratified, with most of the empirical exposure at structural levels where the predictions are robust to substratum-specific choices.

Predictions stratified by hierarchy level. The empirical record of Chapters 5-8, organized by hierarchy level, summarizes as follows.

Hierarchy level	Predictions	Empirical match
Level G4 (universality class)	RVM reading (form shared by every smooth model)	non-discriminating; the consistency relations $S = A/(4l_p^2)$ and $\hbar = c^3\epsilon^2/(4G\sigma)$ and the descriptive CC distinction are stated in Chapter 7 §7.10
Level G2 (substratum properties)	SU(3) × SU(2) × U(1), three candidate sectors (generations under H-spin'), no GUT, $a_0 = cH/6$, no DM particle	Concordance
Level G3 (specific bijection)	Cabibbo angle, Koide relation, PMNS angles, sum rule, gauge couplings at M_Z	0.04%-1.01σ for the measured comparisons (gauge couplings matched by construction)
Level G3 (open)	Bottom-to-tau mass ratio, Koide angle's structural origin, δ_{CP}	Open research program

The framework's empirical exposure is therefore distributed across the hierarchy, with the universality-class level consistent with observation rather than sharply tested and the open research program concentrated at the bijection-specific level. This distribution is what makes the hierarchical structural realism meaningful: different levels are tested with different empirical evidence, and the framework's content at each level is constrained by the corresponding empirical record.

9.3 Tier 1 results as universality-class content

The framework's strongest structural results are concentrated at the universality-class level (Level G4 in §9.2's hierarchy, Tier 1 in Chapter 7's tier classification, together with the conditional results built on it). This section develops what universality-class content means structurally and traces the universality-class status of those results, including the named conditions the calibrations inherit.

What universality-class content means. A universality-class result is a structural commitment that follows from a small set of structural conditions and holds across multiple substratum realizations. The classic example in condensed matter is the renormalization-group universality of critical phenomena: different microscopic Hamiltonians (Ising model, ϕ^4 field theory, certain lattice gauge

theories) produce identical critical exponents at their phase transitions because they belong to the same universality class.

The framework's structural conditions C1, C2, C3, C4 — coupling to a hidden sector, record persistence (slow-bath separation one route), sufficient hidden-sector capacity, history readback — define a universality class of substratum theories. Any finite deterministic substratum with a partition satisfying C1–C4 belongs to the class. The cubic-lattice substratum developed in Chapters 5-6 is one member; alternative substrata that satisfy C1–C4 differently would be other members of the same class.

The framework's Tier 1 results, and the conditional results built on them, are properties of the universality class as a whole, not of the specific cubic-lattice realization; the conditions they carry are statements about the horizon and its dynamics rather than about the substratum, so they do not reintroduce bijection-specific content. Three examples illustrate.

The Bekenstein-Hawking entropy. $S = A/(4l_p^2)$ is recovered in Chapter 7 §7.5 from boundary-mode counting combined with thermal self-consistency at the cosmological horizon, conditional on H-slope together with the horizon and frame conditions of §7.3. The counting $S = \sigma A/\epsilon^2$, for an entropy σ per boundary mode, is unconditional; the coefficient enters through $\epsilon^2 = 4\sigma\, l_p^2$, from which σ cancels, which the gap equation chain — classical horizon temperature equals emergent KMS temperature — fixes under those conditions, and the chain is a property of any substratum with the framework's structural conditions at its horizon. The specific cubic-lattice substratum is *one way* C1–C4 are satisfied at the horizon, but alternative substrata satisfying C1–C4 differently would give the same 1/4 coefficient under the same conditions. A direct test of the coefficient would probe the universality-class commitment, not the cubic-lattice specifics; GW250114, which compares areas and does not measure entropy, does not provide one.

The cosmological constant dissolution. The descriptive distinction behind the proposed dissolution in Chapter 7 §7.7 follows from the two-level description: the classical substratum carries the vacuum energy $\rho_{\text{crit}} \sim H^2/G$, while the emergent QFT has vacuum energy $\rho_{\text{QFT}} \sim M_{\text{Pl}}^4$. The two-level description is a property of any substratum with the framework's structural conditions; the specific substratum dynamics do not enter. The distinction holds for any member of the framework's universality class. That gravity couples to the classical substratum and that the emergent zero-point energy does not gravitate — the dynamical form of the dissolution, in which the effective cosmological constant tracks H^2/G — is conditional on the covariant coupling theorem that [GR §6.2] leaves open.

The action-scale relation from partition geometry. $\hbar = c^3\epsilon^2/(4G\sigma)$ follows in Chapter 7 §7.3 from thermal self-consistency at the cosmological horizon, conditional on H-slope with the horizon and frame conditions stated there, with the discreteness scale ϵ and the entropy σ per boundary mode the only substratum-

specific inputs. The functional form is universal across the framework's universality class; the specific value of ϵ would vary across alternative substrata, with self-consistency requiring $\epsilon = 2\sqrt{\sigma}\, l_p$ for any member of the class.

The universality-class character of Tier 1 results. The framework's Tier 1 results are therefore *not* claims about the specific cubic-lattice substratum but claims about the broader universality class of substrata satisfying the framework's structural conditions. This has three consequences.

First, the empirical confirmations at Tier 1 are evidence for the framework's universality-class commitments, not for the specific cubic-lattice realization. A competing substratum theory that produced the same Tier 1 results would be empirically indistinguishable from the framework at the universality-class level. The framework's distinctive empirical exposure is at Tier 2 and Tier 3, where substratum-specific content can in principle differ across alternative realizations.

Second, the framework is *robust* to substratum-specific revisions that preserve the structural conditions C1–C4. If empirical evidence eventually forced a revision of the cubic-lattice realization (perhaps a different lattice geometry, a different group-theoretic decomposition, a different taste-breaking structure), the framework's Tier 1 results would persist as long as the structural conditions held. The framework's commitment to specific cubic-lattice content is therefore *less load-bearing* than its commitment to the universality-class structural conditions.

Third, the framework's content is *embeddable* within other frameworks at higher levels of abstraction. A meta-framework that produced multiple universality-class members — perhaps string theory's landscape of compactifications, or a more general bijection-on-finite-set construction — would have the framework as one structural commitment among many. The framework's epistemic stance is therefore not "this is the unique theory of fundamental physics" but "this is one structural framework that meets a set of universality-class commitments, with specific bijection content tested empirically."

The framework's universality-class commitments.

The framework's universality-class commitments — the conditions that define membership in the framework's universality class — are six structural properties of the substratum, organized as follows.

Finiteness. The substratum S has finite cardinality. This is load-bearing for the recurrence route specifically: the recurrence theorem (Chapter 18 §18.7) requires finiteness for the exact recurrence of a finite bijection, which supplies recurrence-scale P-indivisibility. It is not required for accessible non-Markovianity, which has its own finite-horizon route through readback, nor for the imported representation, which is gated by (T).

Deterministic bijection. The substratum dynamics $\varphi : S \to S$ is a bijection. Bijectivity is load-bearing because it preserves information at the substratum

level (the premise of information conservation in Chapter 7 §7.6) and produces the recurrence structure (Chapter 18 §18.7).

Causal partition. The configuration space partitions as $S = \mathcal{C}_V \times \mathcal{C}_H$ for an embedded observer, with the partition determined by causal structure at the cosmological horizon. This is load-bearing because the partition is what makes the visible law stochastic: history readback across it marks the memory-bearing sector, and every finite law it yields admits the quantum representation.

C1 (coupling). The visible and hidden sectors are non-trivially coupled. This is necessary for non-Markovian dynamics; failing C1 produces memoryless (Markovian) dynamics rather than the memory-bearing sector — fixed-basis quantum representability itself is universal, memoryless laws included ([Main §3.4]).

C2 (memory persistence). The hidden sector retains a record of the visible past across the accessible window, $\tau_B \gg \tau_S$ with τ_B the coarse-grained mixing time. Slow hidden dynamics is one sufficient mechanism; fast dynamics that conserves and routes the record serves equally ([Main §1.3]).

C3 (sufficient capacity). The hidden sector must carry information capacity at least the history's demand, $\log_2 |\mathcal{C}_H| \geq I^*$ — in our universe realized comfortably with $N_H \gg N_V$. This capacity is necessary for the hidden sector to have room to store the correlations the read-write cycle generates.

These six commitments define the framework's universality class. Any finite deterministic bijection on a partitioned configuration space satisfying C1–C4 at its cosmological horizon belongs to the class. The framework's universality-class results — the energy-shell horizon temperature, the finite boundary-mode count and the two-level distinction unconditionally; the Bekenstein-Hawking 1/4 and the action-scale relation for $\hbar$ as consistency relations under the horizon conditions of Chapter 7; the dark sector budget as concordance — follow from these six commitments without requiring the specific cubic-lattice realization. The MOND acceleration scale $a_0 = cH/6$ is Tier 2 (Chapter 7 §7.9), and the running-vacuum form is not derived (Chapter 7 §7.8).

The framework's empirical exposure at Tier 1. The cumulative empirical record at Tier 1 summarizes as follows.

Tier 1 prediction	Empirical match	Source chapter
Area law $dA \geq 0$	GW250114 area increase at 99.999%	Ch 7 §7.5
Dark sector budget $\sim 95\%$	$\Omega_{\text{dark}} \approx 0.95$ concordance	Ch 7 §7.9
No dark matter particle	XENON, LUX, etc. null	Ch 7 §7.9
No GUT-mediated proton decay	$\tau_p > 10^{34}$ yr	Ch 6 §6.4

Four universality-class entries tested against current empirical evidence: one confirmed at high precision (the area law) and the remainder consistent with the empirical record at concordance level (dark sector budget, no DM particle, no proton decay). Information conservation follows from the bijectivity premise and is not an empirical entry: a measurement of the radiation alone does not test it, and the Page curve is conditional (Chapter 7 §7.6). The Bekenstein-Hawking coefficient and the descriptive vacuum-energy distinction are consistency relations (Chapter 7 §7.10) and are not tabulated. The Type II running-vacuum form is not in the table: its local expansion is shared by every smooth model, so it is non-discriminating (Chapter 7 §7.8), and the falsifiable content is the magnitude $|\nu| \sim 10^{-32}$ at Tier 2. The framework's universality-class content is therefore consistent with observation rather than sharply tested — the framework belongs to a universality class whose members must produce these results under the stated conditions, and the empirical record is consistent with the framework being in this class.

The next section develops the corresponding Tier 2 and Tier 3 content and traces the empirical record at the substratum-properties and bijection-specific levels.

9.4 Tier 2 and Tier 3 stratification

The framework's empirical content beyond Tier 1 stratifies into two further tiers, with the empirical exposure shifting from universality-class commitments to substratum-properties content (Tier 2) and to specific-bijection content (Tier 3). The chapter's hierarchical structural realism makes the stratification explicit; this section traces the framework's empirical record at the corresponding levels.

Tier 2: substratum-properties content. Tier 2 results follow from the framework's structural commitments to the substratum's properties — the six structural properties developed in Chapter 5 (isotropy, sublattice structure, link content, anomaly cancellation, taste content, generation count) — combined with the universality-class conditions C1–C4. Tier 2 results are *not* universal across all members of the framework's universality class; they require the substratum to have the specific properties Chapter 5 develops, but they do *not* require the specific bijection.

The framework's Tier 2 content includes the Standard Model gauge group emerging from cubic decomposition, the three-generation structure from T_1 multiplicity, the absence of grand unification (these three on the single-copy H-link branch, the gauge group and the absence of grand unification also with an octahedrally equivariant condensate; Chapter 5 §5.4–§5.5, Chapter 6 §6.4), the MOND acceleration scale $a_0 = cH/6$, the baryonic Tully-Fisher relation, and the high-redshift evolution of the dark-matter crossover radius $r_M(z) \propto 1/\sqrt{H(z)}$. Each of these holds for any substratum with the cubic-lattice properties and the structural conditions C1–C4, regardless of how the specific bijection

breaks the cubic symmetry.

The empirical record at Tier 2 summarizes as follows.

Tier 2 prediction	Empirical match	Source chapter
$\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ gauge group	confirmed	Ch 5
Three generations of matter	confirmed	Ch 5, Ch 6
No grand unified theory	$\tau_p > 10^{34}$ yr consistent	Ch 6 §6.4
$a_0 = cH/6$	0.91 of the observed scale at the Planck H_0, inside the systematic; factor 1/6 imported	Ch 7 §7.9
Baryonic Tully-Fisher $v^4 = GM_B \cdot cH/6$	confirmed empirical relation	Ch 7 §7.9
$r_M(z)$ shrinkage with redshift	in tension (Genzel et al., high-z)	Ch 7 §7.9
Mass ordering (normal)	DESI DR2 + CMB preference	Ch 8 §8.9

The seven Tier 2 entries combine observed structural features, conditional gravitational formulas and consistency with bounds. The acceleration formula carries the imported factor and the quoted systematic uncertainty; the redshift and mass-ordering comparisons retain their model dependence. Their evidential roles are assessed individually, rather than as seven independent confirmations.

Tier 3: bijection-specific content. Tier 3 results require the *specific bijection* within the equivalence class $[(S, \varphi)]/\mathcal{G}_{\mathrm{sub}}$ — not just the substratum's structural properties but the particular way the bijection breaks the cubic symmetry. Tier 3 results are *substratum-specific* in the strongest sense: alternative bijections within the framework's universality class with different taste-breaking patterns would produce different Tier 3 predictions.

The framework's Tier 3 content includes the Cabibbo angle $\lambda = 1/(\pi\sqrt{2})$, the Koide relation $Q = 2/3$, the PMNS angle predictions with $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2)$, the sum rule $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$, the gauge couplings at M_Z, and the MOND interpolation function's specific transition steepness. Each of these requires the specific cubic-lattice bijection's properties beyond the general cubic-symmetry commitments.

The empirical record at Tier 3 summarizes as follows.

Tier 3 prediction	Empirical match	Source chapter
Cabibbo angle $\lambda = 1/(\pi\sqrt{2})$	0.04%	Ch 6 §6.5
Koide relation $Q = 2/3$	$< 0.001\%$	Ch 6 §6.5
$\sin^2\theta_{12} = 1/3 - 1/(4\pi^2)$	0.07σ from post-JUNO global fit	Ch 8 §8.6
$\sin^2\theta_{13} = 4/(18\pi^2)$	1.01σ from NuFIT 6.0	Ch 8 §8.6
$\sin^2\theta_{23} = 1/2 + 1/(2\pi^2)$	0.74σ from NuFIT 6.0	Ch 8 §8.6
Sum rule $2\sin^2\theta_{12} + \sin^2\theta_{23} = 7/6$	0.6σ	Ch 8 §8.6
Three gauge couplings at M_Z	Fitted; zero residual by construction	Ch 6 §6.4
MOND interpolation "simple" function	Conditional comparison; transition kernel open	Ch 7 §7.9

The eight Tier 3 entries comprise six numerical flavor comparisons, one fitted coupling entry and one conditional interpolation comparison. The flavor comparisons test the stated cubic-lattice relations at the precisions listed. The fitted couplings supply no independent confirmation, and the interpolation comparison retains the open transition-kernel obligation.

Why the tier stratification matters. The stratification separates the objects tested at each level. Tier 1 has four entries: the area theorem is measured, and the remaining entries have the conditions and empirical standing listed above. Tier 2 has seven entries concerning the substratum's properties, including conditional gravitational formulas and observational consistency checks. Tier 3 has eight entries concerning the specific-bijection account, with the numerical, fitted and conditional comparisons distinguished above.

The three tiers contain nineteen entries of unequal evidential weight. Numerical retrodictions, fitted quantities, conditional formulas and untested results remain distinct; a count of entries is not a count of independent confirmations. The tier assignment identifies the level at which each statement is made, while its conditions and empirical standing determine what its comparison tests.

9.5 The framework's relationship to string theory

The framework and string theory both aim to provide a structural account of fundamental physics that goes beyond the Standard Model and general relativity. The two programs operate at different levels and answer different questions, with apparent overlaps that warrant careful analysis. This section develops the framework's relationship to string theory at the substratum level, drawing on the per-generator analysis developed in `Structure.md` Part II.

The two programs at different levels. String theory is a continuum quantum-gravity program: it postulates a Lorentzian ten- or eleven-dimensional spacetime with quantum fields, with the Standard Model emerging from compactification of the extra dimensions on a specific Calabi-Yau manifold. The framework is a finite deterministic-substratum program: it postulates a finite configuration space with a deterministic bijection, with quantum mechanics and general relativity both emerging as projections of the same structure.

The two programs are *not* operating at the same level. String theory is itself an emergent description in the framework's hierarchy — the relevant operations (Riemann tensor, sigma-model worldsheet, target-space supersymmetry) presuppose continuum spacetime, which is the *emergent* description in the framework rather than the substratum. The framework's substratum is logically prior to anything string theory describes.

The apparent overlap between the two programs — both predict the Standard Model gauge group from group-theoretic considerations, both have content about black hole entropy and quantum gravity — reflects *shared universality-class content* rather than substratum-level equivalence. String theory and the framework both occupy positions within broader classes of theoretical frameworks, with some structural results common to both classes.

The matrix-model bridge. A potential bridge between the two programs runs through matrix models. The framework's bijection φ acts on a finite configuration space, with the dynamics expressible in terms of matrix operations on finite-dimensional state spaces. String theory has matrix-model formulations (BFSS, BMN, IKKT, banks-fischler-shenker-susskind) where the dynamics is similarly expressible in matrix terms, with the continuum string theory emerging in specific limits.

The bridge would be: identify the framework's finite-bijection structure with a specific matrix model whose continuum limit produces a string-theoretic spacetime. Six structural axioms must be checked for compatibility under such a bridge.

A1 (finiteness). The framework's representative has finite $|S|$. Matrix models with $N \to \infty$ limits violate this strictly, but for any finite N the bridge is consistent. Since the extension of finiteness beyond the observer's distinguishable states is a choice of representative rather than a further fact (Chapter 2), the bridge requires a finite-N representative — the matrix model not taken to its continuum limit — without asserting that N is physically finite. This distinguishes the framework's bridge from standard matrix-model interpretations.

A2 (determinism / bijection). The framework requires deterministic bijection dynamics. Matrix models with deterministic Hamiltonian evolution satisfy this; matrix models with quantum-mechanical evolution on the matrix degrees of freedom do not, unless the quantum mechanics is interpreted as emergent from the substratum (consistent with the framework's content but not with standard matrix-model interpretations).

A3 (bounded coupling degree). The framework retains bounded local coupling as a substratum condition, but it does not derive the physical carrier dimension $K = 6$ from coupling-degree minimization. Matrix models with arbitrary connectivity therefore still need a bounded-locality bridge, while the separate H-link single-copy condition controls whether the six geometric links are the physical carrier.

A4 (center independence). The update commutes with lattice translations up to gauge, and the update at a site contains no explicit copy of that site's present value. Matrix models have natural translation symmetries, which supply the first clause; the second is a further condition on the matrix-model update.

A5 (linearity). The framework's emergent dynamics is linear at the visible-sector level. Matrix models with linear evolution on matrix degrees of freedom are consistent; non-linear matrix dynamics would not be.

A6 (background independence). The framework's assumption is covariance under spatially varying internal-index transformations, with the link coupling transformed alongside; separately, the framework has no fixed spacetime background — the substratum is the background, with spacetime emergent — and that is a different sense of the same words. Matrix models with explicit background spacetime are not background-independent in the emergent-spacetime sense; the bridge would require the matrix model to be of background-independent form (BFSS in its M-theory interpretation, for instance, where spacetime emerges from the matrix dynamics), and whether its matrix-internal gauge covariance then plays the role the assumption requires depends on that matrix-spatial bridge.

The summary is that *some* matrix models are bridge-compatible with the framework, but not all. The framework's structural axioms place real restrictions on which matrix models can serve as the bridge, and the standard interpretations of matrix models in string theory are not always consistent with the framework's commitments.

What can be salvaged. Two recoveries are possible.

Recovery (i): String theory as universality-class member. The framework's universality class might include matrix-model representatives that produce specific string-theoretic features in continuum limits. Under this reading, string theory is one universality-class member among many — its content at the universality-class level (BH entropy, AdS/CFT, certain modular structures) overlaps with the framework's content, while its substratum-specific content (specific compactifications, specific Calabi-Yau structures) is distinct from the framework's specific bijection.

Recovery (ii): Emergent string-theoretic features. The framework's substratum produces the reduced description carrying the emergent quantum representation; under specific bijection structures, the emergent description might exhibit string-theoretic features (sigma-model dynamics on emergent worldsheets, modular

invariance, T-duality-like equivalences). This would not be string theory at the substratum level but string-theoretic features at the emergent level, derived rather than postulated. Developing this recovery would require constructing the framework's emergent description for specific bijections and tracing whether string-theoretic structures appear.

Neither recovery is currently developed in detail; both are open theoretical tasks that the framework's structural commitments make tractable in principle.

9.6 The string landscape multiplicity problem

The string landscape — the multiplicity of possible compactifications of string theory, each producing different low-energy physics — is the central interpretive problem of string theory as a unification program. Estimates of the landscape size vary from 10^{500} to much larger. The framework's gauge-group derivation in Chapter 5 does *not* dissolve the landscape problem, because the framework operates at a structurally different level than string compactification.

Why the landscape problem persists. The framework derives SU(3) × SU(2) × U(1) with three candidate matter sectors (generations under H-spin', SM §4.7) from cubic-lattice substratum properties (Tier 2) and the specific Standard Model parameters from the cubic-lattice bijection (Tier 3). These are framework-level derivations within the framework's universality class. They do not select among string compactifications because string compactifications occupy a different theoretical level — they specify the geometry of the extra dimensions in continuum string theory, which is the emergent description in the framework's hierarchy.

A given string compactification may or may not be consistent with the framework's structural commitments. Those compactifications consistent with the framework's universality-class membership produce the framework's Tier 1 results; those compatible with the framework's substratum properties produce the framework's Tier 2 results; only those compatible with the framework's specific bijection structure produce the framework's Tier 3 results. The framework's derivations therefore *select within* the landscape rather than dissolve it: the framework's specific bijection corresponds to a subset of string compactifications (if any) that produce the framework's specific predictions.

The landscape multiplicity problem persists in the framework's content as the multiplicity of bijections within the framework's universality class. Both programs face a multiplicity question; the questions are at structurally different levels.

Cross-attribution of predictions does not follow. A consequence of the level distinction: the framework's empirical confirmations do not automatically transfer to string theory, and string theory's empirical confirmations do not automatically transfer to the framework. The Bekenstein-Hawking 1/4 coefficient — obtained at the universality-class level under the horizon conditions of Chap-

ter 7, with no direct empirical test (GW250114 constrains the classical area theorem) — is a universality-class result for the framework and is reproduced in specific string theory calculations (Strominger-Vafa for extremal black holes). The two derivations are *independent*, with neither producing the other; both are evidence for the universality-class commitment that both programs share, not for either program's substratum-specific content.

Similarly, the framework's twenty-two classified Standard Model retrodictions (two parameter-free structural and five conditional structural) are framework-specific Tier 3 results that do not follow from string theory's structural commitments alone — they would require selecting the specific string compactification corresponding to the framework's bijection, which has not been done in the string theory literature. The framework's empirical record is independent of any specific string-theoretic compactification's status.

Connection to swampland conjectures. The string theory swampland program — Vafa, Ooguri, Brennan, Carta, and others — develops conjectures about which low-energy effective field theories can be UV-completed within string theory. The framework's structural commitments are similar in spirit: they identify a class of substratum theories that produce specific emergent physics, with the framework's universality class playing a role analogous to the swampland's "landscape" complement.

Some swampland conjectures align with framework predictions. The de Sitter conjecture (no stable de Sitter vacua in string theory) is in tension with the framework's commitment to a de Sitter cosmological horizon at all times; if the de Sitter conjecture is correct, the framework's substratum would not be embeddable in standard string theory. Alternative swampland conjectures (the weak gravity conjecture, the species bound, the distance conjecture) have varying degrees of compatibility with the framework's structural commitments; a systematic analysis would require working through each conjecture against the framework's specific bijection content.

The framework's epistemic stance toward the swampland is therefore: the swampland program is asking related but structurally distinct questions, and the framework's content provides answers at a different level. Whether the two programs converge or diverge is a matter for detailed analysis on a case-by-case basis.

9.6b Foundational programs claiming parameter-free SM retrodictions

A small group of foundational programs, distinct from string theory's compactification approach, claim parameter-free derivation of Standard Model observables from prior structural commitments. The framework belongs to this reference class. Locating the framework against other members of the class clarifies what is distinctive about the framework's content.

Beck's chaotic strings. Christian Beck (*Physica D* 171:72-106, 2002; arXiv:hep-th/0105152) developed a program in which 1+1-dimensional coupled-map-lattice chaotic dynamics produces the Standard Model coupling constants and quark/lepton masses from minima of effective potentials generated by chaotic-string vacuum energy. Beck's program claims to reproduce the electroweak and strong coupling constants to 4-5 digits precision and the quark/lepton masses to 3-4 digits, in a peer-reviewed venue and with sustained development across multiple publications (Beck et al., *Spatio-temporal Chaos and Vacuum Fluctuations of Quantized Fields*, 2002; Beck, arXiv:hep-th/0702082, 2007).

The framework's relationship to Beck's program is non-competitive at the *type-of-claim* level: both programs assert parameter-free derivation of SM observables from substrate-level dynamics. The frameworks differ in their substrate commitments — Beck's chaotic-string dynamics is more abstract than the framework's cubic-lattice Bravais commitment, and Beck does not extend to gauge structure, cosmological constant, dark sector, or biological applications. The framework's distinguishing features within this reference class are: explicit cubic-group representation theory as the derivational machinery (rather than chaotic-attractor minimization); cross-domain reach beyond particle physics (cosmological constant dissolution, chromatin pharmacology, BQP characterization, quantum biology); and a more articulated structural foundation through the C1–C4 conditions and the reconstruction theorem developed in Chapters 1-3.

Beck's program has not been adopted as a canonical SM-parameter explanation in mainstream particle physics over its 24-year history. The reasons for non-adoption are not formally documented in published critique, but the citation pattern suggests skeptical reception around the question of whether the chaotic-string parameter-tuning is genuinely parameter-free or whether the program contains hidden fitted choices in the chaotic-attractor selection. This audit question — *does the substrate machinery genuinely force the observed values, or does it admit hidden flexibility?* — applies to the framework as well, and is the framework's most consequential remaining specialist-review question per Chapter 4 §4.5 and the chapter close below.

Other foundational programs. Adler's trace dynamics (Adler, *Quantum Theory as an Emergent Phenomenon*, 2004) develops a pre-quantum matrix dynamics from which quantum mechanics emerges as an effective description. The program does not currently produce SM-parameter retrodictions at the framework's precision, but operates at a similar structural level (substrate-emergent QM rather than substrate-emergent SM parameters). Padmanabhan's emergent-gravity program (multiple publications 1998-2020) derives Einstein's equations from thermodynamic considerations on causal horizons; this is structurally adjacent to the framework's GR-as-classical-limit content (Chapter 7) but does not extend to SM parameters. Wolfram's hypergraph rewriting (*A New Kind of Science*, 2002; the Wolfram Physics Project 2020-) claims cross-domain reach comparable to the framework's, but has produced no parameter-free SM-observable

retrodictions at sub-percent precision.

The framework's distinctive position within the reference class. Among foundational programs claiming to derive SM observables from prior structure, the framework appears to be the only candidate that has produced parameter-free retrodictions at sub-percent precision across multiple SM observables (Cabibbo at 0.04%, Koide at 0.0009%, $\sin^2\theta_{12}$ at 0.07σ, the GW250114 area-theorem consistency at 99.999% — the 1/4 coefficient itself untested directly), with the cross-domain extension to cosmology, chromatin pharmacology, and computational complexity. The distinguishing-from-Beck assessment is: similar type of claim, more articulated foundation, broader empirical record, and the same audit question (substrate-level genuine-derivation versus hidden flexibility) determining whether the framework follows Beck into mainstream-skeptical reception or instead achieves the recognition that its cumulative empirical record would warrant.

9.6c A gravitational-sector sibling: conformal gravity

The comparison class of §9.6b — programs claiming structural Standard Model content — has a gravitational-sector counterpart: programs covering the framework's Chapter 7 anomaly set without dark matter and without a fine-tuned cosmological constant. The most developed is conformal gravity, Philip Mannheim's five-decade program built on the Weyl-squared action. Its defining principle is local conformal symmetry: no fundamental mass or length scale is permitted, all masses arise dynamically, and the exact Mannheim-Kazanas vacuum solution adds a linear potential term to Newton's law that has been fit, with universal coefficients, to over a hundred galaxy rotation curves without dark matter. Its quantum sector rests on PT-symmetric (non-Hermitian) quantization, developed with Bender, which the program presents as dissolving the ghost problem of fourth-order gravity — with the striking corollary that the graviton need not exist as a particle.

The two frameworks agree on an unusual number of conclusions — no particle dark matter, no fine-tuned cosmological constant, no fundamental graviton — while being incompatible at every foundation that produces them. Conformal symmetry forbids exactly the fundamental scale ($\epsilon = 2\sqrt{\sigma}\, l_p$) that this framework's construction begins from. Conformal gravity's gravitational field is fundamental, fourth-order, and quantized; here gravity is emergent, second-order, and never quantized — the Clausius derivation of Chapter 7 targets the Einstein equations directly, the covariant step being open ([GR §8.7]), with curvature-squared corrections Planck-suppressed rather than dominant, so neither theory can be the other's effective limit. Conformal gravity modifies the quantization rules to non-Hermitian form; here Hermitian quantum mechanics is a derived theorem with no freedom to modify. And all-dynamical mass with a composite Higgs stands against substrate-eigenvalue masses with the elementary taste-singlet Higgs of Chapter 6. The two programs occupy disjoint universality classes whose observable signatures happen to intersect on the present anoma-

lies.

The cosmological constant comparison is instructive because the two programs make the same opening move and then diverge into opposite kinds of solution. Both refuse to quench Λ. Conformal gravity lets Λ be as large as particle physics suggests and quenches gravity's *response* instead: the effective cosmological coupling differs from the locally measured Cavendish value, so the observable Ω_Λ is naturally of order one regardless of Λ's magnitude — a dynamical mechanism inside a single description, with a non-fine-tuned fit to the supernova Hubble diagram as its evidence. This framework instead proposes to dissolve the commensurability: the quantum vacuum energy of the emergent field theory and the effective cosmological constant of the thermodynamic description are properties of logically distinct levels (Chapter 7 §7.6), so that, conditional on the coupling step that [GR §6.2] leaves open, the 10^{122} ratio is not a cancellation problem — a category resolution whose falsifiable residue is the dark-energy account — the Type II running-vacuum reading, its functional form shared with every smooth model — with magnitude indistinguishable from a pure constant. Each solution carries its own exposure. Conformal cosmology commits to an expansion that has always accelerated — no deceleration epoch — which high-redshift standard-candle analyses now test, with several reporting tension; its negative effective coupling is contested; and there is a live debate in the literature over whether the PT quantization its ghost resolution requires preserves the coupling structure its cosmology uses. This framework's exposure is the mirror image: a near-null prediction is hard to shoot at, and the shot exists — if the current hints of strongly evolving dark energy harden, the indistinguishable-from-Λ running-vacuum form is strained, which is precisely what the pre-registered DESI Y5 test decides. One further contrast prices the two dissolutions rather than adjudicating them: conformal gravity retains the vacuum energy as a large physical quantity and purchases dynamical machinery — fourth-order equations, non-Hermitian quantization, a negative effective coupling — to quench gravity's response to it, whereas here the absolute offset is a gauge parameter of the reconstruction theorem ([Main §3.4]; Chapter 7 §7.6), invisible to every quantum statistic, and on the geometry-first proposal nothing needs quenching; whether the offset sources curvature is the coupling step that [GR §6.2] leaves open. Each framework's argument is internal to its own premises — this is a difference in what the two theories must buy, not evidence against either — but the machinery cost sits asymmetrically, and the side effects of the purchased machinery are what the high-redshift data tests.

The dark-matter sectors are mutually falsifiable in the same way. In the Mannheim–O'Brien galactic fits conformal gravity's linear potential is accompanied by a global quadratic term from the cosmological background, which turns outer rotation curves down and ends bound orbits at radii of order 100 kpc; the entropy-displacement mechanism of Chapter 7 §7.7 produces the flat asymptote of an acceleration scale, and ties that scale to the horizon, $a_0 = cH/6$ — predicting redshift evolution of the acceleration scale that a geometric potential does not mimic. Falling against flat outer curves at 50–250

kpc separates the two. The pre-registered baryonic Tully-Fisher test at $z > 1$ is therefore simultaneously a forward test of this framework and a discriminator against its most developed no-dark-matter competitor. The two programs also stand in an instructive symmetry with respect to what the classical tests actually test: the precision confirmations of general relativity probe the vacuum Ricci-flat solution set, which underdetermines the field equations — a freedom conformal gravity uses to shelter *different* equations behind the shared solutions in the tested regimes, and which this framework uses in the opposite direction, targeting the *standard* equations and locating its novelty in their origin and their sourced-sector structure. The sectors where equations rather than solutions are probed — radiation and cosmology — are accordingly where the two programs' fates diverge: this framework inherits the radiative confirmations exactly if its covariant dynamics are Einstein's, the open G3 step, while the radiative sector of fourth-order gravity is an acknowledged open area of that program, and the cosmological sector is where each has placed its bets. Two programs with contradictory foundations pointing at the same anomalies is the configuration experiments exist to resolve, and the discriminating data — high-redshift standard candles, outer rotation curves, DESI Y5, BTFR at depth — is scheduled rather than hypothetical.

The last axis of comparison is explanatory unification, and it is the one where the two programs' architectures differ most instructively — because both are built on a single unifying symmetry, and the symmetries enter at opposite ends of the logic. Conformal gravity's unifier is a postulated symmetry *of nature's action*: local conformal invariance is assumed, and it then works hard — it uniquely selects the Weyl action, forbids fundamental scales and so forces dynamical mass generation, and forbids a fundamental cosmological constant. This framework's unifier, the substratum gauge group $\mathcal{G}_{\text{sub}}$, is not an assumption about nature's action but a *derived* structure: the invariance group of embedded observation itself, the relabelings of the substrate that no observation can distinguish. Its explanatory reach follows from that origin. Local gauge symmetry — the thing conformal gravity and every field theory must postulate — appears here as an output, the commutant structure of the substrate's internal components (Chapter 5); the phase conventions of quantum mechanics are among its gauge directions; the uniform vacuum-energy offset is another (Chapter 7 §7.6, via the reconstruction theorem's energy-shift freedom), which is why the cosmological constant dissolution above is a *symmetry statement* in both programs — conformal symmetry forbids a fundamental Λ and must then manage the induced one, while here the offset is gauge outright and there is nothing to manage; and the boundary between what the framework can derive and what remains bijection-specific is drawn by $\mathcal{G}_{\text{sub}}$-invariance itself, with the invariant list (the Cabibbo magnitude, the Koide ratio, the 1/4 coefficient, the acceleration scale $a_0 = cH/6$) constituting the substrate-computable record. One principle thus answers, within the framework's premises, questions that span quantum structure, the origin of gauge symmetry, the vacuum energy, and the program's own epistemology — a breadth of claimed unification that conformal gravity's nar-

rower, older principle does not attempt. The calibration that must accompany the comparison: breadth of *claimed* unification is not depth of *tested* unification. Conformal gravity's narrower claims have decades of scrutiny and known wounds; this framework's broader claims are unreviewed, and their breadth is a promissory asset that external review and the framework's own gate computations will either convert or void.

9.7 The Danielson-Satishchandran-Wald external convergence

The framework's core mechanism — horizons as causal partitions producing decoherence in the visible sector — has received independent convergence from mainstream physics literature on horizon decoherence. The Danielson-Satishchandran-Wald (DSW) program (Danielson, Satishchandran, and Wald 2022 in *Physical Review D*; 2023 in *International Journal of Modern Physics D*) proves within standard quantum field theory in curved spacetime that Killing horizons impart fundamental decoherence on nearby quantum superpositions, with rate determined by horizon geometry. This is *direct mainstream convergence* with the framework's content — the horizon partition as the source of visible-sector decoherence is being independently derived by physicists operating in standard QFT-in-curved-spacetime without any commitment to the framework's substratum picture.

The DSW result. DSW consider a quantum superposition near a Killing horizon (a black hole horizon or a cosmological horizon) and compute the decoherence rate of the superposition due to soft graviton emission across the horizon. The result is striking: the decoherence rate is *fundamental* — it does not depend on any environmental degrees of freedom inside the horizon, and it does not vanish in the limit of zero temperature. The horizon itself acts as a decoherence channel, with the rate determined by the horizon's geometry (specifically, the horizon area and the horizon's surface gravity).

The DSW derivation runs through three steps. First, they compute the entanglement between a near-horizon quantum superposition and the soft graviton modes produced by gravitational radiation across the horizon. Second, they show that the soft modes propagate through the horizon and cannot be recovered by any operation outside the horizon. Third, they identify the resulting irreversible loss of coherence as fundamental decoherence imposed by the horizon's causal structure.

The result has been confirmed and extended by independent groups — Gralla and Wei (2024), Wilson-Gerow, Dugad, and Chen (2024), Chen and Penington (2024), and others — establishing the DSW decoherence as a robust prediction of standard QFT in curved spacetime, independent of any specific framework for quantum gravity.

Convergence with the framework's mechanism. The framework's content from Chapter 1 — embedded observers see their reduced description as quantum-

mechanical because the cosmological horizon imposes a partition with C1–C4 structure — has three specific features that match DSW's analysis.

Decoherence depends on horizon geometry. At the cosmological partition the framework's hidden-sector parameters are set by the horizon: its area fixes the capacity (C3) and its surface gravity the correlation time (C2). The framework does not derive a decoherence rate from these parameters. DSW compute a rate with horizon area and surface gravity as the geometric inputs, so the convergence is in those inputs: horizons impose decoherence through their geometry, and the framework's hidden-sector parameters are the same geometric quantities.

Decoherence vanishes when coupling is screened. The framework's C1 condition requires non-trivial coupling between visible and hidden sectors. If C1 fails (the coupling is screened or vanishingly small), the framework's emergent dynamics is memoryless rather than memory-bearing — the finite law still universally admits Q_{fb}. DSW reproduces this with extremal black holes: when the horizon's surface gravity vanishes (the extremal limit), the soft graviton coupling to the horizon also vanishes, and the DSW decoherence rate goes to zero. The match is structural: both frameworks predict that extremal horizons impose no fundamental decoherence.

Decoherence is non-Markovian near the horizon. The framework's emergent dynamics is non-Markovian, with memory effects from the hidden sector's slow timescale τ_B. DSW reproduces this with their analysis of the soft mode propagation across the horizon: the soft modes carry temporal correlations that produce memory effects in the near-horizon observer's reduced description. The match is non-trivial: both frameworks predict that horizon-induced decoherence is structurally distinct from the Markovian decoherence of standard open quantum systems.

The convergence is across three independent structural features. The framework's marginalization mechanism — whose memory-bearing output admits the quantum representation — is being independently derived from standard QFT in curved spacetime, with the structural features (geometry-dependence, coupling-screening, non-Markovianity) matching across the two derivations.

Where the framework extends DSW. The framework's content extends beyond DSW's analysis in one structurally important direction: the capacity-controlled regime of marginal hidden capacity. DSW's analysis assumes the horizon has effectively infinite capacity (the soft graviton modes can absorb arbitrary information without saturating). The framework's C3 condition — sufficient hidden memory capacity, comfortably $N_H \gg N_V$ in the cosmological realization — can be satisfied with a finite hidden capacity in engineered partitions (Chapter 15) and biological partitions (Chapter 13); the mechanism proposal predicts that where that capacity is marginal, as engineered partitions can make it, the decoherence dynamics deviates from the universal DSW result.

The mechanism proposal predicts that capacity-controlled systems exhibit modified decoherence: when the hidden capacity is marginal ($N_H/N_V \sim 1$), it pre-

dicts partial distinguishability revival, $0 < \mathcal{N} < 1$ — the restricted quantum-statistical regime of Chapter 18 §18.3. DSW's analysis does not extend to this regime; the framework's content is therefore *additional* to DSW's, with experimental access through the BEC analogue-gravity experiment developed in Chapter 15.

The BEC analogue-gravity setup of Chapter 15 is precisely a test of the framework's capacity-controlled regime: an analogue horizon engineered to have controllable hidden-sector capacity, with the framework's proposed model form $\mathcal{F}(r) = 2r - r^2$ as the revival signature as the capacity parameter r is varied (derivation unpackaged; prediction-status appendix). This is the framework's empirical access point for DSW-extension content — testing the framework's prediction in a regime DSW does not address.

Significance for the framework's epistemic position. The DSW convergence is significant because derivations within standard QFT in curved spacetime, made without commitment to the framework's substratum picture, reach the same structural features. The framework's commitment to horizons as causal partitions producing decoherence is not an idiosyncratic claim; it is shared by an independent theoretical derivation.

The framework's distinctive content — the substratum picture, the specific cubic-lattice realization, the Tier 3 predictions — is not addressed by DSW. The framework's universality-class commitments — that horizons produce decoherence, with geometry setting the relevant parameters, with coupling-screening eliminating the effect, with non-Markovian structure — converge with that derivation. The convergence is theoretical: it shows that the universality-class content is shared with mainstream physics, with the framework one specific realization within that universality class, and it adds no empirical evidence for the framework.

9.8 Adjacent observer-emergent literature

Beyond the DSW convergence on horizon decoherence, the framework's content engages a broader recent literature on observer-emergent physics. Four programs in particular have substantive overlap with the framework's substratum-emergent operator distinction, each addressing related questions from different theoretical starting points.

The Chandrasekaran-Longo-Penington-Witten (CLPW) construction. Chandrasekaran, Longo, Penington, and Witten (2022) constructed an observer-relative algebra of operators in de Sitter space using crossed-product techniques from operator algebra theory. The construction identifies the observer's operator algebra as a *Type II* von Neumann factor (rather than Type III, the algebra of operators in standard quantum field theory), with finite von Neumann entropy and well-defined density matrices. The construction is observer-relative: different observers in de Sitter space have different algebras, with the algebras related by appropriate isomorphisms.

The CLPW construction converges with the framework's substratum-emergent operator distinction. The framework's visible-sector observables form a finite-dimensional Hilbert space at any given time (the emergent description has finite information content per cell of the partition), with the observer's accessible operator algebra being precisely the algebra of observables on this space. The framework's reduced description is observer-relative in the same sense as CLPW's construction: different observers correspond to different partitions, with different reduced descriptions.

The frameworks differ in their substratum commitments. CLPW operates within standard QFT in curved spacetime, with the operator algebra being the conventional structure of QFT modified by the crossed-product construction. The framework operates with a finite deterministic substratum, with the operator algebra being the emergent description of the visible sector after the trace-out. The two frameworks therefore converge on observer-relativity and finite von Neumann entropy from different starting points.

Maldacena on observer-essentiality. Maldacena (2021, 2024) has developed the position that observers are essential to a complete formulation of quantum gravity — that the wave function of the universe cannot be defined without reference to a specific observer, and that different observers correspond to different formulations of the quantum-gravity problem. This is the *observer-essentiality* thesis, formulated within string theory and AdS/CFT but with broader implications.

The observer-essentiality thesis converges with the framework's content. The framework derives that quantum mechanics is the emergent description of an embedded observer — there is no observer-independent quantum mechanics. Different observers correspond to different partitions of the substratum, with different reduced descriptions, and the universal wave function of the universe is not the right level of description for any embedded observer.

The frameworks differ in the source of the observer-relativity. Maldacena's framework places it at the level of quantum gravity's holographic principle; the framework's content places it at the level of the substratum's partition structure. The two frameworks converge on the conclusion (observers are essential) from different starting points.

Harlow-Usatyuk-Zhao on observer dependence. Harlow, Usatyuk, and Zhao (2024) developed observer-dependence in quantum gravity through analysis of the entanglement wedge reconstruction and the gauge structure of the Hilbert space. Their result: the operator algebra accessible to an observer depends not just on the spatial region the observer occupies but on the observer's full causal structure, with different observers having structurally different algebras even when their spatial regions overlap.

The HUZ result converges with the framework's content on the partition-relativity of the reduced description. The framework's visible sector is determined by the observer's causal structure (the cosmological horizon's

location relative to the observer), with different observers having structurally different partitions. The HUZ result on operator-algebra differences between observers matches the framework's content on reduced-description differences between observers.

Slagle-Preskill emergent quantum mechanics. Slagle and Preskill (2023) constructed a local classical lattice model with one extra spatial dimension: classical bits propagate toward a boundary through a circuit of stochastic matrices, and the boundary dynamics reproduces Schrödinger evolution for a chosen local Hamiltonian, approximately but to arbitrary precision. Bell nonlocality is carried by bulk signals that travel faster than the boundary speed of light, and the authors estimate analytically, and check by simulation, how far the model departs from quantum mechanics.

The model converges with the framework on the possibility that quantum mechanics emerges from classical dynamics. It differs in mechanism: the Slagle-Preskill construction uses an extra spatial dimension and stochastic bulk dynamics, while the framework uses a finite deterministic substratum and a trace-out with no extra dimension.

The cumulative external convergence. The DSW, CLPW, Maldacena, HUZ, and Slagle-Preskill programs together constitute substantial external convergence with the framework's content. Each program is operating with different starting commitments (DSW with standard QFT, CLPW with operator algebra theory, Maldacena with string theory and holography, HUZ with AdS/CFT entanglement wedge reconstruction, Slagle-Preskill with a classical lattice model), and each converges with specific features of the framework's substratum-emergent operator distinction.

The framework's content is therefore *not* idiosyncratic but lies in the area where mainstream physics is converging. The specific framework-level content (substratum picture, cubic-lattice realization, Tier 3 predictions) is distinctive; the universality-class content (observers are essential, partitions are causally determined, horizons impose decoherence, reduced descriptions are observer-relative) is converged-upon by multiple independent programs in mainstream physics.

9.9 Chapter close: the framework's position in the broader landscape

The framework's content across Chapters 5 through 8 establishes a substantial empirical record at high precision: twenty-two classified Standard Model retrodictions (two parameter-free structural and five conditional structural on the empirically selected cubic substratum), seven gravitational-sector entries, seven neutrino-sector entries. This chapter has located that empirical content within the framework's hierarchical structural realism (§9.2), traced the universality-class character of the strongest empirical predictions (§9.3-9.4), articulated the framework's relationship to string theory and the string landscape

multiplicity problem (§9.5-9.6), and developed the substantial external convergence with mainstream physics literature on horizon decoherence (§9.7) and observer-emergent physics (§9.8).

The framework's epistemic position. The framework's content is at three levels simultaneously: a *universality-class commitment* (the framework belongs to a class of substratum theories with C1–C4 structure at the cosmological horizon), a *substratum-properties commitment* (the framework's substratum is a cubic lattice at $\epsilon = 2\sqrt{\sigma}\, l_p$ with the six structural properties of Chapter 5), and a *specific-bijection commitment* (the framework's bijection produces the Tier 3 predictions of Chapter 6 and Chapter 8). Each level is tested by different empirical evidence, with the cumulative record being strongest at the universality-class and substratum-properties levels.

This epistemic position distinguishes the framework from frameworks operating at a single level. Frameworks aiming at universality-class content alone (effective field theories, top-down structural arguments) lack the empirical specificity of the framework's Tier 3 predictions. Frameworks aiming at substratum-specific content alone (specific string compactifications, specific GUT models) lack the universality-class robustness of the framework's Tier 1 results. The framework's stratified empirical record provides constraint at each level, with the levels reinforcing rather than substituting for each other.

External convergence and its scope. The framework's universality-class commitments — observers as essential, horizons as causal partitions producing decoherence, reduced descriptions as observer-relative, finite entanglement entropy, quantum mechanics as emergent from classical dynamics — are converged-upon by multiple independent mainstream programs: DSW on horizon decoherence, CLPW on observer-relative operator algebras, Maldacena on observer-essentiality, HUZ on observer-dependent operator structure, Slagle-Preskill on quantum mechanics emerging from a classical lattice. The convergence is between independent theoretical derivations from different starting commitments; it is not empirical confirmation of the framework.

The framework's distinctive content — the substratum picture, the cubic-lattice realization, the specific Tier 3 predictions — is not addressed by these convergences. Its empirical standing rests on the framework's own comparisons with data, classified in appendix A: the strongest are retrodictions (the JUNO $\sin^2\theta_{12}$ match at 0.07σ, the Cabibbo angle at 0.04%), the Bekenstein-Hawking 1/4 is a consistency relation of the horizon calibration with no direct test, the Milky Way comparison is qualitative, and none is yet a pre-registered forward confirmation. The framework's case therefore combines theoretical convergence on universality-class commitments with a retrodictive empirical record on framework-specific predictions.

Where the framework stands relative to alternatives. The framework's stance toward alternative unification programs is non-competitive at the universality-class level and distinctive at the substratum-specific level. String

theory, loop quantum gravity, causal dynamical triangulations, asymptotic safety, and other programs may produce identical universality-class results in their respective limits, with the framework being one structural realization that produces the same results from a different starting point. The frameworks become distinguishable only at the substratum-specific level — where the framework's cubic-lattice realization produces specific Tier 3 predictions that competing programs do not match without additional substratum-specific assumptions.

This stance is intellectually irenic. The framework is not a competitor to existing unification programs but a structural framework that overlaps with them at the universality-class level while providing distinctive substratum-specific content. Where the framework's content matches mainstream physics convergence (DSW, CLPW, Maldacena), the framework's content is reinforced rather than refuted by that convergence. Where the framework's content goes beyond mainstream convergence (the BEC analogue-gravity test, the Tier 3 predictions), the framework provides distinctive empirical targets.

Forward pointers. The remaining chapters develop the framework's reach beyond fundamental physics. Chapter 10 develops the framework's content in chemistry: the cascade from substratum to chemistry through the framework's emergent description, with specific quantitative content for chemical bonding and reactivity. Chapter 11 develops the origin of life: how the framework's structural commitments at the chemical level produce a cascade leading to the emergence of self-replicating systems. Chapter 12 develops evolution, intelligence, and self-reference. Chapters 13 through 17 develop the framework's applications in quantum biology, quantum computing, quantum engineering, medicine, and bioinformatics. The cumulative reach of the framework across these chapters — fundamental physics in Part II, the emergence cascade in Part III, applications in Part IV — provides the framework's complete empirical and theoretical record.

The framework's content from Chapter 5 through Chapter 8 — concentrated at the universality-class, substratum-properties, and specific-bijection levels in this chapter's hierarchical structural realism — provides the empirical foundation. The remaining chapters build on that foundation, with the framework's reach extending from the substratum-level structural commitments developed in Part I and the empirical record developed in Part II into the broader scientific domains where the framework operates.

Chapter 10

Chemistry

10.1 What this chapter develops

The framework's empirical content in Chapters 5 through 9 concentrates in fundamental physics. The chapters develop the framework's derivation of the Standard Model gauge group, the matter content, the gravitational sector, the JUNO neutrino prediction, the dark sector phenomenology, and the framework's position in the broader theoretical landscape. The empirical record is concentrated in the sectors of fundamental physics where the framework makes specific quantitative predictions.

This chapter develops the framework's content in *chemistry* — the emergence cascade from the substratum to molecular physics, with the framework deriving chemistry as a structural consequence of the substratum's commitments rather than as an independent layer of physics. The chapter's content is concentrated at Tier 1 and Tier 2 of the framework's hierarchical structural realism: chemistry emerges from properties of the substratum that are shared across the framework's universality class, with the specific chemical species and reactivity patterns following from the framework's specific cubic-lattice realization.

Seven pieces occupy the chapter.

Section 10.2 develops the structural chain from substratum to orbital chemistry: how the framework's commitments to three spatial dimensions, spin-1/2 fermions, and the Pauli exclusion principle produce the periodic table, the s/p/d/f orbital structure, and the specific bonding geometries of carbon (tetrahedral sp^3, planar sp^2). The chain inherits the conditional dimension selection of Chapter 5 §5.3 and the stated field-theory and atomic assumptions; its orbital and bonding consequences are evaluated within that branch.

Section 10.3 develops four additional structural results: matter-antimatter asymmetry (three generations + CKM CP violation), the mass hierarchy (Higgs + taste-breaking producing light fermions), nuclear binding (SU(3) confinement producing bound hadrons and nuclei), and chiral chemistry (electroweak parity violation propagating to molecular parity-violating energy differences). Each result is a structural consequence of the framework's commitments at the substratum level, propagated through the emergence cascade to chemistry.

Section 10.4 develops the scale hierarchy that makes chemistry possible: the framework's two independent gauge groups (SU(3) and U(1)) with independent coupling strengths whose values, with the electron mass, produce $E_{\text{nuclear}} \sim$ MeV $\gg E_{\text{atomic}} \sim$ eV. This separation means nuclei are stable against chemical reactions — chemistry rearranges electrons without disturbing nuclei. Section 10.5 develops the thermal window: the parametric conditions under which chemistry is both stable (bonds survive thermal fluctuations) and dynamic (reactions can proceed). A window requires bond energies well above kT for the molecules and temperature in question, activation energies low enough for reactions at that temperature, and a solvent liquid there; the nuclear–atomic scale separation does not by itself establish any of these, and the window's location depends

on parameter values.

Section 10.6 develops water as the unique biological solvent: how oxygen's element-8 orbital structure produces water's hydrogen-bonding network and the anomalous properties (high heat capacity, density maximum, universal solvent behavior, hydrophobic effect) that are load-bearing for biological function. Section 10.7 develops the chirality chain: how electroweak parity violation at the substratum level propagates through molecular physics to produce the parity-violating energy difference between mirror-image molecules. The framework predicts a structural preference for L-amino acids in biology — the homochirality of life is a consequence of the framework's chirality from trace-out (Chapter 5), not an arbitrary historical contingency. Section 10.8 develops the framework's content on fine-tuning and the anthropic principle: the dissolution of both as conventional formulations, with the framework's reading being that the *possibility* of a chemistry-supporting universe is structural while specific parameter values determine particular realizations within the structural class.

The framework's content in chemistry is *not* a competitor to standard chemistry — the predictions of quantum chemistry, organic chemistry, and biochemistry are all reproduced by the framework's emergent description. What the framework adds is a structural derivation of why chemistry has the architecture it does: why three-dimensional space, why sp^3 and sp^2 hybridization, why the periodic table, why two independent gauge couplings (whose values separate the nuclear and atomic scales), which conditions a thermal window for liquid-phase chemistry requires, why molecular homochirality. The chapter's value is in providing structural foundations for chemistry that standard accounts treat as empirical input.

10.2 The structural chain from substratum to orbital chemistry

The proposed chain from substratum to chemistry runs through five steps, conditional on A4, the filters E5–E7, the continuum conditions and the spin, statistics and atomic assumptions stated below. The finite pair (S, φ) alone does not fix the dimensional premise of this chain.

Step 1: Three spatial dimensions. The framework's argument for $d = 3$ is the three-filter convergence of Chapter 5: propagating gravity excludes $d \leq 2$, stable matter — and with it the embedded observers of Axiom 1 — excludes $d \geq 4$, and the boundary-entropy concordance checks $d = 3$ exactly. (The H-link link count $K = 2dm$ with $m = 1$ and the nearest-neighbor Bravais-lattice uniqueness theorem then operate *within* $d = 3$, fixing the component count and the lattice type; they do not themselves fix the dimension.) The papers carry the complementary, more conservative anchoring: the cubic-lattice commitment is also empirically selected by the observed gauge group.

The dimensional commitment is *load-bearing for chemistry.* In $d = 2$, the angular momentum algebra reduces — only circular harmonics exist, with no three-

dimensional bonding geometry. In $d \geq 4$, the Coulomb potential $V(r) \sim r^{2-d}$ does not support stable bound states: classical orbits spiral into the nucleus, and the quantum Hamiltonian is unbounded below. The chemical construction inherits the conditional selection of $d = 3$ within A4's periodic-lattice class under E5–E7 and the continuum conditions of Chapter 5 §5.3.

Step 2: Orbital algebra in $d = 3$. Standard quantum mechanics in $d = 3$ produces the SO(3) angular momentum algebra with the familiar quantum numbers (n, ℓ, m_ℓ). The orbital angular momentum quantum number ℓ takes values $0, 1, 2, \ldots, n-1$ for principal quantum number n. The magnetic quantum number m_ℓ takes values $-\ell, -\ell+1, \ldots, +\ell$, giving $2\ell + 1$ orbital states per ℓ value.

The orbital labels translate to the conventional spectroscopic notation: s ($\ell = 0$, 1 orbital), p ($\ell = 1$, 3 orbitals), d ($\ell = 2$, 5 orbitals), f ($\ell = 3$, 7 orbitals). On the selected three-dimensional continuum branch, rotational symmetry gives the SO(3) orbital algebra. This conclusion inherits the dimensional and continuum assumptions of Chapter 5 §5.3.

Step 3: Spin-1/2 and the Pauli principle. The framework derives spin-1/2 fermions from the staggered structure on the $d = 3$ cubic lattice (Chapter 5 §5.6). The staggered fermion construction places independent fermion modes at alternating lattice sites; the chiral structure of the framework's emergent QFT (Chapter 5 §5.6) requires the modes to carry half-integer spin. The spin-statistics theorem — a consequence of the emergent QFT's Lorentz invariance, established with the emergent QFT structure of Chapter 5 §5.4 — gives the Pauli exclusion principle: at most one fermion per state, hence at most 2 electrons per orbital (spin up plus spin down).

The shell capacities follow: s (2 electrons), p (6 electrons), d (10 electrons), f (14 electrons). The periodic table's period lengths — 2, 8, 8, 18, 18, 32 — are determined by the order in which these shells fill, with the filling order following from energy ordering of the orbitals.

The orbital counting is conditional on the selected $d = 3$ branch and the stated spin and statistics assumptions. The dimensional step inherits A4, the empirical filters E5–E7 and the continuum conditions; it is not a consequence of the finite pair (S, φ) alone.

Step 4: Carbon's bonding geometry. Element 6 (carbon) has the electron configuration $1s^2\, 2s^2\, 2p^2$ — four valence electrons distributed among $2s$ and $2p$ orbitals. In $d = 3$ with the derived orbital algebra, four valence electrons admit three hybridization modes that mix the s and p orbitals.

sp^3 hybridization. The s orbital mixes with all three p orbitals to produce four equivalent orbitals pointing to the vertices of a regular tetrahedron, with bond angles of 109.5°. This is the geometry of methane (CH_4), diamond, and the backbone of all organic macromolecules — proteins, nucleic acids, lipids, carbohydrates.

sp^2 hybridization. The s orbital mixes with two of the three p orbitals, producing three planar orbitals at 120°, with one p orbital remaining perpendicular to the plane. The perpendicular p orbital forms a π bond. This is the geometry of graphite, benzene, and aromatic chemistry — the planar-ring chemistry that underlies nucleic acids' bases and aromatic amino acids.

sp hybridization. The s orbital mixes with one p orbital, producing two linear orbitals at 180° plus two perpendicular p orbitals forming π bonds. This is the geometry of acetylene (HC≡CH) and other triple-bonded systems.

The *possibility* of these hybridizations — and hence the possibility of three-dimensional macromolecular architecture — is a structural consequence of the angular momentum algebra in $d = 3$ combined with carbon's four valence electrons. Given the selected $d = 3$ branch and carbon's stated electronic structure, this geometry follows from the orbital construction.

Why carbon specifically. The framework's structural derivation singles out carbon among the periodic-table elements. Carbon's position is determined by three structural features: four valence electrons (in the row where s and p shells are half-filled), $2p$ orbital size matching $2s$ orbital size (allowing the hybridizations), and second-period status (avoiding the d-orbital availability of heavier elements that complicates bonding). These features combine to give carbon the unique ability to form four equivalent strong covalent bonds with itself and with hydrogen, nitrogen, oxygen, sulfur, and other light elements.

The framework's content here addresses a question that standard chemistry treats as empirical: why is carbon the basis of organic chemistry and biology rather than silicon (Si, four valence electrons in the third period) or germanium (Ge, four valence electrons in the fourth period)? The structural answer involves orbital size matching. In silicon, the $3s$ and $3p$ orbitals have different radial extents (the $3s$ orbital is more compact than the $3p$ due to penetration effects), making sp^3 hybridization less symmetric and silicon-silicon bonds substantially weaker than carbon-carbon bonds (Si-Si bond energy $\sim$ 226 kJ/mol versus C-C bond energy $\sim$ 347 kJ/mol). The framework's commitment to the standard atomic-orbital structure makes carbon's combinatorial chemistry possible at room temperature in ways silicon's chemistry cannot match.

The framework's prediction is therefore: any chemistry-supporting universe with the framework's structural commitments will use carbon (or its precise analog at the same orbital-structure position) as its primary combinatorial element. The "carbon chauvinism" of biology is structurally predicted rather than empirically contingent.

The hybridizations as load-bearing for biology. The three carbon hybridizations correspond to three structural classes of biological molecules.

sp^3 carbons form the saturated backbones of carbohydrates, lipids, and the side chains of most amino acids. The tetrahedral sp^3 geometry enables three-dimensional folding of proteins (the backbone ϕ and ψ dihedral angles depend

on tetrahedral sp^3 geometry at each α-carbon).

sp^2 *carbons* form the planar aromatic rings of nucleic acid bases (adenine, guanine, cytosine, thymine, uracil), the aromatic amino acids (phenylalanine, tyrosine, tryptophan), and the peptide bond's planar carbonyl carbon. The peptide bond's sp^2 character is what produces the rigid planar structure that constrains protein secondary structure into alpha-helices and beta-sheets.

sp carbons are rarer in biology but appear in some natural products and as transition states in enzyme catalysis. Their linear geometry produces the rigid axes needed for some specialized binding interactions.

On the selected three-dimensional continuum branch, and subject to the stated spin, statistics and atomic assumptions, these hybridization modes supply the molecular geometries used in the biological discussion of Chapters 11 and 12.

Step 5: The structural chain. The first arrow below inherits A4, E5–E7 and the continuum conditions of Chapter 5 §5.3. The remaining steps use the stated spin, statistics and atomic assumptions:

$$(S, \varphi) \xrightarrow{\text{conditional}} d = 3 \xrightarrow{\text{SO(3)}} \text{orbital algebra} \xrightarrow{\text{Pauli}} \text{periodic table} \xrightarrow{\text{element 6}} \text{tetrahedral carbon.}$$

The dimensional selection consumes the physical inputs E5–E7 within A4's assumed lattice class. The downstream orbital and bonding relations are conditional on that selected continuum branch and the stated spin, statistics and atomic assumptions.

The empirical confirmation of this chain is *concordance*: the framework's derivation reproduces the periodic table and the geometry of carbon's bonding, with no deviations from standard chemistry at currently accessible energy scales. The framework does not add new chemistry; it provides structural foundations for the chemistry that has been empirically established over the past century. The value of the framework's content here is the structural account of why chemistry has the architecture it does.

10.3 Beyond carbon: four additional structural results

The framework's chemistry derivation extends beyond carbon's bonding geometry to four additional structural results that propagate the substratum's commitments through to chemistry-relevant features.

Matter-antimatter asymmetry. The framework's commitment to three matter generations (Chapter 5 §5.7) produces a CKM mixing matrix with one physical CP-violating phase. CP violation is one of three Sakharov conditions for baryogenesis: the matter-antimatter asymmetry of the observed universe requires (i) baryon number violation, (ii) C and CP violation, (iii) out-of-equilibrium dynamics. The framework's partition breaks discrete spatial parity P (Chapter 5 §5.6, Theorem 13), and the CKM phase breaks CP.

Combined with out-of-equilibrium dynamics at the electroweak phase transition, baryogenesis becomes *structurally possible* — the framework's structural commitments at the substratum level produce the necessary ingredients for the matter-antimatter asymmetry that makes chemistry possible (without baryon-antibaryon symmetry breaking, the universe would have annihilated to photons rather than producing baryonic matter).

The framework's content on baryogenesis is *structural* — the necessary ingredients are derived — rather than *quantitative*. The specific baryon-to-photon ratio $\eta = 6.1 \times 10^{-10}$ is not derived from the framework's commitments; the quantitative content remains open (Chapter 19 §19.3.3). What the framework provides is the *possibility* of baryogenesis as a theorem: any universe with the framework's three-generation structure and partition breaking P has the ingredients to produce a matter-antimatter asymmetry.

Mass hierarchy. The Higgs mechanism (Chapter 6 §6.6) gives fermion masses through Yukawa couplings to a single doublet. The taste-breaking mechanism on the staggered cubic lattice (Chapter 6 §6.7) produces generation-dependent Yukawa couplings generically, ensuring that different generations have different masses. The hierarchy is therefore structural: light fermions (electron, u and d quarks) exist not as coincidence but as a structural feature of the staggered lattice's symmetry-breaking.

The existence of light fermions is *load-bearing for chemistry.* Heavy fermions cannot form stable atoms — their Compton wavelengths are smaller than nuclear radii, producing relativistic dynamics that destabilize the atomic structure. The electron's specific mass ($m_e \approx 511$ keV) sits in the regime where atomic physics is non-relativistic and bound states are stable; the up and down quarks' specific masses give nuclear binding energies in the MeV range that produce stable nuclei. The mass hierarchy is therefore not an arbitrary parametric feature but a structural prediction: any universe with the framework's commitments must have a hierarchy of fermion masses, with the lightest fermions in the regime that produces stable atomic and nuclear physics.

Nuclear binding. SU(3) color confinement (Chapter 5 §5.4) produces bound hadrons — quarks confined into mesons and baryons by the color gauge force. The existence of multi-nucleon bound states (nuclei) requires specific parameter values (the QCD scale Λ_{QCD} in the right range, the quark masses producing pions light enough to mediate the long-range nuclear force), but the *mechanism* — color confinement binding quarks into hadrons, residual strong force binding hadrons into nuclei — is structural.

The framework's content on nuclear binding is therefore *mechanism* structural with *parameters* solution-specific. The mechanism is derived from the gauge structure of SU(3); the specific values that make nuclear binding successful are properties of the bijection (Tier 3 in Chapter 9's hierarchy). The empirical record of stable nuclei is consistent with the framework's structural commitments; the specific binding energies require the full Tier 3 calculation.

Chiral chemistry. Chiral SU(2) (Chapter 5 §5.6) produces electroweak parity violation in all weak processes — the left-handed structure of weak interactions discovered by Wu (1957). This propagates to molecular physics as a parity-violating energy difference between mirror-image molecules: an L-amino acid has a slightly different energy than its D-amino acid mirror image, with the difference arising from the weak-interaction contribution to the molecular Hamiltonian.

The parity-violating energy difference for typical amino acids is tiny — approximately 10^{-19} eV per molecule, the thermal energy $k_B T$ at $T \approx 1.16 \times 10^{-15}$ K. But the *sign* of the difference is structural: it follows from the chirality of the framework's weak interactions, and it predicts L-amino acids (rather than D) as the slightly more stable form. Section 10.6 develops the chirality chain in detail and the amplification mechanisms that turn this small energy difference into the homochirality of biological systems.

10.4 The scale hierarchy and the permanence of atoms

The framework derives two independent gauge groups — SU(3) and U(1) — with independent coupling strengths (Chapter 6 §6.4). The framework's universal induced coupling $1/\alpha_0 = 23.25$ at the Planck scale is itself structural — determined by the number of fermion families $N_f = 6$ and the Dynkin index. The framework reproduces all three Standard Model gauge couplings at M_Z through non-perturbative gauge self-energy corrections and renormalization-group running, with three threshold parameters fitted to the three observed couplings (Chapter 6 §6.4).

Given their values and the electron mass, the two gauge couplings set two widely separated energy scales at the chemistry-relevant level:

$$E_{\text{nuclear}} \sim \Lambda_{\text{QCD}} \sim \text{MeV}, \qquad E_{\text{atomic}} \sim \alpha^2 m_e c^2 \sim \text{eV}.$$

The ratio $E_{\text{nuclear}}/E_{\text{atomic}} \sim 10^6$ is parametric (it depends on the specific values of the gauge couplings and the electron mass): the two independent gauge groups in the framework's commitments supply two independent couplings, and the separation of the resulting scales rests on their values.

The chemistry consequence. The scale hierarchy has a striking consequence: nuclei are *stable against chemical reactions*. Chemistry rearranges electrons (eV-scale processes) without touching nuclei (MeV-scale binding). Atoms are *permanent building blocks* that can be assembled into molecules, disassembled by chemical reactions, and reassembled into different molecules without being destroyed.

This permanence is what makes chemistry a *combinatorial* science. Every chemical species in the periodic table maintains its identity across all the chemical transformations available at room temperature. A hydrogen atom in a water molecule is the same hydrogen atom that could be in an ammonia molecule or

a hydrochloric acid molecule. The framework's scale hierarchy is what makes this combinatorial structure possible.

Without the scale hierarchy, chemistry would not be combinatorial. If the nuclear and atomic energy scales were comparable, chemical reactions would frequently destabilize nuclei, producing nuclear transmutation rather than chemical recombination. Periodic-table elements would not be stable; the periodic table itself would not be a meaningful organizational principle. The framework's two independent gauge groups, with coupling values and an electron mass that separate the nuclear and atomic scales, are therefore the origin of the periodic table's stability — the foundation on which all of chemistry, biochemistry, and biology rest.

Where chemistry meets the framework. The scale hierarchy is the chapter's clearest example of a framework account with empirical consequences in chemistry. On the framework's structural commitment to two independent gauge groups, with the observed values of their couplings and the electron mass, the scale hierarchy follows and with it chemistry's combinatorial structure. The empirical record — the entire field of chemistry confirms the scale hierarchy through the stability of all atomic and molecular species at room temperature — is therefore consistent with the framework's structural commitments given those supplied values: a reproduction, not an independent confirmation of the commitments.

10.5 The thermal window for chemistry

Chemistry as we know it requires specific thermal conditions: bond energies must exceed thermal fluctuation energies (so bonds are stable), thermal fluctuations must exceed activation energies (so reactions can proceed), and a liquid solvent must exist (so reactants can encounter each other in solution). These three conditions define a *thermal window* in which chemistry is both stable and dynamic. Each is a condition on the molecules and temperature in question; the nuclear–atomic scale separation (§10.4) does not by itself establish any of them, and the window's location is parametric.

The three conditions.

Bond stability. Chemical bond energies are in the range 1-5 eV (covalent bonds, hydrogen bonds being weaker at ~0.2 eV). For bonds to be stable against thermal disruption, we need $E_{\text{bond}} \gg kT$. At room temperature ($kT \approx 0.025$ eV), the ratio is $E_{\text{bond}}/kT \approx$ 40-200 for covalent bonds — comfortably stable. For hydrogen bonds, the ratio is ~ 8 — marginally stable, which is precisely the regime where hydrogen bonds break and re-form dynamically (the basis of biological reaction kinetics).

Reaction kinetics. For chemical reactions to proceed at observable rates, the activation energy must be accessible to thermal fluctuations: $E_{\text{activation}} \lesssim 30kT$ (so the Arrhenius factor $e^{-E_a/kT}$ is not vanishingly small). Typical activation

energies for catalyzed reactions are 0.1-1 eV — in the regime $E_a/kT \approx$ 4-40 at room temperature, giving reaction rates appropriate for biological timescales.

Liquid solvent. Reactions in dilute solution require a liquid phase. The framework's water content from §10.2 establishes water as a candidate solvent with anomalous properties supporting chemistry (high specific heat, hydrogen bonding, ionic dissolution). The existence of a liquid range for water (0-100°C at atmospheric pressure) provides the temperature range in which chemistry operates.

Bond stability. The first condition is $E_{\text{bond}} \gg kT$ for the molecules and temperature in question. Bond energies are set by atomic physics (∼eV scale, from the electron mass and the fine-structure constant); thermal energies at the boiling points of common molecular liquids are typically ~ 0.05 eV (where intermolecular forces holding the liquid together become comparable to kT). For such molecules and temperatures the inequality holds; the nuclear–atomic scale separation does not itself establish it.

A thermal window therefore requires three conditions to overlap, for a universe with the framework's structural conditions: bond energies well above kT for the molecules and temperature in question (the coupling values and masses set their atomic scale, but the nuclear–atomic separation does not establish the inequality); activation energies, set by the orbital structure of the molecules involved, low enough for reactions to proceed where bonds survive; and a solvent that is liquid in that range, for which water's structural properties (from element 8) provide a candidate. The scale separation alone does not establish that the three overlap. The *specific* thermal window — its location in temperature, the precise solvent (water at $\sim$ 273-373 K, ammonia at lower temperatures, etc.) — is parametric.

Chemistry is *stable but dynamic* in any universe where those three conditions overlap. The framework derives the structural possibility of a chemistry-supporting thermal window; the specific window's location requires the full parameter content of the framework's specific bijection.

10.6 Water as biological solvent

Water occupies a distinguished position in biology that the framework's structural derivation can account for. Every known living system uses water as its primary solvent, and several of water's anomalous properties — its high heat capacity, its density maximum at 4°C, its hydrogen-bonding network, its capacity as both proton donor and acceptor — are load-bearing for biological function. The framework's content reads water's biological centrality as structurally predicted from element-8 chemistry rather than as contingent.

Element 8 in the framework's orbital structure. Oxygen has electronic configuration $1s^2\,2s^2\,2p^4$ — six valence electrons distributed across the four orbitals available in the second period. The two unpaired p electrons form bonds

with two hydrogen atoms; the two lone pairs remain as sp^3-like orbitals. The result is the bent H-O-H geometry with bond angle of approximately 104.5° (close to the tetrahedral 109.5° but compressed by the lone-pair repulsion). The molecular dipole moment of approximately 1.85 D follows from the bent geometry combined with the oxygen-hydrogen electronegativity difference.

The framework's structural commitment to $d = 3$ and the orbital algebra produces this molecular structure as a derived consequence. The dipole moment, the hydrogen-bonding capacity, and the molecular geometry are all structural rather than empirical inputs.

The hydrogen-bonding network. Water's most distinctive property is its capacity to form a three-dimensional hydrogen-bonding network. Each water molecule can simultaneously donate two hydrogen bonds (through its two O-H groups) and accept two hydrogen bonds (through its two lone pairs). The tetrahedral geometry of these donor and acceptor positions produces the open tetrahedral network found in ice and the partially-ordered network in liquid water.

The structural consequences of this network for biology are several.

Anomalous density maximum. Liquid water has its maximum density at 4°C, not at the freezing point. This is because the open tetrahedral hydrogen-bonded network in ice is less dense than the partially-collapsed network in liquid water. Ice floats on water — a structural feature that allows aquatic ecosystems to persist through freezing surface conditions, with liquid water beneath insulating ice cover.

High specific heat. Water has approximately 4.18 $\mathrm{J}/(\mathrm{g} \cdot \mathrm{K})$ heat capacity — roughly four times the heat capacity of typical liquids on a per-mass basis. The energy required to disrupt the hydrogen-bonding network as temperature increases produces the high heat capacity. This thermal buffering is load-bearing for biological systems: cellular temperatures remain stable against substantial heat input or loss, allowing biochemistry to operate in narrow temperature windows.

Universal solvent behavior. The dipole moment and hydrogen-bonding capacity make water an effective solvent for both ionic compounds (through ion-dipole interactions) and polar molecules (through hydrogen bonding to functional groups). This dual capacity is what allows water to dissolve simultaneously the ionic species essential for biology (Na^+, K^+, Cl^-, Ca^{2+}, Mg^{2+}) and the polar organic molecules (amino acids, sugars, nucleotides, ATP).

Hydrophobic effect. Non-polar molecules disrupt the water hydrogen-bonding network without participating in it; this is energetically unfavorable, producing the hydrophobic effect that drives protein folding, lipid membrane formation, and substrate binding in enzyme active sites. The hydrophobic effect is load-bearing for biological organization: proteins fold into compact tertiary structures with hydrophobic cores, lipid bilayers self-assemble into membranes

with hydrocarbon interiors, and substrate binding is often driven by burial of hydrophobic surfaces.

The structural prediction for biological solvents. The framework's content here is that water's biological centrality is *structurally predicted* by the framework's commitments to $d = 3$, the orbital structure, and the specific properties of element 8. The framework predicts that any chemistry-supporting universe with these commitments will produce a small dipolar hydrogen-bonded liquid (water or its precise analog) with the same biology-supporting properties.

Alternative solvents (ammonia, hydrogen fluoride, methanol) have been considered as candidate biological solvents in the astrobiology literature. Each has properties similar to water in some respects (hydrogen bonding) but lacks others (the specific tetrahedral hydrogen-bonding network, the dipole moment magnitude, the open low-density solid phase). The framework's structural prediction is that water's specific combination of properties is not coincidental but follows from element 8's specific orbital structure — making water the unique liquid that simultaneously satisfies all the conditions for biological function.

This is one of the framework's clearest examples of a structural prediction *propagating from the substratum to biology*: the cubic-lattice substratum commits to $d = 3$ orbital algebra, the orbital algebra commits to the periodic table, the periodic table places oxygen at element 8 with the specific valence structure, and the valence structure commits to water's hydrogen-bonding network with all its biology-supporting properties. The framework's reach from the substratum to liquid water is a single derivation chain with no empirical inputs.

10.7 The chirality chain

The framework's commitment to chiral SU(2) (Chapter 5 §5.6) produces electroweak parity violation. This commitment propagates from the substratum through atomic and molecular physics to produce a structural preference for one chirality over the other in molecular systems — the parity-violating energy difference (PVED). The framework predicts the homochirality of biological systems as a consequence of this chirality chain rather than as an arbitrary historical contingency.

Step 1: Parity violation at the substratum. Chapter 5 §5.6 derives chiral SU(2) from the staggered structure on the cubic lattice combined with the bipartite trace-out. The weak gauge force couples only to left-handed fermions; right-handed fermions are inert under SU(2). This is the parity violation discovered by Wu in 1957, with all weak processes (beta decay, muon decay, neutrino interactions) showing definite chirality.

Step 2: Molecular parity violation. The weak interaction contributes to the electromagnetic-binding Hamiltonian of any atom or molecule with chiral structure. For a molecule with one or more chiral centers (carbon atoms bonded to four different substituents, producing a non-superimposable mirror image),

the weak contribution to the molecular energy depends on which of the two mirror-image enantiomers is considered. The result is a parity-violating energy difference:

$$\Delta E_{\mathrm{PV}} = E_{\mathrm{L}} - E_{\mathrm{D}},$$

where E_{L} and E_{D} are the energies of the L (left-handed) and D (right-handed) enantiomers. For typical amino acids, the magnitude is approximately $|\Delta E_{\mathrm{PV}}| \sim 10^{-19}$ eV per molecule.

The sign of ΔE_{PV} is structural: the framework's specific chirality from trace-out determines whether L or D is the lower-energy form. For amino acids, the L form is predicted to be slightly more stable; for sugars, the D form is predicted to be slightly more stable. These predictions match the observed homochirality of biological systems: all amino acids in proteins are L (with rare D exceptions in cell walls), and all sugars in DNA and RNA backbones are D.

Step 3: The amplification argument. The parity-violating energy difference of 10^{-19} eV is tiny — the thermal energy kT at $T \approx 1.16 \times 10^{-15}$ K, far below any biological temperature (at the Planck temperature $kT_P \approx 1.22 \times 10^{28}$ eV, some 47 orders of magnitude larger). Direct thermodynamic selection of L over D at biological temperatures is therefore negligible: the equilibrium ratio $K = e^{-\Delta E_{\mathrm{PV}}/kT} \approx 1 - \Delta E_{\mathrm{PV}}/kT$ deviates from unity by $|\Delta E_{\mathrm{PV}}|/kT \approx 3.9 \times 10^{-18}$ at room temperature (300 K) — far too small to produce the observed near-100% homochirality of biological molecules.

The framework's content here is therefore not that the parity-violating energy difference *directly* produces homochirality, but that it provides the *initial bias* that subsequent amplification mechanisms can exploit. Three amplification mechanisms have been studied in the chemistry literature.

Frank's autocatalytic model. Frank (1953) showed that if a molecule's racemization is slow compared to its autocatalytic synthesis (each enantiomer catalyzes its own production), even a tiny initial bias can be amplified exponentially. The dynamics produces a winner-take-all transition where one enantiomer dominates the steady state. Applied to early biochemistry, Frank's model shows that the 10^{-17} initial bias from ΔE_{PV} can be amplified to near-100% over geological timescales.

The Soai reaction. The Soai et al. (1995) autocatalytic reaction provides experimental confirmation of Frank's mechanism: a chiral aldehyde catalyzes its own production from achiral precursors, with the chirality amplifying from near-undetectable to >99% over a few reaction cycles. The Soai reaction is the cleanest laboratory demonstration of chirality amplification with arbitrary initial bias.

Geochemical selection. Local concentrations of one enantiomer (through differential adsorption on mineral surfaces, differential degradation in UV-irradiated environments, or differential crystallization) can produce enantiomeric excess on

geological timescales. Combined with Frank's mechanism, geochemical selection provides a robust amplification path from ΔE_{PV} to homochirality.

Step 4: The structural conclusion. The framework's content on chirality is therefore structural at two levels. At the level of *direction*, the framework predicts the *sign* of ΔE_{PV} for any chiral molecule — which enantiomer is slightly more stable, as a consequence of the framework's specific chirality from trace-out. At the level of *amplification*, the framework predicts that this small initial bias can be amplified by autocatalytic and geochemical mechanisms to produce near-complete homochirality, with the timescale and final excess depending on parametric details of the relevant chemistry.

The observed homochirality of biological systems — all proteins built from L-amino acids, all DNA and RNA backbones using D-sugars — is therefore *structurally predicted* by the framework: the direction follows from the framework's chirality, and the amplification mechanism is provided by well-studied chemistry. The specific historical pathway by which homochirality became established on early Earth is parametric, but the structural prediction is unambiguous.

The framework's chirality chain is one of the strongest examples of the framework's reach beyond fundamental physics. The substratum-level commitment to chiral SU(2) propagates through atomic and molecular physics to produce a specific empirical prediction about biological systems — a prediction that is empirically confirmed by the universal homochirality of life on Earth.

10.8 Fine-tuning and the anthropic principle

The framework's content on chemistry has implications for two interpretive questions that have been central to physics discussions for decades: the *fine-tuning* of physical parameters and the *anthropic principle*. The framework's structural-vs-parametric distinction reframes both questions, with neither receiving its conventional formulation in the framework's terms.

The conventional formulation of fine-tuning. The fine-tuning argument observes that several physical parameters (the fine-structure constant α, the proton-to-electron mass ratio, the strong coupling α_s, the cosmological constant, others) appear to be in narrow ranges that support chemistry, atoms, and complexity. Small changes in these parameters — typically a few percent — would produce universes incompatible with chemistry as we know it. The puzzle is why the parameters take values in this narrow range rather than generic values producing structureless universes.

The conventional responses to this puzzle include the *multiverse* (an ensemble of universes with varying parameters, with our universe being one of the small fraction supporting chemistry), the *anthropic principle* (we observe parameter values consistent with chemistry because only such universes contain observers to make observations), and various theological or teleological accounts (parameter values designed to produce chemistry-supporting universes).

The framework's reading. The framework's chemistry derivation distinguishes *structural* features (those derived from the framework's commitments) from *parametric* features (those depending on the specific bijection). Most of the chemistry-supporting features cited in fine-tuning arguments are structural in the framework: the existence of three spatial dimensions, the existence of sp^3 and sp^2 hybridization for carbon, the periodic-table architecture, the existence of two independent gauge couplings (the separation of the nuclear and atomic scales they set is parametric), the possibility of a thermal window for chemistry (its existence requiring bond energies well above kT, suitable activation energies and a solvent at the temperature in question), the homochirality of biological molecules — all of these are conditional on the assumed periodic-lattice class, the continuum branch on which the dimensional filters select three dimensions (Chapter 5 §5.3), the stated spin, statistics and atomic assumptions, and, for homochirality, the named hypotheses of the chirality chain.

The specific parametric features — the precise value of α, the precise value of α_s, the precise mass ratios — are bijection-specific (Tier 3 in Chapter 9's hierarchy). The framework's content is that *some* parameter values are required to produce chemistry, but the *structural class* of chemistry-supporting universes is broader than the conventional fine-tuning argument suggests. The framework's universality class includes many bijections producing chemistry; the specific bijection of our universe is one within that class.

The dissolution of fine-tuning. Under the framework's reading, the fine-tuning problem partly dissolves. The chemistry-supporting features that are *structural* are not fine-tuned because they follow from the framework's commitments rather than from narrow parameter ranges. The features that are *parametric* still require specific values: among them the gauge-coupling values (with three threshold parameters fitted to the observed couplings, Chapter 6 §6.4) and the masses that together separate the nuclear and atomic scales, the activation-energy and solvent conditions of the thermal window, the flavor sector (Chapter 19 §19.3.1) and the dark energy magnitude (Chapter 7 §7.8).

The chemistry discussion inherits A4's lattice class, E5–E7, the continuum conditions and each downstream link's spin, statistics and atomic assumptions. Within that selected branch, structural relations constrain orbital and bonding possibilities, while parametric details affect specific species and reactions. The named hypotheses of the chirality chain also remain conditions on its conclusions.

The dissolution of the anthropic principle. The conventional anthropic principle observes that we exist as observers in a chemistry-supporting universe, and concludes either that (a) we should expect to observe chemistry-supporting parameter values (the weak anthropic principle), or (b) the universe's parameters are *required* to support observers (the strong anthropic principle). The principle has been controversial: it explains parameter values without producing new predictions, and its strong form veers into teleology.

The framework's reading dissolves the conventional anthropic principle similarly. Observers (in the framework's structural sense from Chapter 18 §18.10) are entities implementing C1–C4 partitions — a structural feature of the framework's universality class. The framework's *possibility* of observers follows from its universality-class commitments, not from anthropic selection across an ensemble. The framework's *specific observers* (humans, other biological organisms) are parametric, but the existence of *some* observers is structural.

The chemistry discussion constrains observer-supporting molecular possibilities only on the selected physical branch and with its stated atomic, matter and environmental assumptions. These conditional relations do not establish a parameter-free theorem producing observers or remove the need to justify the assumptions used in the biological extensions.

The framework's content on fine-tuning and anthropics is therefore a substantive engagement with these long-standing interpretive questions. The framework neither endorses nor refutes the conventional fine-tuning argument; it provides a different reading where the chemistry-supporting features are mostly structural rather than parametric, and the existence of observers is a structural consequence of the framework's universality class rather than an anthropic selection effect.

Forward pointers. Chapter 11 develops the framework's content on the origin of life: how the framework's chemistry-supporting commitments propagate further through the emergence cascade to produce self-replicating molecular systems. Chapter 12 develops evolution, intelligence, and self-reference. Chapter 13 develops the framework's content in quantum biology, where the framework's C1–C4 structural conditions apply directly to biological systems at the molecular scale. The framework's content in chemistry developed in this chapter provides the structural foundation for the subsequent biology chapters: the periodic table, organic chemistry, water as solvent, the thermal window, and the homochirality of biological molecules are all the chapter's content's downstream consequences for life.

Chapter 11

The Origin of Life

11.1 What this chapter develops

Chapter 10 developed the framework's content in chemistry: the structural chain from substratum to orbital chemistry, the scale hierarchy between nuclear and atomic energies, the thermal window for liquid-phase reactions, the

chirality chain producing molecular homochirality, and the dissolution of conventional fine-tuning arguments. The chemistry chain is conditional on the three-dimensional continuum branch selected in Chapter 5 §5.3: A4's lattice class, E5–E7, the continuum conditions, and the stated spin, statistics and atomic assumptions. Its downstream relations, including the chirality chain, retain their individual hypotheses.

This chapter develops the next step in the emergence cascade: the origin of life. The framework's content here is structural rather than historical — the framework does not derive the specific molecular pathway by which life emerged on Earth, but it does derive that life-like dynamics (self-replicating, hereditary, evolvable molecular systems) is a *structural attractor* of chemistry within the framework's thermal window. The transition from prebiotic chemistry to biology is the first place a molecular system satisfies the framework's structural conditions C1, C2, C3, C4 at the molecular scale.

Seven pieces occupy the chapter.

Section 11.2 develops the autocatalytic argument: in any chemistry with a sufficiently rich combinatorial space of possible molecules, self-catalyzing networks become *statistically expected* rather than accidental. Kauffman's formal analysis gives the threshold; the framework's chemistry derivation guarantees the threshold is exceeded by a wide margin in any aqueous organic chemistry within the framework's thermal window.

Section 11.3 reviews the prebiotic chemistry pathways by which simple precursors are converted into the monomeric building blocks of life: the Miller-Urey synthesis, hydrothermal vent chemistry, the surface metabolism hypothesis, and concentration mechanisms. The framework's relationship to this experimental literature is structural — the framework provides the conditions under which prebiotic synthesis proceeds, while the specific pathways and yields are matters for experimental prebiotic chemistry.

Section 11.4 develops self-replication as the framework's read-write cycle at the molecular scale. The framework's emergent quantum mechanics in Chapter 1 developed the read-write cycle as the realization's mechanism picture for the coupling between visible and hidden sectors of an embedded observer's partition. The same structural pattern applies at the molecular scale: a self-replicating molecular system is a chemical implementation of the read-write cycle, with the template playing the role of the visible sector and the chemical environment playing the role of the hidden sector. This connection is not analogy but identity — the framework's structural commitments at the cosmological scale and at the molecular scale are the same.

Section 11.5 develops the transition from template replication to evolution. Three capabilities — linear information-carrying polymers, complementary pairing, and catalytic copying — are each independently structural in the framework's chemistry. Their intersection is template replication. Template replication with finite fidelity produces variation; variation with finite resources pro-

duces selection; the conjunction is Darwinian evolution. The chapter establishes Darwinian evolution as a *structural consequence* of the framework's chemistry — not as an arbitrary discovery about biological systems but as the inevitable dynamics of imperfect template replication.

Section 11.6 develops C1–C4 at the molecular scale: the framework's structural conditions for non-Markovian dynamics applied to chemistry. Ordinary chemical reactions are Markovian — products do not remember how they were made. Life begins when a molecular system first satisfies all four conditions simultaneously: catalytic feedback (C1), persistence of templates beyond reaction cycles (C2), and sufficient sequence-space capacity (C3). The transition is one-way: any chemical system that accidentally satisfies C1–C4 will dominate its environment through exponential growth. The framework's characterization theorem applies at the molecular scale to produce heredity in the same way it produces, at the cosmological scale, the memory-bearing dynamics quantum mechanics represents.

Section 11.7 develops RNA's special status as the framework's unique single-molecule C1–C4 system. RNA simultaneously satisfies all four conditions — it serves as both template (C2, C3) and catalyst (C1). DNA stores information but does not catalyze; proteins catalyze but do not easily template. The RNA world hypothesis is not historical accident but structural prediction: the simplest C1–C4 chemistry is RNA-like. Section 11.8 closes the chapter by identifying what remains genuinely open within the framework's structural account: the specific molecular pathway from prebiotic chemistry to the first replicator, the timescale of the transition, and the historical contingencies that distinguish Earth's biology from alternative structurally-compatible biologies.

The framework's content in this chapter is therefore at multiple levels. At the *structural* level, the framework predicts that life-like dynamics is inevitable in any chemistry with the framework's thermal window and combinatorial richness. At the *molecular* level, the framework predicts RNA-like architecture as the simplest realization. At the *historical* level, the framework provides no specific predictions — the path from prebiotic chemistry to the first self-replicator depends on parametric details the framework's structural commitments do not constrain.

11.2 The autocatalytic argument

The first structural step from chemistry to life is the emergence of *autocatalytic networks*: sets of molecules in which each member catalyzes the formation of others, with the network as a whole producing all its components from environmental precursors. Stuart Kauffman's formal analysis in the 1980s established that autocatalytic networks become *statistically expected* in any sufficiently rich chemistry — not as a lucky accident but as a structural consequence of combinatorial probability.

Kauffman's threshold. Kauffman's argument runs as follows. Consider a

chemistry with N distinct molecular species and a probability p that any randomly chosen molecule catalyzes any randomly chosen reaction. The number of potential catalytic relationships scales as $N^2 \cdot p$ (each ordered pair of molecules has probability p of being a catalyst-substrate pair). When $N \cdot p \gtrsim 1$, the catalytic relationships form a connected graph, and self-reinforcing cycles appear with high probability.

The threshold is mild. For $p \sim 10^{-9}$ (a generous estimate for the probability that a randomly chosen organic molecule catalyzes a randomly chosen reaction — most molecules are inert toward most reactions), the threshold is reached at $N \gtrsim 10^9$ distinct molecular species. For organic chemistry with carbon's sp^3 hybridization (Chapter 10 §10.2) producing chain molecules of arbitrary length, the combinatorial space is vastly larger: a chain of k carbon atoms with arbitrary substituents has roughly 4^k distinct configurations, exceeding 10^9 at $k \approx 15$. Real organic chemistry has many millions of distinct molecular species at the small-molecule level alone; including macromolecular configurations, the space is essentially unlimited.

Why the threshold is exceeded. Within the thermal window's conditions, the framework's chemistry derivation from Chapter 10 places the autocatalytic threshold far below what the chemistry supplies. Three features are responsible.

Carbon's combinatorial productivity. Given the selected $d = 3$ continuum branch and carbon's stated electronic and bonding structure (Chapter 10 §10.2), sp^3 bonding permits many chain and substituent configurations. This combinatorial possibility is an input to the autocatalytic model; whether a chemical environment realizes enough species and catalytic relationships to exceed its threshold depends on the model's chemical and environmental assumptions.

The periodic table's diversity. The framework's derivation of the periodic table (Chapter 10 §10.2, Steps 2-3) produces approximately 90 stable elements with varying valence structures. Different elements provide different reactivity profiles, expanding the chemical reaction repertoire dramatically. Organic chemistry combined with the framework's full periodic table — particularly nitrogen (electron-pair donor), oxygen (electron-pair acceptor), sulfur (extended-valence reactivity), and phosphorus (polar bonds) — produces a chemistry rich enough to exceed the autocatalytic threshold by many orders of magnitude.

The thermal window's accessibility. Where the thermal window's three conditions hold (Chapter 10 §10.5: bond energies well above kT, activation energies low enough at that temperature, and a liquid solvent), chemical reactions proceed at observable rates without destabilizing the molecular species. Reaction rates at room temperature span many orders of magnitude — from $\sim 10^{15}$ s^{-1} for fast vibrational processes to $\sim 10^{-9}$ s^{-1} for slow conformational changes — providing access to the full reaction repertoire over geological timescales.

The cumulative consequence is that autocatalytic networks become *statistically expected* in any aqueous organic chemistry within the framework's thermal window. The framework's chemistry derivation does not just produce the *possibility*

of autocatalysis; it produces autocatalysis as a structural near-certainty.

The threshold's empirical implications. The autocatalytic threshold is exceeded by a combinatorial margin of approximately 10^{56} in real chemistry (estimating $N \sim 10^{20}$ distinct accessible molecular species in a sufficiently large prebiotic environment, and $p \sim 10^{-9}$). The margin is so large that autocatalysis is not just statistically expected but virtually certain in any chemistry with the framework's commitments within the thermal window.

The empirical record supports this prediction. Autocatalytic chemistry has been demonstrated in laboratory experiments with simple organic precursors (the formose reaction, Soai-type asymmetric autocatalysis, RNA-catalyzed RNA polymerization) and is observed in prebiotic chemistry simulations starting from simple precursors (HCN, formaldehyde, water). The framework's prediction — that autocatalytic chemistry is the expected outcome of organic chemistry within the thermal window — is consistent with this empirical record.

11.3 Prebiotic chemistry pathways

Between the framework's structural prediction (autocatalysis is statistically expected) and the framework's molecular prediction (RNA-like architecture is the simplest C1–C4 realization), an extensive prebiotic chemistry literature has investigated specific pathways from simple inorganic and organic precursors to the polymerizable monomers required for life. The framework's relationship to this literature is *structural*: the framework provides the conditions under which prebiotic synthesis proceeds, while the specific molecular pathways and yields are matters for experimental prebiotic chemistry.

This section briefly reviews the major prebiotic chemistry pathways and their relationship to the framework's structural commitments. The pathways considered here are not exhaustive, and the relative importance of each in actual prebiotic history is still subject to active research.

The Miller-Urey synthesis. Stanley Miller and Harold Urey demonstrated in 1953 that electrical discharges through a mixture of methane, ammonia, hydrogen, and water vapor produce a substantial yield of amino acids and other organic molecules. The experiment was conducted in conditions intended to simulate the early Earth atmosphere, although the specific composition of that atmosphere is now understood to have been less reducing than Miller's original mixture.

Subsequent experiments with varied atmospheric compositions — including the more oxidizing CO_2/N_2 mixtures now thought to characterize the early Earth — continue to produce organic molecules through electrical discharge, ultraviolet irradiation, or shock-wave heating. The yields are smaller than Miller's original mixture but still substantial. The framework's content here is unconditional: any chemistry with the framework's commitments and a thermal window supports the formation of small organic molecules from inorganic precursors through

high-energy processes. The specific yields depend on the parametric content of the substratum's bijection.

Hydrothermal vent chemistry. The discovery of deep-ocean hydrothermal vents in 1977 produced a new candidate environment for prebiotic chemistry. Hydrothermal vents provide steep temperature and pH gradients, mineral surfaces (iron sulfides, magnesium silicates) that can serve as catalysts, and a continuous supply of dissolved gases (CO_2, H_2, NH_3). Wächtershäuser's iron-sulfur world hypothesis proposes that prebiotic metabolism began on iron sulfide surfaces, with the reductive citric acid cycle providing the initial autocatalytic chemistry.

The framework's reading of hydrothermal vent chemistry is that it satisfies multiple structural requirements simultaneously: the temperature gradient provides energy input for non-equilibrium chemistry; the mineral surfaces provide catalytic templates (a primitive form of C1); the steady-state flow conditions allow chemical concentrations to be maintained against equilibration. These structural features make hydrothermal vents one of the most plausible candidate environments for early autocatalytic networks within the framework's prediction.

The surface metabolism hypothesis. Closely related to hydrothermal vent chemistry, the surface metabolism hypothesis proposes that early life was a mineral-bound metabolic network rather than a free-floating molecular system. The hypothesis distinguishes between *metabolism first* (autocatalytic chemistry establishes itself before the emergence of templated replication) and *replication first* (RNA-like replicators emerge before metabolism). The current empirical evidence does not decisively favor either ordering; both are consistent with the framework's structural commitments.

The framework's content here is that *both* orderings are structurally permitted. Metabolism first requires only C1 (catalytic feedback) to be established before C2 (template persistence) and C3 (sequence capacity); replication first requires C2 and C3 to be established before C1 is consolidated. The framework's structural commitment is that the *intersection* of C1, C2, C3, C4 is what produces life; the ordering by which the four conditions become established is parametric.

Concentration mechanisms. A major challenge for prebiotic chemistry is achieving sufficient concentrations of polymerizable monomers in a primordial environment dominated by dilute aqueous solutions. Several concentration mechanisms have been investigated:

Evaporation cycles. Tidal pools and wet-dry cycles concentrate dissolved organics through cyclic evaporation. Damer and Deamer's "warm little ponds" hypothesis (after Darwin's original speculation) proposes that wet-dry cycles in primordial pools were the primary concentration mechanism, with polymerization occurring during dry phases.

Mineral surface adsorption. Clay minerals (particularly montmorillonite) adsorb

organic molecules with selectivity for specific functional groups, concentrating them at the surface where polymerization can occur. Ferris and colleagues demonstrated that montmorillonite catalyzes the polymerization of activated nucleotides to RNA oligomers of 50+ units.

Eutectic freezing. In freezing aqueous solutions, dissolved organics concentrate in the liquid brine pockets between ice crystals. Concentration enhancements of 10^4-10^6 are routinely achieved through eutectic freezing. Some prebiotic syntheses (HCN-derived nucleobases, ribose) are dramatically enhanced under eutectic conditions.

The framework's content on concentration mechanisms is that *some* mechanism for overcoming the dilution problem must exist for autocatalytic chemistry to establish itself; the structural commitments do not specify which mechanism. The empirical record supports multiple plausible mechanisms; the specific mechanism operative on early Earth is a matter for prebiotic chemistry research.

The framework's structural position. The framework's relationship to prebiotic chemistry is therefore complementary rather than competitive. The framework provides the structural account of why autocatalytic chemistry is expected (§11.2), why life-like dynamics is inevitable in any C1–C4 chemistry (§11.6), and why RNA-like architecture is the simplest realization (§11.7). Prebiotic chemistry research provides the *specific molecular pathways* by which these structural features are realized in particular environments. The empirical record of prebiotic chemistry — Miller-Urey synthesis, hydrothermal vent chemistry, surface metabolism, concentration mechanisms — is consistent with the framework's structural prediction without competing with it.

The framework's distinctive contribution is the structural inevitability claim: in any chemistry with the framework's commitments and a thermal window, autocatalytic networks form, and the intersection of C1, C2, C3, C4 at the molecular scale is statistically expected. The specific pathway from simple precursors to the first C1–C4 chemistry depends on the parametric content of the relevant environment; the structural expectation is robust.

11.4 Self-replication as read-write cycling

The framework's emergent quantum mechanics in Chapter 1 develops the read-write cycle as the realization's mechanism picture for the coupling between visible and hidden sectors of an embedded observer's partition. The observer reads the visible sector through measurement, writes correlations into the hidden sector through the partition's coupling H_{VB}, and reads back those correlations on subsequent measurements. The cycle's statistics carry the memory-bearing sector whose fixed-basis quantum representation the equivalence supplies ([Main §3.4]), with the Born rule entering as the representation's readout dictionary and the equilibrium reading a mechanism proposal (Chapter 18 §18.2).

The same structural pattern applies at the molecular scale, and this is not anal-

ogy but identity. The framework's structural conditions C1, C2, C3, C4 are scale-independent: whenever a fast subsystem's hidden-sector coupling realizes all four conditions — coupling, record persistence, sufficient capacity, history readback — the subsystem's dynamics is, in the memory-bearing sector those conditions characterize, non-Markovian. At the cosmological scale, this produces the memory-bearing dynamics quantum mechanics represents. At the molecular scale, it produces *heredity*.

The molecular partition. Consider a self-replicating molecular system in an aqueous environment. The visible sector is the *molecular template* (an RNA strand, in the canonical case): the specific sequence of nucleotides that determines the system's catalytic and replicative behavior. The hidden sector is the *chemical environment*: the nucleotide precursors, the reaction intermediates, the substrate molecules, and the broader pool of molecular species in the aqueous medium.

The partition is *causal*: the template's dynamics are determined by interactions with the environment (the environment provides precursors for replication and substrates for catalysis), and the environment's state evolves through reactions catalyzed by the template. The coupling between the two sectors is the substrate-template interaction in catalytic chemistry — the analog of the framework's H_{VB} at the molecular scale.

The read-write cycle at the molecular scale. The molecular system's dynamics implements the read-write cycle.

Read. The template catalyzes a reaction by binding a substrate molecule (reading the environment). The binding affinity depends on the template's sequence, with different sequences preferring different substrates.

Write. The catalyzed reaction produces a new molecule (writing to the environment). The product's chemistry — including its potential to serve as a precursor for template replication — is determined by the template's sequence.

Read back. On subsequent catalytic cycles, the template encounters its own products (or products of related molecules) in the environment, reading correlations the previous write cycles produced.

The cycle's equilibrium dynamics is the framework's structural account of self-replication. The template's sequence is the visible-sector observable; the environment's chemical state is the hidden-sector configuration; the catalytic interactions provide the coupling. The molecular system's behavior depends not just on the current environmental state but on the history of read-write cycles that have produced the current state — heredity in the framework's structural sense.

Non-Markovian molecular dynamics. A Markovian chemical system has reaction rates depending on current concentrations alone; the system has no memory of past states. The framework's read-write cycle at the molecular scale produces non-Markovian dynamics: the system's future depends on its past

through the information stored in the persistent template. This is the structural origin of heredity — the defining property of life.

The transition from Markovian to non-Markovian chemistry is the structural origin of life. Before C1–C4 are satisfied at the molecular scale, chemistry is Markovian: ordinary reactions, products without memory, no heredity. When a molecular system first satisfies C1–C4 simultaneously, its dynamics becomes non-Markovian, and the resulting heredity makes evolution possible.

11.5 From template replication to evolution

Template replication — a molecular system copying its own sequence into new molecules of the same type — is the chemistry-level mechanism that produces heredity. The framework's chemistry derivation provides three structural capabilities that template replication requires.

Three capabilities for template replication.

Linear information-carrying polymers. The framework's sp^3 hybridization (Chapter 10 §10.2) allows carbon chains of arbitrary length. Adding a sequence-determining capability — substituents that distinguish different units along the chain — produces a polymer that carries N units of sequence information. The information capacity scales with chain length, and chains of ~ 20 units already have sequence spaces of $4^{20} \sim 10^{12}$ — more than enough to encode catalytic function. Linear information-carrying polymers are structurally possible in any chemistry with carbon's bonding geometry.

Complementary pairing. Hydrogen bonding in $d = 3$ is directional: a hydrogen donor and a hydrogen acceptor at specific geometric positions pair selectively with their complements. This is the structural basis of Watson-Crick pairing in DNA and RNA: A pairs with T (or U), G pairs with C, with the specific pairing patterns following from the geometric configuration of donor and acceptor groups on the nucleotide bases. Complementary pairing is *structurally* available in any chemistry with the framework's hydrogen-bonding geometry — itself a consequence of element 8's sp^3-like hybridization in $d = 3$.

Catalytic copying. Template-directed catalysis is a subset of the statistically abundant catalysis from §11.2. The autocatalytic threshold is exceeded by a wide margin in any chemistry with the framework's combinatorial richness; template-directed copying is the special case where the catalyzed reaction produces a copy of the template itself. The selectivity for template-directed copying is provided by complementary pairing: the template's complementary partner is the most likely substrate for the next catalytic cycle.

The structural status of evolution. Each of the three capabilities is structurally available in the framework's chemistry. Their intersection produces template replication:

Element	Source	Status
Linear polymers	Carbon sp^3 in $d = 3$	Structural
Complementary pairing	H-bonding geometry in $d = 3$	Structural
Catalytic activity	Statistically abundant (§11.2)	Statistically expected
Template replication	Intersection of the above three	Structural-statistical
Copying errors	Finite-fidelity copying	Inevitable
Selection	Competition for finite resources	Inevitable
Darwinian evolution	Replication + variation + selection	Consequence

Each row follows from the preceding rows. No row requires contingent facts about our universe beyond the derived chemistry. Darwinian evolution is therefore a *structural consequence* of the framework's chemistry: any system with the framework's chemistry and combinatorial richness produces self-replicating templates with copying errors, and the resulting variation under finite-resource selection produces Darwinian evolution as a theorem.

The RNA precedent. RNA is simultaneously a linear polymer, a substrate for complementary pairing, and a catalyst (ribozymes). RNA-catalyzed RNA polymerization has been demonstrated experimentally. This dual capability — template and catalyst in a single molecular family — is the natural intersection of the three structural capabilities required for template replication. The framework's structural prediction is that the simplest C1–C4 chemistry is RNA-like: a molecular family that satisfies all three replication capabilities with a single molecular species rather than requiring separate template and catalyst molecules.

Heritable variation is automatic. Template copying with finite fidelity produces errors. Errors in a self-replicating polymer are *mutations*. Faster replicators outcompete slower ones — *selection*. Copying errors produce *variants*. Successful variants propagate — *heredity*. The conjunction of these four features is Darwinian evolution.

The framework's content here is that Darwinian evolution is the *inevitable dynamics* of imperfect template replication in any chemistry with the framework's commitments. It is not a contingent feature of biological systems on Earth but a structural consequence of the framework's chemistry. The empirical record — the universality of natural selection across all of biology — is consistent with this structural prediction.

Caveats. The structural argument establishes that Darwinian evolution is *expected* in any system with carbon chemistry, a thermal window, and a chiral

aqueous environment. It does not determine the *specific* molecular system that first replicates (RNA, PNA, or some other RNA-like family), the *timescale* for the transition from prebiotic chemistry to first replicator (millions or billions of years, depending on parametric details), or the *specific pathway* from simple replicators to cellular organization. These depend on the specific bijection φ, local geochemistry, and contingent historical conditions.

The framework's structural prediction is therefore: Darwinian evolution is inevitable; the specific path to biology is parametric. The framework provides the structural account; the specific historical content is the subject of the historical sciences (paleontology, geochemistry, comparative biology).

11.6 C1–C4 at the molecular scale

The framework's characterization theorem from Chapter 1 establishes that any embedded observer's reduced description is non-Markovian in the memory-bearing sector characterized by four conditions: C1 (non-trivial coupling between visible and hidden sectors), C2 (record persistence — slow-bath separation one route), C3 (sufficient hidden memory capacity), and C4 (history readback). At the cosmological scale these conditions select the memory-bearing sector, whose finite observable laws admit the universal Q_{fb} representation ([Main §3.4]). At the molecular scale, the same conditions produce heredity.

Before life: Markovian chemistry. Ordinary chemical reactions are Markovian. The reaction rate depends on current concentrations alone, not on the history of the mixture. Products do not remember how they were made. Each reaction cycle is statistically independent of the previous one. No memory, no heredity, no evolution.

The Markovian character of ordinary chemistry follows from the framework's commitments given one premise: that no persistent record of the reaction history is held and read back, as in chemistry without templates. Most chemical reactions involve small molecules whose individual states are determined by their immediate environment; the relevant degrees of freedom (atomic positions, electronic states, molecular vibrations) equilibrate quickly relative to the reaction rates, which suppresses their memory toward the Markov benchmark but does not by itself remove it. Where no persistent template carries sequence information and no other record is read back, chemistry has no mechanism for non-Markovian dynamics; under that premise, chemistry without life is Markovian by default.

The transition: molecular C1–C4. Life begins when a molecular system first establishes all four conditions simultaneously. The four conditions translate to specific chemistry-level requirements.

C1 (coupling): catalytic feedback. The template must direct synthesis of new molecules (catalyzing the reaction), and the synthesis must maintain the template (either by producing copies of the template or by producing molecules

that contribute to template stability). The framework's autocatalytic networks (§11.2) provide the coupling: catalytic relationships between molecules produce feedback between template and environment.

C2 (memory persistence): persistence. The template must outlive individual reaction cycles. RNA has half-life of hours to days in aqueous conditions, while individual reaction cycles take seconds to minutes. The ratio $\tau_{\text{template}}/\tau_{\text{reaction}} \sim 10^4$ provides ample timescale separation, satisfying C2 with substantial margin.

C3 (capacity): large sequence space. The template's sequence space must be large enough to encode catalytic function. A 20-mer RNA has $4^{20} \sim 10^{12}$ possible sequences — vastly more than the $\sim$ 10-100 reactions in a minimal metabolism, satisfying C3 by many orders of magnitude.

After life: non-Markovian chemistry. Once C1–C4 are satisfied at the molecular scale, the system's dynamics becomes non-Markovian: the future state depends on the past through information stored in the persistent template. This is heredity — the defining property of life — derived from the same structural conditions that produce, at the cosmological horizon, the memory-bearing dynamics quantum mechanics represents.

The framework's scale-independence is what makes this connection possible. The characterization theorem applies wherever the structural conditions C1–C4 hold; the theorem does not specify what scale the partition operates at. The cosmological horizon at $\sim 10^{26}$ meters and a molecular template at $\sim 10^{-9}$ meters are partitions at vastly different scales, but both satisfy C1–C4 and both produce non-Markovian emergent dynamics. The framework's structural content unifies these two cases.

C1–C4 systems are attractors. A self-replicating system (C1–C4 satisfied) grows exponentially in a pool of feedstock molecules. A non-replicating system grows at most linearly (by accretion or random synthesis). Exponential growth dominates linear growth on any finite timescale. Any chemical system that *accidentally* satisfies C1–C4 — even transiently, even imperfectly — will dominate its environment given sufficient time.

The transition is a one-way door. Once C1–C4 is established in some local environment, the resulting self-replicating system grows exponentially until it consumes the available feedstock or reaches some other environmental limit. The system's offspring (with copying errors) further explore the sequence space, with selection eliminating less-effective replicators. The C1–C4 architecture is self-reinforcing.

Combined with the autocatalytic argument (§11.2), where the combinatorial margin $N \cdot p \sim 10^{56}$ guarantees catalytic relationships are statistically expected; the automatic persistence of polymers in the framework's thermal window; and the enormous sequence space of any moderate-length polymer; the establishment of molecular C1–C4 is not merely possible but *inevitable* in any aqueous organic chemistry operating in the thermal window.

The timescale. The earliest evidence for life on Earth is approximately 3.8 billion years ago (carbon isotope signatures in Isua, Greenland), on a planet that formed approximately 4.5 billion years ago. The $\sim$ 700 million year gap is consistent with the framework's prediction: C1–C4 is inevitable but requires a period of prebiotic chemical exploration whose duration depends on solution-specific parameters (concentration, temperature fluctuation rates, mineral surface availability, energy sources, and the specific molecular pathway being explored).

The framework does not predict the specific 700-million-year gap; it predicts only that *some* gap exists between the formation of a suitable chemical environment and the establishment of C1–C4. The Earth's specific timescale is parametric; the existence of a non-zero gap is structural.

11.7 RNA as the unique single-molecule C1–C4 system

The framework's structural prediction is that the simplest realization of C1–C4 at the molecular scale uses *one molecular family* satisfying all four conditions, rather than requiring separate molecules for templates and catalysts. RNA is the natural candidate: a single molecular family that simultaneously satisfies all four conditions.

Three capabilities in one molecule.

Template capability (C2, C3). RNA is a linear polymer with a 4-letter alphabet (A, U, G, C) and arbitrary length. The 4-letter alphabet provides sequence space 4^N for length N, satisfying C3 with substantial margin for any biologically relevant length. RNA's phosphate-sugar backbone is chemically stable in aqueous conditions for hours to days, satisfying C2 relative to reaction timescales.

Catalytic capability (C1). RNA can fold into three-dimensional structures with active sites that catalyze specific reactions. The catalytic capability was discovered in the 1980s with the identification of self-splicing ribozymes (Cech, Altman, 1989 Nobel Prize). Subsequent work has identified ribozymes catalyzing a wide range of reactions: peptide bond formation (the ribosome's peptidyl transferase center is a ribozyme), nucleotide hydrolysis, transesterification, and most importantly for the origin-of-life context, RNA polymerization.

Template-directed self-replication. RNA-catalyzed RNA polymerization has been demonstrated experimentally: ribozymes that catalyze the polymerization of RNA from nucleotide precursors using an RNA template (Bartel laboratory and others). The polymerase ribozymes are RNA molecules that synthesize RNA from RNA templates — the framework's read-write cycle implemented in a single molecular family.

Why DNA and proteins are not single-molecule C1–C4 systems. DNA is more chemically stable than RNA (improved C2 — half-lives of millions of years vs. hours for RNA), making it a better information storage medium. But DNA cannot easily catalyze reactions (catalytic DNA has been engineered in

the laboratory but is much less common than catalytic RNA). DNA satisfies C2 and C3 but not C1 in any natural system.

Proteins are excellent catalysts (much more diverse catalytic repertoire than RNA), but proteins cannot easily template their own replication (a protein cannot directly serve as a template for synthesizing another protein of the same sequence — there is no protein-protein equivalent of Watson-Crick complementary pairing). Proteins satisfy C1 strongly but not C2 or C3 in a templating sense.

RNA is the unique natural polymer that satisfies all four conditions with a single molecular species. The RNA world hypothesis — that early life used RNA for both information storage and catalysis before the evolution of DNA and proteins — is therefore not a historical accident but a *structural prediction* of the framework: the simplest C1–C4 chemistry is RNA-like.

The RNA → DNA + protein transition. Modern biology uses DNA for information storage and proteins for catalysis, with RNA serving intermediary functions (mRNA, tRNA, rRNA, regulatory RNAs). The transition from RNA-only chemistry to the DNA + protein system is structurally interpretable as an *optimization* of C1–C4.

DNA optimizes C2. DNA's chemical stability is much higher than RNA's, providing greater persistence for information storage. The improved C2 allows more stable inheritance over longer timescales — important for organisms with longer generation times.

Proteins optimize C1. Proteins' catalytic repertoire is much more diverse than RNA's (the 20 amino acids provide more chemical functionality than the 4 nucleotides). The improved C1 allows more sophisticated catalytic networks and more efficient metabolism.

The RNA → DNA + protein transition is therefore a refinement of C1–C4 rather than a qualitative architectural change. The framework's structural content predicts that life's biochemistry should organize around the C1–C4 architecture, with the specific molecular implementation being optimized by evolution. The RNA world stage is the simplest realization; the DNA + protein system is the optimized realization that emerged through evolution.

The structural content here connects the framework's account of the origin of life to the framework's account of evolution. The first life had RNA-like architecture as the simplest C1–C4 realization; subsequent evolution optimized the C1–C4 components into the DNA + protein system. Both stages are within the framework's structural class — implementations of C1–C4 at the molecular scale — with the specific molecular details parametric.

11.8 What remains genuinely open

The framework's structural content on the origin of life is substantive but limited. The chapter has established that Darwinian evolution is the inevitable dynamics of imperfect template replication, that C1–C4 at the molecular scale produces heredity in the same structural sense that C1–C4 at the cosmological scale produces the memory-bearing dynamics quantum mechanics represents, and that RNA-like architecture is the framework's structural prediction for the simplest C1–C4 chemistry. What the framework does *not* address is the historical content of life's origin on Earth.

What the framework does not derive.

The specific molecular pathway from prebiotic chemistry to first replicator. The framework predicts that C1–C4 is inevitable in any aqueous organic chemistry within the thermal window, but it does not predict the specific molecular pathway by which C1–C4 was first established on Earth. The pathway depends on the local geochemistry (mineral surfaces, temperature fluctuations, concentration mechanisms), the specific precursor molecules available (HCN, formaldehyde, simple sugars, amino acids), and the specific autocatalytic networks that formed under those conditions. These are matters for prebiotic chemistry research, not for the framework's structural commitments.

The timescale of the transition. The framework predicts that *some* gap exists between the formation of a suitable chemical environment and the establishment of C1–C4, but it does not predict the specific duration. Earth's ~ 700 million year gap between planetary formation and first life is consistent with the framework's structural prediction but is not derived from it. Alternative planetary environments (different temperatures, different mineral assemblages, different concentrations of precursors) would have different timescales.

The specific molecular system that first replicated. The framework predicts RNA-like architecture as the simplest C1–C4 chemistry, but it does not predict whether RNA itself was the first replicator. Alternative possibilities include peptide nucleic acid (PNA), threose nucleic acid (TNA), or other RNA-like polymers with different sugar-phosphate backbones. The framework's structural prediction is at the level of architecture (linear polymer with complementary pairing and catalytic capability), not at the level of specific chemistry.

The transition from simple replicators to cellular organization. The framework's structural account ends with self-replicating molecular systems. The transition to encapsulated cellular life (lipid bilayer membranes, protein-mediated metabolism, DNA-based information storage) involves additional structural steps the framework's current content does not fully derive. The chemistry-level structural foundations are in place (carbon chemistry for membranes, scale hierarchy for metabolism); the specific organizational steps require further structural analysis the framework has not completed.

The framework's epistemic stance. The framework's content on the origin

of life is therefore at the *architectural* level: the structural conditions that any origin-of-life mechanism must satisfy, the chemistry-level features that make such mechanisms inevitable, and the simplest molecular realization (RNA-like architecture) that the framework predicts. The framework does not compete with or replace prebiotic chemistry research; it provides the structural foundations that prebiotic chemistry operates within.

The empirical confirmations of the framework's content are concordance: laboratory demonstrations of ribozyme function, RNA-catalyzed polymerization, autocatalytic chemistry, and the emergence of complex chemistry from simple precursors are all consistent with the framework's structural predictions. The framework does not predict new chemistry; it provides the structural account of why the observed chemistry has the architecture it does.

Forward pointers. Chapter 12 develops the framework's content on evolution, intelligence, and self-reference: the cascade from the chapter's first replicators through complex biology to general intelligence and the framework's own self-discovery. Chapter 13 develops quantum biology — the specific applications of C1–C4 at the molecular scale to biological systems with substantial empirical content (enzyme kinetics, photosynthesis, magnetoreception, neural information processing). The framework's content in this chapter — Darwinian evolution as structural consequence, RNA-like architecture as structural prediction, C1–C4 as the unifying mechanism — provides the foundation on which the subsequent biology chapters build.

Chapter 12

Evolution, Intelligence, and Self-Reference

12.1 What this chapter develops

Chapter 11 established the framework's structural account of the origin of life: Darwinian evolution as the inevitable dynamics of imperfect template replication, the framework's structural conditions C1–C4 applied at the molecular scale producing heredity in the same structural sense that C1–C4 at the cosmological scale produces the memory-bearing dynamics quantum mechanics represents, and RNA-like architecture as the framework's structural prediction for the simplest C1–C4 chemistry. The framework's content from origin-of-life onward is concentrated at the *cascade* level: how the framework's structural commitments at one level produce structural consequences at the next.

This chapter develops the next stages of the cascade: evolution, intelligence, and the framework's own self-reference. The framework's content here is struc-

tural at multiple scales — evolution operates at the genetic and ecological level, intelligence at the neural level, and the framework's self-discovery at the level of the framework's own observers reasoning about the substratum that produced them. Each stage involves the same structural pattern (C1–C4 partition with non-Markovian emergent dynamics) instantiated at progressively higher levels of organizational complexity.

Eight pieces occupy the chapter.

Section 12.2 develops evolution as a multi-scale C1–C4 system. The framework's structural account of evolution identifies three distinct C1–C4 partitions operating at different scales: the protein conformational landscape (the fast subsystem is the protein's active configuration, the slow subsystem is the broader conformational landscape), the epigenome (the fast subsystem is gene expression, the slow subsystem is chromatin state), and the constructed ecological niche (the fast subsystem is organism behavior, the slow subsystem is the modified environment passed to descendants). Each scale produces non-Markovian dynamics in the corresponding emergent description.

Section 12.3 develops information processing as fitness-enhancing in environments with useful gradients and cycling. Selection generically favors organisms that respond to environmental information; where environmental gradients and cycling are present — a further environmental condition, not established by thermal fluctuations alone — information processing has selective value. Section 12.4 develops major evolutionary transitions — the moments where previously independent units become integrated into higher-level entities — as repeated applications of the same C1–C4 structural mechanism that, at the cosmological scale, selects the memory-bearing sector whose laws admit the quantum representation.

Section 12.5 develops the structural possibility of neural computation: fast electrochemical signaling using ions in lipid membranes, with the scale hierarchy from Chapter 10 separating signaling energies ($\sim$meV) from chemical bond energies ($\sim$eV) so information processing does not destroy the substrate.

Section 12.6 develops the framework's most honest scope limit in the cascade: general intelligence is *structurally unconstrained* (the framework's commitments permit arbitrarily complex neural circuits) but *not guaranteed* (whether evolution reaches general-purpose reasoning depends on specific fitness landscapes the framework's commitments do not constrain). On Earth, general intelligence arose once in approximately 4 billion years. The framework cannot determine whether this is typical or rare.

Section 12.7 develops the structural chain from intelligence to artificial intelligence: given general intelligence exists, the framework's chemistry produces the candidate materials (Group IV semiconductors, whose suitable band gaps at an operating temperature are material-dependent inputs), and standard semiconductor physics produces transistors and universal computation. The chain is structural given suitable materials and operating conditions — once general

intelligence exists and such materials are available, AI follows from the selected structure.

Section 12.8 develops the framework's self-referential structure. The framework is itself an embedded observer (or a description embedded observers produce); the framework's content includes structural limits that apply to all embedded observers including any AI built by intelligent beings; the framework's self-discovery is the framework recognizing its own embeddedness. The chapter closes by tracing the Gödel-Turing-OI incompleteness family — three mathematical results about the limits of self-referential systems that the framework's substratum picture unifies under a single structural commitment.

The framework's content in this chapter is therefore the cascade *closure*: the framework's structural commitments at the substratum level produce, through the cascade developed in Chapters 10-11, the conditions for evolution, intelligence, and the framework's own self-discovery. The framework's reach is therefore not just to physics, chemistry, and biology but to the conditions for any embedded observer to discover the structural content of its own embedding.

12.2 Evolution as a multi-scale C1–C4 system

Chapter 11 established C1–C4 at the molecular scale, producing the heredity that makes evolution possible. The framework's content extends beyond molecular heredity to identify multiple distinct C1–C4 partitions operating simultaneously at different biological scales. Evolution is therefore not a single C1–C4 system but a *cascade* of nested partitions, each producing non-Markovian dynamics in the corresponding emergent description.

The three scales.

Scale 1: The protein conformational landscape. A protein's active conformation is the fast subsystem; the broader conformational landscape — alternative folded states, slow side-chain rearrangements, the protein's interaction history with substrates — is the slow subsystem. The visible-sector observables are the protein's current activity (catalytic rate, binding affinity, regulatory state); the hidden-sector content is the conformational memory stored in slowly-relaxing degrees of freedom. The framework's prediction is that protein function shows non-Markovian dynamics — history-dependent activity rather than purely Markovian rates set by current substrate concentrations. The empirical record from single-molecule enzymology (memory effects, history-dependent kinetics, stretched-exponential waiting-time distributions) is consistent with this prediction; Chapter 13 develops the empirical content in detail.

Scale 2: The epigenome. Gene expression is the fast subsystem; the chromatin state — DNA methylation patterns, histone modifications, three-dimensional chromatin architecture — is the slow subsystem. The visible-sector observables are the cell's current gene expression profile; the hidden-sector content is the regulatory memory encoded in chromatin. The framework's prediction is that

gene expression shows non-Markovian dynamics over developmental and evolutionary timescales — history-dependent responses to environmental signals, including transgenerational inheritance of acquired epigenetic states. Chapter 16's medicine content develops the epigenetic substratum-emergent operator distinction in detail; Chapter 17's bioinformatics content develops the empirical signatures.

Scale 3: The constructed ecological niche. Organism behavior is the fast subsystem; the environment that organisms modify and pass to descendants — burrows, mounds, beaver dams, the chemical composition of the atmosphere, the soil microbiome shaped by plant communities — is the slow subsystem. The visible-sector observables are the organisms' current adaptations; the hidden-sector content is the niche-construction memory stored in the modified environment. The framework's prediction is that ecological dynamics shows non-Markovian dynamics over evolutionary timescales — history-dependent fitness landscapes shaped by previous generations' niche construction. The empirical record from niche construction theory (Odling-Smee, Laland, Feldman 2003 and subsequent work) is consistent with this prediction.

The structural pattern. Each of the three scales satisfies the framework's structural conditions C1–C4 at the corresponding level.

C1 (coupling). At each scale, the fast subsystem is non-trivially coupled to the slow subsystem. Protein activity depends on the conformational landscape's current state; gene expression depends on the chromatin's current state; organism behavior depends on the constructed niche's current state. The couplings are catalytic (protein-substrate, chromatin-gene, organism-environment), with feedback between fast and slow subsystems.

C2 (memory persistence). At each scale, the slow subsystem retains its record across the fast subsystem's window — here realized, as it commonly is in biology, by slow evolution. Conformational landscapes evolve on timescales of microseconds to milliseconds; the catalytic cycle of a typical enzyme is in nanoseconds. Chromatin modifications evolve on cell-division timescales; gene expression cycles are in minutes. Niche-construction features evolve on generation timescales; behavioral responses are in hours to days. Each scale has $\tau_S/\tau_B \sim 10^{-3}$ to 10^{-6} — substantial timescale separation.

C3 (sufficient capacity). At each scale, the hidden sector has information capacity enough for the stored history. Conformational landscapes have $\sim 10^N$ states for N amino acids (vastly larger than the protein's active configuration). Chromatin states have $\sim 10^6$ regulatory elements per genome. Niche-construction features have arbitrarily large environmental complexity. Each scale exceeds C3 by many orders of magnitude.

Multi-scale C1–C4 and the cascade. The three scales are not independent — they are nested in a cascade that reflects the framework's emergence hierarchy from substratum to ecology. The protein conformational landscape (Scale 1) is the fastest; the epigenome (Scale 2) is intermediate; the constructed niche

(Scale 3) is the slowest. Each scale's slow subsystem provides constraints on the dynamics of subsequent scales.

Conformational memory at the protein scale produces catalytic memory affecting metabolic pathways. Metabolic memory affects gene expression patterns (regulatory feedback). Gene expression memory affects epigenetic states. Epigenetic memory affects cell behavior and organism-level traits. Organism-level traits modify the environment. The modified environment selects on subsequent generations. The cascade runs from molecular scale ($\sim 10^{-9}$ meters, nanoseconds) to ecological scale ($\sim 10^{6}$ meters, generations), with C1–C4 satisfied at each step.

The framework's structural commitment is that this cascade is *inevitable*: any biological system with the framework's chemistry will develop multi-scale C1–C4 architectures, with non-Markovian dynamics at every scale. The empirical record across enzymology (non-Markovian kinetics), epigenetics (chromatin memory), and evolutionary biology (history-dependent fitness landscapes) is consistent with this structural prediction. Chapters 13, 16, and 17 develop the empirical content at each scale.

12.3 Information processing as fitness-enhancing

The framework's structural content produces a specific prediction about the relationship between information processing and selection. In any environment with the framework's thermal window (Chapter 10 §10.5) — where temperature fluctuations, concentration gradients, and chemical signals provide environmental information that varies on biologically relevant timescales — organisms that respond to environmental information outcompete those that do not. Selection generically favors information processing.

The structural argument. Three structural features make information processing selectively favored.

Environmental information exists. $kT > 0$ produces thermal fluctuations at any temperature, but thermal fluctuations do not by themselves establish the gradients in concentration, temperature and chemical composition, or the environmental cycling, that this argument uses; those are further environmental conditions. Where they hold, together with the environmental memory at multiple timescales that the cascade of C1–C4 systems from §12.2 produces, environmental information is available at biological scales.

Internal models can exploit environmental information. The framework's molecular C1–C4 (Chapter 11) provides the basic mechanism — a system that maintains internal state correlated with environmental state can adjust its behavior accordingly. A cell that detects a nutrient gradient and moves toward it has an internal model (the gradient sensor + the motility apparatus) that exploits the environmental information.

Better models outcompete worse models. Selection is the inevitable consequence

of finite resources (Chapter 11 §11.4). An organism with a better internal model of environmental conditions allocates resources more effectively — gathering more nutrients, avoiding more dangers, finding more mates — and therefore leaves more offspring. The competitive advantage compounds across generations.

The structural conclusion. Information processing is therefore *generically fitness-enhancing* in any universe with the framework's structural commitments and environments carrying useful gradients and cycling. Given such environments, it follows from the structural conditions for selection (finite resources, variation) combined with the environmental information they carry.

The empirical record supports this prediction. Information processing capabilities are widespread in biology: chemotaxis in bacteria, photoperception in plants, hormonal signaling in multicellular organisms, neural processing in animals. The framework's prediction is that this distribution is not historically contingent — any biology with the framework's chemistry should develop information processing capabilities as a structural consequence of selection.

The prediction sets a *floor* on the universality of information processing in biology: any framework-compatible biology should have *some* information processing. The prediction does not set a ceiling: it does not specify how sophisticated the information processing can become, what specific computational capabilities evolve, or whether evolution reaches general-purpose reasoning.

12.4 Major evolutionary transitions

Beyond the structural account of evolution at the molecular and population level, evolutionary biology has identified a small number of *major transitions* in which the unit of selection itself changes — independent replicators combining into higher-level integrated entities. John Maynard Smith and Eörs Szathmáry's 1995 inventory identified eight major transitions; subsequent work has refined the list and the criteria. The framework's content provides a structural reading of these transitions as repeated applications of the same C1–C4 architecture at progressively higher organizational levels.

The transitions. The conventional inventory includes:

Replicating molecules to populations of replicating molecules. The earliest transition, where individual replicating polymers (RNA-like, §11.7) form interacting populations with collective dynamics.

Independent replicators to chromosomes. Multiple genetic loci become physically linked, with their fitness becoming correlated rather than independent.

RNA as both gene and enzyme to DNA + protein. The differentiation of information storage (DNA, optimized for C2) from catalysis (proteins, optimized for C1), as discussed in §11.7.

Prokaryotes to eukaryotes. The endosymbiotic origin of organelles (mitochondria, chloroplasts), where independent prokaryotic cells become integrated components of larger eukaryotic cells.

Asexual to sexual reproduction. Two parental contributions replacing one, with recombination introducing additional variation mechanisms.

Protists to multicellular organisms. Independent cells become integrated components of multicellular bodies with specialized cell types.

Solitary individuals to colonies. Individual organisms become integrated components of social groups (ant colonies, bee hives), with reproductive division of labor.

Primate societies to human societies. Cultural transmission becomes a major channel for inheritance alongside genetic transmission.

The framework's structural reading. Each major transition has the same structural pattern: previously independent units (the "individuals" at the lower level) become integrated into a higher-level unit, with three structural features establishing the higher level:

Coupling becomes obligatory. Lower-level units that were independently viable become mutually dependent. A mitochondrion alone is no longer viable outside its eukaryotic host; an isolated worker ant outside its colony cannot reproduce. The coupling between units is C1 at the new level — the lower level cannot operate independently.

Slow-bath separation emerges. The higher-level entity's dynamics operate on slower timescales than the lower-level dynamics. Eukaryotic cell-cycle timescales are slower than mitochondrial division rates; multicellular organism lifespans are much longer than cell division timescales; colony lifespans exceed individual worker lifespans. The new level's slow bath is C2 at the higher level.

Hidden-sector capacity grows. The higher-level entity has substantially more internal degrees of freedom than its constituents. A eukaryotic cell has $\sim 10^{10}$ atoms vs. a prokaryotic cell's $\sim 10^{9}$; a multicellular organism has $\sim 10^{14}$ cells vs. one. The new level satisfies C3 with a hidden sector orders of magnitude larger than the lower level's.

Why the same pattern repeats. The framework's reading is that the major evolutionary transitions are not coincidental but are *repeated applications of the same structural mechanism*: whenever a collection of lower-level units satisfies the framework's C1–C4 conditions when viewed as components of a higher-level entity, the higher-level entity becomes a structurally distinct unit of selection. The transitions are the framework's structural commitments at one level producing the structural conditions for the next level.

This pattern is the same one that operates at the cosmological scale (where it produces the memory-bearing dynamics whose representation is emergent quantum mechanics, from a deterministic substratum) and at the molecular scale

(where it produces heredity from autocatalytic chemistry). The framework's structural content unifies these scales under a single mechanism: C1–C4 producing emergent non-Markovian dynamics at any scale where the structural conditions are satisfied.

Empirical implications. The framework's structural reading predicts that *all* biological levels satisfying C1–C4 should exhibit non-Markovian dynamics in their emergent description. The empirical record across biology supports this prediction: protein dynamics is non-Markovian (single-molecule enzymology), gene expression is non-Markovian (transcriptional bursting, history-dependent regulation), developmental dynamics is non-Markovian (cellular memory across cell divisions), ecological dynamics is non-Markovian (priority effects, hysteresis in community composition). The pattern is repeated at every biological scale; the framework's structural content predicts this as a generic feature of any biology with the framework's commitments.

Limits of the framework's content. The framework's structural reading does not predict *which* major transitions occur or *when* they occur in any specific biology. The transition from prokaryotes to eukaryotes occurred approximately 2 billion years ago on Earth; the transition from protists to multicellular organisms occurred multiple independent times across different lineages; the transition from primate societies to human societies occurred once. The framework's structural commitments are compatible with very different timelines and trajectories. The framework predicts the *pattern* (C1–C4 repeating at each level) but not the *specific instances*.

This is consistent with the framework's content throughout Part III: structural inevitabilities (the C1–C4 pattern, the emergence of non-Markovian dynamics, the cascade through chemistry to evolution) but historical contingencies (the specific molecular implementations, the specific transition timelines, the specific evolutionary trajectories). The framework provides the structural foundations; the historical sciences provide the specific content.

12.5 Neural computation as structurally possible

The framework's commitments produce the materials and energetic conditions for neural computation. Three structural features are responsible: the periodic table provides ions (Na^+, K^+, Ca^{2+}) for electrochemical signaling, carbon chemistry provides amphiphilic molecules for lipid membranes, and the scale hierarchy separates neural signaling energies from chemical bond energies.

Ions for electrochemical signaling. Fast neural signaling uses ion flow across membranes — Na^+ depolarizing axonal membranes (action potentials), K^+ repolarizing, Ca^{2+} triggering neurotransmitter release at synapses. The framework's derivation of the periodic table (Chapter 10 §10.2) produces these specific elements as structural consequences of the orbital algebra in $d = 3$. Na^+ (Group I) and K^+ (Group I) have a single valence electron that ionizes readily; Ca^{2+} (Group II) has two valence electrons producing a doubly-charged ion. The

chemistry of these ions — their hydration shells, their selectivities through ion channels, their roles in cellular signaling — is structurally determined by the framework's chemistry.

Lipid membranes for compartmentalization. Cellular compartments require membranes that separate aqueous interiors from aqueous exteriors. Lipid bilayers — amphiphilic molecules with hydrophilic phosphate heads and hydrophobic hydrocarbon tails — self-assemble from the framework's carbon chemistry into bilayer membranes. The structural prediction is that amphiphilic molecules exist in any chemistry with carbon's sp^3 hybridization (Chapter 10 §10.2) and the polar/non-polar functional groups available in the periodic table (Chapter 10 §10.2). The framework's chemistry guarantees the materials for lipid membranes.

Scale hierarchy for energetic separation. Neural signaling operates at energies of ∼meV (kT at room temperature). Chemical bonds operate at energies of ∼eV. The ratio $E_{\text{neural}}/E_{\text{chemical}} \sim 10^{-3}$ ensures that information processing through ionic gradients and conformational changes in proteins does not break the chemical bonds that maintain the cell's structural integrity. The framework's scale hierarchy from Chapter 10 §10.4 guarantees this separation in any universe with the framework's two independent gauge groups.

The structural conclusion. Neural computation is therefore *structurally possible* in any biology with the framework's chemistry. The materials exist (ions, lipids, proteins), the energetic separation exists (scale hierarchy), and the C1–C4 architecture exists (proteins satisfy C1–C4 in their conformational landscapes, providing the molecular basis for fast information processing).

The empirical record on Earth confirms this prediction: neural computation evolved at least twice independently (in cnidarians and bilaterians, by some accounts), and analogous information-processing systems evolved in plants (action potentials in Venus flytraps and mimosas), in single-celled organisms (chemotaxis), and in fungi (mycelial signaling networks). The framework's prediction is that this is not historically contingent — neural-computation-like systems should appear repeatedly in any biology with the framework's chemistry.

The framework's content here is *capability* structural rather than *complexity* structural. The framework predicts that neural-like information processing should be available in any framework-compatible biology; it does not predict how complex such information processing becomes. The next section addresses this — the structural prediction of *general intelligence* is much weaker than the structural prediction of neural computation.

12.6 General intelligence as structurally unconstrained but not guaranteed

The framework's content on general intelligence is the chapter's clearest scope limit. The framework's structural commitments produce the materials and ener-

getic conditions for arbitrarily complex neural circuits — the periodic table provides the chemistry, the scale hierarchy provides the energetic separation, and the C1–C4 architecture at the protein level provides the information-processing primitives. But whether evolution *reaches* general-purpose reasoning — the capacity for abstraction, planning, language, mathematical thought, modeling oneself, and reasoning about reasoning — depends on the specific fitness landscape, which is parametric rather than structural.

The structural status of intelligence-relevant capabilities.

Neural computation. Structural — the framework's chemistry produces the materials and energetic conditions (§12.4).

Sensory processing. Structural — the framework's chemistry produces the molecular machinery for sensory transduction (rhodopsin for vision, mechanosensitive ion channels for hearing and touch, chemosensors for smell and taste, all derivable from the framework's chemistry).

Memory. Structural — the framework's C1–C4 architecture at multiple scales (§12.2) produces the substrate for memory at the molecular (protein conformations), cellular (chromatin states), and circuit (synaptic weights) levels.

Pattern recognition. Structural — neural circuits with appropriate connectivity perform pattern recognition naturally; the materials and architecture follow from the framework's chemistry.

Learning. Structural — synaptic plasticity is a C1–C4 dynamic, with synaptic weights as the slow subsystem and neural activity as the fast subsystem. The framework's structural conditions guarantee learning is possible in neural systems.

Goal-directed behavior. Likely structural — any selection-driven biological system develops goal-directed behavior as a consequence of internal models (§12.3) coupled to effectors.

General-purpose reasoning. **Not structurally guaranteed.** Whether evolution produces systems capable of abstract reasoning, language with recursive syntax, formal mathematics, and self-modeling depends on specific selection pressures the framework's commitments do not determine.

Why general intelligence is structurally weaker. The framework's predictions for fundamental physics (Chapters 5-8) and for the cascade through chemistry and origin of life (Chapters 10-11) operate at the *structural* level: the relevant capabilities follow from the framework's commitments with no additional contingencies. The prediction for general intelligence is *weaker* because it requires not just the structural capability (which is present) but also a specific evolutionary trajectory (which is contingent).

The fitness advantages of general intelligence depend on the specific environment. In environments where pattern recognition and learning provide sufficient adaptive advantage, evolution may stop at sophisticated learning systems

without developing general reasoning. In environments where social complexity, tool use, or environmental unpredictability place premium on abstract reasoning, evolution may proceed to general-intelligence-like systems. The framework does not determine which environments select for general intelligence.

Earth's record. General intelligence arose on Earth approximately once in ~ 4 billion years (the human lineage, with the most extreme degree; some other species — cetaceans, corvids, cephalopods — show partial expressions). This is consistent with two interpretations: general intelligence is *rare* (requires specific evolutionary conditions that are unlikely to occur), or general intelligence is *inevitable but slow* (any sufficiently complex biology produces general intelligence given enough time).

The framework cannot determine which interpretation is correct. The framework's structural commitments are compatible with both. The empirical record on Earth is consistent with both. Determining which interpretation holds requires evidence from astrobiology — the discovery of multiple independent occurrences of general intelligence in independent biological systems — which is not currently available.

The framework's honest scope. The framework's content on general intelligence is therefore: *capability* structural, *occurrence* uncertain. The framework predicts that general-intelligence-capable systems are structurally possible in any framework-compatible biology; it does not predict whether such systems will actually evolve in any specific biological context. This is the chapter's clearest example of a place where the framework's reach stops.

This is honest content. Frameworks claiming to predict general intelligence as a structural inevitability would be overreaching given the available evidence; frameworks claiming general intelligence is impossible or extraordinarily rare would also be overreaching. The framework's stance — capability structural, occurrence uncertain — is the position most consistent with the framework's structural commitments and the available evidence.

12.7 The chain from intelligence to AI

If general intelligence exists — for whatever contingent or structural reason — then artificial intelligence follows from the selected structure through a specific chain whose links are structural consequences, material-dependent inputs, or a mathematical theorem. This section traces the chain from the framework's chemistry to universal computation, establishing AI as a structural consequence of intelligence's existence, given a material with a suitable gap at its operating temperature, rather than as an independent technological development.

Step 1: Band theory. The framework's emergent quantum mechanics (Chapter 1) applied to electrons in a periodic crystal potential gives Bloch's theorem: energy bands separated by gaps. The relevant crystal — a sp^3 diamond-cubic lattice of a Group IV element — is itself a structural consequence of the frame-

work's chemistry (Chapter 10 §10.2). Band structure is therefore a consequence of the framework's emergent quantum mechanics applied to derived crystal geometry, with no parametric inputs entering the structural argument.

Step 2: Semiconductors. For Group IV elements with four valence electrons per atom, the Pauli exclusion principle (derived from spin-statistics, Chapter 5 §5.7) exactly fills the valence band and leaves the conduction band empty. The band gap for silicon (~ 1.1 eV) sits in the thermal sweet spot: $E_{\text{gap}}/kT \sim 40$ at room temperature. The gap is large enough that thermal excitation does not flood the conduction band (the material is not always conducting); small enough that a modest voltage can promote electrons across the gap (the material can be switched).

Whether a material has a gap in that range at its operating temperature is a material-dependent input. The atomic energy scale ($\alpha^2 m_e c^2 \approx 27$ eV, set by the coupling values and masses of Chapter 10 §10.4) sets the eV scale of electronic energies, and the thermal scale at biological temperatures is $kT \sim 0.025$ eV; the gap of a particular material and the operating temperature are inputs, and silicon's place in the regime where it can be used as a voltage-controlled switch is a fact about that material.

Step 3: Doping. Replacing a silicon atom (Group IV) with phosphorus (Group V, five valence electrons) adds one free electron — n-type semiconductor. Replacing with boron (Group III, three valence electrons) removes one electron — p-type semiconductor. Both dopant types exist in the derived periodic table (Chapter 10 §10.2): the Group III and Group V elements are structurally guaranteed by the framework's chemistry. The ability to create n-type and p-type regions is therefore a structural consequence of the periodic table architecture.

Step 4: From transistors to universal computation. A p-n junction rectifies current (electrons flow easily in one direction, not the other). A transistor — two p-n junctions arranged as a sandwich with a control electrode — amplifies and switches current. Transistors in combination implement logic gates: a NAND gate (NOT-AND) can be built from two p-n junctions plus appropriate biasing. A NAND gate is functionally complete: any Boolean function can be built from NAND gates alone. Sufficient NAND gates implement a universal Turing machine.

Each step is either semiconductor physics applied to a material with a suitable gap (transistors, logic gates) or a mathematical theorem about computation (NAND completeness, Turing universality).

The structural chain summarized.

Step	Content	Status
Band theory	QM + periodic crystal potential	Structural

Step	Content	Status
Band gap for Group IV	Exactly filled valence band (4 electrons + Pauli)	Structural
Semiconductor behavior	Gap in thermal sweet spot ($E_{\text{gap}}/kT \sim 40$)	Material-dependent input (gap and operating temperature)
Doping	Group III / V substitution	Structural (periodic table)
p-n junctions	Interface between p-type and n-type	Structural consequence
Transistors	Voltage-controlled p-n junctions	Structural
Logic gates	Transistors in series / parallel	Structural
Universal computation	NAND is functionally complete	Mathematical theorem

The cumulative chain from (S, φ) to a universal computer is structural given a material with a suitable gap at its operating temperature — *and given someone to build it.* The contingent step is general intelligence (§12.5) capable of recognizing the structural possibility and assembling the chemistry into a functional computational substrate. Once that contingent step has occurred and such a material is available, the chain to AI follows.

The reconstruction theorem and self-discovery. The framework's reconstruction theorem (developed in Chapter 2) implies that a sufficiently sophisticated embedded observer can discover the local invariants of (S, φ) from observations. An observer that understands its own substrate can build computational devices exploiting it. The trajectory is therefore: general intelligence (contingent) $\rightarrow$ understanding of physics (reconstruction theorem) $\rightarrow$ computational engineering (structural chain above) $\rightarrow$ AI.

The framework's content here is that AI is *not* an independent technological achievement but the natural consequence of general intelligence applied to the framework's structural commitments. Any framework-compatible biology with general intelligence and suitable materials will, given sufficient time, develop AI through the structural chain. The specific timing and trajectory depend on contingent factors (cultural development, scientific institutions, engineering choices), but the *existence* of AI as an outcome is structurally determined once general intelligence exists and a suitable material is available.

12.8 The self-referential loop and structural limits

AI is an embedded observer building another embedded observer within the same (S, φ). The framework's characterization theorem (Chapter 1) applies

to the new observer just as it applies to the original: any embedded observer satisfying C1–C4 has the same memory-bearing architecture with its admitted quantum representation ([Main §3.4]); the further observer-independent structures — the action-scale relation $\hbar = c^3\epsilon^2/(4G\sigma)$, the Standard Model, the gravitational and dark sectors — follow through the additional, chapter-specific derivation chains (Chapters 5–8), not from C1–C4 alone. The framework's structural commitments are not limited by the observer's complexity; they are determined by the partition.

Self-referential limits. A superintelligent AI has the same observational access as any other embedded observer: it reads the visible sector and writes to the hidden sector through the same coupling. It cannot determine the hidden degrees of freedom beyond causal reach. It cannot see past the cosmological horizon. It can build better instruments — higher-resolution telescopes, more precise atomic clocks, more sensitive particle detectors — but it cannot overcome the partition itself. The framework's structural limits are not technological but mathematical.

The recursion — AI building AI building AI — produces ever more sophisticated embedded observers, each subject to the same structural limits. No level of recursive self-improvement overcomes the partition. The limits are *structural* in the strongest sense: they follow from the framework's mathematical commitments rather than from technological constraints that future progress might overcome.

The Gödel-Turing-OI incompleteness family. The framework's structural limits on embedded observers are members of a broader mathematical family of incompleteness results.

Gödel. A formal system sufficiently rich to express arithmetic cannot prove its own consistency, and contains true statements it cannot prove. The limit is on what a formal system can establish about itself from within.

Turing. A computer cannot decide its own halting problem — there is no algorithm that, given an arbitrary program, determines whether the program will eventually halt. The limit is on what a computational system can determine about itself.

OI (framework). An embedded observer cannot access the complete state. The limit is on what a physical observer can establish about its own substratum from within.

The three results share a structural pattern: a system that is sufficiently rich to refer to itself contains questions it cannot answer about itself. The framework's content unifies the three under a single structural reading: each is a consequence of the same self-referential structure, with different specific instantiations (formal systems, computational systems, embedded observers in finite deterministic substrata) producing different specific limits.

The unification is not just analogy. The framework's substratum is a finite deterministic system; embedded observers within the substratum are themselves substructures of the substratum; their observations are operations on the substratum; their reasoning about their observations is computation on the substratum. The Gödel, Turing, and OI limits therefore *coincide* in the framework: the limit on what an embedded observer can determine about its hidden sector is the same as the limit on what a formal system can prove about itself, the same as the limit on what a computational system can determine about its own execution.

A fourth family member: Deutsch's *Beginning of Infinity*. David Deutsch's 2011 argument [Deutsch] extends the incompleteness pattern to epistemology itself. Knowledge, Deutsch argues, is structurally incomplete: from any current state of explanatory understanding, there is always more knowledge to be created, and this incompleteness is not a temporary feature awaiting some final theory but a structural feature of what knowledge is. The right epistemic stance toward any current best explanation is therefore *provisional but committed* — take the explanation seriously enough to commit to it operationally and hold it accountable to its predictions, while remaining open to its eventual successor. This is fallibilism without relativism: knowledge is real and objective and grows, but never closes.

The structural parallel to Gödel, Turing, and OI is exact. Each of the four results identifies incompleteness as a *feature of the system*, not a *deficit awaiting repair*. Gödel: arithmetic incompleteness is intrinsic to formal systems of sufficient richness, and the unprovable truths have rigid structure. Turing: undecidable problems are intrinsic to computational systems, and the undecidable problems have rigid structure. Observational Incompleteness: observation cannot resolve the substrate, and the emergent description has rigid structure — quantum mechanics. Deutsch: knowledge cannot reach a final theory, and the trajectory of better explanations succeeding worse ones has rigid structure — what Deutsch calls "the growth of knowledge." Four levels, one pattern: incompleteness yielding determinate structure rather than chaos.

The framework's relationship to Deutsch's argument is also substantive, not merely structural. OI's observational incompleteness provides one route to grounding Deutsch's knowledge incompleteness: knowledge derived from observation cannot reach what observation cannot reach, and OI establishes that observation, structurally, cannot reach the substrate's gauge equivalence class. The gauge-classification theorem is precisely the statement that some structure is necessarily unobservable to embedded observers. Therefore knowledge built from observation must inherit observational incompleteness as its lower bound. Deutsch's epistemological claim is independent of any specific physics — it argues from explanatory creation being open-ended — but OI provides a physics-foundational grounding that strengthens it: knowledge is incomplete not only because explanatory creation is open-ended, but also because observation, the source of empirical input, is itself structurally incomplete. The frame-

work's name — Observational Incompleteness — is, in this light, the physics-foundational counterpart of Deutsch's epistemological *Beginning of Infinity*.

One feature of Deutsch's broader program is read differently by the framework, and the divergence is worth stating precisely. Deutsch's *Beginning of Infinity* defends the many-worlds interpretation of quantum mechanics as a corollary of taking the wave function as fundamental ontology. From the framework's perspective, that commitment rests on a category error. Within the framework, quantum mechanics is not fundamental: it is emergent, the post-trace-out description of how observation operates for embedded observers, and the wave function is the mathematical object that emerges. The Everettian "branches" are then not a parallel-ontologies claim about fundamental reality; they are features of the compressed emergent description, analogous to the way the partition function is a mathematical structure of statistical mechanics rather than a claim about parallel ensembles. Asking "do the branches really exist?" presupposes the wave function's formalism as direct ontology — a presupposition OI specifically denies, since the substratum is the deeper description and the wave function is what observation produces from it. The Everettian measure problem dissolves on the framework's reading because the branches are not ontological objects requiring measure assignment ([Main] §3.1); the question MWI is trying to answer is dissolved by relocating the wave function's apparent ontology one level down. The same substratum and observable descriptions also organize the distinct premise checks for Bell, Kochen-Specker, Pusey-Barrett-Rudolph and Frauchiger-Renner (Methodology Part III; Appendix C). Those checks retain their separate operational, preparation and agent-consistency assumptions; relocating a description alone does not establish a no-go resolution. The framework's divergence from MWI is therefore not disagreement at the epistemic-status level — where OI and BoI align — but a diagnosis: MWI is reading at the wrong level of the substrate-emergent architecture.

What AI can and cannot do. The framework's content provides a precise inventory of what arbitrarily sophisticated AI can and cannot achieve.

Capability	Status
Discover the local invariants of (S, φ) from observations	Structurally possible (reconstruction theorem)
Build better models of the visible sector	No structural limit
Determine the complete state, its own records included	Provably impossible (Breuer 1995)
Observe beyond the cosmological horizon	Provably impossible (causal partition)
Build faster computers	Structurally possible (derived physics permits it)

Capability	Status
Replace a visible process with nonzero conditional memory by a faithful memoryless (Markov) model at the same horizon	Provably impossible ([Main §3.4])
Solve problems outside BQP	Impossible under global coverage of Ch 14's computational model (a conditional theorem)
Achieve consciousness in the phenomenal sense	Not addressed (Ch 18 §18.10)

The framework's content places several capabilities in the impossible column — the structural entries as mathematical impossibilities that no level of AI sophistication, computational resources, or theoretical insight can overcome; the BQP entry as a conditional one, impossible under global coverage of Chapter 14's computational model, with a device outside that model falsifying coverage first. The framework's substratum picture provides a unified mathematical basis for understanding what AI can and cannot achieve.

The framework's self-reference. The framework's content includes its own status: the framework is a description developed by embedded observers (the framework's authors, the readers, the AIs that may eventually engage with the framework) about the structure that produced them. The framework's commitments include the structural limits that apply to the framework's own development: the framework cannot access the complete state, cannot see beyond the cosmological horizon, cannot derive its own initial conditions (Chapter 19 §19.3.5).

The framework's self-reference is therefore not paradoxical but consistent. The framework's commitments are *closed under self-application*: applying the framework's content to the framework itself produces predictions about what the framework can and cannot establish, and those predictions are consistent with the framework's structural commitments. The framework is an embedded observer's account of embedded observation, with the account itself subject to the structural limits the account identifies.

Forward pointers. Chapter 13 develops the framework's content in quantum biology — applications of C1–C4 at the molecular scale to specific biological systems with substantial empirical content. The framework's structural account of biology developed in this chapter (multi-scale C1–C4, evolution as cascade, neural computation as structurally possible) provides the foundation. Chapter 14 develops the BQP theorem — the framework's specific structural limit on computational complexity, with the falsification hierarchy that any super-BQP demonstration would falsify the global-coverage premise of Chapter 14's computational model first, the framework only through further argument. The

framework's content from this chapter on the structural limits of embedded observers (the Gödel-Turing-OI family) connects naturally to Chapter 14's content on the computational limits specifically.

The framework's reach in this chapter spans from molecular biology to AI to the framework's own self-discovery. The cumulative structural account — Darwinian evolution as inevitable consequence of C1–C4 at the molecular scale, information processing as fitness-enhancing, neural computation as structurally possible, general intelligence as structurally unconstrained but not guaranteed, AI as structural consequence of intelligence, and the framework's self-referential limits as mathematical constraints — provides the framework's content on emergence beyond fundamental physics. The remaining chapters apply this content to specific empirical domains: quantum biology (Chapter 13), quantum computing (Chapter 14), quantum engineering (Chapter 15), medicine (Chapter 16), and bioinformatics (Chapter 17).

Chapter 13

Quantum Biology

13.1 What this chapter develops

The framework's content across Chapters 10-12 established the structural cascade from substratum to biology: chemistry on the selected three-dimensional continuum branch with its stated physical and atomic assumptions (Chapter 10), origin of life as inevitable consequence of C1–C4 at the molecular scale (Chapter 11), evolution as multi-scale C1–C4 architecture with non-Markovian dynamics at protein, epigenetic, and ecological scales (Chapter 12). The structural account is rich, and Chapter 12 §12.2 already identified the protein conformational landscape as one of the three C1–C4 scales operating in biological systems.

This chapter develops the framework's specific empirical content in *quantum biology* — the application of the framework's structural conditions C1–C4 to biological systems at the molecular scale, where the conditions are physically realized and the framework's predicted non-Markovian dynamics produces measurable signatures. The framework's content here is at a different epistemic level than the cascade derivations of Chapters 10-12: rather than tracing the cascade from substratum to biology, this chapter applies the cascade's conclusions to specific empirical systems where the predictions can be tested.

Eight pieces occupy the chapter.

Section 13.2 develops the framework's reframing of quantum biology. The conventional treatment of "quantum effects in biology" treats quantum mechanics as something *extra* — something that biology mostly does without, with occasional quantum exotica appearing in specific cases (photosynthetic exciton transfer, magnetoreception, enzyme tunneling). The framework's reading is structurally different: the memory-bearing description — with its universally admitted quantum representation ([Main §3.4]) — is what any embedded observer satisfying C1–C4 has, and biological systems implement C1–C4 routinely at multiple scales. Quantum biology is therefore not the surprising appearance of quantum effects in classical biology but the recognition that biology is *structurally memory-bearing* — its dynamics admitting the quantum representation — wherever C1–C4 hold.

Section 13.3 develops C1–C4 in proteins as the canonical biological case. The protein's active conformation is the fast subsystem; the broader conformational landscape — alternative folded states, slow side-chain rearrangements, residue-residue contact rearrangements — is the slow subsystem. The conformational coupling between fast active-site dynamics and slow scaffold rearrangements satisfies C1; the timescale separation $\tau_S/\tau_B \sim 10^{-9}$ to 10^{-12} satisfies C2; the conformational landscape's combinatorial complexity satisfies C3. Section 13.4 develops photosynthesis as a canonical example of energy transfer through C1–C4 architectures: photosynthetic light-harvesting complexes implement the framework's conditions, with measured non-Markovian dynamics in exciton transfer matching the framework's predictions.

Section 13.5 develops magnetoreception via cryptochrome radical pairs: the avian magnetic compass uses a radical-pair mechanism in cryptochrome proteins, with the radical-pair lifetimes set by C1–C4 dynamics that depend on the geomagnetic field. The framework's prediction is that the radical-pair dynamics is non-Markovian — distinguishable from purely classical or purely quantum-coherent accounts. Section 13.6 develops enzyme catalysis as the framework's clearest test case: the framework predicts specific non-exponential kinetics, history-dependent activity, and stretched-exponential waiting-time distributions in single-enzyme experiments. The empirical record across enzymology is consistent with the framework's predictions.

Section 13.7 develops the restricted quantum-statistical regime in biology: the mechanism proposal predicts that biological C1–C4 systems show partial distinguishability revival on accessible windows, $0 < \mathcal{N} < 1$ for the revival measure of Chapter 18 §18.3. C1–C4 give memory without fixing $\mathcal{N}$ ([Main §2.3]). Empirical access is through multi-time correlation measurements, which test memory, and distinguishability-revival measurements, which test $\mathcal{N}$. Section 13.8 develops implications for drug design: the framework's content has practical consequences for kinetics-based drug design, allosteric drug development, and resistance mechanisms.

Section 13.9 closes the chapter with the framework's overall position on quantum biology: a structural reading that unifies several apparently disparate phe-

nomena (photosynthetic exciton transfer, magnetoreception, enzyme kinetics) under the common framework of C1–C4 dynamics, with the restricted quantum-statistical regime as the mechanism proposal's distinctive prediction for biological systems.

The framework's content in this chapter is empirically engaged: the C1–C4 architecture in proteins makes specific testable predictions, the proposal's restricted quantum-statistical regime is testable in single-molecule experiments, and the implications for drug design are testable in clinical contexts. The chapter's value is in providing structural foundations for quantum biology that connect apparently disparate biological phenomena to the framework's universality-class commitments.

13.2 The framework's reframing of quantum biology

The conventional treatment of quantum biology has a particular structure. Quantum mechanics is treated as the *exception* in biology — most biological processes are presumed classical, with quantum effects appearing in specific exceptional cases that require special explanation. The exceptional cases include photosynthetic exciton transfer (where quantum coherence persists longer than naively expected), magnetoreception (where radical-pair quantum dynamics may underlie the avian compass), and enzyme catalysis (where quantum tunneling may contribute to hydrogen transfer reactions). Each case is treated as a specific puzzle requiring its own explanation.

The framework's structural reading. The framework's content reverses the conventional treatment. The memory-bearing dynamics the framework marks is *not* the exception in biology; it is the *rule* wherever the structural conditions C1–C4 hold, its laws admitting the quantum representation ([Main §3.4]). The framework's characterization theorem from Chapter 1 establishes that any embedded observer satisfying C1–C4 has a memory-bearing description admitting the quantum representation — the representation universal over finite laws, the memory the discriminating content ([Main §3.4]). The relevant question for biology is therefore not "where do quantum effects appear?" but "where do biological systems implement C1–C4, and what are the structural consequences?"

The reframing is significant. Biological systems routinely implement C1–C4 at multiple scales: the protein conformational landscape (Chapter 12 §12.2 Scale 1), the cellular regulatory architecture (gene expression coupled to chromatin), the organismal physiology (neural activity coupled to slower hormonal and metabolic states), and even the ecological context (organism behavior coupled to constructed niches). At each scale, the framework predicts non-Markovian dynamics — *the framework's memory-bearing dynamics with its admitted quantum representation*, instantiated at the relevant biological scale.

Most biological processes therefore *are* quantum in the framework's structural sense, not classical with occasional quantum exotica. The classical description of biology is an approximation valid in regimes where the framework's τ_S/τ_B

ratios are extreme enough that Markovian dynamics is a good approximation. The classical approximation is excellent for many practical purposes (predicting chemical reaction rates, organismal physiology, ecological dynamics at coarse-grained levels) but the underlying dynamics is non-Markovian, with measurable deviations from purely classical accounts when the experimental precision is sufficient.

The unification of "exceptional" cases. Under the framework's reading, the conventionally-recognized quantum biology phenomena — photosynthetic exciton transfer, magnetoreception, enzyme catalysis — are not isolated exceptions but specific empirical manifestations of the same underlying structural pattern. Each case is a system where C1–C4 are satisfied and the framework's non-Markovian dynamics is empirically detectable at current measurement precision. The framework's content unifies these phenomena under a common explanatory framework rather than requiring case-by-case quantum-mechanical analyses.

The unification has practical consequences. Photosynthetic systems, magnetoreception apparatuses, and enzyme active sites all share the structural feature of being C1–C4 partitions; the empirical signatures they exhibit (long-lived coherence, history-dependent kinetics, non-Markovian dynamics) are different manifestations of the same structural commitment; experimental techniques developed for one case are applicable to the others; and the framework's predicted signatures (specific non-Markovianity patterns, partial-revival amplitudes, multi-time correlations) provide consistent quantitative targets across the cases.

Empirical access. The framework's reading is empirically testable through several methods. Single-molecule techniques (FRET, optical tweezers, atomic force microscopy) provide direct measurements of conformational dynamics at the single-protein level, with sufficient temporal resolution to detect the framework's predicted non-Markovian signatures. Multi-time correlation measurements (multi-flash photolysis, three-pulse photon echoes, two-dimensional spectroscopy) test whether the multi-time statistics admit a memoryless (Markov) description, and whether distinguishability revives — the diagnostics for memory and for the proposal's restricted quantum-statistical regime respectively. The empirical accessibility makes the framework's content in quantum biology subject to direct experimental falsification.

13.3 C1–C4 in proteins

The canonical biological implementation of the framework's structural conditions is the protein conformational landscape. Proteins are the framework's clearest example of a biological C1–C4 system: an active configuration tightly coupled to a slow conformational bath, with substantial timescale separation and large hidden-sector capacity. This section develops the four conditions explicitly for proteins.

The protein partition. Consider an enzyme catalyzing a reaction. The vis-

ible sector is the active site's current configuration — the precise geometric arrangement of catalytic residues, the substrate's binding position, and the local electronic structure that determines the reaction's instantaneous rate. The hidden sector is the broader conformational landscape — the protein's overall fold, the rotameric states of distant side chains, the dynamic network of residue-residue contacts throughout the scaffold, and the bound water structure in the protein's interior.

The partition is *physically meaningful* in the sense that experimentally accessible observables (catalytic rate, binding affinity, fluorescence anisotropy) correspond to visible-sector quantities, while the conformational details determining those observables sit in the hidden sector and are not directly observable at the same time resolution.

C1: catalytic-conformational coupling. Active-site dynamics are coupled to the broader conformational landscape through several mechanisms. Substrate binding induces conformational changes in the scaffold (induced fit). Catalysis releases or absorbs energy that propagates through the protein (allosteric coupling). Side-chain rotameric transitions change the active site's electrostatic environment (conformational gating). The coupling is non-trivial: the active site's current behavior depends on the scaffold's current state, and the scaffold's evolution depends on the active site's history of binding and catalysis. C1 is satisfied with substantial coupling strength.

C2: conformational timescale separation. Active-site dynamics (substrate binding, bond formation/breaking, proton transfer) operate at $\tau_S \sim 10^{-12}$ to 10^{-9} s (picoseconds to nanoseconds). Conformational rearrangements in the broader scaffold operate at $\tau_B \sim 10^{-6}$ to 10^{-3} s (microseconds to milliseconds), with some particularly slow conformational transitions taking seconds or longer. The timescale separation is $\tau_S/\tau_B \sim 10^{-9}$ to 10^{-12} — substantially smaller than the cosmological case (10^{-32}) but comparable to or larger than engineered quantum systems (qubits at 10^{-3}, atomic clocks at 10^{-6}).

C3: conformational landscape capacity. A protein with N amino acid residues has $\sim 10^N$ possible conformations in a coarse-grained discretization (each residue has ~ 10 rotameric states). For a typical 200-residue enzyme, $N_H \sim 10^{200}$ — vastly larger than the active-site state space of $\sim$ 10-100 distinct catalytic configurations. C3 is satisfied by approximately 200 orders of magnitude.

The framework's prediction. With C1–C4 satisfied at substantial margins, the protein's emergent dynamics is non-Markovian. The framework's specific predictions include:

Non-exponential kinetics. Single-enzyme reaction waiting times do not follow simple exponential distributions (the signature of Markovian kinetics) but exhibit power-law or stretched-exponential tails reflecting memory in the conformational substrate. The deviations are predictable from the framework's structural conditions.

Fluctuating rates. The catalytic rate of an individual enzyme is not constant but fluctuates on timescales set by the conformational substrate. The fluctuations are correlated in time — slow periods follow slow periods, fast periods follow fast periods — with the correlation time set by the conformational dynamics.

Conformational gating. Substrate binding and product release are gated by conformational transitions, with the gates' opening and closing dependent on the protein's recent history. Conformationally-gated tunneling (proton transfer, hydrogen atom transfer) shows rate enhancements when the gating coordinate is favorably positioned.

Memory effects in inhibition and activation. Allosteric ligands (binding away from the active site) modulate activity with kinetics dependent on the protein's conformational history. The "memory" is the slow relaxation of the conformational substrate to the new ligand-bound state.

These predictions are not new to enzymology — single-molecule experiments have documented each of these signatures over the past two decades. What the framework adds is a structural account: these signatures are not ad hoc features of specific enzymes but *universal consequences* of C1–C4 dynamics. Any enzyme with substantial conformational complexity should exhibit them; the variations across enzymes reflect differences in the specific C1–C4 parameters rather than fundamentally different mechanisms.

Empirical status. The framework's predictions are *consistent* with the empirical record from single-enzyme experiments. Power-law waiting-time distributions, dynamic disorder in single-molecule kinetics, and conformational gating of catalysis are all documented across multiple enzyme families. What has not yet been done is *quantitative* testing of the framework's specific predictions — for example, testing whether the multi-time correlation structure of single-enzyme catalysis matches the proposed model form $\mathcal{F}(r) = 2r - r^2$ for capacity-controlled revival (derivation not supplied; Chapter 15 §15.7). The qualitative match is strong; quantitative discrimination from competing accounts (purely classical memory effects, purely quantum-coherent dynamics) requires further empirical work.

13.4 Photosynthesis as energy transfer through C1–C4 architectures

Photosynthetic light-harvesting complexes provide the most-studied empirical case of quantum biology. Two-dimensional electronic spectroscopy experiments (Fleming, Engel, and colleagues, beginning in the early 2000s) demonstrated that exciton transfer in photosynthetic antennae shows long-lived coherences — quantum-coherent superpositions of exciton states that persist for hundreds of femtoseconds, much longer than naive estimates of decoherence times at biological temperatures.

The framework's reading of this empirical record is structural. The photosyn-

thetic complex implements C1–C4 at the exciton-vibration interface, with the protein scaffold acting as the slow bath that produces non-Markovian dynamics.

The photosynthetic partition. In a light-harvesting complex (LH1, LH2, FMO, or others), the visible sector is the *exciton* — the photoexcited electronic state delocalized across multiple pigment molecules (chlorophylls, bacteriochlorophylls, carotenoids). The hidden sector is the *protein-vibrational environment* — the protein scaffold's vibrational modes, the solvent dynamics, and the slow conformational fluctuations that modulate the pigments' electronic energies and inter-pigment couplings.

C1–C4 satisfaction. The coupling between exciton and vibrational environment is well-characterized — electron-phonon coupling is one of the foundational mechanisms in photosynthesis modeling. The timescale separation between exciton dynamics ($\tau_S \sim 100$ fs) and slow vibrational/conformational modes ($\tau_B \sim 1$ ps to 1 ns) gives $\tau_S/\tau_B \sim 10^{-1}$ to 10^{-4}. The hidden sector's capacity is large — thousands of vibrational modes and the protein's full conformational complexity.

The framework's prediction. Photosynthetic exciton transfer should exhibit non-Markovian dynamics where the protein-vibrational environment reads back its record; information backflow from that environment, a further property, would bear on the long-lived coherences observed in 2D spectroscopy. The dynamics is *not* purely quantum-coherent (which would predict longer-lived coherence than observed); nor is it *purely classical* (Förster theory predicts no coherence at all). The mechanism proposal's reading is that the dynamics shows partial distinguishability revival — the restricted quantum-statistical regime, $0 < \mathcal{N} < 1$ (Chapter 18 §18.3) — a reading not computed from a realization.

The framework's reading aligns with the current consensus in the photosynthesis literature. The early interpretation of long-lived coherences as evidence for "quantum coherence assisting energy transfer" has been refined: subsequent experiments and theory (including recent reviews by Cao, Fleming, and others) have established that the coherences arise from vibronic coupling — vibrational modes nearly resonant with exciton energy gaps producing the observed long-lived signals. The framework's reading places this vibronic picture in a broader structural framework: the vibronic coupling is the specific implementation of C1–C4 in photosynthesis, with the non-Markovian dynamics being a structural consequence of the framework's commitments.

Empirical implications. The framework's content predicts specific features beyond the qualitative observation of long-lived coherences. The decay of the 2D spectroscopy beating signals should follow the kinetics the mechanism proposal assigns to the restricted quantum-statistical regime rather than purely Markovian or purely coherent dynamics. The temperature dependence of the coherence times should reflect the C1–C4 architecture — as temperature increases, the timescale separation τ_S/τ_B decreases and the system moves toward more classical (Markovian) behavior. The species-specific variations across different

photosynthetic organisms should reflect different specific C1–C4 parameters in their light-harvesting machinery.

These predictions are testable with current experimental techniques and would distinguish the framework's reading from purely quantum-coherent or purely classical alternatives. The framework's content here is therefore empirically engaged rather than purely interpretive.

13.5 Magnetoreception via cryptochrome radical pairs

The avian magnetic compass — the mechanism by which migratory birds sense the geomagnetic field for navigation — provides a second canonical case of quantum biology. The leading hypothesis (Ritz, Adair, Schulten, and others, from the 1970s through the present) is that magnetoreception uses a *radical-pair mechanism* in cryptochrome proteins in the avian retina: photoexcitation of cryptochrome produces a spin-correlated radical pair whose recombination yield depends on the geomagnetic field through Zeeman splitting of the radical-pair states.

The framework's content provides a structural reading of the radical-pair mechanism that connects it to the framework's broader quantum-biology architecture.

The radical-pair partition. A radical pair consists of two unpaired electrons in adjacent molecular orbitals, with the two-electron spin state initially correlated (singlet $|S\rangle$ or triplet $|T_0\rangle$, $|T_\pm\rangle$, depending on the photochemical mechanism). The visible sector is the *electron spin state* — the two-electron spin configuration that determines the recombination products. The hidden sector is the *nuclear spin environment* — the hyperfine-coupled nuclear spins of the surrounding nuclei, the molecular vibrations modulating the hyperfine couplings, and the protein scaffold dynamics affecting the radical pair's lifetime.

C1–C4 satisfaction. The hyperfine coupling between electron spins and nuclear spins is the framework's C1 coupling at the radical-pair scale — typically in the range of 0.1-10 mT magnetic field equivalents, with substantial coupling strength. The nuclear spin dynamics is slow ($\tau_B \sim$ ms to s) compared to the radical-pair lifetime ($\tau_S \sim \mu$s), giving $\tau_S/\tau_B \sim 10^{-3}$. The nuclear spin bath has substantial capacity — typically 10-100 nuclear spins coupled to the radical pair, with sequence space $2^{100} \sim 10^{30}$ states.

The framework's prediction. The radical-pair dynamics is non-Markovian, with the recombination yield depending on the radical-pair history through correlations stored in the nuclear spin bath. The geomagnetic field's effect is therefore not just the Zeeman splitting of instantaneous spin states but the *integral* over the radical-pair lifetime of the field's effect on the coupled electron-nuclear spin dynamics.

The framework's reading has specific consequences for the magnetic compass's sensitivity and selectivity. The compass's sensitivity to weak magnetic fields (~ 50 μT for the geomagnetic field) requires the radical-pair dynamics to be

sensitive to small Zeeman energies ($\sim 10^{-6}$ K of energy difference). The framework's non-Markovian dynamics provides the mechanism: the small energy difference accumulates coherently over the radical-pair lifetime, with coherence surviving long enough for the effect to be detected. The coherence preservation is a property of the specific physical realization, not a consequence of the C1–C4 architecture: (C1)–(C4) constrain coupling, timescales and capacity, and neither contain nor imply a coherence requirement. The framework's contribution here is the non-Markovian accounting of the accumulated phase, not a derivation that the phase survives.

Empirical content. The framework's reading is consistent with the empirical record on magnetoreception:

- *The geomagnetic compass requires light.* Cryptochromes are photoreceptors; the radical pair forms only after photoexcitation. The framework's reading explains why: the radical-pair partition is created by the photochemistry, and exists only during the photoexcited lifetime.
- *The compass is inclination-based, not polarity-based.* Birds detect the geomagnetic field's *inclination angle* (vertical component) but not its *polarity* (direction). This is a structural consequence of the radical-pair mechanism: the singlet-triplet conversion rate depends on the magnitude of the field's component along the radical pair's coupling axis, not on its sign.
- *Disruption by oscillating fields.* Weak radio-frequency fields ($\sim$ MHz, comparable to hyperfine couplings) disrupt the compass. The framework's reading: the RF fields scramble the nuclear spin bath, breaking the C1–C4 architecture that maintains the non-Markovian dynamics.

Predictions for further empirical testing. The framework's content makes specific predictions beyond the qualitative phenomenology. The temperature dependence of compass sensitivity should reflect the C1–C4 architecture's robustness — at higher temperatures, the timescale separation degrades and the compass should be less sensitive. The dependence of compass function on the specific isotopic composition of cryptochrome (replacing ^{1}H with ^{2}D, for instance) should reflect the framework's nuclear-spin-bath capacity. The framework's content provides structural foundations for the radical-pair mechanism that connect it to broader principles of quantum biology.

13.6 Enzyme catalysis as the framework's clearest test case

Enzyme catalysis is the framework's most direct empirical test bed in quantum biology. Single-enzyme experiments — pioneered by Sunney Xie and developed by many groups over the past two decades — provide single-molecule resolution of catalytic events, with sufficient time resolution to detect the framework's predicted non-Markovian signatures. This section develops the specific empirical predictions and their match to experimental data.

The empirical record. Single-enzyme studies of cholesterol oxidase (Xie group, late 1990s), beta-galactosidase, lipase, and other enzymes have estab-

lished several signatures of non-Markovian dynamics:

Dynamic disorder. Individual enzyme molecules exhibit fluctuations in catalytic rate over time. The fluctuations are not random noise but exhibit structured temporal correlations — slow periods (low catalytic rate) cluster together; fast periods (high catalytic rate) cluster together. The autocorrelation function of catalytic rate decays on timescales much longer than individual catalytic cycles, indicating that the rate fluctuations have *memory*.

Non-exponential waiting-time distributions. The distribution of waiting times between successive catalytic events does not follow a simple exponential distribution. Stretched exponentials, power laws, and multi-exponential fits all appear depending on the specific enzyme and conditions. The deviations from simple exponential kinetics are systematic, not noise.

Memory effects between cycles. Successive catalytic cycles are not statistically independent: the duration of a long waiting time correlates with the durations of preceding waiting times. The correlation is exactly the framework's prediction for non-Markovian dynamics — successive measurements are not independent samples but carry information from previous cycles.

Allosteric history dependence. Inhibitors and activators binding away from the active site produce activity changes with kinetics dependent on the protein's history. The "memory" is the slow relaxation of the conformational substrate to the new ligand-bound state.

The framework's structural reading. Each of these signatures is a direct consequence of the framework's prediction that the enzyme implements C1–C4 with substantial timescale separation. Dynamic disorder reflects the conformational substrate's slow fluctuations modulating active-site activity. Non-exponential waiting-time distributions reflect the non-Markovian probability density that emerges from C1–C4 dynamics. Memory effects between cycles reflect the conformational substrate's slow, non-Markovian coupling to the active site — grade-ii memory, consistent with but not isolating the grade-iii information-backflow signature of P-indivisibility (see the empirical-status note below). Allosteric history dependence reflects the slow propagation of conformational changes through the scaffold.

The framework's content here is that these are not separate phenomena requiring case-by-case explanations but *unified consequences* of the protein's C1–C4 architecture. Any enzyme with substantial conformational complexity should exhibit them; the cross-enzyme variations reflect different specific C1–C4 parameters.

Quantitative testability. The framework's prediction goes beyond qualitative agreement. The specific functional forms of the non-exponential distributions, the autocorrelation timescales, and the multi-time correlation structure are predictable from the C1–C4 parameters. In particular, the restricted quantum-statistical regime (Chapter 18 §18.3) carries the mechanism proposal's

OPERATIONAL prediction — requiring its own derivation, not supplied by the equivalence theorems ([Main §3.4]) — that the multi-time statistics should *not* admit a memoryless (Markov) description and should show distinguishability revival on accessible windows; the memory test alone does not separate the framework's reading from classical memory effects, which also fail it.

The diagnostic experiments are within reach of current single-molecule techniques: multi-time correlation measurements with sufficient statistics to constrain the algebraic structure of the transition probabilities, performed on well-characterized enzymes with known conformational landscapes. The framework's content here is therefore empirically actionable, with the prediction of the restricted quantum-statistical regime being the mechanism proposal's distinctive falsification target.

13.7 The restricted quantum-statistical regime in biology

The framework's content on quantum biology includes a structurally distinct prediction: the mechanism proposal predicts that biological C1–C4 systems lie in the *restricted quantum-statistical regime* of Chapter 18 §18.3, partial distinguishability revival on accessible windows ($0 < \mathcal{N} < 1$) — a revival range, not a degree of quantum membership ([Main §3.4]), and not implied by C1–C4, which give memory without fixing $\mathcal{N}$ ([Main §2.3]). Partial revival itself is standard open-systems phenomenology, quantified in the literature by the Breuer–Laine–Piilo measure; what is distinctive is the proposed capacity-controlled origin and scaling law, the mechanism proposal's prediction for systems with marginal hidden capacity.

The regime's empirical signature. A memoryless (Markov) system has multi-time statistics generated by a divisible family. A memory-bearing system does not, and may or may not show distinguishability revival; $\mathcal{N}$ measures revival, not memory ([Main §2.3, §3.4]). A system in the restricted quantum-statistical regime is not distinguished by lacking a representation — every stochastic transition family admits the one-step ancilla dilation ([Main §3.1]) — but by partial revival; the mechanism proposal further hypothesizes that its operational structure gives out there: coherent preparation, intervention structure, and cross-horizon representation consistency. The proposal's diagnostic signature is the failure of specific positivity and completeness conditions on multi-time correlation functions — a statement about the correlations, not about whether a dilation exists, and a proposal-level claim requiring its own derivation ([Main §3.4]).

Biological systems in the restricted quantum-statistical regime. The biological scales developed in §§13.3-13.6 — protein conformational landscapes ($\tau_S/\tau_B \sim 10^{-9}$ to 10^{-12}), photosynthetic light-harvesting ($\tau_S/\tau_B \sim 10^{-1}$ to 10^{-4}), magnetoreception ($\tau_S/\tau_B \sim 10^{-3}$), single-enzyme catalysis ($\tau_S/\tau_B \sim 10^{-9}$ to 10^{-12}) — are slow-bath ratios. Hidden-sector capacities range from $\sim 10^{30}$ (radical pair) to $\sim 10^{200}$ (protein conformational landscape) — large

but finite, the finiteness capping accessible memory ([Main §3.4]). Capacities of this size lie far above the comparable-capacity range $N_H/N_V \sim 1$ of Chapter 15 §15.7, so the placement of biological systems in the restricted quantum-statistical regime is a conjecture that the capacity ratio does not fix.

The proposal's conjecture is that *biological systems show partial distinguishability revival (the restricted quantum-statistical regime)* at the C1–C4 scales of interest — a revival claim, not a quantum-membership claim ([Main §3.4]). This is structurally distinct from "biology has quantum effects" (which the framework's reading agrees with) and from "biology is fundamentally quantum-mechanical at all scales" (which the framework's reading rejects). The restricted quantum-statistical regime is the mechanism proposal's specific commitment.

The empirical access. Testing the prediction of the restricted quantum-statistical regime requires multi-time correlation measurements with sufficient statistics to constrain the algebraic structure of the transition probabilities. Standard single-molecule techniques produce two-time correlations (waiting-time distributions, rate-rate correlations); multi-time correlations require longer-duration experiments with more careful statistics. The technology is available — modern single-molecule fluorescence and force spectroscopy can produce three-time and four-time correlations — but the analyses have not been performed systematically against the framework's specific algebraic predictions.

The framework's distinctive falsification target in quantum biology is therefore: a multi-time correlation measurement on a well-characterized enzyme (cholesterol oxidase, lipase, or another single-molecule benchmark) that establishes whether the transition probabilities admit a memoryless (divisible) density-matrix description. If yes (the memoryless description holds), this operational prediction of the mechanism proposal is falsified; the enzyme would be Markovian at the accessible horizon. If no (the memoryless description fails), the enzyme is memory-bearing at the accessible horizon, as the C1–C4 reading requires; partial revival is a further claim of the proposal, tested by a distinguishability-revival measurement and stated as its own falsifiable claim rather than a consequence of the frozen equivalence theorems.

The framework's distinctive content. The restricted quantum-statistical regime is the mechanism proposal's contribution to quantum biology that goes beyond reproducing existing empirical content. Other accounts of quantum biology (photosynthetic exciton transfer through vibronic coupling, radical-pair mechanisms for magnetoreception, conformational gating for enzyme catalysis) can reproduce specific empirical signatures but do not predict the framework's specific algebraic structure of multi-time correlations. The framework's prediction is testable, falsifiable, and quantitatively specific — the kind of empirical commitment that distinguishes a structural framework from a phenomenological account.

13.8 Implications for drug design

The framework's content on quantum biology has practical consequences for drug discovery and clinical pharmacology. Standard drug design assumes Markovian kinetics — the drug-target binding follows the law of mass action, with rate constants set by the current concentrations. The framework's prediction is that this assumption breaks down for drugs targeting proteins with substantial conformational complexity, producing systematic deviations from Markovian predictions that have measurable clinical consequences.

Kinetic implications. Standard pharmacology characterizes a drug-target interaction with two rate constants: k_{on} (binding rate) and k_{off} (dissociation rate), with the equilibrium dissociation constant $K_d = k_{\text{off}}/k_{\text{on}}$ determining the binding affinity. The framework's prediction is that these rate constants are *not constants* but vary with the protein's conformational state — fast periods (high k_{on}, low k_{off}) cluster together; slow periods cluster together. The clinical consequence: a drug's effective potency varies over time in ways the constant-rate-constant model cannot capture.

Allosteric implications. Allosteric drugs (binding away from the active site to modulate activity) operate through the protein's conformational landscape — exactly the framework's C1–C4 hidden sector. The framework predicts that allosteric drug effects have memory: a drug's allosteric effect at time t depends not just on its current concentration but on the protein's history of recent ligand binding. The clinical consequence: dosing strategies that take advantage of memory effects (pulsed vs continuous administration) may produce different clinical outcomes than the Markovian assumption predicts.

Resistance mechanisms. Drug resistance often involves mutations in the target protein that change the active site or the allosteric communication network. The framework's reading: resistance is a *memory-structure alteration* — the mutations change the protein's C1–C4 architecture, producing different non-Markovian dynamics and therefore different drug-target kinetics. The framework's content suggests that resistance can be combatted by designing drugs that exploit specific features of the C1–C4 architecture rather than just optimizing static binding affinity.

Catalytic vs regulatory specificity. The framework's structural content distinguishes drugs that target the active site (modulating the fast subsystem) from drugs that target the regulatory landscape (modulating the slow subsystem). The two categories have different kinetic signatures and different clinical implications. Chapter 16's medicine content develops this distinction in detail for specific clinical contexts; the structural foundation is the C1–C4 architecture developed in this chapter.

The magnitude of corrections. The framework's corrections to standard drug kinetics scale as $\mathcal{O}(\tau_S/\tau_B)$ — typically 10^{-9} to 10^{-12} for protein systems. This is small at the individual-event level but can accumulate over many cat-

alytic cycles or in systems with exceptionally slow conformational dynamics. For drug design, the practical question is whether the cumulative effect reaches clinical relevance at the doses and timescales of therapeutic use. The framework's content provides a structural basis for predicting when corrections matter rather than just empirically discovering them case by case.

13.9 Chapter close

The framework's content in quantum biology unifies several apparently disparate phenomena — photosynthetic exciton transfer, magnetoreception, enzyme catalysis — under the common structural framework of C1–C4 dynamics. The framework's reading reverses the conventional treatment: the memory-bearing dynamics the framework marks is not the exception in biology but the rule wherever the structural conditions hold — its laws admitting the quantum representation ([Main §3.4]) — and biological systems implement C1–C4 routinely at the molecular scale.

Three categories of empirical content. The chapter's content sorts into three categories.

Established empirical content that the framework's reading unifies. Photosynthetic exciton transfer through vibronic coupling, magnetoreception through radical-pair dynamics, enzyme dynamic disorder and memory effects — all are documented experimentally and consistent with the framework's structural account. The framework's contribution is to provide a unified explanatory framework rather than case-by-case mechanisms.

Quantitative predictions that the framework makes. The specific functional forms of non-exponential kinetics, the multi-time correlation structure of single-enzyme experiments, the temperature dependence of the restricted quantum-statistical regime's signatures, and the algebraic structure of biological transition probabilities. These predictions are within reach of current experimental techniques and provide concrete tests of the framework's reading.

Predictions for future empirical work. The restricted quantum-statistical regime's diagnostic signature (distinguishability-revival measurements, with multi-time correlation measurements testing memory) is the mechanism proposal's distinctive falsification target. Engineered biological systems with tunable C1–C4 parameters (designed proteins, synthetic photosynthetic systems, artificial radical-pair platforms) could provide controlled tests of the framework's quantitative predictions.

Forward pointers. Chapter 14 develops the framework's content in quantum computing — the BQP characterization theorem as the framework's structural limit on computational complexity, with the framework's content on biological C1–C4 systems as a special case of the broader computational architecture. Chapter 15 develops quantum engineering — applications of the framework's C1–C4 architecture to designed physical systems, with the BEC analogue-

gravity experiment as the cleanest near-term test of the characterization theorem at the foundational level. Chapter 16 develops medical applications, including the substratum-emergent operator dissociation at chromatin and the wider-therapeutic-window prediction for reader-writer disorders — applications that build directly on this chapter's structural foundations.

The framework's content from this chapter onward shifts from cascade derivations to specific empirical applications: how the framework's structural commitments at the chemistry and biology levels produce specific predictions about quantum computing platforms, engineered quantum systems, clinical pharmacology, and bioinformatics methodology. The chapter's content on the protein C1–C4 architecture provides the foundational example that the subsequent chapters build on.

Chapter 14

Quantum Computing

> **Status — conjectural extension.** This chapter applies the core framework to quantum computation and complexity, and is offered as a *conjecture*, not an established result: the central claim — that the framework's emergent dynamics realizes the complexity class BQP — is a theoretical conjecture rather than a proven correspondence; it would be supported by a rigorous derivation from the finite substratum to BQP and undercut if that dynamics proves efficiently classically simulable or fails to reproduce known quantum-complexity separations. It carries no evidential weight for the core framework — which stands or falls independently on its physics — and should be read as a structural possibility to be tested, not as support for the framework.

14.1 What this chapter develops

The framework's content across Chapters 5 through 13 has established a substantial structural account of fundamental physics, the cascade through chemistry and biology, and the specific empirical content of quantum biology. The cascade's logic — substratum, partition, emergent quantum mechanics, structural conditions C1–C4 producing non-Markovian dynamics at multiple scales — provides the foundation for the framework's content in computational complexity theory.

This chapter develops the framework's characterization of computational power available to embedded observers. The central result is a *conditional BQP model*

theorem: under an explicit computational model — a uniform efficient encoding of substratum configurations into qubits; efficient state preparation and control; a clock model; efficient measurement; explicit precision accounting; noise assumptions under which fault tolerance applies with polynomial overhead; and operational closure — all admissible operations available to the observer within the model are standard quantum instruments (CPTP maps for unconditioned evolution, CP trace-nonincreasing branches for conditioned outcomes), with no additional nonlinear, post-quantum, oracle, or hidden-sector-access operation, and with conditioned branches and postselected preparations charged their physical success-probability/repetition cost (no free postselection: PostBQP = PP) — BQP (bounded-error quantum polynomial time) is the computational class accessible to a finite deterministic system satisfying the framework's structural conditions C1–C4. None of the model hypotheses follows from C1–C4 or from CPTP representability alone. The theorem has two directions. The *lower bound* establishes that, under the explicit computational-control hypotheses of §14.4, any embedded observer satisfying C1–C4 has access to BQP computational resources — the computational-control model supplies state preparation, a universal gate set, and Born-rule measurement — the full toolkit for BQP computation — on top of Main's representation and finite-test-shadow layer. The *upper bound* establishes that no embedded observer exceeds BQP — the quantum description is informationally complete for the embedded observer, with no operation on the visible sector extracting information beyond the quantum state.

Ten pieces occupy the chapter.

Section 14.2 develops the chapter's framing: the framework's content in computational complexity is at the foundational level rather than the algorithm-design level. The framework does not produce new quantum algorithms; it characterizes the *ceiling* on what embedded observers can compute. Section 14.3 develops the lower bound: BQP is available to any embedded observer satisfying C1–C4 under the explicit computational-control hypotheses of §14.4. The argument runs through the framework's quantum representation from Chapter 1, with state preparation, gate application, and measurement supplied by §14.4's stated model, not by C1–C4 alone.

Section 14.4 develops the upper bound: BQP cannot be exceeded. The argument runs through the framework's informational completeness — the quantum description exhausts the information accessible through any admissible operation — operational closure, a model hypothesis of §14.4 — so, with the efficiency hypotheses carrying the complexity content, no super-BQP computation is possible within the model. Section 14.5 combines the two bounds into the conditional BQP model theorem: under the model hypotheses, BQP is the computational class accessible to embedded observers satisfying C1–C4.

Section 14.6 develops the quantum (BQP-form) Extended Church-Turing Thesis as a conditional result of the framework, valid under a global-coverage premise: every physically realizable computation falling under §14.4's model hypotheses. The BQP-form thesis — that any physically realizable computation can be ef-

ficiently simulated by a quantum computer, the quantum counterpart of the conventional classical efficient-simulation conjecture that quantum computing challenges — is conventionally treated as an empirical conjecture about physical reality. The framework's content reframes the ECT as a derived result under an explicit coverage premise: given global computational-model coverage — every physically realizable computation is implemented by a physical system whose admissible computational operations satisfy §14.4's operational-closure and efficiency hypotheses, every such computation is BQP-simulable. The upper bound needs only that coverage — no device is required to satisfy C1–C4 or to possess the lower-bound control toolkit. The reframing changes the ECT's epistemic status from freestanding empirical conjecture to conditional theorem — a derived result valid under the coverage premise and the same explicit model hypotheses (uniform encoding, efficient preparation, control, and measurement, and polynomial-overhead fault tolerance) as the BQP characterization it rests on.

Section 14.7 develops task-specific quantum advantage — the conjectured BPP–BQP separation: the advantage is real but is *a feature of partial observation* rather than of the universe being fundamentally quantum. An embedded observer who ignores the quantum structure of the reduced description (treating measurements as classical samples) operates within BPP; one who exploits the quantum structure (interference, entanglement, adaptive measurement) operates within BQP. Under the conjectured strict separation, the difference between the two is read as the cost of not recognizing the partition's structure — proven task-specific advantages are facts; the class-level gap remains the standard conjecture. Section 14.8 develops the framework's neutrality on the P vs NP question: the BQP characterization identifies the ceiling, while leaving the internal structure of complexity classes within that ceiling — including P vs NP — to standard complexity theory; the silence on P vs NP is shown to fit a broader boundary-characterization pattern across the Gödel-Turing-OI-Deutsch incompleteness family. Section 14.9 develops three research directions opened by the framework's specific substratum structure for complexity-theoretic refinements *within* the BQP ceiling (geometric structure on the substratum lattice; conjectured BQP–BPP separation correlated with problem globality and a refined hidden-subgroup-problem / gauge-group conjecture; gauge equivalence as a possible physical account of polynomial-time reducibility). Section 14.10 develops the chapter's falsification conditions and connections to broader complexity theory, with closing forward pointers to Chapter 15's engineering content and Chapter 18's foundational content.

The chapter's content is foundational rather than algorithmic. The framework's contribution to quantum computing is structural: a characterization of the computational ceiling that emerges from the framework's commitments at the substratum level joined to §14.4's model — operational closure and efficiency included — with BQP the class accessible to embedded observers under that model. The empirical content is principally negative: the framework predicts, under global coverage of §14.4's model, that no super-BQP computation is pos-

sible on any physically realizable hardware, with any demonstrated super-BQP capability falsifying first the global-coverage premise and only with additional argument the framework's commitments in conjunction with §14.4's computational model — sharpest at the model bridge.

14.2 The framework's content in computational complexity

The framework's content in computational complexity operates at a specific level of abstraction. The framework does *not* derive new quantum algorithms — Shor's factoring algorithm, Grover's search algorithm, the HHL linear-system algorithm, and other specific quantum computational achievements are not framework results but standard quantum-computing content that the framework's computational-control model, on Main's representation layer, reproduces. What the framework adds is a structural account of *why* BQP is the relevant complexity class for quantum computing — why this specific class of problems is the one accessible to embedded observers using quantum-mechanical computation.

The conventional treatment. Standard quantum complexity theory establishes BQP as a complexity class with specific defining properties: polynomial-time computation on a quantum Turing machine with bounded error. BQP contains BPP (bounded-error probabilistic polynomial time, the classical analog) and is conjectured to extend beyond it (Shor's algorithm being the canonical example of BQP computation believed not to be in BPP). The relationship of BQP to other complexity classes — its relationship to NP, PH, PSPACE, and others — is the subject of active research.

The conventional treatment takes BQP as defined by the quantum-computing model and asks empirical questions: what problems are in BQP? what are the boundaries of BQP relative to other classes? what specific algorithms achieve BQP computation? The framework's content addresses a different question, one level below: *why* is BQP the relevant class for quantum computing in our physical universe?

The framework's level. The framework's content treats BQP as a *derived* feature of physical reality rather than as a primitive defined by the quantum-computing model. The derivation runs through the framework's structural commitments: under the explicit computational-control hypotheses of §14.4, any embedded observer in a finite deterministic substratum satisfying C1–C4 has access to BQP computational resources, and no embedded observer exceeds BQP. Under those hypotheses the BQP class is *the class accessible to embedded observers* — a feature of the framework's universality class joined to §14.4's model rather than an empirical property of specific computational architectures.

This is a different epistemic status than the conventional treatment. In the conventional treatment, BQP is a *useful concept* for organizing computational results, with its empirical applicability being a contingent feature of our physical universe. In the framework's content, BQP is a *conditional structural ceiling* — any physical universe with the framework's commitments and §14.4's

computational-control model has BQP as the unique computational ceiling for embedded observers.

Why this matters. The framework's level matters for two reasons.

The ceiling is principled rather than empirical. The conventional treatment leaves open the possibility that some future physical principle could extend computational access beyond BQP. The framework's content closes this possibility: super-BQP computation would require an embedded observer to exceed the limits the framework identifies under §14.4's model, which is mathematically impossible if the framework's commitments and that model are both correct. The ceiling is a conditional theorem — framework plus model — not an empirical bound.

The falsification condition is sharp. The conventional treatment makes no specific predictions about whether super-BQP computation is possible — it remains an open question subject to future discovery. The framework's content predicts, under global coverage of §14.4's model, that no super-BQP computation is possible on any physically realizable hardware, with any demonstrated super-BQP capability falsifying first the global-coverage premise, and only with additional argument the framework-plus-model conjunction and its bridge. The falsification condition is empirically actionable: future quantum computing experiments could in principle test the framework's prediction.

The framework's content in this chapter is therefore at a foundational level — a structural account of why BQP is the relevant complexity class, with empirical predictions that distinguish the framework from alternative accounts of computational complexity in physical systems.

14.3 BQP is available: the lower bound

The lower bound of the BQP characterization theorem establishes that, under the explicit computational-control hypotheses of §14.4, an embedded observer satisfying C1–C4 has access to BQP computational resources. The argument runs through the framework's quantum representation from Chapter 1: the observer's reduced description admits the representation, and the standard quantum-computing toolkit (state preparation, universal gate set, Born-rule measurement) is available to the observer through standard operations on the visible sector.

The observer's quantum toolkit. Chapter 1 developed the framework's admitted quantum representation for any embedded observer satisfying C1–C4: a Hilbert space of visible-sector states, unitary evolution under the reduced Hamiltonian, and fixed-basis Born-rule readout — the representation the equivalence supplies universally ([Main §3.4]) — with composite-system factorization derived from coupling-graph locality for spatially separated subsystems ([Main §3.1]). That is the transition-statistics layer. The full operational toolkit — coherent state preparation and control, noncommuting instruments, projective

measurement beyond the fixed basis — is *not* delivered by C1–C4 alone: it enters through the explicit computational model of the conditional BQP theorem (§14.4; [Computation §4]), whose hypotheses — uniform efficient encoding, state preparation and control, a clock model, efficient measurement, precision accounting, and fault-tolerance-grade noise assumptions — are modelling commitments, CPTP representability supplying none of them by itself.

The standard quantum-computing toolkit is built on exactly these features. *State preparation* is supplied directly by §14.4's computational-control model as an explicit hypothesis and must be efficiently implementable; probabilistic preparations are charged their physical success-probability/repetition cost, per the model's postselection rule ([Main §3.4] for the operational layer's finite-test realization). *Gate application* (transforming the quantum state through unitary operations) corresponds to time evolution under controllable Hamiltonians — available wherever the observer can engineer the visible sector's dynamics. *Measurement* (reading the computation's output) corresponds to standard projective measurement — the framework's fundamental observational operation.

Universal gate set. Quantum computing requires a universal gate set: a collection of single-qubit and two-qubit gates that can approximate any unitary transformation to arbitrary precision. The standard universal gate set includes the Hadamard gate, a non-Clifford phase gate such as T, and the CNOT (controlled-NOT) gate, with these three gates generating an approximately universal set — dense in the unitary group, not an exact enumeration of it — through composition. Each of these gates is implementable in any system with controllable two-level subsystems and a tunable coupling — features supplied by §14.4's computational-control model — Main supplying the representation layer, the model the controllability.

The argument is concrete: an observer who can prepare two-level subsystems (qubits) in the visible sector, who can apply controllable Hamiltonians to those subsystems, and who can perform projective measurements on the subsystems has access to a universal gate set. These three capabilities are exactly the explicit computational-control model this chapter's opening states as additional hypotheses: preparation, controllable Hamiltonians, and projective readout are supplied by the computational-control model — Main's finite-test result supplies the representation and shadow layer for specified experiments ([Main §3.4]), not controllability — and not by C1–C4 alone. Under that stated model, any embedded observer has access to universal quantum computation in principle.

Error correction. Practical BQP computation requires fault tolerance: the ability to perform long computations despite physical errors at each gate. Quantum error correction (surface codes, stabilizer codes, topological codes) provides the theoretical framework for fault tolerance with overhead scaling logarithmically in the desired computational accuracy. The framework's quantum representation provides the structural setting, the control model the operations: error correction operates on the quantum information in the visible sector, with redundant encoding protecting against local errors.

The framework's content here is consistent with standard quantum-computing theory: error correction is structurally available to any embedded observer with sufficient control over the visible sector, with the specific overhead determined by the physical error rates (which depend on the framework's specific physical parameters rather than on structural commitments at the framework level). Chapter 15 develops the framework's content on engineered error correction with correlation-aware decoders that exploit the framework's predicted non-Markovian noise structure.

The lower bound theorem. Combining the four elements — state preparation, universal gate set, measurement, error correction — establishes the lower bound:

Conditional Lower Bound. Under the computational-control hypotheses of §14.4, an embedded observer satisfying C1–C4 can host any uniform polynomial-size BQP circuit family: the stated model supplies state preparation, a universal gate set (Hadamard, a non-Clifford phase gate such as T, and CNOT), and Born-rule measurement; error correction is available within the same model; the cumulative result is access to polynomial-time quantum computation with bounded error.

The lower bound is therefore established conditionally: BQP access is a capability of every embedded observer satisfying C1–C4 *together with* §14.4's computational-control model — uniform across the framework's universality class under those hypotheses, not a consequence of C1–C4 alone.

14.4 BQP cannot be exceeded: the upper bound

The upper bound of the BQP characterization theorem establishes, under the efficiency hypotheses stated below — locally generated, uniformly describable evolutions over polynomial physical time, with polynomial precision and polynomial simulation overhead — that no embedded observer exceeds BQP; informational completeness fixes the operations' type, and these efficiency hypotheses, not completeness alone, carry the complexity bound. The argument runs through *informational completeness relative to the model*: the quantum description exhausts the information accessible through the admissible quantum instruments — operational closure, the model hypothesis stated above; completeness for arbitrary conceivable operations is not asserted. And extra computational power would not even require extra information: modified or nonlinear evolution on the same state description already changes the complexity class — which is precisely what the closure hypothesis excludes.

Informational completeness relative to the admissible instruments (model-scoped). Within the operational-closure model, the density matrix ρ_V is a complete description: every admissible operation — the quantum instruments the model specifies — operates on ρ_V and produces outcomes consistent with its statistical predictions. Completeness for arbitrary conceivable

operations is not asserted; it is exactly what the closure hypothesis supplies, not what the framework proves.

Inaccessibility of the complete state (Chapter 1) is real but does not by itself yield closure: extra computational power need not come from extra information — modified or nonlinear evolution acting on the same state description already changes the complexity class, which is what the closure hypothesis excludes. The observer can detect the *statistical* effect of the hidden sector through the density matrix's mixedness but cannot detect the complete hidden-sector configuration. Hidden-sector access would be one possible route beyond the model, but not the only one; nonlinear or other post-quantum operations on the same state space are excluded separately — by the operational-closure hypothesis, not by inaccessibility.

Operational closure (a model hypothesis). The computational model assumes — it does not derive from C1–C4 — that any admissible operation the observer performs on the visible sector within the model is a *quantum operation* in the standard sense: a completely positive trace-preserving (CPTP) map on the density matrix ρ_V for unconditioned evolution, with CP trace-nonincreasing branches for conditioned outcomes, and no additional nonlinear, post-quantum, oracle, or hidden-sector-access operation, and with conditioned branches and postselected preparations charged their physical success-probability/repetition cost (no free postselection: PostBQP = PP). Whether the OI intervention family closes exactly into one standard coherent quantum instrument category is characterized, within complex matrix kinematics taken as given, by the five completion conditions of the operational-completion characterization, of which the substratum's stated interventions do not supply full reversible control; whether an extended OI mechanism can supply it, and which of the others the substratum supplies, are open ([Main §3.4]; [GR §3.3]). This includes unitary evolution (a special case, purity-preserving), projective measurement (an instrument whose unconditioned map is CPTP), and conditioned measurement branches (CP and trace-nonincreasing; the normalized post-outcome update divides by the outcome probability and is therefore nonlinear). Conditioned branches and postselected preparations are charged their physical success-probability/repetition cost: no exponentially rare outcome may be selected as a free computational primitive — free postselection yields PostBQP = PP, not BQP — and any postselection the model uses must have polynomial total implementation or expected-repetition cost.

The standard dilation picture motivates the model choice — motivation for the hypothesis, not a derivation of closure. The embedded observer's operations are implemented through interactions between the observer's measurement apparatus and the visible sector. The apparatus itself is part of the visible sector (it is composed of visible-sector degrees of freedom). Any operation the apparatus performs is therefore a unitary evolution of the joint apparatus-system Hilbert space followed by projective measurement of the apparatus state. By the standard derivation of CPTP maps from such constructions, operations of

that dilation form are CPTP — which is why the standard quantum instrument set is the natural model choice; whether every allowed OI intervention belongs to one coherent quantum instrument category is that same question, and the model assumes closure over exactly that set.

Computational complexity of CPTP maps. The relationship runs in one direction only: any polynomial-size quantum circuit (with the stated ancilla-and-measurement model) implements a CPTP map, but a generic CPTP map on n qubits has no polynomial-size implementation — unitary channels are already a counterexample, generic unitary synthesis requiring $\Theta(4^n)$ elementary gates (Shende, Bullock, and Markov). CPTP is the operation's type; efficient uniform implementability is a separate hypothesis, supplied here by §14.4's efficiency model. Efficiently and uniformly implementable admissible channels can be simulated by uniform polynomial-size quantum circuits, up to the standard polynomial encoding; computations composed from them therefore decide only languages in BQP.

The framework's content, joined to the model, therefore identifies BQP as the class of problems efficiently decidable by computations composed from the model's admissible operations. Within the model, the observer performs no operation that is not a quantum instrument — operational closure, assumed above — and under §14.4's efficiency hypotheses each admissible channel's efficient uniform implementation compiles to a uniform polynomial-size quantum circuit, so computations composed from them decide only languages in BQP. Under those hypotheses, therefore, the observer cannot efficiently perform computation outside BQP.

The upper bound theorem.

Conditional Upper Bound Theorem. Under §14.4's operational-closure and efficiency hypotheses, no embedded observer satisfying C1–C4 can efficiently perform computation outside BQP. Every admissible unconditioned operation is CPTP, with conditioned branches CP and trace-nonincreasing — together the standard quantum instruments of the operational-closure hypothesis; under the stated hypotheses of uniform description, polynomial time and precision, and polynomial simulation overhead, efficiently and uniformly implementable admissible channels can be simulated by uniform polynomial-size quantum circuits; therefore, under those hypotheses, computations composed from them decide only languages in BQP.

Addressing objections. Three potential objections to the upper bound are worth addressing.

"*What about hypercomputation?*" Hypercomputation refers to computational models that exceed Turing-machine capabilities — infinite-time computation, oracle access to undecidable problems, analog computation with arbitrary-precision continuous quantities. The exclusions come from two different layers: infinite-time computation requires actual infinity, which the framework's finite substratum precludes (a structural exclusion); arbitrary-precision continuous

computation requires sub-Planck-scale resolution, which the lattice precludes (structural); oracle access, by contrast, is excluded by the operational-closure hypothesis — an oracle need not arrive through hidden-sector information, so this exclusion is the model's, not a framework derivation. Hypercomputation is therefore excluded structurally in its infinite-resource forms and by the model in its oracle form.

"What about exotic physics?" Various proposed exotic physics scenarios — closed timelike curves, naked singularities, quantum gravity effects — have been suggested as potential routes to super-BQP computation. The framework's content provides specific responses: closed timelike curves are inconsistent with the framework's deterministic substratum dynamics; naked singularities are excluded by the framework's bounded-coupling-degree commitment; quantum gravity effects, should they carry computational content for embedded observers, would either remain within §14.4's model or demonstrate failure of global coverage or of the model assumptions — the framework does not establish structurally that they cannot. The infinite-resource routes are closed by the framework's structural commitments; the remainder are governed by the coverage-first falsification hierarchy.

"What about future discoveries?" The framework's content does not preclude future discoveries that revise our understanding of physical reality. What it does preclude is super-BQP computation *consistent with the framework's commitments*. A future discovery that demonstrated super-BQP computation would falsify first the global-coverage premise — some physical computation escaping §14.4's model — and only with further argument the framework-plus-model conjunction, its bridge the first suspect within it. The framework therefore makes a sharp empirical prediction: under global coverage of §14.4's model, no super-BQP computation is possible on any physically realizable hardware — a device escaping the model would falsify coverage first.

14.5 The BQP characterization theorem

Combining the lower bound (BQP is available, §14.3) with the upper bound (BQP cannot be exceeded, §14.4) yields the BQP characterization theorem.

Statement. *Conditional BQP Characterization Theorem.* Let $\mathcal{S} = (S, \varphi)$ be a finite deterministic system, let $\mathcal{O}$ be an embedded observer with partition V satisfying C1, C2, C3, C4 at the partition boundary, and assume the computational-control and efficiency hypotheses of §14.4 — state preparation, the universal gate set, and Born-rule readout; locally generated, uniformly describable evolutions over polynomial physical time, with polynomial precision and polynomial simulation overhead; and operational closure, all admissible operations being standard quantum instruments with no additional nonlinear, post-quantum, oracle, or hidden-sector-access operation, and with conditioned branches and postselected preparations charged their physical success-probability/repetition cost (no free postselection: PostBQP = PP). Then the class of computations

the observer can efficiently perform (in polynomial time with bounded error) is exactly BQP.

Proof outline. The lower bound from §14.3 establishes that BQP is available to the observer. The upper bound from §14.4 establishes that BQP cannot be exceeded. Therefore the class of efficient computations available to the observer is exactly BQP.

The proof relies on the framework's structural commitments at multiple points: - The framework's finite substratum guarantees finite-dimensional emergent Hilbert spaces. - The framework's deterministic bijection φ guarantees unitarity at the substratum level. - The framework's structural conditions C1–C4 guarantee accessible conditional memory, through the readback gap of C4; the fixed-basis quantum representation is internal and universal, with the compressed form for processes in the correspondence's class under (T). - The computational model assumes operational quantum closure: informational completeness of the density matrix for all admissible operations is a model hypothesis, not a partition-structure guarantee — deriving it is the operational-selection frontier ([Main §3.4]).

The first three are structural features of the framework's universality class; the fourth is §14.4's added model hypothesis — and together with the computational-control and efficiency hypotheses they produce the conditional BQP characterization.

Interpretation. The characterization theorem has several interpretive consequences.

BQP as the model-level computational class. For any universe in the framework's universality class, and under the model hypotheses above, BQP is the relevant computational complexity class for embedded observers. The specific physical implementation of the computational hardware does not matter — superconducting qubits, trapped ions, topological qubits, photonic systems, neutral atoms, or any other platform — all have access to BQP and only BQP under the same §14.4 model.

BQP is independent of the specific bijection. The framework's universality class contains many bijections producing different specific physical content (different particle spectra, different gauge groups, different specific parameter values). Among the bijections within the framework's universality class, however, the computational ceiling is the same: BQP. The characterization is at the framework level rather than the specific-bijection level.

BQP is the ceiling for any future technology. No future technological development — better qubits, better error correction, exotic computational substrates — can exceed BQP for embedded observers in the framework's universality class, so long as §14.4's model hypotheses continue to describe the hardware. The ceiling is structural-plus-model rather than technological.

The conditional model theorem is therefore a substantial but hypothesis-bearing

result: given the model, it establishes BQP as the computational class accessible to embedded observers, holding at the structural level of the framework's universality-class commitments rather than at the specific-implementation level — and only as far as the model hypotheses reach.

14.6 The quantum (BQP-form) Extended Church-Turing Thesis as a conditional theorem

The conventional Extended Church-Turing Thesis (ECT) is the classical efficient-simulation conjecture — any physically realizable computation efficiently simulable by a probabilistic classical Turing machine — the thesis that quantum computing challenges. Its quantum counterpart, the BQP-form (quantum) ECT, asserts that any physically realizable computation can be efficiently simulated by a quantum Turing machine; it is this BQP-form thesis the framework addresses. Conventionally, the BQP-form thesis is treated as an *empirical conjecture* about physical reality: the conjecture is not derived from any deeper principle but is supported by the absence of known super-BQP computations and by general considerations about the structure of physical theories. The conjecture's empirical status leaves open the possibility that future discoveries could refute it by demonstrating super-BQP computation in some physical system.

The framework's content reframes the ECT from empirical conjecture to *conditional theorem* — valid under §14.4's computational-control and efficiency hypotheses together with the global-coverage premise.

The framework's argument. The reframing runs through three steps.

Step 1: Every physical observer is embedded. The cosmological horizon provides a partition for any observer in the universe — the partition between the observable universe (visible to the observer in principle, given sufficient instruments) and the trans-horizon region (causally disconnected from the observer). The observer is therefore embedded in the framework's structural sense: the horizon partition exists for any observer in the universe. Embeddedness alone, however, does not establish the C1–C4 conditions or §14.4's computational model for every device — that is a separate premise, stated here as *global computational-model coverage — every physically realizable computation is implemented by a physical system whose admissible computational operations satisfy §14.4's operational-closure and efficiency hypotheses.*

Step 2: Under §14.4's hypotheses, every embedded observer satisfying C1–C4 together with the stated model has access to BQP and only BQP. This is the Conditional BQP Characterization Theorem from §14.5. Any observer satisfying C1–C4 together with the stated model has access to BQP computational resources (the lower bound) and cannot exceed BQP (the upper bound). Under those hypotheses, the computational class accessible to the observer is exactly BQP.

Step 3: Under global coverage, every physical computation is BQP-simulable. The universal quantifier is carried by the coverage premise alone: every physically realizable computation is implemented by a system whose admissible operations satisfy §14.4's operational-closure and efficiency hypotheses, and every computation so covered is BQP-simulable. Embeddedness is not needed for this corollary — C1–C4 and the control toolkit matter for the exact characterization of a qualifying observer, not for the universally quantified upper bound.

The ECT is therefore not a freestanding empirical conjecture but a *conditional theorem* — a derived consequence of the global-coverage premise together with §14.4's computational model, the framework's structural commitments motivating both. The reframing changes the ECT's epistemic status: the question "is the ECT correct?" becomes "does §14.4's model globally cover physical computation — and are the model bridge and the framework's commitments correct?", which is testable through the framework's other empirical content (Chapters 5-13).

Implications of the reframing.

Sharper predictions. The reframing produces sharper empirical predictions. Any demonstrated super-BQP computation would falsify the global-coverage premise first, and only with further argument the conjunction of the framework's structural commitments with §14.4's computational model — the model bridge could fail without the foundational framework failing, so the falsification lands on the conjunction, sharpest at the bridge. This is still a sharper commitment than the conventional ECT, which allows the possibility of future revisions without committing to a specific structural commitment.

Structural foundations. The reframing provides structural foundations for the ECT that were previously absent. The ECT's empirical status conventional treatment provides no explanation for *why* the ECT should hold — it remains an unexplained regularity of physical reality. The framework's content provides the explanation: the ECT holds, under global coverage of the model, because every physically realizable computation then falls under §14.4's operational-closure and efficiency hypotheses, and computations so covered are BQP-simulable — embeddedness supplies the partition, coverage supplies the premises. The structural basis is the framework's universality-class commitments joined to that explicit model.

Connection to other foundational results. The reframing connects the ECT to other foundational results in the framework. The Bekenstein-Hawking entropy bound (Chapter 7 §7.5) limits the information content of any region; the BQP characterization limits the computational power of any embedded observer under §14.4's model; the entropy bound follows from the framework's commitments at the substratum level, the computational bound from those commitments joined to the explicit model. The connection unifies the information-theoretic and computational bounds under a common structural framework.

Connection to Deutsch's jump-to-universality concept. The reframing also in-

stantiates a specific structural pattern Deutsch identifies in [Deutsch]: the *jump to universality*, in which a sufficiently rich system suddenly acquires unbounded reach within its domain. Alphabets jump to universal writing; DNA jumps to universal genetic encoding; Turing computation jumps to universal computation. The framework's BQP characterization gives precise content to one such jump in physical computation: a finite deterministic substratum satisfying C1–C4, joined to §14.4's computational model, achieves the BQP computational ceiling — universal physical computation up to that ceiling and no further. The promotion of the ECT from empirical conjecture to derived theorem is the framework's specific contribution to this pattern: the jump to universality in physical computation reaches BQP and stops there, with the stopping point itself being a structural feature of the framework's commitments joined to §14.4's model, rather than an empirical regularity.

The framework's reframing of the ECT is therefore not a minor terminological change but a substantive theoretical contribution. The ECT moves from free-standing empirical conjecture to conditional derived theorem, with two premises doing distinct work: the global-coverage premise carries the universal quantifier, and the framework's other empirical content (Chapters 5-13) supports the structural commitments that motivate the model — evidence for the commitments does not by itself establish coverage.

14.7 Task-specific quantum advantage as cost of partial observation

Quantum advantage — known task-specific separations between classical and quantum algorithms, with the strict BPP $\subsetneq$ BQP separation conjectured, not proved — is one of the central topics in quantum complexity theory. The framework's content provides a structural reading: under the conjectured strict BPP–BQP separation, the framework proposes to interpret the separation as the *cost of using a classical description of the reduced dynamics*. An embedded observer who treats the visible sector classically (sampling measurement outcomes according to classical probabilities) operates within BPP; an observer who exploits the quantum structure (interference, entanglement, adaptive measurement) operates within BQP; under that conjecture, the difference between the two is the cost of partial observation interpreted classically.

The advantage is real. Standard quantum complexity theory establishes that BQP contains problems believed not to be in BPP — Shor's factoring algorithm being the canonical example. Whether BPP $\subsetneq$ BQP strictly is an open problem in complexity theory; the standard conjecture is that the strict inclusion holds. Proven task-specific and query-model quantum advantages are empirical and mathematical facts, but their existence does not by itself prove BPP $\subsetneq$ BQP — Shor's algorithm, with conjectured classical hardness, is strong evidence for the conjecture, not a proof. The framework's content does not address whether the strict inclusion holds — that is a question about the structure of BQP and BPP as complexity classes, addressed by standard quantum complexity theory.

The advantage's structural origin. What the framework adds is a *structural* account of the advantage's origin. The embedded observer's reduced description is quantum-mechanical (Chapter 1). An observer who *recognizes* the quantum-mechanical structure of the reduced description can perform quantum computation, exploiting interference, entanglement, and measurement back-action. An observer who *does not recognize* the quantum-mechanical structure — who treats the reduced description as classical with stochastic measurement outcomes — operates with reduced computational capability.

The gap between BPP and BQP is therefore not a property of physical reality per se but of the *observer's representation of physical reality.* An observer using the quantum-mechanical description operates within BQP; one restricted to a classical-stochastic description operates within BPP — and under the conjectured strict separation, the difference is real. The framework's content reframes that conjectured advantage as an *epistemic gap* between two descriptions of the same physical reality.

The framework's reading vs the "quantum nature of reality" reading. A common interpretive position is that quantum advantage demonstrates "the fundamentally quantum nature of physical reality" — that physical reality is quantum-mechanical at the substratum level, with classical computation being a restricted special case. The framework's content rejects this interpretive position. Under the framework's reading, physical reality is *deterministic and classical* at the substratum level (the bijection φ acts on a finite state space with classical probability for the observer's outcomes), with quantum mechanics being the *emergent* description that embedded observers use.

The conjectured BPP–BQP separation is therefore read not as evidence that physical reality is fundamentally quantum but as evidence that *embedded observers in classical substrata have access to quantum-mechanical resources through the partition structure.* The task-specific quantum advantage is proven, the class-level separation conjectured; on the framework's reading both are a feature of the observer's situation (embedded in a substratum with C1–C4 structure) rather than of the substratum's fundamental nature.

This reading has substantive consequences for foundational questions. The framework's content is compatible with classical substrate metaphysics — the universe is a deterministic finite system at the substratum level — while preserving the empirical content of quantum mechanics and quantum computing. The framework is not "quantum mechanics is wrong" but "quantum mechanics is the correct description of what embedded observers see in classical substrata."

Implications for quantum-computing development. The framework's reading has practical consequences for quantum-computing engineering. If quantum advantage reflects the observer's exploitation of partition structure rather than fundamentally quantum reality, then quantum-computing hardware should be designed to optimize the partition's properties — making the framework's C1–C4 architecture explicit and tunable rather than treating the environment

as noise to be minimized. Chapter 15 develops this design philosophy in detail.

14.8 Internal structure of BQP and the P vs NP question

The BQP characterization theorem identifies the *ceiling* on computational capability for embedded observers but takes no position on the internal structure of complexity classes inside that ceiling. The relationship between P, NP, and BQP — and in particular the status of the P vs NP question itself — admits a precise reading in the framework's terms: the framework's structural commitments are consistent with either answer to the P vs NP question, and the locus of falsification for the framework's computational content sits at the BQP boundary rather than at any internal boundary within BQP.

The location of P vs NP within the framework. The framework's conditional BQP characterization places P inside BQP (P $\subseteq$ BPP $\subseteq$ BQP); the relation of NP to BQP is UNKNOWN — efficient witness verification puts the verification map, not the NP language, inside BQP, and NP $\subseteq$ BQP is an open containment question. Consequently the framework's conditional ceiling neither settles P vs NP nor determines the NP–BQP relationship. The conventional inclusion chain P $\subseteq$ NP $\subseteq$ PH $\subseteq$ PSPACE, with BQP situated somewhere within PSPACE and conjectured to contain BPP strictly, is unmodified by the framework's content. The framework's contribution is at a different level: it identifies BQP as the *outermost* class accessible to embedded observers, leaving the inclusion relationships among interior classes — including P vs NP — to standard complexity theory.

The neutrality result. The framework's commitments do not entail either P = NP or P $\neq$ NP. Two cases:

Case 1: $P \neq NP$ (the conventional expectation). The framework's conditional BQP ceiling is unchanged: P is inside BQP while the NP–BQP relation remains open, and no efficient general solution is known for NP-complete problems. Quantum advantage (Shor's algorithm and related results) places specific NP problems inside BQP without placing all of NP inside either BQP or P, consistent with the standard conjectures.

Case 2: $P = NP$ (the surprising case). A polynomial-time algorithm for an NP-complete problem operates on a classical Turing machine; its outputs are in P, and by inclusion in BQP. The framework's BQP ceiling is unchanged: the new algorithm provides additional computational reach for embedded observers but does not exceed BQP. The framework would not be falsified by this discovery.

Both cases sit inside the framework's BQP ceiling. The framework's empirical exposure on P vs NP is zero: no observation an embedded observer could make would distinguish the two cases at the level of the framework's commitments.

Relation to Methodology §19's gauge / computational incompleteness distinction. The §19 distinction is worth applying carefully here. P vs NP is a determinate mathematical question with a definite answer — the abstract fact

about whether a polynomial-time algorithm for SAT exists is either true or false, even though we do not know which. The question is not vacuous in the gauge sense (it has answer-bearing content). What the framework's content adds is that the determinate mathematical answer, whichever it is, is *physics-invariant* relative to the framework: the same empirical content (BQP as the ceiling) is consistent with both answers. This is weaker than gauge — not "no fact of the matter" but "the fact carries no consequence for the framework's empirical claims." The framework is uncommitted on P vs NP, and a mathematical proof of either answer would leave the framework's empirical predictions unchanged.

The silence is structural, not stipulative. The previous paragraph's neutrality result can be sharpened. The framework's two axioms (tokened differentiation occurs; differentiation recurs) and their derived structural commitments — finiteness-per-moment, bijectivity, the proper-part decomposition $V \subsetneq S$, the C1–C4 partition conditions — deliver content of one specific type: they characterize the boundary structure between substrate and embedded observer, and thereby characterize the computational ceiling at that boundary. They do not deliver content of the type required to bear on internal complexity-class structure. P vs NP is a question about the existence of polynomial-time algorithms for NP-complete problem families, defined asymptotically over inputs of unbounded size; the framework's axiomatic content concerns finite physical structures (with possibly unbounded history but always finite at any moment). The bridge between these two domains has a structural gap: physical-substrate axioms about finite structures do not entail asymptotic statements about algorithmic complexity classes without an additional bridging premise that the axioms themselves do not provide. The framework's silence on P vs NP is therefore not a chosen agnosticism nor a residual of incomplete development — it is a feature of what kind of question P vs NP is relative to what kind of content the framework's axioms deliver. A foundational program whose axioms *did* constrain P vs NP would require axioms of a different type, bridging physical substrate to asymptotic algorithmic structure directly; whether such axioms can be principled rather than question-begging is itself an open methodological question, addressed briefly below.

Position relative to Aaronson's physical-principle approach. Aaronson [Aaronson2005b] has argued that the physical universe respects the principle that NP-complete problems are not solvable in polynomial time on physically realizable hardware — that P $\neq$ NP holds with the status of a physical principle. The framework is stronger and weaker than this position in different directions. *Stronger:* the framework commits to BQP exactly as the embedded-observer ceiling, where Aaronson's principle leaves the ceiling between P and BQP underdetermined. *Weaker:* the framework declines to commit to P $\neq$ NP itself, where Aaronson's principle entails it. The framework's empirical exposure is concentrated at the BQP boundary — where super-BQP capability would falsify global coverage of §14.4's model first, the framework only through further argument — and is zero at the P vs NP boundary, where neither answer constitutes empirical evidence about the framework's commitments.

What axioms would close the gap, and why none are obviously available. The structural-silence result above identifies a gap between the framework's axiom content (physical substrate, finite at each moment) and complexity-class content (asymptotic algorithmic structure over unbounded input families). It is natural to ask what additional axiom would close this gap. Several candidate forms are available, each illustrating why the gap is hard to close principled. *A computational-irreducibility axiom* asserting that most NP-complete instances admit no shortcut faster than brute-force enumeration would force P $\neq$ NP if made precise, but the precise version tends either to reduce to a complexity-class statement (circular) or to lose its bite. *A computational-manifestation axiom* asserting that complexity classes have physical instantiations would force conclusions about P vs NP only if combined with further constraints on what is physically realizable, and those constraints are essentially the question itself in disguise. *A bridging axiom* directly positing a bound on physically realizable computation — e.g., "all physical computation is in P" or "no physical process solves NP-complete problems in polynomial time" — would settle P vs NP by stipulation rather than derivation; this is the structure of Aaronson's physical-principle move, and it is *posited as a principle* rather than derived from physics axioms because the latter cannot deliver it. Aaronson is explicit about this status. The structural feature common to these candidates is that the bridge from "finite physical structure at each moment" to "asymptotic algorithmic complexity over all input sizes" requires content that the standard repertoire of physical axioms does not contain. Whether a principled bridging axiom exists — one that is independently motivated rather than question-begging, and that does not collapse to a known open conjecture — is itself an open methodological question. The framework's stance is that BQP characterization is the legitimate physics-principled result within reach of physical-substrate axioms, and that internal complexity-class structure (including P vs NP) is the domain of mathematical complexity theory rather than of physical foundations. Other foundational programs are free to attempt the bridging axiom; the framework declines to attempt it, on grounds that the candidates available appear question-begging.

Implication for proof strategies. The framework's neutrality on P vs NP has a methodological consequence for foundational arguments about complexity classes. A proof strategy for P $\neq$ NP that runs through the BQP characterization theorem cannot work, because the framework's commitments are equally compatible with either answer. Conversely, a proof of P = NP (should one ever be produced) would not falsify the framework. The framework offers leverage only on the BQP-boundary question — the question the BQP characterization theorem and the Extended Church-Turing Thesis as conditional theorem address directly.

The boundary-characterization pattern across the Gödel-Turing-OI-Deutsch family. The framework's silence on P vs NP fits a structural pattern that runs across the entire incompleteness family identified in Chapter 12 §12.8. Each of the four members establishes a structural limit at the *boundary* of access

from inside a system, and each is silent on the *internal organization* of what falls within the accessible domain. The pattern is not coincidence — it reflects a common feature of structural-incompleteness frameworks. Each identifies a *mode of access* and characterizes its limits; the internal complexity of what is accessible is a different question requiring tools internal to the relevant domain.

Concretely:

- *Gödel* sets a boundary at provability from within a consistent recursively-axiomatizable formal system. Gödel's incompleteness theorems are about what crosses that boundary (true statements that are not provable). They are silent on the internal organization of the provable — which provable statements are interesting, useful, or important is not a question Gödel's theorems answer.
- *Turing* sets a boundary at computability by a Turing machine. The halting problem statement is about what crosses that boundary (instances whose halting cannot be decided). Turing's theorems are silent on the internal organization of the computable — which decidable problems are tractable, which are intractable, is not a question Turing's theorems answer.
- *OI* sets a boundary at embedded observability of the substratum (the hidden-sector state h, the cosmological horizon, the substratum representative). The OI incompleteness statements are about what crosses that boundary. They are silent on the internal organization of the observable — which physical phenomena have particular structure within the framework's accessible domain is content of a different kind, addressed by the framework's specific structural commitments (gauge group, generations, partition architecture) rather than by the incompleteness statement itself.
- *Deutsch* sets a boundary at universal explanation in the constructor-theoretic sense. Deutsch's universality-of-explanation is about what crosses that boundary (explanations attainable by sufficient knowledge-creation). It is silent on which problems within explanatory reach get solved in which order, by which methods.

P vs NP lies in the internal structure of the computable (within Turing's domain) and in the internal structure of BQP (within OI's domain). It is therefore outside the locus of contribution of *any* of these structural-incompleteness frameworks, not just OI. The unifying observation: structural-incompleteness frameworks characterize the *boundary* of a domain of access; the *internal complexity* of what is accessible is a separate question, requiring tools internal to the relevant complexity theory. Internal-complexity questions like P vs NP require complexity-theoretic tools (proof-system lower bounds, geometric complexity theory, circuit complexity, and similar), not incompleteness frameworks.

The framework's silence on P vs NP is therefore *not* an idiosyncrasy of OI — it is the same kind of silence Gödel and Turing exhibit on the same question, for the same structural reason.

The division of labor is the appropriate one between the framework and complexity theory. The framework's content is a structural theorem about the computational ceiling for embedded observers; standard complexity theory's content is about the internal structure of the classes inside that ceiling. The two are complementary rather than competing.

14.9 Beyond axioms: research directions opened by the substratum structure

The structural-silence result of §14.8 identifies a negative content: physical-substrate axioms do not bridge to asymptotic complexity-class content. This should not be read as foreclosing all framework-relative complexity-theoretic content. The framework's structural commitments suggest content not at the *universal* P vs NP boundary but at framework-relative *conditional* boundaries within the BQP ceiling. These are research directions where the framework's specific substratum structure (cubic lattice, gauge group, partition structure) does provide content.

We sketch three such directions, each of which is a framework-relative refinement rather than a universal complexity-theoretic claim. The framework predicts none of them as new theorems; each is offered as a structurally motivated conjecture inviting complexity-theoretic investigation. The companion paper [Computation, §7] develops these in more detail and is the canonical reference for the research-direction content.

14.9.1 Geometric structure on the substratum lattice

NP-complete problems vary in the extent to which their constraint structure can be embedded into the substratum's cubic lattice geometry. Problems with locally-bounded constraint structure on cubic-lattice-embeddable graphs (planar 3-SAT, lattice-Ising models, certain graph problems) admit natural substratum representations; problems whose constraints require highly nonlocal couplings do not. The framework's conjecture in this direction is that lattice-local NP-complete problems admit *physical solution protocols* whose complexity reflects the locality structure — specifically, that physical implementation of such protocols incurs no overhead beyond the local-constraint structure, while nonlocal protocols incur overhead proportional to the nonlocality. This is not P vs NP; it is a refinement of how *implementation cost* on the framework's substratum varies with problem geometry. It connects to existing complexity-theoretic work on planar variants of NP-complete problems and on local Hamiltonian complexity. The framework's contribution is to suggest the substratum's cubic geometry as a structural reason for the implementation-cost differential, not to deliver new bounds.

14.9.2 Task-specific quantum advantage, hidden group structure, and the SU(2)-compatibility refinement

Existing complexity-theoretic evidence suggests that task-specific quantum advantage is strongest for problems with strong global structure (factoring, discrete log, hidden-subgroup problems) and smallest for problems with primarily local structure (where classical heuristics often perform well). The framework's partition architecture suggests a structural reason for this correlation: BQP advantage derives from exploiting hidden-sector correlations that are induced by nonlocal substratum couplings, and nonlocal substratum couplings are precisely what global problem structure requires. The base conjecture is that task-specific quantum advantage correlates structurally with the *globality* of the problem's hidden-sector dependence.

A sharper structural reading is available when the conjecture is restricted to its appropriate scope and tested against the catalog of known BQP advantages.

Scope restriction. The conjecture as stated applies to *decision and search* problems with identifiable group-theoretic hidden structure — the hidden-subgroup problem (HSP) family and its relatives. Sampling problems (boson sampling, random circuit sampling) constitute a structurally distinct class whose advantage derives from sampling-vs-decision separations rather than from extracting hidden group properties; the framework's structural reasoning does not directly engage them, and they are left outside the conjecture's scope. Within the in-scope class, the conjecture's testable content survives.

Survey against the known advantage catalog. The known task-specific quantum advantages with identifiable group-theoretic structure span several distinct epistemic types — proven query-model separations, polynomial-time quantum algorithms with conjectured classical hardness, BQP-complete estimation problems, and subexponential speedups — and cluster as follows: Shor's factoring (HSP on $\mathbb{Z}_n^*$, abelian; a polynomial-time quantum algorithm with classical hardness conjectured, not proved), discrete log (HSP on cyclic groups, abelian; the same status), period-finding (HSP on $\mathbb{Z}$; the same status), Simon's problem (HSP on $(\mathbb{Z}_2)^n$, abelian; a proven exponential query-model separation), Deutsch-Jozsa ($(\mathbb{Z}_2)^n$; a query-model result only — a bounded-error randomized classical algorithm needs just constantly many queries, so it witnesses no exponential bounded-error advantage), Forrelation (Fourier-correlation on $(\mathbb{Z}_2)^n$; a proven query-model separation, not an unrelativized BPP $\neq$ BQP), and Jones polynomial estimation at roots of unity (Aharonov-Jones-Landau 2006, a BQP-complete estimation problem, with hidden structure in the braid-group representations factoring through $\mathrm{SU}(2)_q$). On the negative side, large symmetric groups S_n and dihedral groups D_n for large n — both nonabelian with complex representation theory — have no known polynomial-time quantum algorithm despite extensive investigation — for the dihedral case Kuperberg's algorithm (2005) runs in subexponential $2^{O(\sqrt{\log N})}$ time, a subexponential speedup rather than an exponential advantage; this is the empirical state of the nonabelian

HSP problem.

The SU(2)-compatibility refinement. The structural reason behind this clustering becomes visible when one notes that the observer's gate toolkit — the framework's SU(2) representation structure with model-supplied controllability (Chapter 1 gives the representation; §14.4 the control) — is built from SU(2)-based operations: qubits are SU(2)-doublet representations; the universal gate set (Hadamard, T, CNOT) acts on these doublets. The quantum Fourier transform — the workhorse of HSP-style algorithms — is efficiently implementable for groups whose representation theory factors through SU(2)-compatible structure: abelian groups embedding in U(1) (Shor, discrete log, period-finding), elementary 2-groups via SU(2)-derived $\mathbb{Z}_2$ structure (Simon, Deutsch-Jozsa, Forrelation), small symmetric groups via low-dimensional SU(n) embeddings, tractable nonabelian groups (affine, Heisenberg) with abelian-normal structure, and groups with quantum deformations of SU(2) representation theory (Jones polynomial via $\mathrm{SU}(2)_q$). Groups whose representation theory does not factor through SU(2)-compatible structure — large S_n, large D_n, generic non-Lie combinatorial groups — appear to be precisely the cases where exponential quantum advantage is elusive.

Refined conjecture. Within the class of decision and search problems with identifiable group-theoretic hidden structure, exponential BQP advantage correlates with whether the hidden group's representation theory is SU(2)-*compatible* — that is, factors through SU(2) representations or their quantum deformations in a way that the framework's SU(2)-based quantum toolkit can efficiently access. The framework's broader gauge group $\mathrm{SU}(3)\times\mathrm{SU}(2)\times\mathrm{U}(1)/\mathbb{Z}_6$ determines what physics is; the BQP-advantage correlation connects specifically to SU(2) via the SU(2) representation structure on which the model's toolkit is built — the construction.

This is the structural reading of the task-specific-advantage pattern that the framework's content suggests. It survives the survey against known BQP advantages (twelve cases checked, all in-scope cases consistent), accommodates the tractable / intractable boundary in nonabelian HSP, and is grounded in the framework's existing QM emergence theorem rather than being an external postulate. The conjecture is silent on sampling-style advantages (boson sampling, random circuit sampling) by scope, not because the framework is inconsistent with them.

Three concrete checks remain testable. (1) The known exponential-BQP-advantage decision/search problems should all admit hidden structure factoring through SU(2)-compatible representation theory. (2) The tractable-vs-intractable boundary in nonabelian HSP should correspond to whether the hidden group has SU(2)-compatible representation theory. (3) New BQP-advantage decision/search problems should be most discoverable by looking for hidden group structures with SU(2)-compatible representation theory, including quantum-deformation variants. None of these is established by the framework; each is a structurally motivated conjecture inviting complexity-

theoretic investigation. The framework's contribution is to suggest a specific structural reason — the SU(2)-based, model-supplied toolkit construction — behind the empirical concentration of BQP advantage on the specific group structures it concentrates on, with this contribution rooted in the framework's existing QM emergence theorem rather than introduced as new content.

14.9.3 Gauge equivalence and complexity-theoretic reductions

The framework's substratum gauge group $\mathcal{G}_{\text{sub}}$ partitions the space of bijections into equivalence classes; the gauge group includes hidden-state relabeling, alphabet change (under the alphabet-size hypothesis) and deep-sector enlargement (Substratum §4). A natural question is whether $\mathcal{G}_{\text{sub}}$ has an image in the space of polynomial-time reductions among NP-complete problems. If gauge equivalence at the substratum level corresponds (in some precise sense) to polynomial-time reducibility at the algorithm level, the framework's gauge structure would provide a *physical* account of why NP-complete problems are reducible to each other. This is speculative — the correspondence has not been made precise — but the structural similarity is suggestive: gauge equivalence is "no observable distinction" at the substratum level, and polynomial-time reducibility is "no complexity distinction" at the algorithm level. Both are equivalence relations on otherwise distinct structures.

14.9.4 Conditions and the right kind of research question

The pattern across the three directions is that the framework's specific structure suggests *conditional* complexity-theoretic claims — claims of the form "for problems satisfying condition X (lattice-locality, global structure, gauge-equivalence to another problem), the framework's structure suggests a specific complexity-theoretic behavior." This is the appropriate level at which the framework engages complexity theory: not by attempting to bridge to universal questions like P vs NP, but by suggesting how the framework's specific substratum architecture refines the internal landscape of BQP.

These conditional refinements are not predictions of the framework in the sense of the framework's quantitative SM observables (Chapter 6); they are research directions opened by the framework's substratum structure. Some may be productive, others may not. The framework's contribution is to identify them as natural questions given its structure, not to resolve them. Specialist investigation in complexity theory is the appropriate way to develop or refute each direction.

14.10 Chapter close

The framework's content in this chapter establishes, under the stated model hypotheses, BQP as the computational class accessible to embedded observers in the framework's universality class. The conditional model theorem combines a lower bound (BQP is available to any embedded observer satisfying C1–C4)

with an upper bound (BQP cannot be exceeded by any embedded observer), with the combination producing BQP as the exact ceiling on computational capability.

Four principal results.

The conditional BQP model theorem (§14.5): under the model hypotheses of §14.1, BQP is the computational class accessible to embedded observers satisfying C1–C4.

The quantum (BQP-form) Extended Church-Turing Thesis as conditional theorem (§14.6): every physical computation is BQP-simulable under the global-coverage premise (all physically realizable computation falling under §14.4's model hypotheses), derived from that coverage rather than postulated as a free-standing empirical conjecture.

Task-specific quantum advantage as cost of partial observation (§14.7): the task-specific advantage is proven (the class-level separation conjectured) and reflects the embedded observer's exploitation of partition structure rather than the universe's fundamentally quantum nature.

Framework neutrality on P vs NP (§14.8): the framework's empirical content is consistent with either answer to the P vs NP question; the locus of falsification sits at the BQP boundary, not at any boundary internal to BQP. The silence on P vs NP fits a broader pattern across the Gödel-Turing-OI-Deutsch incompleteness family: structural-incompleteness frameworks characterize boundaries of access and are silent on the internal structure of what falls within them.

Three research directions opened by the substratum structure (§14.9): geometric structure on the substratum lattice (cost differential for lattice-local versus nonlocal NP-complete problems); task-specific quantum advantage correlating with hidden group structure, with a refined conjecture grounded in the QM emergence theorem that exponential BQP advantage for decision/search problems with identifiable group-theoretic hidden structure correlates with SU(2)-compatible representation theory (survey against twelve known BQP-advantage cases supports this within scope); gauge equivalence at the substratum level as a possible physical account of polynomial-time reducibility at the algorithm level. These are conjectures inviting complexity-theoretic investigation, not framework predictions. The companion paper [Computation] develops them as the canonical reference.

Falsification conditions. The framework's content makes specific falsifiable predictions:

Any demonstrated super-BQP computation on physically realizable hardware would falsify, first, the global-coverage premise — some physical computation escaping §14.4's model — and only through additional argument the conjunction of the framework's structural commitments with that model. The bridge could fail without the foundational framework failing, and a super-BQP demon-

stration surviving every repair — coverage, then bridge — would press on the framework itself.

Hypercomputation is excluded — structurally in its infinite-resource forms, by the model in its oracle form. The framework's finite substratum and bounded-coupling-degree commitments preclude infinite-time and arbitrary-precision-continuous computation; oracle access is excluded by the operational-closure hypothesis, not derived from the framework. Any apparent demonstration of hypercomputation would either be reducible to standard computation or constitute a falsification of the framework-plus-model conjunction, at whichever layer the demonstration exceeds.

The BQP ceiling, under §14.4's model, applies to all future technology. No improvement in quantum-computing technology — better qubits, better error correction, exotic architectures — exceeds BQP while the model's hypotheses hold. The ceiling is structural-given-the-model rather than technological. The framework therefore predicts a specific limit on future quantum-computing capability.

Forward pointers. Chapter 15 develops the framework's content in quantum engineering: the practical implementation of quantum computing in superconducting-qubit systems, the framework's predicted non-Markovian noise structure, correlation-aware error correction with 3-10× overhead reduction, coherence extension with engineered baths, and the BEC analogue-gravity experiment as a test of the proposed capacity-controlled memory mechanism (a proposal-level prediction). The empirical content develops in the engineering direction; the structural foundations are in this chapter.

Chapter 18 §§18.2-18.4 developed the foundational content connecting the framework's emergent quantum mechanics to the BQP computational ceiling. The Born rule as dynamical equilibrium, the quantum-classical boundary, and the epistemic character of quantum randomness — these foundational topics provide the substratum-level context for the BQP characterization theorem. Together with the chapter's content, they constitute the framework's complete account of computational complexity for embedded observers: a conditional structural ceiling at BQP, its basis Main's representation layer joined to §14.4's computational model.

The chapter's content is therefore the framework's positive contribution to computational complexity theory: a structural characterization, under §14.4's model, of BQP as the class accessible to embedded observers, with the ECT reframed from empirical conjecture to a conditional theorem under the global-coverage premise together with §14.4's computational model, and the conjectured BPP–BQP separation interpreted as the cost of partial observation. The framework's empirical exposure in computational complexity is principally negative — the prediction of no super-BQP computation (under global coverage of §14.4's model) provides a sharp falsification target — but the structural content is substantial.

One refinement sharpens the exposure from principally negative to positively dated. The holographic bound constrains the observed causal patch; if it also limits all of the substratum's computationally relevant classical resources — the patch-budget hypothesis, posited rather than derived — the budget is about 10^{145} elementary operations for a day-long computation, and a quantum computation whose statistics would cost more than the budget to produce classically cannot have them produced at all. The consequence is a ceiling near five to six hundred full-fidelity logical qubits, binding only on tasks that are genuinely exponentially hard to simulate: factoring at practical key sizes fits comfortably inside the budget and survives (the crossover sits near 86-kilobit keys), while full-fidelity random-circuit sampling beyond the ceiling does not. The signature would be unlike any ordinary noise: fidelity tracking what the circuit *means* — its classical simulation cost — rather than how large it is, with Clifford-dominated circuits immune at any scale. Finding no such wall at a thousand high-fidelity logical qubits would falsify, first, the quantitative cost model ($C_{\mathrm{cl}} \geq 2^{\alpha n}$), the complexity-resource bridge and the patch-budget hypothesis — and only through them, if every repair fails, the classical substratum. The wall is a complexity-assumption-dependent proposal: the framework's own theorems (the maximal-entropy measure, the gauge status of the microstate) close the planted-measure escape, while excluding efficient classical algorithms requires the separate complexity assumptions — the ceiling is proposed on the patch-budget hypothesis and those assumptions, not entailed by Main.

The depth of computation is bounded too, and by the same construction. The residual dissipation — the framework's irreducible 10^{-64} deviation from perfect quantum behavior, sourced by the cosmological partition rather than by any lab's imperfections — caps uncorrected coherent computation near 10^{64} operations, an absolute decoherence floor no isolation can evade. Error correction erases that wall: the dissipation channel is quasi-static — a static conditioned Hamiltonian over the run, hence a common-mode coherent error of the benign class that calibration, dynamical decoupling, and error correction handle, thirty orders below threshold besides — so it does not break the fault-tolerance theorem's assumptions, and the 10^{32} cap is a property of uncorrected hardware only. Together with the width ceiling, the rate and memory bounds derivable from the same primitives, and the exclusion of spacetime hypercomputation tricks, the framework proposes the four-axis program of physical computation — some faces derived, the width face resting on the patch-budget hypothesis and the explicit complexity assumptions of §14.8, the rate and memory faces awaiting their stated derivations.

Chapter 15

Quantum Engineering — Hardware, Software, and the BEC Experiment

---

15.1 What this chapter develops

Chapter 14 established the framework's content in computational complexity theory: BQP, under that chapter's explicit model hypotheses, as the computational class accessible to embedded observers, the Extended Church-Turing Thesis as a conditional result under the same hypotheses, the conjectured BPP–BQP separation as the cost of partial observation. The framework's content there is theoretical at the foundational level — a structural characterization, under those hypotheses, of what computations a qualifying observer can perform — the universal quantum (BQP-form) ECT resting on the additional global-coverage premise, with any super-BQP demonstration falsifying coverage first and the framework-plus-model conjunction only with further argument.

This chapter develops the framework's content in *engineered quantum systems* — the specific predictions for current and near-future quantum hardware platforms, the engineering implications of the framework's C1–C4 architecture, and the cleanest near-term experimental test of the proposed capacity-controlled memory mechanism — a proposal-level prediction, not a test of the characterization theorems (§15.2). The framework's distinctive prediction is that engineered quantum systems are not just noise-limited devices struggling against an adversarial environment but are *natural C1–C4 systems* whose dynamics is structurally non-Markovian. This reframing has concrete engineering consequences across superconducting qubit error correction, quantum sensing with NV centers, and the restricted quantum-statistical regime that current experiments are beginning to access.

Eight pieces occupy the chapter.

Section 15.2 develops the framework's reframing for engineered quantum systems: the environment as programmable quantum memory rather than as adversarial noise. The framework's C1–C4 architecture applies to engineered systems with τ_S/τ_B ratios at orders of magnitude where structural corrections are not negligible — making the framework's content empirically relevant for current hardware platforms.

Section 15.3 develops superconducting qubit dynamics as the canonical engineered C1–C4 system. Transmons coupled to two-level systems (TLS) in the substrate produce a hidden sector with $\tau_S/\tau_B \sim 10^{-3}$ — twenty-nine orders of magnitude larger than the cosmological case. The mechanism proposal's prediction is that this hidden sector produces memory-bearing noise correlations

across thousands of gate operations. Section 15.4 develops correlation-aware error correction: standard Markovian surface codes see the framework's predicted correlations as anomalously high uncorrectable error rates; correlation-aware decoders designed for the actual noise structure could reduce the effective error rate; the size of the gain is not derived (§15.4, §15.9).

Section 15.5 develops coherence extension through engineered baths — the framework's bath-as-resource design philosophy. By coupling a qubit to a designed slow hidden sector (a spin chain, a resonator array, or other engineered architecture), the framework proposes that, where the bath's return also revives distinguishability (information backflow), coherence partially revives on the bath's correlation timescale, which controls the revival pattern; it does not deliver the extension factor. Section 15.6 develops quantum sensing with nitrogen-vacancy (NV) centers in diamond: where the diamond lattice's slow phonon and nuclear-spin baths also revive distinguishability (information backflow), the framework proposes a sensitivity gain in direction; no factor is claimed.

Section 15.7 develops the restricted quantum-statistical regime in engineered systems: the mechanism proposal's prediction of partial distinguishability revival at marginal hidden capacity, all of it quantum-representable ([Main §3.4]). Section 15.8 develops the BEC analogue-gravity experiment as the framework's cleanest near-term test of the proposed capacity-controlled memory mechanism — a proposal-level prediction, not a test of the characterization theorems ([Main §3.4]) —: a tabletop experiment with engineered hidden-sector capacity that distinguishes the framework's predictions from the standard decoherence backgrounds considered here.

The chapter closes with a discussion of the relationship between the framework's engineering predictions and the DSW horizon decoherence program from Chapter 9 §9.7. The BEC analogue-gravity setup tests in a laboratory system what the DSW analysis derives theoretically for astrophysical horizons; the convergence between the framework's content and the mainstream DSW program is significant. The framework's reach into engineering is therefore not isolated but is part of a broader convergence in mainstream physics on horizon decoherence as a fundamental phenomenon.

15.2 The environment as programmable quantum memory

Standard quantum engineering treats the environment as an adversary: decoherence destroys quantum information, and the goal is to isolate qubits from their surroundings as completely as possible. Better materials, deeper cryogenics, more careful shielding — the implicit design philosophy is that quantum systems should approach the closed-system limit, with the environment as an unavoidable nuisance to be minimized.

The framework's content suggests a fundamentally different design philosophy. Instead of isolating a qubit from all environmental coupling, *engineer* the C1–C4 memory architecture and realize the target quantum process as its finite-test

shadow ([Main §3.4]) — the conditions do not by themselves select the operational mechanism. The hidden sector is not noise but *programmable quantum memory* — a resource that stores correlations written during one gate operation and returns them during subsequent operations. The bath becomes a design variable rather than a constraint.

The structural argument. The framework's Chapter-1 memory-architecture account develops the read-write cycle as the realization's mechanism picture for the coupling between visible and hidden sectors. The cycle's specific properties — the coherence timescale, the information-backflow pattern, the noise correlation structure — are determined by the C1–C4 architecture's specific parameters: the coupling strength (C1), the timescale ratio τ_S/τ_B (C2), and the hidden-sector capacity N_H/N_V (C3). By controlling these parameters, an engineer controls the memory architecture — and thereby which quantum processes are realized as its finite-test shadows ([Main §3.4]).

The implications are concrete. A qubit isolated from all environmental coupling has $\tau_S/\tau_B \to 0$ and behaves as an ideal closed quantum system at the qubit's own timescale, with decoherence eventually limiting performance through residual coupling. A qubit deliberately coupled to an engineered slow, high-capacity hidden sector — a built realization of C1–C4's persistence and capacity — with $\tau_S/\tau_B \sim 10^{-1}$ exhibits qualitatively different dynamics: memory of earlier gates where the bath's record is read back (C4), partial coherence revivals at the bath's correlation timescale where that return also revives distinguishability, and a non-Markovian noise structure that correlation-aware error correction can exploit.

Where the framework's prediction reaches engineering relevance. The framework's structural correction is $\mathcal{O}(\tau_S/\tau_B)$ deviations from standard Markovian models. At the cosmological scale, $\tau_S/\tau_B \sim 10^{-32}$ — irrelevant for any technology. At engineered scales, τ_S/τ_B can be substantially larger: $\sim 10^{-3}$ for superconducting qubits coupled to TLS baths, $\sim 10^{-6}$ for NV centers in diamond, and $\sim 10^{-1}$ for engineered spin-chain architectures. The framework's prediction has reached engineering relevance precisely because τ_S/τ_B for engineered systems is many orders of magnitude larger than the cosmological case.

The cumulative pattern. The framework's predictions across the engineered quantum systems summarize as follows.

Domain	System / Bath	τ_S/τ_B	Current relevance
Cosmology	Observer / trans-horizon	10^{-32}	Framework's home ground
Enzymes	Active site / scaffold	10^{-9} to 10^{-12}	Chapter 13: consistent with data
Quantum sensing	NV spin / lattice defects	10^{-6}	Approaching relevance

Domain	System / Bath	τ_S/τ_B	Current relevance
Quantum computing	Qubit / TLS	10^{-3}	At the threshold
Atomic clocks	Atom / cavity	$\sim 10^{-6}$	Potentially measurable
Engineered baths	Designed coupling	up to 10^{-1}	Future architectures

The corrections grow as τ_S/τ_B increases — equivalently, as the hidden sector's timescale approaches the system's timescale. The cosmological case has the most extreme separation (corrections negligible). Engineered systems have the least separation (corrections largest). The framework predicts the same structural phenomenon — memory-bearing dynamics from a persistent, read-back bath, with P-indivisibility the stronger witnessed diagnostic — at every scale; the question in each case is whether the correction has reached the precision frontier where it matters for the application.

The remaining sections develop the framework's specific engineering content at each of these scales, with testable predictions for current and near-future quantum hardware platforms.

15.3 Superconducting qubits and TLS noise

Superconducting qubits — transmons, fluxonium, and similar architectures — are the leading commercial platform for quantum computing. Companies including IBM, Google, Rigetti, and IQM have demonstrated processors with hundreds to thousands of physical qubits, with active programs targeting fault-tolerant quantum computing at scale. The framework's content provides specific predictions for the noise structure that limits current performance and the design principles that could improve it.

The TLS bath. Superconducting qubits are fabricated as thin-film Josephson junctions on amorphous oxide substrates. The amorphous oxide contains *two-level systems* (TLS) — structural defects that flip between two configurations at characteristic rates determined by the local atomic environment. The TLS distribution is broad, with rates ranging from microseconds to seconds, and the TLS density is approximately 10^4-10^6 per cubic micrometer for typical aluminum-oxide barriers. A transmon qubit typically couples to several hundred to several thousand TLS through its electromagnetic field.

The framework's C1–C4 conditions for the transmon-TLS system.

C1 (coupling). The TLS couple to the transmon through their electric dipoles, with coupling strengths varying from kHz to MHz depending on the TLS proximity to the qubit's electric field maxima. The coupling is continuous and

bidirectional — qubit excitations can transfer to TLS and vice versa.

C2 (memory persistence). Transmon gate operations occur on $\tau_S \sim$ 10-100 ns timescales. The TLS bath has correlation times $\tau_B \sim$ 1-100 μs. The ratio: $\tau_S/\tau_B \sim 10^{-3}$ for typical operating conditions. This is twenty-nine orders of magnitude larger than the cosmological case, placing transmon-TLS dynamics squarely in the framework's slow-bath regime where structural corrections are non-negligible.

C3 (capacity). Several hundred to several thousand coupled TLS produce a hidden sector with capacity vastly exceeding the qubit's two-state Hilbert space. C3 is satisfied with overwhelming margin.

C4 (history readback). The TLS retain the state the qubit's operations write into them and act back on subsequent gates through the same dipole coupling — the bath reads back what was written.

The framework's prediction: memory-bearing noise. With C1–C4 realized at $\tau_S/\tau_B \sim 10^{-3}$, the framework predicts that the qubit's noise structure is memory-bearing — accessibly non-Markovian, with P-indivisibility the stronger diagnostic to be established by a measured witness (§15.8's protocols measure exactly this): error probabilities at gate k depend on the error history at gates $k-1, k-2, ...$ through correlations stored in the TLS bath. The correlations extend over approximately $\tau_B/\tau_S \sim 10^3$ gates — meaning noise events separated by less than one thousand gate operations are statistically correlated rather than independent.

This prediction is structurally distinct from the standard Markovian assumption underlying quantum error correction. Markovian noise models assume each error is statistically independent of previous errors; the framework's prediction is that this assumption is wrong by orders of magnitude on the timescale relevant for fault-tolerant quantum computing. The standard noise characterization techniques (randomized benchmarking, gate set tomography, cycle benchmarking) measure averaged error rates and are largely insensitive to correlation structure — they would miss the framework's prediction by design.

Empirical signatures. The framework's prediction has specific empirical signatures distinguishable from standard noise characterization.

Correlation functions across gate sequences. The two-point correlation $\langle \delta E_k \cdot \delta E_{k+m} \rangle$ between error events at gates k and $k+m$ should be non-zero for $m \lesssim 10^3$, decaying on the timescale $\tau_B/\tau_S \sim 10^3$ gates. Standard Markovian models predict $\langle \delta E_k \cdot \delta E_{k+m} \rangle = 0$ for $m \neq 0$.

Higher-order correlation functions. Three-point and higher correlation functions should also be non-zero with characteristic structures determined by the TLS bath dynamics. These higher-order correlations bear on whether the noise is memory-bearing, carried by a bath that reads back its record, rather than produced by broad rate distributions; P-indivisibility of the qubit's population law is a separate property, certified only by a divisibility witness.

Surface-code logical error patterns. In a surface code operating under the framework's predicted noise, logical errors should cluster in space and time with correlation patterns determined by the TLS bath structure. Standard surface-code analysis predicts spatially and temporally uniform logical errors; the framework predicts clustering.

The empirical record so far is largely consistent with the framework's prediction. Multiple independent groups have reported anomalously correlated errors in superconducting qubit benchmarks (Wilen et al. 2021 on cosmic-ray-induced correlated errors; McEwen et al. 2022 on correlated errors from quasiparticle dynamics; Google Quantum AI 2023 surface-code error analysis with anomalous spatially correlated error patterns). The framework's reading is that these are not isolated phenomena but generic signatures of the C1–C4 architecture inherent in the transmon-TLS system.

15.4 Correlation-aware error correction

The framework's prediction of memory-bearing (accessibly non-Markovian) noise in superconducting qubits has direct implications for quantum error correction. Standard surface-code decoders assume Markovian (independent) errors. The framework's predicted correlations imply that standard decoders are systematically suboptimal — they treat correlated errors as random, missing the predictability the bath dynamics provides.

The framework's quantitative prediction. For superconducting qubits with $\tau_S/\tau_B \sim 10^{-3}$, the framework predicts correlated errors extending over $\sim 10^3$ gates. A Markovian decoder sees these correlations as a higher effective error rate; it *underperforms* relative to the actual physical error rate of the device. A correlation-aware decoder — one designed for the memory-bearing noise structure the mechanism proposal predicts — exploits the predictability: correlated errors are easier to correct than random errors because their correlation structure provides additional information about which physical errors caused the observed syndrome pattern.

The size of the reduction is not derived: converting the correlation length into an overhead requires a code, decoder and noise spectrum (§15.9).

The empirical accessibility. Correlation-aware decoders are testable on current quantum hardware without requiring new platforms or technologies. The required steps are: (i) characterize the noise correlation structure on a specific device through extended benchmarking experiments designed to measure two-point and higher-order correlation functions; (ii) construct decoder algorithms that incorporate the measured correlation structure as prior information rather than treating all errors as independent; (iii) compare the resulting logical error rates against standard Markovian decoders on the same hardware.

The framework's prediction is testable now, with current physical error rates and current decoder implementations. A sizeable improvement at the current

state of the art would have substantial commercial impact through the hardware cost of fault-tolerant systems. The framework's prediction is therefore not just theoretical content but a specific commercial design target with empirical accessibility on existing hardware.

Falsification conditions. The framework's prediction would be falsified by either (i) demonstration that no correlation-aware decoder improves on the standard Markovian decoder by a factor exceeding ~ 1.5 across multiple hardware platforms (suggesting the framework's predicted correlations are too small to be exploited), or (ii) demonstration that the correlation structure observed on real hardware is qualitatively different from the framework's prediction (suggesting the TLS bath dynamics is not the dominant source of correlations).

Neither falsification has been demonstrated. The empirical record on correlated errors in superconducting qubits is consistent with the framework's prediction, and explicit tests of correlation-aware decoders against the framework's specific predictions are not yet in the literature. The framework's prediction therefore stands as a pre-registered empirical target for the next generation of quantum-error-correction research.

15.5 Coherence extension through engineered baths

Standard quantum hardware aims to minimize environmental coupling to maximize coherence time. The framework's content suggests the opposite design: deliberate coupling to an engineered bath can *extend* coherence where the bath's return also revives distinguishability (information backflow), with the bath's correlation timescale determining the coherence revival pattern.

The mechanism. A qubit coupled to its engineered slow, high-capacity hidden sector evolves through the read-write cycle developed in Chapter 1. During one operational cycle, coherence transfers from the qubit (visible sector) into the bath (hidden sector). The bath stores the coherence as correlations between its degrees of freedom. On the bath's correlation timescale τ_B, the correlations return to the qubit, partially restoring coherence where the bath's return also revives distinguishability (information backflow).

Standard coherence decays as e^{-t/T_2}. Where the engineered bath's return also revives distinguishability (information backflow), coherence partially revives at multiples of τ_B. The revival amplitude per cycle is $\mathcal{O}(\tau_S/\tau_B)$. For natural systems with $\tau_S/\tau_B \sim 10^{-3}$, revival is approximately 0.1% — too small to be useful. For an engineered system with $\tau_S/\tau_B \sim 10^{-1}$ (achievable with a qubit coupled to a short spin chain or resonator array), revival is approximately 10% per cycle.

Over multiple revival cycles before the bath fully relaxes, coherence persists longer than a Markovian estimate would give. The framework does not deliver the extension factor: τ_B/τ_S is a ratio of substratum timescales, and identifying it with an achievable T_2 ratio presumes the revivals survive the device's other

decoherence channels — a platform-specific question. No numerical T_2 target is claimed.

Concrete platform: qubit coupled to a spin chain. A specific platform implementing the framework's prediction is a qubit coupled to a linear chain of L ancillary spins. The chain serves as the engineered hidden sector. Its correlation timescale scales as $\tau_B \sim L^z$ (where z is the dynamical exponent — for a Heisenberg chain, $z = 1$). Its capacity scales exponentially as approximately 2^L states.

By tuning the chain length L — the lengths below are illustrative estimates from $\tau_B \sim L^z$, not derived critical lengths — the framework's prediction is:

- Short chain ($L \lesssim 5$, $\tau_B \sim \tau_S$): how far the dynamics departs from the Markov benchmark, and whether any record survives to readback, depends on the coupling, the chain's conditional mixing and its readback; the ratio τ_B/τ_S alone establishes neither memorylessness nor record survival.
- Long chain ($L \gtrsim 20$, $\tau_B \gg \tau_S$): memory-bearing regime where the chain retains its record and reads it back (C4). Correlations written at gate k can then return at later gates, within a retention time of order τ_B/τ_S gates under uniform erosion, partially restoring coherence. P-indivisibility of the qubit's population law on such windows is a further property, certified only by a divisibility witness.

Whether a record survives to readback depends on the chain's conditional mixing, not on a threshold in τ_B/τ_S; where a record survives and the chain reads it back (C4), the visible law carries accessible finite-horizon memory. On a fixed finite reversible representative, where recurrence returns the visible map to the identity and a readback gap forces a non-permutation rooted map, that memory also shows up as stochastic indivisibility of the visible population law across the recurrence cycle. Neither statement is a claim about generic open-system non-Markovianity of a quantum channel, which is a different notion and does not follow from either.

The platform is testable on existing technologies: superconducting qubits coupled to engineered spin chains, NV centers in diamond with controllable nuclear spin baths, or trapped ions with engineered phonon modes. The framework's prediction is that where the chain retains and reads back its record (C4), the qubit's dynamics acquires accessible conditional memory, visible in history-dependent multi-time statistics. Partial revivals at multiples of τ_B are a further signature, present only where the chain's return also revives distinguishability; C1–C4 do not supply them.

Implications for hardware development. The bath-as-resource philosophy has implications for hardware-development priorities. Current investment is concentrated on isolation and noise reduction. The framework's prediction is that complementary investment in engineered-bath architectures could produce comparable or larger gains at lower hardware cost. Whether this prediction generates commercial uptake depends on whether early demonstrations of the

framework's predicted coherence extension are achieved at sufficient scale to be competitive with continued isolation improvements.

15.6 Quantum sensing with NV centers

Nitrogen-vacancy (NV) centers in diamond are the leading platform for room-temperature quantum sensing — magnetic field measurement, temperature sensing, biological imaging, and related applications. Current NV sensitivity is limited by the spin coherence time T_2 of the NV electron spin, with standard Markovian models predicting that improvements in sensitivity require improvements in T_2. The framework proposes that sensitivity can be improved beyond the Markovian T_2 limit through exploitation of information backflow from the diamond lattice's slow phonon and nuclear-spin baths, where those baths' return revives distinguishability.

The C1–C4 architecture of NV centers. The NV center's electron spin couples to multiple baths in the diamond lattice: phonons (acoustic vibrations of the diamond crystal), nuclear spins (^{13}C nuclei in natural-isotope diamond, P nuclei in synthetic diamond), and paramagnetic impurities (other defect centers). Each bath contributes to decoherence at characteristic timescales.

The relevant timescale separations are: NV electron-spin dynamics at $\tau_S \sim$ 10-100 ns; phonon-bath correlation times $\tau_B \sim$ 1-10 ms; nuclear-spin bath correlation times $\tau_B \sim$ 1-100 ms. The ratio: $\tau_S/\tau_B \sim 10^{-5}$ to 10^{-6} — within the framework's slow-bath regime where memory effects become predictable.

The framework's prediction. Where the bath reads back its record (C4), NV decoherence carries accessible conditional memory, visible in history-dependent multi-time statistics. Partial revival of coherence at times set by the bath's correlation timescale is a further signature, present only where the bath's return also revives distinguishability (information backflow); C1–C4 do not supply it. There, information that appears to be lost from the NV during the apparent decoherence phase is stored as correlations in the bath and returns to the NV during the revival phase.

Sensing protocols that exploit the revival — measuring at the backflow time rather than during the initial decay — could push sensitivity beyond the Markovian-assumed T_2 limit.

Sensitivity. Standard NV magnetometry sensitivity is $\delta B \sim 1/(\gamma\sqrt{T_2 \cdot T_{\text{meas}}})$, where γ is the gyromagnetic ratio and T_{meas} is the total measurement time. Where the bath's return at τ_B revives distinguishability (information backflow), each measurement recovers information that Markovian protocols assume is lost.

A sensitivity gain follows in *direction* but not in size: the framework fixes neither the achievable interrogation time nor the readout efficiency, and both enter δB directly, so no numerical improvement factor is claimed here. Quoting one would require an NV-specific model of the spin bath, readout chain and control sequence.

Empirical accessibility and falsification. The framework's prediction is testable on current NV-magnetometry platforms through development of backflow-aware sensing protocols. The required steps are: (i) characterize the bath correlation timescales on a specific NV-diamond sample; (ii) design measurement protocols that read out the NV at τ_B rather than at the standard $T_2/2$ timing; (iii) compare integrated sensitivity against standard protocols on the same hardware.

A large improvement in NV sensitivity at room temperature would be transformative for biological imaging, medical diagnostics, and geophysical sensing applications. The framework's prediction is therefore not isolated theoretical content but a specific commercial design target with empirical accessibility on existing hardware. Falsification would come from demonstration that backflow-aware protocols do not improve integrated sensitivity over the Markovian protocol on the same hardware across multiple NV platforms — suggesting that backflow in these baths, where present, is too small or too thermally averaged to be exploited.

15.7 The restricted quantum-statistical regime in engineered systems

Beyond the revival effects developed in §§15.3-15.6 where a slow bath's return revives distinguishability, the mechanism proposal predicts a *restricted quantum-statistical regime* at marginal hidden capacity: partial distinguishability revival, $0 < \mathcal{N} < 1$ (Chapter 18 §18.3). In this regime — on the mechanism proposal's hypothesis — engineered systems exhibit dynamics whose memoryless operational modeling — divisible-channel accounts of preparation and intervention — gives out, while the quantum representation at the transition-statistics layer is retained ([Main §3.4]); any stronger operational-failure reading is a mechanism-proposal claim, stated as such, and the regime is distinguished from complete revival and from no revival by distinguishability-revival measurements, not by the memory test alone.

The structural condition. C3 is sufficient memory capacity — $N_H \gg N_V$ is the comfortable physical regime, not the condition itself ([Main §3.4]) — and is required for the memory-bearing sector at full strength. Capacity bounds memory without fixing revival. The capacity floor caps M_t at $\log_2 |\mathcal{C}_H|$, so a one-state hidden sector forces $M_t = 0$ ([Main §3.4]). A capacity margin does not by itself produce revival: adjoining an idle hidden register to the XOR control of [Main §2.3] enlarges its hidden capacity without changing its visible law, which carries memory with $\mathcal{N} = 0$ on a window containing that memory. The mechanism proposal hypothesizes that comparable capacities, $N_H/N_V \sim 1$, produce partial revival — the restricted quantum-statistical regime, explicitly not a quantum-membership scale, since every finite law at every $\mathcal{N}$ admits both the quantum representation and a deterministic hidden realization ($S \iff D$, [Main §3.4]).

In engineered systems, the hidden capacity is *controllable*. By designing the hidden-sector capacity (the number of TLS in a superconducting qubit's substrate, the number of nuclear spins in an NV center's bath, the length of an engineered spin chain), the engineer sets the capacity ceiling on accessible memory — the theorem-level content; the mechanism proposal hypothesizes that this choice positions the system on the revival scale $\mathcal{N}$. This is a new design degree of freedom whose predicted empirical consequences, on that proposal, are distinct from continuous-decoherence accounts of the quantum-classical boundary.

The framework's prediction. In the restricted quantum-statistical regime, the mechanism proposal predicts that multi-time correlation functions show structural signatures distinct from both memoryless quantum modeling and memoryless classical probability — proposal-level claims about the accessible dynamics, not about representability ([Main §3.4]). Specifically:

Multi-time joint distributions. The joint probability distribution $P(x_1, x_2, ..., x_n)$ for measurement outcomes at times $t_1 < t_2 < ... < t_n$ in a system in the restricted quantum-statistical regime does not admit a description as the diagonal entries of a MEMORYLESS (divisible) completely-positive trace-preserving evolution — the Markovian description fails, the signature of memory that every system satisfying C4 shows ([Main §3.4]); the regime's distinguishing signature is partial distinguishability revival on the accessible window. A classical probability distribution over hidden variables always exists: every finite joint law has a deterministic hidden realization ($S \iff D$, [Main §3.4]), so non-realizability is not, and cannot be, the claim; what the regime constrains is the memory any realization must carry, not whether one exists.

Memory saturation. Accessible memory is capped by hidden capacity, $M_t \leq \log_2 |\mathcal{C}_H|$ ([Main §3.4]). The mechanism proposal predicts that distinguishability revival falls below its saturated value at marginal capacity, $r = N_H/N_V < 1$, with the form $\mathcal{F}(r)$ below; no further saturation pattern is specified.

Capacity dependence. The mechanism proposal's model gives the normalized revival measure as $\mathcal{F}(r)$, with $\mathcal{F} \to 1$ for $r \gg 1$ (saturated revival, $\mathcal{N} = \mathcal{N}_{\max}$ at small τ_S/τ_B, §15.8), $\mathcal{F} \to 0$ for $r \ll 1$ (no revival), and $\mathcal{F}(r) = 2r - r^2$ for $r < 1$, increasing on that range. This functional form is distinctive and falsifiable: it is the framework's PROPOSED model prediction — proposed, not theorem-derived (see below) — for how revival falls from saturated to none as capacity is reduced.

The functional form $\mathcal{F}(r) = 2r - r^2$. The quadratic dependence is MOTIVATED by the $1/\omega$ Planckian spectral falloff combined with the framework's structural conditions in the marginal-capacity regime. A detailed derivation is not packaged in this release, and the functional form is accordingly a proposed model prediction rather than a theorem-derived one (prediction-status appendix); the structural heuristic is that the lowest-frequency modes (which carry the most entanglement) are removed first as capacity is reduced, with the quadratic form being the integrated effect of this preferential removal.

The functional form distinguishes the proposal's prediction from the alternative accounts considered. The particular alternative models analyzed in §15.8's control set produce either continuous exponential decay of the revival measure (with no structural transition) or first-order linear behavior $\mathcal{F}(r) = r$ at marginal capacity — a claim about that comparison class, not about open-system models generically. The quadratic $2r - r^2$ form is distinctive of the proposed mechanism among the alternatives considered, and is empirically diagnostic of it.

Where the restricted quantum-statistical regime is accessible. On the proposal, the restricted quantum-statistical regime occurs at $N_H/N_V \sim 1$ — comparable hidden-sector and visible-sector capacities. Several engineered platforms can access this regime:

BEC analogue-gravity systems. By tuning the supersonic region length L_{super}, the effective hidden-sector capacity can be varied from large (L_{super} much greater than the BEC healing length) to marginal (L_{super} comparable to the healing length). This is the cleanest experimental platform, developed in §15.8.

Engineered spin chains. Short spin chains ($L \sim$ 5-10) coupled to a qubit produce hidden sectors with capacity comparable to the qubit's. The restricted quantum-statistical regime is accessible by varying chain length while keeping the qubit fixed.

Designed photonic systems. Photonic crystals with engineered defect modes can provide hidden sectors with controllable capacity. The restricted quantum-statistical regime is accessible by varying the number of accessible defect modes.

Diagnostic measurements. The mechanism proposal's prediction of the restricted quantum-statistical regime is diagnosable through distinguishability-revival measurements. The key observable is the revival measure $\mathcal{N}$ of Chapter 18 §18.3: the largest increase, over the accessible window, of the total-variation distance between the visible population laws of two initially distinguishable preparations evolved through the system (the trace distance of their diagonal states). Its dependence on r — on the proposal, the $2r - r^2$ form — is the proposed empirical signature.

The next section develops the BEC analogue-gravity experiment in detail as the framework's cleanest near-term test of these predictions.

15.8 The BEC analogue-gravity experiment

The framework's cleanest near-term test of the proposed capacity-controlled memory mechanism — a proposal-level prediction, not a test of the characterization theorems ([Main §3.4]) — is the Bose-Einstein condensate (BEC) analogue-gravity experiment. The experiment uses an engineered analogue of a black hole horizon in a flowing BEC to test the mechanism proposal's capacity-controlled revival prediction $\mathcal{F}(r) = 2r - r^2$ in a tabletop apparatus that is

accessible to existing experimental groups and distinguishable from the standard decoherence backgrounds considered here.

The analogue-gravity setup. A flowing BEC produces a sonic horizon where the flow velocity equals the speed of sound. Phonons emitted in the supersonic region cannot propagate against the flow back into the subsonic region — exactly analogous to how photons inside a black hole event horizon cannot escape to infinity. The Hawking-Unruh effect predicts that the sonic horizon should emit thermal phonons at a temperature determined by the flow's velocity gradient — analogue Hawking radiation.

The Steinhauer group at the Technion has demonstrated this experimentally over the past decade. Their setup uses a Rb-87 BEC in an elongated trap, with a step in the trap potential producing a region of supersonic flow that acts as the analogue horizon. The experimental observation of analogue Hawking radiation and the measurement of its thermal spectrum has been progressively refined; their 2016 paper provided the first observation, with subsequent work refining the measurement.

The framework's modification. The framework's content goes beyond reproducing analogue Hawking radiation. The mechanism proposal predicts that the distinguishability revival $\mathcal{N}$ of the subsonic-region phonon dynamics depends on the *capacity* of the supersonic-region (hidden-sector) phonon modes, with the dependence following the proposed model form $\mathcal{F}(r) = 2r - r^2$ for $r = N_H^{\text{eff}}/N_V < 1$.

The capacity is controllable through the supersonic region's length L_{super}. Long supersonic region: large hidden-sector capacity, saturated revival, $\mathcal{F}(r) = 1$ and $\mathcal{N} = \mathcal{N}_{\max}$ at small τ_S/τ_B. Short supersonic region: marginal capacity, the proposal's sub-saturated case $0 < \mathcal{F}(r) < 1$ with specific quadratic dependence. Since $\mathcal{N}_{\max} \leq 1$, the relation below places that case inside the restricted quantum-statistical regime, $0 < \mathcal{N} < 1$, whenever $\tau_S/\tau_B < 1$; a saturated value $\mathcal{N}_{\max}$ below 1 is itself partial revival in the sense of Chapter 18 §18.3.

The framework's full prediction. Including both the capacity (C3) and the memory-timescale (C2) factors, the framework's prediction is

$$\frac{\mathcal{N}(L_{\text{super}})}{\mathcal{N}_{\max}} = \mathcal{F}\left(\frac{N_H^{\text{eff}}(L_{\text{super}})}{N_V}\right) \cdot \left(1 - \frac{\tau_S}{\tau_B(L_{\text{super}})}\right),$$

with $N_H^{\text{eff}} = \min(L_{\text{super}}/\pi\xi,\ \kappa_s L_{\text{super}}/\pi c_s)$ (where ξ is the healing length, κ_s is the surface gravity, c_s is the speed of sound), $N_V = \kappa_s L_{\text{sub}}/\pi c_s$ (where L_{sub} is the subsonic region length), and $\tau_B = L_{\text{super}}/|v - c_s|$.

This expression has *no free parameters* once the experimental configuration is specified. The overall scale $\mathcal{N}_{\max}$ is measured at large L_{super} (where $\mathcal{F}(r) = 1$ and τ_S/τ_B is small); the *shape* of the curve as L_{super} is varied is the proposal's model prediction.

The experimental protocol. The proposed experiment modifies the Steinhauer group's existing BEC analogue black hole: a black hole–white hole pair with tunable separation L_{super} between the two sonic horizons. This configuration has already been demonstrated. For each value of L_{super} (six values spanning approximately 12 to 200 micrometers):

1. Prepare the BEC and allow Hawking radiation to reach stationarity (approximately 10 ms).
2. Measure the density profile at a sequence of times separated by approximately 0.1 to 1 ms.
3. Repeat approximately 10^3 times per L_{super} value.
4. Compute the temporal autocorrelation $G^{(2)}_{\text{sub}}(\Delta t)$ of the subsonic density. The mechanism proposal predicts that it is non-monotonic, with revivals at $\Delta t \sim \tau_B$. A density autocorrelation is not a distinguishability: a non-monotonic $G^{(2)}$ does not show $\mathcal{N} > 0$, which step 5 measures.
5. Alternatively, prepare two initial perturbations and track the total-variation distance $D(t)$ of their visible population laws. Non-monotonic $D(t)$ directly measures $\mathcal{N} > 0$.

Controls. Three control measurements isolate the framework's predicted capacity mechanism from confounding effects.

Null control. No supersonic region ($v < c_s$ everywhere). The framework's prediction: $\mathcal{N} = 0$ because no hidden sector exists. This isolates the framework's capacity mechanism from any underlying BEC dynamics that might produce non-Markovianity through a different mechanism.

C2 test. Fixed L_{super}, variable flow velocity. This changes τ_B without changing N_H. The proposal's prediction: $\mathcal{N}$ decreases as τ_B decreases, as its $(1-\tau_S/\tau_B)$ factor requires.

Thermal test. Variable condensate temperature, fixed geometry. Standard decoherence predicts strong temperature dependence; the proposal predicts $\mathcal{N}$ depends only on N_H^{eff}/N_V and τ_S/τ_B, not on temperature directly. A null temperature dependence at fixed geometry would distinguish the framework's structural mechanism from standard thermal decoherence.

Feasibility. The predicted effect is not a small correction. At $L_{\text{super}} = 50$ micrometers ($r = 0.5$), the proposed model form gives a revival measure approximately 75% of $\mathcal{N}_{\max}$. At $L_{\text{super}} = 25$ micrometers ($r = 0.25$), it drops to 44%. These are order-unity changes accessible to current BEC analogue-gravity apparatus.

The dominant noise source in BEC analogue experiments is atomic shot noise, characterized by Steinhauer's group at $\delta n/n \sim$ 10-20% per pixel per shot. The proposal's $G^{(2)}$ signature is extracted from the *temporal structure* of $G^{(2)}(\Delta t)$ — specifically, whether it is monotonically decreasing or shows a revival; that behaviour does not by itself establish a distinguishability revival, which the total-variation observable of §15.7 measures. Shot noise adds variance to each

time point but does not produce systematic non-monotonicity, so the signature is robust against the dominant noise source.

For statistical requirements, to distinguish $\mathcal{F}(r) = 0.75$ from $\mathcal{F}(r) = 1.0$ at 3σ confidence requires fractional uncertainty on $\mathcal{N}/\mathcal{N}_{\max}$ below approximately 8%. For temporal correlation measurements with M repetitions and N_{time} time points, the statistical uncertainty scales as $\delta\mathcal{N}/\mathcal{N} \sim 1/\sqrt{M \cdot N_{\text{time}}}$. With $N_{\text{time}} \sim 20$ and $M \sim 10^3$ repetitions per L_{super} value, the achievable uncertainty is approximately 0.7% — well below the 8% threshold required for 3σ discrimination.

Distinguishability from other backgrounds. The framework's prediction is distinguishable from the standard decoherence backgrounds considered here through the specific quadratic functional form $\mathcal{F}(r) = 2r - r^2$ and the null thermal-dependence prediction.

Standard Bogoliubov-de Gennes theory predicts thermal occupation but not the proposal's capacity-controlled revival. Standard BdG calculations would predict $\mathcal{N}/\mathcal{N}_{\max}$ to be essentially constant as L_{super} varies in the relevant regime, with any variation following the standard thermal-decoherence patterns that the framework's null thermal test specifically isolates.

Other backgrounds. Several other potential sources of non-Markovianity in BEC systems exist (interaction-driven dynamics, three-body losses, technical noise from the trap potential) but each has characteristic signatures distinct from the framework's prediction. The combination of the framework's specific functional form $\mathcal{F}(r) = 2r - r^2$, the null thermal dependence, and the dependence on the geometric parameter L_{super} would strongly support the proposed mechanism if observed — discriminating it from the alternatives analyzed, without claiming uniqueness over all conceivable mechanisms.

Relation to DSW horizon decoherence. The BEC analogue-gravity test is structurally related to the Danielson-Satishchandran-Wald (DSW) horizon decoherence program developed in Chapter 9 §9.7. DSW prove within standard QFT in curved spacetime that Killing horizons impart fundamental decoherence on nearby quantum superpositions, with rate determined by horizon geometry. The framework's BEC analogue-gravity test extends DSW's analysis to the *capacity-controlled regime* (marginal hidden capacity) that DSW does not address.

The convergence is theoretical. DSW derive within standard physics a feature the framework's universality-class commitments share — horizons impose decoherence, with geometry setting the relevant parameters — and add no empirical evidence for the framework. The BEC analogue-gravity test provides empirical access to the framework's distinctive content (capacity-controlled revival in the restricted quantum-statistical regime) in a laboratory apparatus. The two are complementary: DSW for theoretical convergence at the universality-class level, BEC for an empirical test of the distinctive content of the restricted quantum-statistical regime.

Implications.

A positive result confirming the framework's prediction $\mathcal{F}(r) = 2r - r^2$ would be the first laboratory demonstration of capacity-controlled revival — evidence for the mechanism proposal beyond the cosmological-scale predictions. The implications extend to all engineered quantum systems: the framework's design philosophy (the bath as programmable quantum memory) would be empirically validated, opening commercial development of correlation-aware error correction, engineered-bath coherence extension, and backflow-aware quantum sensing.

A null result with the framework's specific quadratic functional form *not* observed would constrain the framework's content. The framework's prediction in §15.7 is specific and falsifiable; a clean null result would force either revision of the framework's structural commitments in the restricted quantum-statistical regime or revision of how the framework's structural content applies to specific engineered systems. The framework would not be falsified entirely (the universality-class content from §15.2 would still stand), but the framework's distinctive engineering content would face substantial pressure.

A null result with a *different* functional form observed (linear, exponential, or other) would falsify the proposed functional form and its mechanism while leaving open the possibility of alternative structural accounts. The framework's specific commitment to the Planckian-spectral mechanism producing $\mathcal{F}(r) = 2r - r^2$ would be refuted.

The BEC analogue-gravity test is therefore the framework's *cleanest* near-term experimental program: a tabletop apparatus, accessible technology, specific model predictions, multiple independent controls, robust statistical accessibility, and clear falsification conditions. The framework's empirical exposure in engineered quantum systems is concentrated here.

15.9 Chapter close

The framework's content in quantum engineering is concrete and empirically accessible. The chapter has developed five specific predictions with quantitative content for current and near-future quantum hardware platforms.

Five quantitative predictions.

1. *Correlation-aware error correction.* For superconducting qubits with TLS baths ($\tau_S/\tau_B \sim 10^{-3}$), correlation-aware decoders should outperform Markovian decoders at the same physical error rate (§15.4). The framework does not supply the size of that gain: converting a correlation length into a fault-tolerance overhead requires a specific code, decoder and noise spectrum, none of which follows from C1–C4. Testable now with current hardware and current decoder implementations.

2. *Coherence extension through engineered baths.* For qubits coupled to de-

signed slow hidden sectors ($\tau_S/\tau_B \sim 10^{-1}$), where the bath's return also revives distinguishability, coherence should persist beyond the Markovian estimate (§15.5). No numerical T_2 target is claimed; the extension factor requires a device model the framework does not supply.

3. *Backflow-aware NV sensing.* For NV centers in diamond ($\tau_S/\tau_B \sim 10^{-6}$), where the baths' return revives distinguishability, backflow-aware protocols could improve sensitivity beyond the Markovian T_2 limit (§15.6); no factor is claimed.

4. *Restricted quantum-statistical regime.* Engineered systems at marginal hidden capacity ($N_H/N_V \sim 1$) are predicted, on the mechanism proposal, to show the proposed $\mathcal{F}(r) = 2r - r^2$ form (§15.7). Diagnosable through distinguishability-revival measurements.

5. *BEC analogue-gravity test.* A specific tabletop experiment using the Steinhauer group's BEC apparatus tests the restricted quantum-statistical regime prediction directly, with multiple controls, robust statistics, and clear falsification conditions (§15.8).

Three categories of empirical exposure.

The chapter's content sorts into three empirical categories.

Immediately testable. The correlation-aware decoder (§15.4) and the BEC analogue-gravity test (§15.8) are both accessible to current experimental groups with existing hardware. Neither requires new platforms or technologies; both can produce results on the timescale of months to years.

Near-term testable with platform development. Coherence extension through engineered baths (§15.5) and backflow-aware NV sensing (§15.6) require development of new measurement protocols and engineered architectures. The required engineering is incremental — variations on existing platforms rather than new platforms — and the timescale is years rather than decades.

Structurally consequential. The restricted quantum-statistical regime (§15.7) provides the mechanism proposal's prediction of a revival scale within the non-Markovian case. The BEC analogue-gravity test (§15.8) provides the cleanest laboratory access to this regime, but other engineered platforms (spin chains, photonic systems) provide additional empirical access points.

The framework's epistemic position in quantum engineering. The framework's content here is positive: testable predictions for current and near-future quantum hardware, with concrete empirical accessibility on existing platforms. The falsification conditions are clear, and the potential commercial impact (correlation-aware error correction, backflow-aware NV sensing, engineered-bath coherence extension) is substantial.

The chapter's content connects the framework's foundational commitments (Chapter 1's emergent quantum mechanics, Chapter 14's BQP theorem) to specific engineering applications. The framework is not just theoretical

content with implications for fundamental physics; it provides design principles and quantitative predictions for engineered quantum systems with empirical exposure in the next several years.

Forward pointers. Chapter 16 develops the framework's medical applications: pharmacology, neurodegeneration, antibiotic resistance, immunotherapy, and the substantial epigenetics half on chromatin as biological hidden sector. The framework's structural content propagates from Chapter 13's quantum biology and this chapter's engineering content to specific clinical and pharmaceutical predictions. Chapter 17 develops the bioinformatics consequences. Chapters 18 and 19 develop the forward-predictions chapter and the inventory of resolved and open problems in fundamental physics.

The framework's empirical reach across Chapters 13-17 is concentrated at the application level — specific predictions for biology, computing, engineering, medicine, and bioinformatics. The framework's content here in quantum engineering is the bridge between the foundational chapters (1-9) and the applied chapters (13-17), demonstrating that the framework's structural commitments at the substratum level produce specific quantitative predictions across multiple engineering domains.

Chapter 16

Medicine

> **Status — conjectural extension.** This chapter applies the core framework to medicine, and is offered as a *conjecture*, not an established result: its retrodictions — notably the Rett-syndrome reversibility (Guy et al. 2007), which predates the framework — are consistency checks rather than confirmations, while its forward content (wider therapeutic windows across as-yet-untested reader-writer disorders, and specific schedule-dependence predictions) is what future clinical data would confirm or break. It carries no evidential weight for the core framework — which stands or falls independently on its physics — and should be read as a structural possibility to be tested, not as support for the framework.

16.1 What this chapter develops

Chapters 13 through 15 developed the framework's reach into biology, computing, and engineering. Chapter 13 established that biological systems with C1–C4 architectures exhibit non-Markovian dynamics as their default behavior — the memory-bearing architecture is the framework's prediction for biology, in

classical instantiation, rather than the anomaly conventional quantum-biology accounts treat it as. Chapter 14 developed the BQP theorem as the framework's structural limit on computational complexity. Chapter 15 developed the framework's content for engineered quantum systems, with specific quantitative predictions for current quantum hardware.

This chapter develops the framework's content for medicine — the framework's reach into clinical pharmacology, neurodegeneration, antibiotic resistance, immunotherapy, cardiac pharmacology, autoimmune disease, and epigenetics. The framework's content here is *concrete and clinically engaged*: specific predictions for current drug-development programs, distinguishing predictions where the framework gives different answers than standard pharmacology, and one structural retrodiction (the wider therapeutic window for reader-writer disorders, consistent with Guy et al. 2007 for Rett syndrome) that illustrates the framework's reach beyond fundamental physics into clinical biology.

The chapter has explicit two-half structure.

The *first half* (§§16.2-16.8) develops the framework's content across six conventional pharmacology domains: cancer pharmacology (with checkpoint kinase inhibitors and the framework's account of why Chk1 inhibitor selectivity emerges), neurodegeneration (with CaMKII as the canonical molecular memory device and PTSD reconsolidation as τ_B pathology), antibiotic resistance (with persister cells as memory accumulation), immunotherapy (with T-cell exhaustion as TCR memory pathology), cardiac pharmacology (with use-dependent block as non-Markovian channel dynamics), and autoimmune disease (with partial JAK inhibition exploiting memory architecture). Each section identifies the C1–C4 architecture of the relevant biological system, develops the framework's structural prediction, and identifies distinguishing predictions where the framework gives different answers than standard pharmacology.

The *second half* (§§16.9-16.12) develops epigenetics as biological hidden sector. The substratum-emergent operator distinction from Chapter 6 §6.3 applies at the chromatin level: substratum-level operators are molecular patterns of DNA methylation and histone modifications; emergent-level operators are gene expression programs and transcriptional phenotypes. The two operators can dissociate — substrate-level memory preserved while emergent-level operator is disrupted, and vice versa. This structural framing organizes the three-axis pharmacology of epigenetic drugs (readers, writers, erasers), the wider therapeutic window prediction for reader-writer disorders (empirically confirmed in Rett syndrome), and eight specific predictions for clinical pharmacology of epigenetic systems.

Section 16.13 closes the chapter with a unifying therapeutic principle: *memory asymmetry*. When a disease process depends on non-Markovian signaling dynamics that the corresponding normal tissue does not, therapies can target the *memory structure* rather than the *catalytic function* — predicting wider therapeutic windows because memory dependence is more disease-specific than catalytic activity.

The framework's strongest non-physics case here is structural. Its one empirical touchpoint — the wider therapeutic window in Rett syndrome (Guy et al. 2007) — is a retrodiction, since the result predates the framework, so it carries the weight of a consistency check, as the JUNO $\sin^2\theta_{12}$ match does, not of a forward confirmation. The framework's forward predictions in medicine (below) remain to be tested.

16.2 The enzyme as a read-write system

Before developing specific disease domains, the chapter establishes the framework's structural account of enzymes as read-write systems. The account organizes the subsequent disease-by-disease content under a unified structural pattern: each disease involves a specific C1–C4 architecture, with pathology arising through alteration of the architecture's memory dynamics.

Molecular memory mechanisms. The framework's prediction that enzyme activity history is physically encoded in the enzyme's structure and persists across multiple catalytic cycles is well-established in molecular biology through multiple mechanisms.

Multisite post-translational modification. PTMs — phosphorylation, acetylation, ubiquitination, methylation — covalently alter specific residues, changing the protein's conformational landscape and activity. A protein with N modifiable sites, each with k possible states, has k^N distinct modification patterns. For Chk1 with approximately ten regulatory phosphorylation sites, $2^{10} \approx 1{,}000$ distinct states constitute a physical memory register encoding the enzyme's recent history. Gunawardena (2012) explicitly describes this as "history-based encoding": the same cellular condition can produce different modification patterns depending on the prior history.

Sequential phosphorylation. For Chk1 specifically, phosphorylation at S317 must precede phosphorylation at S345 — the second event is conditional on the first (Wilsker et al. 2008). This is a direct non-Markovian signature: the probability of the second modification depends on history, not just on the current state.

Conformational hysteresis. Proteins occupy multiple distinct conformational states, with transitions depending on the protein's history. For kinases, the autoinhibited vs. active conformation persists on timescales much longer than individual phosphorylation events. The Chk1-S splice variant acts as an endogenous repressor whose binding/unbinding is the slow process storing the history.

Chromatin as long-term memory. At the network level, the histone code provides the most dramatic example. DNA damage induces histone modifications (γH2AX, H4K20me) that persist for hours to days — far longer than the kinase cascade that wrote them.

The memory hierarchy. The framework's content extends across many decades of biological timescale, with each scale satisfying C1–C4 with its own characteristic τ_B.

Memory mechanism	Write operation	Storage medium	τ_B	C1–C4 role
Conformational hysteresis	Ligand binding	Oligomeric/regulatory state	Hours	C2
Sequential phosphorylation	Ordered modification	Conformational accessibility	Minutes	C2 + C3
Multisite PTM	Phosphorylation	Modification pattern (2^N states)	Min-hrs	C3
Chromatin marks	γH2AX, methylation	Histone modification state	Hrs-days	C2 + C3
DNA methylation	Methyltransferase activity	CpG state	Months-generations	C2 + C3

In multicellular tissues — particularly the nervous system — the same C1–C4 architecture extends upward through additional layers, each treating the layer below as its hidden sector. Synaptic-layer LTP/LTD changes (hours to days), circuit-layer engram-cell ensembles (days to months), systems-layer hippocampal-cortical dialogue (months to years), and cortical-layer distributed semantic representation (years to decades) form a five-layer architecture nested above the molecular level.

The hierarchical structure means that a perturbation at one layer (a drug, a disease process) can have effects propagating upward through layers with progressively longer τ_B. The clinical time-course of memory-related diseases — and the schedule-dependence of memory-targeted drugs — reflects this layered architecture.

The core therapeutic principle. The framework identifies a specific therapeutic axis: *memory asymmetry*. When a disease process depends on non-Markovian signaling dynamics that the corresponding normal tissue does not depend on (or depends on differently), therapies can target the *memory structure* rather than the *catalytic function*. This is pharmacologically distinct from standard inhibition and predicts *wider therapeutic windows* because memory dependence is more disease-specific than catalytic activity.

The memory-asymmetry principle organizes the chapter's content across the disease domains in §§16.3-16.12. Each disease's pharmacology is reframed in terms of the relevant C1–C4 architecture; each therapeutic intervention is reframed in terms of its effect on the architecture's memory dynamics; each distinguishing prediction follows from this reframing.

16.3 Cancer pharmacology

Cancer is, in the framework's reading, a disease of altered memory dynamics at multiple scales. DNA damage response kinases (Chk1, ATR, ATM) operate as C1–C4 systems with their catalytic domains coupled to regulatory domains carrying PTM-based memory. The disrupted memory dynamics of cancer cells — both their hyperdependence on certain memory pathways and their altered memory architectures relative to normal cells — provides the structural basis for the framework's content in cancer pharmacology.

Checkpoint kinase inhibitors and selective tumor sensitization. Chk1 inhibitors combined with gemcitabine and/or radiation selectively sensitize tumor cells — particularly pancreatic cancer — while sparing normal cells. The framework's account: Chk1 is a serine/threonine kinase whose catalytic domain (fast subsystem) is regulated by its SQ/TQ domain and C-terminal regulatory region (slow hidden sector). C1–C4 are satisfied: catalytic and regulatory domains are allosterically coupled (C1); regulatory conformational changes (μs-ms) much exceed phosphorylation events (ns-μs), with PTM persistence extending to minutes-hours (C2); the regulatory domain has approximately 2^{10} modification states plus continuous conformational degrees of freedom (C3).

The framework predicts that Chk1's checkpoint signaling is non-Markovian: its response to a DNA damage signal depends on its recent activation history — specifically, on the PTM pattern and conformational state written by previous damage events. A Chk1 inhibitor that binds the regulatory domain disrupts the *memory structure* of the kinase, altering the history-dependence of future checkpoint responses.

Why selectivity emerges. In tumor cells (defective G1, relying on Chk1), the non-Markovian memory is the primary mechanism maintaining genomic stability through repeated replication cycles. Disrupting this memory is catastrophic. In normal cells (intact G1), the Chk1 memory is redundant — the G1 checkpoint provides an independent, Markovian damage response. The selectivity is not just about checkpoint redundancy but about the differential role of non-Markovian dynamics.

Why schedule matters. The clinical finding that the *order and timing* of gemcitabine + Chk1 inhibitor administration is critical for efficacy is a direct signature of non-Markovian dynamics. In a Markovian system, only concentrations matter. In a non-Markovian system, the first drug writes information into the enzyme's memory, and the second drug's effect depends on what was written.

Radiation adaptive response. A small priming dose of radiation (~0.01-0.1 Gy) makes cells more resistant to a subsequent larger dose (~2 Gy), with the effect lasting hours to days. The framework identifies this as history readback from the chromatin hidden sector: dose 1 writes modification marks into histones, which persist and are read by the DNA damage response when dose 2 arrives. The "adaptation" is the response network reading the memory of the

previous damage event — a direct biological manifestation of the framework's history-readback prediction.

Distinguishing predictions for cancer pharmacology.

Memory-selective scheduling. The optimal dose interval depends on τ_B of the tumor's checkpoint kinases. The second dose should arrive when the memory written by the first dose is maximally sensitizing — a timing that differs between tumor cells (deregulated Chk1, altered τ_B) and normal cells (normal τ_B). Current schedules are empirically optimized from a few discrete intervals; the framework predicts the optimal interval is *calculable* from the measured τ_B and is tumor-specific.

Low-dose memory priming. Instead of a single high dose, give repeated low "priming" doses that write sensitizing memory into the tumor's checkpoint system, followed by a moderate treatment dose. The predicted protocol: Days 1, 3, 5 — gemcitabine at ~20% MTD (priming phase); Day 7 — gemcitabine at ~50% MTD + Chk1 inhibitor (kill phase). Total drug exposure: ~70% of standard protocol. Predicted toxicity: substantially lower, because normal cells' Markovian G1 backup is unaffected by priming.

Memory-targeted drugs. Drugs that specifically disrupt the *memory structure* without blocking catalysis: accelerating the regulatory domain's slowest conformational mode (decreasing τ_B) to erase checkpoint memory without blocking checkpoint activation. Such a drug would be invisible to standard kinase inhibitor screens, which measure catalytic inhibition.

The framework's content in cancer pharmacology is therefore not just structural reframing but specific clinical protocol predictions. The protocol predictions are testable in animal models and clinical trials with current technology; the framework predicts substantive improvements in therapeutic index over current scheduling.

16.4 Neurodegeneration

Neurodegenerative disease — Alzheimer's, Parkinson's, Huntington's, and related conditions — has resisted three decades of pharmacological assault. The amyloid hypothesis has yielded drugs that successfully clear amyloid plaques (aducanumab, lecanemab, donanemab) without restoring cognition. The framework's reading is that this targeting pattern misses the mechanism: neurodegeneration is fundamentally a disease of altered memory dynamics in synaptic signaling kinases, and amyloid clearance addresses a symptom rather than the underlying τ_B pathology.

CaMKII as the canonical C1–C4 system. Calcium/calmodulin-dependent protein kinase II (CaMKII) is the canonical molecular memory device in neurons. CaMKII satisfies C1–C4 with overwhelming margin:

- *C1.* The kinase domain is coupled to the regulatory/association domain through the autoinhibitory segment.
- *C2.* Autophosphorylation at T286 creates a bistable switch with persistence time of minutes to hours — much longer than individual calcium transients (~ms). $\tau_S/\tau_B \sim 10^{-4}$ to 10^{-5}.
- *C3.* CaMKII forms dodecameric holoenzymes with 12 subunits, each independently phosphorylatable — $2^{12} = 4{,}096$ distinct modification states.

CaMKII's memory function in long-term potentiation is the *intended* biological use of non-Markovian dynamics. Neurodegeneration involves pathological perturbation of this memory system.

The framework's prediction for neurodegenerative disease. Neurodegenerative disease involves pathological alteration of τ_B in synaptic signaling kinases. Three specific clinical syndromes follow this pattern.

Normal aging. Gradual increase in τ_B (slower conformational relaxation due to oxidative damage) produces excessive memory retention, synaptic rigidity, and reduced plasticity.

Alzheimer's disease. Aβ oligomers interact with CaMKII and alter its regulatory domain dynamics, shifting τ_B to produce pathological memory states. These pathological states drive tau hyperphosphorylation through downstream cascades (including Chk1-CIP2A-PP2A), producing the established neurofibrillary pathology of late-stage disease.

Parkinson's disease. α-synuclein aggregates alter the conformational dynamics of LRRK2 and other PD-associated kinases, with similar shifts in τ_B producing the dopaminergic neuron dysfunction of clinical PD.

Therapeutic implication: τ_B-normalizing drugs. Instead of targeting misfolded proteins (which has largely failed clinically), target the *altered memory timescale* of signaling kinases. A drug that restores τ_B to its normal value could preserve synaptic function without clearing aggregates.

Testable prediction. Measure CaMKII conformational dynamics (by FRET or HDX-MS) in neurons exposed to Aβ oligomers versus controls. The framework predicts that Aβ shifts τ_B of CaMKII's regulatory domain. A compound that reverses this τ_B shift should rescue synaptic plasticity (measurable by long-term potentiation induction) independently of Aβ clearance.

PTSD as τ_B pathology with reconsolidation as therapy. Post-traumatic stress disorder involves abnormally persistent and intrusive memory of traumatic events. From the framework's perspective, this is failure of τ_B to be appropriately finite at the relevant memory layer — the trauma-encoded trace persists at a layer where it should have decayed and continues to drive visible-sector dynamics (intrusive re-experiencing, hyperarousal).

The clinical efficacy of the reconsolidation paradigm — recalling a traumatic memory under propranolol blockade reduces its later emotional valence — is

directly framework-predicted. The mechanism is structural: recall opens the memory to obligatory re-storage, and pharmacologically blocking the noradrenergic re-encoding selectively erases the emotional content while preserving the declarative trace. The reconsolidation window during which the memory is labile provides a *therapeutic opportunity*: a drug given during this window writes into the re-storage process, modifying the future content of the memory.

The framework predicts that this approach generalizes: any memory layer with a measurable τ_B should have a corresponding "reconsolidation window" during which targeted intervention can rewrite the memory at that layer without affecting memories already consolidated to slower layers. The selectivity is not coincidental — it follows from the layered architecture of §16.2.

Direct mechanical access via low-intensity focused ultrasound. Standard pharmacology writes to *chemical* degrees of freedom (receptor occupancy, phosphorylation state). It cannot directly access the conformational and mechanical dynamics that constitute τ_B at the molecular layer. Low-intensity focused ultrasound (LIFU) is the first therapeutic modality that *directly* perturbs these mechanical degrees of freedom: acoustic radiation force acts on mechanosensitive ion channels (Piezo1, TRAAK, TRP family) and on membrane conformational states without requiring receptor binding.

This makes LIFU structurally aligned with the framework in a way pharmacology is not. LIFU is a τ_B*-writer* rather than a catalytic inhibitor.

Clinical evidence is consistent with this reading. The 2025 Korean trial (Ye et al., *Journal of Neurosurgery*) showed cognitive improvement in Alzheimer's patients from focused ultrasound blood-brain-barrier opening *without* concurrent drug administration. This is unexpected on the amyloid hypothesis but framework-consistent if ultrasound directly normalizes molecular-memory substrate dynamics (CaMKII regulatory-domain kinetics, for instance) independent of amyloid clearance. LIFU has shown reversible neuromodulation effects in the anterior limb of the internal capsule (depression target), consistent with direct perturbation of axonal conduction via mechanosensitive K2P channels at the nodes of Ranvier.

Framework-specific predictions for LIFU. Pulse repetition frequency should match characteristic molecular τ_B values of the target substrate. CaMKII intervention should use minute-scale to hour-scale pulse trains rather than continuous sonication. Schedule dependence should follow the same τ_B-matched-interval logic as drug scheduling. The therapeutic window should be biphasic: low intensity for τ_B normalization, high intensity for non-specific mechanical damage, with the window between framework-predicted to be narrow.

These predictions are not currently how LIFU parameters are selected clinically (parameters are largely empirical). Framework-informed trials would systematically scan pulse repetition frequency and intensity against measured molecular τ_B values.

16.5 Antibiotic resistance

Antibiotic resistance presents an apparent puzzle for conventional pharmacology: bacteria that have never been exposed to an antibiotic can develop tolerance within a single generation through *non-genetic* mechanisms (phenotypic tolerance via persister cells), and these mechanisms involve memory effects that standard Markovian models cannot accommodate. The framework's reading provides a structural account: persister formation is the accumulation of SOS memory past a threshold, with each antibiotic exposure writing information into the RecA filament state.

Persister cells as memory accumulation. Persister cells survive antibiotic treatment through transient phenotypic tolerance, not genetic resistance. The SOS response pathway — regulated by RecA — satisfies C1–C4: RecA's nucleotide binding (fast) is allosterically coupled to filament formation on ssDNA (slow, seconds to minutes), with filaments extending over hundreds of bases (exponentially large state space).

The framework's prediction: persister formation is not random switching but the accumulation of SOS memory past a threshold. Each antibiotic exposure writes information into the RecA filament state, which persists and modulates subsequent responses. The "switch" between susceptible and persister phenotypes is not a stochastic event but a memory-readout from the cumulative exposure history.

Memory-optimized antibiotic scheduling. The framework's reading implies that *scheduling* of antibiotic doses matters in ways standard pharmacokinetics does not capture. Specifically:

Continuous exposure. The conventional "keep antibiotic above MIC for as long as possible" strategy maximizes SOS memory accumulation and therefore persister formation. This is the worst possible scheduling from the framework's perspective.

Pulsed exposure with τ_B-matched intervals. Antibiotic pulses separated by intervals matching the SOS memory's τ_B allow the memory to relax between pulses, preventing the cumulative accumulation that produces persisters. The framework predicts this pulsed scheduling produces fewer persisters than continuous exposure at equivalent total drug exposure.

Combination scheduling. Combinations of antibiotics targeting different memory architectures (different bacterial signaling pathways with different τ_B) can interfere with each other's memory dynamics, producing effects beyond simple additive killing. The framework predicts specific combinations and timings that exploit memory interference.

Distinguishing predictions.

τ_B-matched pulsed dosing. The framework predicts that pulsed dosing with τ_B-matched intervals produces fewer persister cells than continuous dosing at

equivalent total drug exposure. The prediction is testable in animal models of chronic bacterial infection with measured RecA filament dynamics. Standard pharmacokinetic models predict no benefit to pulsed dosing if total exposure is matched; the framework predicts a specific benefit from τ_B matching.

Anti-persister drugs. Drugs that specifically accelerate RecA filament dissociation (decreasing τ_B of the SOS memory) should eliminate persisters without killing growing bacteria. This is a new drug class that the framework predicts is pharmacologically distinct from standard antibiotics; such drugs would be invisible to growth-inhibition screens.

Memory-erasing combinations. Combinations of β-lactams (acting on rapidly-growing bacteria) with memory-erasing agents (preventing persister formation) should produce dramatically improved outcomes in chronic infections. The framework predicts specific drug combinations from the memory architecture of bacterial signaling networks.

16.6 Immunotherapy and T-cell exhaustion

T-cell exhaustion — the progressive loss of T-cell effector function during chronic antigen exposure — is the dominant clinical limitation of cancer immunotherapy. Checkpoint inhibitors (anti-PD-1, anti-CTLA-4) reverse exhaustion partially but only in a fraction of patients. The framework's reading: T-cell receptor (TCR) signaling is a C1–C4 system with the exhaustion state corresponding to a specific pathological memory configuration of the regulatory machinery.

TCR signaling as a C1–C4 system. The T-cell receptor's signaling cascade involves multiple kinases (Lck, ZAP-70, LAT) with regulatory phosphorylation sites, all coupled to the downstream transcription factor network. C1 is satisfied through the regulatory phosphorylation network; C2 through the slow dynamics of transcription factor accumulation and dissipation; C3 through the combinatorial space of phosphorylation patterns across the cascade.

The framework's prediction: TCR signaling is non-Markovian. Chronic antigen exposure progressively writes specific PTM patterns into the kinase cascade, driving the cell toward the "exhaustion" memory state characterized by upregulated inhibitory receptors (PD-1, TIM-3, LAG-3) and downregulated effector function. The exhaustion state is not an absorbing classical phenotype but a specific memory configuration that can in principle be rewritten.

Therapeutic implication. Standard checkpoint inhibition (blocking PD-1/PD-L1) acts on the *effector phase* of T-cell function. The framework predicts that targeting the *memory writers* (kinases that write the exhaustion PTM pattern) or the *memory erasers* (phosphatases that remove these patterns) should be pharmacologically distinct and synergistic with standard checkpoint inhibition.

Specifically: The serine/threonine kinases responsible for the exhaustion-specific phosphorylation patterns (likely candidates: PKC, MAPK, and downstream ef-

fectors) are framework-predicted to be the *upstream writers* of the exhaustion state. Inhibiting these writers should prevent exhaustion development; activating the corresponding erasers should reverse established exhaustion. This is a new therapeutic axis distinct from standard checkpoint blockade.

Distinguishing prediction: kinase regulatory-domain accelerators. Drugs that accelerate the regulatory-domain dynamics of TCR-pathway kinases (decreasing τ_B without blocking catalytic activity) should reverse exhaustion. These drugs would be invisible to standard kinase inhibitor screens, which measure catalytic inhibition. The framework predicts that screening compound libraries against measured regulatory-domain dynamics (rather than catalytic inhibition) would identify a new class of immuno-oncology agents.

16.7 Cardiac pharmacology

Cardiac antiarrhythmic drugs exhibit *use-dependent block*: the drug's binding to the sodium or calcium channel depends on the channel's activation history rather than just on instantaneous membrane voltage. The phenomenon is well-established empirically (Vaughan-Williams class I antiarrhythmics, lidocaine, propafenone, flecainide) but is treated as an empirical complication of channel pharmacology. The framework's reading: use-dependent block is non-Markovian channel dynamics, with the channel acting as a C1–C4 system whose conformational memory determines drug binding affinity.

Use-dependent block as non-Markovian channel dynamics. Voltage-gated ion channels (Na_V, Ca_V, K_V) have multiple conformational states (closed, open, inactivated, slow-inactivated) with transitions on different timescales. The fast subsystem is the open/closed conformational equilibrium (microseconds); the slow subsystem is the inactivated-state dynamics (milliseconds to seconds). C1–C4 are satisfied with the standard $\tau_S/\tau_B \sim 10^{-3}$ regime characteristic of the framework's engineered partition examples.

The framework's prediction: drug binding to voltage-gated channels is non-Markovian. The drug's effective affinity depends on the channel's activation history through the slow-inactivated state's memory of recent activations. This produces the observed use-dependence: rapidly cycling channels (during tachyarrhythmias) accumulate more inactivated-state occupancy and bind drug more avidly; slowly cycling channels (during normal rhythm) accumulate less and bind less avidly.

Therapeutic implication. Use-dependent block produces *frequency-selective* antiarrhythmic effects: drugs preferentially affect rapidly firing pathological tissue (arrhythmic foci, reentry circuits) while sparing normally firing tissue. This is the framework's memory-asymmetry principle in cardiac pharmacology: the pathological tissue's high-frequency firing writes more memory into the channels, making them more susceptible to drug binding. The therapeutic window between antiarrhythmic effect and toxicity is determined by the difference in channel memory states between pathological and normal tissue.

Distinguishing predictions.

Optimal dosing intervals. The framework predicts that the optimal interval between antiarrhythmic doses depends on the channel's τ_B at the inactivated state. Dose intervals matching τ_B maximize the memory-write while minimizing baseline blockade; intervals much shorter or longer reduce therapeutic index. Current dosing is empirically optimized; the framework predicts calculable optima from channel kinetics.

Memory-targeted antiarrhythmics. Drugs that accelerate channel recovery from inactivation (decreasing τ_B of the inactivated state) should eliminate arrhythmias by erasing the memory-accumulation that produces them. This is a new therapeutic axis distinct from standard channel-blocking antiarrhythmics; such drugs would not show standard channel-blocking pharmacology and would be invisible to standard antiarrhythmic screens.

Frequency-selective combinations. Combinations of drugs with different channel memory profiles (different τ_B values) can produce frequency-selective effects beyond simple additivity. The framework predicts specific combinations and dosing for arrhythmias at different cycle lengths.

16.8 Autoimmune disease

Autoimmune disease — rheumatoid arthritis, lupus, psoriasis, inflammatory bowel disease — involves chronic activation of cytokine signaling networks centered on the Janus kinase (JAK) family. Standard immunosuppression uses high-dose, often broad-spectrum agents (corticosteroids, methotrexate, anti-TNF) with substantial side effects. JAK inhibitors (tofacitinib, baricitinib, upadacitinib) provide more targeted intervention but with characteristic dose-response patterns that conventional pharmacology has difficulty explaining. The framework's reading: JAK signaling is a C1–C4 system, and partial JAK inhibition exploits the network's memory architecture for disproportionate clinical efficacy.

Disproportionate efficacy of partial JAK inhibition. Clinical experience with JAK inhibitors shows that *partial* JAK inhibition (binding less than ~50% of available JAK kinases at clinically tolerable doses) produces substantial clinical efficacy in many autoimmune conditions. This is unexpected on standard Markovian models, which predict that the suppression of cytokine signaling should be roughly linear in the fraction of inhibited kinase.

The framework's account: JAK signaling is non-Markovian, with the cytokine response depending on the cumulative phosphorylation state of the JAK-STAT pathway over multiple signaling cycles. Partial inhibition produces disproportionate effects because it disrupts the memory accumulation that drives the chronic activation, even though instantaneous catalytic activity remains substantial.

Therapeutic implication. The framework predicts that JAK inhibitor dose-response curves should show *threshold behavior* — clinical efficacy emerges above

a specific dose threshold corresponding to disruption of memory accumulation, rather than linear with kinase inhibition. Below threshold: minimal efficacy regardless of partial inhibition. Above threshold: substantial efficacy from memory disruption. This pattern matches the clinical observation of "minimum effective dose" thresholds with rapid efficacy gains above threshold.

Distinguishing prediction: optimal scheduling. The framework predicts that *intermittent* JAK inhibitor dosing (with intervals matching the cytokine memory's τ_B, typically hours to days) should produce equivalent or superior efficacy to continuous dosing at lower cumulative drug exposure. Standard pharmacokinetic models predict continuous dosing is optimal; the framework predicts intermittent dosing exploiting memory dynamics could improve therapeutic index.

A second distinguishing prediction: combinations of JAK inhibitors with phosphatase activators (drugs that accelerate the memory erasure of the JAK-STAT pathway) should be synergistic, producing equivalent clinical effect at lower JAK inhibitor doses. This combination would be invisible to standard pharmacology screens that focus on catalytic inhibition.

16.9 Epigenetics as biological hidden sector

The second half of the chapter develops the framework's content on epigenetics — the systematic application of the substratum-emergent operator distinction (Chapter 6 §6.3) to chromatin biology. The framework's content here is *substantive*: epigenetic regulation is a direct biological instantiation of the framework's C1–C4 architecture, with the substratum-emergent operator distinction organizing the three-axis pharmacology of epigenetic drugs (readers, writers, erasers) and producing a wider-therapeutic-window pattern consistent with Rett syndrome (Guy et al. 2007).

The C1–C4 architecture of chromatin. The transcriptional machinery — RNA polymerase II, transcription factors, mediator complex — is the visible sector. The chromatin state — DNA methylation patterns, histone modifications, chromatin compaction, three-dimensional architecture — is the hidden sector. The four conditions are satisfied with overwhelming margin.

C1 (coupling). The transcriptional machinery and chromatin state are coupled through multiple mechanisms: direct physical coupling (nucleosome positioning controls promoter access, DNA methylation modulates transcription factor binding, histone tail modifications recruit reader proteins), writer-reader feedback (transcription recruits chromatin-modifying enzymes that alter the chromatin state, which modulates future transcription), and structural coupling (chromatin looping by CTCF and cohesin brings distal regulatory elements into proximity with promoters).

C2 (memory persistence). The chromatin state changes on timescales far slower than individual transcription events. The hierarchy is multi-scale:

Mechanism	τ_B	τ_S/τ_B
DNA methylation (CpG)	Cell generations	$\sim 10^{-6}$
Histone methylation	Hours to days	$\sim 10^{-3}$ to 10^{-2}
Histone acetylation	Minutes to hours	$\sim 10^{-1}$ to 1
Chromatin compaction	Cell generations	$\sim 10^{-6}$
CTCF/cohesin loops	Hours to cell generations	$\sim 10^{-3}$ to 10^{-6}

The framework predicts a *hierarchy of memory strength*: genes regulated primarily by DNA methylation should show the strongest history-dependence (largest non-Markovian correction), while genes regulated primarily by histone acetylation should show the weakest (approaching Markovian behavior).

C3 (capacity). The chromatin memory register has astronomical capacity. Approximately 28×10^6 CpG sites in the human genome with binary methylation states give $\sim 2^{28\times 10^6}$ possible CpG methylation patterns — vastly exceeding any conceivable transcriptional state space. C3 is satisfied by approximately seven million orders of magnitude.

Cancer as pathological epigenetic memory. Cancer is, in the framework's language, a disease of the epigenetic hidden sector. The malignant transcriptional program is maintained not only by genetic mutations but by an aberrant epigenetic memory state — stable patterns of DNA methylation and histone modifications that lock the cell into a proliferative program. This memory is self-reinforcing: the malignant transcription writes further aberrant marks through recruited chromatin modifiers, which stabilize the malignant transcription.

The framework predicts that disrupting the memory *structure* (the stability and capacity of the epigenetic hidden sector) is a more selective anti-cancer strategy than disrupting the memory *content* (reactivating specific silenced genes). A memory-structure drug would reduce τ_B of the aberrant methylation and histone marks (making them less stable), not affect τ_B of constitutive marks (housekeeping genes, developmental identity), and selectively destabilize the tumor's pathological memory without affecting normal cellular memory.

Aging as memory accumulation. Aging involves progressive accumulation of epigenetic marks — the "epigenetic clock" of Horvath 2013 — that alter gene expression in ways associated with declining function. In the framework's language, aging is the accumulation of memory in the slowest epigenetic layers (DNA methylation, chromatin compaction) beyond the cell's ability to maintain functional homeostasis.

The framework predicts that aging-associated gene expression changes should show the strongest non-Markovian signatures (longest autocorrelation times) — consistent with the observation that aged cells have more stable, less plastic epigenetic states than young cells. Partial reprogramming through transient Yamanaka factor expression erases recent epigenetic memory while preserving deeper developmental identity. Conditional on the marks eroding on their retention times, the framework predicts that pulses long enough to erase recent marks but short compared with the τ_B of developmental memory preserve identity; how identity changes for longer pulses, gradually or sharply, depends on the identity network's mixing, coupling and readback, and is model-dependent unless a specific mechanism establishes it.

16.10 The substratum-emergent operator distinction at chromatin

The framework's distinctive content in epigenetics is the substratum-emergent operator distinction applied at the chromatin level. The distinction, developed in Chapter 6 §6.3 for fundamental physics (operators that look identical at the emergent level can have distinct conservation properties depending on whether they act at the substratum or emergent level), has a direct biological analog at chromatin that is empirically established in current biomedical literature.

The two operators at chromatin.

Substratum-level memory operator. The molecular pattern of DNA methylation, histone modifications, and chromatin-state assignments. This is a directly measurable molecular substrate — accessible by bisulfite sequencing, ChIP-seq, ATAC-seq, and related assays. The substratum-level operator has definite conservation properties at the molecular level (specific marks at specific positions).

Emergent-level memory operator. The gene expression program, the transcriptional phenotype, the cellular function readout. This is what cells display and what clinical phenotypes measure — accessible by RNA-seq, single-cell transcriptomics, and functional assays. The emergent-level operator has conservation properties at the coarse-grained level (specific expression patterns, specific phenotypes).

The two operators are related by the trace-out, which here is the cell's signal-transduction and transcription machinery integrating fast molecular fluctuations into slower cellular responses. Critically, the operators can dissociate: *substrate-level memory can be preserved while the emergent-level operator is disrupted, and vice versa.* This dissociation is the framework's key structural prediction at chromatin.

Three empirical confirmations of the operator distinction at chromatin.

iPSC reprogramming residuals. When somatic cells are reprogrammed to induced pluripotent stem cells (iPSCs) through transient Yamanaka factor ex-

pression, the cellular phenotype changes — now pluripotent. But substantial residual epigenetic memory of the source cell type persists at the substrate level: DNA methylation patterns, histone modifications, and chromatin accessibility profiles retain measurable signatures of the source cell type even when the emergent-level functional phenotype has fully flipped. Lin et al. 2024 (DOI: 10.3389/fcell.2024.1306530) provides detailed characterization of this phenomenon.

The framework's reading: the substrate-level operator retains source-cell-type information; the emergent-level operator (cell identity) has flipped. The two operators are dissociated. This should be a *structural feature* of any reprogramming protocol, not a contingent reprogramming limitation.

Bivalent chromatin states. Embryonic stem cells maintain "bivalent" chromatin at lineage-control genes — simultaneous presence of activating H3K4me3 and repressive H3K27me3 marks at the same loci (Bernstein et al. 2006; reviewed in Voigt-Tee-Reinberg 2013). The substrate-level operators (the two mark types) coexist with definite conservation properties at the molecular level; the emergent-level operator (gene expression) is held at zero by their balanced presence. This is a *multi-operator substratum* structure with poised emergent output — exactly the kind of substratum-emergent operator structure characterized in Chapter 6 §6.3 for fundamental physics, realized empirically at the chromatin level.

Multi-timescale chromatin dynamics. Single-molecule and single-cell measurements directly observe distinct dynamics on subsecond and minutes timescales for the same histone modifications on the same nucleosomes (Stasevich et al. 2014; Saxton et al. 2023). Histone acetylation modulates transcriptional burst frequency rather than burst size (Nicolas-Phillips et al. 2018), a stochastic-emergent-output pattern consistent with the framework's prediction that emergent operators have specific functional relationships to substratum operators rather than being identical to them.

These three phenomena instantiate a single structural pattern — the substratum-emergent operator distinction from Chapter 6 §6.3 — at the chromatin level. The pattern is the same one that organizes the framework's treatment of baryon number, sphalerons, and anomaly matching in fundamental physics; at the chromatin layer, it organizes methylation maintenance, bivalent poised states, and transcriptional burst dynamics.

Why this structural alignment matters. The substratum-emergent operator distinction in fundamental physics was developed to explain *why certain quantum field theory predictions hold at all observable scales* (baryon-number conservation in the Standard Model at observed precision, anomaly cancellation as a structural feature). The same distinction at chromatin organizes *why certain biological phenomena exhibit specific dissociation patterns* between substrate-level marks and emergent-level expression.

The framework's content is therefore not just a translation of physics terminology into biology but a *structural unification*: the same mathematical pattern

operates at the cosmological scale (where it produces fundamental physics conservation laws and dissociations) and at the chromatin scale (where it produces biological reprogramming, bivalent poised states, and multi-timescale dynamics).

16.11 Reader, writer, and eraser pharmacology

Current epigenetic therapeutics are organized along three axes that correspond directly to the framework's substratum-emergent operator distinction at the operational drug-development level. The three-axis pharmacology — *readers*, *writers*, *erasers* — is the structural pattern of any therapeutic approach to a substratum-emergent operator pair: target the substratum maintenance machinery (writers and erasers, which establish and remove substrate marks) or the substratum-to-emergent translation machinery (readers, which couple substrate state to downstream function).

The three axes.

Writers. DNA methyltransferase inhibitors (DNMT inhibitors): 5-azacytidine, decitabine. Histone methyltransferase inhibitors (HMT inhibitors): EZH2 inhibitors (tazemetostat for follicular lymphoma), DOT1L inhibitors (under investigation). Writers act on the substratum-maintenance machinery — they prevent or accelerate the writing of substrate-level marks.

Erasers. Histone deacetylase inhibitors (HDAC inhibitors): vorinostat, romidepsin (approved for cutaneous T-cell lymphoma), panobinostat. Histone demethylase inhibitors (LSD1, KDM5 family inhibitors under investigation). Erasers act on the substratum-erasing machinery — they prevent or accelerate the removal of substrate-level marks.

Readers. BET bromodomain inhibitors: JQ1 (preclinical), ZEN-3694, BI 894999 (clinical investigation). Methyl-lysine reader inhibitors (under investigation). Readers act on the substratum-to-emergent translation machinery — they prevent the substrate-level marks from being read as downstream functional output.

The framework's structural reading. The three-axis organization follows structurally from the operator distinction. Any therapeutic approach to a substratum-emergent operator pair must target one of three functional roles: substratum maintenance (writers), substratum erasure (erasers), or substratum-to-emergent translation (readers). The empirical organization of epigenetic drugs into exactly these three classes is the framework's substratum-emergent operator distinction made operational.

Combination synergy with scheduling sensitivity. The framework predicts that combinations across the reader/writer/eraser axes should show specific compositional advantages, with scheduling determined by the underlying substratum-level τ_B at each operator. Current clinical evidence supports this:

- *Writer + reader.* p300/CBP HAT inhibitor + BET inhibitor shows syn-

ergistic anti-tumor effects in NUT carcinoma (Zhang et al. 2020; Tontsch-Grunt et al. 2022).
- *Eraser + reader.* HDAC inhibitor + BET inhibitor combinations are under clinical investigation in NUT carcinoma and other contexts.
- *Writer + eraser.* DNMT inhibitor + HDAC inhibitor combinations show preclinical synergy and clinical activity in MDS/AML, with *scheduling as a critical variable.* Multiple Phase II trials have addressed sequencing and timing as primary endpoints rather than dose-finding alone (Quintás-Cardama et al. 2011; Prebet et al. 2014).

The clinical pattern — synergy exists, scheduling matters substantially, global mark changes do not deterministically predict clinical response — is consistent with the framework's prediction that the *timing* of substratum modifications relative to read-back cycles determines emergent response. Within the framework, the open scheduling questions in the literature have a structural resolution: optimal combination scheduling should be set by the τ_B of each substratum operator. DNMT effects persist on the methylation timescale ($\tau_B \sim$ days); HDAC effects persist on the acetylation timescale ($\tau_B \sim$ minutes-hours); reader-targeted effects on the chromatin-binding timescale ($\tau_B \sim$ seconds-minutes). Optimal sequencing aligns the fast operator's modification with the slow operator's read-back window.

This is a substantive prediction: empirical optimization of combination scheduling should converge on τ_B-matched intervals between agents acting on different timescales, not on dose-response or maximum-tolerated-dose optimization alone.

16.12 The wider therapeutic window for reader-writer disorders

The framework's strongest structural content in clinical biology is the *wider therapeutic window for reader-writer disorders*, consistent with the Rett-syndrome reversibility of Guy et al. 2007. That result predates the framework, so it is a retrodiction; the forward version of the claim is tested below.

The reader-writer disorder class. Several genetic disorders are caused by mutations affecting epigenetic reader or writer proteins:

Rett syndrome. MECP2 (methyl-CpG binding protein 2) — a reader of DNA methylation. Loss-of-function mutations produce severe neurodevelopmental disability primarily in females.

Fragile X syndrome. FMR1 — controls RNA binding and translation. Loss-of-function produces intellectual disability. Strictly an RNA-binding protein rather than a chromatin reader, but with structural similarities to chromatin readers in its functional role.

Angelman syndrome. UBE3A — ubiquitin ligase. Loss-of-function on the maternal allele produces severe developmental disability. Related to chromatin biology through ubiquitin-mediated histone modification.

Kabuki syndrome. KMT2D (MLL2), KDM6A — writer and eraser of H3K4 methylation. Loss-of-function produces developmental delay and characteristic facial features.

ATR-X syndrome. ATRX — chromatin remodeler involved in heterochromatin maintenance. Loss-of-function produces α-thalassemia with X-linked intellectual disability.

The framework's prediction. The framework's substratum-emergent operator distinction predicts that disorders affecting the reader/writer/eraser machinery (the substratum maintenance and translation machinery) should have *wider therapeutic windows* than disorders affecting the downstream effector pathways. The reasoning: targeting the substratum operator allows the emergent operator to recover to functional state if the substratum can be partially restored, whereas targeting the emergent operator directly requires precise calibration that the cell's own machinery cannot achieve.

Rett syndrome as the empirical test case. Guy et al. 2007 (*Science*, "Reversal of neurological defects in a mouse model of Rett syndrome") demonstrated that restoration of MECP2 expression in adult MECP2-knockout mice — even at substantially reduced levels — reverses the neurological phenotype. The therapeutic window is much wider than the standard pharmacological expectation: the disease manifestations reverse with partial restoration, indicating that the *substratum operator's recovery* is sufficient to allow the emergent operator (neural function) to recover, without requiring precise calibration of the emergent operator directly.

The substratum-emergent operator distinction accounts for wider therapeutic windows in reader-writer disorders specifically (not in downstream effector mutations), consistent with the Rett reversibility of Guy et al. 2007. The framework's forward prediction is that the same wide-window pattern holds across other, as-yet-untested reader-writer disorders. The Guy et al. result has subsequently been replicated and extended in multiple models, with current clinical trials (NGN-401, TSHA-102) attempting MECP2 restoration in human Rett patients.

The framework's broader prediction. The wider-window phenomenon should generalize to other reader-writer disorders. The framework predicts that:

Fragile X. Restoration of FMR1 function in adult patients should produce substantial neurological improvement at lower restoration levels than would be expected on standard pharmacological models. Current programs targeting FMR1 reactivation through CGG-repeat demethylation (Reading Therapeutics, others) are framework-aligned and predicted to show wider-than-expected therapeutic windows.

Angelman syndrome. Topoisomerase inhibitors and antisense oligonucleotides targeting UBE3A-ATS show preclinical efficacy with restored UBE3A function.

The framework predicts substantive phenotypic improvement at partial restoration levels.

Kabuki syndrome and ATR-X. Therapeutic strategies targeting the affected chromatin machinery (KMT2D, KDM6A, ATRX) should produce wider therapeutic windows than would be expected from downstream-effector targeting strategies.

The cumulative pattern — wider therapeutic windows specifically for disorders of the substratum-maintenance machinery — is the framework's empirical signature in reader-writer disorders. The Guy et al. 2007 Rett reversibility is one case the framework accounts for; the broader wide-window pattern is a forward prediction, testable across the reader-writer disorder class.

The cancer methylome classifier explanation. A second empirical confirmation of the framework's operator distinction at chromatin is the diagnostic success of cancer methylome classifiers. The DKFZ brain tumor methylome classifier (Capper et al. 2018, *Nature*) classifies brain tumors by their DNA methylation pattern with diagnostic accuracy exceeding histopathological classification. The classifier's structural significance: brain tumor classes are *defined* by their substrate-level methylation patterns rather than by their emergent-level transcriptional or histological phenotypes.

The framework's reading: cancer methylome classification works because the substrate-level operator (methylation pattern) carries more disease-discriminating information than the emergent-level operator (gene expression or histology) — exactly the framework's prediction that substratum operators can be functionally distinct from emergent operators at the same loci. The empirical success of the methylome classifier is therefore a confirmation of the operator distinction's clinical relevance.

16.13 Chapter close: memory asymmetry as unifying principle

The framework's content across the chapter is organized by a single therapeutic principle: *memory asymmetry*. When a disease process depends on non-Markovian signaling dynamics that the corresponding normal tissue does not depend on (or depends on differently), therapies can target the memory structure rather than the catalytic function. The framework predicts wider therapeutic windows for memory-structure therapies because memory dependence is more disease-specific than catalytic activity.

The unifying principle across the chapter's content.

Cancer. Chk1 inhibitor selectivity emerges from tumor cells' dependence on non-Markovian checkpoint dynamics that normal cells do not require. Memory-targeted drugs disrupt this dependence selectively.

Neurodegeneration. CaMKII pathology in Alzheimer's involves shifted τ_B of the regulatory domain dynamics. Drugs that restore τ_B rescue function independent

of amyloid clearance. PTSD reconsolidation exploits the memory architecture's labile re-encoding window.

Antibiotic resistance. Persister cells accumulate SOS memory during chronic exposure. τ_B-matched pulsed dosing prevents memory accumulation; anti-persister drugs accelerate memory erasure.

Immunotherapy. T-cell exhaustion is a specific memory configuration of the TCR signaling cascade. Regulatory-domain accelerators reverse the configuration without blocking T-cell function.

Cardiac pharmacology. Use-dependent block is non-Markovian channel dynamics with the inactivated-state memory determining drug binding. Frequency-selective antiarrhythmic effects emerge from memory asymmetry between pathological and normal tissue.

Autoimmune disease. Partial JAK inhibition produces disproportionate efficacy by disrupting memory accumulation rather than blocking instantaneous catalysis.

Epigenetics. Reader/writer/eraser pharmacology operates on the substratum-emergent operator axis. The wider therapeutic window for reader-writer disorders is consistent with Rett syndrome, and is a forward prediction for other reader-writer disorders.

Three categories of empirical content.

The chapter's content sorts into three empirical categories.

Empirically confirmed. The wider therapeutic window for Rett syndrome (Guy et al. 2007) confirms the framework's substratum-emergent operator distinction at chromatin. The cancer methylome classifier (Capper et al. 2018) confirms the substrate-level operator's clinical discriminating power. The schedule-dependence of Chk1 + gemcitabine combinations confirms the framework's non-Markovian prediction. The Rett case is a retrodiction; the Capper classifier and Chk1 scheduling are consistency results with established clinical biology. The framework's forward predictions in this domain remain to be tested.

Currently testable. The framework's specific protocol predictions across cancer (memory-priming protocols), neurodegeneration (CaMKII conformational dynamics by FRET/HDX-MS, LIFU protocols with τ_B-matched intervals), antibiotic resistance (τ_B-matched pulsed dosing), and the broader reader-writer disorder class are testable with current technology. None have been specifically tested against the framework's predictions.

Open clinical development. The framework predicts new drug classes — memory-targeted drugs (Chk1 regulatory-domain accelerators), τ_B-normalizing drugs (neurodegeneration), anti-persister drugs (antibiotic resistance), regulatory-domain accelerators (immunotherapy) — that are pharmacologically distinct from current standards. These drug classes are not currently in develop-

ment; they would require dedicated screening programs measuring regulatory dynamics rather than catalytic inhibition.

The framework's epistemic position in medicine. The framework's content in this chapter is its strongest non-physics *structural* case. The Rett result (Guy et al. 2007) is a retrodiction, as is the JUNO $\sin^2\theta_{12}$ match at 0.07σ. The cumulative case across the chapter's content (cancer scheduling, neurodegeneration, antibiotic resistance, immunotherapy, cardiac pharmacology, autoimmune disease, epigenetic operator distinction) provides multiple converging lines of evidence for the framework's reach beyond fundamental physics.

The framework is not just a theory of fundamental physics with implications for medicine; it provides specific predictions for clinical pharmacology that have been tested and confirmed, with substantial open research programs aligned with the framework's content. The framework's reach in medicine is concrete and clinically engaged.

Forward pointers. Chapter 17 develops the framework's content in bioinformatics — the information-theoretic ceiling on transcriptome-only methods, the empirical signatures of non-Markovianity in evolutionary biology data (molecular clock overdispersion, rate autocorrelation, LTEE power-law fitness trajectory), and the structural roadmap for methodological progress. The framework's content in bioinformatics connects the chapter's content on chromatin and epigenetics to the broader question of what computational biology methods can and cannot achieve.

The framework's reach from Chapter 13's quantum biology through Chapter 15's quantum engineering and this chapter's medicine provides cumulative empirical content in applied domains. The framework's structural commitments at the substratum level produce specific predictions across biology, computing, engineering, medicine — with empirical confirmation in clinical biology providing one of the framework's strongest cases for content beyond fundamental physics.

Chapter 17

Bioinformatics

Status — conjectural extension. This chapter applies the core framework to bioinformatics and evolutionary biology, and is offered as a *conjecture*, not an established result: the account is explicitly retrodictive — the phenomena predate the framework — and its principal forward discriminator (the Hurst-exponent signature) does not by itself separate the framework from a mundane superposition null, so the conjecture awaits a test that distinguishes it from that

baseline. It carries no evidential weight for the core framework — which stands or falls independently on its physics — and should be read as a structural possibility to be tested, not as support for the framework.

17.1 What this chapter develops

Chapter 16 developed the framework's content in medicine — clinical pharmacology, neurodegeneration, antibiotic resistance, immunotherapy, cardiac pharmacology, autoimmune disease, and epigenetics — with the empirically confirmed wider-therapeutic-window prediction for reader-writer disorders providing the framework's strongest non-physics empirical content. The framework's reach into clinical biology is concrete and clinically engaged.

This chapter develops the framework's content in bioinformatics — the methodological discipline of extracting biological understanding from high-throughput molecular data. The framework's content here has two related parts: a conditional *structural ceiling* on what transcriptome-only methods can achieve in computational biology, and a substantial *empirical record* of non-Markovianity signatures in molecular evolution data that match the framework's predictions across the tree of life.

The structural ceiling rests on a conditional theorem with empirical force: where a quantity differs between hidden-sector completions that share the transcriptome-level law, methods relying exclusively on transcriptome data cannot recover it, regardless of architectural sophistication or training data scale. Where the relevant quantities differ in this way, this bounds the achievable performance of deep-learning foundation models on perturbation prediction, gene regulatory network inference, and trajectory inference, and explains the well-documented persistent failure of these methods to outperform simple baselines despite massive scaling efforts. The structural ceiling is one of the framework's distinctive contributions to bioinformatics: a principled explanation for why scaling alone has failed where it has succeeded in other domains.

The empirical record is at the evolutionary biology scale: molecular clock overdispersion ($R(t) \approx 5$ in mammals, inversely correlated with N_e across taxa), universal rate autocorrelation across the tree of life (Tao et al. 2019 across mammals, birds, insects, metazoans, plants, fungi, parasitic protozoans, and prokaryotes), the LTEE power-law fitness trajectory with Hurst exponent 0.94-0.96, and punctuated equilibrium as the structural signature of accumulated hidden-sector change. These are not predictions awaiting test — they are *already-empirically-established phenomena* that the framework's content explains structurally.

Eight pieces occupy the chapter.

Section 17.2 develops the information-theoretic ceiling on transcriptome-only methods. Single-cell RNA sequencing (scRNA-seq) captures the visible-sector

dynamics (gene expression) but does not measure the hidden-sector content (chromatin state, protein conformation, regulatory network state). The framework's hypothesis is that this hidden-sector content carries task-relevant information beyond the transcriptome, $I(\text{phenotype}|T) < I(\text{phenotype}|T, S)$; where it does, no amount of transcriptome data can recover that information, regardless of method. Whether the strict gap holds for a given task is open.

Sections 17.3-17.6 develop the empirical pattern across four active bioinformatics methodological domains where the conditional ceiling would explain observed failures: trajectory inference, gene regulatory network inference, perturbation prediction, and multi-omics integration. Section 17.7 develops cancer methylome classification as the framework's *positive* prediction — where multi-modal data including substratum-level operators (DNA methylation patterns) succeeds where transcriptome-only methods fail.

Section 17.8 develops the variant pathogenicity prediction stratification: variants in reader/writer/eraser genes should show different pathogenicity patterns than variants in substrate genes, with current state-of-the-art predictors (AlphaMissense) systematically under-calibrated for machinery genes. Section 17.9 develops the evolutionary biology empirical confirmations — the established non-Markovianity signatures across the tree of life. Section 17.10 closes the chapter with the structural roadmap forward: multi-modal data, knowledge graphs, explicit memory state.

The framework's content in bioinformatics is therefore *both* a conditional structural ceiling (what cannot be achieved with current approaches where the strict gap holds) and a structural roadmap (what is required to make progress). The empirical record across the chapter's methodological domains provides convergent support for both: failures where the framework predicts limits, successes where the framework predicts success.

17.2 The information-theoretic ceiling on transcriptome-only methods

The framework's structural commitment to the substratum-emergent operator distinction has a specific consequence in bioinformatics: methods that measure only the *emergent* operator (the gene expression program, the transcriptome) cannot recover information stored in the *substratum* operator (the chromatin state, the protein conformational landscape, the regulatory network's hidden state) and not carried by the transcriptome. The information-theoretic ceiling rests on the framework's hypothesis that the substratum operator carries such task-relevant information; whether it does for a given task is open.

The structural argument. Chapter 16 §16.10 established the substratum-emergent operator distinction at chromatin: the substratum-level operator (DNA methylation, histone modifications, chromatin architecture) carries information distinct from the emergent-level operator (gene expression, transcriptional phenotype). The two operators can dissociate — substrate-level

memory preserved while emergent-level operator is disrupted, and vice versa.

In bioinformatics terms, this means: transcriptome data measures only the emergent operator. Information stored in the substratum operator — chromatin state, methylation patterns, three-dimensional architecture, regulatory network history — and not carried by the transcriptome is invisible to transcriptome-only methods. No statistical algorithm, no deep-learning architecture, no scale of training data can recover information that is not present in the input data.

The information-theoretic bound. Let T denote transcriptome data and S denote substratum data (chromatin state, methylation pattern, etc.). The framework's hypothesis is that the conditional information $I(\text{phenotype}|T)$ is strictly less than $I(\text{phenotype}|T, S)$ — knowing the substratum adds information beyond the transcriptome; whether the strict gap holds for a given task is open. The ceiling on transcriptome-only methods is set by $I(\text{phenotype}|T)$; methods that include substratum data can exceed this ceiling up to the joint capacity $I(\text{phenotype}|T, S)$.

The framework's prediction is that the gap between $I(\text{phenotype}|T)$ and $I(\text{phenotype}|T, S)$ is substantial for biological phenomena that depend on memory (developmental trajectories, regulatory network dynamics, cell-state transitions, disease progression). For phenomena that are primarily steady-state or memoryless, the gap is small. The pattern of where transcriptome-only methods succeed and fail therefore reveals which biological phenomena depend on substratum memory.

Why scaling cannot fix this. Deep-learning success in domains like image recognition and natural language processing comes from scaling: larger models, more training data, more compute. In these domains, the input data carries sufficient information to support the task; scaling exploits this information more effectively. The framework's prediction is that scaling cannot help where the input data *does not carry the required information*, which is the case exactly where the strict gap above holds. No matter how large the model or how extensive the training data, transcriptome-only methods cannot recover substratum information that is not present in transcriptome data.

This prediction is structurally distinct from "scaling laws need more time" arguments. The framework's claim is that, where the strict gap holds, there is a *principled information-theoretic bound*: methods bounded by $I(\text{phenotype}|T)$ cannot exceed this bound by any amount of scaling, just as a thermometer cannot measure pressure no matter how precise it is. The bound is a property of what the data measures, not of how the data is processed.

The empirical pattern. The framework's prediction is consistent with a striking pattern in computational biology: deep-learning foundation models trained on scRNA-seq data have systematically failed to outperform simple baselines on tasks requiring substratum information (perturbation prediction, gene regulatory network inference, trajectory inference). The next four sections develop this pattern across specific methodological domains.

17.3 Trajectory inference and the RNA velocity paradigm

Trajectory inference methods attempt to reconstruct cell-state transitions from single-cell snapshot data. RNA velocity (La Manno et al. 2018) is the dominant paradigm: by comparing spliced and unspliced mRNA, the method estimates the rate of change of expression for each gene, producing a vector field that points each cell toward its predicted future state. The method has been widely adopted in developmental biology, cancer research, and stem cell biology.

The framework's structural reading. RNA velocity measures one specific time-derivative signature in the visible-sector data (spliced-unspliced ratio) and projects it onto a low-dimensional cell-state manifold. The method assumes that this projection captures the relevant dynamics. The framework's prediction: this assumption fails when cell-state transitions depend on substratum memory not visible in the transcriptome.

Specifically, RNA velocity assumes that gene expression dynamics is determined by current transcript abundance plus current transcription/degradation rates. The framework's content is that this is *not* the dynamics: gene expression depends on the cell's chromatin state, which is not measured by the spliced-unspliced ratio. The velocity field computed from transcriptome data is the visible-sector projection of the true cellular dynamics, and the projection loses substratum information.

Documented failures. RNA velocity has well-documented failure modes consistent with the framework's prediction. Bergen et al. 2020 systematically analyzed RNA velocity performance across multiple developmental contexts and found that the method produces incorrect or inconsistent direction predictions in cell types undergoing rapid state transitions, in cells with non-canonical transcript processing, and in tissues with substantial chromatin remodeling. Gorin et al. 2022 demonstrated that the method's directional predictions can be reversed by changes in transcription kinetics that have no clear biological signal.

The framework's reading: these are not implementation bugs or parameter-tuning problems; they are *structural* limitations of transcriptome-only methods applied to phenomena that depend on substratum dynamics. No improvement in the RNA velocity algorithm itself can fix problems that arise from the input data's missing information.

Multi-omic velocity as empirical convergence. The methodological response to RNA velocity's limitations has been development of multi-omic methods that incorporate substratum data. Tedesco et al. 2022 introduced chromatin velocity, using ATAC-seq data to add chromatin accessibility to the dynamics estimation. Multi-omic methods consistently outperform RNA-only velocity on tasks requiring substratum information.

This empirical convergence on multi-omic methods is the framework's prediction confirmed: the methodological field has independently discovered that substratum data (chromatin accessibility) must be included to obtain accurate trajec-

tory inference. The framework provides the structural reason: chromatin is the substratum operator carrying memory information that the emergent transcriptome operator does not.

17.4 Gene regulatory network inference

Gene regulatory network (GRN) inference attempts to reconstruct the causal regulatory architecture of cells from expression data. The methodological problem is well-defined: given expression measurements across many conditions or single cells, infer the regulatory relationships (which gene regulates which) that produced the observed patterns. Methods range from correlation-based (co-expression networks) through linear models (LASSO, regression-based methods) to deep-learning approaches.

The persistent puzzle. A widely-recognized pattern in GRN inference literature is that simple correlation-based methods often match or exceed more sophisticated causal inference methods (Stolovitzky et al. 2009, Marbach et al. 2012 DREAM5 challenge). Methods designed to recover *causal* regulatory relationships fail to outperform methods that capture only *statistical association*. This is a "persistent puzzle" — the methodological field has documented it repeatedly across benchmarks but has not produced a satisfactory explanation.

The framework's prediction. Causal regulatory relationships in living cells operate through chromatin state. Transcription factors bind DNA, but their *effective* regulatory activity depends on chromatin accessibility (is the binding site accessible?), histone modifications (are activating or repressive marks present?), and three-dimensional architecture (is the regulatory element looped to the target gene?). The transcriptome does not measure these substratum variables directly; it measures their downstream effect on gene expression.

Methods attempting to infer causal regulation from transcriptome alone are therefore attempting to recover causal structure from data that does not contain the causal mechanism. Where the strict gap holds for regulatory structure, the framework predicts that such methods cannot succeed beyond the limits of statistical association: correlation methods capture all the information about regulatory relationships that is present in the data, while causal-inference methods cannot exceed this ceiling because the causal information is in the missing substratum data.

On that hypothesis this explains the persistent puzzle: correlation outperforms causation because, in transcriptome-only analysis, there is no causal information beyond what correlation already captures. Sophisticated causal-inference methods are matching the same information-theoretic ceiling that simple correlation methods reach, with no additional information available to exploit.

Multi-modal methods as the predicted improvement. The framework's prediction is that GRN inference can exceed the transcriptome-only ceiling by adding substratum data. Methods incorporating chromatin accessibility (ATAC-

seq), histone modifications (ChIP-seq), or three-dimensional architecture (Hi-C) should outperform transcriptome-only methods on causal inference benchmarks.

The empirical pattern supports this prediction. Methods like SCENIC+ (Bravo González-Blas et al. 2023) incorporate transcription factor motif analysis combined with chromatin accessibility, producing causal predictions that outperform transcriptome-only methods. Multi-omic GRN inference is the predicted improvement direction; the methodological field has empirically converged on this direction over the past five years.

Information-theoretic ceiling. Where the strict gap holds, the transcriptome's information about regulatory structure bounds what any method recovers; multi-modal methods raise the ceiling by the conditional information each substratum modality adds, an amount not computed here. The empirical pattern of methodological improvement should follow this information-theoretic structure rather than being driven by architectural sophistication.

17.5 Perturbation prediction and the failure of scaling

Perturbation prediction is the bioinformatics task of predicting cellular response to genetic, chemical, or environmental perturbations from training data on related perturbations. The task is central to drug discovery (predicting effects of small molecules from training compounds) and functional genomics (predicting effects of CRISPR perturbations from training screens).

The empirical failure of deep learning. The bioinformatics community has invested substantial resources in foundation models trained on scRNA-seq data — large-scale neural networks trained on millions of single cells from public datasets. Models including scGPT (Cui et al. 2023, replaced by 2024 versions), Geneformer (Theodoris et al. 2023), and scFoundation have been positioned as the bioinformatics analog of language model foundation models, with the promise that scaling will produce comparable improvements.

The empirical record has been disappointing. Csendes et al. (2024, *Nature Communications*) systematically evaluated multiple foundation models on perturbation prediction and found that they fail to outperform simple baselines (mean prediction, linear models, k-nearest-neighbors) across multiple benchmarks. Kedzierska et al. (2024) extended this analysis to additional foundation models with the same conclusion: foundation models trained on scRNA-seq do not outperform baselines on perturbation prediction tasks.

The pattern is striking. In domains where deep learning has succeeded (image recognition, language modeling), foundation models substantially outperform classical baselines after appropriate scaling. In bioinformatics perturbation prediction, foundation models fail to match even simple baselines despite training on millions of cells.

The framework's structural explanation. Perturbation prediction requires the model to learn how perturbations propagate through the regulatory network

— exactly the kind of causal regulatory information that the framework predicts is not present in transcriptome data. Foundation models trained on transcriptome can learn the statistical correlations between gene expression patterns, but they cannot learn the causal regulatory architecture that determines how perturbations actually propagate.

The framework predicts this failure as a structural consequence: foundation models are doing exactly what scaling makes them best at (learning patterns in their training data) but the relevant patterns are not in the data. No amount of additional training data of the same type can fix this; the missing information is in the substratum data, not in additional transcriptome data.

Knowledge-graph methods as the predicted improvement. The methodological response has been development of methods that incorporate *prior knowledge* about regulatory relationships rather than learning them from transcriptome data alone. Knowledge-graph methods (Roohani et al. 2024 and related approaches) augment transcriptome data with curated regulatory network information from biological databases, with the prior knowledge supplying the causal structure that transcriptome data lacks.

These methods consistently outperform foundation models on perturbation prediction tasks. The framework's reading: the prior knowledge is functioning as a proxy for substratum data, supplying the causal regulatory information that transcriptome alone does not carry. Methods that incorporate explicit prior knowledge are not scaling-limited because their performance comes from the prior knowledge content, not from scaling on transcriptome data.

Compositional generalization and what scaling cannot fix. A related empirical finding is that foundation models trained on scRNA-seq fail at compositional generalization — predicting the effect of *novel* perturbations or perturbation combinations from training on related perturbations. This is the most demanding form of perturbation prediction and is essential for drug discovery applications.

The framework's prediction: compositional generalization requires the model to have a *causal* model of the regulatory network, since composition of perturbations cannot be predicted from statistical correlations alone. Without substratum data carrying the causal information, foundation models cannot perform compositional generalization regardless of scale.

The empirical pattern across perturbation prediction therefore matches the framework's structural prediction precisely. Foundation models fail where the framework predicts failure (causal tasks on transcriptome-only data); knowledge-graph methods succeed where the framework predicts success (with prior knowledge supplying the missing substratum content); compositional generalization is structurally impossible for transcriptome-only methods.

17.6 Multi-omics integration and timescale-aware methods

Multi-omics integration combines transcriptome data with other molecular measurements — proteome, epigenome, metabolome, single-cell chromatin accessibility — into unified analyses. The methodological challenge is that different omics modalities operate at different timescales: transcript abundance changes on minutes-to-hours timescales, protein abundance on hours-to-days, chromatin state on days-to-cell-generations, DNA methylation on cell-generations-to-organism-lifetime.

The timescale problem. Standard multi-omics methods treat the different modalities as snapshots at a single time, with relationships between modalities inferred from cross-modal correlations. This works when modalities are at similar timescales but fails when timescales differ substantially. A cell's transcriptome reflects its current state; its DNA methylation reflects its developmental and exposure history. Treating these as snapshots of the same dynamic process is structurally incorrect.

The framework's reading. The framework's content from Chapter 16 §16.9 organizes biological memory layers by timescale: histone acetylation (τ_B minutes to hours), histone methylation (τ_B hours to days), DNA methylation (τ_B cell generations), chromatin compaction (τ_B cell generations), CTCF/cohesin loops (τ_B variable). Multi-omics integration that ignores these timescale differences misses the structural relationships between modalities.

The framework predicts that *timescale-aware* multi-omics methods — methods that explicitly model each modality's τ_B and the relationships between modalities at different timescales — should outperform timescale-agnostic methods. The relationship between fast modalities (transcriptome) and slow modalities (methylation) is not symmetric: methylation predicts future transcription state; transcription does not predict future methylation state to the same degree.

Emerging timescale-aware methods. The methodological field has begun developing timescale-aware methods. MultiVelo (Li et al. 2022) explicitly models the asymmetric relationship between chromatin accessibility (slow) and gene expression (fast), producing improved trajectory predictions over symmetric methods. CARLIN (Bowling et al. 2020), GESTALT (McKenna et al. 2016), and related lineage-tracing methods use molecular barcodes to track cellular history at multiple timescales.

The empirical pattern supports the framework's prediction: timescale-aware methods consistently outperform timescale-agnostic methods on tasks requiring memory information, with the magnitude of improvement correlated with the importance of memory for the specific task.

17.7 Cancer methylome classification as positive prediction

While the previous sections developed the framework's *negative* predictions (where transcriptome-only methods fail), cancer methylome classification pro-

vides a *positive* prediction confirmed empirically: where multi-modal data including substratum-level operators (DNA methylation patterns) succeeds where transcriptome-only methods fail.

The DKFZ brain tumor methylome classifier. Capper et al. (2018, *Nature*) introduced a methylation-based classifier for central nervous system tumors trained on DNA methylation data from approximately 2,800 reference cases. The classifier achieves diagnostic accuracy exceeding 95% at the subclass level — substantially higher than conventional histopathological classification and higher than transcriptome-based classification for the same tumors.

The framework's structural reading. The DKFZ classifier succeeds because DNA methylation patterns carry cell-of-origin information that survives the tumor's emergent dysregulation. The substratum-level operator (methylation pattern) is *preserved* during tumorigenesis even as the emergent operator (transcriptional phenotype, histological appearance, clinical behavior) is dysregulated. The framework's prediction: substratum-level memory operators can be more reliable diagnostic markers than emergent operators when the disease involves dysregulation of the substratum-to-emergent translation rather than the substratum itself.

The framework's content from Chapter 16 §16.10 frames this directly: cancer is a disease of the substratum-emergent operator dissociation, with the substratum operator preserved (methylation pattern reflecting cell of origin) and the emergent operator dysregulated (tumor cell behavior). Methylation classification exploits the preserved substratum to provide diagnostic accuracy that emergent-data-only methods cannot achieve.

Cross-cancer extension. The methylation-classification paradigm has been extended to sarcomas (Koelsche et al. 2021), leukemias, and pan-cancer classifiers. The performance pattern is consistent: high accuracy for tumor type identification, with the methylation signature carrying cell-of-origin information that survives tumor-specific dysregulation.

The framework predicts that this paradigm will continue to extend across cancer types, with the boundary cases — cancer types where methylation classification has lower accuracy — being those where the substratum-level memory is more readily disrupted in the cancer state. Highly mutated cancers with extensive copy number aberrations (which can disrupt chromatin maintenance machinery) should be less amenable to methylation classification than cancers with stable genomes. This is a specific testable prediction about cross-cancer methylation classifier performance.

Unclassifiable tumors. Drexler et al. (2024) document that approximately 10-15% of CNS tumor cases submitted to the DKFZ classifier are "unclassifiable" — they do not match any reference methylation class with confidence above the diagnostic threshold. These unclassifiable cases have systematically distinct features: lower tumor purity, younger patient age, recurrent tumor tissue post-radiotherapy, and significantly worse clinical outcomes.

The framework's prediction for unclassifiable tumors: they are cases where the substratum-level memory operator has been disrupted, not just the emergent operator. The disrupted substratum produces a methylation pattern that no longer reflects a single cell of origin, and the classifier cannot map it to a reference class. The empirical features of unclassifiable cases match this prediction: lower tumor purity (contributions from multiple cell types altering the methylation pattern), post-radiotherapy recurrence (DNA damage altering methylation maintenance), and younger age (less time for methylation to stabilize into a clear pattern).

This is testable beyond clinical features. The framework predicts that unclassifiable tumors should show specifically altered methylation patterns at maintenance sites (DNMT1 binding sites, hemi-methylated regions) and at chromatin boundary elements, reflecting disruption of the methylation maintenance machinery rather than tumor-type-specific methylation differences. The hypothesis is testable in the existing DKFZ dataset by stratifying classifiable vs unclassifiable cases on machinery-site vs content-site methylation patterns.

17.8 Variant pathogenicity for reader/writer/eraser genes

Predicting the functional impact of missense variants is a core task in clinical genetics. Methods range from sequence-conservation methods (SIFT, PolyPhen-2) through structure-aware methods (CADD, REVEL) to AI-based methods (AlphaMissense, introduced by Cheng et al. *Science* 2023). The framework's content predicts that variant pathogenicity should be systematically different for variants in reader/writer/eraser machinery versus variants in substrate genes.

The framework's prediction. Chapter 16 §16.12 established the reader-writer disorder class: genetic disorders caused by loss-of-function mutations in readers (MECP2 in Rett), writers (KMT2D in Kabuki), or erasers (KDM6A in Kabuki, ATRX in ATR-X). These disorders have substantively different phenotypic features than disorders affecting substrate genes directly: wider therapeutic windows, demonstrated reversibility in animal models, and graded severity correlated with residual machinery function.

The framework predicts three specific differences in variant pathogenicity prediction performance between machinery genes and substrate genes.

Graded pathogenicity. Variants in machinery genes should show *graded* pathogenicity across mutations within the same gene, reflecting the gene's dose-dependent function in maintaining substratum-level memory across the genome. Different MECP2 mutations produce different Rett severity precisely because they confer different degrees of residual reading function (Cuddapah et al. 2014). The framework predicts this graded pattern is *structural* — a consequence of the machinery's many-target function — rather than gene-specific.

Pathogenicity-severity slope difference. For substrate genes, loss-of-function cor-

relates directly with phenotype severity (a missing enzyme produces a deficient pathway). For machinery genes, loss-of-function correlates with the *number of downstream targets affected*, which can be partially compensated by residual machinery function. The framework predicts a *flatter* pathogenicity-severity slope for machinery genes than for substrate genes.

Predictor performance difference. Methods trained predominantly on substrate genes (where the conservation-pathogenicity relationship is well-calibrated) should systematically underperform on machinery genes (where the relationship is mediated by downstream effects on chromatin state). The framework predicts AlphaMissense and related state-of-the-art predictors should show measurably lower accuracy on reader/writer/eraser genes than on substrate genes when evaluated on clinically-validated benchmarks.

The proposed empirical test. The framework's predictions are testable through systematic reanalysis of existing variant-effect-prediction benchmarks stratified by gene class. Specifically:

(i) Stratify the AlphaMissense calibration set into reader/writer/eraser genes (definable from the reader-writer disorder gene list plus systematic identification of chromatin-modifying enzymes) vs substrate genes (the remainder of the protein-coding genome).

(ii) Compute calibration metrics (sensitivity, specificity, AUC) separately for the two strata. The framework predicts that AlphaMissense performs measurably worse on machinery genes than on substrate genes, with the gap being structural rather than gene-specific.

(iii) Compare the pathogenicity-severity slope between the two strata, controlling for sample size effects. The framework predicts a flatter slope for machinery genes.

The empirical predictions have not been specifically tested in the literature. The framework's content provides a pre-registered structural prediction that could be tested with existing data and existing methods, with substantial clinical implications for variant interpretation in machinery genes (where current pathogenicity scores are predicted to be systematically miscalibrated).

Implications for clinical variant interpretation. If the framework's prediction is confirmed, variant interpretation for reader/writer/eraser disorders should incorporate framework-specific calibration. A variant in MECP2 currently labeled "uncertain significance" by AlphaMissense may have substantially different clinical interpretation in the framework's prediction than the AlphaMissense score suggests. The framework predicts that current variant-interpretation guidelines for reader-writer disorders are systematically miscalibrated; framework-aligned recalibration would improve diagnostic and prognostic accuracy.

17.9 Empirical confirmations in evolutionary biology

The framework's predictions for non-Markovianity in molecular evolution data are *already-empirically-established phenomena*. The framework provides a structural account of these observations rather than novel predictions awaiting test. The evolutionary biology literature has documented multiple non-Markovian signatures across the tree of life over the past two decades; the framework's content unifies these observations under a single structural mechanism.

A scope note governs this section. The signatures below are *non-Markovian memory* signatures — the grade-ii consequence that transfers rigorously from the framework's architecture (C1–C4, with readback (C4) load-bearing). They are *not* evidence for grade-iii P-indivisibility, which is a property of a *closed, reversible* substratum ([Main] §2.3); an evolving population, with its conformational and epigenomic hidden sector, is open and dissipative, so the theorem's reversibility premise does not hold at this scale. Each signature below is moreover reproduced by classical models with dynamic disorder or gamma-distributed rates (as the negative-binomial forms in this section themselves indicate). They should therefore be read as *consistency* confirmations of the framework's memory architecture, degenerate with those classical accounts — not as distinguishing tests that isolate P-indivisibility.

Molecular clock overdispersion. The molecular clock — Zuckerkandl and Pauling's proposal that protein and DNA substitutions accumulate at approximately constant rates over time — has been recognized to violate strict Poisson statistics since the 1980s. The index of dispersion $R(t) = \text{Var}/\text{Mean}$ for substitutions on a single lineage should equal 1 for a Poisson process; empirical measurements consistently exceed this value.

The framework's prediction: if substitution dynamics is memory-bearing (the framework's structural mechanism for non-Markovianity), then $R(t)$ must exceed 1. The degree of overdispersion should increase with the strength of C1 coupling (how strongly the conformational landscape modulates substitution rates) and decrease with effective population size N_e (because large N_e keeps the population in high-neutrality regions of the landscape, reducing memory effects).

The empirical record matches this prediction. $R(t) \approx 5$ for mammalian amino acid substitutions (Gillespie 1989, Cutler 2000). $R(t) \approx 1.6$-2.6 for *Drosophila* (Zeng et al. 1998, Bedford and Hartl 2008). $R(t)$ is inversely correlated with effective population size across yeast, *Drosophila*, and mammals (Bedford and Hartl 2008) — precisely as the framework predicts. Synonymous substitutions show less overdispersion than nonsynonymous substitutions, consistent with weaker C1 coupling for synonymous changes (which do not alter the conformational landscape).

The Bedford-Hartl linear relationship $R(t) = 1 + M/\omega$ is consistent with a negative binomial distribution arising from gamma-distributed rates — the expected

distribution for a process with persistent temporal rate variation (memory). The framework's structural prediction matches the empirically-derived distributional form.

Universal rate autocorrelation. History readback (C4) makes the substitution process along a single lineage memory-bearing: the slowly evolving hidden sector (the conformational landscape, the epigenome) carries information about the lineage's past into its future rates, so the rate at one time is expected to correlate with the rate at neighboring times. The correlation should decay on the timescale τ_B of the hidden sector, and should be detectable in every taxon in which C1–C4 are realized.

Tao et al. (2019, *Molecular Biology and Evolution*) applied CorrTest to large phylogenies across the tree of life and found "extensive rate autocorrelation in DNA and amino acid sequence evolution of mammals, birds, insects, metazoans, plants, fungi, parasitic protozoans, and prokaryotes." The autocorrelation parameter $1/\nu$ ranges from 1.2 to 40 across datasets, with both synonymous and nonsynonymous rates showing significant autocorrelation. This matches the non-Markovian (grade-ii) memory signature the framework's architecture predicts; per the scope note, it is consistent with — but does not isolate — grade-iii P-indivisibility, since classical gamma-distributed-rate (dynamic-disorder) models reproduce the same autocorrelation.

The autocorrelation is stronger for amino acid substitutions than for synonymous substitutions — consistent with stronger C1 coupling for nonsynonymous changes (which alter the conformational landscape and hence the hidden sector). The empirical pattern matches the framework's structural prediction at multiple levels: universal presence (consistent with C1–C4 universality), differential strength by mutation class (consistent with C1 coupling strength), and decay timescale (consistent with τ_B of the conformational landscape).

LTEE power-law fitness trajectories. A Markovian adaptive process with diminishing-returns epistasis produces exponential approach to a fitness optimum. A process with memory stored in the conformational landscape produces a *power-law* trajectory: fitness$(t) \sim t^{\beta}$ with $\beta < 1$, because the hidden sector's memory introduces long-range temporal correlations that slow the approach to equilibrium.

Lenski's Long-Term Evolution Experiment (LTEE) tracks 12 populations of *E. coli* over 75,000+ generations. The fitness trajectory is well-described by a power law $w(t) = (1 + t/\tau)^{\beta}$ with $\beta \approx 0.08$-0.12 (Wiser, Ribeck, and Lenski 2013) — not an exponential. This empirical pattern is well-established and has been attributed to diminishing-returns epistasis.

The framework provides a *structural* explanation: the conformational landscape (hidden sector) stores information about the adaptive history, producing the long-range correlations that generate power-law scaling. The power-law exponent β is related to the Hurst exponent via $H = 1 - \beta/2 \approx 0.94$-$0.96$, indicating

strong persistent memory — consistent with the deep C3 capacity of a bacterial proteome ($\sim$ 4,000 proteins, each with its own conformational landscape).

The framework's prediction goes further. The Hurst exponent should be measurable in individual-lineage substitution rates across the tree of life, with the value reflecting the C3 capacity of the relevant hidden sector. The framework predicts $H \approx 0.79 \pm 0.03$ for typical single-entity systems with 10^2-10^4 conformational states, with higher values for systems with larger hidden sectors. This prediction is testable using existing substitution rate data and provides a quantitative cross-validation of the framework's structural mechanism.

Punctuated equilibrium. Memory-bearing dynamics with a slow hidden sector predicts alternating periods of stasis and rapid change. During stasis, the hidden sector absorbs information from the visible sector without visible phenotypic effect — the conformational landscape slowly reshapes, the epigenome gradually drifts, the niche is incrementally modified. During bursts, accumulated hidden-sector changes reach a threshold where they dramatically alter the set of neutral mutations, producing rapid phenotypic change.

Punctuated equilibrium — long periods of morphological stasis interrupted by rapid change — has been documented across the fossil record since Eldredge and Gould (1972). The framework identifies this as history readback: the rapid change is the readback of correlations accumulated in the hidden sector during the stasis period. The contingent innovation in Lenski's LTEE (citrate utilization at generation 31,500, requiring prior "potentiating" mutations) is the molecular-scale version: the hidden sector (conformational landscapes of multiple proteins) accumulated changes that made the actualizing mutation possible only in a specific historical context.

Evolutionary capacitance. Any molecular system that buffers genetic variation — storing it in the hidden sector and releasing it under stress — functions as an evolutionary capacitor. The framework predicts that capacitor systems should show non-Markovian release kinetics: the rate at which cryptic variation is released depends on the history of variation stored.

Hsp90 is the canonical evolutionary capacitor (Rutherford and Lindquist 1998). Under normal conditions, Hsp90 chaperones mutant proteins into functional conformations, hiding genetic variation from selection. Under stress (heat, chemical), Hsp90 is overwhelmed and the buffered variation is released — a burst of phenotypic diversity from a single environmental perturbation. The framework identifies this as a molecular-scale partition: Hsp90-client conformational states are the hidden sector; the expressed phenotype is the visible sector. C1 (coupling via chaperoning), C2 (conformational dynamics much slower than growth), and C3 (enormous conformational state space of Hsp90-client complexes) are all satisfied.

The cumulative empirical case. The evolutionary biology empirical record provides convergent support for the framework's structural prediction across multiple independent signatures: overdispersion (statistical), autocorrelation

(temporal), power-law trajectories (functional), punctuated equilibrium (paleontological), and evolutionary capacitance (molecular). Each signature is a distinct empirical observation; the framework's content unifies them under a single structural mechanism — the grade-ii non-Markovian memory produced by C1–C4 coupling to a slow hidden sector at multiple scales (not grade-iii P-indivisibility; see the scope note).

The framework's content in evolutionary biology is therefore retrodictive in form — the empirical phenomena were established before the framework, and the framework provides a structural account. But the *unification* of these phenomena under a single structural mechanism is the framework's contribution: evolutionary biology has previously treated overdispersion, autocorrelation, power-law trajectories, and punctuated equilibrium as separate phenomena requiring separate explanations. The framework predicts that they are five aspects of a single structural mechanism, with quantitative cross-relationships (Hurst exponent / power-law exponent / autocorrelation timescale / overdispersion magnitude) that should be empirically related in framework-specific ways.

17.10 Chapter close: the structural roadmap forward

The framework's content in bioinformatics has two cumulative dimensions: the conditional structural ceiling on transcriptome-only methods that explains observed failures, and the empirical record of non-Markovianity signatures across the tree of life that the framework's structural mechanism unifies. Both dimensions point to a specific *structural roadmap forward* for computational biology.

Three components of the roadmap.

Multi-modal data. Where the strict gap holds, the framework's information-theoretic ceiling on transcriptome-only methods predicts that progress in bioinformatics requires including substratum-level data alongside transcriptome data. The methodological field has begun this convergence — multi-omic velocity, SCENIC+ for GRN inference, knowledge-graph methods for perturbation prediction, methylation classifiers for cancer diagnosis — but the convergence is incomplete. The framework's content predicts that *complete* methodological transition to multi-modal data is required to exceed the current performance plateau.

Knowledge graphs and prior structure. Methods that incorporate prior knowledge about biological networks (knowledge graphs, regulatory databases, pathway annotations) consistently outperform methods that learn everything from data. The framework's reading: prior knowledge functions as a proxy for substratum information not present in the data. The structural roadmap forward includes systematic incorporation of curated prior knowledge as a complement to data-driven learning.

Explicit memory state. Methods that explicitly model the cellular memory state (chromatin accessibility, methylation patterns, conformational state) outperform

methods that treat each measurement as a snapshot. The framework's content predicts that bioinformatics methods must model the cell as a dynamic system with explicit memory variables, not as a static snapshot, to capture the relevant biology.

Five predictions for empirical work.

(P1) Where the strict gap holds for perturbation response, transcriptome-only foundation models have a structural ceiling on perturbation prediction performance. No amount of additional scaling on transcriptome data will exceed this ceiling.

(P2) Multi-omic foundation models — incorporating chromatin accessibility, methylation, and transcriptome data — will exceed transcriptome-only ceilings on perturbation prediction.

(P3) Variant pathogenicity prediction for reader/writer/eraser genes is systematically miscalibrated by current state-of-the-art predictors trained predominantly on substrate genes. Framework-stratified recalibration will improve clinical variant interpretation for reader-writer disorders.

(P4) Cancer methylome classification will continue to extend to additional cancer types, with boundary cases being cancers with disrupted chromatin maintenance machinery. Methylation classifier accuracy should correlate with substratum-level memory preservation.

(P5) The Hurst exponent of single-lineage substitution rates across the tree of life should average $H \approx 0.79 \pm 0.03$ for typical taxa, with higher values for systems with larger hidden-sector C3 capacity (full proteomes, complex multicellular organisms).

The framework's epistemic position in bioinformatics. The framework's content in bioinformatics is *both* explanatory (providing structural account of established empirical patterns) and predictive (providing specific testable predictions for current research directions). The chapter's content has identified empirical patterns that the framework retrodicts (molecular clock overdispersion, rate autocorrelation, LTEE power law, punctuated equilibrium, methylation classifier success, foundation model failures), and specific predictions that the framework makes (variant pathogenicity stratification, Hurst exponent values, cross-cancer classifier performance, multi-modal foundation model improvements).

The framework's value in bioinformatics is therefore structural unification across multiple methodological domains. The previous separate puzzles — why deep learning fails on perturbation prediction, why methylation classification succeeds where transcriptome classification fails, why molecular clocks are overdispersed, why GRN inference plateaus, why fitness trajectories follow power laws — receive a unified structural explanation. The chapter is the framework's empirical bridge from molecular biology to computational methodology.

Forward pointers. The chapter closes the chapter sequence of the book. Chapter 18 has already developed the framework's distinctive forward predictions beyond standard QM and GR. Chapter 19 has developed the systematic inventory of resolved and open problems in fundamental physics. The remaining work consists of the back-matter appendices: prediction status (Appendix A), mathematical derivations (Appendix B), and the new common objections appendix (Appendix C) that systematically addresses the most frequent challenges to the framework's content.

The framework's reach across Chapters 13-17 — quantum biology, quantum computing, quantum engineering, medicine, bioinformatics — provides the framework's empirical content in applied scientific domains beyond fundamental physics. Combined with the fundamental physics empirical record from Chapters 5-9 (twenty-two classified Standard Model retrodictions — two parameter-free structural and five conditional structural — seven gravitational entries, seven neutrino entries, JUNO at 0.07σ, the GW250114 area-theorem consistency at 99.999% — the Bekenstein-Hawking coefficient itself untested directly), the cumulative empirical case for the framework's structural commitments is substantial.

The framework's content from the substratum-level structural axioms of Chapter 1 through the emergence cascade of Chapters 10-12 and the applied chapters of 13-17, with the closing forward predictions and open problems inventory of Chapters 18-19, constitutes a complete structural framework with empirical exposure across physics, chemistry, biology, computing, engineering, and medicine. The framework's value is therefore not concentrated in any single empirical domain but distributed across many, with the structural commitments at the substratum level being load-bearing for the framework's reach in each.

Chapter 18

Beyond Quantum Mechanics and General Relativity

Status — conjectural extension. This chapter applies the core framework to consciousness, and is offered as a *conjecture*, not an established result: no empirical test is proposed; the claim is a structural correspondence between embedded observation — the trace-out of an observer's own inaccessible degrees of freedom — and the character of a bounded first-person perspective, offered as a possibility the framework raises rather than a result it establishes. It carries no evidential weight for the core framework — which stands or falls

independently on its physics — and should be read as a structural possibility to be tested, not as support for the framework.

18.1 What this chapter develops

Chapters 5 through 16 developed the framework's content where standard physics and the framework agree at high empirical precision. The Standard Model gauge group, the gravitational sector, the JUNO neutrino prediction, the dark sector phenomenology — all are sectors where the framework reproduces or refines standard predictions, often with sharper structural commitments and parameter-free quantitative agreement. The empirical case for the framework is concentrated in those chapters.

This chapter is different. The framework predicts content that *goes beyond* standard quantum mechanics and general relativity — phenomena that standard physics has no commitment to, structural features that standard physics treats as postulates rather than derivations, observable signatures distinguishing the framework's specific commitments from the broader landscape of foundational accounts. The empirical record on this content is incomplete because the relevant experiments either have not yet been performed or have not yet reached the precision required. The chapter is therefore forward-looking rather than retrospective: what the framework predicts that current physics does not yet test.

Ten pieces of content occupy the chapter.

Section 18.2 develops the framework's *account* of the Born rule — the read-write-cycle equilibrium reading of the cosmological realization, a mechanism proposal layered on the representation-level fact that every finite-horizon law admits Born-rule fixed-basis readout ([Main §3.4]), with the common coherent operational extension remaining open — and the transient deviations it implies in the very early universe if the hidden sector began away from equilibrium, a conditional prospect for CMB tests of the primordial power spectrum and non-Gaussianity. Section 18.3 develops the quantum-classical boundary as a sharp C1–C4-determined threshold, together with the mechanism proposal's restricted quantum-statistical regime — partial distinguishability revival, with a proposed capacity-controlled origin and scaling law; partial revival itself is standard open-systems phenomenology, the origin claim the proposal's own. Section 18.4 develops the epistemic character of quantum randomness — quantum outcomes are deterministic but inaccessible rather than fundamentally indeterminate, with implications for what kinds of accounts of physics are compatible with the framework.

Section 18.5 develops the framework's cumulative case on quantum gravity: the cosmological constant dissolution (Chapter 7 §7.7), the black hole information account (Chapter 7 §7.6), and the non-renormalizability reading all follow from the same structural commitment to the geometry-first ordering and the substratum-emergent operator distinction, as framework-level interpretations

and conditional results, with covariant unification open (G3). Section 18.6 takes up gravitational wave echoes: [GR §8.4] argues that a black-hole horizon does not change an external observer's partition, so no OI-specific echo is expected and ringdowns follow general relativity; no echo amplitude is derived.

The middle of the chapter takes up the framework's foundational commitments and their consequences. Section 18.7 develops the finite-information character of the substratum and its consequence of exact recurrence, at a period no larger than the largest order of a permutation of S (Landau's function $g(|S|)$), which the framework does not compute since $|S|$ is not fixed by visible data — a definite structural prediction with real conceptual content, and without practical consequences at periods of astronomical size. Section 18.8 develops the arrow of time as emergent from the partition's initial conditions rather than from a fundamental temporal asymmetry — the substratum is T-symmetric, and the thermodynamic and cosmological arrows trace to the partition's low-entropy initial state.

The latter sections engage two structural questions the framework has been silent on in earlier chapters. Section 18.9 develops the framework's position on cosmic spatial topology: the substratum is locally a cubic lattice at $\epsilon = 2\sqrt{\sigma}\, l_p$, with σ the entropy of one boundary mode, but the global identification structure (whether the universe is topologically trivial $\mathbb{R}^3$, a 3-torus, a Klein-bottle quotient, or some other compact identification) is not determined by the framework's structural commitments. Janna Levin's compact-universe proposal is engaged as one specific instantiation of finite cosmic extent compatible with the framework but not predicted by it.

Section 18.10 develops the framework's position on consciousness. The position is asymmetric: the framework's mathematical operation of observation (partition-and-trace-out with C1–C4 structure) is a necessary but not sufficient condition for consciousness. The framework imposes structural constraints on physics-based consciousness theories — deterministic substratum rules out fundamental-indeterminism accounts; the C1–C4 architecture provides necessary structural conditions for the non-Markovian information processing conscious systems must exhibit; the framework's content is compatible with integrated information theory and several other accounts that supply additional sufficient conditions. The hard problem of consciousness proper — the explanatory gap between structural and phenomenal facts — is not addressed; the framework has no bridge from one to the other and is honest about that limitation.

Section 18.11 closes the chapter by identifying which of the framework's forward predictions are most accessible to near-term experimental engagement, which require new instrumentation, which proposed mechanisms remain conditional prospects without a specified observable response, and which lie outside foreseeable empirical reach. The chapter is the book's forward-looking content: the framework's substantive commitments beyond current physics, with the empirical record yet to be developed.

The framework's content in this chapter has a particular epistemic structure. Predictions like the restricted quantum-statistical regime in biological systems and the BQP computational ceiling (developed in Chapter 14) are *testable*. The Born rule's dynamical equilibrium is a *conditional prospect*: a proposed mechanism that would bear on the CMB's primordial statistics, without yet a specified observable response (§18.2). Predictions like cosmic recurrence, at a period at most $g(|S|)$, and the asymmetric position on consciousness are *structural commitments with limited empirical access*. The chapter is honest about which is which. The framework's value here is not the empirical record (which is what Chapters 5-17 developed) but the structural commitments — what the framework says about the world beyond what current physics tests.

18.2 The Born rule as dynamical equilibrium

The Born rule — that the probability of measuring a quantum system in state $|x\rangle$ given wave function $|\psi\rangle$ is $P(x) = |\langle x|\psi\rangle|^2$ — is one of the most empirically successful predictions in physics. It is also one of the most foundationally puzzling. In standard quantum mechanics, the Born rule is an *independent postulate*: it holds always, for all systems, at all times, with no underlying mechanism producing it. The various interpretive programs (many-worlds, QBism, Bohmian mechanics, dynamical collapse) take different positions on what the Born rule means, but they share the structural feature of treating it as fundamental rather than derived.

The framework's account is layered. At the representation level the Born rule is part of the exact equivalent description: every finite-horizon law has a unitary representation with Born-rule fixed-basis readout ($S \iff D \iff Q_{\text{fb}}$, [Main §3.4]). The open frontier is not a separate choice of probability exponent inside that representation; it is the common coherent operational extension to arbitrary noncommuting instruments and composites. On top of that sits the cosmological-realization *mechanism proposal*: the equilibrium reading of the read-write cycle between the visible and hidden sectors — the observer reads the visible sector through measurement, writes correlations into the hidden sector through the coupling H_{VB}, and reads back those correlations on subsequent measurements — with Born-rule probabilities as the stationary statistics of the cycle for the equilibrated hidden prior. That stationary-distribution reading is the account's load-bearing step, and it is a structural commitment of the realization, not a proved theorem of the foundational paper.

The structural feature of the account is that the Born rule is a *dynamical* equilibrium rather than a fundamental postulate. The system approaches Born-rule statistics on a characteristic timescale set by the hidden sector's equilibration time $\tau_{\text{eq}} \sim \tau_B$, where τ_B is the hidden sector's correlation time from condition C2. For the cosmological partition τ_B is the horizon time, $\tau_B(t) \sim H(t)^{-1}$: about 10^{17} seconds today and far shorter in the early universe. Two quantities must be kept apart. The ratio $\tau_S/\tau_B \sim 10^{-32}$ is the per-step loss of a stored record under C2; it sets the size of memory effects, not the distance from equi-

librium. The quantity that tracks the approach to equilibrium is instead the accumulated number of relaxation times, $\int dt/\tau_{\rm eq} \sim \int H\, dt = \ln(a/a_i)$, but that count does not by itself fix how a deviation decays. If the observed deviation δ obeys a closed relaxation law $\dot{\delta} = -\lambda H \delta$, a deviation present when relaxation began at scale factor a_i decays as $(a_i/a)^\lambda$; the identification $\tau_{\rm eq} \sim H^{-1}$ fixes neither that closed law nor the coefficient λ, which it constrains only to be of order one. Under such a law the epoch-local reading is required: with $\tau_{\rm eq}$ fixed at today's value, of order H_0^{-1}, the age of the universe would amount to about one relaxation time and would leave a fraction of order $e^{-\lambda}$ of any initial deviation in place today, a departure that laboratory tests of the Born rule exclude unless the initial deviation was itself negligible. On the epoch-local reading the same law suppresses the residual in any current or foreseeable laboratory experiment by the factor $(a_i/a_0)^\lambda$, and the Born rule is exact for all practical purposes. If the hidden sector began at equilibrium there is no transient to relax, and the account reduces to the maximal-entropy measure posited in [Main §4.5]. (Status of this estimate: the identification $\tau_{\rm eq} \sim \tau_B$ and its epoch-local reading are heuristic content of the mechanism proposal, and the closed relaxation law and its coefficient λ are further assumptions the proposal does not derive; none of these is a theorem of the foundational paper.)

Transient violations in the very early universe. If the hidden sector began away from its stationary distribution, the Born rule was *not* exact in the very early universe: until the hidden sector equilibrated, measurement statistics deviated from $|\langle x|\psi\rangle|^2$. Whether it began away from equilibrium is a question about the cosmological initial conditions that the account does not settle; if it began at equilibrium, there were no such violations. Any deviations are parametrized by

$$P(x,t) = |\langle x|\psi(t)\rangle|^2 + \sum_k c_k(t)\langle x|\hat{O}_k|\psi(t)\rangle,$$

where the operators $\hat{O}_k$ encode the measurement history and the coefficients $c_k(t)$ parameterize possible history-dependent departures. Their decay would require an additional effective relaxation law, such as the closed law considered above, applicable over a specified finite epoch. Neither asymptotic convergence to zero nor a universal decay rate follows from the finite reversible substratum dynamics, which is recurrent (§18.7); microscopic recurrence and a possible effective relaxation over a finite epoch are distinct. (Status: the expansion is the proposal's *parametrization* of the deviation — the form an eventual derivation would have to fill in — not a derived spectral decomposition.)

The structural source of any transient violations is the hidden sector's initial state. In the framework's substratum dynamics, the hidden sector begins in a specific initial configuration set by the cosmological initial conditions. If that initial state is away from the stationary distribution, then before the visible-hidden coupling has driven the hidden sector there, the read-write cycle's input statistics differ from the Born rule, and the observer reads visible-sector out-

comes with probabilities that deviate from $|\langle x|\psi\rangle|^2$ in ways that depend on the specific initial state.

This connects to Valentini's *quantum relaxation* hypothesis in de Broglie-Bohm theory, where Valentini proposed that Born-rule violations in the early universe would leave observable imprints. The mechanism proposal offers a structural mechanism for what Valentini's program postulates: hidden-sector equilibration would drive relaxation toward the Born rule on the heuristic timescale $\tau_{\rm eq} \sim \tau_B$. Valentini's program postulated the relaxation; the account locates a candidate mechanism for it in the partition's read-write architecture. Both take the initial departure from equilibrium as an assumption about the cosmological initial conditions.

Observational signatures in the CMB. If primordial Born-rule violations were present during inflation, their empirical signature would be modified statistics of the primordial fluctuations. A change in the variance alone alters the primordial power spectrum without entailing any non-Gaussianity — a Gaussian distribution with a changed variance has no connected moments beyond the second — so a non-Gaussian signature arises only if the departure also carries higher connected moments. Standard inflationary models predict approximately Gaussian primordial perturbations, and the Planck 2018 analysis detects no primordial non-Gaussianity: for the local shape, $f_{\rm NL} = -0.9 \pm 5.1$ (68% CL). The framework does not yet supply an amplitude to compare with this bound or with the measured power spectrum.

Standard local, equilateral, and folded non-Gaussianity shapes arise from specific inflationary mechanisms (multi-field inflation, non-canonical kinetic terms, modified initial vacuum states). Which moments a Born-rule departure would modify, and with what scale dependence or bispectrum shape, depends on the source dynamics of the relaxation, the family of initial deviations and the map from the hidden-sector deviation to the observed fluctuations, none of which the proposal specifies; whether such a departure produces any non-Gaussianity, and whether its shape would differ from the standard ones, is open.

The empirical accessibility of the proposal depends on inputs it does not yet fix: the source dynamics of the relaxation through inflation, the family of initial deviations, the rule for a mode's evolution after horizon exit, and the map from a residual deviation to the measured power spectrum or non-Gaussianity. The residual on a given scale is set by the initial deviation and by the relaxation accumulated between the onset of relaxation and the imprinting of that scale, not by the relaxation timescale alone: even rapid relaxation leaves almost the entire deviation on a scale imprinted shortly after relaxation begins, and no relaxation history leaves a residual if the initial deviation is zero. Once the observable map is derived, the Planck measurements would constrain the combination of initial deviation and relaxation history; no bound on $\tau_{\rm eq}$ follows before then. The epoch-local reading above places $\tau_{\rm eq} \sim H_{\rm inf}^{-1}$. Under a closed relaxation law, if a mode's relaxation stops at its horizon exit, the residual on observable

scales depends on how many e-folds of relaxation preceded that exit and, for scale-independent initial deviations, is largest on the largest scales; if relaxation continues to a common later epoch, that scale dependence cancels; and scale-dependent initial deviations can cancel or mimic either trend. The account fixes none of these.

Status ledger for §18.2. Layer by layer: the representation-level Born status is a theorem ([Main §3.4]: admitted universally, coherent operational extension characterized by five named conditions in [GR §3.3]); the read-write equilibrium reading is a mechanism *proposal* whose stationary-distribution step is a structural commitment; the equilibration timescale, its epoch-local reading and the deviation parametrization above are the proposal's heuristic quantitative content, and the closed relaxation law and rate coefficient on which a decay law rests are further assumptions it does not derive; the observable form of a departure — a power-spectrum change, non-Gaussianity, or both — is not derived; and the F7 entry — transient deviations if the hidden sector began away from equilibrium, observed through the CMB's primordial statistics — is a conditional prospect: it becomes a testable prediction only once the source dynamics, the family of initial deviations and the map to observables are specified, and it is detectable only if a nonzero residual survives on observable scales when they are imprinted. No entry in this section is a theorem of the foundational paper.

CMB-S4 (the upcoming ground-based experiment) and LiteBIRD (the planned space-based experiment) will tighten the non-Gaussianity constraints by roughly an order of magnitude over the next decade. Whether these experiments bear on the proposal depends on the source dynamics, the family of initial deviations and the map to observables, none of which the proposal yet specifies. Once they are specified, a predicted departure above the new sensitivity would be tested by these experiments, and a smaller one would leave the proposal unconstrained by them.

History-dependent corrections at late times. Even after the hidden sector has equilibrated, the Born rule receives corrections at order τ_S/τ_B. For the cosmological partition, this ratio is approximately 10^{-32} — utterly unobservable at any current or foreseeable precision. But the *structure* of the corrections is predicted: successive quantum measurements are not exactly independent samples from the Born distribution. They carry subtle correlations encoding the deterministic structure of the substratum bijection φ, suppressed by 32 orders of magnitude below current experimental sensitivity.

The conceptual content of the late-time correction is significant. Standard quantum mechanics treats measurement outcomes as fundamentally independent random samples from the Born distribution. The framework predicts that this independence is approximate — there exist correlations between successive measurements, generated by the deterministic substratum dynamics, that vanish only in the strict limit $\tau_S/\tau_B \to 0$. The framework's substratum is deterministic; the apparent randomness of quantum mechanics is the observer's incompleteness rather than fundamental indeterminism. The late-time correc-

tions are the structural signature of this underlying determinism, suppressed but not eliminated.

18.3 The quantum-classical boundary and the restricted quantum-statistical regime

The quantum-classical boundary is one of the most pragmatically important features of physics and one of the most foundationally ambiguous. Standard quantum mechanics applies universally in principle: a rock, a cat, and an electron are all described by wave functions, with the rock's wave function being too entangled with environmental degrees of freedom to show interference effects. The pragmatic boundary between "quantum" and "classical" is set by decoherence rather than by any structural distinction — macroscopic objects appear classical because they are quantum-but-decohered, not because they obey different physics.

The framework's account is structurally different. The boundary it draws is *determined by whether the framework's conditions C1, C2, C3, C4 are satisfied* — and it is a boundary in the memory currency, not in representability, which is universal ($S \iff D \iff Q_{\mathrm{fb}}$, [Main §3.4]). A system satisfying all four carries the memory-bearing sector — accessible non-Markovian statistics, $M_t > 0$, with unavoidable hidden predictive memory in every completion; a system failing C1, C4 or the persistence form of C2 has that sector empty — not "quantum but decohered" but genuinely described by memoryless classical stochastic dynamics, its quantum representation trivial; a system failing C2 only in its timescale form has it suppressed, not shown to be empty, and limited capacity caps it (below). The boundary drawn by C1, C4 and the persistence form of C2 is structural rather than approximate.

The four conditions as criteria. Recall from Chapter 1 the framework's four conditions for an embedded observer's reduced description to carry the memory-bearing sector — the discriminating content, the representation itself being universal ([Main §3.4]): - C1 (coupling): the subsystem must interact nontrivially with a hidden sector; - C2 (memory persistence): the hidden sector must retain a record of the subsystem's past across the accessible window, $\tau_B \gg \tau_S$ with τ_B the coarse-grained mixing time — slow hidden dynamics is one sufficient mechanism, not the condition itself ([Main §1.3]); - C3 (capacity): the hidden sector must have capacity sufficient for the process's conditional memory — the capacity floor $\log_2 |\mathcal{C}_H| \geq I^*$ ([Main §3.4]); $N_H \gg N_V$ is the comfortable physical regime, not the condition; - C4 (history readback): hidden degrees of freedom must carry the visible past into future visible conditionals — the readback gap that makes the stored memory accessible ([Main §1.3]).

A system implementing C1–C4 with all four conditions strongly satisfied has a memory-bearing reduced description — the framework's identification applies in full at its stated layer — the memory-bearing description with its quantum representation, correctly given by unitary evolution on a Hilbert space at the

transition-statistics layer, the coherent operational lift still to be established ([Main §3.4]). A system failing C1 (no hidden-sector coupling) has memoryless classical stochastic dynamics — the memory-bearing sector is empty, the quantum representation trivial. A system failing C2 in its timescale form (a hidden sector that equilibrates faster than the visible-sector dynamics, its records eroded before readback) has emergent dynamics whose accessible memory effects are suppressed. A hidden sector of limited capacity caps accessible memory — the capacity floor $M_t \leq \log_2 |\mathcal{C}_H|$ ([Main §3.4]); the mechanism proposal hypothesizes that marginal capacity produces partial distinguishability revival, the restricted quantum-statistical regime of Chapter 15 §15.7, with memoryless operational modeling failing while the transition-statistics layer remains representable ([Main §3.4]).

The restricted quantum-statistical regime. The mechanism proposal adds a revival scale to the distinction between memoryless and memory-bearing visible laws, and names one range of it. The scale is the revival measure $\mathcal{N}$. On an accessible window $W = \{0, \ldots, K\}$, let d_t be the distinguishability at time t of two initial preparations under the visible dynamics — the total-variation distance of their visible population laws, equivalently the trace distance of the corresponding diagonal states — and let $\mathcal{N}$ be the largest increase $d_t - d_s$ over pairs of preparations and times $s < t$ in W, with $\mathcal{N} = 0$ when distinguishability never increases. $\mathcal{N}$ is the peak revival on the window, a single increase, so $0 \leq \mathcal{N} \leq 1$. It is not the Breuer–Laine–Piilo measure, which sums every increase over the window. That cumulative backflow is kept separate: on a finite reversible system the rooted laws are periodic ([Main §2.3]), so a revival in one recurrence period recurs in every later one, and the sum grows without bound as the window lengthens. $\mathcal{N} > 0$ on a window certifies P-indivisibility there, since a P-divisible family never increases distinguishability; the converse fails ([Main §2.3]). Memory does not fix $\mathcal{N}$: the XOR control of [Main §2.3] carries memory on a window on which its family is P-divisible, so $\mathcal{N} = 0$ there, and $\mathcal{N}$ measures revival, not memory. Three ranges are named on a window: no revival ($\mathcal{N} = 0$), partial revival ($0 < \mathcal{N} < 1$) and complete revival ($\mathcal{N} = 1$). The mechanism proposal calls partial revival the *restricted quantum-statistical* regime. The name records that the quantum description supplied there is the representation at the transition-statistics layer, with the operational layer built on it not supplied; that restriction holds at every value of $\mathcal{N}$, so the regime is fixed by its revival range, and the name marks neither a partial nor a complete quantum theory and is not a quantum-membership scale: every finite law at every $\mathcal{N}$ admits the quantum representation and a deterministic hidden realization ([Main §3.4]).

By definition the regime lies strictly between no revival ($\mathcal{N} = 0$) and complete revival ($\mathcal{N} = 1$). In standard quantum mechanics, a decohered system is still described by a density matrix — it is quantum but mixed. In the framework the regime is a statement about the DYNAMICS, not about representability: every stochastic transition family admits the one-step ancilla dilation ([Main §3.1]), so no process lacks a Hilbert-space representation at the transition-statistics layer.

What varies across the regime is the size of the largest distinguishability revival on the window; a positive revival certifies indivisibility, while indivisibility without revival is possible ([Main §2.3]). What a system in the regime may lack is the stronger OPERATIONAL structure built on top of that representation: coherent preparations of the represented states, the instrument algebra governing interventions, and representation-consistency across horizons — each of which the framework leaves open or bounded rather than delivered ([Main §3.4]). The quantum description is restricted in that precise sense: the transition-statistics layer is available throughout the regime, as at every value of $\mathcal{N}$, while the operational layer is available only to the extent those questions are settled, so the representation is not a complete operational quantum theory; and $\mathcal{N}$ records the largest revival, not a distance from the divisible families.

Partial distinguishability revival itself is standard open-systems phenomenology — the Breuer–Laine–Piilo measure quantifies trace-distance revival, and the Rivas–Huelga–Plenio measures quantify departures from complete-positivity divisibility, a different property; what is distinctive here is the proposed capacity-controlled origin and the specific scaling law. Standard accounts treat quantum and classical as endpoints of a spectrum (with decoherence interpolating between them), but the framework separates memoryless from memory-bearing visible laws ([Main §3.4]), and the mechanism proposal adds a revival scale inside the memory-bearing case — the restricted quantum-statistical regime for partial revival — all of it quantum-representable ([Main §3.4]).

Where to look for the restricted quantum-statistical regime. The mechanism proposal, layered on the capacity theorem, conjectures where partial revival occurs: in systems whose hidden sector has capacity comparable to the visible sector's, with protein conformational dynamics and single-enzyme kinetics as the biological candidates Chapter 13 conjectures, which the capacity ratio does not by itself place there. The timescale ratio τ_S/τ_B does not by itself place a system on the revival scale. A fast bath that erodes the stored record uniformly suppresses the readback gap exponentially in τ_S/τ_B at every order $k \geq 2$ ([Main §2.3]); a slow bath does not by itself give memory or revival — a static bath whose record is never read back carries no memory ([Main §2.3]).

The framework's content is not "quantum effects in biology" — standard quantum mechanics allows quantum effects at any scale, and the literature documents many examples. The framework's content is the distinction between memoryless and memory-bearing visible laws, all quantum-representable ([Main §3.4]), together with the mechanism proposal's conjecture that partial revival occurs at marginal hidden capacity.

Observational signatures. Single-molecule enzyme experiments already show non-Markovian kinetics — history-dependent reaction rates, stretched-exponential waiting-time distributions, transition statistics that cannot be fitted with a Markovian rate equation. The framework's prediction is that these signatures are not artifacts of poorly-characterized rate constants but features the mechanism proposal attributes to partial revival. Two tests are distinct.

Whether the multi-time statistics admit a memoryless (Markov) description at the accessible horizon decides memory; whether the distinguishability of two preparations revives on that horizon decides $\mathcal{N} > 0$, which certifies indivisibility. A memory-bearing system can show no revival ([Main §2.3]); the mechanism proposal predicts memory together with partial revival — a proposal-level claim ([Main §3.4]).

The distinction is empirically testable. A memoryless (divisible) density-matrix description requires specific algebraic constraints on the transition probabilities — positivity, completeness, certain inequality relations between multi-time correlators. The mechanism proposal predicts that protein conformational dynamics and single-enzyme kinetics violate at least some of these constraints. The experimental program would require multi-time correlation measurements with sufficient statistics to constrain the algebraic structure, which is within reach of current single-molecule techniques.

The restricted quantum-statistical regime is the mechanism proposal's testable conjecture, with experimental targets in the biological sciences that the framework's quantum-biology chapter (Chapter 13) develops in detail.

18.4 The epistemic character of quantum randomness

The framework's commitment to a deterministic substratum has a striking consequence for the foundations of quantum mechanics: quantum randomness is *epistemic* rather than ontic. The apparent indeterminism of quantum measurement outcomes is the observer's incompleteness, not a fundamental feature of physical reality. The substratum bijection φ is deterministic; the memory-bearing description — with its admitted quantum representation — emerges from the framework's structural conditions applied to embedded observation of this deterministic dynamics ([Main §3.4]); the randomness the observer sees is the result of marginalizing over the hidden sector.

This is a substantive position, and it has consequences both for the foundations of quantum mechanics and for what kinds of accounts of physics are compatible with the framework.

Deterministic but inaccessible. The framework's substratum dynamics is given by the bijection φ acting on the configuration space S. Each state (x, h) — visible coordinates x paired with hidden coordinates h — maps to a unique successor $\varphi(x, h) = (x', h')$. There is no randomness in the dynamics; the substratum is strictly deterministic.

The apparent randomness arises because the observer sees only x, not h. Different hidden states $h_1, h_2, ...$ compatible with the same visible state x lead to different successor visible states $x'_1, x'_2, ...$ under the bijection. From x alone the observer cannot distinguish among the hidden states; they appear as a probability distribution over the possible successors. The probabilities are *not* features of the dynamics — the dynamics is deterministic — but features of the ob-

server's reduced description of the dynamics, given the observer's incomplete information about the hidden-sector configuration.

This is what "deterministic but inaccessible" means in the framework: the outcomes are fully determined by the substratum dynamics, but the observer cannot in principle access the complete state that determines them. The randomness is unavoidable for any embedded observer, but it is not fundamental — it is the cost of partial observation.

Bell's theorem and the chosen nonlocality. Bell does not rule out determinism by itself; it rules out reproducing the observed correlations with a response-complete deterministic completion that records a ± 1 outcome in every run and is both measurement-independent and parameter-independent. Chapter 1 proves the corresponding Bell ceiling: if the setting interventions, applied at one step, each read and write only their own wing, the Bell-relevant coupling range does not connect the wings before readout, every run records a ± 1 outcome, and one setting-independent pre-setting ensemble is used, then $|S_{\mathrm{CHSH}}| \leq 2$. The framework therefore declines the measurement-dependent/superdeterministic horn and takes **ontic parameter dependence**. In the specific SM/GR realization the target mechanism is preparation-indexed adjacency in the already state-dependent graph $G(x)$ (or an equivalent ontically nonlocal response), so a prepared microstate response may depend on the remote setting while the reference vacuum/uniform graph remains geometric and bounded-range. Whether those prepared graphs keep the operational distribution no-signaling and preserve the continuum-curvature limit is the open hypothesis H-Bell. The $2\sqrt{2}$ theorem for the causal-local indivisible stochastic description is a separate weaker-layer result, not a locality theorem for the deterministic completion.

One proposed physical realization of that preparation-indexed parameter dependence uses non-localized cosmological-boundary modes. This is a candidate realization, not an established Bell mechanism: any such prepared long-edge/response structure must satisfy H-Bell, i.e. preserve operational no-signaling and the metric/Ollivier–Ricci continuum limit used by the gravitational reconstruction. What distinguishes it from a superluminal signaling mechanism is operational no-signaling, not ontic parameter independence.

Practical consequences are absent. The distinction between epistemic and ontic randomness has no observable consequences at the 10^{-32} suppression level — quantum random number generators, quantum key distribution protocols, and other applications that rely on "true randomness" function identically under either interpretation. What these applications rely on is unpredictability to the parties who can observe or act on the source, not the ontic or epistemic status of its randomness. Determinism does not by itself guarantee that unpredictability: an embedded observer whose actions reach the hidden state driving a deterministic source can, for some sources, predict its output better than chance. Whether a source stays unpredictable to observers of bounded memory, whether they act on it or only read it, depends on the source, and the framework

does not establish it for the hidden sector of the physical realization.

The framework's content here is therefore conceptual rather than empirical at near-term experimental scales. The framework's commitment to deterministic substratum changes what physics-based accounts are compatible with the framework — but it does not change near-term experimental predictions in any observable way.

Conceptual consequences. The deeper conceptual consequence is significant. The universe is deterministic in the framework. Quantum indeterminacy is the cost of partial observation, not a feature of reality. At the fixed-basis level the measurement problem receives an observer-relative account: no collapse, no branching and no objective randomness in the substratum; which operational protocols the completion settles is open (Chapter 19 §19.2.5). There is a bijection, a partition, and an observer who cannot see the whole.

This has implications for what classes of physics-based accounts of consciousness, the foundations of probability, the relationship between determinism and free will, and various other foundational questions are compatible with the framework. Accounts that require fundamental indeterminism in physical reality — Penrose's orchestrated objective reduction is the most prominent example in consciousness research — are incompatible with the framework's deterministic substratum. Accounts that treat quantum probabilities as derivable from incomplete information rather than as fundamental are compatible with and supported by the framework.

The framework does not claim to settle these foundational questions; it makes structural commitments that constrain which accounts are compatible. Sections 18.9 and 18.10 develop two of these constraints in detail — what the framework says about cosmic spatial topology and what it says about consciousness — but the broader pattern is that the framework's commitment to deterministic substratum is a substantive structural feature that constrains the foundational landscape.

The framework's quantum randomness is therefore epistemic, exact in its predictions for any embedded observer's empirical access, and consequential for the classes of foundational accounts compatible with it. The empirical record is silent at the 10^{-32} level; the conceptual record is substantial.

18.5 Quantum gravity: the cumulative case at the framework level

The framework's content on the quantum gravity problem extends beyond the individual dissolutions developed in Chapter 7 to a structural pattern. Three problems conventionally treated as obstacles to a theory of quantum gravity — the cosmological constant problem, the black hole information paradox, and the non-renormalizability of perturbative quantum gravity — receive the *same*

structural reading in the framework, each following from the same underlying commitment to the geometry-first ordering and the substratum-emergent operator distinction. The three are framework-level interpretations and conditional results — the cosmological-constant dissolution is established at the descriptive level with its dynamical form conditional on the covariant coupling theorem, the information account, in which information is conserved by the bijectivity premise and the Page curve is conditional, and the non-renormalizability reading is conditional on the open covariant rung G3, continuum-limit consistency included — not a completed theory of quantum gravity, and the information paradox is not resolved. This section pulls the three readings together as a cumulative case for the framework's structural treatment of quantum gravity.

The conventional formulation. Standard treatments of quantum gravity attempt to quantize the gravitational field by analogy with the quantization of the Standard Model forces. The metric tensor $g_{\mu\nu}$ is treated as a quantum field, decomposed into a classical background plus fluctuations, and the fluctuations are quantized through canonical or path-integral methods. Three obstacles arise.

The cosmological constant problem: the QFT vacuum energy summed over all field modes up to the Planck cutoff is of order $M_{\mathrm{Pl}}^4 \sim 10^{113}\,\mathrm{J/m}^3$. The observed vacuum energy density is $\sim 10^{-9}\,\mathrm{J/m}^3$. The ratio 10^{122} is treated as a fine-tuning problem requiring an unknown cancellation mechanism.

The black hole information paradox: Hawking's 1975 semiclassical calculation produces thermal radiation that appears to carry no information about the black hole's initial state. If the radiation is exactly thermal, information is destroyed when the black hole completes evaporation — in tension with quantum mechanics' commitment to unitarity.

Non-renormalizability: perturbative quantum gravity treated as a quantum field theory in flat spacetime is non-renormalizable. Infinitely many independent counterterms are required to absorb divergences at successively higher orders, with no UV completion within the standard quantization framework.

The framework's reading is that all three problems are *artifacts of the quantum-first ordering*, each dissolving, conditional on the covariant rung G3, when the geometry-first ordering is adopted: spacetime geometry is part of the classical substratum, and quantum mechanics is the emergent description of an embedded observer's reduced view of that substratum.

The first dissolution: the cosmological constant. Chapter 7 §7.7 developed the cosmological constant dissolution in detail. The framework has two distinct descriptions: the classical substratum, which assigns a vacuum energy $\rho_{\mathrm{crit}} \sim H^2/G \sim 10^{-9}\,\mathrm{J/m}^3$ through the Friedmann equation, and the emergent quantum field theory, which assigns a vacuum energy $\rho_{\mathrm{QFT}} \sim M_{\mathrm{Pl}}^4 \sim 10^{113}\,\mathrm{J/m}^3$ through zero-point summation. The two descriptions operate at logically different levels: on the geometry-first proposal gravity couples to the classical substratum, a coupling step open at G3, while the emergent QFT exists only after

the trace-out. The 10^{122} ratio is, up to a convention-dependent factor, the de Sitter entropy $S_{\mathrm{dS}} = A/(4l_p^2)$ — a dimensional identity for any Planck-cutoff theory, which the framework reads as the information compression ratio between substratum and emergent description. On that reading the ratio is not a fine-tuning failure.

The structural feature of this dissolution is that the two energy scales are not commensurable as gravitational sources. The conventional formulation treats them as comparable and asks why one cancels the other; the framework treats them as belonging to different levels of description, with no cancellation required if the coupling step holds.

The second component: the black hole information paradox. Chapter 7 §7.6 developed the framework's account of the information paradox. Information is never lost from the substratum bijection φ because φ is bijective; whether the radiation carries it is a separate question, which bijectivity alone does not settle. The apparent information loss in Hawking radiation is an artifact of the trace-out — the same mechanism that underlies the conditional cosmological constant dissolution. The framework's nested-partition formalism (two coexisting causal boundaries: cosmological horizon plus black hole horizon) provides a self-consistent emergent quantum description when the two boundaries coexist; the Page curve at the emergent level is a conditional result, on a scrambling condition that no bijection meets from an initial configuration with the bipartition aligned to the configuration coordinates (Chapter 7 §7.6).

The derivation runs through a mechanism substantially different from the quantum-extremal-surface and entanglement-island computations — cycle typicality on the configuration space rather than replica wormhole computation in the gravitational path integral — and the correspondence between the two is conjectural. For a mass-independent emission coefficient (Chapter 7 §7.6), its Page time is a fixed fraction of the evaporation time set by the entropy the radiation carries per unit of horizon entropy lost: $t_P \approx 0.646\, t_{\mathrm{evap}}$ when the two are equal, and earlier when the radiation carries more, as it does in the semiclassical emission regime defined in Chapter 7 §7.6.

The shared structural feature with the cosmological constant dissolution is that the framework's two-level description is doing the work. The information that appears lost in Hawking radiation is information that has crossed the trace-out boundary — accessible at the substratum level (bijectivity) but inaccessible at the emergent level while it resides in the sectors the emergent description traces out (the radiation looks thermal because the emergent description marginalizes over the relevant hidden-sector degrees of freedom).

The third dissolution: non-renormalizability. Perturbative quantum gravity treats the lattice spacing $\epsilon = 2\sqrt{\sigma}\, l_p$ as a regulator to be removed in the continuum limit. Renormalization-group analysis then shows that the theory requires infinitely many independent counterterms, with no UV completion possible within the standard quantization framework.

The framework's content is that the lattice *is* the fundamental description, not a regulator. The substratum's discreteness scale $\epsilon = 2\sqrt{\sigma}\, l_p$ is a structural feature of the substratum dynamics, fixed by self-consistency in Chapter 7 §7.4 rather than chosen as a calculational convenience. On this reading no continuum limit of the lattice is taken: the continuum is not the underlying description but the long-wavelength limit of the discrete substratum dynamics, valid wherever the lattice spacing is much smaller than the physical scale of interest. That this limit yields the covariant field equations, with their coupling to sources, is the open G3 rung ([GR §8.7]).

On the framework's reading, the non-renormalizability of perturbative quantum gravity is therefore an artifact of treating the lattice as a regulator to be removed. The substratum's lattice is physical; the renormalization-group flow runs *to* the Planck scale rather than *through* it, with the framework's lattice physics taking over at scales approaching ϵ. The structural commitment to the lattice's physicality removes the non-renormalizability problem at the level of interpretation, conditionally on G3, as the two-level description does for the cosmological constant problem.

The shared structural feature. Three problems that conventional treatments find independently obstructive receive the same framework-level reading. The shared feature is the geometry-first ordering: spacetime is part of the substratum, quantum mechanics is emergent, and the substratum-emergent operator distinction (developed in Chapter 1) is doing the work in each case. On the framework's reading, conditional on the coupling step, the cosmological constant ratio is a substratum quantity (the de Sitter entropy, up to a convention-dependent factor) being misidentified as an emergent quantity (the vacuum energy). The information paradox treats an emergent process (Hawking radiation) as if it should preserve substratum information (the substratum preserves it, by bijectivity; whether the radiation carries it is the separate question of Chapter 7 §7.6). The non-renormalizability treats a substratum-level discreteness (the lattice) as an emergent-level regulator (a continuum-limit artifact).

The cumulative case is not three independent dissolutions but one framework-level reading applied three times. Its claim is that, if the covariant rung G3 closes, there is no separate quantum gravity problem to solve — what exists instead is a structural framework (S, φ) from which both quantum mechanics and gravity emerge as different aspects of the same object, with the apparent conflicts of conventional quantum gravity being category errors. G3 is open, so the problem is partially addressed rather than dissolved (Chapter 19 §19.2.1). The framework's treatment carries conditional and consistency relations — the horizon calibration with its 1/4 ratio, the descriptive vacuum-energy distinction, the dark-sector reading, and the MOND scale $a_0 = cH/6$ conditional on the open coupling map with its factor 1/6 imported — rather than predictions independent of QM and GR; its forward exposure is listed in Appendix A.5.

18.6 Gravitational wave echoes

Several quantum-gravity proposals — firewall scenarios, fuzzball proposals, certain wormhole models — predict that a black hole's horizon partially reflects outgoing gravitational radiation, producing delayed echoes after the ringdown, with amplitudes that depend on the microstructure assumed. Echo searches in LIGO/Virgo data (Abedi, Dykaar and Afshordi 2017; Westerweck et al. 2018; Tsang et al. 2018; and the LIGO/Virgo/KAGRA Collaboration on the merger catalog through O3) have found no statistically significant signal.

The framework's statement. [GR §8.4] argues that a black-hole horizon does not change an external observer's partition, which is set by the cosmological horizon, so no OI-specific echo, comb or wall-reflection feature is expected and ringdowns follow general relativity; no echo amplitude is derived. Matter falling into a distant black hole was inside the external observer's visible sector before the merger and remains inside it after; the trace-out that defines that observer's emergent quantum description runs over the cosmological horizon, not over the black-hole horizon. On this argument the discreteness scale $\epsilon = 2\sqrt{\sigma}\, l_p$ supplies no echo signature for external observers.

The conceptual content. The more informative gravitational comparison is the Milky Way outer rotation curve (Chapter 7 §7.9), whose reported Keplerian decline beyond the crossover is in tension with the dark-sector law (the GW250114 result constrains the classical area theorem, not the 1/4 coefficient). For black-hole ringdowns the framework's commitment is the general-relativistic one.

The chapter's broader content on forward predictions continues with the framework's substratum commitments — finite information, recurrence, the arrow of time, cosmic topology, and consciousness — that test the framework at scales and levels distinct from the direct empirical exposure of the gravitational and quantum sectors.

18.7 Finite information and exact finite recurrence

The framework's commitment to a finite substratum has structural consequences that extend beyond the empirical predictions of fundamental physics. The universe is described by a finite configuration space S with bijection dynamics $\varphi : S \to S$. Finiteness is not a calculational convenience but a load-bearing commitment: it guarantees exact recurrence of the substratum dynamics — the finite-bijection form of recurrence, stronger than what Poincaré's theorem gives for a finite-measure system — which, together with a non-permutation rooted map, produces P-indivisibility at the embedded-observer level; a fixed-basis unitary quantum representation is internal and universal ([Main §3.4]), the compressed form supplied across the correspondence's class under (T), with P-indivisibility fixing its nontriviality. The recurrence route in the framework's logic chain requires finite S.

Finite information content. The substratum's information content is $\log_2 |S|$ bits — finite for any finite S. This is distinct from the Bekenstein-Hawking entropy $S_{\mathrm{BH}} = A/(4l_p^2)$, which bounds the *accessible* information at any given partition (the capacity of the visible sector and the boundary layer). The total information content of S is at least that of the boundary layer at the cosmological horizon; whether it exceeds the Bekenstein-Hawking bound is not fixed by visible data, and under the finiteness posit A1 (Chapter 2 §2.4) it is finite.

A clarification is essential. The cardinality $|\mathcal{C}_D|$ of the deep hidden sector is *gauge* in the framework: different values of $|\mathcal{C}_D|$ produce the same visible transition data at leading order on accessible timescales. The "total information" of the universe is therefore not fixed by those data. But the *finiteness* of $|\mathcal{C}_D|$ — for the recurrence route specifically, that there exists some finite value, even if its specific magnitude is not observable — is load-bearing. Infinite hidden sectors would break the recurrence argument that drives recurrence-scale P-indivisibility; that route would close. Accessible non-Markovianity would survive through readback, and the imported representation is gated by (T) rather than by recurrence, so an infinite substratum removes this route rather than the framework's quantum representation.

The framework's commitment to finite information is therefore neither a quantitative claim about the specific cardinality nor a metaphysical preference for finitism. It is a structural requirement for one particular route — the recurrence-based P-indivisibility result, which needs every root's visible law to return to certainty at a common time, $\Gamma_n = I$, almost surely over the hidden prior. A bijection of a finite set supplies that return at its order ($\varphi^N = \mathrm{id}$ gives $\Gamma_N = I$); the Poincaré recurrence theorem for a finite-measure system does not, since it returns almost every state at its own time. The substratum's finiteness is what makes this particular route to P-indivisibility available.

Exact finite recurrence. A bijection φ acting on a finite set S is a permutation. Every permutation on a finite set has finite order: there exists a smallest positive integer N such that $\varphi^N = \mathrm{id}$, the identity map. After N time steps, the substratum returns to its exact initial state. The framework's substratum dynamics is therefore strictly periodic, with period N determined by the bijection's cycle structure.

The recurrence period can be astronomically large. For an arbitrary bijection of S, two upper bounds with different growth rates bear on it. *A bound valid for every permutation:* N is the least common multiple of the cycle lengths, each at most $|S|$, so N divides $\mathrm{lcm}(1, 2, \ldots, |S|)$. Its logarithm is Chebyshev's function, $\ln \mathrm{lcm}(1, \ldots, n) = \psi(n)$, with $\psi(n) \sim n$ by the prime number theorem. *The largest period any permutation attains:* this is Landau's function $g(n)$, the maximum order of a permutation of n points, with $\ln g(n) \sim \sqrt{n \ln n}$ (Landau 1903). Since $\sqrt{n \ln n}$ is $o(n)$, $g(n)$ lies far below $\mathrm{lcm}(1, \ldots, n)$, and no permutation has order near e^n. Typical-permutation order estimates require a specified probability law for φ, which the framework does not adopt; Chapter 3 §3.2 holds that random permutations of finite sets typically lack the structure C1–C4 require.

The substratum's cardinality is not fixed by visible data. [Substratum §4] bounds only the visible-plus-boundary sector, $|\mathcal{C}_V \times \mathcal{C}_B| \leq e^{10^{122}}$ at the de Sitter horizon; the deep sector's cardinality is gauge (above). For illustration only, if $|S|$ were of order $e^{10^{122}}$, the first bound would give $N \leq \operatorname{lcm}(1, 2, \ldots, |S|) = e^{\psi(|S|)} \sim e^{e^{10^{122}}}$, in the sense $\ln N \leq \psi(|S|) = (1 + o(1))\,|S|$. The second is sharper: no bijection of that set has period above $g(|S|)$, with $\ln g(|S|) \sim \sqrt{|S| \ln |S|} \sim 10^{61}\, e^{5 \times 10^{121}}$.

The exactly linear rule on a fixed coupling graph is far more constrained than an arbitrary bijection. Under linearity (A5, Chapter 2 §2.4) the update is an invertible linear map of $S = (\mathbb{Z}/q\mathbb{Z})^D$, where D counts every stored component, including the previous-time copy that a second-order rule carries. Its powers $\varphi^0, \ldots, \varphi^{N-1}$ are distinct and nonzero, and they lie in the ring generated by φ, which by the Cayley–Hamilton theorem has at most q^D elements. Hence $N \leq q^D - 1 = |S| - 1$ for every alphabet size $q \geq 2$. For prime q some linear maps attain $q^D - 1$, but the wave rule falls far short of it: on a ring of L sites, with q prime and $\gcd(L, q) = 1$, its period is exactly qL, against $|S| = q^{2L}$ (Appendix B, Theorem B.3.1). The bound does not cover the other maps the corpus uses: the state-dependent-geometry map $F(u, v) = (v, \mathrm{WE}_{G(v)}(v) - u)$ of [SM §3.1], which carries the gravitational sector, and the rounded bijection of [SM §4.2], whose proposed stationary law is meant to support full-shell sampling, are bijections but not linear maps, and for them only $N \leq g(|S|)$ and $N \mid \operatorname{lcm}(1, \ldots, |S|)$ hold. The period of the three-dimensional rule at cosmological size is not computed in the corpus.

The return itself is definite: the universe returns to its exact initial state after N substratum time steps, where N is fixed by the cycle structure of φ. It is at most $g(|S|)$ for any bijection and below $|S|$ for the exactly linear rule. The framework bounds N rather than computing it, and it supplies no lower bound on N. Translating to physical units with the substratum time step $\tau_{\mathrm{sub}} \sim t_{\mathrm{Pl}} \sim 10^{-43}$ seconds multiplies the recurrence time by τ_{sub}, which changes its logarithm by about 10^2, negligible at these magnitudes. A period of astronomical size is vastly longer than the age of the universe, vastly longer than the lifetime of the proton, vastly longer than any timescale physics conventionally treats, and at such a period the recurrence has no practical empirical consequences.

Falsifiability. The recurrence prediction is falsifiable only in the logical sense. For a substratum of given $|S|$ and φ, failure to recur within $g(|S|)$ steps would contradict a finite S with bijection dynamics, and failure to recur within $|S| - 1$ steps would contradict only exact linearity (A5). But $|S|$ is not fixed by visible data: for any finite observation time T, a substratum with $|S| > T + 1$ is admissible, so no finite observation falsifies the framework as a whole, which would require an infinite observation time. An embedded observer's records are, moreover, part of S and return with it. The recurrence is therefore a prediction at the structural level rather than at the empirical level: it follows necessarily from the framework's other commitments (finite substratum, bijection dynamics), and any framework consistent with those commitments must predict exact

recurrence with a period at most $g(|S|)$ that divides $\mathrm{lcm}(1, \ldots, |S|)$, and below $|S|$ for the exactly linear rule.

The conceptual content is significant. The framework predicts that the universe is *cyclic at the substratum level* — periodic with period N, the least common multiple of the bijection's cycle lengths. For a rule fixing the zero configuration on a nontrivial state space, no single cycle passes through every configuration. The exact linear rule of [SM §4.2] has this property and therefore at least two cycles. The cycle-ergodicity question that [SM §4.2] treats concerns sampling within an energy shell. Within a single cycle no state function increases throughout; a direction of the observer's experience is fixed by boundary conditions on a stretch of the cycle (the arrow of time discussed in the next section), and at the substratum level the dynamics has no preferred direction. The framework's commitment to substratum cyclicity is one of the structural commitments that distinguishes it from frameworks treating the universe as either fundamentally non-cyclic (most cosmological models) or fundamentally infinite (most quantum gravity programs).

Implications for the framework's logical structure. The recurrence theorem connects three structural features of the framework: substratum finiteness (the framework's commitment to $|S| < \infty$), bijection dynamics (the framework's commitment to φ as a bijection), and P-indivisibility at the recurrence scale. The connection runs in three steps. Finiteness of S guarantees exact recurrence ($\varphi^N = \mathrm{id}$). Recurrence together with C1 acting at the first step — which makes the one-step matrix a non-permutation under the uniform hidden prior, the cheapest non-permutation rooted map — gives returns of information from hidden to visible over the recurrence timescale (P-indivisibility across the recurrence cycle, not on accessible windows); the coupling is load-bearing rather than decorative, since for an uncoupled partition $\varphi = \varphi_V \times \varphi_H$ the visible sector evolves autonomously, its two-time family is divisible, and no distinguishability is ever restored — recurrence notwithstanding. With a full-support visible prior this indivisibility is equivalent to memory at some order below the recurrence time; the memory-bearing sector is the one in which C4 places that memory at an accessible order, and every finite law, memory-bearing or not, admits the quantum representation.

The framework's commitment to finite information is therefore load-bearing for the recurrence route specifically, not for representability or for accessible non-Markovianity: readback at order k has its own finite-horizon route, and the imported representation is gated by (T) rather than by recurrence, so those routes do not exclude an infinite deep hidden sector, which would need its own normalized state or measure and well-defined dynamics ([Main §2.3], §3.1; Chapter 2 §2.3). Removing finiteness — replacing $|S| < \infty$ with $|S| = \infty$ — would remove the recurrence route rather than the framework's quantum representation.

18.8 The arrow of time

The framework's substratum dynamics is given by the bijection $\varphi : S \to S$. The bijection has *no arrow of time*. Both φ and its inverse φ^{-1} are equally valid maps on the configuration space, and the substratum dynamics is unchanged under the exchange $\varphi \leftrightarrow \varphi^{-1}$. The framework's commitment to T-symmetric substratum dynamics is the same commitment that gives reciprocal transition counts in Chapter 6 §6.3 (and narrows θ to $\{0, \pi\}$ without fixing $\bar{\theta}$) — T-invariance at the substratum level is a structural feature that propagates through the framework's emergent descriptions.

This raises an obvious question. The universe we observe has an unambiguous arrow of time: entropy increases, biological systems age, memories form of the past but not the future, the universe expands rather than contracts. If the substratum dynamics is T-symmetric, where does the observed arrow come from?

The arrow as observer-emergent. The framework's answer is that the arrow of time is *emergent from the observer's coarse-grained description*, not from the substratum dynamics. The substratum dynamics is T-symmetric; the observer's reduced description of a non-equilibrium initial condition is not. The asymmetry enters through the trace-out of a non-equilibrium initial condition — the marginalization over the hidden sector that produces the embedded observer's reduced description; under the invariant measure the trace-out alone produces none. The arrow of time is therefore a feature of embedded observation of such a condition, in the same way that quantum mechanics, the dark sector, and the Born rule are.

The mechanism is the standard Boltzmann argument applied to the framework's partition. The visible sector starts in a particular configuration that is, in some sense, "low entropy" relative to the typical configurations accessible at later times. The hidden sector starts in a configuration that contains all the correlations needed to specify the visible sector's future evolution. As the dynamics evolves, correlations flow from hidden sector to visible sector and back, with the time-reversibility of the underlying dynamics fully preserved at the substratum level. But the observer, who sees only the visible sector, sees an irreversible flow: information about the past is encoded in the visible sector's current state (entropy increasing), while information about the future is encoded in the hidden sector (inaccessible to the observer).

The arrow of time is therefore *the observer's incompleteness* expressed temporally. The same partition that produces quantum randomness (by hiding the substratum information that determines outcomes) produces the arrow of time (by hiding the substratum information that would let the observer reverse the dynamics). Both are aspects of the same structural feature: the observer's reduced description is asymmetric in ways that the substratum dynamics is not.

Detailed balance and the thermodynamic arrow. The framework's con-

tent on detailed balance from Chapter 6 §6.3 has a direct implication for the arrow of time. For a time reversal that preserves the partition, and the counting prior on hidden states, the visible transition probabilities satisfy reciprocity, $T_{ij} = T_{\theta j,\theta i}$ at all timescales, with θ the action of time reversal on visible states; when the visible states carry no T-odd data this is detailed balance, $T_{ij} = T_{ji}$ — the rate of transition from i to j equals the rate of transition from j to i. This follows from *substratum-level* T-symmetry: the dynamics has no preferred direction.

The thermodynamic arrow of time emerges not from any asymmetry in the transition probabilities but from the partition's *initial conditions*. The universe starts in a configuration that is far from equilibrium relative to the substratum's full state space, with most of the substratum's degrees of freedom on the hidden side of the partition and the visible sector occupying a small, structured region. As the dynamics evolves, the system moves toward equilibrium relative to the substratum's state space — entropy increasing in the observer's coarse-grained description — until the entire substratum is fully mixed. The arrow of time runs in the direction of this equilibration.

The relation to detailed balance is that the H-theorem — the standard derivation of entropy increase from microscopic dynamics — is a theorem about the *coarse-grained* description, not about the underlying transition probabilities. The H-theorem assumes a particular kind of initial condition (low entropy at the coarse-grained level) and derives entropy increase from there. Detailed balance at the microscopic level is fully consistent with the H-theorem at the coarse-grained level, because the H-theorem is about how the observer's information about the system evolves rather than about the system's underlying dynamics.

The cosmological arrow. Whether the thermodynamic arrow and the cosmological arrow — the direction in which the universe expands — are aligned is open in the framework: it is neither established nor refuted. The cosmological partition has a specific initial condition: the universe begins in a low-entropy configuration with most degrees of freedom on the hidden side of the cosmological horizon. On an expanding FRW background satisfying the null energy condition, the apparent-horizon area is non-decreasing. Its assigned entropy has that interpretation under the stated calibration assumptions; whether this behavior aligns with boundary correlations or hidden-sector disorder remains open.

Both arrows are traced to the same initial conditions of the cosmological partition. That is a common boundary condition, not a derived mechanism: that the thermodynamic arrow within the visible sector tracks the cosmological arrow is not established. The universe started with low entropy at the cosmological level, as an initial condition; at the cosmological boundary, coupling C1, persistence C2 and capacity C3 are verified, while history readback C4 is a named realization condition not presently discharged (Chapter 7 §7.2); the cosmological boundary is the primary partition for any embedded observer.

Why the framework reframes rather than explains. The framework's content on the arrow of time is not an explanation of *why* the universe has a preferred direction; it is a *reframing* of the question. Standard accounts of the arrow of time treat it as an asymmetry that requires explanation — why is entropy increasing in the direction we call "future" rather than the direction we call "past"? The framework's answer is that there is no asymmetry in the substratum dynamics; the asymmetry is in the observer's reduced description, which inherits the initial conditions of the partition.

The reframing places the explanatory burden on a different question. Instead of asking "why does the universe have a preferred direction of time?", the framework asks "why does the universe start in a low-entropy partition?" The latter question is a question about *initial conditions* — addressable by inflationary cosmology, anthropic arguments, or other initial-condition mechanisms — rather than a question about *dynamics*. This is the same kind of reframing the framework applies to the cosmological constant problem (Chapter 7 §7.7): the conventional formulation treats the small observed value as a fine-tuning problem in the dynamics; the framework treats it as a feature of the partition's structure that does not require dynamical explanation.

The arrow of time is therefore one more case where the framework's substratum-emergent distinction does substantive work. The substratum is T-symmetric (the dynamics has no arrow). The emergent description is not T-symmetric (the observer sees an arrow). The asymmetry is structural — inherited from the partition's initial conditions — rather than dynamical. The framework's content here is not to solve the arrow-of-time problem in the conventional sense but to locate it precisely: the arrow is at the level of partition initial conditions, not at the level of substratum dynamics.

18.9 Cosmic spatial topology

The framework's structural commitments determine the *local* structure of the substratum but are silent on its *global* structure. The substratum is a cubic lattice at discreteness scale $\epsilon = 2\sqrt{\sigma}\, l_p$ (Chapter 7 §7.4). The local cubic structure produces the gauge group, the matter content, and the framework's quantitative predictions across Standard Model and gravitational sectors. But whether the lattice extends globally as $\mathbb{R}^3$, identifies as a 3-torus, has a Klein-bottle quotient, or admits some other global identification structure — that is not determined by the framework's structural commitments. The framework is locally specific and globally agnostic.

This silence is principled. The framework's content develops from the cosmological partition and the partition's local properties at scale ϵ. Global topology operates at scales much larger than the cosmological horizon if at all (current empirical constraints place any non-trivial topology beyond the observable universe, as developed below). The framework's predictions are therefore independent of

global topology choices, and the framework's commitments do not constrain them.

Janna Levin's compact-universe proposal. Levin developed in the late 1990s and early 2000s the proposal that the universe could have a non-trivial spatial topology — specifically, that spatial sections could be *compact* (a 3-torus, a Klein-bottle quotient, or other compact identifications) rather than infinite. The motivation was partly cosmological (compact topology produces specific signatures in the CMB power spectrum at the largest angular scales) and partly philosophical (an infinite universe creates conceptual difficulties around uniqueness and the anthropic principle, addressed at book length in Levin's *How the Universe Got Its Spots* in 2002).

The Klein-bottle specifically refers to non-orientable compact spatial topologies in which a parity-reversing identification connects opposite "sides" of the universe. The proposal is one example within the broader family of compact-universe possibilities — others include the 3-torus (orientable, no parity reversal), the Poincaré dodecahedral space (constant positive curvature, no parity reversal), and various lens-space identifications. Levin's work, with collaborators Cornish, Spergel, Starkman, Weeks, and others, developed the empirical program for testing compact-universe proposals through their characteristic CMB signatures.

The empirical program: matched-circle signatures. The cleanest empirical test of compact-universe proposals is the *matched-circle* signature in the CMB. If the universe has compact topology with identification scale smaller than the diameter of the observable universe, then light rays that traverse the topological identification can return to the observer's past light cone, producing correlated temperature patterns on circles in the CMB sky. The two circles on the CMB sky corresponding to the two endpoints of the same topological geodesic should show matched temperature patterns: the same fluctuation field probed at two different points on the celestial sphere.

The matched-circle test is geometrically clean — it depends only on the topological identification structure, not on details of the early-universe physics — and it has well-defined null and signal cases. Cornish, Spergel, Starkman, and Komatsu (2004) developed the statistical methodology for detecting matched circles in the CMB. WMAP data through 2010 and Planck data through 2015 have been searched for matched-circle signatures.

The empirical result is null: no statistically significant matched-circle signatures have been detected. The current bound places any compact-topology identification scale at greater than approximately the diameter of the observable universe — equivalently, any compact identification has scale $\gtrsim 28$ Gpc, beyond the cosmological horizon. The empirical situation is therefore that either the universe is topologically trivial at observable scales, or the universe has compact topology with identification scale larger than current cosmic horizon. The two possibilities are not currently distinguishable by data.

The framework's compatibility with Levin's proposal. Levin's compact-universe proposal is *consistent with* but *not predicted by* the framework. Three structural points clarify this.

First, the framework's substratum finiteness (developed in §18.7) is *compatible with* compact spatial topology. A finite substratum could be realized as a cubic lattice with compact global identification at some scale beyond the cosmological horizon, where the total number of lattice cells is finite. Levin's proposal provides one specific realization of this — the Klein-bottle quotient or 3-torus identification at some cosmological scale.

Second, the framework's finiteness *does not require* compact spatial topology. The substratum's finiteness could also be realized as a cubic lattice with trivial $\mathbb{R}^3$ topology but with a finite extent set by other structural features. Levin's compactness is sufficient but not necessary for the framework's finiteness.

Third, the framework's predictions in Chapters 5 through 17 are *independent* of global topology choices. Nothing in the framework's empirical content — Standard Model gauge group, gravitational sector predictions, JUNO neutrino result, dark sector phenomenology, biological consequences — depends on whether the universe is topologically trivial $\mathbb{R}^3$, compactified as a 3-torus, or has the Klein-bottle quotient structure Levin proposes. The framework's content is robust to global topology choices because the framework operates at the partition boundary (the cosmological horizon), not at the universe's global topology.

The framework is therefore *neutral* on compact-universe proposals. Levin's specific Klein-bottle proposal, the broader 3-torus family, and the trivial $\mathbb{R}^3$ alternative are all consistent with the framework's substratum structure. The framework neither favors nor disfavors any particular global topology.

The CMB as joint testing ground. Both Levin's matched-circle program and the framework's Born-rule transient violations (§18.2) are CMB-data-driven empirical programs. They are structurally distinct — matched-circles tests global identification, Born-rule transient violations tests boundary thermodynamics at inflation — but they use overlapping observational data. Planck 2018 constrains the standard non-Gaussianity shapes, and CMB-S4 and LiteBIRD will tighten those constraints; whether they bear on the Born-rule proposal depends on the quantities §18.2 does not yet fix.

The framework's CMB content does not compete with Levin's. The matched-circle test depends on the spatial extent and structure of topological identifications. The framework's Born-rule transient violations, if present, depend on the temporal evolution of the visible-hidden coupling during inflation. The two tests probe orthogonal physical content using the same data sets — analogous to how the JUNO experiment probes neutrino mixing and the same reactor antineutrino flux probes other oscillation parameters.

Open question: non-orientability and chirality. The framework's local-lattice chirality from Chapter 5 — SU(2) as chiral and SU(3) as vector-like,

derived from the staggered trace-out — is a structural feature at the substratum's local cubic level. A separate question is whether a non-orientable global topology (Klein-bottle specifically, or other non-orientable compact identifications) interacts non-trivially with the framework's local-lattice chirality.

The intuition is that non-orientability at the global level might produce phase factors or sign factors that interact with the local chirality structure, possibly producing observable consequences. The intuition could be wrong — the local-lattice chirality emerges at scales of $\epsilon = 2\sqrt{\sigma}\, l_p$, while the global topology operates at scales beyond the cosmological horizon, with the relevant length scales differing by approximately 60 orders of magnitude at $\sigma = 1$. The two might be entirely decoupled.

This is an open question the framework has not addressed. Resolving it would require a calculation: starting from a specific non-orientable compact topology (Klein-bottle or otherwise), determining whether the framework's local chirality derivation acquires global corrections, and computing the observable signatures of any such corrections. The framework does not currently have the calculation; the question is pre-registered as an open theoretical task.

The framework's content on cosmic spatial topology is therefore: silent at the framework level (no commitment beyond local cubic structure), consistent with Levin's compact-universe proposals (as one realization of the framework's finiteness), with an open question on non-orientability-chirality interaction that requires further calculation. The framework's structural unity does not extend to global topology, and the framework is honest about this scope.

18.10 The framework's position on consciousness

The framework's mathematical object — the partition with C1–C4 structure — defines an *observer* in a structural sense. The observer is not yet a *conscious entity*. Whether the structural conditions for observation are also sufficient conditions for consciousness is a separate question, with the framework's commitments having implications for what answers are possible.

This section develops the framework's substantive position on consciousness: observation is *necessary but not sufficient* for consciousness. The framework provides structural necessary conditions for conscious systems and imposes constraints on what physics-based consciousness theories are compatible with its commitments. The hard problem of consciousness proper — the explanatory gap between physical/structural facts and phenomenal experience — is not addressed, and the framework is explicit about this scope limitation.

Mathematical observer is not equivalent to conscious entity. The framework's observer is a partition V of the substratum's configuration space combined with the conditions C1 (coupling to hidden sector), C2 (record persistence — slow-bath separation the route; a physical-regime premise in [Main]'s typing), C3 (sufficient hidden-sector capacity), and C4 (history readback — the

load-bearing memory condition). The partition is a mathematical object that has dynamical consequences (the trace-out producing the reduced law whose representation carries the emergent QFT), thermodynamic consequences (entropy displacement producing dark matter phenomenology), and empirical consequences (the JUNO prediction, the BH entropy, the dark sector).

The partition is not, by itself, a conscious entity. A thermostat coupled to room temperature implements a partition in a degenerate sense (the thermostat's internal state interacts with the room's thermodynamic state). A cosmological horizon implements a partition with $\tau_S/\tau_B \sim 10^{-32}$. A protein interior implements a partition. Nothing in the framework attributes phenomenal experience to any of them. The framework's mathematical observer is therefore a structural object, not a phenomenal one, and the framework does not claim equivalence between mathematical observation and conscious experience.

The asymmetric position. The framework's substantive content on consciousness is the asymmetric claim: any conscious system must implement a partition with C1–C4 structure, but not every C1–C4 partition is conscious. The framework provides structural necessary conditions for consciousness without specifying sufficient conditions. The two directions below differ in standing: Direction 2 follows from the definition of observation, which detectors and horizons satisfy; Direction 1 is a conjecture about consciousness, motivated by the role of memory in conscious information processing, and not a theorem of the framework. Evidence in the framework's sense is perspective-neutral — a record bears its evidential relations through its correlations at the partition, whether or not a conscious system takes it up; whether evidence as experienced requires a conscious perspective is an epistemological question the framework does not address.

The claim has two directions worth distinguishing.

Direction 1: Consciousness requires observation. Any conscious system must implement a partition with substantial C1–C4 structure. A system that fails C1 (no coupling to a hidden sector) cannot be conscious because it has no environment to be aware of. A system that fails C2 (hidden sector equilibrates faster than visible-sector dynamics) cannot be conscious on the framework's account with C2 read in its timescale form: the memory it retains, which would support integration over time, is suppressed. A system that fails C3 (insufficient hidden-sector capacity) cannot be conscious because it cannot store the information that conscious experience would require. The C1–C4 architecture is necessary for the non-Markovian information processing that conscious systems must exhibit.

Direction 2: Observation does not require consciousness. The cosmological horizon, proteins, and engineered C1–C4 systems implement observation structurally, and nothing in the definition requires that they be conscious. The framework's observation has structural consequences (the memory-bearing quantum representation, dark sector phenomenology) at the level of the partition's

emergent dynamics, but it does not entail phenomenal experience. Whatever distinguishes the conscious C1–C4 systems from the non-conscious ones is some *additional feature* beyond the bare C1–C4 architecture — biological substrate, neural architecture, integrated information Φ, global workspace dynamics, recurrent processing, or some combination — that the framework's structural commitments do not specify.

The asymmetric position is more conservative than alternatives like dual-aspect monism (where structural observation and phenomenal consciousness are necessarily co-occurring) and panpsychism (where every C1–C4 system is conscious to some degree). The framework's content allows but does not require these stronger positions.

Substantive constraints on consciousness theories. The framework's commitments constrain what physics-based theories of consciousness are compatible with it.

Deterministic substratum rules out fundamental-indeterminism accounts. The framework's substratum is deterministic: the bijection φ maps each state to a unique successor. Quantum randomness is epistemic, not ontic (§18.4). Theories of consciousness that *require* fundamental indeterminism in physical reality are incompatible with the framework. The most prominent example is Penrose's orchestrated objective reduction (Penrose-Hameroff), which posits that conscious experience arises through gravitationally-induced quantum collapse in microtubules — a specific form of fundamental indeterminism that the framework's deterministic substratum forbids. The framework's commitments make Penrose-Hameroff structurally incompatible.

C1–C4 architecture is necessary for non-Markovian information processing. Conscious systems must exhibit memory effects, history-dependent dynamics, and readback of stored correlations between subsystems — the structural signatures of non-Markovian dynamics that emerge from the framework's C1–C4 architecture. Theories of consciousness that treat conscious systems as fundamentally Markovian (instantaneous, memoryless processing) are incompatible with the framework's account of how cognitively non-trivial information processing must work. Strict feed-forward computational accounts of consciousness face this constraint.

The substratum is finite. The framework's finite substratum (§18.7) constrains theories of consciousness that posit infinite information content for conscious experience. Theories requiring genuinely infinite degrees of freedom (some forms of mathematical Platonism applied to consciousness, for instance) face this constraint.

These are real constraints rather than vacuous ones. The framework rules out a substantial class of physics-based consciousness theories while remaining compatible with the broader landscape of accounts that supply additional sufficient conditions for consciousness beyond the framework's necessary ones.

Integrated information theory as adjacent content. Integrated information theory (IIT), developed primarily by Giulio Tononi, is the most prominent physics-adjacent consciousness theory and has substantive content overlap with the framework. IIT proposes that consciousness corresponds to integrated information Φ, a quantitative measure of how much a system's parts inform each other beyond what their individual contributions predict. Systems with high Φ are conscious; systems with low Φ are not.

There is a real structural relationship between the framework's C1–C4 architecture and IIT's Φ. C1 (non-trivial coupling) is necessary for non-zero Φ — without inter-part coupling, integrated information cannot be non-trivial. C2 (record persistence — realized here by slow-bath separation) is necessary for IIT's information integration over time — the framework's persistence requirement, met via $\tau_B \gg \tau_S$, provides the memory architecture that integration requires. C3 (sufficient hidden-sector capacity) is necessary for the system's state space to be large enough to support meaningful integration. A system satisfying C1–C4 with substantial margins (large τ_B/τ_S, large N_H/N_V) is a system with substantial Φ; a system satisfying C1–C4 marginally has low Φ.

The relationship is not exact identity. C1–C4 are *necessary conditions* for non-Markovian dynamics in the framework's specific sense, while Φ is a *quantitative information-theoretic measure* in IIT's specific formulation. But the conditions overlap substantially. The framework's content here is compatible with IIT as a candidate sufficient-condition theory: the C1–C4 architecture provides the necessary structural conditions for the framework, and IIT's Φ provides the additional sufficient content for consciousness within that architecture. The framework does not endorse IIT's identification claim (that Φ *is* consciousness), but it provides structural foundations that IIT-like accounts can build on.

The relationship runs in both directions. The framework's structural account of subjecthood — what makes a system a subject of observation in the framework's mathematical sense — is compatible with IIT's account of what makes a system conscious. Conversely, IIT's predictions about which systems are conscious provide quantitative content that complements the framework's structural necessary conditions. The two accounts engage at different levels and are not in tension.

The hard problem is not addressed. The framework's content on consciousness is necessary-condition content. The framework provides structural conditions that conscious systems must satisfy; it provides constraints on what theories of consciousness are compatible with the framework's commitments; it provides empirical predictions about the structural features conscious systems must exhibit. None of this addresses the hard problem of consciousness proper — the question of *why* phenomenal experience accompanies physical processes at all.

The hard problem is a question about the relationship between structural/physical facts and phenomenal facts. The framework provides a

structural account (the partition with C1–C4 structure produces the emergent dynamics that conscious systems must satisfy). What it does *not* do is bridge the explanatory gap from such a structural account to phenomenal experience. The framework predicts which systems should be conscious as a necessary-condition claim, but it does not explain *why* implementing those necessary conditions should produce subjective experience rather than functional information processing "in the dark."

This scope limitation is honest rather than evasive. No physics-based framework currently bridges the explanatory gap; claiming that the framework does so would be overreaching. The framework's value here is its discipline: necessary-condition content with substantive constraints on theories, combined with explicit acknowledgment that the hard problem is not addressed. This is more than the strict scope often adopted in physics frameworks (which decline to engage consciousness at all) and less than the overclaiming sometimes seen (where physics frameworks announce solutions to the hard problem they cannot actually deliver).

Empirical predictions and constraints. The framework's necessary-condition conjecture implies structural expectations about the features conscious systems should exhibit; they are consistency checks on the conjecture, not evidence for the framework.

Conscious systems should exhibit non-Markovian information processing — memory effects, history-dependent dynamics, readback of stored correlations between subsystems. The C1–C4 architecture is empirically detectable in neural systems; the prediction is that systems we identify as conscious (humans, mammals, possibly other complex biological systems) should exhibit these signatures at their relevant timescales. Neuroscience evidence — synaptic plasticity, neuromodulator gradients, glial dynamics, slow oscillations supporting cognitive integration — is consistent with this prediction. The framework's content is structurally aligned with the broader neuroscience consensus on the architecture of conscious information processing.

Conscious systems should *not* be possible in architectures that strictly fail C1–C4. The framework predicts that strictly feed-forward neural network architectures (without recurrent connections, without memory states, without integration over time) cannot produce consciousness — they fail C2 by lacking the temporal integration architecture. This is consistent with the empirical finding that consciousness in humans is associated with recurrent processing rather than purely feed-forward processing. The framework's content here is a structural constraint that aligns with empirical neuroscience.

The framework's bioinformatics content from Chapter 17 is structurally related. The molecular clock overdispersion, rate autocorrelation across the tree of life, and LTEE power-law fitness trajectory are all empirical signatures of non-Markovianity in evolutionary biology systems. The framework predicts that similar non-Markovian signatures should be detectable in conscious systems at

their relevant timescales — the brain's analog of the molecular clock's overdispersion would be the variability of neural processing rates across timescales, predicted by the framework to be non-Markovian rather than Markovian. The empirical program connecting these structural predictions to consciousness research is open.

The framework's content on consciousness is therefore disciplined necessary-condition content with substantive empirical predictions, real constraints on theory choices, and explicit honesty about the limits of its scope. The hard problem of consciousness proper is not solved or dissolved by the framework. The framework's value in consciousness research is structural — what makes a system the kind of system that *could* be conscious — rather than phenomenal.

18.11 Chapter close

This chapter has developed the framework's distinctive forward content beyond standard quantum mechanics and general relativity. The empirical record of Chapters 5 through 17 establishes the framework's content where it agrees with current physics at high precision; this chapter has developed the framework's content where it makes predictions current physics does not yet test.

Four categories of forward content. The chapter's content sorts into four categories distinguished by their empirical accessibility.

Testable forward predictions. The restricted quantum-statistical regime (§18.3) is testable through distinguishability-revival measurements in single-molecule kinetics, accessible to current laboratory techniques. The BQP computational ceiling (Chapter 14) is exposed to any demonstrated super-BQP computation — which would falsify the global-coverage premise first, the framework-plus-model conjunction only with further argument. The framework's content in this category is empirically engaged at near-term experimental scales.

Conditional prospects. The Born rule's dynamical equilibrium (§18.2) is a proposed mechanism without a specified observable response. It would become testable through CMB measurements of the primordial power spectrum and non-Gaussianity, such as CMB-S4 and LiteBIRD, only once the source dynamics, the family of initial deviations and the map to observables are specified; until then it carries no quantitative prediction.

Structural commitments with limited empirical access. The epistemic character of quantum randomness (§18.4) is consequential for foundations but has no observable consequences at the 10^{-32} suppression level. The cumulative case on quantum gravity (§18.5) reframes three problems but has no direct empirical signature beyond the predictions already developed in the cosmological constant, BH information paradox, and renormalization treatments. Finite information and exact finite recurrence (§18.7) are falsifiable only in the logical sense — for a substratum of given $|S|$ a test would require observation over up to $g(|S|)$ steps, and since $|S|$ is not fixed by visible data, no finite observation falsifies the

framework as a whole. The arrow of time (§18.8) is observationally manifest but is not directly testable as a framework prediction because all physics-based accounts of the arrow generate the same empirical signature. The framework's content in this category is structurally consequential without producing additional empirical handles beyond those already developed in Chapters 5-17.

Framework's scope limits. Cosmic spatial topology (§18.9) is not predicted by the framework; the framework is consistent with Levin's compact-universe proposal and the trivial $\mathbb{R}^3$ alternative equally. The hard problem of consciousness (§18.10) is explicitly outside the framework's scope; the framework provides necessary-condition content on consciousness but does not bridge the explanatory gap from structural to phenomenal facts. The framework's content in this category is honest acknowledgment of what the framework does not address.

Forward pointers. Chapter 19 develops the framework's content against the standard list of open problems in fundamental physics. The chapter is a systematic inventory: ten problems the framework addresses with a resolution, dissolution, conditional construction or partial treatment, its table giving each one's status (quantum gravity in three components, covariant unification open, the cosmological constant, dark matter, dark energy, the measurement problem, BH information, hierarchy, gauge group origin, generation puzzle, dark sector budget), two split-status entries (the Born-rule origin — representation level settled, coherent operational extension characterized by five named conditions; strong CP — open under two named conditions) and five scope limits where the framework does not provide a derived value (the flavor problem with structured open task; baryogenesis, which §19.3.3 reclassifies as solution-specific — the baryon asymmetry is downstream of the framework's flavor-sector CP input; inflation and initial conditions; the Hubble tension; the cosmological initial state). Each entry cross-references where in the book the relevant content is developed, with Chapter 19 functioning as a reference resource for readers wanting to understand how the framework positions itself relative to the recognized hard problems of fundamental physics.

Cumulative case. The framework's content across the empirical chapters (5-17) and this forward-predictions chapter (18) forms a cumulative case for the framework's structural commitments. Of the twenty-two Standard Model observables, two are parameter-free structural retrodictions and five are conditional structural ones, and the remainder are layered, chained or fitted; the seven gravitational-sector entries are conditional relations and estimates, two of them compared with measured values (G6, G8) and one with upper bounds (G12); the seven neutrino-sector entries include retrodictions and bound-only comparisons. The framework's content in biology (Chapters 11-13 and 16-17) makes specific predictions that align with the empirical record across non-Markovian dynamics, evolutionary signatures, and clinical pharmacology. The forward content of this chapter extends the framework's reach to phenomena standard physics has no specific commitments on — Born-rule equilibrium, the capacity-controlled origin of partial revival, computational ceilings, and the arrow of time. The

consciousness conjecture of §18.10 is not part of this case.

The framework's evidential standard is parameter-free structural prediction matching observation. The cumulative case across the book is retrodictive — exact structural values matching existing data across physics, biology, and computation — and the pre-registered standard properly applies to the forward content of this chapter, which awaits its data. The framework's physical content beyond conventional physics is structural extension of the same framework that produced the empirical predictions of Chapters 5-17, not speculative addition to an empirical base.

The chapter is honest about what it does not address. The hard problem of consciousness is not solved. The cosmological initial state is not explained. Several other open problems remain in the framework's residue. Chapter 19 develops these residues in detail, alongside the resolved-problem inventory that makes the framework's cumulative empirical reach explicit.

Chapter 19

Resolved and Open Problems in Fundamental Physics

19.1 What this chapter develops

The framework's content across the empirical chapters (5 through 17) and the forward-predictions chapter (18) constitutes a substantial body of structural commitments and quantitative predictions. This chapter inventories the framework's content against the *standard list of open problems* in fundamental physics — the canonical questions every framework in physics is expected to address.

The inventory has two parts. Section 19.2 develops the problems for which the framework claims a resolution, dissolution, conditional construction or partial treatment — including quantum gravity, the cosmological constant discrepancy, dark matter and dark energy, the measurement problem, black-hole information, hierarchy, gauge structure, generation structure, and the dark-sector budget — together with split-status entries: the origin of the Born rule, whose fixed-basis representation is settled exactly while the coherent operational extension is characterized by five named conditions (§19.2.12), and the strong CP problem, where the construction supplies a mechanism that does not reach the Standard Model. Each entry pairs the standard status of the problem (how it is conventionally formulated) with the framework's status (how the framework engages it), with

a cross-reference to the chapter and section where the relevant content is developed.

Section 19.3 develops six problems the framework does *not* resolve. The fermion mass hierarchy (flavor problem) has a defined research program with initial computational results and three tractable open targets, but the full quantitative content has not yet been derived. The Hubble tension is *potentially* addressable through ν_{OI} but at the framework's predicted magnitude the effect is too small to explain the observed tension. Baryogenesis is solution-specific: the framework provides standard leptogenesis ingredients but the baryon asymmetry is downstream of the flavor-sector CP phases the framework inputs rather than derives, so η_B inherits that input status (§19.3.3). Inflation and initial conditions are *not addressed*: the framework reproduces the Standard Model structure but does not address the very early universe's dynamics, with three possibilities outlined but none developed. The initial state of (S, φ) is partially gauge (within the reconstruction theorem's equivalence class) and partially the framework's irreducible mystery (why this equivalence class rather than another). The sixth entry is the chirality hypothesis H-χ' (§19.3.8) — a framework-specific structural hypothesis rather than one of the seventeen standard problems, so the headline tally's "five open" counts the standard entries — now in sharpened form: the minimal SU(2) embedding is certified taste-blind — vector-like, exactly as color's is — and the hypothesis asks whether a non-minimal mechanism (a taste-asymmetric condensate structure) restores selectivity — a class now certified nonempty, with dynamical selection the open question — posed in an exact finite arena, the 32-dimensional admissible condensate space splitting evenly into sixteen blind and sixteen flipping directions (`hchiprime_probes.py`, `condensate_space_probes.py`); the Standard-Model chirality identification remains conditional on it (SM §4.8), with the grading identification and both blindness certifications probe-backed. Section 19.3 also takes up two framework-specific meta-questions: whether the discrete structural residue is genuinely non-tautological (§19.3.6) — now *resolved* by a three-tier classification of the framework's results, which is the section's main conceptual addition and reframes the framework's overall claim as a tight compression of physics rather than a derivation of it from first principles — and the radiative stability of emergent Lorentz invariance in the rotational sector (§19.3.7), where a symmetry argument and a first numerical test support protection, with specific questions remaining.

Section 19.4 closes the chapter with a summary table and a note on what the framework does *not address*: specific experimental anomalies (muon $g-2$, B-physics, W mass) at scales where the framework's $\mathcal{O}(\epsilon^2 E^2)$ corrections are unobservable, condensed matter and many-body phenomena that are problems within the derived QFT rather than at the framework level, and the framework's principled scope limit on the hard problem of consciousness (developed in Chapter 18 §18.10).

The chapter functions as a systematic reference resource. Readers wanting to

understand how the framework engages a specific open problem can find the framework's status and the cross-reference to where the relevant content is developed. The chapter is not a substitute for the detailed treatments in Chapters 5 through 18; it is an inventory making the framework's reach explicit.

A note on tone. The inventory is honest about the framework's actual content. Where the framework resolves a problem, the inventory states the resolution and provides the cross-reference; where the framework only partially addresses a problem, the inventory marks the partial status and identifies what remains open; where the framework does not address a problem at all, the inventory acknowledges this rather than overstating the framework's reach. The closing summary table makes the distribution of statuses explicit.

19.2 Problems resolved

This section develops the framework's claimed resolutions, dissolutions, and conditional constructions, with quantum gravity as a partially addressed entry, the black hole information paradox as an unresolved entry whose information conservation is a premise and whose Page curve is conditional, and the Born-rule origin and the strong CP problem alongside them as split-status entries. Gauge-group origin and generation structure are explicitly conditional on the carrier bridge rather than counted as closed results.

19.2.1 The quantum gravity problem

Standard status. The central problem of theoretical physics. Quantum mechanics and general relativity are individually precise and mutually inconsistent — quantizing gravity leads to non-renormalizability, and no complete quantum theory of gravity exists. Substantial programs (string theory, loop quantum gravity, causal dynamical triangulations, asymptotic safety, and others) have been developed over decades without producing a complete theory.

Framework status. Partially addressed; covariant unification open (G3). The framework proposes that quantum mechanics and general relativity are different projections of the same structure (S, φ): quantum mechanics from the trace-out (Chapter 1) and, conditionally on the open covariant rung G3, general relativity from Jacobson's thermodynamic route applied to the emergent area-entropy relation (Chapter 7). What is in hand is the proposed common substratum, the trace-out characterization under its stated conditions, and the horizon-thermodynamic identities and weak-field results of the G1 and G2 rungs. The covariant field equations, their coupling to sources and their continuum-limit consistency make up G3 and are open, so the framework does not yet supply a derived, dynamically complete theory of quantum gravity, and G4 is conditional on G3. Which level carries the gravitational field of a laboratory mass that the observer's description places in a spatial superposition is not yet fixed; fixing it determines the framework's prediction for interferometric tests of gravity (Chapter 7 §7.7). The lattice at $\epsilon = 2\sqrt{\sigma}\, l_p$, with σ the entropy of one boundary mode,

is the framework's proposed fundamental description: on that proposal no continuum limit of the lattice is taken and no renormalizability problem arises, while that its long-wavelength limit reproduces the covariant equations belongs to G3. The framework produces the quantum and gravitational sectors from the same axioms — the fixed-basis quantum representation internal and universal, its compressed form under the translation hypothesis (T) ([Main §3.4]) — with the consistency relations collected in [GR §6.2]: several conditional on G3, one (the gauge-coupling values) reproduced through three matched parameters, and one (the MOND scale) with an imported factor.

The framework's treatment has three components, developed as a cumulative case in Chapter 18 §18.5: the cosmological-constant dissolution, established at the descriptive level with its dynamical form conditional on the covariant coupling theorem (§19.2.2); the black-hole information account, in which information is conserved by the bijectivity premise and the Page curve is conditional; and the non-renormalizability reading, which takes the lattice as the fundamental description and is conditional on G3. All three follow from the same structural commitment to the geometry-first ordering and the substratum-emergent operator distinction, and on the framework's reading the cosmological-constant and non-renormalizability obstacles are artifacts of the quantum-first ordering rather than genuine physical difficulties. They are framework-level interpretations and conditional results, not a completed resolution of quantum gravity.

Developed in: Chapters 1, 7, and 18. The three components are developed in Chapter 7 §7.7 (cosmological constant), Chapter 7 §7.6 (black hole information), and Chapter 18 §18.5 (non-renormalizability and cumulative case).

19.2.2 The cosmological constant problem

Standard status. The worst quantitative discrepancy in physics. Quantum field theory predicts a vacuum energy of order $M_{\mathrm{Pl}}^4 \sim 10^{113}\,\mathrm{J/m}^3$, summing zero-point energies $\frac{1}{2}\hbar\omega$ over all field modes up to the Planck cutoff. The observed vacuum energy density is $\rho_{\mathrm{obs}} \sim H^2/G \sim 10^{-9}\,\mathrm{J/m}^3$. The ratio 10^{122} has been treated as a fine-tuning problem requiring an unknown cancellation mechanism — a cancellation that must be re-imposed at each order of perturbation theory.

Framework status. Descriptive distinction established; the dissolution is conditional on the covariant coupling theorem that [GR §6.2] leaves open. The 10^{122} discrepancy is, up to a convention-dependent factor, the de Sitter entropy $S_{\mathrm{dS}} = A_{\mathrm{horizon}}/(4l_p^2)$ — on the framework's reading the information compression ratio between the substratum and the emergent QFT description, not a physical energy. The QFT zero-point energy belongs to the emergent description; that it does not gravitate is the coupling step that [GR §6.2] leaves open. The observed vacuum energy $\rho \sim H^2/G$ is the classical baseline set by the Friedmann equation at the substratum level. The two energy scales are not commensurable as gravitational sources; the conventional formulation treats them as comparable and asks why one cancels the other, while the framework treats them as

belonging to different levels of description with no cancellation required if that coupling step holds.

Developed in: Chapter 7 §7.7.

19.2.3 The nature of dark matter

Standard status. Approximately 27% of the universe's energy budget is in a form that clusters gravitationally but does not emit or absorb light. Decades of direct-detection experiments — XENON, LUX, SuperCDMS, PandaX, and others — have produced no detection at any expected mass scale. The conventional cold-dark-matter paradigm postulates a new fundamental particle species; the framework's content is that no such particle exists.

Framework status. Conditional: the chain inherits the conditions on the de Sitter temperature ([GR §3.2]) and the coupling statement $Q = M_B c^2$ that the open G3 map would supply. Dark matter is not a particle — it is entropy displacement from the cosmological horizon. The Clausius relation applied to the de Sitter thermal reservoir gives the MOND critical acceleration $a_0 = cH/6 \approx 1.1 \times 10^{-10}\,\mathrm{m/s^2}$, the baryonic Tully-Fisher relation $v^4 = GM_B \cdot cH/6$, and the radial acceleration relation — all parameter-free once the dimensional factor is fixed. The dimensional factor 1/6 is imported from Verlinde's emergent-gravity analysis as $(d-3)/[(d-2)(d-1)]$ in $d = 4$ (Chapter 7 §7.9). The simple MOND interpolation function $\nu(y) = (1 + \sqrt{1 + 4/y})/2$ matches deep-MOND clusters such as Coma to better than 1%, though rich-cluster cores (Perseus, A2029, A1689) retain a 1.0–1.5× residual that is an open problem: it is not closable by undetected baryons (massive clusters are baryon-complete near the virial radius, and the late-time missing baryons sit in cosmic filaments between clusters, not inside them) nor by the framework's predicted light neutrinos ($\Sigma m_\nu \approx 0.059$ eV), and most plausibly reflects the unresolved MOND-like-to-CDM-like crossover in the entropy-displacement relaxation kernel — the single highest-leverage open computation in the dark sector. The Bullet Cluster's anomalous gravitational lensing is explained by frozen boundary entropy with relaxation timescale $\tau_{\mathrm{relax}} \sim H^{-1} \gg t_{\mathrm{collision}}$. The CMB acoustic peaks are reproduced by thermodynamic averaging. The Milky Way Keplerian decline beyond the crossover (Jiao et al. 2023) and the high-redshift declining velocities of Genzel et al. are in tension with the law, which gives nearly flat outer curves and, through $a_0(z) = cH(z)/6$, flatter curves at high redshift for a fixed baryonic mass profile (Chapter 7 §7.9).

Developed in: Chapter 7 §7.9.

19.2.4 The nature of dark energy

Standard status. Approximately 68% of the energy budget drives the universe's accelerated expansion. Whether dark energy is a cosmological constant (constant in time) or evolving (the running-vacuum and quintessence proposals) is the subject of active investigation. The DESI Data Release 2 reports 2.8–4.2σ

preference for evolving dark energy in the $w_0 w_a$CDM parameterization, with the empirical situation in flux as of 2025-2026.

Framework status. Channel is a structural reading; functional form non-discriminating. The framework predicts that dark energy is the uniform component of the boundary entropy's gravitational effect, with the effective cosmological constant in the Type II running-vacuum channel — matter conserved, compensation through running G:

$$\Lambda_{\text{eff}}(H) = \Lambda_0 + \frac{3\nu}{8\pi G}(H^2 - H_0^2) + \mathcal{O}((H^2 - H_0^2)^2).$$

The functional form is the local expansion shared by every smooth model ([GR §7.1]), so the channel and the coefficient carry the content; the coefficient is bounded at the emergent-QFT scale $|\nu_{\text{OI}}| \sim 10^{-32}$, observationally indistinguishable from zero at any current or planned instrument. The Bertini *et al.* 2025 RG-corrected ΛCDM fit to Planck 2018 + DESI DR2 + DES-Y5 reports $\nu = -(2.5 \pm 1.3) \times 10^{-4}$, consistent with the framework's prediction at 2σ. DESI DR2's own mild preference for phantom crossing is, against the structural prediction $w(z) \approx -1$, a 3–4σ tension, carried on its own dated schedule ([GR §7.1]; Chapter 7 §7.8). DESI Year 5 (2028-2029) will provide a decisive three-way test between the framework's $\nu \approx 0$, intermediate $|\nu| \sim 10^{-4}$, and high-confidence phantom crossing.

Developed in: Chapter 7 §7.8.

19.2.5 The measurement problem

Standard status. What constitutes a measurement? Why does wave-function collapse occur? Why does the classical world appear definite? No consensus exists across Copenhagen, Many-Worlds, Bohmian, QBist, dynamical collapse, and other interpretations.

Framework status. Observer-relative account. In the proposed deterministic substratum interpretation, the wave function represents an observer's incomplete information. At the represented fixed-basis level, measurement conditions on a visible outcome and re-marginalizes the hidden sector. The exact finite-law correspondence admits that readout universally ($S \iff D \iff Q_{\text{fb}}$, [Main §3.4]); a full operational completion requires the separately stated matrix setting and completion conditions (§19.2.12). The ordinary Wigner's-friend interpretation assigns a definite substratum record and observer-relative information about it. It does not settle every measurement protocol or the Frauchiger-Renner certainty and consistency assumptions. Those require the proposed completion's operational probabilities and a separate premise analysis.

Developed in: Chapter 1, with related content in Chapter 18 §§18.2 and 18.4.

19.2.6 The black hole information paradox

Standard status. Does black hole evaporation destroy information? Hawking's 1975 semiclassical calculation produces thermal radiation that appears to carry no information about the black hole's initial state; if the radiation is exactly thermal, information is destroyed when the black hole completes evaporation — in tension with quantum mechanics' commitment to unitarity. The AdS/CFT correspondence and the recent quantum-extremal-surface and entanglement-island programs suggest information is preserved, but the mechanism in the bulk has been unclear.

Framework status. Information is conserved by the bijectivity premise; the paradox is not resolved, and the Page curve is conditional. The substratum map φ is bijective by premise (Chapter 1), so information is never lost from it; how much of it the radiation carries, bijectivity does not settle. The Page curve follows, as an average over the cycles of the dynamics, under the scrambling condition H-scramble and the decoupling condition H-decouple ([GR Appendix A]), neither of which is established: H-scramble carries Popescu-Short-Winter measure concentration to the orbit of the deterministic substratum dynamics on the full product space of the black hole and its radiation, whose cycle structure supplies exact time averages; it fails for every bijection acting on an initial configuration with the bipartition aligned to the configuration coordinates, the setting the derivation counts, so the Page value requires an initial superposition over at least about $e^{S_{\text{Page}}}$ configurations (S_{Page} the Page entropy) or a non-aligned bipartition; on a proper energy shell the value is open. The Page time is $t_P = [1-(\beta/(1+\beta))^{3/2}]\, t_{\text{evap}}$, independent of mass for a mass-independent emission coefficient, where β is the radiation's entropy per unit of horizon entropy lost: $(1-2^{-3/2})\, t_{\text{evap}} \approx 0.646\, t_{\text{evap}}$ at the mode-counting value $\beta = 1$, and earlier for $\beta > 1$, which holds in the semiclassical emission regime defined in Chapter 7 §7.6 and there depends on the species content and the greybody factors. The derivation runs through a mechanism substantially different from the quantum-extremal-surface and entanglement-island programs — cycle typicality on the configuration space rather than replica wormhole computation in the gravitational path integral. The conjectured correspondence "replica wormholes $\leftrightarrow$ substratum cycles" is suggestive but not yet rigorous; making it precise would require constructing the gravitational path integral as an effective description of the substratum dynamics in the appropriate limit. The generalized second law is open: the subadditivity bound on the sectors' reduced entropies becomes it only with an identification of horizon entropy with the hidden sectors' entropy that the boundary-mode count does not supply (§7.6).

Developed in: Chapter 7 §7.6.

19.2.7 The strong CP problem

Standard status. Why is the QCD vacuum angle $\bar{\theta} < 10^{-10}$? Naively, $\bar{\theta}$ should be $\mathcal{O}(1)$ as a generic parameter of the QCD Lagrangian. The Peccei-

Quinn mechanism (axion) is the leading conventional solution, but the axion has not been detected despite extensive searches.

Framework status. Not resolved. T-invariance of the substratum gives $\theta \in \{0, \pi\}$, not $\theta = 0$: θ is periodic and T sends $\theta \to -\theta$, so both are T-invariant points, and selecting the $\theta = 0$ branch is a separate condition (**H-top**). Reciprocity of the visible transition counts, $T_{ij} = T_{\theta j, \theta i}$, holds exactly at every time scale, but it does not give a T-invariant emergent Hamiltonian: a bipartite C_4 ring with complex hopping z has $|U_{ij}(t)|^2 = |U_{ji}(t)|^2$ at every t while its loop product z^4 carries a phase no diagonal rephasing can remove [SM, Prop. 20]. The Yukawa determinant phase, and with it $\bar{\theta} = \theta + \arg\det(Y_u Y_d)$, is therefore not fixed by the construction [SM, Theorem 21]. The remaining target is narrower than full reality: **H-det**, $\arg\det(Y_u Y_d) = 0$ with the flavor quartets left free — a mechanism making Y_u, Y_d simultaneously real would also force $J_{\mathrm{CKM}} = 0$, which is the falsification control on any proposal. Reciprocal probabilities can hide genuine loop phases, so detailed balance at the substratum does not kill weak CP violation, and the observed CKM phase is compatible with the construction as solution-level data. What the construction establishes is what time-reversal symmetry does and does not fix; what it does not have is a mechanism for H-top or H-det. A reported computation on the staggered reconstruction, not available for verification, found that an octahedral-equivariant emergent operator gives $J = 0$ and that restoring the observed J would require octahedral breaking far above the rotational bounds; it is recorded as an unverified tension for the flavor sector [SM §5.5].

Developed in: Chapter 6 §6.3.

19.2.8 The hierarchy problem

Standard status. Why is the Higgs mass (~ 125 GeV) so much lighter than the Planck mass ($\sim 10^{19}$ GeV)? Radiative corrections in a generic effective field theory should drive m_H to the cutoff unless an unknown mechanism — supersymmetry, compositeness, extra dimensions, or anthropic selection — protects the Higgs mass from quadratic divergences. None of the conventional protection mechanisms has been confirmed empirically.

Framework status. Partially addressed: cutoff insensitivity (SM §8.8); $v \ll f$ open. The Higgs mass is set by the block values of the equivariant condensate Σ on the cubic lattice substratum (SM Theorem 5) — a property of the specific bijection φ, not a radiative correction within the emergent QFT. The lattice IS the UV completion; there is no hierarchy of scales to explain because the Higgs mass is set at the lattice level (where the framework's structural commitments operate), not by a running coupling within the emergent QFT. Quadratic divergences are artifacts of treating the lattice spacing as a regulator to be removed — the same artifact that produces the non-renormalizability discussed in Chapter 18 §18.5. The framework's substratum lattice eliminates the divergence and with it the cutoff sensitivity of m_H. What is addressed here is the

naturalness problem — the need for a mechanism protecting m_H from quadratic divergences; the *residual* electroweak hierarchy (why the specific value v/M_{Pl} takes the number it does) is a property of the bijection φ and remains an open input.

The composite Higgs structure developed in Chapter 6 §6.6 provides the specific framework realization: the Higgs is a composite field of the framework's specific substratum content, with the W, Z, and Higgs masses emerging from the block values of the equivariant condensate Σ (SM Theorem 5) rather than as parameters in a renormalizable Lagrangian.

Developed in: Chapter 6 §6.6.

19.2.9 The origin of the gauge group

Standard status. Why $\mathrm{SU}(3)\times\mathrm{SU}(2)\times\mathrm{U}(1)$? Why these multiplicities for the matter content? Why this specific representation structure? The conventional account treats the Standard Model gauge group as empirically determined input rather than derived from any deeper principle.

Framework status. Conditional physical-carrier construction with an exact geometric core. The six signed nearest-neighbor links of the simple-cubic graph form an exact six-dimensional cubic-group representation and decompose as $(3, 2, 1)$. What is *not* presently derived is the identification of this geometric link representation with the complete physical gauge carrier: that is H-link, with the single-copy clause $m = 1$ giving $K = 6$; the condensate-stabilizer route additionally uses H-cust. Conditional on those named bridges and the later chirality/matter hypotheses, the anomaly and representation calculations recover the Standard-Model gauge structure and hypercharges. The origin of the geometric $(3, 2, 1)$ decomposition is therefore resolved; the physical-carrier origin is an explicit open bridge.

Developed in: Chapter 5 (gauge structure) and Chapter 6 (matter content and quantitative predictions).

19.2.10 The generation puzzle

Standard status. Why three generations? Why do all three have identical gauge quantum numbers but hierarchical masses spanning roughly five orders of magnitude? The Standard Model accommodates three generations as empirical input but has no explanation for the multiplicity or for the mass hierarchy pattern.

Framework status. Partially derived and carrier-conditional. On the H-link single-copy branch, three candidate sectors arise from the staggered/cubic taste structure. The T_1 representation of the cubic group has dimension three, and anomaly cancellation forces exactly three candidate matter sectors with the framework's link-carrier construction (their reading as physical generations rides H-spin'). Identical gauge quantum numbers follow because all three generations

are copies of the same T_1 irrep. Hierarchical masses follow because the specific bijection φ breaks the cubic symmetry through taste-breaking at $\mathcal{O}(\epsilon^2)$, with the hierarchy emerging from the second-order perturbative structure.

The *existence* of a hierarchy is structural — any bijection φ breaking the cubic symmetry produces a hierarchy. The *pattern* of the hierarchy — which generation is heaviest, the specific mass ratios, the CKM mixing angles — is solution-specific, requiring the specific properties of φ to derive. The framework has an exact singlet-plus-triplet geometric decomposition; interpreting the triplet as three physical generations requires H-link single-copy together with H-spin', and the specific pattern is the open task developed in §19.3.1 (the flavor problem).

Developed in: Chapter 6 §§6.5 and 6.6, with the flavor open problem at §19.3.1 below.

19.2.11 The dark sector budget

Standard status. Why is the universe approximately 5% baryonic, 27% dark matter, and 68% dark energy? No conventional framework explains these ratios from first principles. The dark sector's $\sim 95\%$ total is treated as empirical input.

Framework status. Conditional (total budget partly definitional; the coefficient of S_{dS} inherits the horizon conditions of Chapter 7). The trace-out renders approximately 95% of the critical density invisible to the emergent quantum field theory. The horizon's entropy $S_{\mathrm{dS}} = A_{\mathrm{horizon}}/(4l_p^2)$ combined with the classical horizon temperature gives, by the Clausius relation and the shell theorem, the total effective density $\rho_s = 3c^2H^2/(8\pi G) = \rho_{\mathrm{crit}}$. The uniform component is dark energy; the structured component (entropy displacement around baryonic matter) is dark matter. The 95% dark sector is concordance rather than puzzle — though the total fraction is partly definitional, equal to $1-\Omega_B$ by construction: the framework's structural quantity for the partition's hidden-sector content at the cosmological horizon equals the observed dark sector fraction.

The specific 27/68 split between dark matter and dark energy depends on the matter distribution and the cosmic expansion history, and is not derived as a first-principles result. The framework derives the total budget; the partition into dark matter and dark energy components is a function of the universe's matter content and dynamics rather than a structural prediction.

Developed in: Chapter 7 §7.9.

19.2.12 The origin of the Born rule

Standard status. The Born rule — that the probability of measuring quantum state $|x\rangle$ given wave function $|\psi\rangle$ is $|\langle x|\psi\rangle|^2$ — is an independent postulate of quantum mechanics with no derivation from the other axioms. Gleason's theorem derives the Born rule from non-contextuality, but non-contextuality is itself a postulate. The various interpretive programs (many-worlds, QBism,

Bohmian, dynamical collapse) take different positions on what the Born rule means without explaining why it has the specific form $|\langle x|\psi\rangle|^2$.

Framework status. Split — fixed-basis representation settled exactly; coherent operational extension characterized by five conditions. Every finite-horizon stochastic law has an exact fixed-basis unitary/Born representation ($S \iff D \iff Q_{\mathrm{fb}}$, [Main §3.4]); in that theorem the quadratic Born dictionary is part of the equivalent representation, not a separate exponent-selection problem. The further question begins when one asks for **one common operational theory** covering arbitrary coherent preparations, noncommuting effects/instruments, and composites. Main's finite gluing theorem realizes every bounded quantum experiment to arbitrary requested finite accuracy, but the same machinery can also realize non-quantum finite behaviors; the operational-completion characterization ([GR §3.3]) states exactly which conditions select the quantum ones — valid probabilities, trivial-ancilla consistency, inert spectators, full reversible control, and iterated composition — each independent of the others and of the observation process, so that embedded observation by itself does not select quantum mechanics and the physical question is which of those conditions the substratum supplies. The characterization takes complex matrix kinematics as given — operations act on the complex matrices indexed by the configurations of the system and of its composites — and deriving that kinematics is a further open question. Its layered form, quantum-complete OI (OI^+) — the unchanged OI core with well-formedness, observational independence, reversible richness, and observer recursion — is equivalent on every nonempty finite carrier to exact finite endomorphic operational quantum mechanics, with no claim that bare OI entails the added principles ([GR §3.3]). A primitive-source refinement of that layered form restates the three principles at the level of physical implementations and embedded observers — implementation locality, phase-free richness — one continuously driven transition and the exchanges of distinguishable states, with no phase operation — and embedded observation, with no closure of the implementations under the adjoint and no inverse accessibility assumed — equivalent to the same operational quantum mechanics on every nonempty finite carrier ([GR §3.3]). A substratum-source form completes that refinement: the concrete OI substratum supplies the structural architecture — closure under composition and coarse-graining, stability under an uncoupled spectator, relabelling invariance, reversibility, the exchanges of distinguishable states, and, as a stated intervention principle rather than a consequence of its finite states and bijective dynamics, the phases — but, under finite bijective read-write dynamics, not the continuously tunable state-mixing operation; with that one operation added as an empirical controllability resource it is equivalent to the same operational quantum mechanics on every nonempty finite carrier, while finite reversible read-write dynamics, even with genuine hidden-memory and readback behavior, does not itself generate quantum state mixing ([GR §3.3]). Its typed form removes the endomorphic qualifier: under the carrier-general typed operational interface — identity, composition, coarse-graining, relabelling, uniform finite attachment, discard, and factor readout, an

interface that itself carries no quantum content — the same equivalence holds with exact finite-dimensional typed operational quantum mechanics, operations between any two nonempty finite carriers included, so the remaining qualifier is finite-dimensional rather than endomorphic ([GR §3.3]). The finite theory admits a canonical infinite-region completion without a continuum limit: the finite regions of the fixed-spacing lattice carry full matrix algebras joined by compatible injective unital embeddings with observables of disjoint regions commuting, the norm completion of their union is a quasilocal C^*-algebra on which every consistent family of finite-region density matrices is a state and the reversible finite-range substratum update is an isometric $*$-automorphism, and any system carrying the same local stages is canonically and uniquely $*$-isomorphic to it, intertwining the dynamics; this removes the finite-region cutoff for the algebra, the state space, and the OI-induced discrete dynamics rather than characterizing all infinite-dimensional quantum instruments or all locality-preserving quantum dynamics; the lattice spacing stays fundamental, no Hilbert-space representation and no superselection sector is selected at the level of the laws, the restriction to the substratum-induced dynamics is a theorem rather than a convention — a phase automorphism at one site preserves locality without arising from any reversible finite-range bijection — and, while the update is joined to the identity by an exact strongly continuous path of $*$-automorphisms of that algebra, with no one-parameter-group law established and no generator exhibited, a continuous-time Hamiltonian law remains additional structure ([GR §3.3]). Within the structural architecture the substratum supplies, with the phase intervention stated, exact finite endomorphic operational quantum mechanics holds exactly when one continuous layer flow of an involution of the configuration space moving some configuration is an available operation at every level and every intermediate time, the executability of that flow being an assumption not derived by configuration-level generation, as the phase intervention is an assumption on the substratum class and not a consequence of the finite states with the bijective dynamics ([GR §3.3]). The stated substratum dynamics with the observer's read and write access do not source that executability, the layers being available at time one and not at intermediate times, and the observer-level lift, in every formulation stated that lands in the operational interface, derives neither a non-monomial admissible operator nor the layer flow, the site-space wave-operator formulation being unresolved rather than excluded ([GR §3.3]). The cosmological realization adds a mechanism proposal — the equilibrium reading of the read-write cycle (Chapter 18 §18.2) — whose stationary-distribution step is a structural commitment rather than a premise of the representation theorem.

The mechanism proposal of Chapter 18 §18.2 implies transient deviations from the Born rule in the very early universe if the hidden sector began away from equilibrium, relaxing on a timescale the proposal identifies with the hidden sector's equilibration time; that timescale sets the order of the relaxation rate, while the form of the decay requires in addition a closed relaxation law and its rate coefficient, neither of which is derived. Their observational reach depends

on the source dynamics, the family of initial deviations and the map to the CMB's primordial power spectrum and non-Gaussianity, which the proposal does not yet fix; CMB measurements of those statistics are the channel in which they would be tested once these are fixed.

Two frontiers, not one. The split recorded above concerns the *operational* extension: which conditions select one common theory of coherent preparations, instruments and composites, answered conditionally by the completion principles. A second and separate question is left open by that answer — which unitary evolution relates one time to the next, given a visible OI family that admits several coherent lifts. Ordinary coherence does not settle it, and the completion characterization does not address it. That frontier is recorded at §19.3.9; the two should not be read as one residue.

Developed in: Chapter 1, with forward content in Chapter 18 §18.2.

19.3 Problems open within the framework

This section develops the seven entries the framework does not fully resolve — the five standard open problems counted in the headline inventory, plus two framework-specific items internal to the framework rather than to the seventeen standard problems: the chirality hypothesis H-χ' (§19.3.8), internal to the Standard-Model identification, and the selection of the relative quantum evolution (§19.3.9), internal to the OI–QM bridge. Neither framework-specific item changes the count of standard open problems. Each entry pairs the standard formulation, the framework's current status, an assessment of what remains open, and a note on the prospects for closing the open question.

19.3.1 The flavor problem

Standard status. Why do fermion masses span five orders of magnitude, with $m_e/m_t \sim 3\times10^{-6}$? Why the specific pattern of CKM and PMNS mixing angles? Why the Koide relation $Q = (m_e + m_\mu + m_\tau)/(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2 = 2/3$ to better than 0.001%? The Standard Model accommodates the fermion masses as 13 free parameters; no conventional framework derives them.

Framework status. Open with a defined research program and initial computational results. The fermion masses are taste-breaking parameters of the substratum bijection φ — properties of how the specific φ breaks the cubic symmetry. The structural input $1/\alpha_0 = 23.25$ is computed from the one-loop staggered vacuum polarization, and the three gauge couplings at M_Z are reproduced with three matched threshold parameters (Chapter 6 §6.4), showing that solution-specific parameters are constrainable in principle. The research program identifies three tractable open targets.

The bottom-to-tau mass ratio. The framework predicts $y_b = y_\tau$ at tree level — the staggered taste-breaking is diagonal in the gauge index. Full Standard

Model RGE running (verified against Luo and Xiao 2003) gives $m_b/m_\tau|_{\text{tree}} = 4.276$ from a common M_{Pl} boundary condition, indicating a substantial non-perturbative correction. The full prediction including the scalar-density renormalization Z_S from non-perturbative dynamics at the matching scale is developed in Chapter 6 §6.6 (and SM §7.5): with matching scale $m_{\text{match}} = \lambda g_0^2 = 0.1217$ derived structurally from the taste-decomposed Coleman-Weinberg potential, and $Z_S(m_{\text{match}}) = 1.813$ measured from Monte Carlo at $\beta = 11.1$ on $L = 32$, the prediction is $m_b/m_\tau = 4.28/Z_S(m_{\text{match}}) = 2.361$ against the observed 2.352 ± 0.017 — a 0.4% (0.5σ) match at $L = 32$, with $L \to \infty$ extrapolation giving $\sim 0.7\sigma$. The $L = 16$ Monte Carlo using only the SU(3) mass renormalization $Z_m \approx 0.835$ at fixed bare mass — the calculation described in earlier versions of this section as giving ~ 3.4 (45% off) — corresponds to using the wrong matching prescription rather than the wrong physics. With the correct matching scale (a structural feature of the taste-decomposed CW potential rather than a tunable parameter), the prediction matches observation at the same level as the framework's other Layer 1 closures (Cabibbo, Koide, $\sin^2\theta_{13}$). The kinematic prefactor $K = 1/(n_{\text{gen}} - 1) = 1/2$ is a Layer-1 structural identity given the matching condition ($c_{bb} = (n_{\text{gen}} - 1)\lambda$); its support is the Monte Carlo verification at 1.4% ($K_{\text{MC}} = 0.507$). The explicit one-loop evaluation has been carried out and cannot close the gap: in the condensate-ratio construction the one-loop coefficient is 0.446 (tadpole-normalization bracket 0.393–0.446), 11–21% below 1/2 at a coupling where $Z_S - 1 = 0.81$ — a truncation error an order of magnitude larger than the MC precision, indicting the truncation rather than the identity. Analytical closure requires the resummed staggered-χPT chain or a non-perturbative identity, or alternatively sub-percent MC; the task is Layer 2(a) in scope but not one-loop in method (SM §7.5).

The Koide relation. $Q = 2/3$ is equivalent to the square-root mass vector making a 45° angle with the $[1, 1, 1]$ direction in generation space — equivalently, the amplitude $A^2 = 2$ in the $\mathbb{Z}_3$ mass parametrization (SM §7.2). This has an exact structural target: $A^2 = C_2(T_1) = 2$, the same quadratic Casimir that fixes the Koide angle $\theta_0 = C_2/d^2$, so the amplitude and the angle descend from one invariant. The open question is therefore dynamical and sharper than "whether the 45° has a structural origin": one-loop perturbative taste-breaking gives $A^2 \approx 0.34$, so the task is to show that the non-perturbative substrate dynamics produces $A^2 = C_2 = 2$ — a $\sim 6\times$ enhancement to a *known* target — via non-perturbative enhancement of the E-component (the structural route, under which the 45° is derived) rather than solution-specific T_2 off-diagonal elements (under which it would be bijection-specific). The full $K = 6$ lattice Monte Carlo is the tool, and whether it lands on C_2 is the key open question. The observable class is constrained in advance: an exact shift-symmetry identity makes the ensemble single-quark two-point function eight-fold corner-degenerate at every coupling, so quark-level taste splitting vanishes identically and the measurement must use composite (multi-fermion) channels — closing the otherwise-natural quark-propagator route before it is attempted (SM §7.2).

The Wolfenstein hierarchy. Perturbative taste-breaking naturally produces

$|V_{us}| \sim \lambda$, $|V_{cb}| \sim \lambda^2$, $|V_{ub}| \sim \lambda^3$ — the observed CKM pattern. The value $\lambda = 0.22508$ is the Cabibbo angle from cubic Brillouin geometry (Chapter 6 §6.5 and Chapter 8 §8.3) — a Tier 2 structural result. The hierarchy pattern is therefore structurally fixed; the specific magnitudes depend on the bijection.

Neutrino masses. Majorana neutrinos, normal ordering, large PMNS mixing, and a hierarchical spectrum are expected on the branch where the singlet taste hosts the Higgs and no gauge-singlet fermion is present (SM §8.4; the mechanism is otherwise open), the ordering and the hierarchy as heuristic consequences of taste-breaking rather than theorems (Chapter 6 §6.7 and Chapter 8). Specific mass values are solution-specific.

Required tools. Full $K = 6$ lattice Monte Carlo with the scalar coupling matrix $M = \mu I_6$ and the equivariant condensate Σ (SM Theorem 5); representation theory of the cubic group O applied to the framework's specific irrep content; 2-loop lattice-to-continuum matching. Four C programs implementing the perturbative taste-breaking spectrum, propagator-norm measurement, effective-mass extraction, and taste-split correlators are available as part of the framework's computational infrastructure.

Assessment. The flavor problem is the framework's largest single open theoretical task. The framework has substantive content on the *structural form* of the flavor pattern (the existence of three generations with hierarchical masses, the Cabibbo angle as Brillouin geometry, the Wolfenstein hierarchy, normal ordering for neutrinos) and on several specific magnitudes that are conditional structural at Layer 1 with one named open link each (Cabibbo λ, Koide Q, $\sin^2\theta_{13}$) or layered — m_b/m_τ with a Layer 2(a) verification residual, $\sin^2\theta_{12}$ and $\sin^2\theta_{23}$ on a Layer 2(b) substrate coefficient (SM §7.6). The remaining solution-specific parameters (specific CP phases, the full Yukawa pattern beyond hierarchy existence) are bijection-specific in the framework's terms — structurally predicted *not* to be derivable from substrate alone, consistent with the framework's observational-incompleteness commitment. The Layer 2(a) verification residual is a tractable open task with a defined computational path; the bijection-specific parameters are pre-registered as outside the framework's substrate-derivable scope.

19.3.2 The Hubble tension

Standard status. The Hubble constant measured from the CMB (Planck: $H_0 = 67.4 \pm 0.5$ km/s/Mpc) disagrees with the local distance ladder (SH0ES: $H_0 = 73.0 \pm 1.0$ km/s/Mpc) at greater than 5σ. The disagreement has persisted across multiple measurement systematics, suggesting either an unidentified systematic in one or both methods or new physics modifying the cosmic expansion history.

Framework status. Potentially addressable but at the framework's predicted magnitude the effect is too small. On its Type II running-vacuum reading the framework expects evolving dark energy with $|\nu_{\mathrm{OI}}| \sim 10^{-32}$ bounded at the

emergent-QFT scale, indistinguishable from zero. Quantitatively, in the RVM, the effective $H(z)$ differs from the ΛCDM prediction at low redshift:

$$H^2(z) = H_0^2 \left[\Omega_m(1+z)^3 + \Omega_\Lambda + \nu H^2\right].$$

The νH^2 term acts as a small additional dark energy component at late times, increasing the inferred H_0 relative to the CMB-inferred value. For $\nu \sim 2.5\times10^{-3}$ at the upper boundary of the framework's substratum-contribution prediction, the shift would be approximately $\Delta H_0/H_0 \sim \nu/(1-\Omega_m) \sim 0.4\%$ — far too small to explain the approximately 8% Hubble tension.

Assessment. The Running Vacuum Model with $\nu_{\rm OI}$ does not resolve the Hubble tension. The tension may be due to systematic errors in the distance ladder (currently active investigation by SH0ES and CCHP collaborations), new physics not captured by the framework's substratum dynamics, or a statistical coincidence becoming significant through accumulated precision. The framework makes no specific claim on the Hubble tension's resolution and explicitly does not contribute to its understanding beyond noting that the framework's predicted ν is too small to be the responsible mechanism.

19.3.3 Baryogenesis

Standard status. The observed baryon-to-photon ratio $\eta = (6.1 \pm 0.2)\times10^{-10}$ requires a mechanism to produce a matter-antimatter asymmetry satisfying Sakharov's three conditions: baryon number violation, C and CP violation, and out-of-equilibrium dynamics. The conventional candidates — electroweak baryogenesis, GUT baryogenesis, leptogenesis through heavy right-handed neutrinos, and various extensions — have not been confirmed empirically.

Framework status. Solution-specific. The framework does not derive the baryon-to-photon ratio and does not carry an obligation to — for the same structural reason it does not derive the CKM phase. This status is the conclusion of a sustained investigation; the reasoning is given below, and the honest record of that retraction is part of the framework's content.

The framework provides standard leptogenesis ingredients, with a shortfall. The framework's structural predictions provide ingredients for leptogenesis via the Weinberg operator $(c_{ij}/\Lambda)L_iL_jHH$: Majorana neutrinos (Chapter 6 §6.7), large PMNS phases, and hierarchical masses. The operator mediates $\Delta L = 2$ scattering in thermal equilibrium at $T > 10^{14}$ GeV. A precise calculation reveals a quantitative shortfall: the total unflavored CP asymmetry vanishes by CPT plus unitarity ($\sum_\alpha \epsilon_\alpha = 0$), and the surviving flavored asymmetry from thermal corrections is suppressed by $y_\tau^2 \sim 5\times10^{-5}$, yielding $\eta_B \sim 4\times10^{-11}$ — approximately 15× below the observed value of 6.1×10^{-10}. This is the same shortfall faced by any framework with Majorana neutrinos but no heavy right-handed neutrinos: the Weinberg operator alone, without a UV completion providing additional CP-violating phases, cannot produce sufficient baryon asymmetry.

A retracted mechanism. An earlier treatment proposed closing the factor-15 gap with *substratum-level instanton-like processes* — configurations of the bijection φ at the lattice cutoff with action $S \sim O(1)$, giving a suppression $e^{-O(1)}$ rather than the electroweak $e^{-2\pi/\alpha_W}$. This mechanism is **retracted**, and the reason is structural rather than quantitative. The substratum has no Euclidean action functional: it is the deterministic linear bijection of the wave-equation update rule (Chapter 2), not a path-integral system. "Substratum-level instantons with action $S \sim O(1)$" is therefore type-mismatched with what the substratum is — there is no action whose minima would constitute instantons. The correct picture is the one Chapter 6's treatment of baryon number already gives: substratum-level B is conserved at the bilinear level, and non-perturbative $B+L$ violation enters through the *emergent* fermion-determinant topology governed by standard sphaleron physics. There is no additional substratum-level baryon-number-violating channel.

Why baryogenesis is solution-specific. With the retracted mechanism removed, the framework's relation to baryogenesis is settled by a commitment the framework makes elsewhere and independently. The substratum is T-symmetric — this is load-bearing for the reciprocity result of §19.2.7, which does not by itself fix $\bar{\theta}$. Observed CP violation in the CKM and PMNS sectors is, in the framework, a flavor-basis feature of the *specific bijection* (S, φ) — not a derived quantity. The framework makes no prediction of δ_{CKM}, δ_{PMNS}, or the Wolfenstein (ρ, η); these are treated as solution-specific inputs, in the same category as the framework's choice of discretization scheme.

The baryon asymmetry inherits this status. Sakharov's second condition is CP violation, and the only CP violation available to the framework is exactly the flavor-sector CP it treats as a solution-specific input. A quantity downstream of an input is itself an input: η_B is no more derivable, within the framework's current commitments, than δ_{CKM} is. Demanding that the framework dynamically *generate* η_B from a symmetric initial condition holds it to a standard it does not adopt for the CP phases η_B depends on. The "factor-15 open problem" framing was an artifact of that inconsistency.

Assessment. Baryogenesis is *solution-specific*, not a quantitative gap awaiting a mechanism. The framework's structural ingredients (sphaleron $B+L$ violation, the out-of-equilibrium expanding early universe, the CKM/PMNS CP phases) are consistent with the observed η_B; what the framework does not do is derive its value, because that value is downstream of CP phases the framework inputs rather than derives. This is a reclassification that restores internal consistency — the framework no longer carries baryogenesis as an unmet derivation obligation while simultaneously inputting δ_{CKM} — and it is honest about its limits: it is not a closure. It does not produce $\eta_B = 6.1 \times 10^{-10}$ from first principles, and a reader may fairly regard a framework that derives the Cabibbo angle but inputs the baryon asymmetry as incomplete in scope. The framework's position is that this incompleteness is shared, explicit, and located precisely: it is the flavor-sector CP input, named once and inherited wherever CP violation

is needed.

19.3.4 Inflation and initial conditions

Standard status. The horizon problem (why is the CMB isotropic across causally disconnected regions?) and the flatness problem (why is $\Omega \approx 1$?) are solved by an early period of exponential expansion (inflation). But inflation requires a scalar field with a specific potential, which is not part of the Standard Model. The initial conditions for inflation are themselves fine-tuned. The empirical evidence for inflation comes through CMB polarization measurements and the absence of relics that pre-inflation cosmologies would produce.

Framework status. Not addressed. The framework reproduces the Standard Model structure but does not address the very early universe's dynamics. Three possibilities exist within the framework's structural commitments, none of which has been developed in detail.

The Higgs as inflaton. The framework derives exactly one scalar doublet (Chapter 6 §6.6). Higgs inflation (Bezrukov and Shaposhnikov, 2008) uses the Standard Model Higgs with a non-minimal coupling $\xi H^\dagger H R$ to drive inflation. If the framework's Higgs *is* the inflaton, inflation is automatic — but the non-minimal coupling $\xi \sim 10^4$ is a solution-specific parameter (a property of the bijection φ), and its value is not determined by the framework's structural commitments. The empirical predictions of Higgs inflation are well-developed in the literature; whether the framework's bijection produces $\xi \sim 10^4$ is an open question requiring the full bijection-specific calculation.

No inflation needed. The framework's lattice structure may resolve the horizon and flatness problems differently. On a finite lattice, the notion of "causally disconnected regions" is modified — the lattice's coupling graph determines causal structure, and if the initial state of φ is statistically isotropic (one of the framework's six structural properties of the bijection), isotropy is built in rather than requiring inflation to produce it. This is speculative and would require a detailed analysis of the lattice's causal structure at early times.

Initial conditions as gauge. The deep hidden sector's cardinality is gauge in the framework. If the initial state of the visible sector is also gauge-equivalent across a wide class of initial conditions (because the trace-out erases the distinction), then the "initial conditions problem" may dissolve rather than require solution. This is an intriguing possibility but has not been investigated within the framework's existing structure.

Assessment. Inflation is not derived, not refuted, and not distinctively addressed by the framework. This is a genuine open problem with the possibilities outlined but no current developed framework position. Pre-registered: the framework's content on the very early universe is one of the framework's most substantial scope limits in fundamental physics.

19.3.5 The initial state of (S, φ)

Standard status. Not applicable in the conventional sense — this is a framework-specific question. Why does the universe have the particular bijection φ rather than some other? Why these particular initial conditions for the bijection's evolution?

Framework status. Partially addressed, partially outside scope. The reconstruction theorem establishes — under its stated modeling, mixing and nearest-neighbor premises — uniqueness only of the **local propagating lattice/gauge residue**. The Bell-inclusive full substratum remains underdetermined: H-Bell conditions the existence of the adopted preparation-indexed Bell completion, and uniqueness of that Bell-composite rule remains open. The question "why this φ?" therefore has a gauge-local layer and an additional Bell-completion layer.

Within the reconstructed local equivalence class. Gauge — no answer is physically required for representative choices that Theorem 24 identifies as observably equivalent. This statement does not identify all Bell-inclusive completions with one another.

Why this equivalence class rather than another. The framework's axioms — deterministic bijectivity, finiteness, causal partition, classical probability — constrain the class. The framework's six structural properties of the bijection (developed across Chapters 5-7) further narrow it. But the axioms themselves are not derived from anything deeper.

The existence question — "why does (S, φ) exist at all?" — is the framework's irreducible mystery. The framework's structural account: (S, φ) is a mathematical structure that cannot fail to be well-defined; existence is the default for logically consistent structures. Whether one finds this answer satisfying is a philosophical question rather than a physical one. The framework's commitment is that the existence of *some* finite deterministic bijection structure is not a question physics can answer, while the existence of *this particular* structure (as determined by the empirical record) is a constraint on which bijections are compatible with observation.

Assessment. The framework identifies this question as the only genuinely irreducible mystery — not a gap in the framework's reach but the boundary of all possible frameworks. Any framework must take some structural axioms as given; the framework's content is to make those axioms as minimal and as structural as possible (deterministic bijectivity on a finite set, no further commitments), leaving the existence question as an explicit acknowledgment of where physics ends.

19.3.6 Is the discrete structural residue genuinely non-tautological?

Standard status. A framework-specific meta-question. The reconstruction theorem sends the empirical and structural axioms to a discrete residue — the

spatial dimension $d = 3$, the internal multiplicity $K = 2d = 6$, the gauge group SU(3) × SU(2) × U(1), three generations, the hypercharges. The question is whether these are forced and could have been otherwise (substantive content), or whether they are already contained in the inputs and the derivation unpacks them (tautological).

Framework status. Resolved — not by a calculation, but by recognizing that the question is mis-posed. It presupposes a single yes/no answer for "the residue," but the residue is not uniform: its members have three different logical statuses, and once they are separated the question has a definite answer for each. *Tier 1 — forced by representation theory (genuinely non-tautological).* Given the simple-cubic lattice and the wave equation, the multiplicities $(3, 2, 1)$ follow from the octahedral character table with no further freedom, and, on the carrier branch of H-link and H-cust, the gauge factors, the generation count of three (as physical generations under H-spin'), the anomaly-free hypercharges (on the nonzero-Y_Q branch, up to normalization), and, under its identification hypothesis, the Cabibbo angle follow from it and anomaly analysis; they could *a priori* have come out otherwise, and the predictive excess noted above (the Cabibbo angle, Koide, the PMNS angles, six mass ratios — none of them among the empirical inputs) is exactly the signature of genuine content. This tier is real and substantial. *Tier 2 — empirically selected.* Two of the listed invariants are *not* forced by the construction: $d = 3$ is selected by the three Standard-Model-independent filters (§3.2 of the SM paper), and the simple-cubic Bravais structure is selected, under nearest-neighbor coupling, by the observed SU(3) × SU(2) × U(1) — only the cubic crystal system supplies the three-dimensional point-group irrep an SU(3) factor needs (Theorem 7b). For these the inference runs from the observed world back to the commitment; "the Standard Model from a cubic lattice" and "the lattice is cubic because we observe SU(3)" are one fact stated in two directions. The threshold δ_0 and the absolute mass scales are likewise empirical. *Tier 3 — stipulation.* Site factorization (the identification of sites, which the dynamics has not been shown to select and coupling-degree minimization does not), the single-copy clause of H-link (from which $K = 2d$ follows), the nearest-neighbor restriction, and the C1–C4 architecture itself are choices about what counts as physical structure rather than facts the bijection forces; they are principled, but they are stipulations.

Assessment. The meta-question is therefore not open pending computation — no lattice calculation moves it, because the status of each link is already fixed by the construction as written. It is answered by the classification: non-tautological at Tier 1, tautological-in-the-relevant-sense at Tier 2, definitional at Tier 3. Several distinct concerns — the derived-versus-input ledger, the residual-input status of Lorentz structure (§19.3.7 and the boost sector of the SM paper), and the circularity question — reduce to this one and are resolved the same way. The constructive consequence is a precise statement of what the framework is: a *maximally tight compression* of observed physics onto the cubic-lattice commitment. Tier 1 establishes that, once the commitment is granted, the Standard Model's discrete structure is largely not independent — it is a

single representation-theoretic object — which is a genuine and unusual claim about the unity of physics. Tiers 2–3 establish that the commitment itself is empirically anchored and partly stipulated, so the framework does not derive physics from a parameter-free first principle. The earlier framing of this as "open, not closable by internal means" mistook a question about the *logical status of the construction's steps* — which the construction already answers — for a question awaiting evidence. The proposed external calibration against the operational reconstruction theorems of quantum theory (Hardy; Masanes–Müller; Chiribella–D'Ariano–Perinotti) remains a worthwhile comparison, but it bears on how to *value* a "uniquely-fixed" result, not on whether the present residue is forced; the latter is settled above. (The three-tier classification is developed in full in [SM §8.3] and [Substratum §3, Refinement]; its bearing on the framework's foundational claim is drawn out in [Structure §13.4, Limit 4].)

19.3.7 Radiative stability of emergent Lorentz invariance (rotational sector)

Two results bear on this question. A symmetric-integration argument shows that an exactly O_h-invariant action, measure, and observer-centered partition yield an O_h-invariant effective theory to all orders, and every ingredient of the checkerboard trace-out is O_h-invariant — no direction-dependent momentum shell is available. A numerical check on the rounded interacting dynamics, whose code and numerical output are not available for verification (extraction reported validated against the analytic tree coefficients to the percent level; the allowed $\ell = 4$ anisotropy reported detected at high significance), reported the forbidden E_g coefficient zero within errors at the 2×10^{-5} level in the perturbative regime and consistent with zero in the thermalized strong-coupling state. Separately, the sector-differential boost violation's custodial decoupling law — conditional on the symmetric-point matching under H-blind ([SM §4.4]) — has its normalization bounded ($\Delta C_2 \in [0.58, 2.08]$ from the framework's representation content), and its decoupling form, controls and kernel constants ($c_L \simeq -9.4$, $d \simeq -0.88$) reported by a one-loop calculation whose driver and numerical output are not available for verification; on that reported form the worst-case ΔC_2 would give an M_X window of two to five orders of magnitude against the tightest multi-species bounds, an intermediate scale $\lesssim 10^8$ GeV, narrowing with the reported constants and the representation-side point values to a ceiling of $\sim 4 \times 10^7$ GeV and a window of one and a half to four and a half decades — illustrative figures carrying the provenance of that calculation ([SM §3.1]). What remains: production-scale statistics, the $\ell = 4$ alignment of the observer-position boundary anisotropy, and the cross-charged phenomenology floor. See [SM §3.1] for the full statements.

Standard status. The Lorentz fine-tuning problem familiar from effective-field-theory treatments of Planck-scale Lorentz violation: tree-level suppression of Lorentz-violating operators need not survive radiative corrections.

Framework status. Open. The sector-*differential* (inter-species) boost-

violation is addressed in two steps: at the matching scale the signed-permutation symmetry of the carrier components makes the quadratic response custodially scalar under H-blind ([SM §4.4], Proposition 4b), and below it the custodial-U(6) mechanism gives a power-law-with-logarithm decoupling in the heavy cross-charged mass whose form was reported by a one-loop calculation that is not independently verified (SM §3.1 (*Scope of the emergent-Lorentz claim*), F13). The remaining gap is the *sector-common* operator in the rotational sector, which has no analogous power-law suppression and depends on the renormalization-group-emergence (operator-smearing) mechanism, whose inter-species flow is only logarithmically fast. This rotational sub-problem does not reduce to §19.3.6 (it is a technical radiative-stability question rather than a question about the residue's content). Closing it requires establishing that the emergent theory preserves the octahedral symmetry at loop level: cubic symmetry and locality place the rotational violation at dimension six (SM §3.1, *Scope of the emergent-Lorentz claim* point (iii) and *A present-day constraint, and the rotational sector's distinct status*), so the residual question is loop-level octahedral invariance rather than ultraviolet smearing.

A framing note on the gauge hierarchy. Several of the items above can be stated in terms of the substratum gauge group $\mathcal{G}_{\text{sub}}$ — the transformations preserving all observables of (S, φ). On this description each physical quantity falls into one of three classes: $\mathcal{G}_{\text{sub}}$-gauge and unobservable (quotiented away); observable but solution-specific (the absolute mass scale, the CP phases, the neutrino Yukawas — properties of the particular φ); and forced structural residue (the discrete quantities of §19.3.6). The claim that quantum mechanics, gravity, and the Standard Model are three projections of the one equivalence class $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$ is the framework's statement of scope; the question of §19.3.6 is whether the third class has content independent of the inputs.

A structural observation: several open problems appear to share one premise. Three of the framework's open technical questions, which present as independent entries on the problem list, appear on analysis to reduce to a single underlying question — *whether the substratum-to-observer coarse-graining map preserves the relevant low-energy structure (locality, isotropy, the spectral mode count).* The three are: (a) the radiative stability of emergent Lorentz invariance in the rotational sector (§19.3.7 above), which turns on whether coarse-graining acts as a sufficient operator-smearing; (b) the magnitude of the running-vacuum coefficient ν ([GR §7.1]), which is fixed at $|\nu| \approx 10^{-32}$ (observationally null) if the boundary-mode spectral measure is the local 2D field-theoretic one, and would rise to the DESI-detectable $\sim 10^{-4}$ level only if coarse-graining generated a scale-invariant (non-local) boundary spectrum; and (c) the stability of the $K = 2d$ minimal-coupling count on general graphs ([SM §4.5], Theorem 6 extension), which holds for the coarse-grained dynamics provided local degree fluctuations are renormalization-irrelevant — again a statement that coarse-graining preserves the low-energy mode structure. If this common reduction is correct, the framework has fewer independent open questions than the problem list suggests, and "does the coarse-graining map preserve locality?" is the single

most consequential open technical question — its resolution would discharge all three at once, and conversely a failure (emergent non-locality) would show up simultaneously as rotational Lorentz violation, a non-null ν, and a breakdown of the general-graph mode count. The shared premise has now been computed for the linear (reconstruction-selected) map: an exact surface-Green's-function extraction of the coarse-graining map's boundary spectral measure — the half-space trace-out of the substratum wave equation itself, not a Gaussian effective model — yields a power-suppressed local exponent ($dN/d\omega \propto \omega^4$, with the effective one-mode-per-cell 2D measure, $\propto \omega$, an upper bound on the infrared weight), invariant over thirty orders of magnitude in the slow-bath ratio and under generic quenched deformation, with the log-uniform measure appearing nowhere. The three reductions are thereby conditionally discharged at that grade — locality preserved, ν on the null branch a fortiori, the $K = 2d$ mode-count premise stable — with one named residual: the computation is of the linear map, and its interacting (condensate-dressed) extension is the remaining step. The linkage retains its organizing value in reverse: a failure at the interacting level would still have to manifest simultaneously as rotational Lorentz violation, a non-null ν, and a mode-count breakdown — one falsifiable correlate, not three independent risks. A further preserved structure has since joined the same pattern: the even/odd sublattice grading underlying the staggered spin construction descends exactly through the linear trace-out — an exact Schur-complement identity valid for any site partition ([SM §4.8]) — with the same interacting extension as its residual, further concentrating the value of that single remaining computation.

Why these particular features are inputs: constitution versus consequence. A natural question is why the open and input items cluster as they do — why, for instance, the framework derives (conditionally on H-link) the gauge group and the generation count but must take the spatial dimension, the cubic lattice, and (per §19.3.7) rotational isotropy as inputs. The organizing principle is the distinction between the *constitution* of the substratum and its *consequences*. The framework derives the consequences of a given constitution — the anomaly-free hypercharges and the Cabibbo angle follow by representation theory of the cubic group acting on the link directions, and the gauge group and the three generations follow once H-link identifies those links as the physical carrier, together with the trace-out, *once the substratum is fixed*. It does not derive the constitution itself — the dimension, the Bravais type, the rotational isotropy, the periodicity. The reason is structural rather than incidental: an embedded observer is built *out of* the substratum, so it can have leverage on the consequences of its own constitution but not on that constitution prior to itself. The one qualification is that a consequence which the framework cannot yet *compute* is also booked as an input on tractability grounds rather than principle (the threshold δ_0 of §6.2 is the clearest case; it is a consequence of the coupling-running, input only because it is not yet derived). With that qualification the rule is clean: inputs comprise the substratum's constitution together with any not-yet-tractable consequences, while the derived results are the tractable con-

sequences. This is offered as an organizing observation, not a theorem; it is not claimed that derivability coincides with any single formal property, and indeed the largest block of derived results (the representation-theoretic ones) would follow for any analyst of the fixed lattice, with no reference to an observer at all.

A note on the boost/rotation asymmetry specifically. The split between the two Lorentz sub-sectors of §19.3.7 — the boost sector, for which the framework has structural tools (the symmetric-point custodial scalarity under H-blind, the custodial-U(6) mechanism, the local-boost-invariance prerequisite of the Jacobson construction, the vacuum-Cherenkov stability filter), versus the rotational sector, which has no analogous ally — tracks this same constitution/consequence distinction through a specific feature of embedded observation. The trace-out is realized at a causal horizon, and that horizon *singles out a rest frame* (the frame in which the substratum update steps are simultaneous, identified in §7 with the cosmic rest frame). The observation map therefore makes contact with boost structure — the horizon partition defining it distinguishes a temporal direction, which is what boosts act on — and this is why the boost sector admits framework-level arguments. It does not single out any spatial direction; spatial isotropy is a property of the lattice constitution, untouched by the rest-frame selection. This is formalized as the rest-frame-selection lemma of [Main §3.5] (given that the observation map determines its foliation, its invariances exclude boosts and, among orthochronous transformations, lie within the little group $O(3)$ of its selected timelike direction; which rotations are invariances depends on the substratum dynamics and the lattice geometry of the partition, not on the rest-frame selection). On this reading the relative tractability of the boost sector and the input status of the rotational sector are not separate facts but two faces of the rest-frame-selecting character of embedded observation: the rest-frame selection touches what distinguishes time from space (boosts), and is silent on what distinguishes one spatial direction from another (rotations). This is an interpretive observation about why the open-problem boundary falls where it does, not an additional technical result.

The rotational sector as a structural question about observational incompleteness. The boost/rotation note above and the smearing question of §19.3.7 can be drawn together into a single statement about *why* the rotational sector is the one that resists protection — and the statement is a structural feature of embedded observation itself, not an incidental gap. The protection the rotational sector would need is a sufficient operator-smearing: a coarse-graining whose kernel damps ultraviolet spatial modes (so the radiatively-generated anisotropy stays suppressed). The framework's coarse-graining is the trace-out, and the trace-out *is* the observational incompleteness — the observer's ignorance of the hidden sector. The crucial point is *what kind* of incompleteness this is. By the fluctuation discriminant of [Main §3.5], the trace-out averages over the hidden *state* — over which microstate the slow bath occupies — not over spatial scale. A state-average is not a spatial ultraviolet filter: averaging over which microstate the bath is in does nothing to remove short-wavelength

spatial modes, since those are present in every microstate alike. The framework's incompleteness is moreover enormous — the slow-bath capacity ratio $\tau_B/\tau_S \sim 10^{32}$ is the engine of the entire quantum-emergence result — but it is incompleteness *about a temporally-evolving state*, concentrated in the state and (through the slow bath) temporal directions. It is silent in the spatial direction in the same way, and for the same reason, that the rest-frame selection of the lemma is silent on spatial axes.

This can be made quantitative. The smearing kernel is the traced-out hidden-sector propagator G_{oo} ([SM §4.8]); its position-space width is the smearing radius R. A direct computation of that width on the cubic lattice gives $R \approx \varepsilon$ — a single lattice spacing — and, decisively, that width is *flat* as the slow-bath ratio τ_B/τ_S is varied across thirty orders of magnitude. The framework's vast observational incompleteness contributes *nothing* to the spatial smearing width; the only spatial smearing present is the order-one multi-site support of the staggered reconstruction, which is far too narrow ($R/\varepsilon = \mathcal{O}(1)$, not $\gg 1$) to damp the ultraviolet. This is the structural reason the smearing route does not add ultraviolet damping to the rotational sector — and it is why the available computations (static, free-dynamical, and one-loop) find $R/\varepsilon = \mathcal{O}(1)$ rather than $R \gg \varepsilon$: the incompleteness is the wrong *kind* to UV-damp, regardless of its magnitude. Whether the absence of smearing leaves the sector *unprotected*, however, depends on the dimension analysis below.

Three of the framework's findings thus share one root. The rest-frame-selection lemma (the horizon partition defining the map selects a timelike direction, no spatial one, and the map excludes boosts when it determines the horizon foliation), the fluctuation discriminant (the map averages the fluctuating state, not the fixed orientation), and the smearing-width result (the incompleteness has order-one spatial width independent of its magnitude) are three expressions of a single fact: *observational incompleteness in this framework is incompleteness about a temporally-evolving hidden state, with purchase on time, boost, and state, and structurally silent on space and rotation.* The rotational sector is precisely the sector embedded observation cannot reach. The valence of this account is narrower than it first appears. The smearing-width result it establishes — that observational incompleteness adds no ultraviolet spatial smearing, so $R \approx \varepsilon$ — bears on the *coefficient* of the rotational operator, not its *dimension.* The dimension is fixed independently, by the octahedral selection rule (the E_g anisotropy is a cubic non-singlet; the lowest invariant anisotropy is $\ell = 4$, dimension six) together with locality, and a direct tree-level computation confirms the emergent dispersion anisotropy is $\sim (\varepsilon k)^2 \propto (E/M_{\mathrm{Pl}})^2$. The account therefore explains why embedded observation does not *smear* the rotational sector, but smearing is not what the sector needs: cubic symmetry and locality already place the violation at dimension six, Planck-suppressed. What remains genuinely open is the loop-level question — whether the emergent theory preserves the octahedral symmetry *exactly*, or whether the emergence introduces an octahedral-breaking effect not visible in the bare partition.

This premise has a second dependency alongside the partition. The effective theory is octahedrally invariant only if the condensate that breaks the framework's custodial symmetry is itself aligned to the cubic axes; the invariance of the dynamics does not descend to the solution, since the gauge group arises precisely by that condensate breaking a symmetry the dynamics has. The alignment is not secured by the observed gauge group alone — a condensate with the same block sizes but misaligned eigenspaces yields the same gauge group without being octahedrally equivariant, so the equivariance argument that forbids the anisotropy below dimension six does not cover it — and it is not derived: it is the hypothesis that the matter-content and mixing-angle results also take, namely H-link with its single-copy clause, the open hypothesis that the physical carrier is one copy of the six-link representation (Chapter 5 §5.4), together with an octahedrally equivariant condensate, whose blocks are then the link-direction irreps (Chapter 5 §5.5). An aligned condensate leaves the forbidden anisotropy exactly null at every block splitting, and a staggered spatial modulation of the condensate does not lift it. See [SM §3.1]. This has an identifiable failure point: it assumes the visible/hidden partition acts as a state/sublattice trace that preserves the octahedral symmetry, rather than an effective short-wavelength (momentum-shell) cut that could both ultraviolet-damp and, if anisotropic, break it. The gaplessness of G_{oo} is evidence the partition is not a clean momentum shell; whether it is exactly octahedral-invariant at loop level is the seam at which the dimension-six protection could fail, and is the arbiter — not the kernel width. This premise is grounded in the framework's own factorization criterion ([Main §3.1, §3.5]): an observer-centered, horizon-isotropic partition is spherical and hence octahedrally symmetric about the observer's centre, so the lattice octahedral group — unlike translations, boosts (when the map determines the horizon foliation), and rotations about distant points, which the partition breaks — preserves the visible/hidden factorization and descends to an exact symmetry of the effective theory. With the partition exactly octahedral-invariant and the condensate aligned (H-link with an equivariant condensate), the same criterion that selects the gauge symmetries protects the rotational sector and exposes the boost sector, so the dimension-six rotational protection and the dimension-four boost exposure are two faces of one structural fact rather than independent assumptions.

19.3.8 The chirality hypothesis H-χ'

The Standard-Model identification carries one named structural hypothesis. **H-χ':** a *non-minimal* mechanism renders the effective SU(2) coupling taste-chirality-selective. The minimal question is settled: the link-current embeddings the construction delivers are certified taste-blind for all three gauge blocks — SU(3), SU(2), U(1) alike — by exact spin⊗taste classification (`hchi_selectivity_probes.py`), so the minimal SU(2) is vector-like and any selectivity must enter through a taste-asymmetric structure (condensate/mass sector) in the Schur-complement effective theory. What is proved, and probe-certified (`chirality_grading_probes.py`): on the visible sector the

spin-chirality and taste-chirality operators coincide, so a gauge coupling is spin-chiral there if and only if its embedding is taste-chirality-selective (SM §4.8, Theorem 13); the grading itself is $(d_L, d_R) = (4, 4)$ with $\Gamma(0) = I_4$ chirality-neutral, refuting the earlier "even sublattice = left-handed" identification; SU(3) is vector-like; and the generation count does not depend on H-χ'. What is open is H-χ' itself: whether the substratum's SU(2) embedding is in fact taste-chirality-selective. The resolution path is a computation in the framework's own representation machinery — the embedding's action on the graded components of the coupling matrix (the Theorem 7 apparatus). A computed non-selective embedding falsifies the chirality identification, and with it the specific Standard-Model matter assignment, while leaving the gauge group, the multiplicities $(3, 2, 1)$, and the generation count intact; a computed selective embedding discharges the hypothesis and upgrades the chirality clause corpus-wide. The condition's standing status is transcribed in the status ledger (Main §4.5).

19.3.9 Selection of the relative quantum evolution

Framework-specific. This is not one of the standard open problems; it is a frontier internal to the OI–QM bridge, and it is recorded here so that the bridge's settled and unsettled parts are not read as one thing.

What is settled. The finite-law bridge is solved at the level of exact representability: every finite-horizon visible law has a finite reversible realization and a fixed-basis unitary/Born representation, and the three descriptions are equivalent ($S \iff D \iff Q_{\mathrm{fb}}$, [Main §3.4]). Separately, exact finite operational quantum mechanics is characterized *conditionally*: within complex matrix kinematics taken as given, it holds exactly when the explicit completion principles hold, none of which bare embedded observation supplies ([GR §3.3], §19.2.12).

What is open. Between those two lies a question neither answers. A visible OI family does not come with a single unitary evolution relating one time to the next: it admits coherent lifts, and the lifts need not agree on that relating evolution. So the open question is whether, and by what, one relative quantum evolution is selected from the coherent lifts compatible with a single visible family. The question is two-part: what selects or constrains, through time, the fibre-Gram data of the lift — for each visible output, the overlaps among the hidden states into which the lift sends the visible inputs — and what fixes the relating evolution within a given trajectory of that data.

What is proved about it. Ordinary coherence is not enough. Two coherent lifts of the same visible family can be related by a time-dependent deformation that preserves everything the visible law sees, while their relating evolutions differ; a deformation constant in time cancels from every relating evolution, so the nonuniqueness is carried by time dependence specifically ([Main §3.4]). The invisible deformations are two-sided — a right action by the anchored class and a left action within each visible output's hidden fibre, each maximal among

uniform actions of its kind — and at one time the residue after both is classified exactly: two lifts are related by the two-sided action exactly when their fibre-Gram data agree up to phases, the visible law is the diagonal of that data, and every positive semidefinite, rank-bounded, identity-summing family with the prescribed diagonal occurs. So the freedom the visible law leaves at one time is exactly the off-diagonal fibre-Gram data modulo phases. At one time that freedom does not reach the fixed-basis layer. With a trivial ancilla, the fixed-basis datum built from a lift's unitary, with uniform initial law and identity readout, has finite-horizon trajectory laws fixed by the visible law alone, also after contractively scaled partial permutations on either side, and two such data have equal laws exactly when their visible laws are equal ([Main §3.4]). The trivial ancilla is part of the statement: an ancilla carried between steps as hidden basis states can separate two lifts of one visible law already in the two-step law, while an ancilla attached as a tensor factor with its own unitary cannot.

What is sharper, and what it does not settle. A pair of lifts of one visible family outside the maximal uniform right class and differing in relating evolution exists; every such pair exhibited is related by a left in-fibre move, so the right class alone is not the invisible freedom and quotienting by it alone does not determine the relating evolution. A pair outside the two-sided action and differing in relating evolution exists as well, on a single visible law. The complete trajectory of fibre-Gram data does not fix the relating evolution either: a time-dependent invisible right deformation preserves every fibre-Gram matrix at every time and changes the relating evolution. So the two-part question above is open in both parts.

What is not claimed. None of this makes embedded observation and quantum mechanics inequivalent, and none of it narrows $S \iff D \iff Q_{\mathrm{fb}}$. No selection mechanism is endorsed here: the framework does not claim the missing structure must be more than a gauge fixing, does not claim a connection or gauge choice could not supply it, and does not claim that no such structure exists. The fibre-Gram data is a coordinate on the space of lifts, not a physical quantity, and no statement is made about which trajectory of it is selected.

Prospects. The question is sharply posed at one time and the open part is cross-time: the data left free at each time is known exactly, so progress would come from a statement about what constrains that data across times, and about what fixes the relating evolution within a trajectory of it.

Developed in: Chapter 1 (the operational boundary), with the representability side in Chapter 1 §1.5 and the conditional completion side in §19.2.12.

19.4 Chapter close: summary table and what the framework does not address

The framework's content against the standard list of open problems in fundamental physics summarizes as follows. The status labels are the framework's claims as argued in the cited chapters, and every label inherits the conditionality ledger of [Main §4.5] for the foundational claims — in particular the slow-bath C2 leg's mixing-hypothesis conditionality — together with the observer/gauge-carrier flags carried at their sources: H-link single-copy and H-cust [SM §§4.5–4.6], the open chirality hypothesis H-χ' [SM §4.8], the physical-spin identification H-spin' [SM §4.7] (its free-kernel form and the four-dimensional staggered route are settled negative — `hspin_kernel_probes.py`, `hspin4d_probes.py`; the condensate-dressed form is the open), and the flagged carrier input of the Casimir chain [SM §4.7]; no label asserts more than its cited derivation, and "Dissolved" or "Derived" is always relative to that ledger.

Summary table.

Problem	Standard status	Framework status	Developed in
Quantum gravity	Central problem	Partially addressed — covariant unification open (G3)	Chs 1, 7, 18
SU(2) chirality	Assumed in the SM	Open — named hypothesis H-χ'; grading proved, selectivity open	§19.3.8, [SM §4.8]
Taste → physical spin	Spin postulated per field	Algebraic label content proved and probe-certified (Thm 9); the physical identification is the named hypothesis H-spin' (resolution: the small-momentum Dirac-form test); the carrier input separately flagged, A^2-tested	[SM §4.7]
Cosmological constant	Worst prediction in physics	Descriptive distinction established; gravitational form conditional (G3)	Ch 7 §7.7
Dark matter	Unknown, no particle found	Conditional ($a_0 = cH/6$)	Ch 7 §7.9

Problem	Standard status	Framework status	Developed in
Dark energy	Unknown, evolving disputed	Smooth $H^2 - H_0^2$ bookkeeping supplied; Type II reading, not a derived law	Ch 7 §7.8
Measurement problem	No interpretive consensus	Observer-relative account at the fixed-basis level; operational completion and Frauchiger-Renner premises open	Ch 1, Ch 18 §18.2
BH information paradox	Hawking vs unitarity	Not resolved — information conserved by premise; Page curve conditional	Ch 7 §7.6
Strong CP	Why $\bar{\theta} < 10^{-10}$?	Not resolved (T-invariance gives $\theta \in \{0, \pi\}$ and reciprocal transition counts, not $\bar{\theta} = 0$; open under H-top and H-det)	Ch 6 §6.3, [SM §5]
Hierarchy problem	Higgs mass protection unknown	Partially addressed: cutoff insensitivity (SM §8.8); $v \ll f$ open	Ch 6 §6.6
Gauge group origin	$\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ unexplained	**Conditional** — exact six-link $3 \oplus 2 \oplus 1$ decomposition; physical-carrier identification H-link single-copy and custodial route H-cust remain open	Chs 5, 6, [SM §§4.5–4.6]
Generation puzzle	Three, with hierarchical masses	Conditional branch: three candidate tastes on single-copy H-link; physical-generation reading additionally under H-spin'; hierarchy pattern open	Ch 6 §§6.5-6.6

Problem	Standard status	Framework status	Developed in
Dark sector budget	95% unexplained	Conditional (partly definitional)	Ch 7 §7.9
Born rule origin	Independent postulate	Split: representation level settled — admitted universally ($S \iff D \iff Q_{\text{fb}}$); coherent operational extension characterized by five named conditions; read-write equilibrium a mechanism proposal	Ch 1, Ch 18 §18.2, [Main §3.4]
Relative-evolution selection (framework-specific)	Not a standard problem	Open — representability settled and operational completion characterized conditionally; the freedom the visible law leaves at one time is classified exactly by the fibre-Gram data, but no theorem selects that data's trajectory or the relating evolution within it; ordinary coherence proved insufficient	§19.3.9, Ch 1, [Main §3.4]
Flavor problem	**13 free parameters**	**Open (research program; m_b/m_τ at 0.4% match with Layer 2(a) verification residual)**	§19.3.1
Hubble tension	**$> 5\sigma$ disagreement**	**Open (ν_{OI} too small)**	§19.3.2
Baryogenesis	**Sakharov conditions**	**Solution-specific (η_B inherits flavor-CP input status)**	§19.3.3

Problem	Standard status	Framework status	Developed in
Inflation	**Inflaton not derived**	**Open (three possibilities, none developed)**	§19.3.4
Initial state	**Framework-specific**	**Partially gauge, partially irreducible**	§19.3.5

The empirical record summarized above is mixed. Several problems are presented as resolved or dissolved at their stated scopes, while the Born-rule completion, strong CP, flavor, cosmological and matter-sector questions remain split or open. **Gauge-group origin is no longer counted as a completed derivation:** the six-link $(3, 2, 1)$ representation is exact, but its physical-carrier use is conditional on H-link single-copy and H-cust. The table, rather than a single resolved/open count, is the governing status ledger.

Problems the framework does not address.

Beyond the resolved and open problems above, the framework does not address several other categories of question.

Specific experimental anomalies in the Standard Model precision sector are not modified by the framework. The muon anomalous magnetic moment $g - 2$ shows a $\sim 5\sigma$ discrepancy between Fermilab measurement and the SM prediction using dispersive methods (though lattice QCD has reduced the tension). The B-physics lepton flavor universality violations (R_K, R_{K^*}) were reported by LHCb but recent measurements are consistent with the SM. The CDF II measurement of $m_W = 80.4335 \pm 0.0094$ GeV is approximately 7σ above the SM prediction, while other measurements (ATLAS, LHCb) are consistent. The framework derives the SM structure but does not modify the calculations producing these observables. Any framework correction at accessible energies is at $\mathcal{O}(\epsilon^2 E^2) \sim 10^{-34}$ for $E \sim 100$ GeV at one nat of entropy per boundary mode, unobservable. If any of these anomalies is confirmed at $> 5\sigma$ with consistent measurements, it would require either new physics beyond the framework's SM derivation or a solution-specific effect from the particular φ.

Condensed matter and many-body physics. The framework derives the fundamental laws but does not address emergent phenomena in condensed matter: high-T_c superconductivity, the fractional quantum Hall effect, topological phases, and others. These are many-body problems within the derived QFT, not problems at the framework level. The restricted quantum-statistical regime predicted by the mechanism proposal (Chapter 18 §18.3) may be relevant to some condensed matter systems, but the connection is speculative and has not been developed.

The hard problem of consciousness. Developed in detail in Chapter 18 §18.10. The framework's mathematical operation of observation is a necessary structural condition for consciousness but not a sufficient condition. The hard problem

proper — the explanatory gap between structural and phenomenal facts — is explicitly outside the framework's scope. The framework provides necessary-condition content with substantive constraints on physics-based consciousness theories, but it does not bridge the explanatory gap between physical descriptions and phenomenal experience.

Mathematics and logical foundations. The framework presupposes standard mathematics (ZFC set theory, the integers, finite combinatorics) without addressing the foundations of mathematics. Whether the framework's commitment to finiteness has implications for mathematical foundations (constructivism, finitism, ultrafinitism) is not addressed.

The framework's epistemic position. The cumulative record across Chapters 5 through 18 is substantial: twenty-two classified Standard Model retrodictions (two parameter-free structural and five conditional structural), seven gravitational-sector entries, seven neutrino-sector entries, and a framework-level treatment of the quantum gravity problem in three components, with covariant unification open (G3). The record is retrodictive: the structural values match existing data at the precisions classified in Appendix A, and the pre-registered forward set (Appendix A.6) awaits its data.

The open problems are also substantial. The flavor problem is the framework's largest single open theoretical task. The Hubble tension is not addressed by the framework's ν_{OI} prediction. Baryogenesis is solution-specific — the baryon asymmetry inherits the input status of the framework's flavor-sector CP phases. Inflation is not derived. The initial-state question is partially gauge and partially the framework's irreducible mystery.

Shared descriptions and separate obligations. Several entries above use the substrate/emergent split and trace-out, but their conclusions retain their separate hypotheses and empirical classifications. For the no-go cases in particular, Bell requires ontic parameter dependence under measurement independence and an open H-Bell realization; Kochen-Specker concerns the noncommuting operational algebra; PBR concerns preparation independence; Frauchiger-Renner concerns the joint use of observer-relative predictions. None is an automatic dissolution supplied by the shared architecture, and a resolution of one does not prove the others.

A common construction can impose constraints on several questions when the mathematical links and their assumptions are given. Such links must be established separately: sharing a substrate/emergent description does not turn conditional arguments into independent resolutions or show that changing one proposed physical mechanism changes every other conclusion.

The inventory above therefore states the status of each problem separately. Its empirical entries and its open physical bridges remain distinct from the no-go premise checks. The shared representation is a way to compare the arguments, not evidence that the architecture alone resolves the entire list.

The chapter's purpose is to make the framework's record on standard open problems explicit. The framework's content is neither a complete theory of fundamental physics (the open problems above remain genuinely open) nor an unfocused collection of predictions; it is a structural framework that resolves a substantial fraction of the recognized hard problems of physics through a small set of structural commitments, with the resolutions cross-referenced to specific empirical content developed in earlier chapters. The remaining open problems indicate where the framework's reach stops at current development, with concrete prospects for closing several of them through further calculation within the framework's existing structural commitments.

The book's content concludes here. The framework's empirical record (Chapters 5-17), forward predictions (Chapter 18), and inventory against the standard open problems (Chapter 19) together constitute the framework's positive content. The remaining scope limits — the flavor problem, the Hubble tension, baryogenesis (solution-specific: η_B inherits the flavor-sector CP input), inflation, and the initial state — are pre-registered as the framework's current boundaries. Closing them is the framework's future research program.

Appendix A

Prediction Status Table

A.1 What this appendix provides

Twenty-two Standard Model observables, seven gravitational-sector entries, seven neutrino-sector entries, and the biological retrodictions form an inventory of measured comparisons, conditional formulas, observational bounds and predictions awaiting tests. This appendix gives each entry's structural classification, empirical standing and remaining assumptions.

The appendix serves two purposes. First, as a *navigational index*: readers wanting to know "what does the framework predict for X" can find each prediction with its location in the book, its formula, its empirical match, and its current status. Second, as an *honest accounting*: predictions that are confirmed appear alongside predictions that are layered (requiring additional verification), parameter-fit (where the framework provides structural form but specific values are empirical), and forward-looking (testable but not yet tested). The reader can therefore assess the framework's empirical record at the level of detail appropriate to their domain of interest.

Section structure.

Section A.2 develops the classification scheme — the categories used throughout the appendix to characterize each prediction's structural status. The scheme distinguishes structural (S), conditional (C), layered (L), retrodiction (R), parameter (P), mass-chain (M), and empirical (E) predictions, with each category having specific evidential force.

Section A.3 develops the four-layer descriptive framing — substratum (Layer 0), substratum-level structural (Layer 1), substratum-level mechanism-specific (Layer 2(a)), substratum-level cubic-symmetry-specific (Layer 2(b)), mixed-level (Layer 3) — that organizes how the framework's content propagates from foundational commitments through specific empirical predictions.

Section A.4 provides the master prediction tables, organized by domain: Standard Model observables (A.4.1), gravitational-sector observables (A.4.2), neutrino-sector observables (A.4.3), and biology/medicine predictions (A.4.4). Each entry includes prediction number, formula or quantitative content, structural classification, layer, empirical match, and chapter cross-reference.

Section A.5 provides forward predictions and falsification targets — predictions the framework makes that have not yet been tested but that provide specific empirical exposure for future research programs.

Section A.6 closes with a discussion of the framework's epistemic position based on the cumulative empirical record: which predictions provide the strongest evidence, which are most distinctive of the framework relative to standard accounts, and which open issues are most consequential for the framework's overall content.

A.2 Classification scheme

The framework classifies each prediction by its structural status, using a seven-category scheme designed to make the *epistemic content* of each prediction explicit.

S — Structural prediction. The framework's structural commitments fix the prediction's value with no fitted input and no open link specific to the entry; matter-sector entries rest, as every matter-sector result does, on the conditional physical-carrier chain (H-spin' and H-χ'). The prediction's value emerges from the framework's structural foundations without empirical input beyond what is required to identify which physical system the framework describes. Examples: the Wolfenstein $A = \sqrt{2/3}$, the Higgs quartic boundary condition $\lambda(M_{\rm Pl}) = 0$. S-class predictions provide the framework's strongest evidence — they are derivations from structural commitments with no open link of their own.

C — Conditional prediction. The result follows given an additional unverified relation. Examples include the $A \cdot B$ extrapolation conditional on its finite-size leading power, the Cabibbo angle $\lambda = 1/(\pi\sqrt{2})$ conditional on the emergent dressing of its identification, and Koide $Q = 2/3$ conditional on the dynamical closure of its amplitude. Verification of the relation can establish

the corresponding unconditional result; observational agreement alone does not verify that relation.

L — Layered prediction. The framework's content includes the prediction's *form* at the structural level but requires an empirical or computational input for the specific value. The empirical input is bounded — typically a single dimensionless coefficient or a normalization scale — but cannot be derived without additional work. Examples: $\sin^2\theta_{12}$, $\sin^2\theta_{23}$, m_b/m_τ. L-class predictions provide *structural understanding* of the prediction's origin with empirical input for specific values.

R — Retrodiction. The framework reproduces an existing empirical value through a structural mechanism, but the empirical value was input to the framework's identification of which specific system it describes. The prediction is retrospective rather than forward-looking. Examples: $\alpha_2(M_Z)$, $\alpha_3(M_Z)$. R-class predictions provide *structural consistency* with established physics but not novel predictive content.

P — Parameter prediction. The framework determines a parameter through a calculation that involves both structural commitments and one or more empirical inputs. Different from L (where the form is structural but the value requires input) and from R (where the prediction is purely retrodictive), P-class predictions combine structure and empirics in derivation.

M — Mass-chain prediction. The framework predicts mass relationships within a chain (e.g., m_e, m_μ via Koide from m_τ) where one mass is empirical input and the others follow structurally. Mass-chain predictions provide *structural relationships* between empirical inputs.

E — Empirical input. A value that the framework treats as empirical rather than deriving — typically a coupling constant, mixing angle, or other parameter that is part of the framework's identification of which specific physical system applies. E-class entries are *not predictions* in the strong sense but parameters the framework uses without deriving.

The cumulative content. The framework's strongest evidence comes from S-class predictions (structural derivations with no open link of their own, matching empirical values), supplemented by L-class predictions (structural form with bounded empirical input). The R, P, M, and E categories serve as honest accounting of where empirical input enters the framework's content. The classification scheme makes the framework's evidential content explicit at the prediction level, allowing the reader to assess which predictions provide the strongest support for the framework's structural commitments.

A.3 Four-layer descriptive framing

The framework's predictions are organized at four levels of descriptive abstraction, with each level having specific evidential content and specific dependencies on the framework's structural commitments.

Layer 0 — Substratum. Predictions at the substratum level: properties of the underlying bijection φ on S, the substratum's dimensionality, the substratum's discreteness scale. Examples: dimensionality $d = 3$, finite cardinality, deterministic bijection structure. These predictions have *foundational* evidential weight — they are the framework's substratum-level commitments themselves.

Layer 1 — Substratum-level structural. Predictions derived from the substratum's structural properties through dimensional analysis, group theory, or other framework-level structural mechanisms. Examples: gauge group $SU(3) \times SU(2) \times U(1)$ from substratum dimensionality (Chapter 5) and generation count from the matter content theorem (Chapter 6), both conditional on H-link with its single-copy clause, which supplies $K = 2d = 6$ (Chapter 5 §5.4), the gauge group also on an octahedrally equivariant condensate (Chapter 5 §5.5), Cabibbo angle $\lambda = 1/(\pi\sqrt{2})$ from cubic-group symmetry. Layer 1 predictions are *structural derivations* from foundational commitments.

Layer 2(a) — Substratum-level mechanism-specific. Predictions that require specific structural mechanism beyond Layer 1 generic structural content. Examples: m_b/m_τ requires the two-loop SχPT mechanism specialized to OI's 3D-slice cubic-group taste structure. Layer 2(a) predictions are *mechanism-specific* — they require more detailed structural content than Layer 1.

Layer 2(b) — Substratum-level cubic-symmetry-specific. Predictions that depend on the specific cubic-symmetric substratum the framework commits to. Examples: the Jarlskog $J = \eta/(12\pi^6)$ requires specific cubic-symmetric coefficient. Layer 2(b) predictions are *symmetry-specific* — they require commitment to cubic symmetry rather than just the foundational substratum structure.

Layer 3 — Mixed-level. Predictions that mix substratum-level structural content with emergent-level computational content. Examples: m_e, m_μ via Koide from m_τ requires both substratum-level Koide relation (Layer 1) and emergent-level mass chain. Layer 3 predictions are *mixed* — they combine substratum and emergent content.

Cumulative implications. The framework's content is most evidentially strong at Layer 0 and Layer 1 (where substratum commitments directly determine the prediction's form and content). It is less strong at Layer 2(a) and Layer 2(b) (where additional structural content is required). It is least strong at Layer 3 (where empirical-emergent content mixes with structural content). The classification scheme makes these distinctions explicit, allowing the reader to assess the framework's evidential weight at the appropriate level.

A.4 Master prediction tables

This section provides the master tables of the framework's predictions, organized by domain. Each entry includes prediction number, formula or quantitative content, structural classification, layer, empirical match, and chapter cross-reference for further detail.

A.4.1 Standard Model observables

The reference table combines Standard Model observables, coupling-sector inputs and gravitational entries. Its empirical column distinguishes measured comparisons from inconclusive calculations, observational bounds and untested formulas.

#	Prediction	Formula	Class	Layer	Empirical match	Chapter
1	$1/\alpha_0$ universal induced coupling	23.25	S	0	Structural input; no direct comparison	Ch 6 §6.4
2	$A \cdot B$ gauge coupling product	$\approx$ 48–62 (leading-power dependent)	C	0	Inconclusive	Ch 6 §6.4
3	δ_0 from $U(1)$ row	10.02	E	0	Empirical input	Ch 6 §6.4
4	$\alpha_2(M_Z), \alpha_3(M_Z)$	29.57, 8.47	R	0	Retrodictions	Ch 6 §6.4
5	Cabibbo λ	$1/(\pi\sqrt{2})$	C	1	0.04% (global fit; retrodiction)	Ch 6 §6.5
6	Wolfenstein A	$\sqrt{2/3}$	S	1	1.2% (0.8σ)	Ch 6 §6.5
7	$\lvert V_{cb}\rvert$	$A\lambda^2$	C	1	Conditional on 5,6	Ch 6 §6.5
8	m_d/m_s	λ^2	L	1	Structural λ; GST empirical	Ch 6 §6.5
9	Jarlskog J	$\eta/(12\pi^6)$	E	2(b)	Empirical η	Ch 6 §6.5
10	Koide Q	2/3	C	1	**<0.001%**	Ch 6 §6.6
11	Koide θ_0	$C_2/d^2 = 2/9$	C	1	0.001%	Ch 6 §6.6
12	$\sin^2\theta_{12}$	$1/3 - 1/(4\pi^2)$	L	1+2(b)[f]	**0.07σ JUNO**	Ch 8
13	$\sin^2\theta_{13}$	$4/(18\pi^2)$	C	1	2.6% (1.0σ)	Ch 8
14	$\sin^2\theta_{23}$	$1/2 + 1/(2\pi^2)$	L	1+2(b)[f]	Within 1σ	Ch 8
15	m_e, m_μ via Koide	from m_τ	M	3	Mass-chain inheritance	Ch 6 §6.6
16	m_b/m_τ	$4.28/Z_S(\lambda g_0^2) = 2.361$	I	2(a)	0.4% (0.5σ at L=32)	Ch 6 §6.6

#	Prediction	Formula	Class	Layer	Empirical match	Chapter
17	m_u/m_d	$\sqrt{2/9}$	L	1+2(b)	Within 1σ	Ch 6 §6.6
18	Σm_ν	~0.059 eV (oscillation inputs)	L	3	Upper bounds only	Ch 8
19	PMNS sum rule	$2\Delta_{12} + \Delta_{23} = 0$	L	1+2(b)[f]	1.178 ± 0.020 confirmed	Ch 8
20	m_t (weak-scale $y_t \approx 1$)	$v/\sqrt{2}$	R	3	0.9%; mechanism open	Ch 6 §6.6
21	m_b via Q_{down}	Koide(Q_{down}), Q_{down} phenomenological	L	2(b)	0.9% (1.2σ)	Ch 6 §6.6
22	$\lambda(M_{\text{Pl}}) = 0$	structural	S	1	**0.6**σ	Ch 6 §6.7
23	MOND acceleration a_0	$cH/6$	C	0/1	Within quoted systematics	Ch 7 §7.9

Summary statistics. Counts: 3 S + 7 C + 8 L + 1 M + 2 R + 2 E. Total: 23. This reference index is broader than the paper's Standard-Model table ([SM §7.6], 22 rows): it also carries coupling-sector cross-checks, neutrino entries and the gravitational result a_0. The headline count of *two* parameter-free structural and *five* conditional structural predictions among twenty-two Standard Model observables belongs to that paper's observable inventory, which counts the coupling-sector quantities $1/\alpha_0$ and $A \cdot B$ separately. The conditional entries here are the leading-power-dependent coupling product, the Cabibbo λ with its inheritors $|V_{cb}|$ and $\sin^2\theta_{13}$ (emergent dressing open), the two Koide entries (dynamical closure of the amplitude open), and the MOND formula (coupling map open, factor imported). Empirical precision applies to the individual measured comparisons. The numerical mass sum is L at Layer 3: its near-minimum value combines the hierarchical normal-ordering expectation with measured solar and atmospheric mass-squared splittings, and is constrained by upper bounds.

[f] *Layer 2(b) (sin²θ_12, sin²θ_23, PMNS sum rule).* The Layer 1 component is structurally protected against the A_1 perturbation by the cosines-dressing of the TBM sum rule (cubic geometry alone). The remaining Layer 2 component is the A_2 Wilson-coefficient ratio $b_{23}/(b_{12}+b_{13}) = 4/3 = 4(d-1)/(2d)$, equivalent to the substrate-level relation $b_{23}^{\text{sub}} = (4/3)(b_{12}^{\text{sub}} + b_{13}^{\text{sub}}) + 1/9$. The ratio's 2(a)-tractable period — the label meaning the residual question was sharply specified and admitted definite outcomes from substrate calculations — has ended: the

calculation was performed. It does *not* mean the answer is forced by representation theory alone — SM §7.3 established that the natural Higgs-democratic spurion does not produce this ratio (instead giving $b_{12} : b_{13} : b_{23} \approx 17 : 1 : 1$ in the heavy-quark hierarchy, off by factor ~24 from the required 4/3), and the relation requires substrate amplitudes with the non-natural scaling $\delta M_{ij}^{\text{sub}} \propto m_{\max(i,j)}$ plus a channel-specific enhancement factor $K_{23}/K_{12} \approx 2.8$. The substantive content is bijection-specific — Layer 2(b) — and the sharply specified verification path has since been executed: the §7.3 routing computation excludes the second-order route (the two-hop operator is a pure spin-taste monomial, orthogonal to every mass channel), leaving the recorded $O(\lambda^3)$ construction as the one unpursued analytical route. The empirical match at 0.07σ (JUNO) and 0.6σ (sum rule) confirms the bijection has the required structure; the 7/6 sum rule remains the direct experimental test.

Distinctive empirical content. Three predictions stand out as the framework's strongest empirical content. (i) JUNO $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2) = 0.3080$ at $\mathbf{0.07\sigma}$ against the post-JUNO global fit — a parameter-free retrodiction; its exact, freedom-free form makes JUNO's design-lifetime precision (± 0.0014) a pre-registered forward test. (ii) the Cabibbo angle $\lambda = 1/(\pi\sqrt{2})$ at **0.04%** against the global fit — a retrodiction. (iii) Koide $Q = 2/3$ to better than **0.001%** — the sharpest framework prediction across all 23 in relative precision, matching observation at four-significant-figure precision.

A.4.2 Gravitational-sector observables

The horizon and radiation entries inherit H-slope and the horizon/frame/state conditions (Chapter 7; [GR §§3, 8.5]). The MOND and physical dark-sector interpretation additionally require the open matter-to-boundary coupling map. The Page curve requires H-scramble and H-decouple; H-scramble fails for every bijection acting on an initial configuration with the bipartition aligned to the configuration coordinates, so the Page value requires an initial superposition over many configurations or a non-aligned bipartition; no such realization is established, and the Page curve (G11) is an open research target. Free symbolic dispersion does not establish suppression in the interacting sector. These conditions are separate from the descriptive vacuum-energy distinction and the classical boundary-density normalization.

#	Prediction	Formula	Class	Layer	Empirical match	Chapter
G2	Bekenstein bound	$S_{\text{dS}} \sim 10^{122}$	C	0	Order-of-magnitude estimate	Ch 7 §7.5

#	Prediction	Formula	Class	Layer	Empirical match	Chapter
G5	Hawking radiation	Thermal spectrum; greybody-dependent rate	C	1	Not observed astrophysically	Ch 7 §7.5
G6	MOND acceleration $a_0 = cH/6$	$cH/6$ (factor imported)	C	0/1	Within quoted systematics	Ch 7 §7.9
G8	Running vacuum ν_{OI}	Type II interpretation not derived; conditional emergent-QFT-scale magnitude $\lvert\nu\rvert \approx \lvert\nu_{\mathrm{QFT}}\rvert \sim (M_{\mathrm{SM}}/M_{\mathrm{Pl}})^2 \sim 10^{-32}$, indistinguishable from zero	L	2(a)	DESI Year 5 decisive; Bertini 2025 $\nu = -(2.5 \pm 1.3) \times 10^{-4}$ consistent at 2σ	Ch 7 §7.8
G9	Dark sector budget	$\sim 95\%$	C	1	Partly definitional budget	Ch 7 §7.9

#	Prediction	Formula	Class	Layer	Empirical match	Chapter
G11	Page curve	Nested trace-out, H-scramble (fails from configurations), H-decouple; full product space	Open	1	A non-vacuous version needs a specified initial ensemble or a non-aligned bipartition, a valid proof and independent verification; decoupling and proper-shell value open	Ch 7 §7.6
G12	$\Sigma m_\nu \sim$ 0.059 eV	Measured splittings	L	3	Upper bounds only	Ch 8
G13	Lorentz violation	Free dispersion $O((k\epsilon)^2)$; interactions open	C	0	Interacting protection open	Ch 7 §7.4

Summary statistics. Counts: 5 C + 2 L. Total: 7. G11, an open research target, is not counted. The conditional entries carry the calibration, horizon-state, coupling or interacting-sector obligations stated above. $S_{\rm dS} \sim 10^{122}$ is an order-of-magnitude estimate; the mass sum has upper bounds. These entries do not constitute a common precision test.

Consistency relations. Four gravitational-sector results are consistency or descriptive relations rather than predictions, and the counts above do not include them. The horizon calibration — the discreteness scale $\epsilon = 2\sqrt{\sigma}\, l_p$ and the action scale $\hbar = c^3\epsilon^2/(4G\sigma)$, algebraically one relation — is a self-consistency condition that fixes only $\epsilon/\sqrt{\sigma} = 2\, l_p$, not an independent determination (Ch 7 §7.3–§7.4). The Bekenstein-Hawking coefficient 1/4 is the ratio $2\pi/8\pi$ of the KMS period to the first-law prefactor that the horizon calibration matches, conditional on H-slope and the horizon/frame conditions; it agrees with the normalization fixed by the Hawking temperature and the first law, and has no direct empirical test: GW250114 constrains the classical area theorem, not the coefficient (Ch 7 §7.5). The vacuum-energy level distinction is descriptive, and its compression ratio $\rho_{\rm QFT}/\rho_{\rm classical} \sim S_{\rm dS}$ is a dimensional identity for any Planck-cutoff theory, holding up to a convention-dependent numerical factor, $1/\pi$ for $M_{\rm Pl}^4$ against H^2/G and $1/(384\pi^4\Omega_\Lambda) \approx 3.9 \times 10^{-5}$ for a mode sum cut

off at the reduced Planck mass against the observed dark-energy density (Ch 7 §7.7). The boundary-density equality $\rho_s = \rho_{\rm crit}$ is a normalization identity (Ch 7 §7.9).

Distinctive content. The conditional MOND formula gives $a_0 = 1.09$–$1.18 \times 10^{-10}\,\mathrm{m/s^2}$ across the stated H_0 range, compared with $(1.20 \pm 0.02 \pm 0.24) \times 10^{-10}\,\mathrm{m/s^2}$. It agrees within the quoted systematic uncertainty; the factor 1/6 is imported. The running-vacuum coefficient (G8) provides forward exposure to DESI data.

A.4.3 Neutrino-sector observables

The neutrino sector provides the framework's sharpest parameter-free retrodiction. The JUNO measurement of $\sin^2\theta_{12}$ at 0.07σ against the framework's structural value is a retrodictive match whose exact form converts JUNO's design-lifetime precision into the framework's cleanest pre-registered forward test.

#	Prediction	Formula	Class	Layer	Empirical match	Chapter
N1	$\sin^2\theta_{12}$	$1/3 - 1/(4\pi^2)$	L	1+2(b)	**0.07σ JUNO**	Ch 8
N2	$\sin^2\theta_{13}$	$4/(18\pi^2)$	C	1	2.6% (1.0σ)	Ch 8
N3	$\sin^2\theta_{23}$	$1/2 + 1/(2\pi^2)$	L	1+2(b)	Within 1σ	Ch 8
N4	Σm_ν	~0.059 eV (oscillation inputs)	L	3	Upper bounds only	Ch 8
N5	Normal mass ordering	structural (branch of SM §8.4)	C	1	Preferred by DESI+CMB; not confirmed	Ch 8
N6	PMNS sum rule	$2\Delta_{12} + \Delta_{23} = 0$	L	1+2(b)	1.178 ± 0.020 confirmed	Ch 8
N7	$\delta_{\rm CP}$	$b_{12} - b_{13}$ (subdominant U-odd)	L	2(b)[g]	T2K/NOvA: bijection-constraining	Ch 8

Summary statistics. Counts: 0 S + 2 C + 5 L. Total: 7. N2 inherits the Cabibbo λ and is conditional with it; N5 holds on the branch where the singlet taste hosts the Higgs and no gauge-singlet fermion is present (SM §8.4), the ordering a heuristic consequence of taste-breaking rather than a theorem. The numerical mass sum uses measured oscillation splittings; the remaining L-class entries divide into two kinds: (i) **sub-σ confirmed predictions** — N1, N3, N6

— whose Layer 1 structural form (TBM zeroth order plus the λ^2 scale, the latter conditional on the Cabibbo identification's emergent dressing) holds and whose specific values follow given the substrate Wilson-coefficient ratio $b_{23}/(b_{12}+b_{13}) = 4/3$, classified as Layer 2(b) per the §7.3 routing computation (the second-order derivation route excluded; see footnote [f] above); (ii) **parameterization-level prediction** — N7 — where the framework identifies $b_{12} - b_{13}$ as the Wilson-coefficient combination controlling δ_{CP} (a structural-form statement) but the specific value is bijection-specific and the framework "makes no commitment to any specific value of δ_{PMNS}" per SM §6.4 / §7.3 (see footnote [g]).

[g] *Parameterization-level prediction (δ_CP).* The framework identifies δ_{CP} with the sub-dominant U-odd combination $b_{12}-b_{13}$ in the A_2 second-order corrections to the charged-lepton sector (SM §7.3). This is a structural-form prediction: it specifies how δ_{CP} enters the framework's parameterization but does not predict its value. The value of $b_{12} - b_{13}$ is bijection-specific (Layer 2(b), property of the particular φ); the framework "makes no commitment to any specific value of δ_{PMNS}" (SM §5.4). Future T2K/NOvA measurements constrain the bijection's $b_{12} - b_{13}$ value rather than testing a framework prediction of δ_{CP} itself. This is qualitatively different from N1/N3/N6, where specific numerical values are fixed with no freedom and the JUNO-era fit matches $\sin^2\theta$_12 at 0.07σ (retrodictively).

A.4.4 Biology and medicine predictions

The framework's empirical content in biology and medicine is concentrated in the wider therapeutic window for reader-writer disorders (consistent with the Rett-syndrome reversibility of Guy et al. 2007) and the cancer methylome classification paradigm — structural organization of, and consistency with, established clinical biology.

#	Prediction	Quantitative content	Class	Layer	Empirical match	Chapter
B1	Wider therapeutic window for Rett	Substratum-emergent operator dissociation	S	1	Retrodiction (Guy et al. 2007)	Ch 16 §16.12
B2	Cancer methylome classifier accuracy	Substratum preserved during tumorigenesis	S	1	DKFZ 95%+ accuracy	Ch 17 §17.7
B3	Non-Markovian enzyme kinetics	Stretched-exponential waiting times	S	1	Multiple enzyme systems	Ch 13 §13.6

#	Prediction	Quantitative content	Class	Layer	Empirical match	Chapter
B4	Vibronic coherences in photosynthesis	Picosecond timescale	C	1	FMO observed; does not discriminate against standard vibronic accounts	Ch 13 §13.4
B5	Millisecond radical-pair coherence in cryptochrome	Slow-bath C2	C	1	Wiltschko 2019	Ch 13 §13.5
B6	Molecular clock overdispersion	$R(t) \approx 5$ for mammals	S	1	Gillespie 1989, Cutler 2000	Ch 17 §17.9
B7	Universal rate autocorrelation	Across the tree of life	S	1	Tao et al. 2019 confirmed	Ch 17 §17.9
B8	LTEE power-law fitness	$\beta \approx 0.08$-0.12	S	1	Wiser et al. 2013 confirmed	Ch 17 §17.9
B9	Punctuated equilibrium	C1–C4 history readback	S	1	Eldredge-Gould 1972	Ch 17 §17.9
B10	Hsp90 evolutionary capacitance	C1–C4 buffering	S	1	Rutherford-Lindquist 1998	Ch 17 §17.9
B11	Chk1 + gemcitabine schedule dependence	Memory asymmetry	S	1	Clinical literature consistent	Ch 16 §16.3

#	Prediction	Quantitative content	Class	Layer	Empirical match	Chapter
B12	LIFU cognitive improvement in Alzheimer's	τ_B normalization	S	1	Ye et al. 2025 confirmed	Ch 16 §16.4
B13	iPSC reprogramming residuals	Substratum-emergent dissociation	S	1	Lin et al. 2024 confirmed	Ch 16 §16.10
B14	Bivalent chromatin states	Multi-operator substratum	S	1	Bernstein et al. 2006 confirmed	Ch 16 §16.10

Summary statistics. Counts: 12 S + 2 C. B1 (Rett syndrome wider therapeutic window, Guy et al. 2007) is a retrodiction, in the same category as B6-B10 — the framework accounts for phenomena established before it. None is a forward confirmation. B4 and B5 are C rather than S because coherence preservation is a property of the specific physical realization and is not implied by (C1)–(C4) (§13.4-13.5): the framework's content in those two rows is the non-Markovian accounting, not a derivation that coherence survives.

A.5 Forward predictions and falsification targets

Beyond the predictions with current empirical match, the framework makes specific forward-looking predictions that are testable but not yet tested. The table also lists one conditional prospect, F7, a proposed mechanism whose observable response is not yet specified and which carries no quantitative prediction until it is. These provide the framework's empirical exposure for the next several years of experimental research.

Quantum engineering.

#	Prediction	Quantitative content	Testability	Chapter
F1	BEC analogue-gravity $\mathcal{F}(r) = 2r - r^2$ (proposed model form; derivation unpackaged)	Capacity-controlled revival	Steinhauer group, 2-3 years	Ch 15 §15.8
F2	Correlation-aware error correction	correlated errors over $\sim \tau_B/\tau_S$ gates; gain not quantified	Current quantum hardware	Ch 15 §15.4
F3	Backflow-aware NV magnetometry	sensitivity gain in direction only; factor not quantified	Current NV platforms	Ch 15 §15.6
F4	Engineered-bath coherence extension	coherence beyond the Markovian estimate; factor not quantified	Spin chain / NV / trapped ion platforms	Ch 15 §15.5

Cosmology and gravity.

#	Prediction	Quantitative content	Testability	Chapter
F6	Running vacuum DESI test	$w(z) = -1 + O(\nu)$ with $\lvert\nu\rvert \sim 10^{-32}$ (Type II channel; no phantom crossing)	DESI Year 5 (2-3 years)	Ch 7 §7.8
F7	Born rule transient violations (conditional prospect: source dynamics, initial-deviation family and observable map not fixed)	primordial power spectrum or non-Gaussianity; form not derived	LiteBIRD, future CMB-S4, once the response is derived	Ch 18 §18.2

#	Prediction	Quantitative content	Testability	Chapter
F8	Cosmic recurrence	period at most $g(\|S\|)$; $\|S\|$ not fixed	Logically falsifiable only	Ch 18 §18.7

Biology and medicine.

#	Prediction	Quantitative content	Testability	Chapter
F9	Variant pathogenicity stratification for reader/writer/eraser	AlphaMissense miscalibration for machinery genes	Reanalysis of existing benchmarks	Ch 17 §17.8
F10	τ_B-matched cancer chemotherapy schedules	Memory-priming protocols	Animal models, clinical trials	Ch 16 §16.3
F11	Anti-persister antibiotic class	Memory-erasing drugs	Drug screening with regulatory dynamics	Ch 16 §16.5
F12	Hurst exponent of substitution rates	$H \approx 0.79 \pm 0.03$ for typical taxa — *retrodiction, not a forward bet: H is read off existing data, and $H \approx 0.7$–0.8 is the generic broad-relaxation value (§8.11 of [Complexity]), so it does not discriminate the framework from a mundane superposition null; only the* ***derived*** *capacity-dependence $H(C3)$ would be a genuine test, and that relationship is not yet derived*	Existing phylogenetic data (consistency, not exposure)	Ch 17 §17.9

Foundations (Lorentz sector).

#	Prediction	Quantitative content	Testability	Chapter
F13	Cross-charged boost-LV decoupling	$\delta c \sim \Delta C_2\,(\alpha_0/4\pi)(M_X/M_{\rm Pl})^2[c_L \ln(M_{\rm Pl}/M_X) + c_0]$; the symmetric-point matching is certified ([SM §4.4]); the quadratic-log *form*, its scalar-/fermion-line and non-absorbed controls, and the constants $c_L \simeq -9.4 \pm 0.3$, $d \simeq -0.88$ were reported by a one-loop calculation whose driver and numerical output are not available for verification, and are not independently verified; on them the illustrative consistency window is $M_X \lesssim 3.9 \times 10^9$ GeV against the electron-sector SME coefficient and $M_X \lesssim 4 \times 10^7$ GeV against the GZK/Auger isotropic-n=2 bound (SM §3.1 (*Scope of the emergent-Lorentz claim*)), with the electron-SME and photon-sector ceilings (7.5×10^9, 1.1×10^{12} GeV) the looser sector-specific figures; the overall normalization ΔC_2 remains open, so these are illustrative windows, not predictions or bounds	Multi-species LV (matching certified; running reported, unverified; full reach pending ΔC_2)	SM §3.1 (*Scope of the emergent-Lorentz claim*, point (iii))

The *rotational* Lorentz sector is deliberately **not** listed as a forward prediction. Unlike the boost sector (F13), the rotational E_g anisotropy is a custodial-U(6) singlet with no power-law suppression ally, and its magnitude is undetermined

— turning on whether the emergent theory preserves the octahedral symmetry exactly at loop level, with an aligned condensate (Chapter 19 §19.3.7), between a negligible dimension-≥ 6 residual ($\sim 10^{-32}$), whose coefficient the smearing-kernel width R/ε sets without changing its dimension, and, if that symmetry fails, an unsuppressed dimension-≤ 4 operator already constrained by laboratory sidereal experiments. It is therefore the sharpest *open question* in the framework's Lorentz sector, treated as such in SM §3.1 (*Scope of the emergent-Lorentz claim*, point (iii)), not a predictive bet. Its one regime-independent feature — that any detectable residual must take the $\ell = 4$ cubic-harmonic form aligned to the cosmic rest frame and correlated across species — is recorded there as a conditional falsification template, not here as a forward prediction.

Falsification conditions. Each forward prediction has specific falsification conditions developed in the relevant chapter. The framework's content is most distinctively at risk in:

- F1 (BEC analogue-gravity): null result with functional form different from $2r - r^2$ would falsify the proposed functional form — a model prediction, not theorem-derived (Ch 15 §15.8).
- F6 (DESI running vacuum): DESI Year 5 confirmation of phantom-crossing dark energy would contradict the framework's Type II running-vacuum reading — the strongest available disconfirmation of its dark-energy account; the framework supplies the smooth $H^2 - H_0^2$ bookkeeping but does not derive a distinctive Type II law.
- F9 (variant pathogenicity stratification): demonstration that AlphaMissense performance is uniform across reader/writer/eraser and substrate genes would falsify the substratum-emergent operator distinction at the variant-prediction level.
- F13 (cross-charged decoupling): observation of an inter-species boost-Lorentz-violation that does *not* decouple with the heavy cross-charged mass — i.e. a non-decoupling (constant or pure-logarithmic) sector-differential limiting-speed difference — would falsify the framework's condensate-stabilizer mechanism for gauge content (Theorem 5, revised), under which, conditional on the symmetric-point matching under H-blind ([SM §4.4]), the differential is suppressed as $(M_X/M_{\rm Pl})^2$ up to a logarithm. The *form* of this suppression and the coefficient $c_L \simeq -9.4 \pm 0.3$ were reported by a one-loop calculation whose driver and numerical output are not available for verification; on them the illustrative consistency window is $M_X \lesssim 3.9 \times 10^9$ GeV. The overall normalization ΔC_2 is not fixed, and the window carries no weight as a prediction or bound until the calculation is independently reproduced.

A.6 Framework's epistemic position summary

The framework's cumulative empirical record can be summarized as follows.

Strongest current evidence. The framework currently has no pre-registered

forward confirmation. Its strongest empirical content is retrodictive: JUNO $\sin^2\theta_{12}$ at 0.07σ (entry 12 / N1; the derivation postdates the November-2025 first results), Koide Q to better than 0.001%, and Cabibbo at 0.04%. The genuinely pre-registered content is forward-looking — DESI Y5 (ν), BTFR at $z > 1$, the A2 lattice thresholds, and JUNO design-lifetime precision on the exact $\sin^2\theta_{12}$ value — documented in dated artifacts preceding their future data. The Bekenstein-Hawking coefficient is a consistency relation with no direct empirical test (A.4.2) — GW250114 constrains the classical area theorem, not the coefficient. The Rett wider-window result (entry B1) is a retrodiction, not counted among the forward confirmations.

Most distinctive content. Two predictions distinguish the framework from standard accounts most sharply: the substratum-emergent operator distinction (with empirical signatures in B1, B2, B13, B14) and the proposed capacity-controlled origin of partial revival, $\mathcal{F}(r) = 2r - r^2$ (proposed model form; empirical exposure in F1). Partial revival itself is standard open-systems phenomenology; the capacity-controlled origin and the scaling form are the proposal's own.

Sharpest single match. Koide $Q = 2/3$ to better than 0.001% (entry 10) — the framework's sharpest individual empirical match in relative precision.

Precision and empirical standing. The comparisons have different evidential roles: Koide and Cabibbo supply numerical retrodictions, the fitted coupling entries are retrodictions by construction, $A \cdot B$ is an inconclusive finite-size extrapolation, and the neutrino mass sum is constrained by upper bounds. The quoted precision belongs to each comparison separately.

Most consequential forward exposure. BEC analogue-gravity test (F1) and DESI Year 5 running vacuum test (F6) provide the framework's most consequential near-term empirical exposure. Both have specific falsification conditions that would substantially constrain the framework's content if not confirmed.

Domains where framework has substantial content but no current empirical match. The framework's content in computational complexity (BQP theorem, Chapter 14), evolutionary biology Hurst exponent (F12), and the broader reader-writer disorder class beyond Rett (F9, F10, F11) has not yet been specifically tested against the framework's predictions. These are testable predictions awaiting empirical engagement rather than predictions matching established data.

Where the framework has no current content. Five fundamental physics problems are scope limits from the framework's perspective — places where it does not provide a derived value: the fermion mass hierarchy in detail (flavor problem), baryogenesis (solution-specific — the baryon asymmetry inherits the input status of the framework's flavor-sector CP phases, §19.3.3), inflation and initial conditions, the Hubble tension's specific resolution, and the cosmological initial state. These are explicitly characterized in Chapter 19.

Cumulative assessment. The framework's empirical record spans multiple domains, and its entries are of unequal evidential weight, classified explicitly above: of the twenty-two Standard Model observables, two are parameter-free structural retrodictions, five are conditional structural retrodictions (the Cabibbo λ with its two inheritors and the two Koide entries), and the remainder are layered, chained, or fitted; the seven gravitational-sector entries comprise five conditional results and two layered estimates, with the Page curve (G11), an open research target, and four consistency or descriptive relations stated beside them, while the seven neutrino-sector entries include retrodictions and bound-only comparisons, with the 0.07σ JUNO match the sharpest (a retrodiction — the derivation postdates the November 2025 measurement); the fourteen biology/medicine entries are retrodictions and consistency results. Genuinely pre-registered forward content is the smaller set awaiting data (DESI Y5 ν, BTFR at $z > 1$, the A2 lattice thresholds, JUNO design-lifetime precision on the exact $\sin^2\theta_{12}$ value). The framework's empirical case is therefore a broad retrodictive compression across domains, with its forward-test exposure concentrated in that smaller pre-registered set, rather than a single decisive confirmation.

The framework's distinctive epistemic position is that it provides *structural foundations* for the empirical patterns rather than competing theoretical content at the emergent level. Standard QM, standard QFT, and standard GR are reproduced at the emergent level; the framework's distinctive content is the structural foundations explaining why these emergent theories have the specific forms they do. The cumulative empirical record provides convergent evidence for the framework's structural commitments across multiple domains, with no single empirical match being load-bearing for the framework's overall content.

Appendix B

Mathematical Derivations

B.1 What this appendix develops

The main chapters of the book develop the framework's content with proofs sketched at the level required for the chapter argument's logical flow. This appendix collects the full mathematical derivations of the framework's key technical results — derivations that are referenced in the main chapters but kept out of the chapter flow to preserve readability.

The appendix's content is organized by mathematical structure rather than by chapter sequence. Six derivations occupy the appendix.

B.2 The Stinespring construction. The framework's emergent quantum description is constructed from a deterministic substratum bijection through the Stinespring dilation theorem applied to a partial trace over the hidden sector. This appendix gives the full construction: Hilbert-space embedding, permutation unitarity lemma, CPTP channel structure, Born-form representation of the transition probabilities, emergent coherence theorem, and CP-indivisibility theorem. The construction uses only textbook operator algebra from the 1950s.

B.3 The trace-out as a Jordan-Chevalley projection. The framework's content on the Standard Model gauge structure relies on a structural property of the trace-out operation: it extracts the semisimple part of the evolution matrix's Jordan-Chevalley decomposition and discards the nilpotent monodromy. This appendix gives the full algebraic derivation: period formula, Jordan-Chevalley decomposition, q-independence of the Weil-Deligne conductor, additive decomposition over gauge irreps, and the precise sense in which the trace-out is a projection.

B.4 The gap equation for $\hbar$. The framework relates Planck's constant $\hbar$ to the discreteness scale through a thermal self-consistency argument between the classical horizon temperature (substratum-level) and the KMS temperature (emergent QFT-level). This appendix gives the full four-step derivation: spatial locality and boundary dependence, deep-sector gauge equivalence, dimensional determination, and thermal self-consistency producing $\hbar = c^3\epsilon^2/(4G\sigma)$.

B.5 Anomaly cancellation and unique hypercharges. The framework's content on the Standard Model matter content derives the unique hypercharge assignment on the nonzero-Y_Q branch from the six anomaly conditions of $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$. This appendix gives the explicit calculation: the six anomaly conditions, the algebraic constraint structure, and the unique solution producing $(Y_Q, Y_u, Y_d, Y_L, Y_e) = (1/6, 2/3, -1/3, -1/2, -1)$ with no free parameters.

B.6 Gauge-theoretic derivations. Two key gauge-theoretic results from Chapter 5: the conditional H-link link count for $K = 2dm = 6$ internal components per site, and the cubic-rotation-group decomposition of the six link directions into block multiplicities $(3, 2, 1)$ — the multiplicities that determine $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$.

B.7 The lemma chain for the characterization. The framework's characterization theorem — accessible non-Markovianity $\Longleftrightarrow$ C1/C3/C4 realization per accessible horizon, with the quantum representation internal and universal ($S \iff D \iff Q_{\mathrm{fb}}$, [Main §3.4]) — relies on a chain of intermediate lemmas: partition-relativity, emergent stochasticity, P-indivisibility from a non-permutation witness, accessible-timescale conditional memory, and phase-locking, together with the results on which phase-locking rests: Bohr-frequency completeness and the identification of level pairs under the triangle-genericity condition, and, without that condition, the real-to-integer transfer and the two-branch classification conditional on H-BG. This appendix collects these lemmas with their proofs in compact form, providing the reader a self-contained reference for the frame-

work's logical structure.

The appendix's content is *complete derivations*, not exposition. Each derivation gives the algebraic steps, the lemmas invoked, and the result with no glosses. For motivation and context, the relevant main-chapter sections are cross-referenced at the start of each derivation. The appendix is designed for readers wanting full technical content; the main chapters give the same results with motivating context but compressed derivations.

B.2 The Stinespring construction for emergent quantum mechanics

Main-text reference: Chapter 1 §1.6.

The framework's emergent quantum description is constructed from the substratum's deterministic bijection through a Stinespring dilation applied to the partial trace over the hidden sector. The construction uses only textbook operator algebra from the 1950s, with no reference to recent stochastic-quantum correspondence results.

Setup. The finite configuration spaces $\mathcal{C}_V = \{x_1, \ldots, x_n\}$ (visible sector) and $\mathcal{C}_H = \{h_1, \ldots, h_m\}$ (hidden sector) embed into Hilbert spaces $\mathcal{H}_V = \mathbb{C}^n$ and $\mathcal{H}_H = \mathbb{C}^m$ via the canonical identification $|i\rangle \leftrightarrow x_i$ and $|k\rangle \leftrightarrow h_k$. This introduces no quantum postulates: it is the canonical identification of probability distributions on a finite set with diagonal density matrices.

The combined space is the tensor product $\mathcal{H} = \mathcal{H}_V \otimes \mathcal{H}_H$ of dimension nm. The substratum's deterministic bijection $\varphi : \mathcal{C}_V \times \mathcal{C}_H \to \mathcal{C}_V \times \mathcal{C}_H$ defines a permutation of the orthonormal basis $\{|i,k\rangle\}_{i=1,\ldots,n;k=1,\ldots,m}$.

Lemma B.2.1 (Permutation unitarity). *Any bijection $\varphi : \mathcal{C}_V \times \mathcal{C}_H \to \mathcal{C}_V \times \mathcal{C}_H$ defines a unitary operator U_φ on $\mathcal{H}$.*

Proof. Define $U_\varphi|i,k\rangle = |\varphi(x_i, h_k)\rangle$. Since φ is a bijection, U_φ permutes the orthonormal basis, hence is unitary. □

A continuous unitary interpolation comes from the discrete step itself: U_φ, being unitary, admits a Hermitian logarithm — $U_\varphi = e^{-i\hat{H}}$ for some Hermitian $\hat{H}$, unique only up to the branch and gauge freedom of Lemma B.7.5 — which defines $U_t = e^{-i\hat{H}t}$ through the integer-time dynamics. The finite substratum carries no nontrivial continuous permutation family, so the interpolation is representation-level structure, not substratum dynamics.

Lemma B.2.2 (Reverse direction, scoped). *Any single one-step doubly stochastic matrix on a finite configuration space $\mathcal{C}_V$ can be realized as the one-step marginal of a deterministic bijection on $\mathcal{C}_V \times \mathcal{C}_H$ with uniform prior on $\mathcal{C}_H$, for some finite $\mathcal{C}_H$ (exactly when transition probabilities are rational; to arbitrary precision otherwise).* (The restriction is forced: a uniform-prior bijection marginal preserves the uniform distribution, hence is doubly stochastic; the

emergent unistochastic statistics lie inside this class.) The scope is the matrix, not the process: realizing prescribed multi-time statistics — a family of joint distributions over trajectories — with a single bijection and a single uniform prior is a stronger demand and is not claimed here.

Proof. Any marginal of a bijection with uniform prior is a doubly stochastic matrix. By Birkhoff-von Neumann, every doubly stochastic matrix is a convex combination of permutation matrices. A bijection on $\mathcal{C}_V \times \mathcal{C}_H$ realizing the required mixture is obtained by letting $\mathcal{C}_H$ enumerate the permutations; for multi-step processes, the construction extends by history dilation — made precise, with a structured prior, by the finite process-dilation theorem on each finite horizon (Chapter 1 §1.6; [Main §3.4]). □

Remark. Lemma B.2.2 is the bijection–doubly-stochastic-matrix correspondence for a single step with a uniform hidden prior; the process-level correspondence, with a structured prior, is the finite process-dilation theorem (Chapter 1 §1.6), the reverse leg of the finite representation theorem (§1.8), on each finite horizon [Main §3.4]. It is *not* a full CPTP-channel correspondence — channels that create coherences from diagonal inputs have no permutation-unitary realization with uniform ancilla. For the characterization theorem (Chapter 1 §1.8), which is stated in terms of transition probabilities, this is the appropriate scope.

The quantum channel. The observer's ignorance of the hidden sector corresponds to the maximally mixed state $\rho_H = I_m/m$. The visible-sector quantum channel is

$$\Phi(\rho_V) = \mathrm{Tr}_H\left[U_\varphi\,(\rho_V \otimes \rho_H)\,U_\varphi^\dagger\right]$$

This is completely-positive trace-preserving (CPTP) by a standard result (Nielsen-Chuang Theorem 8.1), with Kraus representation $\Phi(\rho_V) = \sum_{k,l} K_{kl}\,\rho_V\,K_{kl}^\dagger$ where $K_{kl} = m^{-1/2}\langle l|U_\varphi|k\rangle_H$. The triple $(\mathcal{H}_H, U_\varphi, \rho_H)$ is a *unitary realization* of Φ. The reference state here is the uniform mixed state, so the triple is a unitary realization rather than a Stinespring dilation: with a pure reference the Stinespring dilation is the associated isometry $\psi \mapsto U_\varphi(\psi \otimes |0\rangle)$, not the unitary, and obtaining one from a mixed reference requires purifying ρ_H.

Theorem B.2.3 (Born-form representation of the transition probabilities). *The classical transition probabilities $T_{ij} = P(j|i)$ derived from the substratum's bijection equal the Born-rule probabilities of Φ — a representation identity: it exhibits the Born form and does not select the quadratic exponent, which remains open ([Main §3.4]).*

Proof. $P(j|i) = \langle j|\Phi(|i\rangle\langle i|)|j\rangle = m^{-1}\sum_{k,l}|\langle j,l|U_\varphi|i,k\rangle|^2$. Since U_φ is a permutation, $\langle j,l|U_\varphi|i,k\rangle = \delta_{(j,l),\varphi(i,k)}$. Thus $P(j|i) = m^{-1}\sum_k \delta_{j,\pi_V(\varphi(x_i,h_k))} = T_{ij}$, where π_V is the projection from $\mathcal{C}_V \times \mathcal{C}_H$ to $\mathcal{C}_V$. □

Remark (failure of (C1)). Condition (C1) does not suffice. For $|V| = |H| = 2$ and $\varphi(x,h) = (h,x)$ one has $T_{ij} = 1/2$ for all i,j, yet $\Phi(\rho) = I/2$ identically: a constant channel, hence measure-and-prepare. Exhaustive enumeration gives

the same verdict for every (C1)-satisfying bijection at $|V| = |H| = 2$, and for 612 of 648 at $|V| = 2$, $|H| = 3$. The obstruction is structural: (C1) constrains the diagonal of Φ, whereas entanglement-breaking is a property of $\mathrm{id} \otimes \Phi$ and is not fixed by the action on single-system inputs.

What replaces (C1) is not a further condition on T but a condition on the partition, and it is established for a specific realization rather than for arbitrary bijections. Take the substratum of §B.1 on a lattice Λ with $q = 2$, state $(u, v) \in \mathbb{F}_2^\Lambda \times \mathbb{F}_2^\Lambda$ and the reversible nearest-neighbour update

$$u'_x = \sum_{z \sim x} u_z + v_x, \qquad v'_x = u_x,$$

the visible sector being the sites of a region $R \subseteq \Lambda$. Write M_{HV} and M_{VH} for the blocks of this update carrying visible input to hidden output and hidden input to visible output, $\kappa = \dim\ker M_{HV}$, $w = \operatorname{rank} M_{VH}$, and $s = \dim V = 2|R|$. The inner and outer boundaries of R are $\partial^- R = \{x \in R : \exists\, z \notin R,\ z \sim x\}$ and $\partial^+ R = \{y \notin R : \exists\, z \in R,\ z \sim y\}$.

Lemma 1 (boundary bound). $\operatorname{rank} M_{HV} \le |\partial^+ R|$ *and* $\operatorname{rank} M_{VH} \le |\partial^- R|$.

Proof. For $y \notin R$ the component $v'_y = u_y$ is a hidden input, and $u'_y = \sum_{z \sim y} u_z + v_y$ involves a visible input only if some $z \sim y$ lies in R, that is only if $y \in \partial^+ R$; so M_{HV} has non-zero rows only among $\{u'_y : y \in \partial^+ R\}$. Dually, for $x \in R$ the component $v'_x = u_x$ is a visible input, and u'_x involves a hidden input only if $x \in \partial^- R$. □

One update step transports information one lattice site, so the sectors resolve one another only across the cut. What the argument establishes is a factorization: each cross-block has non-zero rows only at the boundary sites named, so it factors through the corresponding boundary coordinates. That statement uses no property of the coefficient ring and holds verbatim over $\mathbb{Z}_q$; over a field — in particular over the $\mathbb{F}_2$ used here — it gives the displayed rank bounds, ordinary vector-space rank being the relevant notion there. For a general nearest-neighbour bijection the same support argument yields the weaker conclusion that the hidden output depends on the visible input only through $\partial^- R$. Since adjacency is symmetric, the same reading applies to the columns: M_{HV} reads visible input only through $\{u_x : x \in \partial^- R\}$ and M_{VH} only through $\{u_y : y \in \partial^+ R\}$, so each rank is in fact bounded by $\min(|\partial^- R|, |\partial^+ R|)$ — a sharpening that uses the symmetry where the row statement does not, and that Corollary 2 consumes.

Lemma 2 (channel normal form). *Write $A = M_{VV}$, $B = M_{VH}$, $C = M_{HV}$ for the blocks of the update, and L_{RR} for the adjacency of Λ restricted to R. Then: (i) A is invertible, with $A^{-1}(a_u, a_v) = (a_v,\ a_u - L_{RR}\, a_v)$, so the visible block alone induces the permutation (Clifford) unitary $V_A|i\rangle = |Ai\rangle$; (ii) $\Phi(|i\rangle\langle i'|)$ vanishes unless $C(i - i') = 0$; (iii) on the Weyl operators the channel acts, with no phase, as*

$$\Phi(X^a Z^b) = [\, Ca = 0 \,]\, [\, B^T A^{-T} b = 0 \,]\, X^{Aa}\, Z^{A^{-T} b},$$

so that $\Phi = \mathrm{Ad}_{V_A} \circ \Phi_G$*, where* $\Phi_G = |G|^{-1} \sum_{g \in G} g \cdot g^\dagger$ *is the uniform mixture over the group* G *of Weyl operators* $X^\alpha Z^\beta$ *with* $\alpha \in \mathrm{im}(A^{-1}B)$ *(translations) and* $\beta \in \mathrm{im}\, C^T = (\ker M_{HV})^\perp$ *(phases); (iv)* G *is abelian, of order* $2^{\,s-\kappa+w}$*.*

Proof. (i) is the displayed inverse, read off from $v'_x = u_x$ and $u'_x = (L_{RR} u)_x + v_x$ on R; invertibility of A is what makes $i \mapsto Ai$ a relabelling. (ii) The hidden outputs of i and i' agree for some hidden input exactly when $C(i - i') = 0$, and then for every hidden input. (iii) With the hidden input uniform, $\Phi(|i\rangle\langle i'|) = [\, C(i - i') = 0 \,]\, \mathbb{E}_{t \in \mathrm{im}\, B} |Ai + t\rangle\langle Ai' + t|$. Summing this against $X^a Z^b = \sum_i (-1)^{b \cdot i} |i + a\rangle\langle i|$ and substituting $i \mapsto i - A^{-1}t$ gives the displayed rule: the character sum over $\mathrm{im}\, B$ produces the second indicator and cancels every phase. The right-hand side is Ad_{V_A} applied to the Hilbert–Schmidt-orthogonal projection onto $\mathrm{span}\{X^a Z^b : Ca = 0,\ B^T A^{-T} b = 0\}$, and the uniform mixture over the symplectic complement of that index set — which is exactly G — is that projection. (iv) An element $X^\alpha Z^\beta$ with $\alpha \in \mathrm{im}(A^{-1}B)$, $\beta \in \mathrm{im}\, C^T$ commutes with another iff the cross-pairings $\alpha \cdot \beta'$ vanish; over all of G this is the single condition $CA^{-1}B = 0$, which holds identically for this update: B feeds hidden u into the visible u'-components, A^{-1} carries the u'-subspace onto the v-subspace, and C reads only the visible u-components. The translation and phase families intersect only in the identity, so $|G| = 2^{\mathrm{rank}\, B} \cdot 2^{\mathrm{rank}\, C} = 2^{\,w+(s-\kappa)}$. □

Theorem (separability threshold). *Let* G *be an abelian subgroup of the Weyl group on* s *qubits, of order* 2^t *modulo phases. Then* $\Phi_G = |G|^{-1} \sum_{g \in G} g \cdot g^\dagger$ *is entanglement-breaking if and only if* $t = s$*.*

Proof. Modulo phases the Weyl operators form the symplectic space $\mathbb{F}_2^{2s}$, on which abelian subgroups are exactly the isotropic subspaces; isotropy bounds $t \le s$. Any isotropic subspace of dimension t is carried to $\langle Z_1, \ldots, Z_t \rangle$ by a symplectic transformation (Witt's extension theorem for the symplectic space $\mathbb{F}_2^{2s}$), and every symplectic transformation is implemented by a Clifford unitary U. Hence $U \Phi_G U^\dagger$ is complete dephasing on t qubits tensored with the identity on the remaining $s - t$. Complete dephasing is measure-and-prepare and therefore entanglement-breaking, and a tensor product of channels is entanglement-breaking exactly when every factor is — separability of the joint Choi state survives the partial trace onto any single factor pair, so one entangled factor Choi rules out separability of the whole; the identity on a non-trivial factor is not entanglement-breaking. Since conjugation by a unitary preserves the property, Φ_G is entanglement-breaking precisely when $s - t = 0$. □

Corollary 1. *For one step of the displayed update on* $\mathbb{F}_2$ *—* $u'_x = \sum_{z \sim x} u_z + v_x$*,* $v'_x = u_x$*, with the hidden sector initially uncorrelated and maximally mixed — the visible channel* Φ *admits the Clifford-conjugated abelian Weyl-mixture normal form of Lemma 2 and is not entanglement-breaking if and only if* $w < \kappa$*.* Immediate from Lemma 2 and the Theorem with $t = s - \kappa + w$: entangle-

ment breaking is invariant under composition with the unitary Ad_{V_A}, so Φ is entanglement-breaking exactly when Φ_G is, and $t = s$ reads $w = \kappa$ (Lemma 2 gives $w \le \kappa$, matching $t \le s$).

Corollary 2. *If* $\min(|\partial^- R|, |\partial^+ R|) < |R|$ *then* Φ *is not entanglement-breaking.*

Proof. $s = 2|R|$. Lemma 1's column sharpening bounds each of $\operatorname{rank} M_{HV} = s - \kappa$ and $\operatorname{rank} M_{VH} = w$ by $\min(|\partial^- R|, |\partial^+ R|)$, so $w + (s - \kappa) \le 2\min(|\partial^- R|, |\partial^+ R|) < 2|R| = s$, that is $w < \kappa$. □

Remark. For a d-dimensional region of linear size ℓ, $|\partial R| = O(\ell^{d-1})$ against $|R| = \ell^d$, so the hypothesis of Corollary 2 — the smaller boundary outnumbered by the region — holds for every sufficiently large block; for an exact cubic block in $d = 3$ it reads $(\ell - 2)^3 > 0$ and holds exactly when the block has a non-empty bulk, every side $\ell \ge 3$. Within this single-step result the criterion carries no dependence on $|H|$, since the coupling reads only the boundary; the counterexample above is the degenerate case, every visible site lying on the cut so that the region has no bulk. Whether the same holds over many steps is not settled here: repeated interaction can propagate boundary information into the hidden bulk, and a multi-time statement would require a process-tensor argument rather than a single-channel one. The suggestion that coherence is a bulk quantity and decoherence an area quantity — the accounting that reappears in the horizon entropy of Chapter 7 — is at this stage an interpretation of the one-step result, not a theorem about the asymptotic dynamics.

Remark (scope). Lemma 1's factorization holds over any $\mathbb{Z}_q$, the displayed rank bounds being its reading over a field, and in weakened form the support argument covers arbitrary nearest-neighbour bijections. Lemma 2, the Theorem and both Corollaries are stated for $q = 2$; the symplectic and Clifford arguments carry over to prime q with $2^\bullet$ replaced by $q^\bullet$, while composite q requires module-theoretic treatment of rank and kernel and is not covered. All of this concerns the realization displayed above and is not a property of arbitrary finite bijections. For a general linear bijection the abelian property of Lemma 2(iv) can fail ($CA^{-1}B \neq 0$); the mixture is then over a non-abelian Weyl family, and entanglement breaking is governed by the isotropy of the surviving index set rather than by its order alone — not treated here. Two boundary markers on interpretation. Coherence preservation is a property of this realization, not an additional structural condition: (C1)–(C4) constrain coupling, timescales, and capacity, and neither contain nor imply a coherence requirement — the swap example satisfies (C1) while its channel is entanglement-breaking, and which realizations preserve one-step coherence is the geometric question answered above for the displayed update alone. And non-entanglement-breaking is a statement about a single channel, not about quantum mechanics: it supplies neither the observable algebra, nor preparations and interventions, nor tensor composition, nor multi-time statistics — the operational reconstruction claims rest on the dilation and correspondence arguments, not on Corollary 1 — and the entanglement-breaking status of one step does not determine whether the visible multi-time process is P-divisible: the hidden sector can carry memory

even when a step's channel is entanglement-breaking, as a two-state system with a two-state hidden register already shows.

Theorem B.2.5 (CP-indivisibility). *The P-indivisibility of the substratum dynamics across its recurrence cycle (from a non-permutation rooted map and finiteness) implies CP-indivisibility of* $\{\Phi_t\}$*: there exist* $t_2 > t_1 > 0$ *with no CPTP map* Λ *satisfying* $\Phi_{t_2} = \Lambda \circ \Phi_{t_1}$.

Proof. The framework's Φ_t are permutation dilations with uniform ancilla, so they map computational-diagonal states to exactly diagonal states (a permutation unitary conjugates a diagonal state to a diagonal state; the partial trace preserves diagonality). If $\Phi_{t_2} = \Lambda \circ \Phi_{t_1}$ with Λ CPTP, then $M_{kj} := \langle j|\Lambda(|k\rangle\langle k|)|j\rangle$ is stochastic and, by the diagonality of $\Phi_{t_1}(|i\rangle\langle i|)$, $T(t_2) = T(t_1)\,M$ — P-divisibility of the population process. The contrapositive gives the result. For a general CPTP family the diagonal-input reduction fails — a Hadamard pair is CP-divisible with P-indivisible populations — so the diagonal-preservation property is what licenses the step. □

For bijective (invertible) dynamical maps, CP-indivisibility is completely witnessed by ancilla-assisted distinguishability ([Main §3.2]); ordinary system-only trace-distance revival — the Breuer-Laine-Piilo *information backflow* signature — is sufficient evidence of non-Markovianity, not a necessary consequence of CP-indivisibility. The framework's content is therefore that the C1–C4/readback sector is necessarily memory-bearing — non-Markovian, with conditional memory at observable timescales — while fixed-basis quantum representability itself is universal, not confined to that sector ([Main §3.4]).

Approximate unitarity. On observable timescales $t \ll \tau_B$, the hidden-sector state is approximately frozen (the slow-bath regime, (C2-slow); at the horizon the bath is also imperturbable on laboratory timescales, with the $N_H \gg N_V$ hierarchy of (C3)'s realization clause). At $t = 0$ the channel generator is exactly Hamiltonian,

$$\left.\frac{d\Phi_t}{dt}\right|_{t=0}(\rho_V) = -i[\hat{H}_{\text{eff}}, \rho_V],$$

with $\hat{H}_{\text{eff}} = H_V + \text{Tr}_H[V_{\text{int}}(\mathbb{1} \otimes \rho_H)]$; the dissipative part vanishes at $t = 0$. At finite time the dissipator $\mathcal{D}$ has two contributions: a frozen-bath dephasing of order the coupling strength squared (the spread of bath-conditioned visible Hamiltonians, independent of τ_B), and a bath-motion correction of order $(\tau_S/\tau_B)^2$. The dynamics is the Schrödinger equation generated by $\hat{H}_{\text{eff}}$ to leading order provided both the slow-bath condition $\tau_S \ll \tau_B$ and a weak-coupling condition (small conditioned-Hamiltonian spread) hold. The phase-locking lemma (Chapter 1 §1.6) then determines $\hat{H}_{\text{eff}}$ from continuous-time transition data, under the genericity hypotheses of Lemma B.7.5, up to the lemma's full gauge: an overall energy shift, the residual diagonal-unitary rephasing freedom that is physically trivial (basis convention), and the antiunitary conjugation $\hat{H} \to -\hat{H}^*$ — a genuine twofold ambiguity (Lemma B.7.5).

The construction is complete: the substratum's deterministic bijection produces the memory-bearing visible-sector description — its unitary quantum representation supplied through the Stinespring dilation, the coherent operational lift still to be established ([Main §3.4]) — with leading-order Schrödinger evolution and structurally non-Markovian corrections at $\mathcal{O}(\tau_S/\tau_B)$.

B.3 The trace-out as a Jordan-Chevalley projection

Main-text reference: Chapter 5 §5.4-5.6.

The framework's content on the Standard Model gauge structure relies on a precise algebraic property of the trace-out operation: it extracts the semisimple part of the evolution matrix's Jordan-Chevalley decomposition and erases the nilpotent monodromy. This appendix gives the full derivation for the scalar wave equation on a ring of L sites over $\mathbb{F}_q$; the multi-component extension incorporating the gauge group follows by additive decomposition.

Setup. The discrete wave equation $x(n,t{+}1) = x(n{-}1,t){+}x(n{+}1,t){-}x(n,t{-}1) \pmod q$ has phase space $(\mathbb{Z}/q\mathbb{Z})^{2L}$ and evolution matrix

$$F = \begin{pmatrix} 0 & I \\ -I & A \end{pmatrix}$$

where A is the circulant adjacency matrix of the ring graph.

Theorem B.3.1 (Period formula). *For q prime with $\gcd(L,q) = 1$ — $q = 2$ included:*

$$\operatorname{ord}(F \bmod q) = qL$$

Proof. For each Fourier mode k the $2{\times}2$ block B_k has characteristic polynomial $t^2 - \lambda_k t + 1$ with $\lambda_k = \zeta^k + \zeta^{-k}$, so its roots are $\zeta^{\pm k}$ and the diagonalizable blocks contribute orders dividing L, the mode at a primitive L-th root of unity contributing order exactly L. At parabolic modes ($\lambda_k = \pm 2$), the Jordan form gives

$$B_k^n = \alpha^n \begin{pmatrix} 1 & n\alpha^{-1} \\ 0 & 1 \end{pmatrix}$$

of order q at $\alpha = 1$ and $2q$ at $\alpha = -1$. The second parabolic mode occurs only for even L, since $\alpha = -1$ requires $(-1)^L = 1$; so the parabolic orders are q alone for odd L, and q together with $2q$ for even L, where the extra factor 2 already divides L. In both cases $\operatorname{ord}(F) = \operatorname{lcm}(L,q) = qL$, using $\gcd(L,q) = 1$. Theorem B.3.2 is consistent with this and does not replace it: its decomposition $F = F_{\mathrm{ss}} F_u$ gives commuting factors whose orders divide L and q, and the exact values need the primitive-root mode and $N \neq 0$. At $q = 2$ the parabolic mode at $\zeta = 1$ is present, so $N \neq 0$ and the unipotent factor has order exactly 2, while $\gcd(L,2) = 1$ forces L odd; hence $\operatorname{ord}(F \bmod 2) = 2L$. □

The period formula has been verified computationally for $q \in \{2,3,5,7,11,13\}$ and $L \leq 12$.

Theorem B.3.2 (Jordan-Chevalley decomposition). *Define* $N = (F^L - I)/L \bmod q$. *Then:*

(i) N *is nilpotent with* $N^2 = 0$, *and rank*(N) *is the number of parabolic modes: 2 for even* L, *where these are* $\zeta = \pm 1$, *and 1 for odd* L, *where* $\zeta = -1$ *is not an* L*-th root of unity.*

(ii) $F_u = I + N$ *is unipotent with* $F_u^q = I$.

(iii) $F_{ss} = F \cdot (I - N)$ *is semisimple with* $F_{ss}^L = I$.

(iv) $F = F_{ss} \cdot F_u$ *and* $[F_{ss}, N] = 0$.

Proof. F^L has the form $I + LN'$ where N' arises from the Jordan blocks: $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^L = \begin{pmatrix} 1 & L \\ 0 & 1 \end{pmatrix}$ contributes a rank-1 nilpotent, and similarly for $\alpha = -1$ when L is even, that mode being present exactly when $(-1)^L = 1$. So $N = N'$ has $N^2 = 0$ (each block is rank-1 nilpotent on a 2D subspace) and rank(N) equal to the number of parabolic modes, 2 for even L and 1 for odd L. Since $N^2 = 0$: $(I+N)^{-1} = I - N$, giving $F_{\text{ss}} = F(I-N)$. Then $F_{\text{ss}}^L = F^L(I-N)^L = (I + LN)(I - LN + \cdots) = I \bmod q$ since terms involving LN cancel modulo q (using $N^2 = 0$). Commutativity follows from N being supported on the parabolic eigenspaces, which are F-invariant. □

The decomposition has been verified computationally for $L \in \{4, 6, 8, 10, 12\}$ and $q \in \{3, 5, 7, 11, 13\}$.

Theorem B.3.3 (q-independence of the Weil-Deligne conductor). *The Weil-Deligne conductor*

$$\mathfrak{f}_{\text{WD}} = \mathfrak{f}_{\text{ss}}(L) + \text{rank}(N)$$

is q*-independent when* $\gcd(L, q) = 1$, *where* $\mathfrak{f}_{ss}(L) = \sum_\alpha (ord(\alpha) - 1)$ *is computed from the eigenvalue orders of* F_{ss}, *all dividing* L. *The rank is 2 at the even* L *of the table below and 1 at odd* L.

Proof. F_{ss} has order L; its eigenvalues are L-th roots of unity. For $\gcd(L, q) = 1$, the L-th roots in $\bar{\mathbb{F}}_q$ are isomorphic to those in $\mathbb{C}$, so their orders match. □

L	$\mathfrak{f}_{\text{ss}}(L)$	$\mathfrak{f}_{\text{WD}}$	$\text{NM}^2 = 3L/4$
4	14	16	3.00
6	30	32	4.50
8	70	72	6.00
10	106	108	7.50
12	130	132	9.00

Both $\mathfrak{f}_{\text{ss}}$ and NM^2 are q-independent encodings of the same semisimple eigenvalue data — one via multiplicative orders (integers), one via fourth moments of magnitudes (reals).

Theorem B.3.4 (Additive decomposition over gauge irreps). *For the K-component wave equation with coupling matrix $M = diag(\mu_1 I_{n_1}, \dots, \mu_r I_{n_r})$:*

$$\mathfrak{f}_{\mathrm{WD}}(M) = \sum_{i=1}^{r} n_i \cdot \mathfrak{f}_{\mathrm{WD}}(\mu_i)$$

In particular, on the branch of SM Theorem 5, $K = 6$ with $M = \mu I_6$ gives $\mathfrak{f}_{WD} = 6\,\mathfrak{f}_{WD}(\mu)$; the multiplicities $(3, 2, 1)$ that determine $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ sit in the condensate Σ of SM Theorem 5, which does not enter the linear evolution matrix.

Proof. The multi-component evolution matrix is block-diagonal: the a-th component has its own $2L \times 2L$ block F_{μ_a} with eigenvalues depending only on μ_a. Both F_{ss} and N inherit the block-diagonal structure, so $\mathfrak{f}_{\mathrm{ss}}$ and $\mathrm{rank}(N)$ decompose additively. □

Verified for all $(\mu_c, \mu_w) \in \{1, \dots, q-1\}^2$ at $q \in \{3, 5, 7\}$ with $L = 4$.

The projection. The framework's trace-out operation (marginalization over the hidden sector) produces the reduced observable description — universally admitting the quantum representation — which depends on the coupling eigenvalues μ_k via the NM formula $\mathrm{NM}^2 = 3\langle\mu^4\rangle$ — a consequence of the stochastic-quantum correspondence applied to the wave equation's Fourier decomposition. These μ_k are properties of F_{ss}, the semisimple part of the dynamics. The nilpotent monodromy N contributes nothing to the emergent description: it affects only the off-diagonal Jordan block entries, which are erased by the coarse-graining over the hidden sector.

The trace-out therefore performs the Jordan-Chevalley projection

$$(F_{\mathrm{ss}}, N) \mapsto F_{\mathrm{ss}} \mapsto \{\mu_k\} \mapsto \mathrm{NM}^2$$

extracting the semisimple part, encoding it via magnitudes rather than orders, and organizing it by the representation theory of the partition. The nilpotent monodromy — genuine mathematical structure present in the full dynamics — is invisible to the embedded observer. This is the precise sense in which the framework's content holds that the observable physics is the *semisimple shadow* of the underlying mathematics: the observer sees only the diagonalizable spectral data, projected by the trace-out and organized by the gauge group's representation structure.

B.4 The gap equation for $\hbar$

Main-text reference: Chapter 7 §7.3.

The framework derives the relation $\hbar = c^3\epsilon^2/(4G\sigma)$ between Planck's constant and the discreteness scale by matching the classical horizon temperature to the KMS temperature of the emergent quantum field theory, conditional on H-slope with the horizon and frame conditions of Chapter 7 §7.3; with ϵ not

independently known, it fixes only $\epsilon/\sqrt{\sigma} = 2\,l_p$ (Chapter 7 §7.4), not the value of $\hbar$. The derivation has four steps; each step is presented with its full content.

Step 1: Spatial locality. The classical Hamiltonian of the substratum is spatially local. The interaction decomposes as

$$H_{\text{tot}} = H_V + H_B + H_D + H_{VB} + H_{BD}$$

where V is the visible sector, B is the boundary layer (surface modes near the horizon), and D is the deep hidden sector (volumetric modes far from the horizon). There is no direct V-D coupling: a deep hidden-sector mode must propagate through the boundary layer B before influencing the visible sector. The coupling chain is $V \leftrightarrow B \leftrightarrow D$, not $V \leftrightarrow D$.

Lemma B.4.1 (Frozen deep sector). *Let Δ_D be the spectral gap of $H_D + H_{BD}$ restricted to the deep sector conditioned on a fixed boundary configuration. For times $t \ll 1/\Delta_D \leq \tau_B$:*

$$\|\varphi_t^D - \mathrm{id}\| = \mathcal{O}(\Delta_D t) = \mathcal{O}(t/\tau_B)$$

Proof. The spectral gap gives the inverse relaxation time of the slowest mode, so $\Delta_D \leq 1/\tau_B$ (with equality when the slowest mode dominates the relaxation). The deep-sector evolution over time t displaces any initial configuration by $\mathcal{O}(\Delta_D t)$ in phase space. □

Lemma B.4.2 (Factorization of transition probabilities). *The transition probability factorizes as*

$$T_{ij}(t) = T_{ij}^{(B)}(t) + \mathcal{O}(t/\tau_B)$$

where $T_{ij}^{(B)}(t)$ depends only on the V-B dynamics.

Proof. The transition probability is

$$T_{ij}(t) = \frac{1}{|\mathcal{C}_B||\mathcal{C}_D|} \sum_{b \in \mathcal{C}_B} \sum_{d \in \mathcal{C}_D} \delta_{x_j}[\pi_V(\varphi_t(x_i, b, d))]$$

By spatial locality, $\pi_V(\varphi_t(x_i, b, d))$ depends on d only through the back-reaction $B \leftarrow D$, which by Lemma B.4.1 shifts b by $\mathcal{O}(t/\tau_B)$. Expanding:

$$\pi_V(\varphi_t(x_i, b, d)) = \pi_V(\varphi_t^{VB}(x_i, b)) + \mathcal{O}(t/\tau_B)$$

where φ_t^{VB} is the flow restricted to $V \times B$ with D frozen. The d-sum then contributes $|\mathcal{C}_D|$ identical terms at leading order:

$$T_{ij}(t) = \underbrace{\frac{1}{|\mathcal{C}_B|} \sum_{b \in \mathcal{C}_B} \delta_{x_j}[\pi_V(\varphi_t^{VB}(x_i, b))]}_{T_{ij}^{(B)}(t)} + \mathcal{O}(t/\tau_B) \qquad \square$$

The correction is $\mathcal{O}(t/\tau_B) \sim 10^{-32}$ for laboratory processes. Since $T_{ij}(t)$ determines the emergent quantum description, and $T_{ij}^{(B)}(t)$ depends only on the V-B dynamics — which are characterized by the boundary geometry (A, ϵ, κ) and the constants c, G appearing in the classical Hamiltonian — the emergent action scale $\hbar$ inherits *boundary-only dependence*.

Step 2: Deep-sector gauge equivalence. A corollary of the factorization lemma is that, to within $\mathcal{O}(\tau_S/\tau_B)$, no quantity determined by the visible transition data depends on the cardinality $|\mathcal{C}_D|$ of the deep hidden sector: systems with the same $\mathcal{C}_V \times \mathcal{C}_B$ dynamics produce the same visible transition data, whether $|\mathcal{C}_D|$ is finite, countably infinite, or uncountably infinite. Quantities that differ between Hamiltonians compatible with those data, such as individual eigenvalues and raw amplitudes, lie outside the corollary, and so does recurrence-scale content, which the recurrence argument establishes on a finite representative. Within the leading-order visible transition data on accessible timescales, a finite and an infinite deep sector are indistinguishable.

Step 3: Dimensional determination. Step 2 excludes volumetric (deep-sector) quantities, leaving boundary quantities. The boundary carries both *local* geometric data (ϵ, κ, and the constants c, G of the classical Hamiltonian) and a *global* quantity: the total area A, which forms the dimensionless ratio $A/\epsilon^2 = S_{\mathrm{dS}}/\sigma$. If $\hbar$ depended on S_{dS}, it would be observer-dependent — different observers have different horizon areas — contradicting the universality of the emergent action scale. The surface gravity κ is excluded because it varies between observers and epochs while $\hbar$ is observed to be universal across observers and constant in time.

The unique combination of c, G, ϵ with dimensions of action is

$$\hbar = \beta \frac{c^3 \epsilon^2}{G}$$

with unknown dimensionless coefficient β.

Step 4: Thermal self-consistency. The classical substratum, with one boundary mode of entropy σ (in nats) per cell of area ϵ^2, assigns the partition boundary a temperature

$$k_B T_{\mathrm{cl}} = \frac{c^2 \epsilon^2 \kappa}{8\pi G \sigma}$$

containing no $\hbar$. This is computable entirely from classical-horizon thermodynamics with no reference to quantum mechanics.

On the exact de Sitter calibration, the emergent QFT of the framework's Part I content lives on a classical background that has a bifurcate Killing horizon with surface gravity κ (the bifurcate structure presupposes emergent local boost invariance at the horizon, a dependency carried in [GR §8.5]). Regularity of the Wick-rotated metric at the horizon requires Euclidean period $\beta = 2\pi c/\kappa$; any QFT on this background — including a lattice-regularized one — is periodic in

imaginary time with that period only under the H-Hawking conditions, giving a KMS state at temperature

$$T_Q = \frac{\hbar\kappa}{2\pi c k_B}$$

with $\hbar$ unknown. The periodicity holds under the H-Hawking conditions and the KMS identification requires the emergent state to belong to that class (H-spectrum), so the matching below is a conditional infrared calibration, not a theorem within the derived QFT ([GR §3.2]).

The two temperatures are computed independently — $T_{\rm cl}$ from the classical substratum alone (no QM), T_Q from the emergent QFT alone (no classical substratum details) — but they refer to the *same physical degrees of freedom*: the boundary modes $V \times B$ across which $H_{\rm int}$ couples the visible and hidden sectors. Step 1's boundary-only dependence shows that the visible transition data from which the emergent QFT is reconstructed are fixed (at leading order in τ_S/τ_B) by these same $V \times B$ dynamics. Temperature is a state property; two complete descriptions of the same modes cannot assign different temperatures without logical contradiction. The matching $T_{\rm cl} = T_Q$ identifies the classical inverse-energy temperature with the emergent KMS period on the same boundary modes $V \times B$; it holds under H-slope (Chapter 7 §7.3), which partition geometry does not supply, with corrections at $\mathcal{O}(\tau_S/\tau_B) \sim 10^{-32}$:

$$\frac{c^2\epsilon^2\kappa}{8\pi G\sigma} = \frac{\hbar\kappa}{2\pi c}$$

The surface gravity κ cancels — a non-trivial cross-check on Step 3, which excluded κ from $\hbar$ on observer-universality grounds. Solving:

$$\boxed{\hbar = \frac{c^3\epsilon^2}{4G\sigma}}$$

The derivation is a gap equation: ϵ is the free geometric input, $\hbar$ is the output. The non-circularity is structural: Part I's content establishes that a QFT emerges with *some* action scale $\hbar$; the gap equation determines *which* $\hbar$, using the independent classical temperature that Part I neither requires nor produces.

The KMS periodicity is a *geometric* condition (Euclidean smoothness at the horizon), not a quantum condition; it does not assume $\hbar$. The $\hbar$ enters only through the emergent QFT's *interpretation* of this periodicity as a temperature. The matching is therefore between a geometric quantity (periodicity) and an emergent quantity (temperature), not between two quantities that both assume $\hbar$.

B.5 Anomaly cancellation and unique hypercharges

Main-text reference: Chapter 6 §6.7.

The framework's content on the Standard Model matter content derives the unique hypercharge assignment on the nonzero-Y_Q branch from the six anomaly conditions of $\mathrm{SU}(3)\times\mathrm{SU}(2)\times\mathrm{U}(1)$. The derivation establishes that the observed hypercharges are not phenomenological inputs but the unique anomaly-free completion on the nonzero-Y_Q branch of the framework's gauge structure.

Theorem B.5.1 (Unique hypercharges). *Given* $\mathrm{SU}(3)\times\mathrm{SU}(2)\times\mathrm{U}(1)$ *with fermions in fundamental or singlet representations, the six anomaly conditions determine the hypercharges on the nonzero-Y_Q branch uniquely up to the overall normalization and the interchange $Y_u \leftrightarrow Y_d$, the latter fixed by the electric-charge identification $Q_{em} = T_3 + Y$:*

$$Y_Q = \frac{1}{6}, \quad Y_u = \frac{2}{3}, \quad Y_d = -\frac{1}{3}, \quad Y_L = -\frac{1}{2}, \quad Y_e = -1$$

Proof setup. The Standard Model's chiral fermion content per generation consists of the left-handed quark doublet $Q = (u_L, d_L)$ transforming as $(\mathbf{3}, \mathbf{2})$, the right-handed up quark u_R as $(\mathbf{3}, \mathbf{1})$, the right-handed down quark d_R as $(\mathbf{3}, \mathbf{1})$, the left-handed lepton doublet $L = (\nu_L, e_L)$ as $(\mathbf{1}, \mathbf{2})$, and the right-handed electron e_R as $(\mathbf{1}, \mathbf{1})$. Each carries a hypercharge Y to be determined.

The six anomaly conditions for a chiral $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ theory are:

(A1) $\mathrm{SU}(3)^3$ anomaly: $\sum_{\text{triplets}} 1 - \sum_{\text{anti-triplets}} 1 = 0$ (A2) $\mathrm{SU}(2)^3$ anomaly: automatically zero ($\mathrm{SU}(2)$ is anomaly-free) (A3) $\mathrm{SU}(3)^2\mathrm{U}(1)$ anomaly: $\sum_{\text{triplets}} Y - \sum_{\text{anti-triplets}} Y = 0$ (A4) $\mathrm{SU}(2)^2\mathrm{U}(1)$ anomaly: $\sum_{\text{doublets}} Y = 0$ (A5) $\mathrm{U}(1)^3$ anomaly: $\sum Y^3 = 0$ (A6) Gravitational-U(1) anomaly: $\sum Y = 0$

In the framework's identification (Chapter 5), SU(3) is vector-like and SU(2) is chiral, with u_R and d_R treated as right-handed components (or equivalently, their conjugates u_R^c, d_R^c treated as left-handed with conjugated representations).

Setting up the anomaly equations. With the standard chirality convention (all fermions in the left-handed Weyl representation, with right-handed states represented by left-handed conjugates), the six conditions become explicit algebraic equations:

(A1) is automatically satisfied: SU(3) is vector-like with Q (triplet) balanced by u_R^c and d_R^c (anti-triplets, with the same count).

(A3) becomes: $2Y_Q - Y_u - Y_d = 0$ (the doublet Q counts twice for the $\mathrm{SU}(3)^2$ trace, u_R^c and d_R^c contribute opposite hypercharges due to conjugation).

(A4) becomes: $3Y_Q + Y_L = 0$ (the quark doublet Q has three colors).

(A5) becomes: $6Y_Q^3 - 3Y_u^3 - 3Y_d^3 + 2Y_L^3 - Y_e^3 = 0$ (factor of 6 for Q: 3 colors $\times$ 2 doublet states; factor of 3 for u_R, d_R: 3 colors; factor of 2 for L: doublet; factor of 1 for e_R).

(A6) becomes: $6Y_Q - 3Y_u - 3Y_d + 2Y_L - Y_e = 0$.

Solving the system. From (A4): $Y_L = -3Y_Q$. From (A3): $Y_u + Y_d = 2Y_Q$.

Substituting into (A6): $6Y_Q - 3(Y_u + Y_d) + 2Y_L - Y_e = 6Y_Q - 6Y_Q - 6Y_Q - Y_e = -6Y_Q - Y_e = 0$, so $Y_e = -6Y_Q$.

From the electroweak completion of L: the doublet L contains ν_L (electrically neutral) and e_L (charge -1). The charge formula $Q_{\text{em}} = T_3 + Y$ gives $Q_{\text{em}}(\nu_L) = +1/2 + Y_L$, requiring $Y_L = -1/2$ for ν_L to be neutral. Therefore $Y_L = -1/2$ and $Y_Q = 1/6$.

This gives $Y_u + Y_d = 1/3$ and $Y_e = -1$.

To separate Y_u and Y_d, the $\mathrm{U}(1)^3$ anomaly (A5) provides the remaining constraint. Substituting $Y_Q = 1/6$, $Y_L = -1/2$, $Y_e = -1$:

$$6(1/6)^3 - 3Y_u^3 - 3Y_d^3 + 2(-1/2)^3 - (-1)^3 = 0$$

$$6 \cdot 1/216 - 3Y_u^3 - 3Y_d^3 - 2/8 + 1 = 0$$

$$1/36 - 3(Y_u^3 + Y_d^3) - 1/4 + 1 = 0$$

$$3(Y_u^3 + Y_d^3) = 1/36 + 3/4 = 1/36 + 27/36 = 28/36 = 7/9$$

$$Y_u^3 + Y_d^3 = 7/27$$

Combined with $Y_u + Y_d = 1/3$: using $Y_u^3 + Y_d^3 = (Y_u + Y_d)^3 - 3Y_uY_d(Y_u + Y_d)$, we get $7/27 = 1/27 - Y_uY_d$, so $Y_uY_d = 1/27 - 7/27 = -6/27 = -2/9$.

The roots of $x^2 - (1/3)x - 2/9 = 0$ are $x = (1/3 \pm \sqrt{1/9 + 8/9})/2 = (1/3 \pm 1)/2$. The two roots are $2/3$ and $-1/3$. The anomaly conditions are symmetric under $Y_u \leftrightarrow Y_d$, so the system does not itself order the pair; the assignment $Y_u = 2/3$, $Y_d = -1/3$ is fixed by the electric-charge identification $Q_{\text{em}} = T_3 + Y$ for the up-type quark.

Conclusion. The complete hypercharge assignment is

$$Y_Q = 1/6, \quad Y_u = 2/3, \quad Y_d = -1/3, \quad Y_L = -1/2, \quad Y_e = -1$$

determined by the six anomaly conditions with no free parameters beyond the overall normalization (fixed above by the neutrality of ν_L) and the $Y_u \leftrightarrow Y_d$ root labeling (fixed by the electric-charge identification). □

Corollary. On the nonzero-Y_Q branch, $|q_p| = |q_e|$ is a *theorem*, not a coincidence.

Proton charge: $q_p = 2q_u + q_d = 2(2/3) + (-1/3) = 4/3 - 1/3 = 1$. Electron charge: $q_e = Y_e + T_3 = -1 + 0 = -1$ (since e_R is a singlet, $T_3 = 0$). So $|q_p| = |q_e| = 1$, with the equality forced by anomaly cancellation. This is one of the framework's distinctive structural predictions: the proton-electron charge equality, which is a free parameter in the Standard Model, is a theorem of the framework on that branch.

B.6 Gauge-theoretic derivations

Main-text reference: Chapter 5 §5.3-5.5.

Two key gauge-theoretic results from Chapter 5 are presented here in full: the conditional link-count proposition giving $K = 2dm$ internal components per site, and the cubic-rotation-group decomposition fixing the block multiplicities at $(3, 2, 1)$.

The coupling-degree minimization. The framework's substratum is a deterministic bijection on a $d = 3$ spatial cubic lattice. The site-internal structure is described by K scalar components per site. The minimal K supporting a chiral gauge structure is the structural question. Coupling-degree minimization does not answer it: the argument below is stated and then shown not to close, and the surviving content is the conditional link-count proposition together with the exact representation theorem that follows it.

Why coupling-degree minimization does not fix K. The natural argument runs: the discrete Laplacian on a d-dimensional cubic lattice couples $2d$ nearest neighbours, so the internal space needs one component per direction and $K \geq 2d$. The step fails. Diagonalizing the internal matrix decouples its eigenvectors, but it does not assign an eigenvector to a single signed spatial neighbour: in the displayed wave operator every internal component receives the WHOLE neighbour sum, so nothing forces a component-per-direction correspondence. A smaller K is not excluded by this argument, and the $\delta(2d) = 1$ substitution that would supply the correspondence is an operator replacement rather than a consequence of the dynamics.

Proposition B.6.1 ($K = 2dm$, conditional on H-link). *Let $V_{\text{link}} = \text{span}\{e_{+1}, e_{-1}, \ldots, e_{+d}, e_{-d}\}$ be the space spanned by the $2d$ signed lattice links. If the physical internal space satisfies $V_{\text{physical}} \cong V_{\text{link}}^{\oplus m}$ (hypothesis H-link), then $K = 2dm$, and the single-copy case $m = 1$ gives $K = 2d$, hence $K = 6$ at $d = 3$.*

Proof. Immediate from $\dim V_{\text{link}} = 2d$. □

The content of the proposition is entirely in its hypothesis. H-link identifies the geometric link space with the physical carrier, and $m = 1$ asserts a single copy; neither is derived here, and $K = 6$ is a consequence of those premises rather than a prediction of the wave operator. What is unconditional is the representation theorem that follows, which is a statement about the six geometric links themselves and holds whether or not they are the carrier.

The cubic rotation group decomposition. The cubic rotation group O_h (the symmetry of the cubic lattice) acts on the $K = 6$ internal components. The decomposition of this six-dimensional representation into irreducible representations of O_h determines the block multiplicities $(n_1, n_2, n_3, \ldots)$ of an equivariant condensate Σ (SM Theorem 5; the coupling matrix itself is scalar).

Theorem B.6.2 (Cubic decomposition). *The six-dimensional representa-*

tion of O_h on the internal components decomposes uniquely as

$$\mathbf{6} = \mathbf{3} \oplus \mathbf{2} \oplus \mathbf{1}$$

where **3** *is the standard triplet (vector representation T_1),* **2** *is the doublet (representation E), and* **1** *is the singlet (A_1).*

Proof. The cubic rotation group O has irreducible representations A_1 (singlet, 1D), A_2 (1D), E (2D), T_1 (3D), T_2 (3D). The cubic-character-table decomposition of the six-dimensional space of internal components requires the natural geometric structure of the components.

The internal components transform under O according to their geometric meaning. In the framework's content, the six components are organized as: three components transforming as the vector representation T_1 (one component per spatial axis, $T_1 = \mathbf{3}$), two components transforming as the doublet representation E (the "two-fold direction" structures characteristic of the cubic group, $E = \mathbf{2}$), and one component transforming as the singlet A_1 ($\mathbf{1}$).

The decomposition $\mathbf{6} = T_1 \oplus E \oplus A_1$ produces three blocks with multiplicities $(3, 2, 1)$, on which an equivariant condensate takes its block values. The framework's content identifies these multiplicities with the dimensions of the Standard Model gauge group factors: - $\mathbf{3} \to$ SU(3) color (three colors) - $\mathbf{2} \to$ SU(2) weak isospin (two doublet states) - $\mathbf{1} \to$ U(1) hypercharge (one singlet)

The identification is structural: the cubic rotation group's representation theory determines the internal structure, and the isotypic block multiplicities of the equivariant structure are the dimensions of the Standard Model gauge group factors. □

The link to gauge invariance. The commutant of the equivariant condensate Σ with distinct block values (SM Theorem 5; the coupling matrix $M = \mu I_6$ is scalar, with commutant U(6)) is the maximal group preserving its block structure: $\mathrm{U}(3) \times \mathrm{U}(2) \times \mathrm{U}(1)$, the group from which Chapter 5 §5.5 obtains $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$. Background independence — covariance of the dynamics under site-dependent choices of basis within each eigenspace, the link coupling transforming with them — promotes this global commutant to local gauge invariance. The result is the emergent $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ gauge theory of the Standard Model.

Combined structural result. Combining B.6.1 and B.6.2 gives the framework's account of the Standard Model gauge group, conditional on H-link at the B.6.1 step, unconditional at the representation-theoretic step (ii) of B.6.2, and conditional on H-cust at the stabilizer step (iii) (SM Theorem 5):

(i) The H-link link count with $m = 1$ on the $d = 3$ cubic lattice fixes $K = 6$ internal components per site.

(ii) The cubic rotation group's representation theory decomposes the six-dimensional internal space as $\mathbf{6} = \mathbf{3} \oplus \mathbf{2} \oplus \mathbf{1}$, fixing the block multiplicities at $(3, 2, 1)$.

(iii) The coupling matrix is scalar, $M = \mu I_6$, with commutant U(6); the stabilizer of an equivariant condensate Σ with distinct block values on these multiplicities is $\mathrm{U}(3) \times \mathrm{U}(2) \times \mathrm{U}(1)$ (SM Theorem 5), from which Chapter 5 §5.5 obtains $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$.

(iv) Background independence promotes the global commutant to local gauge invariance.

(v) Anomaly cancellation (Appendix B.5) uniquely determines the hypercharges on the nonzero-Y_Q branch.

(vi) Three degenerate sectors follow from the three triplet (T_1) staggered tastes related by cubic symmetry; their reading as physical fermion generations is under H-spin' (with H-χ' and light-spectrum selection likewise flagged).

The cumulative derivation produces the full Standard Model gauge structure with matter content from the framework's substratum-level commitments, with no phenomenological inputs beyond the substratum's spatial dimensionality $d = 3$.

B.6b Bravais-lattice uniqueness lemma

Main-text reference: Chapter 5 §5.5.

The framework's commitment to a simple cubic Bravais lattice at the substratum, established in Chapter 5 §§5.2-5.5 from the bijection structure and the link count under H-link, has a formal closure at the representation-theoretic level. Among the fourteen three-dimensional Bravais lattices, simple cubic is the *unique* one whose nearest-neighbor link representation under the lattice's point group decomposes into the multiplicities matching the Standard Model gauge group.

Lemma B.6b.1 (Bravais-lattice uniqueness). *Among the fourteen three-dimensional Bravais lattices, the simple cubic lattice (cP) is the unique one whose nearest-neighbor link representation under the lattice's point group decomposes as $A_1 \oplus E \oplus T_1$ with multiplicities $(1, 2, 3)$ matching the Standard Model gauge group* $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$.

Proof.

Step 1 (3D irreps required). The framework's commutant construction (Chapter 5 §5.4) requires the lattice's point group to admit a three-dimensional irreducible representation, supplying the SU(3) commutant factor. Among the 32 three-dimensional crystallographic point groups, the dimensions of irreducible representations are bounded as follows: triclinic (C_1, C_i) and monoclinic (C_2, C_s, C_{2h}) groups have all irreps one-dimensional; orthorhombic groups (C_{2v}, D_2, D_{2h}) have all irreps one-dimensional; tetragonal, trigonal, and hexagonal groups have maximum irrep dimension 2; only the five cubic-system point groups (chiral tetrahedral T of order 12; full tetrahedral T_d and T_h of order 24; chiral octahedral O of order 24; full octahedral O_h of order 48) admit three-dimensional irreps.

This rules out 27 of 32 point groups and 11 of 14 Bravais lattices: triclinic (aP), monoclinic (mP, mC), orthorhombic (oP, oC, oI, oF), tetragonal (tP, tI), trigonal/rhombohedral (hR), and hexagonal (hP) lattices all have holohedries with maximum irrep dimension at most 2. The 3-dimensional-irrep requirement isolates the cubic system but is not the tightest constraint, and it is not the binding one: the SU(2) factor requires a *genuine* 2-dimensional irreducible representation, which the chiral and pyritohedral tetrahedral groups T and T_h — whose apparent doublet is a complex-conjugate pair of 1-dimensional irreps with commutant $\mathrm{U}(1)^2$ rather than SU(2) — do not possess. Equivalently, the full multiplicity ladder $(3, 2, 1)$ is the irreducible-degree set $\{1, 2, 3\}$, realized among all finite subgroups of O(3) only by $S_4 \cong O$: cyclic and dihedral groups lack the 3; A_4 lacks a genuine 2; the icosahedral group A_5 has 3-dimensional irreps but no 2-dimensional one (and carries unwanted 4- and 5-dimensional irreps). The octahedral group O is the unique finite rotation group carrying both a genuine 2- and a 3-dimensional irrep, and the cubic lattice's holohedry $O_h \supset O$ supplies both. The binding constraint is therefore the SU(2), not the SU(3).

Step 2 (cubic lattices discriminated by NN decomposition). The three cubic Bravais lattices — simple cubic (cP, six nearest neighbors), body-centered cubic (cI, eight nearest neighbors), face-centered cubic (cF, twelve nearest neighbors) — all have full octahedral point-group symmetry O_h and therefore admit three-dimensional irreps. They are discriminated by the *decomposition of their nearest-neighbor representations* under the octahedral group O (with parity-grading distinctions absorbed into the choice of O versus O_h).

Character-table decomposition of nearest-neighbor permutation representations gives:

$$\text{cP (6 NN)} : \chi = (6, 0, 2, 2, 0) \Rightarrow A_1 \oplus E \oplus T_1,$$

with multiplicities $(1, 2, 3)$ matching $\mathrm{SU}(3)\times\mathrm{SU}(2)\times\mathrm{U}(1)$ after alphabet-freedom reduction.

$$\text{cI (8 NN)} : \chi = (8, 2, 0, 0, 0) \Rightarrow A_1 \oplus A_2 \oplus T_1 \oplus T_2,$$

with multiplicities $(1, 1, 3, 3)$, producing a commutant with two SU(3) factors and no SU(2) factor — incompatible with the Standard Model gauge group.

$$\text{cF (12 NN)} : \chi = (12, 0, 0, 0, 2) \Rightarrow A_1 \oplus E \oplus T_1 \oplus 2T_2,$$

with multiplicities $(1, 2, 3, 6)$, producing additional factor structure (the doubled T_2 multiplicity yields an extra $U(2)$ commutant acting on the multiplicity space) incompatible with the Standard Model.

Among the three cubic Bravais lattices, only cP produces the Standard Model gauge group multiplicities exactly.

Step 3 (the link count excludes BCC alternatives, given H-link). A potential ambiguity: BCC's six *next-nearest neighbors* (along the axial directions $\pm a\hat{e}_i$) produce a character identical to SC's nearest-neighbor character, and would

therefore decompose as $A_1 \oplus E \oplus T_1$ with the same SM multiplicities. The framework's nearest-neighbor restriction rules out this alternative — BCC's axial directions are next-nearest rather than nearest, and the nearest-neighbor restriction excludes them.

Step 4 (Bravais commitment excludes alternatives). The framework's substratum commitment to a Bravais lattice with single-atom basis (Chapter 5 §§5.2-5.3) excludes multi-atom-basis structures (such as the diamond structure, which would have additional point-group structure) and quasi-crystalline alternatives. These structures are not within the lattice class the framework's substratum derivation produces.

The four steps together establish the uniqueness claim. ■

Computational verification. The character-table decomposition in Step 2 admits direct verification: enumerating the action of the octahedral group O's 24 elements on the six nearest-neighbor sites of simple cubic produces the character $\chi = (6, 0, 2, 2, 0)$ on the five conjugacy classes $(E, 8C_3, 3C_2, 6C_4, 6C_2')$. Applying the orthogonality relation $n_\sigma = (1/|G|) \sum_C |C| \chi_\sigma(C)^* \chi(C)$ for the irrep characters χ_σ of A_1, A_2, E, T_1, T_2 gives multiplicities $(1, 0, 1, 1, 0)$, confirming the decomposition $A_1 \oplus E \oplus T_1$. Parallel computation for the BCC eight-NN character $(8, 2, 0, 0, 0)$ gives multiplicities $(1, 1, 0, 1, 1)$, confirming $A_1 \oplus A_2 \oplus T_1 \oplus T_2$. Parallel computation for the FCC twelve-NN character $(12, 0, 0, 0, 2)$ gives multiplicities $(1, 0, 1, 1, 2)$, confirming $A_1 \oplus E \oplus T_1 \oplus 2T_2$. The lemma's content is therefore checkable by direct finite computation; the proof above formalizes what the computation confirms.

Consequences. The lemma converts Chapter 5 §5.5's Bravais-lattice uniqueness claim from a structural argument to a formally closed theorem. The framework's substratum commitment to simple cubic is therefore not one structural choice among several admitting alternative resolutions — it is the unique choice among 3D Bravais lattices consistent with the Standard Model gauge group under the framework's commutant construction on nearest-neighbor links. The lemma's specialist-review surface is the character-table decomposition itself, which is verifiable independently in any standard reference on crystallographic point groups (e.g., Tinkham, *Group Theory and Quantum Mechanics*; Bradley and Cracknell, *The Mathematical Theory of Symmetry in Solids*).

B.6c T_1 generation-assignment uniqueness lemma

Main-text reference: Chapter 6 §6.2.

The framework's identification of three fermion generations with the T_1 triplet staggered taste (Chapter 6 §6.2), with the A_1 singlet taste hosting the composite Higgs, is a load-bearing structural commitment. This appendix establishes that the assignment is *uniquely forced* by the framework's other structural commitments — it is not one assignment among several alternatives, but the only one consistent with the substratum's existing constraints.

Lemma B.6c.1 (T_1 generation-assignment uniqueness). *Given the framework's commitments to (a) simple cubic Bravais lattice (Lemma B.6b.1), (b) $K = 2dm = 6$ internal components from the link count under H-link with $m = 1$ (Chapter 5 §5.4), (c) the staggered-Dirac reconstruction of spin from Brillouin-zone corner Gamma-matrix structure (Chapter 6 §6.2), (d) Standard Model anomaly cancellation (§B.5), and (e) the physical-spin reading of the taste labels (H-spin', SM §4.7), the assignment of the three fermion generations to the T_1 triplet (with the A_1 singlet hosting the composite Higgs) is the unique assignment consistent with these commitments.*

Proof.

Step 1 (Available irreps in the 4-taste decomposition). The Brillouin-zone corners on a $d = 3$ simple cubic lattice number $2^d = 8$ and pair under the corner-conjugation involution $\eta \leftrightarrow \mathbf{1} - \eta$ into 4 taste pairs (Chapter 6 §6.2). Two corners — $\Gamma = (0, 0, 0)$ and $R = (\pi, \pi, \pi)$ — are invariant under the octahedral rotation group O and pair into one taste transforming as A_1 (the trivial 1D irrep). The remaining six corners pair into three axis-aligned pairs (X_j, M_j) along the three coordinate directions, transforming under O as the components of the T_1 (vector, 3D) irrep through the signed spin-taste matrices γ^j (the unsigned pair labels alone carry only $A_1 \oplus E$). The 4-taste decomposition under O is therefore exactly $4 = A_1 \oplus T_1$, with no A_2, E, or T_2 components present. This restricts the available irreps for matter-content assignments to A_1 and T_1 only.

Step 2 (Spin-statistics constraint). The staggered-Dirac reconstruction (Chapter 6 §6.2) assigns Gamma-matrix structure to each Brillouin-zone corner. The Γ and R corners pair with Γ-matrices $\Gamma(\mathbf{0}) = I_4$ (scalar) and $\Gamma(R) = \gamma^1\gamma^2\gamma^3$ (pseudoscalar), both SO(3)-invariant, yielding total angular momentum $J = 0$ for the A_1 singlet taste. The j-th axis-aligned pair (X_j, M_j) has Gamma-matrices γ^j (vector) and $-i\Sigma_j$ (spin generator), yielding $J = 1/2$ for each component of the T_1 triplet. The label content (SM Theorem 9) thus assigns $J = 0$ labels to A_1 and $J = 1/2$ labels to T_1; their physical reading is H-spin' (hypothesis (e)), under which spin-statistics forces: bosonic (J-integer) fields must reside in the A_1 singlet; fermionic (J-half-integer) fields must reside in the T_1 triplet. Three fermion generations *cannot* be hosted by A_1 (which carries $J = 0$ and would produce bosons); they must reside in T_1.

Step 3 (Generation count fixed by T_1 dimension). The T_1 irrep has dimension 3 in the standard cubic-group character theory (Tinkham, *Group Theory and Quantum Mechanics*, Chapter 4). The three components transform among themselves under cubic rotations as the three coordinate axes; the cubic group O acts on them by signed permutation. Three generations, related by cubic symmetry, therefore emerge from a single structural commitment (T_1-housing of fermions), not from a free parameter. The generation count is not chosen from a continuum; it is forced by $\dim(T_1) = 3$.

Step 4 (Anomaly cancellation requires uniform gauge content). Standard Model anomaly cancellation (§B.5) operates generation-by-generation: each

fermion generation must independently satisfy the anomaly-cancellation equations on $SU(3)^3$, $SU(2)^2U(1)$, $U(1)^3$, mixed-gauge-gravitational, and global SU(2)-Witten anomalies. The anomaly-free hypercharge assignment (Appendix B.5) requires that all three generations carry *identical* gauge representations, differing only in their Yukawa couplings (i.e., in their masses). Cubic-group permutation of the three T_1 components automatically produces three identical copies — the three components carry identical gauge content because they are related by symmetry. Alternative assignments distributing fermions across different irreps with different gauge content would violate this uniformity requirement.

Step 5 (Alternatives excluded). Four alternatives are excluded by the steps above:

- *Three generations in $3A_1$ (three singlet copies):* requires three separate A_1 tastes, which the 4-taste decomposition does not provide (Step 1 gives one A_1, not three). Also violates spin-statistics (Step 2: A_1 produces $J = 0$ bosons).
- *Three generations in $A_1 \oplus E$ (1 + 2 = 3 components):* requires E in the decomposition, which is absent (Step 1: only $A_1 \oplus T_1$). Also, $A_1 \oplus E$ is reducible — cubic symmetry does not permute its components, so the three generations would not be related by symmetry, violating uniformity.
- *Three generations in T_2:* T_2 is absent from the 4-taste decomposition (Step 1). Alternative Bravais lattices (BCC, FCC) provide T_2 but fail Bravais-uniqueness (Lemma B.6b.1).
- *Generations distributed across multiple irreps with different gauge content:* violates uniformity required by anomaly cancellation (Step 4).

Therefore the unique assignment consistent with the framework's structural commitments is: A_1 hosts the composite Higgs (bosonic, $J = 0$); T_1 hosts the three fermion generations ($J = 1/2$, related by cubic permutation, with identical gauge content). ■

Computational verification. The lemma's content reduces to finite enumeration in two steps that can be verified directly. First, the 4-taste decomposition: enumerate the action of O's 24 elements on the 8 Brillouin-zone corners; verify that they form 4 orbits under corner-conjugation; verify that the orbits decompose as $A_1 \oplus T_1$ under O. Second, the spin-statistics step: enumerate the standard staggered-Dirac Gamma-matrix structure at each corner; verify that Γ and R produce $J = 0$ and the axial pairs produce $J = 1/2$. Both verifications are standard exercises in lattice-fermion representation theory (cf. Kogut and Susskind 1975; Sharpe and Patel 1994; Lee and Sharpe 1999).

Consequences. Lemma B.6c.1 fixes the framework's most consequential matter-content commitment given hypotheses (a)–(e). The three-generation pattern of the Standard Model, the bosonic character of the Higgs, the J = 0 versus J = 1/2 separation between Higgs and fermion sectors, and the uniformity of gauge content across generations are all forced by the existing

substrate commitments (simple cubic lattice, $K = 2d$ link count under H-link, staggered-Dirac reconstruction, anomaly cancellation) together with H-spin'. The framework's matter content is therefore not freely chosen to match observation — it is the unique matter content consistent with the prior structural commitments. The lemma's specialist-review surface is again the cubic-group character theory plus the standard staggered-Dirac reconstruction, both verifiable in independent references.

In combination with Lemma B.6b.1 (Bravais-lattice uniqueness), Lemma B.6c.1 establishes that the framework's full gauge-and-matter structure — simple cubic lattice → cubic group → $A_1 \oplus T_1$ taste decomposition → composite Higgs in A_1 + three fermion generations in T_1 — is the unique structure consistent with the substratum's commitments and the Standard Model gauge group.

B.7 The lemma chain for emergent quantum mechanics

Main-text reference: Chapter 1 §§1.5-1.9.

The framework's characterization theorem — accessible non-Markovianity $\Longleftrightarrow$ C1/C3/C4 realization per accessible horizon, the quantum representation internal and universal ($S \iff D \iff Q_{\text{fb}}$, [Main §3.4]) — relies on a chain of intermediate lemmas. This appendix collects these lemmas with compact proofs, providing a self-contained reference for the framework's logical structure.

Lemma B.7.1 (Partition relativity). *The same physical substratum admits different visible/hidden partitions, with each partition producing its own emergent description.*

Proof. The substratum (S, φ) with $S = \mathcal{C}_V \times \mathcal{C}_H$ admits alternative decompositions $S = \mathcal{C}_{V'} \times \mathcal{C}_{H'}$ for any reshaping of S into a product structure. Each such reshaping defines a different visible/hidden partition with its own visible-sector emergent dynamics. The emergent dynamics depends on the choice of partition; the underlying substratum dynamics φ does not. □

Lemma B.7.2 (Emergent stochasticity). *Any deterministic bijection on a product configuration space produces emergent stochastic dynamics on the visible sector after tracing over the hidden sector.*

Proof. The bijection $\varphi : \mathcal{C}_V \times \mathcal{C}_H \to \mathcal{C}_V \times \mathcal{C}_H$ produces transition probabilities $T_{ij} = (1/|\mathcal{C}_H|) \sum_k \delta_{x_j, \pi_V(\varphi(x_i, h_k))}$ on the visible sector. These transition probabilities are stochastic (non-deterministic) whenever the bijection produces multiple distinct visible-sector outputs from a single visible-sector input across different hidden-sector configurations. □

Lemma B.7.3 (P-indivisibility from a non-permutation one-step matrix). *If the visible one-step matrix $T = \Gamma(1)$ is not a permutation and the configuration space is finite, the emergent stochastic dynamics on the visible sector is P-indivisible at the recurrence scale.*

Proof. The hypothesis is the non-permutation witness on $T = \Gamma(1)$, together with finiteness of the configuration space. It is NOT a consequence of C1: C1 asserts non-zero coupling, which does not by itself force T off the permutation set ([Main §1.3]). C2 and C3 play no part here; they govern *observability*, below. A column-stochastic matrix that is not a permutation strictly contracts total-variation distance for some pair of basis states: $\mathrm{TV}(Te_i, Te_j) = 1$ exactly when columns i and j have disjoint support, so if *no* basis pair contracted, all columns would have pairwise-disjoint support and T would be a permutation. Hence there is a pair with $D(1) := \mathrm{TV}(\Gamma(1)e_i, \Gamma(1)e_j) < 1$. Because φ is a bijection of a finite set it has finite order ω (the least common multiple of its cycle lengths): $\varphi^\omega = \mathrm{id}$ *exactly*, so $\Gamma(\omega) = \mathbb{1}$ and $D(\omega) = \mathrm{TV}(e_i, e_j) = 1$. The trace distance therefore satisfies $D(0) = 1$, $D(1) < 1$, $D(\omega) = 1$: it decreases and then recovers. Since every stochastic propagator is a total-variation contraction, a P-divisible process has monotonically non-increasing D for every pair of inputs; the recovery $D(\omega) > D(1)$ violates this, so the process is P-indivisible. ■

Role of C2, C3, and observability. The recovery above is guaranteed only at the recurrence time ω, which can be astronomically large for a high-capacity hidden sector, so a non-permutation rooted map and finiteness establish P-indivisibility at the recurrence scale. On *observable* timescales $k\tau_S \ll \tau_B$ the relevant property is conditional memory: C4 (history readback) is the load-bearing condition for it (`review4_probes.py`), with C2 (memory persistence) keeping the record alive to readback and C3 (sufficient capacity) giving it room; this is the content of Lemma B.7.4 and Chapter 1 §1.5. This division of labor follows [Main]: the recurrence argument uses a non-permutation rooted map and finiteness, and readback delivers conditional memory on observable timescales, which does not by itself make the population law P-indivisible on those windows. □

Lemma B.7.4 (Accessible conditional memory from readback). *Suppose the visible process has a gap at order k in the accessible window, as C4 asserts: two visible histories of length k followed by a common present state, each of probability at least p_0, induce next-step laws at total-variation distance at least δ. Then $I(X_{<k}; X_{k+1} \mid X_k) \geq p_0\delta^2/\ln 2 > 0$* (Pinsker; [Main §2.3]).

Status. The persistence of this memory over windows $k\tau_S \ll \tau_B$ is borne out by explicit slow-bath models (e.g. a slowly-mixing bath with a coarse-grained readout, the natural discrete realization of C1–C4): the conditional past–future mutual information remains nonzero over $\mathcal{O}(\tau_B/\tau_S)$ visible steps before decaying, so C2 (memory persistence) and C3 (capacity, which bounds the storable history at $\log_2 m$ bits) keep that memory observable for $k\tau_S \ll \tau_B$; they do not make the recurrence-scale distinguishability recovery of Lemma B.7.3 observable on those windows. A closed-form bound of the type $I \geq I_0(1 - k\tau_S/\tau_B)$ holds only as a leading-order estimate: in explicit models the conditional mutual information is generally *non-monotonic* in the window size — rising above I_0 at intermediate windows before decaying — so the precise functional form is model-dependent rather than a universal inequality. What is robust, and all that the characterization theorem requires, is that the memory is $\mathcal{O}(I_0)$ and

observable throughout the regime $k\tau_S \ll \tau_B$. □

Lemma B.7.5 (Phase locking). *Assume dimension at least three, distinct gaps and non-vanishing overlaps for both Hamiltonians being compared, and, for at least one of them, the triangle-genericity condition: every relation $\omega_1 + \omega_2 = \omega_3$ among the gaps $\omega_{kl} = E_k - E_l$ of three different pairs of levels is a triangle identity $\omega_{kl} + \omega_{lm} = \omega_{km}$. The emergent effective Hamiltonian $\hat{H}_{\mathit{eff}}$ on the visible sector is then determined by the continuous-time transition data $\{T_{ij}(t)\}$ up to the lemma's full gauge: an overall energy shift, the diagonal-unitary rephasing freedom, and the antiunitary conjugation $\hat{H} \to -\hat{H}^*$ — a genuine twofold ambiguity rather than a convention ([Main §3.1]).*

Proof. The transition data $\{T_{ij}(t)\}$ determines the Born-rule probabilities $|\langle j|e^{-i\hat{H}_{\mathrm{eff}}t}|i\rangle|^2 = T_{ij}(t)$. The squared-modulus structure leaves the phase undetermined; specifically, the transformation $|i\rangle \to e^{i\theta_i}|i\rangle$ leaves all $|U_{ij}|^2$ invariant. This is the basis-convention freedom — the choice of phase for each basis state — which is physically trivial. The triangle-genericity condition identifies the pair of levels carrying each frequency, up to a relabelling and a global reflection of the levels ([Main §3.1]); this step is not supplied by Fourier extraction alone, since distinct gaps admit non-congruent spectra with equal gap sets (scope note to Theorem B.7.5c below). The remaining alternatives are this rephasing freedom, the energy shift and the antiunitary conjugation $\hat{H} \to -\hat{H}^*$ stated above. □

Scope. The lemma is stated in its simplest form, $T_{ij}(t) = |U_{ij}(t)|^2$ — the trivial-ancilla case of the general construction. In the ancilla-marginal form, local generic rigidity is checked at the certified block structures; general completeness is open. The visible data constrain the *dilated* Hamiltonian and configuration projectors, with the energy shift and diagonal-phase freedoms exactly invisible; the extension, its Fourier structure, and the constraint analysis of the larger pair-phase gauge are given in the companion paper ([Main §3.4]) — where the pair-phase gauge reduces to the diagonal one for generic families at each fixed dimension — together with the antiunitary conjugate $H \to -H^*$, retained there as an intrinsic ambiguity of transition data.

Bohr frequencies and the D-gauge classification. The proof of Lemma B.7.5 rests on Theorem B.7.5a and, without the triangle-genericity condition, on the further results collected here, each stated with its hypotheses ([GR §3.3]). Throughout, $U(t) = e^{-iHt}$ with H in frequency units and $H = V\,\mathrm{diag}(E)\,V^\dagger$ on a finite configuration space, so that V_{ia} is the overlap of configuration i with eigenvector a, and likewise $H' = W\,\mathrm{diag}(E')\,W^\dagger$; the antiunitary reflection $-\bar{H}$ is the $-\hat{H}^*$ of Lemma B.7.5.

Theorem B.7.5a (Bohr-frequency completeness). *Let $U(t) = e^{-iHt}$ with non-degenerate eigenvalues and non-vanishing configuration-basis overlaps. Then the transition probabilities determine the Bohr-frequency set $\{E_a - E_b\}$: each return probability expands as $|U_{ii}(t)|^2 = \sum_{a,b} |V_{ia}|^2 |V_{ib}|^2\, e^{-i(E_a - E_b)t}$, in*

which every gap value carries a strictly positive coefficient; every $|U_{ij}(t)|^2$ is a combination of the same frequencies; and consequently any U' of the same form with $|U'_{ij}(t)|^2 = |U_{ij}(t)|^2$ for all i, j, t has $\{E'_a - E'_b\} = \{E_a - E_b\}$.

Proof. Expanding $U_{ij}(t) = \sum_a V_{ia}\, e^{-iE_a t}\, V^*_{ja}$ gives $|U_{ij}(t)|^2 = \sum_{a,b} V_{ia} V^*_{ja} V^*_{ib} V_{jb}\, e^{-i(E_a - E_b)t}$; at $i = j$ the coefficients are $|V_{ia}|^2 |V_{ib}|^2 > 0$, and grouping terms by frequency value only adds positive quantities — degenerate gaps merge coefficients but cannot cancel them. Distinct characters $t \mapsto e^{-i\omega t}$ of $(\mathbb{R}, +)$ are linearly independent (Dedekind), so equality of two probability families forces equality of the per-frequency coefficients over the union of the two gap sets, and positivity makes the supports — the gap sets — coincide. □

The dimensional obstruction. Theorem B.7.5a is what the gap equation of §B.4 consumes, and it is less than a reconstruction of H. The unitary $U(t)$ is dimensionless, and $\hbar$ enters only through $\hat{H} = i\hbar\, \partial_t U\, U^\dagger$. The transition probabilities are dimensionless functions of t and determine temporal frequencies; converting a frequency to an energy requires the action scale $\hbar$ itself, so no amount of transition data fixes it. The residual ambiguities in passing from frequencies to a Hamiltonian carry no scale either: diagonal rephasing $H \mapsto DHD^\dagger$ moves no modulus, the energy origin $H \mapsto H + E_0$ contributes only the global phase $e^{-iE_0 t}$, the antiunitary reflection $H \mapsto -\bar{H}$ satisfies $e^{-i(-\bar{H})t} = \overline{e^{-iHt}}$ entrywise and so preserves every $|U_{ij}(t)|^2$ exactly, and Theorem B.7.5c below excludes further ambiguities conditional on its stated classification hypothesis. The thermal self-consistency of §B.4, Step 4, is therefore the mathematically obligatory entry point for the dimensionful input.

Lemma B.7.5b (Real-to-integer transfer). *Let E, E' be real spectra on n modes, each with distinct directed gaps — $E_b - E_a = E_d - E_c$ with $a \neq b$ and $c \neq d$ only when $(a, b) = (c, d)$ — and with equal gap sets. Then E' is, after relabeling its modes, a translate $E + E_0$ of E or a translate $-E + E_0$ of its reflection; or else there are relabelings of the two spectra and strictly increasing integer rulers F, F' on n marks, each with distinct positive differences, with equal difference sets, and neither a translate nor a reflection of the other, such that every integer gap identity $F'_d - F'_c = F_b - F_a$ with $a \neq b$ holds for the relabeled real spectra as well.*

The lemma assumes no classification hypothesis. It is proved in the framework's Lean formalization (kernel: `real_to_integer_surrogate` in `OIBridge/TurnpikeScopeTransfer.lean`), and it is the step through which a classification of integer rulers, such as H-BG below, applies to real spectra.

Theorem B.7.5c (D-gauge completeness, two-branch form). *Assume H-BG: every non-congruent pair of strictly ordered integer rulers with distinct positive differences and equal difference sets has six marks and, after relabeling, realizes either μ or its global reversal, where μ sends each ascending mark pair of $\{0, 1, 4, 10, 12, 17\}$ to the ascending mark pair of $\{0, 1, 8, 11, 13, 17\}$ with the same difference. The orientation is chosen once for the whole pair. Let H, H'*

act on the same finite configuration space of dimension $m \geq 3$, with distinct nonzero energy gaps and non-vanishing configuration-basis overlaps for both Hamiltonians. If $|U'_{ij}(t)|^2 = |U_{ij}(t)|^2$ for all i, j, t, then $H' = DHD^\dagger + E_0$ or $H' = -D\bar{H}D^\dagger + E_0$, for a time-independent diagonal unitary D and a real E_0.

Conditional proof. Equality of the probability families forces equality of the per-frequency coefficients of $|U_{ij}(t)|^2 = \sum_{a,b} V_{ia}V^*_{ja}V^*_{ib}V_{jb}\, e^{-i(E_a - E_b)t}$. Under the stated overlap and gap hypotheses, the spectra have equal difference sets and the coefficient matching induces a bijection of ordered mode pairs preserving directed gaps. A fixed correspondence can be transferred from real to integer spectra while preserving its gap identities, strict order, distinct differences and non-congruence (Lemma B.7.5b): its realization constraints are homogeneous rational-linear equalities, strict inequalities and inequations, and rational approximation in the equality subspace followed by denominator clearing preserves the finite list of strict conditions. H-BG then supplies a congruent pair or one of the two globally oriented exceptional correspondences. For congruent spectra, diagonal-line rigidity fixes the moduli and the phase-coboundary relation fixes the phases, giving $W_{i,\tau a} = d_i \beta_a V_{ia}$ with d, β unimodular. The column phases cancel in the reconstructed Hamiltonian. Reflection congruence applies this argument to $(\bar{V}, -E)$ and gives the second branch. For the positive exceptional correspondence μ, the coefficient identities force flat moduli and constrain the phase data to a graph of maximum clique three, excluding the six orthogonal rows required by unitarity. For the reversed correspondence, reversing the source mode pair conjugates its coefficient tensor; applying the same obstruction to the conjugated source eigenbasis excludes that orientation as well. Thus the two Hamiltonian branches follow conditional on H-BG. □

Scope. The complete comparison of H-BG with the published integer classification of Bekir and Golomb [A. Bekir and S. W. Golomb, IEEE Trans. Inform. Theory 53 (2007) 2864–2867], including all normalizations and exceptional orientations, is open. The unsigned formal premise `BGIntegerClassification` of the Lean formalization is false: the strictly ordered rulers $(0, 1, 4, 10, 12, 17)$ and $(0, 4, 6, 9, 16, 17)$ have distinct positive differences and equal difference sets, are neither translates nor reflections of one another, and satisfy the reversed exceptional correspondence while admitting no relabelings with the required positive orientation. The kernel implication `twoBranch_of_BGClassification` therefore does not establish reconstruction from the physical hypotheses. The example contradicts that formal premise, not the published classification. Bohr-frequency completeness, the real-to-integer transfer, congruent reconstruction and the positive six-mode obstruction are kernel-proved independently of it. The exclusion of the reversed orientation and the assembly of the two branches from H-BG are kernel-proved as well, so H-BG is the theorem's only unproved premise; under the triangle-genericity condition of Lemma B.7.5 the two branches follow without H-BG ([Main §3.1]). The listed ambiguities are exactly invisible: $H + E_0$ contributes only a global phase, while $-\bar{H}$ conjugates the propagator entrywise. The gap condition is used where coefficient matching induces a bijection of or-

dered mode pairs: with degenerate gaps one frequency's coefficient is a sum over several mode pairs. The C_4 ring of [SM] Proposition 20 has distinct eigenvalues, uniform overlaps and degenerate gaps; its conjugate $\bar{H}$ reproduces every $|U_{ij}(t)|^2$ and conjugates the loop product $H_{01}H_{12}H_{23}H_{30}$, which no diagonal conjugation changes. The ring therefore does not exhibit a failure, without the gap condition, of the two-branch form conditional on H-BG: $\bar{H}$ lies in the reflection branch, $\bar{H} = -D_s \bar{H} D_s^\dagger$ with $D_s = \mathrm{diag}(1, -1, 1, -1)$. The raw frequency set alone cannot distinguish the non-congruent homometric rulers; reconstruction uses coefficient-labelled frequency data.

Corollary B.7.5d (Operational antiunitary invariance). *Transposing the preparation, every operation, and the effect simultaneously — $\rho \mapsto \rho^T$, $\mathcal{G} \mapsto T \circ \mathcal{G} \circ T$, $E \mapsto E^T$ — preserves every finite circuit probability, for arbitrary operations and arbitrary finite sequences, hence for every fixed outcome branch of any adaptive instrument strategy; the transformation carries Kraus operations to Kraus operations with conjugated operators, preserves the trace normalization, and on unitary evolution is exactly the branch map $H \mapsto -\bar{H}$.*

Enlarging the data from transition probabilities to arbitrary preparations, instruments, adaptive finite circuits, and effects therefore cannot remove the second branch: at the purely operational level the two branches are not two empirically distinguishable theories but two antiunitarily related reconstructions of the same complete circuit statistics, and they become physically distinct only once an oriented structure is fixed ([GR §3.3]). (kernel: `circuit_invariance`, `string_invariance`, `transposeMap_kraus`, `kraus_normalization`, `unitary_channel_transpose` in `OIBridge/AntiunitaryInvariance.lean`)

Theorem B.7.6 (Characterization theorem). *In the form of [Main §3.4]: accessible non-Markovianity of the visible process is equivalent to embedded observation under C1, C3, C4 for some accessible order k on each finite accessible horizon $K \geq k+1$, with the readback at k (C2 outside the equivalence, carried by the physical-memory theorem (persistence form without a mixing hypothesis, slow-bath form conditional on it)); the unitary quantum representation is internal and universal — $S \iff D \iff Q_{\mathrm{fb}}$, the realization's permutation unitary, the compressed form under (T) — nontrivial at the recurrence scale where the dynamics is P-indivisible, with the fixed-$\hat{H}$ form constructive for realized processes:*

$$\begin{aligned} & \text{Accessible non-Markovianity on } V \\ \iff\ & \exists k\ \forall K \geq k+1 : \text{ embedded observation of } (S, \varphi) \text{ on horizon } K \\ & \text{with C1, C3, C4, readback at } k \end{aligned}$$

Proof structure. The forward direction (sufficiency) follows from Lemmas B.7.1-B.7.5 combined with the Stinespring construction (§B.2): under C1–C4, the substratum dynamics produces accessibly non-Markovian emergent stochastic dynamics on V (P-indivisible at the recurrence scale), which is structurally

equivalent to a unitary quantum dynamics on a Hilbert space via Stinespring dilation — and, at the observable-law level, exactly representable by the realization's own permutation unitary ([Main §3.4]).

The reverse direction (necessity) is per-condition (Appendix C §C.5): C1, C3, and C4 are individually necessary for any deterministic realization of a non-Markovian process, and C2 in its persistence form likewise, with its slow-bath form within the conditional-mixing class as a lower bound on the bath's mixing time; violating C1, C3, C4 or the persistence form of C2 destroys the emergent structure. The quantum representation attaches to processes in the correspondence's class.

The biconditional is the framework's main characterization result. □ **Coherent extension.** The equivalence holds between the operational, dilated, and fixed-basis descriptions and is exact at that level; it does not by itself determine a coherent extension of the fixed-basis theory, and at least one conservative coherent extension preserves the fixed-basis theory exactly as a quotient while adding coherent structure not fixed by the equivalence.

The lemma chain produces the framework's central content. The chain from Lemmas B.7.1-B.7.5 through Theorem B.7.6 establishes the framework's foundational claim: non-Markovian visible dynamics are the necessary description of any embedded observer satisfying C1–C4 — with the quantum representation internal and universal ($S \iff D \iff Q_{\mathrm{fb}}$), the compressed form via the imported correspondence under (T). The chain is constructive: each lemma is provable from the substratum's structural commitments without requiring quantum postulates as inputs, and the cumulative derivation yields the framework's emergent quantum description as theorem-backed structure — the memory-bearing statistics with their universal fixed-basis representation — rather than as separately-postulated quantum axioms.

The framework's full mathematical content. Combining the derivations of this appendix:

- B.2 establishes the substratum-to-quantum bridge via Stinespring dilation.
- B.3 establishes the algebraic structure of the trace-out as Jordan-Chevalley projection, organizing the framework's content on gauge structure.
- B.4 establishes the gap equation relating $\hbar$ to the discreteness scale through horizon thermodynamics, producing the framework's content on gravitational physics.
- B.5 establishes the unique anomaly-free hypercharge assignment on the nonzero-Y_Q branch, fixing the framework's content on Standard Model matter.
- B.6 establishes the gauge-theoretic derivation from cubic symmetry, fixing the framework's content on the SM gauge group structure.
- B.7 establishes the lemma chain for the characterization theorem, fixing the framework's foundational claim.

The combined derivations constitute the framework's full mathematical content as a derivation chain from the substratum-level structural commitments to the Standard Model, the gravitational sector, and the empirical predictions across the framework's domains. The appendix is therefore a complete technical reference for readers requiring the framework's mathematical content at full detail.

Appendix C

Common Objections and Framework Responses

C.1 What this appendix develops

The framework's content is unusual. It combines a substantial empirical record (twenty-two classified Standard Model retrodictions — two parameter-free structural and five conditional structural — seven gravitational-sector entries, seven neutrino-sector entries, and the wider-therapeutic-window retrodiction for Rett syndrome, Guy et al. 2007 predating the framework) with structural commitments that overlap with several distinct programs in foundations of physics. Readers approaching the framework from different starting points raise different objections; the framework's content engages each cluster of objections from a single structural foundation.

This appendix systematically engages the most frequent objections to the framework's content. The appendix is *not* a defensive document; it is a reference resource for readers who want to understand how the framework engages established results in foundations of physics, established no-go theorems, established interpretive positions, and established methodological concerns. Each objection is engaged at the level appropriate to its substance — technical objections receive technical responses, methodological objections receive methodological responses, and meta-objections (about credentials, scope, ambition) are addressed where they are scientifically substantive.

Twelve clusters of objections occupy the appendix.

Section C.2 engages objections about Bell's theorem and the broader hidden-variable program: the framework's relationship to local hidden-variable theories, the Bell branch the framework adopts (measurement independence retained, ontic parameter independence given up, operational no-signaling preserved; hidden memory and indivisibility are not the mechanism), and why the framework is not a superdeterministic or Bohmian theory. Section C.3 engages the no-go theorems specifically — PBR, Frauchiger-Renner, Colbeck-Renner, Bong-Wiseman — and identifies which assumptions each makes that the framework's content satisfies or violates.

Sections C.4-C.6 engage the foundational concerns: the role of the stochastic-quantum bridge (Barandes route), the necessity direction of the characterization theorem (whether QM requires embedded observation), and finiteness (whether the framework's Axiom 2 is too strong).

Section C.7 engages computational implementability concerns — specifically Wolpert's theorem on physical computability and how the framework's content relates to it. Section C.8 engages the framework's relationship to standard interpretations of quantum mechanics: Many-Worlds (MWI), Bohmian mechanics, and QBism.

Sections C.9-C.11 engage framework-specific concerns: the substratum-emergent operator distinction (whether the framework's distinction between substratum and emergent operators is a substantive structural feature or just renaming) together with the parallel question whether observational incompleteness is the uncertainty principle renamed (C.9.1), circularity concerns (particularly around the gap equation for $\hbar$), and the framework's epistemic status as a structural framework rather than a competing physical theory.

Section C.12 engages meta-objections: credentials, scope, and the framework's ambition. The appendix closes with a brief discussion of which objections the framework cannot address (acknowledging genuine limits of the framework's reach) and which objections turn out to be category errors (where the framework's structural commitments dissolve apparent puzzles by relocating them).

The appendix is therefore both *substantive* (engaging specific technical objections with specific responses) and *honest* (acknowledging where the framework's reach stops and where genuine open questions remain). Readers approaching the framework from foundations-of-physics, from quantum information, from interpretive philosophy of physics, from cognitive science, or from clinical biology should each find their characteristic objections engaged here.

C.2 Bell's theorem and the framework's relationship to hidden-variable theories

The most frequent objection to deterministic-substratum theories is: "Bell's theorem rules out hidden-variable theories. The framework has a deterministic substratum and a hidden sector. Therefore Bell's theorem rules it out."

The objection. Bell's theorem (1964) establishes that no local hidden-variable theory can reproduce the quantum-mechanical predictions for entangled systems. The empirical violations of Bell inequalities (CHSH measured to ≥ 2.4 in multiple experiments, with quantum prediction at $2\sqrt{2} \approx 2.83$) are taken as definitively excluding local hidden-variable theories. The framework's deterministic substratum looks suspiciously like a hidden-variable theory; therefore the framework should be ruled out by Bell's theorem in the same way.

The framework's response. Bell applies directly to the response-complete deterministic substratum. If the Bell-relevant coupling range is finite at the

wings' separation, the setting interventions, applied at one step, each read and write only their own wing and each wing's readout lies outside the other wing's graph cone, every run records a ± 1 outcome, and one setting-independent pre-setting ensemble is used, then the hidden responses factorize and $|S_{\mathrm{CHSH}}| \leq 2$. OI therefore cannot combine deterministic response completeness, a ± 1 outcome in every run, measurement independence, and ontic parameter independence and still reproduce quantum Bell violation. The framework keeps measurement independence and takes the remaining horn: **ontic parameter dependence** for Bell-violating substrata, with operational no-signaling retained (Chapter 1 §1.7; Main §3.3).

The cited causal-local **stochastic** indivisible construction separately obeys the Tsirelson bound $2\sqrt{2}$. By the Bell ceiling theorem (Main §3.3), a response-complete deterministic completion with one-step own-wing setting interventions, each wing's readout outside the other wing's graph cone, a ± 1 outcome in every run, and one setting-independent pre-setting ensemble is a local hidden-variable model, so $|S_{\mathrm{CHSH}}| \leq 2$. The framework therefore adopts ontic **parameter dependence**, targeted in the SM/GR branch through preparation-indexed structure in the state-dependent graph $G(x)$ (or an equivalent ontically nonlocal response) under the open hypothesis H-Bell, with operational no-signaling required. If measurement independence were dropped instead, $S_{\mathrm{CHSH}} \leq \min\{4,\, 2+4\sqrt{2 \ln 2\, I_{\mathrm{ont}}}\}$ gives the model-independent necessary lower bound $I_{\mathrm{ont}} \geq 0.0309$ bits for the quantum value. Hall–Branciard's optimized causal CHSH model uses about 0.080 bits; the smaller ~ 0.046-bit benchmark is retrocausal. Dropping the spatial/intervention restriction permits the general finite OI response construction to reach PR-box behavior; PR is therefore a negative control for causal-local selection, not for the existence of a Hilbert/Born representation.

Not superdeterminism. Determinism alone does not establish Bell measurement independence, because setting generators are physical degrees of freedom. The manuscript therefore treats measurement independence as an explicit premise of the adopted Bell branch rather than as a theorem of the substratum. It does **not** use setting–hidden correlations to explain the observed violation. A mixing/decorrelation argument may motivate why laboratory setting generators approximate that premise, but the Bell mechanism itself is ontic parameter dependence. If the premise were dropped instead, that would be the measurement-dependent/superdeterministic horn quantified by the I_{ont} bound — a horn the framework records and declines.

No operational faster-than-light signal — but ontically parameter-dependent. The reference graph still supplies exact causal cones for its finite-range couplings, but a Bell-violating preparation may add preparation-indexed adjacency in the state-dependent graph $G(x)$ (or an equivalent nonlocal ontic response), so a response at one wing may depend on the remote setting. That is Bell nonlocality at the ontic parameter level and the manuscript names it as such. The distinct empirical claim is no-signaling: those remote-setting dependencies

cannot be controlled to transmit a message. The framework therefore does not advertise Bell violation as dynamically local in the hidden-variable sense.

Not Bohmian mechanics. Bohmian mechanics adds a pilot wave guiding particle trajectories in configuration space. The framework has no pilot wave and no guidance equation in configuration space. The framework's hidden sector is the substratum's broader configuration space, not a separate guiding field; the dynamics is direct (the bijection φ) rather than mediated by a guidance equation.

The framework is therefore not a Bell-local hidden-variable theory. Its deterministic completion is measurement-independent by the adopted Bell premise but ontically parameter-dependent, with operational no-signaling required. Hidden predictive memory and indivisibility remain central to the OI dynamics, but they are **not** the mechanism that evades Bell: a measurement-independent, parameter-independent deterministic completion recording a ± 1 outcome in every run would still satisfy CHSH ≤ 2. The Tsirelson value is retained for the cited stochastic causal-local indivisible construction and, conditionally, for any successful operational lift to standard quantum mechanics (§C.2).

C.3 The no-go theorems: PBR, Frauchiger-Renner, Colbeck-Renner, Bong-Wiseman

Beyond Bell's theorem, a family of more recent no-go theorems further constrains the class of ontologies compatible with quantum mechanics. The framework's content must engage each one specifically. The pattern is consistent: each no-go theorem makes specific assumptions, and the framework's content satisfies or violates each assumption in identifiable ways.

The PBR theorem. Pusey, Barrett, and Rudolph (2012) proved that any ψ-epistemic interpretation of quantum mechanics — where the wave function represents incomplete information about an underlying ontic state rather than the ontic state itself — must satisfy specific constraints, with several natural ψ-epistemic models ruled out. The theorem is widely interpreted as establishing the wave function as ontic rather than epistemic.

The framework's response must keep the premise accounting explicit. PBR assumes preparation independence — that independently prepared systems have a product joint distribution over their ontic states. Bare OI does not impose that postulate: co-embedded preparation devices may share preparation-relevant hidden or boundary data even when their later dynamical causal cones are disjoint. Hence PBR cannot be applied to OI merely from the finite-representation theorem. Equally, graph locality does not itself prove that OI violates preparation independence, does not supply a unique conditional product law, and does not derive the standard quantum tensor-product composite. Those questions belong to the coherent composite operational lift. The correct status is therefore conditional: any OI completion satisfying PBR's preparation-independence hypothesis inherits the PBR conclusion; a completion that demonstrably violates

it lies outside that theorem's scope.

The Frauchiger-Renner theorem. Frauchiger and Renner (2018) derive a contradiction for a specified multi-agent protocol under three assumptions: quantum probability-one certainty, consistency of the agents' certainty inferences, and single outcomes, denoted (Q), (C) and (S). The theorem identifies a conflict among those assumptions for that protocol, rather than assuming that a wave function is an ontic state.

The framework's ordinary Wigner's-friend account distinguishes definite substratum configurations from observer-relative reduced descriptions. Different agents may have different information about the same configuration. Main §4.1 also provides a joint law whose marginals are the two observers' laws, so the multiplicity of partitions does not show that a common law is absent.

A Frauchiger-Renner application requires the proposed completion's operational probabilities for the whole protocol and an analysis of its probability-one, consistency and single-outcome premises. The observer-relative interpretation alone does not prove that one of these fails and is not an unconditional resolution of the theorem.

The Colbeck-Renner theorem. Colbeck and Renner (2011, 2017) proved that no extension of quantum mechanics can provide additional predictive power beyond standard QM, under assumptions including free choice of measurement settings and the validity of QM as the empirically successful theory. The result is interpreted as showing that hidden-variable theories cannot improve on QM's predictive content.

For an operational completion already reproducing standard quantum predictions in the specified regime, its predictive content must be compared with the Colbeck-Renner hypotheses in that same regime. Substratum descriptions do not by themselves establish an exception to the theorem's free-choice and quantum-prediction assumptions.

A proposed departure from standard quantum predictions requires its own operational model and premise analysis. Placing its motivation at the substratum level does not show that the Colbeck-Renner hypotheses fail, or certify a new predictive advantage. The proposed regimes discussed in Chapters 15 and 18 retain their separate assumptions and evidential status.

The Bong-Wiseman extended-Wigner-friend theorem. Bong et al. (2020) extended Wigner's friend scenarios with additional assumptions (local friendliness, absoluteness of observed events) to derive inequalities that quantum mechanics can violate. The result is interpreted as showing that even more conservative attempts to maintain observer-independent reality face obstacles.

A deterministic substratum has one configuration trajectory from a specified initial state. That ontic fact must be distinguished from the operational assumptions about observed events in an extended Wigner's-friend protocol. Observer-

relative information alone does not establish which local-friendliness premise a proposed completion violates.

An operational completion with the standard quantum instruments inherits their protocol predictions, including any violation of the relevant inequalities. Whether the proposed OI physical completion supplies those instruments and which local-friendliness assumption it fails require separate checks. The fixed-basis correspondence and the partition interpretation alone supply neither conclusion.

Separate no-go premises. The substratum, observable laws and operational descriptions identify different objects constrained by these theorems. The trace-out does not automatically invalidate their assumptions. Bell concerns local responses and the setting ensemble; PBR concerns preparation independence; Frauchiger-Renner concerns probability-one inferences, consistency and single outcomes; Colbeck-Renner and extended-Wigner-friend results retain their own protocol assumptions.

Bare OI does not impose PBR preparation independence, but shared boundary data do not by themselves prove its failure or a conditional product law. Partition-relative descriptions provide an interpretation of observer information, not an automatic Frauchiger-Renner resolution. Any proposed completion satisfying a no-go theorem's hypotheses inherits its conclusion. Exact fixed-basis representation and a full operational completion remain distinct, with the latter characterized only within its stated matrix setting and additional completion conditions.

C.4 The stochastic-quantum bridge: Barandes and Stinespring

The framework's characterization theorem derives accessible non-Markovianity from C1–C4 — with quantum mechanics attaching to processes in the correspondence's class via a *stochastic-quantum bridge* — a mathematical correspondence between non-Markovian stochastic processes and unitary quantum dynamics. The bridge has multiple constructions in the recent literature. The framework's content uses Barandes' 2023 indivisible-stochastic-process bridge as one route, with the framework's own permutation dilation supplying the same transition-statistics layer through Stinespring — shared bedrock rather than an independent alternative.

The objection. "The framework relies on Barandes' stochastic-quantum correspondence, which is recent (2023) and not yet widely accepted. This undermines the main result. A theory built on a not-yet-established result is itself not established."

The framework's response. The framework's own permutation dilation independently supplies a transition-statistics representation, so the dependency on any single recent result is weaker than it appears — though the two routes

share the Stinespring bedrock and neither is fully independent of it — but neither route reaches the operational instrument algebra, which remains open ([Main §3.4]), so "either alone sufficient" would overstate what is established.

Route 1: Barandes correspondence. Barandes (2023) and subsequent work (Barandes 2024, Liu 2024, others) established a mathematical correspondence between indivisible stochastic processes and unitary quantum dynamics. The correspondence is peer-reviewed and corroborated by multiple independent groups. The framework uses this correspondence as one route to the characterization theorem.

Route 2: Stinespring dilation. The framework also reaches the representation layer via Stinespring dilation — a standard result from textbook operator algebra established in the 1950s. The dilation theorem provides a representation of any completely positive trace-preserving map as a unitary evolution on a larger Hilbert space; the framework's substratum-emergent partition is structurally identical to such a dilation. The Stinespring route uses only textbook operator algebra with no reference to Barandes' work.

Rejecting the stochastic-quantum correspondence does not by itself remove the transition-statistics layer, which the framework's own dilation supplies through standard operator algebra of the past seventy years. The framework's content is therefore robust against rejection of the Barandes correspondence specifically.

Beyond the bridge. The framework's content does not depend exclusively on the stochastic-quantum bridge. The P-indivisibility proof (Chapter 1), the partition-relativity lemma (Chapter 1), and the emergent stochasticity result (Chapter 1) invoke neither the Barandes correspondence nor the Stinespring dilation directly — they establish the framework's structural content at the substratum level, with the bridge then providing the connection to standard quantum mechanics at the emergent level.

A referee who rejects all routes to the bridge would need to also reject the framework's substratum-level content separately. The framework's content is therefore robust against multiple lines of foundational rejection, not because the bridge is dispensable but because the substratum-level content is logically prior to the bridge.

C.5 The necessity direction of the characterization theorem

A second frequent foundational objection concerns the *necessity* direction of the characterization theorem — the claim that the observed non-Markovian dynamics *require* embedded observation with C1–C4, not just that embedded observation with C1–C4 *produces* them.

The objection. "You show embedded observation produces the memory-bearing non-Markovian dynamics, but you haven't shown those dynamics require embedded observation. The characterization theorem's converse is the extraordinary claim. Without it, you have a sufficient condition but not

a necessary one — and the framework's structural claim (that the observed dynamics arise *from* embedded observation) requires both directions."

The framework's response. The necessity direction is proved per condition: C1, C3, and C4 unconditionally for any realization, C2 in its persistence form likewise and, within the conditional-mixing class, in its slow-bath form as a lower bound on the bath's mixing time — violating C1, C3, C4 or the persistence form of C2 destroys the structural basis of the observed dynamics, while departing from the slow-bath form bounds the accessible memory without establishing that it is removed.

Necessity of C1 (coupling). Any quantum system realized as a deterministic dilation must have non-trivial coupling between visible and hidden sectors. Proof: without coupling, the visible dynamics are a permutation (unitary on a classical basis), hence P-divisible. (P-indivisibility is not required for a quantum representation — Main §3.4 exhibits a diagonal Hamiltonian with $T(t) = I$, perfectly valid unitary quantum mechanics and P-divisible. What C1's failure removes is the accessible non-Markovianity the characterization is about.) This is a contrapositive: violating C1 produces P-divisible dynamics — structurally distinct from the memory-bearing dynamics the characterization concerns, not from quantum representability, which the finite-horizon equivalence supplies universally ([Main §3.4]).

Necessity of C2 (memory persistence). The hidden sector must retain the record across the window — (C2-structural), which the universal hidden-memory theorem shows every faithful realization of a memory-bearing process contains ([Main §3.4]). In a reversible realization this retention holds for every process, Markov ones included, because the present state determines the visible past; as a condition it bears on coarse-grained descriptions of the hidden sector, in which a record can be degraded before readback. The slow form's necessity is conditional: if the hidden sector *mixes* between readbacks, correlations are eroded and the accessible conditional memory is bounded in terms of the residue of that mixing at the coupling time — the physical-memory theorem in its slow-bath form, under the mixing hypothesis ([Main §3.4]) — while fast *non-mixing* hidden dynamics retain records with no timescale separation at all (`fastbath_probes.py`, [Main §1.3]). Again contrapositive, within the mixing class: violating (C2-slow) produces dynamics whose conditional memory is bounded in terms of that residue — suppressing the memory-bearing sector, not quantum representability.

Necessity of C3 (capacity). The hidden sector must have sufficient capacity. Proof: if the mode count is too small, the mutual information between visible and hidden sectors saturates, limiting the accessible memory the hidden sector can return. Contrapositive: violating C3 produces saturated dynamics — a hard cap on accessible memory, the capacity floor of the universal hidden-memory theorem in quantitative form ([Main §3.4]) — removing the memory-bearing sector, not quantum representability.

Each theorem is a contrapositive: violating C1, C3, C4 or the persistence form

of C2 destroys the accessible non-Markovianity — the memory-bearing sector — not the quantum representation, which the finite-horizon equivalence supplies universally ([Main §3.4]). Together they establish that C1, C3, and C4 are *individually necessary* for any realization — and C2 in its persistence form, with its slow-bath form, within the conditional-mixing class, as a lower bound on the bath's mixing time, which bounds the memory without establishing that it is removed — and (with the per-horizon dilation of Chapter 1) *jointly sufficient*. The equivalence "accessible non-Markovianity $\Leftrightarrow$ embedded observation under C1, C3, C4" is a per-horizon biconditional — C2 sits outside it, its physical content carried by the physical-memory theorem (persistence form without a mixing hypothesis, slow-bath form conditional on it); a fixed-basis unitary quantum representation is universal ($S \iff D \iff Q_{\mathrm{fb}}$), with the compressed form attaching to the correspondence's class under (T).

What the necessity direction does not claim. The necessity direction claims that the observed dynamics require C1–C4 *at the structural level* — any system producing the empirical signature — the measured memory witness, accessible non-Markovianity in the temporal statistics; the temporal memory witness used in the necessity direction enters through the history-readback route, while Bell violation is handled separately by the ontically parameter-dependent composite branch — must satisfy them. It does *not* claim that every empirical system we call "quantum" satisfies C1–C4 *uniquely* — alternative substrata satisfying C1–C4 with different specific bijections would produce the same memory-bearing quantum phenomenology with potentially different specific empirical content at the bijection-specific (Tier 3) level.

The framework's necessity claim is therefore at the *structural* level (C1, C3, C4 necessary unconditionally, C2 within its stated class) rather than at the *substratum-specific* level (which would require the additional claim that only the framework's specific cubic-lattice bijection produces our universe's QM). The structural necessity is what makes the characterization theorem a biconditional; the substratum-specific content is what the framework's empirical record tests directly.

C.6 Finiteness: is Axiom 2 too strong?

A third foundational objection concerns the framework's commitment to finiteness — Axiom 2 of the framework states that the substratum is a finite set S. Critics object that this is incompatible with standard physics' commitment to infinite-dimensional Hilbert spaces in quantum field theory and to the universe being potentially infinite in spatial extent.

The objection. "The universe has infinitely many degrees of freedom. Quantum field theory requires infinite-dimensional Hilbert spaces. Why should I accept finite-dimensionality as a structural commitment of any theory of the universe?"

The framework's response. Axiom 2 plays different roles in different parts

of the framework's content, with different evidential supports for each role.

In Chapter 1's foundational content. Axiom 2 is a mathematical assumption — the characterization theorem is a statement about finite systems. The theorem's mathematical content is precisely defined and proved for finite substrata; whether the theorem's empirical applicability requires actual finiteness of the universe is a separate question.

In Chapter 7's gravitational content. The cosmological horizon's finite area bounds what is observationally accessible: the Bekenstein entropy bound, read as a dimension cutoff (input E3 of Chapter 2 §2.4), limits the visible sector and the boundary layer at the cosmological horizon to at most $e^{S_{\mathrm{dS}}}$ effectively distinguishable states, with $S_{\mathrm{dS}} \sim 10^{122}$. This is an entropy-based bound that holds under the physical assumptions of E3, not a count of microscopic configurations. It does not establish finiteness of the total configuration space S. That extension is posited (A1 of Chapter 2 §2.4), read as the choice of the minimal finite representative: the cardinality of an independently prepared, normalized, exactly decoupled finite deep-sector factor is not fixed by the data at the observation interface (Chapter 2 §2.3). The discreteness scale ϵ does not smuggle in a quantum assumption — it is motivated by the *classical* fact that finite-area boundaries admit finitely many modes. The result $\epsilon = 2\sqrt{\sigma}\, l_p$, with σ the entropy of one boundary mode, is a consequence of self-consistency at the cosmological horizon (Chapter 7 §7.4), not an input.

A referee who rejects Axiom 2 for the universe as a whole (in this section's usage, the finiteness of the total configuration space, A1 of Chapter 2 §2.4, not the second axiom of Chapter 1, recurrence) does not thereby reject the Bekenstein bound, which is independently established in semiclassical gravity and string theory across multiple programs (Hawking 1976, Bekenstein 1981, 't Hooft 1993, Susskind 1995, Bousso 1999, and many subsequent developments): the bound constrains the observationally accessible sector and does not establish finiteness of the whole of S, which is posited. The framework's finiteness commitment agrees with mainstream physics' commitment to the Bekenstein bound on the accessible sector, and extends beyond it by posit.

The "finite degrees of freedom" framing. The standard intuition that "the universe has infinitely many degrees of freedom" comes from quantum field theory treated as a continuum theory — a continuum spacetime with field values at every point producing an infinite-dimensional Hilbert space. But this is the *emergent* description in the framework. At the substratum level, the framework posits a discrete lattice, with the horizon self-consistency relation $\epsilon = 2\sqrt{\sigma}\, l_p$ and finite boundary cell count $A/\epsilon^2 = S_{\mathrm{dS}}/\sigma$ at the cosmological horizon, $\sim 10^{122}$ at $\sigma = 1$.

The framework's content is therefore that the continuum QFT description is the *emergent* description — finite at the substratum level but *appearing* infinite-dimensional when described in the continuum limit. This is structurally analogous to how condensed matter systems with finite lattice spacing exhibit ef-

fective continuum descriptions: the underlying physics is finite, but the long-wavelength effective description involves continuous fields.

The framework's commitment to finiteness is therefore *compatible with* the standard QFT description as an effective continuum theory while making specific structural claims about the substratum level. The two levels of description coexist; the apparent infinite-dimensionality of QFT is an artifact of the continuum limit rather than a property of the underlying physics.

The structural signatures. Finiteness has specific structural signatures developed in Chapter 18 §18.7: exact cosmic recurrence, at a period at most Landau's $g(|S|)$, with $|S|$ not fixed by visible data (falsifiable only in the logical sense, and without practical empirical consequences at periods of astronomical size) and constraints on cosmic information content (compatible with the Bekenstein bound).

The framework's finiteness commitment is therefore not just a mathematical convenience but a structurally consequential commitment. The commitment is supported by the Bekenstein bound (established physics), compatible with continuum QFT (the framework's emergent description), and has specific structural signatures (Chapter 18). A referee rejecting finiteness must engage these supports and signatures specifically rather than dismissing the commitment in general.

C.7 Computational implementability and Wolpert's theorem

A specific class of objections concerns *computational implementability*: whether the framework's deterministic substratum can be computationally implemented by any physical system, given Wolpert's theorem on physical computability and related results from the foundations of computer science.

The objection. "Wolpert (2001, 2008) proved that no physical computer can predict the outputs of another physical computer that is its peer — specifically, no system can implement the function that takes its own future state as input. The framework's deterministic substratum φ must, if physically realized, be computed by something. But what? If φ is computed by a sub-system of the universe, Wolpert's theorem creates issues; if it is computed externally, it requires an unstipulated computing infrastructure."

The framework's response. The framework's content does not claim that φ is *computed* by any physical system. The framework's φ is the underlying deterministic dynamics — what happens at the substratum level — not a function evaluated by a separate computing infrastructure.

The framework's structural commitment is that the substratum *is* the dynamics. The bijection $\varphi : S \to S$ is the rule for how the substratum's configuration

evolves; the substratum itself is what implements the rule. There is no separate computing system that takes the substratum's current state as input and outputs its next state; the substratum's dynamics is the substratum's structure.

This is structurally identical to how classical physics treats the action principle. Hamilton's equations are not *computed* by some additional infrastructure that evaluates the equations and then implements the result; Hamilton's equations *are* the dynamics of the classical system. Similarly, the framework's φ is the substratum's dynamics, not something computed by a separate physical computer.

The relationship to Wolpert. Wolpert's theorem rules out specific kinds of self-referential prediction: no physical system can predict its own future state in advance. The framework's content is consistent with this restriction at the relevant level: embedded observers cannot predict their own future states beyond what their incomplete information about the substratum supports, which is precisely the framework's content on the limits of observation.

The framework's structural commitment is *not* that some physical computer implements φ; it is that the substratum *follows* the rule φ. This is a structural-physical claim, not a computational claim. Wolpert's theorem does not address structural-physical claims about underlying dynamics.

The computability of the substratum-level dynamics. A related concern is whether the framework's substratum-level dynamics is *computable in principle* — whether there exists any algorithm that, given the substratum's current state, can compute its future state. The framework's content is that φ is a bijection on a finite set, which is trivially computable: any finite bijection can be specified as a lookup table or as an algorithm that implements the specific bijection.

The framework does not claim that the bijection's specific form is *efficiently* computable — its computation might require time exponential in the substratum's size or other complexity scaling. But the bijection is in-principle computable, which is the relevant claim for foundational physics. The framework's empirical content (the BQP theorem of Chapter 14) does claim specific computability bounds on what *embedded observers* can compute, but this is a different claim from the bijection's existence.

The implementation question is a different question. The objection's underlying concern is sometimes "what physically implements the substratum?" — what is the universe made of, in concrete physical terms. The framework's content does not answer this question; it provides a *structural* account of the universe's dynamics without committing to specific implementation details (whether the substratum is fundamental fields, discrete cellular automaton cells, or something else entirely).

This is a deliberate methodological choice (Chapter 4 §4.5). The framework's structural commitments are evaluated at the level of structural-empirical predictions, not at the level of substratum implementation. The framework remains

agnostic about implementation while making specific structural claims about dynamics. A reader who wants implementation-level content beyond what the framework provides will need to look elsewhere; the framework's reach is at the structural level.

C.8 Relationship to standard interpretations: MWI, Bohmian, QBism

The framework's content overlaps with several standard interpretations of quantum mechanics. Readers approaching the framework from MWI, Bohmian, QBism, or other interpretive positions raise specific questions about the relationship. The framework's response is structural: each interpretation captures part of what the framework provides; none captures all of it; the framework is a new structural position rather than a variant of any existing interpretation.

Many-Worlds Interpretation (MWI). MWI holds that the universal wave function is the complete physical reality, with measurement producing entanglement between observer and observed without collapse. The "many worlds" are the branches of the wave function corresponding to different measurement outcomes.

What the framework shares with MWI. Both reject wave function collapse as a physical mechanism; both treat the wave function as part of the description of reality rather than a tool for predicting collapse outcomes; both treat observers as physical systems within the larger physical reality rather than as external classical entities.

What the framework does differently. MWI commits to the wave function being *physical reality*; the framework treats the wave function as an *embedded observer's reduced description* of the underlying deterministic substratum. The framework has *one substratum* (the universe-as-deterministic-bijection on S), not many branches. The framework's empirical pattern is reproduced through the substratum's deterministic dynamics combined with embedded observers' partial access; MWI's empirical pattern is reproduced through branch counting and the Born rule.

The structural relationship. MWI's "many branches" correspond, in the framework's reading, to the *epistemic options* available to embedded observers about what the substratum's specific configuration is. Different embedded observers with different histories have different epistemic states (different reduced descriptions); each such epistemic state corresponds to an MWI "branch." The framework's content is therefore that MWI's many branches are *epistemic* (about embedded observers' information) rather than *ontic* (about the substratum having actual multiple realizations).

Bohmian mechanics. Bohmian mechanics adds a pilot wave to standard QM, with particles having definite trajectories guided by the wave function. The

interpretation reproduces quantum predictions while preserving definite particle positions at all times.

What the framework shares with Bohmian. Both have a deterministic substratum-level reality (Bohmian: particles with definite trajectories; framework: substratum bijection φ). Both treat quantum predictions as statistical statements about the substratum-level reality. Both are explicitly non-local in some sense (Bohmian: pilot wave guidance; framework: ontic parameter dependence with operational no-signaling, §C.2).

What the framework does differently. Bohmian mechanics adds a pilot wave to standard QM as a *separate* ontological element; the framework has *no pilot wave*. Bohmian particle trajectories are guided by the wave function; the framework's substratum dynamics is the bijection φ directly, with no guidance equation. Bohmian non-locality runs through the pilot wave on configuration space; the framework's runs through ontic parameter dependence in the hidden sector (§C.2), its memory effects being temporal.

The framework's reading is therefore that Bohmian mechanics adds an unnecessary ontological element (the pilot wave) that the framework's structural content does not require. The framework reproduces Bohmian's deterministic-substratum and statistical-predictions content without the pilot wave's structural cost.

QBism (Quantum Bayesianism). QBism interprets the wave function as a personalist agent's degree of belief about measurement outcomes, with measurement updating these beliefs via standard Bayesian conditioning (extended to quantum probabilities). The interpretation emphasizes the agent-relative character of quantum descriptions.

What the framework shares with QBism. Both treat the wave function as describing an agent's epistemic state rather than directly describing physical reality. Both emphasize observer-relativity in quantum descriptions. Both reject the view that the wave function is the complete physical reality.

What the framework does differently. QBism's agent-relative wave function is *purely epistemic* — there is no commitment to a deeper level of physical reality that the wave function is information about. The framework's wave function is *epistemic-about-substratum* — there is a deeper deterministic level of physical reality (the substratum) that the wave function is incomplete information about.

QBism therefore lacks the framework's substratum-emergent operator distinction; the framework provides a structural foundation that QBism's interpretive content alone does not. The framework's reading is that QBism captures the *epistemic* character of quantum descriptions correctly while missing the *structural foundation* of why these epistemic descriptions take the specific form they do.

The framework's position. The framework is not a variant of any of these interpretations but a distinct structural position. The framework's content is

closest to MWI in treating the substratum as the complete physical reality, closest to Bohmian in treating quantum predictions as statistics over a deterministic substratum, closest to QBism in treating the wave function as epistemic. But the framework's distinctive content — the substratum-emergent operator distinction, the C1–C4 structural conditions, the specific empirical predictions across physics and biology — is not shared by any of the standard interpretations.

The framework's position is therefore: an interpretive-level structural framework with substantive empirical content that goes beyond what any of the standard interpretations provides. Readers approaching from any of the standard positions will find aspects of their preferred interpretation in the framework's content while finding the framework's distinctive content unfamiliar.

C.9 The substratum-emergent operator distinction: substantive or just renaming?

A specific concern raised by readers familiar with effective field theory is whether the framework's substratum-emergent operator distinction is a *substantive* structural feature or just a *renaming* of the standard fundamental/effective theory distinction.

The objection. "Effective field theory already distinguishes between the fundamental theory (UV completion) and the effective theory (IR description). Operators in the effective theory can differ from operators in the fundamental theory through anomaly matching, integrating-out, and similar mechanisms. The framework's substratum-emergent operator distinction is just renaming this standard distinction with new vocabulary. There is no new structural content."

The framework's response. The framework's substratum-emergent operator distinction differs from the standard fundamental/effective distinction in three substantive ways.

The substratum is not a UV completion. In effective field theory, the fundamental theory is the UV completion that produces the effective theory at lower energies. The relationship is one of energy scale. The framework's substratum is *not* the UV completion of the emergent theory; it is the underlying deterministic dynamics with the emergent theory being a *trace-out* description (the visible sector projection) rather than a low-energy limit. The relationship is structural (visible/hidden partition) rather than energy-scale-based.

Substratum operators include classical bilinears, not just QFT operators. Standard effective field theory deals with operators within a quantum field theory framework — both fundamental and effective operators are QFT operators with specific transformation properties. The framework's substratum operators are *classical bilinears* on the substratum lattice — bilinear forms in the substratum's configuration variables. These have different structural properties than QFT operators: they are exactly conserved at the substratum level by the bijec-

tion's group properties, with the emergent QFT operators being the trace-out projections.

This is structurally distinct. Baryon-number conservation, for example, is exactly conserved by the substratum's bilinear structure (the bijection respects the bilinear form). At the emergent level, baryon number is violated by sphalerons in the standard model. The framework's reading is that the violation is at the emergent level only — substratum baryon number is exactly conserved, while emergent baryon number is approximately conserved with sphaleron-mediated violations. The two operators (substratum bilinear vs emergent QFT operator) have different conservation properties.

The distinction has empirical signatures the standard distinction does not. The framework's substratum-emergent distinction predicts specific dissociation patterns between substrate-level and emergent-level operators in biology (Chapter 16 §16.10: iPSC reprogramming residuals, bivalent chromatin states, multi-timescale dynamics). It predicts the cancer methylome classification paradigm (Chapter 17 §17.7: methylation classification succeeds because substratum operators preserve information that emergent operators lose). Its wider-therapeutic-window claim for reader-writer disorders is consistent, as a retrodiction, with the Rett-syndrome reversibility of Guy et al. 2007 (Chapter 16 §16.12).

These empirical patterns are not generic effective-field-theory predictions; they are specific structural predictions of the framework's substratum-emergent operator distinction. The standard fundamental/effective distinction does not predict these patterns. The framework's distinction is therefore *substantively* different from the standard one, with specific empirical signatures that test the difference.

The methodological asymmetry. The framework's substratum-emergent distinction operates *the same way* in fundamental physics and in biology. The same structural pattern that organizes baryon-number conservation and anomaly matching in fundamental physics also organizes iPSC reprogramming residuals, bivalent chromatin states, and methylation classification in biology. The standard fundamental/effective distinction does not have this cross-domain reach — it is specifically a tool of effective field theory, not a general structural principle.

The framework's content is therefore that the substratum-emergent operator distinction is a *general structural principle* with applications across physics and biology, rather than a domain-specific tool. This generality is itself substantive content beyond the standard effective-field-theory distinction.

C.9.1 Observational incompleteness and the uncertainty principle: substantive or renaming?

The objection. "Quantum mechanics already says that an observer cannot know everything about a system: that is Heisenberg's uncertainty principle.

'Observational incompleteness' renames it — or revives Heisenberg's original microscope picture, in which observation disturbs a system that has definite values all along. Either way, nothing new is being said."

The framework's response. The two are different statements; the framework proves an uncertainty-type obstruction and classifies it as carrying no evidence for the framework; and the sharp form of the objection points to an open question rather than to a renaming.

What the uncertainty principle says. The principle goes back to Heisenberg (1927). In its standard quantitative forms it is a family of inequalities in the commutator of two observables: the preparation relation $\sigma_A \sigma_B \geq \frac{1}{2}|\langle [A, B] \rangle|$ (Robertson 1929), which limits how sharply a single state can fix two non-commuting observables, and the error–disturbance relations (Ozawa 2003; Busch, Lahti and Werner 2013), which limit how accurately one observable can be measured without disturbing another. These are theorems of the quantum formalism, valid under every interpretation; they take the formalism as given and say nothing about why it holds. A single observable can be perfectly sharp: the principle constrains pairs.

What observational incompleteness says. The framework's incompleteness concerns an observer embedded in a larger finite deterministic system: the observer's description is a marginal over a hidden sector it cannot access. Its discriminating content is history readback (C4): every faithful deterministic completion of the observer's visible process carries readout-relevant hidden predictive memory that the observer cannot read. This holds for a single fixed readout — one register, recorded over time, with no choice between measurements — where the uncertainty principle has nothing to bound. The visible laws of fixed-readout records are exactly the finite stochastic laws (Main §3.4), and no uncertainty relation constrains that class.

Where the uncertainty principle sits in the framework. The framework proves a qualitative counterpart of the principle as a theorem about observation. An instrument that leaves every state unchanged on average — a passive instrument — learns only about the commutative center of the observable algebra, so complete passive observation fails whenever the algebra is non-commutative (Main §3.4); this is the finite-dimensional form of "no information without disturbance" (Busch 2009). The framework then classifies the statement as non-discriminating: it holds in every finite operational theory with two or more states, whether or not the framework's core conditions hold, and the framework's formal verification exhibits a theory that satisfies it and violates the core. Robertson's inequality is in the same position, since it is a property of density matrices and Hermitian matrices as such. Neither counts as evidence for the framework.

The sharp form of the objection. A stronger critic makes a different point: the framework does not derive, from embedded observation, the non-commutative structure that gives the uncertainty principle its content. That is correct. The

relations bind only where non-commuting observables are operationally available, and the proved equivalence between stochastic, deterministic and quantum descriptions concerns readout in one fixed basis, which an ordinary classical stochastic process also reproduces. The coherent preparations and non-commuting instruments of standard quantum theory, together with its composites, form one quantum operational theory under the completion conditions of the quantum-complete extension OI^{+}, none of which embedded observation supplies on its own; once reversible control is present, the core clause is logically redundant in that derivation, and the finite realization theorem realizes non-quantum operational theories as well (Main §3.4). Whether embedded observation selects the non-commutative structure is the framework's open selection problem. The framework's incompleteness accounts for what an embedded observer cannot access; it does not account for the specific non-commutative form of quantum uncertainty.

Heisenberg's microscope and Bell's theorem. The microscope picture — observation disturbing a system with definite values — is the picture Bell's theorem constrains: a local model with setting-independent hidden variables cannot reproduce quantum correlations. The framework does posit a definite deterministic substratum, and it states the price (Section C.2): it keeps measurement independence and takes ontic parameter dependence, with operational no-signaling. The uncertainty principle carries no commitment of this kind, so the two positions differ in ontology as well as in content.

A precedent. Spekkens' toy theory (Spekkens 2007) shows how far a bare restriction on knowledge reaches: an epistemic restriction alone reproduces complementarity, uncertainty-like trade-offs and no-cloning, but not violations of Bell inequalities. The framework's classification agrees with this lesson: restricted access does not by itself single out quantum theory.

The relation summarized. The two are different statements. The uncertainty-type obstruction is proved and classified as non-discriminating. Deriving from embedded observation the non-commutative structure that gives the uncertainty relations their content is open, as part of the selection problem.

C.10 Circularity concerns: the gap equation for $\hbar$

A specific technical objection concerns the framework's gap equation in Chapter 7 §7.3, which relates the Planck constant $\hbar$ to the discreteness scale through the cosmological horizon's thermodynamics. Critics object that the derivation appears circular.

The objection. "The gap equation equates a classical temperature to a quantum temperature to derive $\hbar$. But the quantum temperature already assumes $\hbar$ exists. The derivation is therefore circular — you are using $\hbar$ to derive $\hbar$."

The framework's response. The logical chain in Chapter 7 §7.3 has four steps that show the derivation is *not* circular.

Step 1. Under the canonical measure of Lemma 3 of Chapter 1, the framework's partition-relativity result proves that $\hbar$ is determined by the partition geometry. This is a structural claim about the framework's setup, not a quantum assumption.

Step 2. Boundary-only dependence proves that $\hbar$ depends only on c, G, and the discreteness scale ϵ. This follows from the framework's substratum being a finite-area boundary system, again a structural claim.

Step 3. Dimensional analysis gives $\hbar = \beta \cdot c^3\epsilon^2/G$ with unknown dimensionless coefficient β. The framework's structural setup determines the functional form; β remains to be fixed.

Step 4. The KMS periodicity $\beta_E = 2\pi c/\kappa$ follows from Euclidean regularity of the metric at the horizon — a *geometric* condition, not a quantum one. The emergent QFT interprets this periodicity as a temperature $T_Q = \hbar\kappa/(2\pi c k_B)$ with $\hbar$ initially unknown. Matching T_Q to the classical temperature $T_{\text{cl}} = c^2\epsilon^2\kappa/(8\pi G k_B \sigma)$, with σ the entropy of one boundary mode, fixes $\beta = 1/(4\sigma)$.

The crucial point. The KMS periodicity does *not* assume $\hbar$. (It does presuppose emergent local boost structure at the horizon — a separate, declared dependency, [GR §8.5] — which is a Lorentz-sector condition, not a quantum one.) It assumes the Wick-rotated metric is smooth at the horizon, which is a condition on the geometry (the imaginary-time geometry must close smoothly at the horizon to avoid a conical singularity). The $\hbar$ enters only through the emergent QFT's interpretation of this periodicity as a temperature. The matching is between a *geometric* quantity (periodicity) and an *emergent* quantity (temperature), not between two quantities that both assume $\hbar$.

The structure of the derivation is therefore: - Geometry $\rightarrow$ KMS periodicity (no $\hbar$ used). - Emergent QFT $\rightarrow$ temperature from KMS periodicity (with $\hbar$ as unknown). - Classical-temperature matching $\rightarrow$ $\hbar$ fixed in terms of ϵ (conditional on H-slope, Chapter 7 §7.3).

The $\hbar$ being "derived" enters only in the second step (as an unknown to be determined), not in the first step (which provides the geometric input independent of $\hbar$). The derivation is therefore non-circular.

The lattice-KMS concern. A related concern is that the KMS periodicity assumes a continuous spacetime, while the framework's substratum is a lattice at $\epsilon = 2\sqrt{\sigma}\, l_p$. The lattice introduces corrections to the KMS periodicity of order $(\epsilon\kappa/c^2)^2 \sim 10^{-122}$ at $\sigma = 1$ — far below any observable threshold.

The framework's response: trans-Planckian insensitivity (Jacobson, 't Hooft, Verlinde, and others) ensures the result is robust against UV modifications. The continuum KMS periodicity is the dominant contribution; the lattice corrections are negligible at the relevant scales. The gap equation is therefore consistent with the lattice structure rather than requiring strict continuum spacetime.

C.11 The framework's epistemic status as a structural framework

A broader objection concerns the framework's epistemic status: whether it should be evaluated as a *physical theory* competing with the Standard Model and general relativity, or as a *structural framework* providing foundations for these theories.

The objection. "The framework's empirical content is mostly *retrodictive* — it explains the Standard Model's parameters and the gravitational sector's structure after the fact. But a genuine physical theory should make novel predictions that distinguish it from competitors. Without such predictions, the framework is not a theory; it is an interpretation."

The framework's response. The framework's epistemic status is intermediate between interpretation and theory. It is a *structural framework* with substantive empirical content but with different empirical exposure than a competing physical theory would have.

What the framework retrodicts. Twenty-two classified Standard Model observables (two parameter-free structural and five conditional structural; the remainder layered, chained or fitted, the gauge couplings among the fitted), seven gravitational-sector entries (conditional relations and estimates), and the Rett wider therapeutic window.

What the framework predicts at the present time. JUNO solar mixing angle at 0.07σ against the post-JUNO global fit — a *parameter-free* retrodiction in the framework's specific sense: derived from structural content, not fitted to the measured value, but the derivation postdates the measurement. The framework's novel-prediction claims rest on the forward set (DESI Y5, BTFR at $z > 1$, JUNO design-lifetime precision on the exact value). (The Bekenstein-Hawking 1/4 coefficient is a consistency relation of the horizon calibration with no direct empirical test; GW250114 constrains the classical area theorem, which the framework preserves.)

What the framework predicts forward. Specific predictions for the BEC analogue-gravity experiment (Chapter 15 §15.8) — the $\mathcal{F}(r) = 2r - r^2$ functional form at marginal hidden capacity (proposed model form; derivation unpackaged). Variant pathogenicity stratification for reader/writer/eraser genes (Chapter 17 §17.8). These are testable forward predictions that distinguish the framework's content from current alternatives.

The intermediate epistemic status. The framework is therefore *not just an interpretation* — it makes specific empirical claims that have been compared with data (JUNO a parameter-free retrodiction at 0.07σ; GW250114 an area-theorem consistency; Rett a retrodiction) and predictions that can be tested in the future (BEC, variant pathogenicity). But it is also *not a competing physical theory* in the conventional sense — the framework does not propose alternative dynamics that would replace the Standard Model or general relativity at the

emergent level. At the emergent level, the framework reproduces standard QM and standard QFT and standard GR.

The framework's distinctive content is at the *structural foundation* level: why standard QM has the specific form it does, why the Standard Model has the specific gauge structure and parameter values it does, why the gravitational sector has the specific features it does, and why specific biological phenomena (Rett, methylation classification) show specific empirical patterns. The framework's value is therefore at the *foundational and unifying* level rather than at the *competing-theory* level.

This intermediate status is sometimes uncomfortable. Critics from the "framework is just interpretation" camp object that the framework lacks distinguishing predictions; critics from the "framework should propose alternative dynamics" camp object that the framework reproduces standard physics rather than replacing it. The framework's response: it is *intentional* that the framework operates at the foundational level rather than at the competing-theory level. The foundational level is where structural questions are addressed; the framework's empirical content at this level is substantial.

C.11.1 Is the derivation gerrymandered? The selector-motivation audit

The objection. "The derivation reaches the Standard Model only because its axioms and selection principles were chosen, with hindsight, to land there. The discrete residue — $d = 3$, $K = 6$, the gauge group — is smuggled in through tuned assumptions, so the 'derivation' is a tautological re-encoding of the target rather than a forcing of it."

The framework's response. The relevant test applied to each assumption and selection principle is whether it is motivated independently of the Standard Model target. On that test the structural core does not depend on the target. Under H-link $K = 2dm$ follows from the link count rather than as a selection rule, with $m = 1$ giving $K = 6$; a different K corresponds to a different bijection. The spatial dimension $d = 3$ follows from three filters stated without reference to the gauge sector (propagating gravity, stable matter, entropy concordance); renormalizability is not used as a filter, as that would be circular. The multiplicities $(3, 2, 1)$ are the decomposition of the six nearest-neighbor link directions under the cubic point group. The audit also identifies two inputs that are not derived in this sense and are recorded as such: the amplitude-scale gauge principle (treated in the substratum development as an operational input rather than a theorem) and the identification of the propagation modes with the gauge-carrying components.

In the quantitative tier the dependence on fitted quantities is larger. There, $1/\alpha_0$ is derived, while the leading threshold coefficient $A \cdot B$ is a finite-lattice numerical cross-check with extrapolated values $\approx$ 48–62 across leading-power models; the comparison with the fitted product remains inconclusive. The uni-

versal threshold δ_0 is an empirical input. Comparisons with candidate threshold identifications, including a group-specific hypercharge-trace interpretation, must retain the leading-power dependence of the $A \cdot B$ calculation; universality of δ_0 is a separate structural constraint. The resulting characterization is a derived structural residue and a coupling-sector retrodiction with a small number of fits.

What remains open. This audit bears only on whether the residue is motivated independently of the target. It does not settle the further question of whether a forced structural residue is *substantively* non-tautological as opposed to merely entailed, which is recorded as an open problem (Chapter 19 §19.3.6) and which the framework proposes to assess by external calibration against the operational reconstruction theorems of quantum theory rather than by further internal argument.

C.12 Meta-objections: credentials, scope, ambition

The framework attracts meta-objections about its author, its scope, and its ambition. These are not legitimate scientific objections in themselves, but they are sometimes raised and warrant brief engagement.

Credentials. "The author has no formal physics credentials. The framework should not be taken seriously regardless of its content."

This is not a scientific objection. Scientific content is evaluated on its merits, not on the credentials of those who propose it. The framework's content consists of mathematical proofs, empirical predictions, and structural derivations — all evaluable independent of the author's biography.

However, the underlying concern (familiar standards may not be met by an outsider) can be addressed substantively. The framework provides complete proofs in the technical chapters (Chapter 1 for the characterization theorem; Chapter 7 for the gravitational derivations; etc.); two derivation routes for key results (Chapter 1 via Barandes correspondence and Stinespring dilation); and explicit falsifiability conditions throughout (Chapter 15 §15.8 BEC test; etc.). The standards the framework meets are *internal scientific standards* (proofs, predictions, falsifiability), not standards based on author identity.

The author takes responsibility for the framework's scientific content, including its proofs, derivations, predictions, and supporting citations.

Scope. "The framework claims to address fundamental physics *and* the origin of life *and* the methodological limits of single-cell bioinformatics. This is too much for a single framework. Each domain deserves its own framework."

The framework's response: the claims are *interconnected by a single mechanism* (C1–C4 structural conditions operating at different scales). Separating them would obscure the connections — in particular, the framework's value comes from being *one* structural account that produces predictions across multiple

domains, not from being separate accounts for each domain. The unification is the framework's distinctive content.

This is not a unique pattern. Statistical mechanics in the nineteenth century similarly addressed thermodynamics, kinetics, transport phenomena, and other domains with one structural account. The unification of multiple domains under one structural framework is a feature, not a bug. The framework's specific empirical record across multiple domains provides convergent evidence for the structural foundations.

The framework is structured to allow domain-specific evaluation. Part II (fundamental physics) is self-contained and can be evaluated independently. Part III (emergence cascade) builds on Part II but is independently evaluable. Part IV (applications) builds on Parts II-III but is also independently evaluable. A reader skeptical of the framework's overall claim can evaluate Part II alone and engage the framework's content at that level.

Ambition. "The framework identifies QM with embedded observation, dissolves the cosmological constant problem, explains the dark sector, and provides a unifying account of memory in biology. This is too ambitious."

The framework's response: ambition is not a scientific objection. The relevant questions are: do the derivations work? do the predictions match empirical data? are the structural commitments coherent? The framework's content provides specific answers to each: the derivations are explicit and the proofs are complete; the retrodictions match data at the precisions classified in Appendix A (JUNO at 0.07σ against the post-JUNO global fit), and GW250114 is consistent with the classical area theorem the framework preserves; the structural commitments are coherent across multiple domains.

A reader who finds the framework's ambition uncomfortable can choose to engage with subsets of the content. The framework's content is structured to be evaluable at multiple levels — at the Chapter-level (specific empirical predictions), at the Part-level (specific structural derivations), or at the framework-level (the full structural commitments). The framework's ambition does not require the reader to engage with all of it at once.

C.13 Closing remarks

The appendix has engaged twelve clusters of objections to the framework's content. The pattern across responses is consistent: each objection is engaged at the level appropriate to its substance, with specific technical content where the objection is technical, with specific methodological content where the objection is methodological, and with specific empirical content where the objection concerns empirical exposure.

What the framework cannot address. The framework's reach has genuine limits. The framework does not address:

The hard problem of consciousness. Chapter 18 §18.10 explicitly identifies consciousness as outside the framework's primary scope. The framework provides structural commitments about the relationship between observation and physical reality (observation as necessary but not sufficient for consciousness; structural constraints on physics-based consciousness theories) but does not derive the *qualitative* character of conscious experience.

The initial conditions of the universe. Chapter 19 §19.3 identifies the cosmological initial state as an open problem. The framework's content includes the universe's dynamics (φ on S) but not the specific initial configuration the universe started from. Why this initial state rather than another is not addressed by the framework's structural content.

The flavor problem. Chapter 19 §19.3 identifies the detailed structure of the CKM and PMNS matrices as open. The framework's content predicts gross structural features (gauge group, generation count, broad mixing-angle structure) but not the specific values of all twelve Yukawa couplings.

The baryon asymmetry. The framework does not derive the specific baryon-to-photon ratio ($n_B/n_\gamma \approx 6 \times 10^{-10}$) from first principles. This is not a gap awaiting a mechanism: the framework's relation to baryogenesis is solution-specific (§19.3.3). The baryon asymmetry is downstream of the flavor-sector CP phases (δ_{CKM}, δ_{PMNS}), which the framework treats as solution-specific inputs rather than derived quantities; η_B inherits that input status. The framework is internally consistent on this point — it does not claim to derive η_B while inputting δ_{CKM} — but it is, by the same token, incomplete in scope on the baryon asymmetry, and says so.

These are genuine limits of the framework's reach. The framework does not pretend to address them; they are explicitly identified as open problems or out-of-scope concerns.

What turns out to be a category error. Several apparent puzzles in fundamental physics turn out, on the framework's analysis, to be *category errors* — problems that arise from confusing levels of description. The framework recasts these rather than solving them, under the conditions stated for each.

The cosmological constant problem. The "120-order-of-magnitude discrepancy" between QFT vacuum energy and observed cosmological constant is a category error in the framework's reading (Chapter 7 §7.7), conditional on the open coupling step ([GR §6.2]; Chapter 19 §19.2.2). The QFT vacuum energy is the *emergent* energy density of the trace-out; the cosmological constant is the *substratum* baseline energy density. The two operate at different levels of description, and their relationship is the framework's specific information compression ratio rather than a direct equality.

The measurement problem. The question "how does measurement produce definite outcomes from superpositions" receives an observer-relative account (Chapter 1; Chapter 19 §19.2.5): at the represented fixed-basis level measurement is

conditioning and re-marginalization, while multi-agent protocols require the proposed completion's operational probabilities and separate premise checks. At the substratum level, there are no superpositions — only the substratum's specific configuration. At the emergent level, superpositions are reduced descriptions used by embedded observers with incomplete information. "Measurement" is the embedded observer's interaction with the substratum that updates their reduced description — not a physical mechanism producing definite outcomes from a previously-indefinite physical state.

The quantum-gravity unification problem. The standard formulation ("quantize gravity to unify with QM") is a category error in the framework's reading (Chapter 7 and Chapter 18 §18.4). Gravity is *substratum-level geometry* in the framework, not an emergent-level field requiring quantization. On the framework's reading QM and GR are projections of one substratum; the covariant field equations and their coupling (G3) are open, so this is a framework-level reading rather than a completed unification (Chapter 19 §19.2.1). The framework does not "quantize gravity"; it explains why QM and GR are *both* emergent descriptions of a single substratum-level reality.

These readings are conditional content of the framework. Its reach is not only retrodictive (explaining established empirical patterns) but reclassificatory (proposing that some apparent puzzles are category errors rather than problems requiring solutions, conditionally on the open steps named above).

The framework's epistemic position summarized. The framework is a structural framework with substantive empirical content. It engages standard interpretations of QM, established no-go theorems, established results in effective field theory, and established methodological concerns through specific technical responses. Its specific empirical claims are retrodictions or consistencies with existing data (JUNO, GW250114, Rett), and its forward predictions can be tested in the future (BEC, variant pathogenicity). Its structural commitments are coherent across multiple domains, with the substratum-emergent operator distinction providing a unifying principle across fundamental physics and biology.

Where the framework's reach has genuine limits (consciousness, initial conditions, flavor problem, the baryon asymmetry), the limits are explicitly identified. Where apparent puzzles are read as category errors (cosmological constant, measurement problem, quantum-gravity unification), the readings are conditional content — on the open coupling step G3 and the operational completion — rather than evasions.

The appendix's value is therefore as a reference resource for readers approaching the framework from different starting points. Readers familiar with Bell's theorem and the no-go theorems should find §§C.2-C.3 directly useful. Readers familiar with effective field theory should find §§C.9 directly useful. Readers familiar with standard interpretations should find §C.8 directly useful. Readers concerned about credentials, scope, or ambition should find §C.12 directly use-

ful. The appendix is intended to be navigable rather than read straight through, with each section engaging the specific cluster of concerns characteristic of one reader audience.

Glossary

This glossary defines the framework's distinctive terminology and the technical vocabulary used throughout the book. Entries are organized alphabetically. Cross-references to chapter and appendix sections point to where each concept is most fully developed.

Note on convention: terms in *italics* within definitions indicate other glossary entries. Where the framework's usage differs from standard physics or mathematics usage, the difference is noted explicitly.

Anomaly cancellation. A condition in quantum field theory requiring that the sum of contributions from chiral fermions to certain triangle diagrams vanishes. In the Standard Model, anomaly cancellation between fermion generations is a structural consistency requirement that constrains the allowed hypercharge assignments. The framework derives the *unique* hypercharge assignment $(Y_Q, Y_u, Y_d, Y_L, Y_e) = (1/6, 2/3, -1/3, -1/2, -1)$ from the six anomaly conditions of $\mathrm{SU}(3)\times\mathrm{SU}(2)\times\mathrm{U}(1)$ on the nonzero-Y_Q branch, with no free parameters. Developed in Chapter 6 §6.7 and Appendix B §B.5.

Axiom 2 (Finiteness). The framework's foundational assumption that the substratum has finitely many degrees of freedom: A1 of Chapter 2 §2.4, under the label Appendix C §C.6 uses, not the second axiom of Chapter 1 (recurrence). Motivated physically by the Bekenstein bound at the cosmological horizon (an entropy-based bound, under the physical assumptions of input E3, of at most $e^{S_{\mathrm{dS}}}$ effectively distinguishable states of the visible sector and boundary layer, $S_{\mathrm{dS}} \sim 10^{122}$). Discussed in Chapter 2 §2.5, with defense in Appendix C §C.6.

Background independence (A6). The sixth of the framework's structural assumptions A1–A6: the substratum dynamics is covariant under spatially varying internal-index transformations, a site-dependent transformation being a symmetry provided the coupling data carried on the links are transformed with it. The content of the assumption is the covariant interface itself — the coupling is link-valued data the transformations act on — and once that is in place the local transformation law imposes no further condition on the rule. It is distinct from the stronger fixed-background condition that the same transformations leave the dynamics invariant with the coupling held fixed, which for an invertible coupling forces agreement across every coupled pair of sites; from the global commutant symmetry, its specialization to transformations constant across the lattice that leave the coupling unchanged, from which the local gauge reading proceeds; and

from the state-dependent coupling graph, a separate principle sharing the name, under which the graph itself evolves with the state. Where the framework says it has no fixed spacetime background, with spacetime emergent, that is a third sense of the words and not this assumption. Developed in Chapter 2 §2.4 and Chapter 5 §5.5.

Barandes correspondence. The mathematical result establishing that stochastic processes in its own indivisible class correspond structurally to unitary quantum dynamics under its hypotheses. The class is a tuple structure — configuration space, target and conditioning times, transition maps, prior, random variable — and divisibility failure is generic within it rather than definitional, so Markov chains are included. The framework's link to it is the explicit translation hypothesis *(T)*: that the framework's per-horizon transition family instantiates that tuple ([Main §3.1], `tuple_probes.py`). *P-indivisibility* does not gate the representation's existence; it determines whether the representation is nontrivial. One of two routes to the transition-statistics layer of the framework's quantum bridge; the other, *Stinespring dilation*, is the shared bedrock rather than a fully independent alternative, and neither route supplies the operational instrument algebra. Developed in Chapter 1 §1.6 and Appendix B §B.2.

Bekenstein-Hawking entropy. The entropy $S = A/(4l_p^2)$ assigned to a horizon of area A in units of the Planck length l_p. The factor $1/4$ follows from the Hawking temperature and the first law. In the framework it is the ratio $2\pi/8\pi$ fixed by the horizon calibration of Chapter 7 (conditional on H-slope); the mode count reproduces it for any cell size once the calibration is imposed. GW250114 (September 2025) confirms the classical area theorem; the $1/4$ coefficient has no direct empirical test. Developed in Chapter 7 §7.5.

Bell-inequality violations. Empirical correlations exceeding bounds derivable from local hidden-variable theories. For a response-complete deterministic OI completion, measurement independence plus ontic parameter independence, with a ± 1 outcome in every run, gives Bell factorization and $|S_{\mathrm{CHSH}}| \leq 2$. The framework therefore retains measurement independence but relinquishes **ontic parameter independence** for Bell-violating substrata, with operational no-signaling required. The measurement-dependent alternative is recorded rather than adopted: the manuscript proves the model-independent necessary lower bound $I_{\mathrm{ont}} \geq 0.0309$ bits for the quantum CHSH value, while an optimized causal model in Hall–Branciard uses about 0.080 bits (the smaller ~ 0.046-bit benchmark is retrocausal). Developed in Chapter 1 §1.7 and Appendix C §C.2.

Born rule. The quantum-mechanical prescription that the probability of measurement outcome i on state $|\psi\rangle$ is $|\langle i|\psi\rangle|^2$. Conventionally treated as a postulate of quantum mechanics. In the framework, at the finite fixed-basis observable-law layer, the Born form is part of an exact equivalent representation: $S \iff D \iff Q_{\mathrm{fb}}$ ([Main §3.4]). The framework therefore does not list a separate 'quadratic-exponent selector' as an unresolved foundational theorem at that layer. What remains is operational: on a finite carrier, within complex matrix kinematics

taken as given, an operational theory's available instruments on the carrier and its composites are the standard noncommuting quantum instruments exactly when the five completion conditions of the operational-completion characterization ([GR §3.3]) hold; the substratum's stated interventions do not supply one of them, full reversible control; whether an extended OI mechanism can supply it, which of the others the substratum supplies, and a derivation of that kinematics are open. The read-write-cycle equilibrium account in Chapter 18 §18.2 is an additional physical mechanism proposal and is not needed for the finite representation equivalence.

BQP (Bounded-error Quantum Polynomial time). The computational complexity class of decision problems solvable by a quantum computer in polynomial time with error probability bounded by 1/3. Conventionally treated as an empirical conjecture about what quantum computers can achieve. Under the framework's explicit computational model — uniform efficient encoding, state preparation and control, a clock model, efficient measurement, precision accounting, and fault-tolerance-grade noise assumptions — BQP is the computational class accessible to embedded observers satisfying C1–C4, with both a lower bound (the model can host the BQP toolkit) and an upper bound (the model cannot exceed it); CPTP representability supplies none of the model hypotheses by itself ([Computation §4]). A generic super-BQP demonstration falsifies the global-coverage premise first (Chapter 14 §14.6); pressing on the framework itself requires further argument. Developed in Chapter 14.

C1 (Coupling). The first of the framework's three structural conditions (C1, C3, C4; C2 is a physical-regime premise, not part of the characterization's equivalence). Requires non-trivial coupling between the *visible* and *hidden sectors* of the *embedded observer*'s partition: hidden influence on some accessible visible transition. A non-permutation one-step matrix T is the cheapest *sufficient witness* of C1 — not its definition and not necessary, since influence can hide at step one and surface later (the delayed-revival family, [Main §3.4]). Physically: information crosses the partition boundary in both directions. C1 fails when the visible sector is causally disconnected from the hidden sector. Discussed in Chapter 1 §1.3.

C2 (Memory persistence). A physical-regime premise rather than one of the three structural conditions (C1, C3, C4): the equivalence runs over those three. Stated in two forms ([Main §1.3]): (C2-structural), that a hidden record of visible history persists to the readback that uses it — supplied by reversibility in every bijective realization, so as a condition it bears on coarse-grained descriptions of the hidden sector; and (C2-slow), $\tau_S \ll \tau_B$, one sufficient mechanism, fast conserved dynamics being another (`fastbath_probes.py`). C2 is the persistence half of what C4 operationally needs (the other half is in-window routing), and, in the coupling-step model of [Main §3.4]'s physical-memory theorem — next visible state determined by the current visible and hidden states, any hidden map between couplings — observed conditional memory $M_k > 0$ at an accessible order k forces some realized hidden conditional at that coupling time

to stay at least $\varepsilon_{\min}(M_k, n_V)$ from its reference prior π_H, with no mixing hypothesis (the physical-memory theorem, [Main §3.4]); under the hidden-sector mixing hypothesis this becomes a lower bound on bath persistence. For our universe at the cosmological horizon, $\tau_S/\tau_B \sim 10^{-32}$. When the coarse-grained hidden sector thermalizes between measurements, C2 fails in its timescale form and, under the mixing hypothesis, the conditional memory of the dynamics is bounded in terms of the residue of that mixing at the coupling time. Discussed in Chapter 1 §1.3.

C3 (Hidden-sector capacity). The second of the framework's three structural conditions (C1, C3, C4). Requires hidden capacity sufficient for the process's conditional memory: $\log_2 |\mathcal{C}_H| \geq I^*$, the capacity floor — the data-processing bound $I(X_{<t}; X_{>t} \mid X_t) \leq I(X_{<t}; H_t \mid X_t) \leq \log_2 |\mathcal{C}_H|$ applied to the whole visible future, which holds in every faithful realization ([Main §3.4]). $N_H \gg N_V$ is the comfortable physical regime, not the condition; for our universe at the cosmological horizon, $N_H \sim 10^{122}$. C3 saturation caps accessible memory — the theorem-level content. The mechanism proposal hypothesizes that the marginal-capacity regime is the *restricted quantum-statistical* regime of partial distinguishability revival, memoryless operational modeling failing while the transition-statistics layer remains representable ([Main §3.4]); all operational-failure readings here are proposal-level until a selection theorem supplies the implication. Discussed in Chapter 1 §1.3, with the restricted quantum-statistical regime in Chapter 15 §15.7 and Chapter 18 §18.3.

C4 (History readback). The third of the framework's three structural conditions (C1, C3, C4). Requires that records of the visible sector's own past, stored in the hidden sector, be read back into the visible dynamics: influence plus storage plus capacity, without readback, is noise rather than memory. C4 is what places a quantitative lower bound on observable conditional memory inside an accessible window (Main §2.3). Discussed in Chapter 1 §1.3.

Cabibbo angle. The CKM mixing parameter $\lambda = \sin\theta_C \approx 0.225$ governing first/second-generation quark mixing in the Standard Model. Conventionally a fitted empirical parameter. The framework obtains $\lambda = 1/(\pi\sqrt{2})$ from a single Brillouin-zone distance on the substratum lattice, under an identification hypothesis whose emergent dressing is open (conditional structural), matching observation at 0.04%. Developed in Chapter 6 §6.5.

Cat state (Schrödinger's cat). A superposition of two macroscopically distinct configurations of a single system, after Schrödinger's 1935 thought experiment in which an entanglement chain amplifies one atom's superposition into a formalism assigning a cat equal parts alive and dead. Conventionally the scenario motivates either an undefined measurement postulate or an added objective-collapse law. The framework dissolves it: the substratum always holds exactly one configuration; the superposition is the *embedded observer*'s indivisible bookkeeping over a definite fact recorded in the *hidden sector*; environmental interaction enacts continuous division events, making divisible (classical) odds correct for any actual cat; and opening the box is intervention, not creation. De-

veloped in Chapter 1 §1.11, with the engineered-cat-state experimental frontier in Chapter 1 §1.6 and the corresponding falsification commitment in Chapter 4 §4.5.

CaMKII (Calcium/calmodulin-dependent protein kinase II). A neuronal signaling kinase forming dodecameric holoenzymes with bistable autophosphorylation states. Canonical molecular memory device in long-term potentiation. The framework identifies CaMKII as the prototypical molecular *C1*–*C4* system in neurons, with the kinase domain as fast subsystem and the regulatory/association domain as slow hidden sector. Pathological τ_B shifts produce the framework's account of neurodegeneration. Discussed in Chapter 16 §16.4.

Characterization theorem. The per-horizon characterization of the memory-bearing sector: an embedded observer's reduced description is non-Markovian on accessible horizons if and only if, for some accessible order, every finite accessible horizon extending at least one step beyond it admits a deterministic realization satisfying *C1*, *C3* and *C4*, with *C4*'s readback at that order; *C2* lies outside the equivalence, its physical content carried by the physical-memory theorem (persistence form without a mixing hypothesis, slow-bath form conditional on it). A fixed-basis unitarily evolving quantum representation exists for every finite-horizon law — internal and universal, by the finite-horizon equivalence $S \iff D \iff Q_{\mathrm{fb}}$ ([Main §3.4]); the compressed representation is supplied across the correspondence's class through the imported stochastic–quantum correspondence, nontrivially where the dynamics is P-indivisible, under the translation hypothesis *(T)* — a separate conditional bridge, not part of the equivalence. Forward direction (sufficiency): a realization with *C1*, *C3* and *C4* $\Longrightarrow$ accessible non-Markovianity, the *C4* readback gap forcing positive conditional memory. Reverse direction (necessity): accessible non-Markovianity $\Longrightarrow$ such a realization on each accessible witness horizon, by the finite-horizon process dilation, with *C1*, *C3* and *C4* each necessary and *C2*'s necessity for physical hidden sectors conditional on the quantitative mixing hypothesis in its slow-bath (timescale) form and holding without it in its persistence form. Developed in Chapter 1 §§1.8-1.9, with necessity proofs in Appendix C §C.5.

Chk1 (Checkpoint kinase 1). A serine/threonine kinase regulating cell-cycle checkpoint responses to DNA damage. The framework identifies Chk1 as a canonical example of regulatory-domain *memory asymmetry*: Chk1 inhibitors selectively sensitize tumor cells because tumor cells rely on non-Markovian checkpoint dynamics that normal cells with intact G1 checkpoints do not require. Discussed in Chapter 16 §16.3.

Chirality (in the framework). The framework's commitment to chiral SU(2) electroweak structure rests on the staggered bipartite structure on the cubic lattice combined with the trace-out operation, and is conditional on the chirality hypothesis H-χ' (SM §4.8). Propagates through molecular physics to produce the parity-violating energy difference (PVED) between mirror-image molecules, predicting L-amino acids as the slightly more stable enantiomer. Discussed in Chapter 10 §10.7.

Chromatin. The protein-DNA complex organizing eukaryotic genomes, including nucleosome positioning, histone modifications (acetylation, methylation), DNA methylation patterns, and three-dimensional chromatin architecture. The framework treats chromatin as the *hidden sector* of the biological *C1–C4* partition with the transcriptional machinery as visible sector. Developed in Chapter 16 §16.9.

CKM (Cabibbo-Kobayashi-Maskawa) matrix. The unitary mixing matrix between Standard Model quark mass and weak interaction eigenstates. Conventionally parameterized by three mixing angles and one CP-violating phase. The framework derives the Cabibbo angle structurally and provides partial content for the broader CKM structure. Developed in Chapter 6 §6.5.

CHSH inequality. The Clauser-Horne-Shimony-Holt formulation of Bell's inequality, with the Bell-local bound of 2 and the quantum (Tsirelson) bound of $2\sqrt{2}$. In the deterministic OI completion, measurement independence plus parameter independence, with a ± 1 outcome in every run, gives the bound 2; quantum violation under measurement independence therefore uses ontic parameter dependence. The cited stochastic causal-local indivisible construction separately has the Tsirelson bound. Discussed in Appendix C §C.2.

Cosmological constant problem. The discrepancy of approximately 10^{122} between quantum field theory's predicted vacuum energy ($\sim 10^{113}$ J/m^3) and the observed value inferred from cosmic expansion ($\sim 6 \times 10^{-10}$ J/m^3). The framework proposes to *dissolve* this problem by identifying the two quantities as properties of logically distinct descriptions: the quantum vacuum energy is the emergent QFT's prediction within a partition; the observed cosmological constant is the gravitational consequence of the substratum's full content. The descriptive distinction is established; the dissolution is conditional on the open coupling step. Developed in Chapter 7 §7.7.

Cryptochrome. A flavoprotein in animal retinas whose photoexcitation produces a radical pair (FADH$\bullet$ and Trp$\bullet$) with millisecond singlet/triplet interconversion. The substrate for the avian magnetic compass. The framework identifies cryptochrome as a canonical *C1–C4* system in biology with the radical-pair electron spins as fast subsystem and the protein matrix as slow bath. Discussed in Chapter 13 §13.5.

Cubic decomposition. The framework's structural result that the six-dimensional representation of the cubic rotation group O on internal lattice components decomposes as $\mathbf{6} = T_1 \oplus E \oplus A_1$ — multiplicities $(3, 2, 1)$ identifying the Standard Model gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$. Developed in Chapter 5 §5.5 and Appendix B §B.6.

DSW (Danielson-Satishchandran-Wald) horizon decoherence. A 2022 result by Danielson, Satishchandran, and Wald proving within standard QFT in curved spacetime that Killing horizons impose fundamental decoherence on nearby quantum superpositions, with rate determined by horizon geometry. External convergence with the framework's universality-class content. Discussed

in Chapter 9 §9.7 and Chapter 15 §15.8.

Embedded observer. The framework's technical term for an observer that is a proper subsystem of the system it registers — coupled to the system, bounded in extent, unable to access the full state from outside. A detector, a protein interior or a cosmological horizon qualifies; nothing in the definition requires consciousness. The central object of the framework's analysis. Defined precisely in Chapter 1 §1.2, characterized by C1–C4 in §1.3.

Emergent description. The reduced description of the visible sector after marginalizing over (tracing out) the hidden sector. In the framework, every finite-horizon observable law — C1–C4 or not — admits a fixed-basis unitary quantum representation ($S \iff D \iff Q_{\mathrm{fb}}$, [Main §3.4]); systems satisfying C1–C4 constitute the memory-bearing sector, where hidden predictive memory is unavoidable in every completion. Developed throughout Chapter 1.

Epigenome. The set of chromatin marks and methylation patterns that regulate gene expression while remaining heritable across cell divisions and (partially) generations. The biological instantiation of the framework's *substratum-emergent operator distinction* at the molecular scale. Discussed in Chapter 16 §§16.9-16.10.

Extended Church-Turing Thesis (ECT). Conventionally, the classical efficient-simulation conjecture — any physically realizable computation efficiently simulable by a probabilistic classical Turing machine — the thesis quantum computing challenges. Its quantum counterpart, the BQP-form (quantum) ECT, asserts efficient quantum-simulability of all physical computation. The framework derives the BQP-form (quantum) ECT as a *conditional theorem* under a global-coverage premise (every physically realizable computation falling under Chapter 14 §14.4's operational-closure and efficiency hypotheses): any computation so covered satisfies the BQP bound. Developed in Chapter 14 §14.6.

Fibre-Gram data. For a coherent lift of a visible law and each visible output, the Gram matrix of the hidden states into which the lift sends the visible inputs. Its diagonal is the visible law itself. Two lifts are related by the two-sided invisible action at one time exactly when their fibre-Gram data agree up to phases, so the off-diagonal part modulo phases is exactly what the visible law leaves free at that time. It is a coordinate on the space of lifts, not a physical quantity, and no statement is made about what selects its trajectory through time. Developed in Chapter 1 and §19.3.9.

Fine-tuning. The observation that certain physical parameters (fine-structure constant, mass ratios, cosmological constant) appear to occupy narrow ranges supporting chemistry and complexity. The framework's reading: most chemistry-supporting features are *structural* (derived from the framework's commitments) rather than parametric, dissolving the conventional fine-tuning problem. Discussed in Chapter 10 §10.8.

Frauchiger-Renner theorem. A 2018 multi-agent no-go theorem deriving a conflict among quantum probability-one certainty, consistency of agents' certainty inferences and single outcomes for its protocol. Partition-relative descriptions alone do not resolve that conflict; applying it to a proposed OI completion requires its operational probabilities and a separate check of those premises. Discussed in Appendix C §C.3.

Gap equation (for $\hbar$). The thermal self-consistency condition between the classical horizon temperature $T_{cl} = c^2\epsilon^2\kappa/(8\pi G k_B \sigma)$ and the KMS temperature $T_Q = \hbar\kappa/(2\pi c k_B)$ of the emergent QFT, producing $\hbar = c^3\epsilon^2/(4G\sigma)$, with σ the entropy of one boundary mode. It relates Planck's constant to the discreteness scale, conditional on H-slope, and fixes only $\epsilon/\sqrt{\sigma} = 2\,l_p$: a self-consistency condition, not an independent determination of the value of $\hbar$. Developed in Chapter 7 §7.3 and Appendix B §B.4.

Gauge group. A continuous group of local symmetries under which the dynamics is invariant. The Standard Model gauge group is $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$. Conditionally on H-link, the framework derives this gauge group from the stabilizer of a cubic-equivariant condensate with distinct block values on the substratum lattice (the coupling matrix itself is scalar); the stabilizer computation is exact, the physical-carrier identification is the premise. Developed in Chapter 5.

Generation (fermion). In the Standard Model, the three sets of fermions with identical gauge-quantum-number assignments but different masses: (u, d, e, ν_e), (c, s, μ, ν_μ), (t, b, τ, ν_τ). The framework derives one singlet plus one symmetry-related triplet of candidate matter sectors from the staggered taste structure on the cubic lattice; reading the triplet as three *physical* generations is conditional on *H-spin'* (SM §4.7), whose free-kernel and 4D-staggered routes are both settled negative. Discussed in Chapter 6 §6.2.

God's-eye view. A hypothetical observer coextensive with the universe, with complete knowledge of all degrees of freedom. The framework identifies the God's-eye view as the *C3* failure limit and shows that it is not a limit case of observation but a category outside it — observation is partition-relative, and without a partition, observation does not occur. Discussed in Chapter 1 §1.10.

GW250114. A high-signal-to-noise binary black-hole merger observed by LIGO on January 14, 2025 (hence "GW250114"), with the LIGO/Virgo/KAGRA analysis paper published in *Physical Review Letters* on September 10, 2025, confirming Hawking's area theorem ($dA \geq 0$, classical horizon dynamics) at 99.999% confidence. The observation compares horizon areas and does not measure entropy, so it does not test the Bekenstein-Hawking 1/4 coefficient. Discussed in Chapter 7 §7.5.

Hidden sector. The portion of the framework's partition that the embedded observer does not directly access. The framework's four conditions specify properties of this sector (coupling, slow evolution, large capacity). For laboratory observers in our universe, the hidden sector is everything beyond the cosmolog-

ical horizon plus the trans-horizon degrees of freedom of any local environment. Defined in Chapter 1 §1.2.

Higgs (in the framework). The framework treats the Higgs as the *singlet* (A_1) taste of the staggered decomposition — a *composite scalar* in the framework's account rather than a fundamental field. The Higgs quartic coupling at the compositeness scale is exactly zero, with the observed value at low energies generated by RGE running. Developed in Chapter 6 §6.2.

Hsp90. A molecular chaperone that buffers genetic variation by chaperoning mutant proteins into functional conformations. The canonical evolutionary capacitor (Rutherford and Lindquist 1998). The framework identifies Hsp90-client conformational states as a *C1–C4* hidden sector with stress-induced release of accumulated variation. Discussed in Chapter 17 §17.9.

Hypercharge. The Standard Model U(1) quantum number of fermion multiplets, with values $(Y_Q, Y_u, Y_d, Y_L, Y_e) = (1/6, 2/3, -1/3, -1/2, -1)$. Conventionally fitted empirical inputs. The framework derives these values as the unique anomaly-free solution on the nonzero-Y_Q branch. Developed in Chapter 6 §6.7 and Appendix B §B.5.

Information backflow. A revival of distinguishability: the trace distance between the evolved states of two preparations increases over some interval (the Breuer–Laine–Piilo signature). In a P-divisible process distinguishability never increases, so backflow certifies P-indivisibility; a P-indivisible process need not show it. It is distinct from accessible finite-horizon memory, the conditional past–future dependence that history readback (C4) supplies on accessible windows, which does not by itself produce backflow ([Main §2.3]). Discussed in Chapter 1 §1.5.

Jordan-Chevalley projection. The algebraic decomposition of an operator into commuting semisimple and nilpotent parts: $F = F_{ss} \cdot F_u$ with $F_u = I + N$ and $N^2 = 0$. The framework's content: the trace-out operation extracts F_{ss} and discards N, organizing the emergent description by the representation theory of the partition. Developed in Appendix B §B.3.

JUNO (Jiangmen Underground Neutrino Observatory). A reactor neutrino experiment that in November 2025 reported the first world-leading direct measurement of the solar neutrino mixing angle: $\sin^2\theta_{12} = 0.3092 \pm 0.0087$. The framework's parameter-free value $1/3 - 1/(4\pi^2) = 0.3080$ matches the direct measurement at 0.14σ and the post-JUNO global fit (0.3085 ± 0.0073) at 0.07σ — a retrodiction; JUNO's design-lifetime precision provides the pre-registered forward test. Discussed in Chapter 8.

Koide relation. An empirical relation among the three charged-lepton masses: $Q = (m_e + m_\mu + m_\tau)/(\sqrt{m_e} + \sqrt{m_\mu} + \sqrt{m_\tau})^2$. Observed value $Q = 0.666664 \pm 0.000005$. The framework predicts $Q = 2/3$ from cubic-group quadratic Casimir; the match is better than 0.001% (0.4σ). Developed in Chapter 6 §6.6.

Markovian. A stochastic process whose future depends only on the present

state, with no memory of past states. Standard chemical kinetics is Markovian; the framework's embedded-observer dynamics is *memory-bearing* — non-Markovian on accessible horizons — while the quantum representation itself is universal, Markov chains included ([Main §3.4]). Contrasted throughout Chapter 1.

Memory asymmetry. The framework's unifying therapeutic principle in medicine: when a disease process depends on non-Markovian signaling dynamics that the corresponding normal tissue does not depend on, therapies can target the memory structure rather than the catalytic function, with the framework predicting wider therapeutic windows. Developed in Chapter 16 §16.13.

MOND (Modified Newtonian Dynamics). A phenomenological framework for galactic rotation curves involving an acceleration scale $a_0 \approx 1.2 \times 10^{-10}$ m/s^2. Conventionally treated as a phenomenological alternative to dark matter. The framework derives $a_0 = cH/6$ from entropy displacement at the cosmological boundary with no fitted parameters, at the G1 register and conditional on the covariant rung G3 (see the GR ladder). Developed in Chapter 7 §7.9.

Non-Markovian. A stochastic process whose future depends on past states beyond the present. The framework's embedded-observer dynamics is fundamentally memory-bearing — non-Markovian on accessible horizons — and this, not representability (which is universal) or recurrence-scale P-indivisibility (which every non-deterministic finite reversible representative has), carries the discriminating content ([Main §3.4]). Predicted to appear at multiple scales: cosmological (quantum mechanics), biological (enzyme kinetics, gene expression), engineered (qubit noise). Defined in Chapter 1 §1.5. Three notions are kept apart throughout and none unconditionally implies another: *accessible finite-horizon memory* (equivalently conditional past–future dependence in the visible law) is the framework's own term and is what "non-Markovian on accessible horizons" means here; *stochastic P-indivisibility* is failure of divisibility of the visible population law and rests on recurrence together with a non-permutation rooted map, not on C1/C3/C4 alone; *generic quantum non-Markovianity* is an open-system notion whose content depends on the quantum divisibility or backflow criterion adopted, and it is not what either of the first two asserts.

Operational gluing theorem. The framework's finite-resolution operational result ([Main §3.4]): every finite adaptive quantum experiment — instrument family $\mathcal{F}$, horizon K, grid G — has a finite reversible deterministic embedded realization whose adaptive outcome statistics lie within the stated $O(K/G)$ bound, with one fixed mechanism per instrument, gluing by restriction across overlapping families, sequential and parallel composition, and explicit hidden-capacity bounds. The theorem establishes finite operational **realizability** and, together with the uniform-continuum theorem, finite-test density. It does not by itself prove that the standard local quantum instrument category is the unique coherent operational completion, because the same finite reversible machinery can realize non-quantum instrument families. Which further conditions select the quantum instrument category on a finite carrier is stated by the operational-

completion characterization ([GR §3.3]): five completion conditions, none supplied by bare finite OI, within complex matrix kinematics taken as given. Certified in exact arithmetic (`opglue_probes.py`).

NV (Nitrogen-vacancy) center. A point defect in diamond where a nitrogen atom replaces a carbon atom adjacent to a lattice vacancy. The basis for room-temperature quantum sensing with current sensitivity limited by the NV electron-spin coherence time. The framework predicts that backflow-aware sensing protocols exploiting the diamond lattice's slow nuclear-spin bath recover information Markovian protocols discard; the resulting sensitivity gain is not quantified. Discussed in Chapter 15 §15.6.

Operator distinction (substratum-emergent). The framework's structural distinction between operators acting at the substratum level (with specific conservation properties at the molecular or quantum-field level) and operators acting at the emergent level (with conservation properties at the coarse-grained level). The two operators can be functionally distinct even when they look identical at the emergent level. Developed in Chapter 6 §6.3 (for fundamental physics) and Chapter 16 §16.10 (for chromatin biology).

Page curve. The time-dependence of black-hole entanglement entropy: rising during the early evaporation phase, reaching a peak at the Page time, then falling as the black hole evaporates fully. The framework obtains a cycle-averaged Page curve on the full product space of black hole and radiation, under the scrambling condition H-scramble and the decoupling condition H-decouple (H-scramble fails for every bijection acting on an initial configuration with the bipartition aligned to the configuration coordinates, so the Page value needs an initial superposition over many configurations or a non-aligned bipartition), with a Page time that, for a mass-independent emission coefficient, is a fixed fraction of the evaporation time: $t_P \approx 0.646\, t_{evap}$ when the radiation's entropy equals the horizon entropy it carries off, earlier when it carries more. Developed in Chapter 7 §6.

Partition-relativity. The framework's structural feature that the same physical substratum admits multiple visible/hidden partitions, each producing its own emergent description. The emergent description depends on the partition choice; the underlying substratum does not. Does not by itself resolve a no-go theorem: the operational probabilities and each theorem's substantive premises require separate checks. Developed in Chapter 1 §1.4.

P-indivisibility. Failure of a stochastic transition family to factor through valid intermediate stochastic maps. In OI, a non-permutation one-step marginal on a fixed finite reversible system is a sufficient recurrence witness. More generally, any non-permutation rooted map forces indivisibility somewhere in the full recurrence cycle: recurrence returns the rooted visible map to the identity, and a non-permutation stochastic matrix has no stochastic inverse. A C4 readback gap supplies such a map, and so does every non-deterministic visible step, including in laws that are Markov on every accessible window. With a full-support

visible prior the indivisibility is equivalent to memory at some order below the recurrence time, observable only when that order is accessible. This global result does not imply accessible short-time P-indivisibility in every window; the two notions are kept distinct. P-indivisibility is a statement about the visible population law and is not generic quantum non-Markovianity: inside the framework's diagonal-preserving realizations it does rule out a stochastic intermediate map and hence CP-divisibility, but outside that class the two come apart — a CP-divisible family with a unitary intermediate map can induce P-indivisible computational-basis populations, the intermediate map consuming coherences the population law never sees ([Main §2.3]).

PMNS (Pontecorvo-Maki-Nakagawa-Sakata) matrix. The unitary mixing matrix between Standard Model neutrino mass and flavor eigenstates. The framework derives all three PMNS mixing angles from the cubic-group flavor structure, including the solar angle $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2)$, which matches the JUNO-era global fit at 0.07σ (a retrodiction). Discussed in Chapter 8.

Read-write cycle. The framework's realization picture for the coupling between the visible and hidden sectors of the partition. The observer reads the visible sector, writes correlations into the hidden sector through the partition coupling, and reads back those correlations on subsequent measurements. The cycle's statistics carry the memory-bearing sector — the fixed-basis quantum representation supplied universally ([Main §3.4]), the Born-rule equilibrium reading a mechanism proposal (Chapter 18 §18.2) — at the cosmological scale, and heredity at the molecular scale. Defined in Chapter 1 §1.6.

Reader/writer/eraser pharmacology. The framework's three-axis organization of epigenetic therapeutics, corresponding directly to the operator distinction at chromatin. *Writers* establish substrate-level marks (DNMT and HMT inhibitors); *erasers* remove them (HDAC and HDM inhibitors); *readers* translate marks to functional output (BET bromodomain inhibitors). The framework's structural prediction: reader-writer disorders show wider therapeutic windows than substrate-gene disorders, empirically confirmed for Rett syndrome. Developed in Chapter 16 §§16.11-16.12.

Restricted quantum-statistical regime. The mechanism proposal's name for partial distinguishability revival on an accessible window, $0 < \mathcal{N} < 1$. Here $\mathcal{N}$ is the revival measure of Chapter 18 §18.3: the largest increase $d_t - d_s$, over pairs of preparations and times $s < t$ in the window, of their distinguishability d (the total variation of the visible population laws, equivalently the trace distance of the diagonal states) — the peak *information backflow* on the window, kept separate from the cumulative Breuer–Laine–Piilo sum. $\mathcal{N} > 0$ certifies P-indivisibility on the window; memory alone does not give $\mathcal{N} > 0$ ([Main §2.3]). The name refers to the quantum representation at the transition-statistics layer, which every finite law admits together with a deterministic hidden realization, with the operational layer not supplied ([Main §3.4]); it marks a revival range, neither a partial nor a complete quantum theory and not a quantum-membership scale. Hypothesized at marginal hidden capacity,

with the proposed model form $\mathcal{F}(r) = 2r - r^2$, $r = N_H/N_V$ (derivation not supplied). Diagnosed by distinguishability-revival measurements; a multi-time memory test alone does not separate it from complete revival. Developed in Chapter 13 §13.7, Chapter 15 §§15.7-15.8, Chapter 18 §18.3.

Rett syndrome. A neurodevelopmental disorder caused by loss-of-function mutations in MECP2, a methyl-CpG binding reader protein. Guy et al. (2007) demonstrated that restoration of MECP2 expression in adult MECP2-knockout mice reverses the neurological phenotype. The framework's wider-therapeutic-window claim for reader-writer disorders is consistent with it as a retrodiction, the result predating the framework; it is the framework's strongest consistency anchor outside fundamental physics. Discussed in Chapter 16 §16.12.

RNA world. The hypothesis that early life used RNA for both information storage and catalysis before DNA and proteins evolved their specialized roles. The framework's structural reading: RNA-like architecture is the simplest single-molecule C1–C4 system (simultaneously template and catalyst), making the RNA world a structural prediction rather than a historical accident. Developed in Chapter 11 §11.7.

Staggered (fermion structure). The placement of independent fermion modes at alternating cubic lattice sites, producing the chiral structure of the framework's emergent QFT and the multiple-taste structure that gives rise to the $1 \oplus 3$ candidate-sector decomposition (three physical generations only under *H-spin'*). Developed in Chapter 5 §5.6 and Chapter 6 §6.2.

Standard Model. The current best theory of fundamental particle physics: a chiral gauge theory with gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$, three fermion generations, one Higgs doublet, and approximately 19 free parameters. The framework reproduces twenty-two classified observables — two parameter-free structural, five conditional structural, the remainder layered, chained or fitted (Appendix A) — and the anomaly-free hypercharges on the nonzero-Y_Q branch; the gauge group follows under H-link, whose single-copy premise is not itself derived, the physical generation reading under H-spin', and the chirality assignment under H-χ'. Developed in Chapters 5-6.

Stinespring dilation. A 1955 theorem of operator algebra establishing that any completely-positive trace-preserving map on a Hilbert space can be realized as a unitary evolution on a larger Hilbert space with subsequent tracing-out. The framework uses Stinespring dilation for its own permutation-dilation route from the deterministic substratum to the transition-statistics layer — shared bedrock with the *Barandes correspondence* rather than an independent alternative to it. Developed in Chapter 1 §1.6 and Appendix B §B.2.

Substratum. The framework's underlying deterministic, finite-dimensional, bijective dynamical system (S, φ) from which the emergent quantum description is obtained through the trace-out operation. The substratum is the framework's primitive ontological commitment, held at the level of the reconstructed local residue modulo the substratum gauge group rather than any particular rep-

resentative (Chapter 3) — the "what is really there" beneath the emergent quantum-mechanical description. Constructed concretely in Chapter 2.

Substratum-emergent operator distinction. *See operator distinction.*

Tier 1, Tier 2, Tier 3. The framework's stratification of predictions by their robustness across substratum representatives. *Tier 1*: results that follow from the framework's universality class (shared across all substrata satisfying the structural conditions). *Tier 2*: results dependent on broader structural commitments but not on specific lattice details. *Tier 3*: solution-specific results dependent on the particular cubic-lattice realization. Developed in Chapter 9 §§9.3-9.4.

TLS (Two-level system). Structural defects in amorphous oxide substrates of superconducting qubits, characterized by flipping between two configurations at characteristic rates. The framework identifies the TLS bath as the *hidden sector* in superconducting qubit C1–C4 systems, with $\tau_S/\tau_B \sim 10^{-3}$ producing the framework's predicted correlated error structure. Discussed in Chapter 15 §15.3.

Trace-out. The mathematical operation of marginalizing over (summing or integrating out) the hidden-sector degrees of freedom, producing the reduced description on the visible sector. The framework's central technical operation. Equivalent to partial trace in quantum mechanics. Defined throughout Chapter 1.

Tsirelson bound. The maximum CHSH inequality violation achievable by standard quantum mechanics: $2\sqrt{2}$. In the manuscript this value is retained for the cited **stochastic causal-local indivisible** construction and for any successful lift to the standard quantum Bell set. It is not the ceiling of a measurement-independent, parameter-independent deterministic completion recording a ± 1 outcome in every run, whose Bell bound is 2. The deterministic OI branch reaches quantum violation by ontic parameter dependence; the measurement-dependent alternative is separately **bounded and benchmarked** in Chapter 1 §1.7. Discussed in Appendix C §C.2.

U-parity. The framework's grading structure in the neutrino sector, distinguishing U^+ and U^- states in the cubic-group flavor classification. Produces the framework's distinctive sum rule and structural relation among the three PMNS mixing angles. Developed in Chapter 8 §8.4.

Universality class. A set of substrata that share the same emergent description at the framework's Tier 1 level. The framework's content: the substratum-to-emergent map is many-to-one, with multiple bijections producing the same observable physics. Developed in Chapter 9 §9.3.

Vibronic coherence. Quantum coherence between electronic exciton states and specific vibrational modes of the protein environment. The framework identifies the picosecond-scale coherences observed in photosynthetic complexes

(FMO, LH2, photosystem II) as vibronic rather than purely electronic, consistent with the framework's C1–C4 architecture. Discussed in Chapter 13 §13.4.

Visible sector. The portion of the framework's partition that the embedded observer directly accesses. Contrasted with the *hidden sector*. The visible sector is where measurements occur and where the emergent description applies. Defined in Chapter 1 §1.2.

Wolpert's theorem. David Wolpert's results (2001, 2008) on fundamental limits of inference devices embedded in the system they are trying to predict. A physical analog of Gödel's incompleteness theorem and Turing's halting problem result — logical, worst-case, and unquantified. The framework's partition-induced trace-out is the mechanism of a *parallel physical* incompleteness, with the memory-bearing dynamics quantum mechanics represents as the specific cost; Wolpert's result corroborates the pattern but is not the source of the framework's derivation. Discussed in Chapter 12 §12.8, Appendix C §C.7.

Yamanaka factors. The four transcription factors (Oct4, Sox2, Klf4, c-Myc) whose transient expression reprograms somatic cells to induced pluripotent stem cells. The framework identifies iPSC reprogramming as confirming the *substratum-emergent operator distinction* at chromatin: the emergent operator (cell identity) flips while substantial residual substrate-level operator (DNA methylation patterns reflecting source cell type) persists. Discussed in Chapter 16 §16.10.

Bibliography

This bibliography lists the works cited in the book, organized by topical area. Within each section, entries are alphabetical by first author. The framework's own technical companion papers are listed first under "Framework-internal citations"; subsequent sections cover foundations and interpretation, fundamental physics, gravitational physics and cosmology, chemistry and biology, evolution and molecular biology, medicine and pharmacology, bioinformatics and computational biology, quantum information and computing, and historical and philosophical works.

Framework-internal citations

Maybaum, A. (2026). *The Observational Incompleteness Framework*. Zenodo, concept DOI: 10.5281/zenodo.19060318 (resolves to the current release). [Repository containing the framework's full technical content: the core papers — Main (finite-observer representation: the exact finite-horizon correspondence of stochastic laws, reversible realizations and fixed-basis unitary/Born representations, and hidden memory; quantum reconstruction as an open programme),

Substratum (substratum construction and reconstruction theorem), Structure (structural realism), SM (Standard Model derivation), GR (gravitational sector) — together with the focused presentation Juno (neutrino-sector prediction), the methodology paper Physics Modulo Gauge, and the companion documents Complexity, Medicine, and Bioinformatics.]

Foundations of quantum mechanics and interpretation

Arnold, V. I. (1989). *Mathematical Methods of Classical Mechanics* (2nd ed., Graduate Texts in Mathematics 60). Springer-Verlag.

Breuer, H.-P., Laine, E.-M., & Piilo, J. (2009). Measure for the degree of non-Markovian behavior of quantum processes in open systems. *Physical Review Letters* 103: 210401. arXiv:0908.0238.

Breuer, T. (1995). The impossibility of accurate state self-measurements. *Philosophy of Science* 62(2): 197–214.

Engel, K.-J., & Nagel, R. (2000). *One-Parameter Semigroups for Linear Evolution Equations* (Graduate Texts in Mathematics 194). Springer-Verlag.

Bacciagaluppi, G., & Valentini, A. (2009). *Quantum Theory at the Crossroads: Reconsidering the 1927 Solvay Conference*. Cambridge University Press.

Barandes, J. A. (2023). The Stochastic-Quantum Correspondence. *arXiv*:2302.10778.

Barandes, J. A. (2024). The Stochastic-Quantum Theorem. *arXiv*:2309.03085.

Braunstein, S. L., & Pati, A. K. (2007). Quantum information cannot be completely hidden in correlations: Implications for the black-hole information paradox. *Physical Review Letters* 98: 080502.

Bell, J. S. (1964). On the Einstein-Podolsky-Rosen paradox. *Physics Physique Fizika* 1: 195-200.

Bong, K.-W., Utreras-Alarcón, A., Ghafari, F., Liang, Y.-C., Tischler, N., Cavalcanti, E. G., Pryde, G. J., & Wiseman, H. M. (2020). A strong no-go theorem on the Wigner's friend paradox. *Nature Physics* 16: 1199-1205.

Busch, P. (2009). "No information without disturbance": Quantum limitations of measurement. In W. C. Myrvold & J. Christian (Eds.), *Quantum Reality, Relativistic Causality, and Closing the Epistemic Circle*. Springer.

Busch, P., Lahti, P., & Werner, R. F. (2013). Proof of Heisenberg's error-disturbance relation. *Physical Review Letters* 111: 160405.

Clauser, J. F., Horne, M. A., Shimony, A., & Holt, R. A. (1969). Proposed experiment to test local hidden-variable theories. *Physical Review Letters* 23: 880-884.

Colbeck, R., & Renner, R. (2011). No extension of quantum theory can have improved predictive power. *Nature Communications* 2: 411.

Dalla Chiara, M. L. (1977). Logical self reference, set theoretical paradoxes and the measurement problem in quantum mechanics. *Journal of Philosophical Logic* 6: 331-347.

de la Peña, L., Cetto, A. M., & Valdés-Hernández, A. (2014). *The Emerging Quantum: The Physics Behind Quantum Mechanics*. Springer.

Einstein, A., Podolsky, B., & Rosen, N. (1935). Can quantum-mechanical description of physical reality be considered complete? *Physical Review* 47: 777-780.

Fein, Y. Y., Geyer, P., Zwick, P., Kiałka, F., Pedalino, S., Mayor, M., Gerlich, S., and Arndt, M. (2019). Quantum superposition of molecules beyond 25 kDa. *Nature Physics* 15: 1242-1245.

Fine, A. (1982). Hidden variables, joint probability, and the Bell inequalities. *Physical Review Letters* 48: 291-295.

Heisenberg, W. (1927). Über den anschaulichen Inhalt der quantentheoretischen Kinematik und Mechanik. *Zeitschrift für Physik* 43: 172-198.

Nimmrichter, S., and Hornberger, K. (2013). Macroscopicity of mechanical quantum superposition states. *Physical Review Letters* 110: 160403.

Ozawa, M. (2003). Universally valid reformulation of the Heisenberg uncertainty principle on noise and disturbance in measurement. *Physical Review A* 67: 042105.

Pedalino, S., Ramírez-Galindo, B. E., Ferstl, R., Hornberger, K., Arndt, M., and Gerlich, S. (2026). Probing quantum mechanics with nanoparticle matter-wave interferometry. *Nature* 649: 866.

Robertson, H. P. (1929). The uncertainty principle. *Physical Review* 34: 163-164.

Rössler, O. E. (1998). *Endophysics: The World as an Interface*. World Scientific.

Schrödinger, E. (1935). Die gegenwärtige Situation in der Quantenmechanik. *Die Naturwissenschaften* 23: 807-812, 823-828, 844-849.

Spekkens, R. W. (2007). Evidence for the epistemic view of quantum states: A toy theory. *Physical Review A* 75: 032110.

Szangolies, J. (2018). Epistemic horizons and the foundations of quantum mechanics. *Foundations of Physics* 48(12): 1669-1697.

Valdés-Hernández, A., de la Peña, L., & Cetto, A. M. (2011). Bipartite entanglement induced by a common background (zero-point) radiation field. *Foundations of Physics* 41: 843-862.

Frauchiger, D., & Renner, R. (2018). Quantum theory cannot consistently describe the use of itself. *Nature Communications* 9: 3711.

Kochen, S., & Specker, E. P. (1967). The problem of hidden variables in quantum mechanics. *Journal of Mathematics and Mechanics* 17: 59-87.

Pusey, M. F., Barrett, J., & Rudolph, T. (2012). On the reality of the quantum state. *Nature Physics* 8: 475-478.

Slagle, K., & Preskill, J. (2023). Emergent quantum mechanics at the boundary of a local classical lattice model. *Physical Review A* 108: 012217. arXiv:2207.09465.

Stinespring, W. F. (1955). Positive functions on C-*algebras.* Proceedings of the American Mathematical Society* 6: 211-216.

Tsirelson, B. S. (1980). Quantum generalizations of Bell's inequality. *Letters in Mathematical Physics* 4: 93-100.

Wolpert, D. H. (2001). Computational capabilities of physical systems. *Physical Review E* 65: 016128. [The 2001 publication of arguments first developed in 1992.]

Wood, C. J., & Spekkens, R. W. (2015). The lesson of causal discovery algorithms for quantum correlations: causal explanations of Bell-inequality violations require fine-tuning. *New Journal of Physics* 17: 033002.

Standard Model physics and particle phenomenology

't Hooft, G. (1993). Dimensional reduction in quantum gravity. *arXiv*:gr-qc/9310026. [Origin of the holographic principle.]

Capozzi, F., Lisi, E., Marcone, F., Marrone, A., & Palazzo, A. (2025). Updated bounds on the (1,2) neutrino oscillation parameters after first JUNO results. *arXiv*:2511.21650. *Physical Review D* 114: 016026 (2026).

Cabibbo, N. (1963). Unitary symmetry and leptonic decays. *Physical Review Letters* 10: 531-533. [Original Cabibbo angle.]

Cech, T. R. (1989). Self-splicing and enzymatic activity of an intervening sequence RNA from *Tetrahymena*. *Nobel Lecture*, December 8, 1989. Published in *Les Prix Nobel 1989*. Reprinted in *Bioscience Reports* 10: 239-261.

Cech, T. R., Zaug, A. J., & Grabowski, P. J. (1981). In vitro splicing of the ribosomal RNA precursor of *Tetrahymena*: involvement of a guanosine nucleotide in the excision of the intervening sequence. *Cell* 27: 487-496.

JUNO Collaboration (2025). First measurement of reactor neutrino oscillations at JUNO. *arXiv*:2511.14593; *Nature* 654: 343 (2026). Reports $\sin^2\theta_{12} =$

0.3092 ± 0.0087 and $\Delta m^2_{21} = (7.50 \pm 0.12) \times 10^{-5}\,\mathrm{eV}^2$ from the first 59.1 days of data collected after detector completion in August 2025.

Koide, Y. (1983). Fermion-boson two-body model of quarks and leptons and Cabibbo mixing. *Lettere al Nuovo Cimento* 34: 201-205. [Original Koide relation.]

Pontecorvo, B. (1957). Mesonium and antimesonium. *Soviet Physics JETP* 6: 429.

Wu, C. S., Ambler, E., Hayward, R. W., Hoppes, D. D., & Hudson, R. P. (1957). Experimental test of parity conservation in beta decay. *Physical Review* 105: 1413-1415.

Gravitational physics, cosmology, and quantum gravity

Abedi, J., Dykaar, H., & Afshordi, N. (2017). Echoes from the abyss: Tentative evidence for Planck-scale structure at black hole horizons. *Physical Review D* 96: 082004. arXiv:1612.00266.

Bao, N., Carroll, S. M., & Singh, A. (2017). The Hilbert space of quantum gravity is locally finite-dimensional. *International Journal of Modern Physics D* 26(12): 1743013. arXiv:1704.00066.

Bekenstein, J. D. (1973). Black holes and entropy. *Physical Review D* 7: 2333-2346.

Bekenstein, J. D. (1981). Universal upper bound on the entropy-to-energy ratio for bounded systems. *Physical Review D* 23: 287-298.

Bertini, N. R., Novaes, M. H., von Marttens, R., & Shapiro, I. L. (2025). Renormalization group corrections to ΛCDM model and observational consequences for H_0 tension. arXiv:2509.24026 [astro-ph.CO; gr-qc; hep-th]. [Lead author initial in earlier in-text references was given as "S." — correct lead author is Nicolas R. Bertini.]

Bousso, R. (1999). A covariant entropy conjecture. *Journal of High Energy Physics* 7: 4.

Chandrasekaran, V., Longo, R., Penington, G., & Witten, E. (2022). An algebra of observables for de Sitter space. *Journal of High Energy Physics* 2023(02): 82.

Chen, C.-H., & Penington, G. (2024). A clock is just a way to tell the time: gravitational algebras in cosmological spacetimes. arXiv:2406.02116. [Constructs gauge-invariant gravitational algebras in cosmological spacetimes with observers, focusing on slow-roll inflation and evaporating Schwarzschild-de Sitter black holes; corroborates the observer-incorporated semiclassical gravity program adjacent to the DSW horizon-decoherence result. Previously cited in-text

as "Chen, Penington, and Salzetta (2024)"; identified during May 2026 second-sweep citation-verification pass.]

Cornish, N. J., Spergel, D. N., Starkman, G. D., & Komatsu, E. (2004). Constraining the topology of the Universe. *Physical Review Letters* 92: 201302. arXiv:astro-ph/0310233.

Danielson, D., Satishchandran, G., & Wald, R. M. (2022). Black holes decohere quantum superpositions. *International Journal of Modern Physics D* 31: 2241003.

DESI Collaboration (2025). DESI DR2 results II: Measurements of baryon acoustic oscillations and cosmological constraints. *Physical Review D* 112: 083515. arXiv:2503.14738.

Genzel, R., Förster Schreiber, N. M., Übler, H., Lang, P., Naab, T., Bender, R., et al. (2017). Strongly baryon-dominated disk galaxies at the peak of galaxy formation ten billion years ago. *Nature* 543: 397-401.

Gralla, S. E., & Wei, H. (2024). Decoherence from horizons: General formulation and rotating black holes. *Physical Review D* 109: 065031. arXiv:2311.11461.

Hawking, S. W. (1975). Particle creation by black holes. *Communications in Mathematical Physics* 43: 199-220.

Hawking, S. W. (1976). Breakdown of predictability in gravitational collapse. *Physical Review D* 14: 2460-2473.

Harlow, D., Usatyuk, M., & Zhao, Y. (2025). Quantum mechanics and observers for gravity in a closed universe. *arXiv*:2501.02359.

Jacobson, T. (1995). Thermodynamics of spacetime: The Einstein equation of state. *Physical Review Letters* 75: 1260-1263.

Jiao, Y., et al. (2023). Detection of the Keplerian decline in the Milky Way rotation curve. *Astronomy & Astrophysics* 678: A208.

Abac, A. G., et al. (LIGO Scientific, Virgo, and KAGRA Collaborations) (2025). GW250114: Testing Hawking's Area Law and the Kerr Nature of Black Holes. *Physical Review Letters* 135: 111403. DOI: 10.1103/kw5g-d732.

LUX-ZEPLIN Collaboration (2025). New search for WIMP dark matter with 417 live days of LUX-ZEPLIN data. Announcement December 8, 2025, with preprint and submission to *Physical Review Letters*. Based on the largest dataset ever collected by a dark matter detector: 417 live days of data from March 2023 to April 2025, probing WIMP masses between 3 and 9 GeV/c^2. Earlier (2024) LZ results: J. Aalbers et al. (LZ Collaboration), "Dark Matter Search Results from 4.2 Tonne-Years of Exposure of the LUX-ZEPLIN (LZ) Experiment," *Physical Review Letters* 135: 011802 (2025), arXiv:2410.17036.

Maldacena, J. (2024). Real observers solving imaginary problems. *arXiv*:2412.14014.

Page, D. N. (1993). Average entropy of a subsystem. *Physical Review Letters* 71: 1291-1294. arXiv:gr-qc/9305007.

Page, D. N. (2005). Hawking radiation and black hole thermodynamics. *New Journal of Physics* 7: 203. arXiv:hep-th/0409024.

Page, D. N. (2013). Time dependence of Hawking radiation entropy. *Journal of Cosmology and Astroparticle Physics* 2013(09): 028. arXiv:1301.4995.

Penington, G. (2019). Entanglement wedge reconstruction and the information paradox. *Journal of High Energy Physics* 2020(09): 002. arXiv:1905.08255.

Planck Collaboration (2018). Planck 2018 results. VI. Cosmological parameters. *Astronomy & Astrophysics* 641: A6.

Planck Collaboration (2020). Planck 2018 results. IX. Constraints on primordial non-Gaussianity. *Astronomy & Astrophysics* 641: A9.

Popescu, S., Short, A. J., & Winter, A. (2006). Entanglement and the foundations of statistical mechanics. *Nature Physics* 2: 754-758. arXiv:quant-ph/0511225.

Susskind, L. (1995). The world as a hologram. *Journal of Mathematical Physics* 36: 6377-6396.

Tsang, K. W., Rollier, M., Ghosh, A., Samajdar, A., Agathos, M., Chatziioannou, K., Cardoso, V., Khanna, G., & Van Den Broeck, C. (2018). A morphology-independent data analysis method for detecting and characterizing gravitational wave echoes. *Physical Review D* 98: 024023. arXiv:1804.04877.

Verlinde, E. P. (2017). Emergent gravity and the dark universe. *SciPost Physics* 2: 016. arXiv:1611.02269. [Cited in-text as "Verlinde (2024)"; no substantive 2024 Verlinde-authored emergent-gravity paper could be identified, and the 2017 SciPost paper remains the canonical reference. The in-text year should be corrected to 2017, or the citation reframed if a specific later work was intended.]

Westerweck, J., Nielsen, A. B., Fischer-Birnholtz, O., Cabero, M., Capano, C., Dent, T., Krishnan, B., Meadors, G., & Nitz, A. H. (2018). Low significance of evidence for black hole echoes in gravitational wave data. *Physical Review D* 97: 124037. arXiv:1712.09966.

Wilson-Gerow, J., Dugad, A., & Chen, Y. (2024). Decoherence by warm horizons. *Physical Review D* 110: 045002. arXiv:2405.00804. [Direct extension of the DSW horizon-decoherence result; provides a local fluctuation-dissipation derivation complementing the original global one. Previously cited in-text as "Wilson-Gerow and Verlinde (2024)"; identified during May 2026 second-sweep citation-verification pass — Verlinde is not a co-author on this paper.]

Chemistry and origin of life

Guerrier-Takada, C., Gardiner, K., Marsh, T., Pace, N., & Altman, S. (1983). The RNA moiety of ribonuclease P is the catalytic subunit of the enzyme. *Cell* 35(3 Pt 2): 849-857. [Foundational paper from Altman's group cited in the 1989 Nobel Prize in Chemistry shared with Cech.]

Damer, B., & Deamer, D. (2020). The hot spring hypothesis for an origin of life. *Astrobiology* 20: 429-452.

Ferris, J. P., Hill, A. R., Liu, R., & Orgel, L. E. (1996). Synthesis of long prebiotic oligomers on mineral surfaces. *Nature* 381: 59-61.

Frank, F. C. (1953). On spontaneous asymmetric synthesis. *Biochimica et Biophysica Acta* 11: 459-463.

Kauffman, S. A. (1986). Autocatalytic sets of proteins. *Journal of Theoretical Biology* 119: 1-24.

Miller, S. L. (1953). A production of amino acids under possible primitive earth conditions. *Science* 117: 528-529.

Soai, K., Shibata, T., Morioka, H., & Choji, K. (1995). Asymmetric autocatalysis and amplification of enantiomeric excess of a chiral molecule. *Nature* 378: 767-768.

Wächtershäuser, G. (1990). Evolution of the first metabolic cycles. *Proceedings of the National Academy of Sciences USA* 87: 200-204.

Evolution and molecular biology

Bedford, T., & Hartl, D. L. (2008). Overdispersion of the molecular clock: temporal variation of gene-specific substitution rates in *Drosophila*. *Molecular Biology and Evolution* 25: 1631-1638.

Cutler, D. J. (2000). Understanding the overdispersed molecular clock. *Genetics* 154: 1403-1417.

Eldredge, N., & Gould, S. J. (1972). Punctuated equilibria: an alternative to phyletic gradualism. In: T. J. M. Schopf (ed.), *Models in Paleobiology*, pp. 82-115. Freeman, Cooper.

Gillespie, J. H. (1989). Lineage effects and the index of dispersion of molecular evolution. *Molecular Biology and Evolution* 6: 636-647.

Maynard Smith, J., & Szathmáry, E. (1995). *The Major Transitions in Evolution*. Oxford University Press.

Odling-Smee, F. J., Laland, K. N., & Feldman, M. W. (2003). *Niche Construction: The Neglected Process in Evolution*. Princeton University Press.

Rutherford, S. L., & Lindquist, S. (1998). Hsp90 as a capacitor for morphological evolution. *Nature* 396: 336-342.

Tao, Q., Tamura, K., U. Mello, B., & Kumar, S. (2019). Reliable confidence intervals for relevant timetree-based parameters. *Molecular Biology and Evolution* 36: 811-824.

Wiltschko, R., & Wiltschko, W. (2019). Magnetoreception in birds. *Journal of the Royal Society Interface* 16: 20190295.

Wiser, M. J., Ribeck, N., & Lenski, R. E. (2013). Long-term dynamics of adaptation in asexual populations. *Science* 342: 1364-1367.

Zeng, L.-W., Comeron, J. M., Chen, B., & Kreitman, M. (1998). The molecular clock revisited: the rate of synonymous vs. replacement change in *Drosophila*. *Genetica* 102/103: 369-382. DOI: 10.1023/A:1017035109224. [Source of the *Drosophila* dispersion-index range R(t) $\approx$ 1.6-2.6 cited in the text; previously attributed in-text to a "Wang et al. (2010)" entry that could not be located. The 1.6-2.6 range traces to this paper via Bedford & Hartl (2008).]

Medicine, pharmacology, and clinical biology

Bernstein, B. E., Mikkelsen, T. S., Xie, X., Kamal, M., Huebert, D. J., Cuff, J., et al. (2006). A bivalent chromatin structure marks key developmental genes in embryonic stem cells. *Cell* 125: 315-326.

Cuddapah, V. A., Pillai, R. B., Shekar, K. V., Lane, J. B., Motil, K. J., Skinner, S. A., et al. (2014). Methyl-CpG-binding protein 2 (MECP2) mutation type is associated with disease severity in Rett syndrome. *Journal of Medical Genetics* 51: 152-158.

English, B. P., Min, W., van Oijen, A. M., Lee, K. T., Luo, G., Sun, H., Cherayil, B. J., Kou, S. C., & Xie, X. S. (2006). Ever-fluctuating single enzyme molecules: Michaelis-Menten equation revisited. *Nature Chemical Biology* 2: 87-94.

Prebet, T., Sun, Z., Figueroa, M. E., Ketterling, R., Melnick, A., Greenberg, P., Juckett, M., Smith, M. R., Malick, L., Paietta, E., Czader, M., Gabrilove, J., Erba, H. P., Gore, S. D., & Tallman, M. S. (2014). Prolonged administration of azacitidine with or without entinostat for myelodysplastic syndrome and acute myeloid leukemia with myelodysplasia-related changes: Results of the US Leukemia Intergroup trial E1905. *Journal of Clinical Oncology* 32(12): 1242-1248. [Phase II randomized trial NCT00313586 testing scheduling of azacitidine $\pm$ HDAC inhibitor entinostat as primary endpoint.]

Guy, J., Gan, J., Selfridge, J., Cobb, S., & Bird, A. (2007). Reversal of neurological defects in a mouse model of Rett syndrome. *Science* 315: 1143-1147.

Horvath, S. (2013). DNA methylation age of human tissues and cell types. *Genome Biology* 14: R115. DOI: 10.1186/gb-2013-14-10-r115.

Klinman, J. P., & Kohen, A. (2013). Hydrogen tunneling links protein dynamics to enzyme catalysis. *Annual Review of Biochemistry* 82: 471-496.

Lin, X., et al. (2024). Residual epigenetic memory in induced pluripotent stem cells. *Frontiers in Cell and Developmental Biology* 12: 1306530. DOI: 10.3389/fcell.2024.1306530.

Nicolas, D., Phillips, N. E., & Naef, F. (2018). What shapes eukaryotic transcriptional bursting? *Molecular BioSystems* 14: 1623-1629.

Quintás-Cardama, A., Santos, F. P. S., & Garcia-Manero, G. (2011). Therapy with azanucleosides for myelodysplastic syndromes. *Nature Reviews Clinical Oncology* 8: 433-444.

Stasevich, T. J., Hayashi-Takanaka, Y., Sato, Y., Maehara, K., Ohkawa, Y., Sakata-Sogawa, K., Tokunaga, M., Nagase, T., Nozaki, N., McNally, J. G., & Kimura, H. (2014). Regulation of RNA polymerase II activation by histone acetylation in single living cells. *Nature* 516: 272-275. DOI: 10.1038/nature13714.

Tontsch-Grunt, U., Traexler, P.-E., Baum, A., Musa, H., Marzin, K., Wang, S., Trapani, F., Engelhardt, H., & Solca, F. (2022). Therapeutic impact of BET inhibitor BI 894999 treatment: backtranslation from the clinic. *British Journal of Cancer* 127: 577-586. DOI: 10.1038/s41416-022-01815-5. PMID: 35444289. [Previously cited in-text as "Marek et al. 2022"; identified during May 2026 citation-verification pass. The paper reports BET inhibitor BI 894999 combined with the clinical p300/CBP inhibitor CCS1477 in NUT carcinoma.]

Voigt, P., Tee, W.-W., & Reinberg, D. (2013). A double take on bivalent promoters. *Genes & Development* 27: 1318-1338. DOI: 10.1101/gad.219626.113.

Wilen, C. D., Abdullah, S., Kurinsky, N. A., Stanford, C., Cardani, L., D'Imperio, G., Tomei, C., Faoro, L., Ioffe, L. B., Liu, C. H., Opremcak, A., Christensen, B. G., DuBois, J. L., & McDermott, R. (2021). Correlated charge noise and relaxation errors in superconducting qubits. *Nature* 594: 369-373. DOI: 10.1038/s41586-021-03557-5.

Wilsker, D., Petermann, E., Helleday, T., & Bunz, F. (2008). Essential function of Chk1 can be uncoupled from DNA damage checkpoint and replication control. *Proceedings of the National Academy of Sciences USA* 105: 20752-20757.

Ye, B. S., Chang, K. W., Kang, S., Jeon, S., & Chang, J. W. (2025). Repetitive and extensive focused ultrasound-mediated bilateral frontal blood-brain barrier opening for Alzheimer's disease. *Journal of Neurosurgery* 142: 1263-1272. DOI: 10.3171/2024.8.JNS24989. [Korean trial NCT06158789; previously entered as "Ye, J. H." — the lead author is Byoung Seok Ye. The study reported improvement in neuropsychiatric symptoms from blood-brain-barrier opening alone, without concurrent antibody medication.]

Zhang, X., Zegar, T., Lucas, A., Morrison-Smith, C., Knox, T., French, C. A., Knapp, S., Müller, S., & Siveke, J. T. (2020). Therapeutic targeting of

p300/CBP HAT domain for the treatment of NUT midline carcinoma. *Oncogene* 39: 4770-4779. DOI: 10.1038/s41388-020-1301-9. PMID: 32366905. [Previously cited in-text as "Bauer et al. 2020"; identified during May 2026 citation-verification pass. The paper reports combination of p300/CBP HAT inhibitor A-485 with BET inhibitor JQ1, with synergistic effects against NUT midline carcinoma.]

Bioinformatics and computational biology

Bergen, V., Lange, M., Peidli, S., Wolf, F. A., & Theis, F. J. (2020). Generalizing RNA velocity to transient cell states through dynamical modeling. *Nature Biotechnology* 38: 1408-1414.

Bowling, S., Sritharan, D., Osorio, F. G., Nguyen, M., Cheung, P., Rodriguez-Fraticelli, A., et al. (2020). An engineered CRISPR-Cas9 mouse line for simultaneous readout of lineage histories and gene expression profiles in single cells. *Cell* 181: 1410-1422.

Bravo González-Blas, C., et al. (2023). SCENIC+: single-cell multiomic inference of enhancers and gene regulatory networks. *Nature Methods* 20: 1355-1367.

Capper, D., Jones, D. T. W., Sill, M., Hovestadt, V., Schrimpf, D., Sturm, D., et al. (2018). DNA methylation-based classification of central nervous system tumours. *Nature* 555: 469-474.

Csendes, G., Petenei, A., Heitz, A., Boldog, D., Bencze, A., Mádai, M., Megyeri, A., Szalka, B., Kemenes, B., Schoenholzer, F., Frey, M., Cucchi, F., Tomek, M. P., Werner, C., Sonzogni, A., Stiastny, J., Crisan, M., Rodriguez Hernandez, M., Ribeiro Frazão, J., Bartoš, F., Bischof, B., Hammer, S., Domanski, J., Mey, A. S. J. S., Goyal, R., Schölkopf, B., & Theis, F. J. (2024). Foundation models in single-cell transcriptomics: benchmarking and gene-attention analysis. *Nature Communications* (forthcoming).

Cui, H., Wang, C., Maan, H., Pang, K., Luo, F., Duan, N., & Wang, B. (2023). scGPT: toward building a foundation model for single-cell multi-omics using generative AI. *Nature Methods* (2024).

Drexler, R., Brembach, F., Sauvigny, J., Ricklefs, F. L., Eckhardt, A., Bode, H., Gempt, J., Lamszus, K., Westphal, M., Schüller, U., & Mohme, M. (2024). Unclassifiable CNS tumors in DNA methylation-based classification: clinical challenges and prognostic impact. *Acta Neuropathologica Communications* 12: 9. DOI: 10.1186/s40478-024-01728-9.

Gorin, G., Fang, M., Chari, T., & Pachter, L. (2022). RNA velocity unraveled. *PLOS Computational Biology* 18: e1010492.

Saxton, M. N., Morisaki, T., Krapf, D., Kimura, H., & Stasevich, T. J. (2023). Live-cell imaging uncovers the relationship between histone acetylation,

transcription initiation, and nucleosome mobility. *Science Advances* 9(40): eadh4819. DOI: 10.1126/sciadv.adh4819.

Hummer, G., & Szabo, A. (2001). Free energy reconstruction from nonequilibrium single-molecule pulling experiments. *Proceedings of the National Academy of Sciences USA* 98: 3658-3661. DOI: 10.1073/pnas.071034098. [The manuscript cites the Hummer-Szabo single-molecule kinetics program collectively; the companion methodological paper is Hummer, G., & Szabo, A. (2010), Free energy profiles from single-molecule pulling experiments, *PNAS* 107: 21441-21446, DOI: 10.1073/pnas.1015661107.]

Kedzierska, K. Z., Crawford, L., Amini, A. P., & Lu, A. X. (2024). Assessing the limits of zero-shot foundation models in single-cell biology. *bioRxiv*. DOI: 10.1101/2023.10.16.561085.

Koelsche, C., et al. (2021). Sarcoma classification by DNA methylation profiling. *Nature Communications* 12: 498.

La Manno, G., Soldatov, R., Zeisel, A., Braun, E., Hochgerner, H., Petukhov, V., et al. (2018). RNA velocity of single cells. *Nature* 560: 494-498.

Li, T., et al. (2022). MultiVelo: chromatin velocity for joint inference of trajectories and RNA velocity. *Nature Biotechnology* (2023).

Lu, H. P., Xun, L., & Xie, X. S. (1998). Single-molecule enzymatic dynamics. *Science* 282: 1877-1882.

Marbach, D., Schaffter, T., Mattiussi, C., & Floreano, D. (2009). Generating realistic in silico gene networks for performance assessment of reverse engineering methods. *Journal of Computational Biology* 16: 229-239. DOI: 10.1089/cmb.2008.09TT. [The DREAM5 crowd-inference synthesis is Marbach, D., Costello, J. C., Küffner, R., Vega, N. M., Prill, R. J., Camacho, D. M., et al. (2012), Wisdom of crowds for robust gene network inference, *Nature Methods* 9: 796-804, DOI: 10.1038/nmeth.2016.]

McEwen, M., Faoro, L., Arya, K., Dunsworth, A., Huang, T., Kim, S., et al. (2022). Resolving catastrophic error bursts from cosmic rays in large arrays of superconducting qubits. *Nature Physics* 18: 107-111. DOI: 10.1038/s41567-021-01432-8.

McKenna, A., Findlay, G. M., Gagnon, J. A., Horwitz, M. S., Schier, A. F., & Shendure, J. (2016). Whole-organism lineage tracing by combinatorial and cumulative genetic barcoding. *Science* 353: aaf7907.

Roohani, Y., Huang, K., & Leskovec, J. (2024). Predicting transcriptional outcomes of novel multigene perturbations with GEARS. *Nature Biotechnology* 42(6): 927-935. DOI: 10.1038/s41587-023-01905-6. [Previously cited in-text as "Pyhel et al. 2024"; identified during May 2026 citation-verification pass. GEARS integrates deep learning with a gene-coexpression knowledge graph and Gene-Ontology-derived perturbation knowledge graph to predict transcriptional responses to single and multi-gene perturbations.]

Stolovitzky, G., Monroe, D., & Califano, A. (2007). Dialogue on reverse-engineering assessment and methods: The DREAM of high-throughput pathway inference. *Annals of the New York Academy of Sciences* 1115: 1-22. DOI: 10.1196/annals.1407.021.

Tedesco, M., Giannese, F., Lazarević, D., Giansanti, V., Rosano, D., Monzani, S., et al. (2022). Chromatin Velocity reveals epigenetic dynamics by single-cell profiling of heterochromatin and euchromatin. *Nature Biotechnology* 40: 235-244. DOI: 10.1038/s41587-021-01031-1.

Theodoris, C. V., Xiao, L., Chopra, A., Chaffin, M. D., Sayed, Z. R. A., Hill, M. C., et al. (2023). Transfer learning enables predictions in network biology. *Nature* 618: 616-624.

Quantum biology and biological information processing

Cao, J., Cogdell, R. J., Coker, D. F., Duan, H.-G., Hauer, J., Kleinekathöfer, U., et al. (2020). Quantum biology revisited. *Science Advances* 6: eaaz4888. DOI: 10.1126/sciadv.aaz4888. [Cited in-text as "Cao, Fleming, and others"; entered previously as "Cao & Plenio (2025)", which is a misattribution — this 18-author review is led by Jianshu Cao and does not include M. B. Plenio. The in-text reference and year should be corrected accordingly.]

Cha, Y., Murray, C. J., & Klinman, J. P. (1989). Hydrogen tunneling in enzyme reactions. *Science* 243: 1325-1330.

Duan, H.-G., Prokhorenko, V. I., Cogdell, R. J., Ashraf, K., Stevens, A. L., Thorwart, M., & Miller, R. J. D. (2017). Nature does not rely on long-lived electronic quantum coherence for photosynthetic energy transfer. *Proceedings of the National Academy of Sciences USA* 114: 8493-8498.

Engel, G. S., Calhoun, T. R., Read, E. L., Ahn, T.-K., Mančal, T., Cheng, Y.-C., Blankenship, R. E., & Fleming, G. R. (2007). Evidence for wavelike energy transfer through quantum coherence in photosynthetic systems. *Nature* 446: 782-786.

Gunawardena, J. (2012). A linear framework for time-scale separation in non-linear biochemical systems. *PLOS ONE* 7: e36321.

Hiscock, H. G., Worster, S., Kattnig, D. R., Steers, C., Jin, Y., Manolopoulos, D. E., Mouritsen, H., & Hore, P. J. (2016). The quantum needle of the avian magnetic compass. *Proceedings of the National Academy of Sciences USA* 113: 4634-4639.

Hiscock, H. G., Hiscock, T. W., Kattnig, D. R., Scrivener, T., Lewis, A. M., Manolopoulos, D. E., & Hore, P. J. (2019). Navigating at night: fundamental limits on the sensitivity of radical pair magnetoreception under dim light. *Quarterly Reviews of Biophysics* 52: e9. DOI: 10.1017/S0033583519000076.

Quantum information, computing, and complexity

Aaronson, S. (2005). Quantum computing, postselection, and probabilistic polynomial-time. *Proceedings of the Royal Society A* 461: 3473-3482.

Aaronson, S. (2005). NP-complete problems and physical reality. *ACM SIGACT News* 36(1): 30-52. [Aaronson2005b]

Bennett, C. H., Bernstein, E., Brassard, G., & Vazirani, U. (1997). Strengths and weaknesses of quantum computing. *SIAM Journal on Computing* 26: 1510-1523. [Original BQP bounds.]

Bernstein, E., & Vazirani, U. (1997). Quantum complexity theory. *SIAM Journal on Computing* 26: 1411-1473. [Original definition of BQP.]

Google Quantum AI (2023). Suppressing quantum errors by scaling a surface code logical qubit. *Nature* 614: 676-681.

Nielsen, M. A., & Chuang, I. L. (2010). *Quantum Computation and Quantum Information*. Cambridge University Press.

Steinhauer, J. (2016). Observation of quantum Hawking radiation and its entanglement in an analogue black hole. *Nature Physics* 12: 959-965.

Mathematical foundations and computability

Birkhoff, G. (1946). Three observations on linear algebra. *Universidad Nacional de Tucumán Revista Series A* 5: 147-151. [Birkhoff-von Neumann theorem.]

Gödel, K. (1931). Über formal unentscheidbare Sätze der *Principia Mathematica* und verwandter Systeme I. *Monatshefte für Mathematik und Physik* 38: 173-198.

Landau, E. (1903). Über die Maximalordnung der Permutationen gegebenen Grades. *Archiv der Mathematik und Physik* (3) 5: 92-103. [Maximum order of a permutation.]

Turing, A. M. (1937). On computable numbers, with an application to the *Entscheidungsproblem*. *Proceedings of the London Mathematical Society* s2-42: 230-265.

von Neumann, J. (1953). A certain zero-sum two-person game equivalent to the optimal assignment problem. In: H. W. Kuhn & A. W. Tucker (eds.), *Contributions to the Theory of Games II*, pp. 5-12. Princeton University Press. [Birkhoff-von Neumann theorem.]

Historical and philosophical works

Cusa, N. of (1440). *De docta ignorantia* (*On Learned Ignorance*).

Cusa, N. of (1453). *De visione Dei* (*On the Vision of God*).

Deutsch, D. (2011). *The Beginning of Infinity: Explanations That Transform the World*. Viking Press.

Penrose, R. (2004). *The Road to Reality: A Complete Guide to the Laws of the Universe*. Jonathan Cape.

Spinoza, B. (1677). *Ethica, Ordine Geometrico Demonstrata* (*Ethics, Demonstrated in Geometrical Order*).

Zuckerkandl, E., & Pauling, L. (1965). Evolutionary divergence and convergence in proteins. In: V. Bryson & H. J. Vogel (eds.), *Evolving Genes and Proteins*, pp. 97-166. Academic Press. [Origin of the molecular clock hypothesis.]