The Substratum Construction: Reconstruction, the Substratum Gauge Group, and the QM-GR Synthesis

Observer geometry and physical carrier. SM §4.1 distinguishes two claims. First, finite equivariant projection is classical and exact: cubic symmetry is inherited by the projected Markov part and every memory kernel, forbidding quadratic anisotropy in the scalar symbol — so whatever nondegenerate quadratic structure survives is isotropic — while allowing cubic anisotropy at quartic order. The particular normalized nearest-neighbor operator $A/(2d)$ additionally requires an **observer-level** center-free kernel; microscopic center independence alone does not imply that after correlated hidden variables are marginalized. Second, the six signed simple-cubic links and their exact $3 \oplus 2 \oplus 1$ representation are unconditional geometry, but identifying that link space with the complete physical carrier is **H-link**; the single-copy clause gives $K = 6$, and the custodial condensate-stabilizer reading is **H-cust**. Accordingly, any downstream statement here that consumes $K = 6$, $N_f = 6$, the physical SM gauge carrier, or a fixed three-sector count inherits those named bridge conditions unless it is explicitly referring only to the six geometric links.

Author: Alex Maybaum
Date: April 2026
Status: DRAFT PRE-PRINT
Classification: Theoretical Physics / Foundations

Abstract

Quantum mechanics and general relativity are conventionally treated as two fundamental theories awaiting unification. This paper develops the alternative that they are two projections of a single finite deterministic construction, joined by a third projection that supplies a time-symmetric observer-selection rule, read against the area of the apparent horizon, which on an expanding homogeneous general-relativistic background satisfying the null energy condition is non-decreasing as a matter of background kinematics. The underlying object is the bijection (S, φ) on a finite configuration space. Its visible-sector reduction has the exact finite stochastic/quantum-representation equivalence established in [Main]: finite observable laws, finite reversible realizations, and fixed-basis unitary/Born representations coincide, while genuine history readback forces hidden predictive memory. Its spatial coupling graph supplies finite causal cones; equality with the complete standard quantum instrument theory holds, on a finite carrier with complex matrix kinematics taken as given, exactly under the five completion conditions characterized in [GR §3.3]; under the substratum's stated interventions, condition (iv), full reversible control, is not supplied [GR §3.3], and whether an extended OI mechanism supplies it, and which of the other

conditions the substratum supplies, are open. The boundary-thermodynamic projection yields the gravitational description developed in [GR], and the lattice dynamics yields the Standard-Model structure developed in [SM].

The two substratum-level results that make these emergences one construction rather than independent applications are the reconstruction theorem (Theorem 23) and the substratum gauge group (Theorem 24). Theorem 23 reconstructs the **local propagating lattice/gauge residue** from its stated empirical inputs and structural assumptions, conditional on the named modeling and nearest-neighbor premises; Bell-inclusive existence additionally requires H-Bell and Bell-inclusive uniqueness is not established. Theorem 24 identifies generators of the observables-preserving kernel of the dilation-datum/local reconstruction; their completeness is open. The synthesis claim is therefore structural rather than rhetorical: the same finite deterministic object supports the visible stochastic/quantum projection, the boundary-thermodynamic gravitational projection, and the lattice gauge projection. The traditional QM–GR "unification problem" is correspondingly recast as a category error between two projections of one object, while the theorem statuses and remaining conditional steps are kept explicit.

1. Introduction

The standard physical picture of the world contains two fundamental theories. Quantum mechanics describes the microscopic regime in terms of unitary evolution on a Hilbert space, superposition, and Born-rule probabilities. General relativity describes the gravitational regime in terms of a deterministic, classical, dynamical metric obeying Einstein's equations. The two are mathematically and conceptually incompatible. Quantum mechanics has no preferred temporal foliation and no observer-independent state; general relativity has both. Quantum mechanics treats measurement as a primitive operation; general relativity treats it as a process within the dynamics. Quantum mechanics is linear; general relativity is not. Every attempt to unify them — to quantize gravity, to geometrize quantum mechanics, to merge the two into a deeper underlying theory — has produced either inconsistencies, ambiguities, or empirically unfalsifiable conjectures. The unification problem has resisted solution for nearly a century, and the most active current programs (loop quantum gravity, string theory, causal dynamical triangulations, asymptotic safety) have produced neither a quantitative empirical confirmation nor a clean conceptual resolution.

This paper argues that the QM-GR incompatibility is not a technical problem to be solved by finding the right mathematical framework. It is a category error: an attempt to merge two descriptions that turn out to be projections of the same underlying object viewed from different perspectives, rather than two competing accounts of the same regime. The framework developed in this four-paper synthesis to which this paper belongs — [Main], [SM], [GR], and the present paper — provides the underlying object and the projection map. The underlying object is a finite deterministic bijection (S, φ) on a configura-

tion space partitioned into a visible sector accessible to an embedded observer and a hidden sector beyond the observer's causal reach. The projection from the underlying object to the observer's reduced description is the trace-out over the hidden sector. [Main] proves the exact finite observable-law equivalence $S \iff D \iff Q_{\mathrm{fb}}$: finite stochastic laws, finite reversible realizations, and fixed-basis unitary/Born representations coincide. The representation itself is universal; the memory-bearing OI sector is singled out instead by history read-back, which forces hidden predictive memory. For spatial systems the same construction supplies graph causal cones for finite-range couplings. What remains conditional on the quantum side is the strongest operational reading — one common standard quantum instrument/composite theory for the whole coherent intervention family. Bell is separate: the response-complete deterministic layer keeps measurement independence and reproduces quantum Bell violation only on the ontically parameter-dependent branch of [Main §3.3], with operational no-signaling. The projection from the same underlying object to general relativity is the thermodynamic limit at the partition boundary, formalized in [GR] as the cosmological-horizon application: applied to the cosmological horizon as a causal partition, the framework relates the action scale to the discreteness scale through the calibration $\hbar = c^3\epsilon^2/(4G\sigma)$, which fixes only $\epsilon/\sqrt{\sigma} = 2\,l_p$ for an entropy σ per boundary mode, and recovers the Bekenstein-Hawking entropy with the $1/4$ coefficient as a consistency relation of that calibration, both conditional on H-slope [GR §3.2], and proposes a dissolution of the cosmological constant problem conditional on the covariant coupling step [GR §6.2]. The Standard Model emerges from the same bijection on a cubic lattice with the wave equation as substratum dynamics: the cubic lattice is selected by the observed gauge group (the SU(3) factor requires the cubic crystal system), and — given that lattice — three generations (physical reading conditional on H-spin', [SM §4.7]; chirality on H-χ', [SM §4.8]), the Higgs as composite, and the $(3, 2, 1)$ gauge structure follow as forced consequences [SM]; strong CP is open there (H-top, H-det). The framework's hierarchical structural realism and its relationship to other unification programs are developed in the fifth core paper [Structure].

The present paper develops the substratum-level results that make these three emergences — emergent QM in [Main], the Standard Model derivation in [SM], and the gravitational sector in [GR] — a single construction rather than three separate applications of the same formalism. The two technical results that do this work are the *reconstruction theorem* (§3) and the *substratum gauge group* (§4). The reconstruction theorem shows — under its stated conditions (M1-T; C2-mixing; the Stage-2 structural inputs, including the nearest-neighbor premise) — that observed physics fixes the **local propagating lattice/gauge residue**. Bell violation enters separately through M1-B: a common measurement-independent deterministic completion exists only conditional on H-Bell, and no theorem here makes the Bell-inclusive completion unique. The substratum gauge group is the kernel of this inverse map; §4 identifies three families of transformations of (S, φ) in it, one of them (alphabet change) under the alphabet-size hypothesis of [SM §2.7], whose exhaustion of

all observables-preserving operations of the dilation-datum substratum is open. The combination of these two results turns the framework's three derivations into a single object: the SM gauge group is not chosen from a landscape but follows from the construction once the cubic lattice is selected from observation (§3.6) and conditionally on H-link and H-cust [SM §4.5–§4.6], the gravitational thermodynamics are not appended to a quantum description, and the emergent QM is not postulated as a starting point. All three emerge from the same (S, φ), and the substratum gauge group is the symmetry that makes the relationship between them exact rather than approximate at the level of the reconstructed local residue (§7).

The synthesis claim that follows from these results — developed in §5 — is that quantum mechanics and general relativity are not two theories awaiting unification but two projections of the same construction, viewed at different levels of description. Quantum mechanics lives at the level of the visible-sector trace-out: the embedded observer sees the bijection through the hole of finite causal access, and the resulting compressed description admits the universal unitary quantum representation. General relativity lives at the level of the boundary thermodynamics: the same causal partition that produces, at the visible-sector level, the reduced description admitting the quantum representation produces classical horizon thermodynamics at the boundary level, and Jacobson's thermodynamic argument promotes this to Einstein's equations. The two descriptions are not in tension because they refer to different objects in the construction — not different *aspects of the same regime*, which would invite unification, but different *projections of the same substratum*, which makes unification a category error. The substratum gauge group makes this precise: at the level of (S, φ), there is no quantum and no classical, no metric and no wave function; there is only the bijection and its symmetries. The trace-out at the visible-sector level produces one set of derived structures (the wave function, the SM gauge group, particle content), and the thermodynamic limit at the boundary level produces another (the metric, $\hbar$, the BH entropy, the dynamical dark energy), and the framework provides the construction that makes both of these descriptions exact projections of the same object. The same construction gives a third projection (§5.4): the C1–C4 observer-selection theorem of [Main, §4.6], which is time-symmetric and selects where observers can exist, read against the apparent-horizon area, which is non-decreasing on an expanding homogeneous general-relativistic background satisfying the null energy condition (background kinematics, not a substratum prediction); a common mechanism aligning the cosmological, thermodynamic, memory and measurement arrows is not established.

The paper is organized as follows. §1.1 states the framework's primary metaphysical commitment and its relation to explanatory and empirical claims. §2 develops the interpretation of (S, φ) as a finite lossless memory with partial read-write access and distinguishes the physical implementation question from the fixed-interface invariances of §4. §3 states the reconstruction theorem (Theorem 23) with its conditions. §4 states the substratum invariance results (Theorem 24), their open completeness question and the three-level gauge comparison.

§5 develops the synthesis claim, citing [Main], [SM] and [GR] for the empirical content. §6 discusses structural realism, the ontological hierarchy and the measurement problem. §7 concludes.

1.1 The metaphysical commitment

The framework's primary metaphysical claim is *ontic structural realism applied to the reconstructed local-sector equivalence class*. What is real at this level is the gauge-invariant local lattice/gauge structure fixed by Theorem 23, not any specific substratum representative. The substratum gauge group (§4) identifies representative freedoms that are observably irrelevant. H-Bell is a separate compatibility condition for embedding the Bell-violating composite structure into the same deterministic ontology; Theorem 23 does not prove that the Bell-inclusive completion is unique. A stronger claim — realism about a particular full Bell-complete bijection — therefore outruns the reconstruction theorem.

Stated at its most distilled, the framework's foundational commitment — what [Main §1.2] gives as "observation occurs" — is itself structured: it is two axioms. The first, that tokened differentiation occurs, is indubitable and conceptually primitive. The second, that differentiation recurs — across more than one domain, minimally configuration and time — is a substantive posit: not indubitable, not derivable from the first (a no-go result establishes this), though presupposing it. The commitment is single in the sense that the two axioms invoke one principle; it is, in count, two, and the second is the framework's one non-evidential foundational posit. Part I of the companion paper *Physics Modulo Gauge* itemizes what the first axiom forces as lemmas and what the second adds. (One item of that itemization is a reading of the primitive, adopted here explicitly: the observer's embeddedness — the proper-part decomposition $V \subsetneq S$ — belongs to the first axiom, because "tokened differentiation" is read perspectivally, as a registering from a locus against a remainder. On a thin reading it would instead be a third axiom; the framework adopts the perspectival reading, so the count is two. The choice is presentational, not empirical, and changes nothing downstream of (S, φ, V).)

The structural-realism commitment operates at multiple levels. Within OI's specific universality class, the framework's theorems license realism about $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$. Across structural classes — at the level of partial-trace observational features that any embedded-observer system must respect — broader structural realism applies, with the framework's universality-class equivalence ([Structure §9]) being the appropriate gauge equivalence. The full multi-level structure is articulated in [Structure], which canonicalizes both the observation hierarchy and the gauge hierarchy and their orthogonality, and develops the cross-framework analysis through universality classes of embedded observers. The substratum-class commitment of this paper is one level of a broader hierarchical structural realism: realism at multiple combinations of observation-hierarchy depth and gauge-hierarchy breadth, with the framework's empirical content concentrated at the deepest observation level (Level D in

[Structure §2]) and the most class-specific gauge level (Level G3 in [Structure §2]). The articulation here focuses on this concentrated level because that is where the reconstruction theorem and substratum gauge group operate; the broader hierarchical content is developed in [Structure].

Two positions often proposed as alternatives to structural realism are, in this framework, consequences of it. *Explanatory economy*: if the equivalence class is real, the emergence of quantum mechanics in [Main §3] is a theorem about representations of a real structure, not a coincidence requiring additional interpretation, and the framework's reduction of postulates follows from that uniqueness — under Theorem 23's stated conditions — rather than standing as an independent virtue. *Empirical predictions*: if the class's structure is real, its structural features — the dimension $d = 3$, the $K = 2d = 6$ internal components, the cubic decomposition $(3, 2, 1)$, the wave equation, the cosmological-horizon partition — imply specific numbers at the observable level, giving the twenty-two quantitative predictions of [SM §7] and the gravitational predictions of [GR §7]. The three framings (realist, explanatory, empirical) answer three different questions — what is real, what is gained over standard QM, what is testable beyond QM — rather than providing competing answers to the same question. The rest of this paper develops the structural and explanatory consequences; [SM] and [GR] develop the empirical consequences.

---

2. The Physical Interpretation of (S, φ)

Storage and memory. S is the set of all distinguishable states: *finite capacity*. φ is a bijection: *perfect memory* — information is never created or destroyed. Together, (S, φ) is a finite lossless memory. The partition V defines the observer's *access* — both read and write. The observer reads the visible sector and, through the coupling H_{int}, visible operations write correlations into the hidden sector. The slow-bath form of (C2) is one mechanism that preserves those writes until readback. C4 then makes the visible law genuinely history-sensitive; on a fixed finite recurrent representative any non-permutation rooted map, which that readback supplies, forces indivisibility somewhere in the recurrence cycle. The resulting finite observable law has an exact fixed-basis unitary/Born representation by the foundational equivalence theorem. The unresolved step is not the existence or quadratic form of that representation, but its simultaneous extension to the full coherent local instrument/composite algebra. Interference in the mechanism picture is write-then-read: information deposited in the hidden sector during one transition is retrieved later and changes the visible transition statistics.

The 10^{122} CC discrepancy is the compression ratio between total storage and readable storage. The dark sector is the gravitational effect of the unreadable storage. The Bekenstein-Hawking entropy is the storage capacity of the partition boundary. The action scale $\hbar$ is the conversion factor between storage

geometry and read-write statistics.

Remark (the domain of the local lattice dynamics). The wave equation of [SM §4.1] — the unique second-order reversible nearest-neighbor dynamics, up to the coefficient α, compatible with center independence, isotropy, and linearity — governs the full **local lattice sector** used by the SM/GR construction, not the visible degrees of freedom alone. Visible and hidden lattice degrees of freedom in that sector run the same wave equation; the partition V imposes observer access without changing the local update. This is enough for (i) the trace-out of [Main] over lattice-hidden degrees of freedom, (ii) the nested local trace-out of [GR §8.4], and (iii) the link-carrier construction of [SM §4.7.1.2]. It is not enough for the Bell-violating deterministic completion under measurement independence: a nearest-neighbor update, whose cone advances one edge per update, together with interventions each reading and writing only its own wing and readouts outside the remote setting's graph cone, falls under [Main §3.3]'s Bell ceiling and gives $|S_{\mathrm{CHSH}}| \leq 2$. The one-object synthesis therefore additionally requires H-Bell (§3.3): the state-dependent graph must realize Bell-relevant preparation-indexed parameter dependence with operational no-signaling while preserving the metric/Ollivier–Ricci continuum limit required by the GR projection.

The substrate objection. A structural description (S, φ) specifies dynamics rather than the material implementing it. If two implementations induce identical data for specified preparations and readouts, every decision based only on those data gives the same answer for both; this is the factoring argument of [GR §6.3]. An independently prepared, normalized, exactly decoupled finite ancillary factor leaves the original reduced channel unchanged by the partial-trace identity in Theorem 24(iii). This is a fixed-interface invariance, not a proof that every deeper implementation has identical observations or that the explanatory regress terminates. A complete observable-to-substratum classification, with its admissible interfaces and their correspondence specified, remains open.

Relation to computation. A Turing machine has storage, a read/write head and an update rule. These provide a vocabulary for the finite OI configuration, its visible access and its bijection. The exact OI recurrence argument uses finiteness of the total carrier, whereas the idealized Turing tape is unbounded. Reversible Turing machines provide lossless-update examples, but this comparison does not identify a finite hidden-state inference problem with the halting problem. An infinite OI extension requires a separately specified normalized state or measure and well-defined dynamics. The boundary-only dependence approximation alone does not supply such an extension.

Remark (effective finiteness and the computation comparison). The finite visible sector and boundary layer can determine the leading-order accessible-time law under [GR §3.2]'s hypotheses. This is a comparison of observed laws, not a proof that total-carrier finiteness is physically unnecessary or that every infinite tape realizes the same law. The exact normalized-product-ancilla identity of Theorem 24(iii) concerns finite added factors. The representation result

of [Main §3.4] covers finite observable laws, while the history-readback sector retains its separate memory hypotheses.

The arrow of time. The substratum has no arrow of time — φ and φ^{-1} are equally valid. Observers satisfying C1–C4, with C2 read in its timescale form, are nonetheless structurally restricted to the non-equilibrium phase of (S, φ): the theorem in [Main, §4.6] shows, under its effective-mixing hypothesis and for the coarse-grained description generated by iterating the one-step effective channel, that $\tau_B(V)$ is bounded by the local mixing time on equilibrium microstates, so C2 fails there in its timescale form, record persistence being degraded rather than shown to fail, wherever that mixing time is short compared with τ_S — and the readback gap of C4, the condition the characterization registers, is suppressed exponentially in τ_S/τ_B at every order $k \geq 2$ by the erasure lemma of [Main §2.3]. In that description, this replaces the low-entropy "past hypothesis", in its role of selecting observer-compatible configurations, with a structural observer-selection rule and excludes Boltzmann brains structurally rather than anthropically. The rule is time-symmetric: the direction in which coarse-grained entropy increases rests on a low-entropy or product boundary condition, not on the rule (§5.4).

The incompleteness family. The framework's central result belongs to a family of impossibility-with-structure theorems. Gödel: a formal system cannot prove all truths about itself — the unprovable truths have rigid structure. Turing: a computer cannot decide all questions about its own behavior — the undecidable problems have rigid structure. OI: an observer embedded in a deterministic system cannot access the complete state — the emergent description has rigid structure (quantum mechanics). The common structure is *self-reference under finite resources*.

For a fixed model, a hidden-sector state h is a determinate quantity, and its effect on later observations depends on the dynamics and the allowed records. In a known finite model, retained records identify a hidden quantity exactly when it is constant on the compatible states (Proposition 24.5); for $\varphi(v, h) = (h, v)$ with readout v, two readings determine h_0. This differs from redundancy between descriptions relative to specified observations. Theorem 24(iii)'s independently prepared finite decoupled enlargement changes the ancillary cardinality while preserving the original coherent channel family. Alphabet change preserves observables only under the alphabet-size hypothesis of [SM §2.7], which separates results proved for every $q \geq 2$ from those that carry q. A slowly coupled deep sector instead preserves only the leading-order law in its stated regime. None of these comparisons makes every inaccessible quantity physically empty or classifies all underlying implementations.

Mathematics and physics. The trace-out performs a Jordan-Chevalley projection ([SM, Appendix A]): it extracts the semisimple part of the dynamics and erases the nilpotent monodromy. Physics is the semisimple shadow of mathematics — the diagonalizable spectral data, projected by the trace-out and organized by the gauge group's representation structure. The reconstruction theorem (be-

low) proves — under its stated conditions — that the **local lattice/gauge residue** is determined up to gauge. It does not prove uniqueness of the Bell-inclusive full substratum.

---

3. The Reconstruction Theorem

The forward direction — from (S, φ) to observed physics — is established by [SM §§3.1, 4–5] and the companion paper [Main]. The converse question is which parts of (S, φ) the observed physics fixes. The reconstruction proceeds in three stages. The composite theorem (Theorem 23) aggregates the stages and separates the uniquely reconstructed local lattice/gauge residue from the Bell-inclusive completion, whose existence is conditional on H-Bell and whose uniqueness is open.

3.1 Inputs to the reconstruction

The reconstruction takes two kinds of inputs: empirical observations and structural assumptions. Being explicit about both is essential to the uniqueness claim.

Empirical inputs (facts about the observed universe):

(E1) **Unitary quantum mechanics.** The observed physics is described, to the precision tested, by quantum mechanics: states as vectors in a complex Hilbert space, unitary time evolution, self-adjoint observables, Born-rule outcome statistics.

(E2) **Bell violations.** The observed correlations violate Bell inequalities, ruling out local hidden-variable theories with factorizable distributions.

(E3) **Finite boundary entropy.** The entropy of any bounded region is finite and scales as the area of its boundary. This is a theoretical input — the holographic bounds [1, 2], which are not themselves observed — applied to the cosmological horizon, whose finite area makes the bound apply.

(E4) **Spatial isotropy.** Observed physics is rotationally invariant; no spatial direction is preferred.

(E5) **Propagating gravity.** Gravitational degrees of freedom propagate (gravitational waves are observed). This requires $d \geq 3$, since the Weyl tensor vanishes for $d \leq 2$. (Used in Stage 2(a) below; one of the physical inputs to the conditional dimensional selection.)

(E6) **Stable matter.** Atoms, planets, and bound states exist on observed timescales. This requires $d \leq 3$, since the Coulomb potential in d dimensions gives unstable atoms for $d \geq 4$ (Ehrenfest [8]) and gravitational orbits are unstable for $d \geq 4$ (Tangherlini [9]). Without stable matter, no embedded

observers exist to define the partition. (Used in Stage 2(a); second forward filter for $d = 3$.)

(E7) **Boundary-entropy concordance.** The cosmological parameter inferred from observation under the standard cosmological model satisfies $\rho_s/\rho_{\text{crit}} \approx 1$ (spatial flatness near critical density); it is an inferred quantity, not a directly measured one. In d spatial dimensions, the general-d form of this ratio is $\rho_s/\rho_{\text{crit}} = 2/(d-1)$, which equals unity only for $d = 3$. (Used in Stage 2(a); third forward filter for $d = 3$. See [SM §3.2] and [GR §7.2] for the derivation.)

Structural assumptions (restrictions on the class of candidate substrates (S, φ)):

(A1) **Finiteness.** The configuration space S is finite. E3, read as a Hilbert-space dimension cutoff $\dim \mathcal{H} \leq e^{A/4}$, bounds the observationally relevant visible sector and boundary layer $\mathcal{C}_V \times \mathcal{C}_B$ under the boundary-only dependence hypothesis of [GR §3.2]. This bound alone does not establish finiteness of every underlying sector. Theorem 24(iii) gives an exact freedom to adjoin independently prepared finite decoupled factors at a fixed interface. Infinite extensions require specified normalized measures or states and well-defined dynamics; the finite identity does not supply that construction. A slowly coupled deep sector changes the transition law by the stated $\mathcal{O}(t/\tau_B)$ correction, so its comparison is only at leading order on accessible timescales. A1 remains the finite-carrier premise for the recurrence arguments.

(A2) **Determinism.** $\varphi : S \to S$ is a bijection (deterministic, reversible dynamics). (Two-part status, per the dilemma argument of [Main §2.2]: a non-injective descent either leaves a statistical trace — excluded where the observed statistics are of the doubly stochastic class that the uniform-prior marginal of a bijection produces — or leaves none, in which case it is removable, every map on a finite set restricting to a bijection on its eventual image, where all recurrent registered history lives. Bijectivity for the dynamics governing registered history is thus partly structural and partly anchored by one empirical input, the observed statistics in that scope; Main's honest-accounting ledger records the same division.) The anchor is empirically live: its current frontier is nanoparticle matter-wave interference at macroscopicity $\mu = 15.5$ — sodium clusters above 1.7×10^5 amu held in 133-nm path superpositions [4] — an order of magnitude beyond the prior benchmark, and the direct one-sided test against the intrinsic-collapse alternatives that would break it.

(A3) **Bounded coupling degree.** Each site is coupled to a bounded number of neighbors in φ's action. (Required for locality and the emergence of a coupling graph with well-defined dimension. The physical content is boundedness itself; bounded-degree graphs with the same growth exponent, spectral mode count and statistical isotropy each admit a finite-law representation and share the growth exponent (Theorem 24(iv)), but need not give the same $T_{ij}(t)$, and the continuum-curvature step requires its own conditions [SM §3.1].)

(A4) **Center independence.** The dynamics φ does not depend on a choice of

"center" site, in two senses used below: φ commutes with lattice translations up to gauge (the coupling graph is a finite periodic lattice, the quotient of a rank-d lattice by a finite-index sublattice, d being its dimension), and the update at a site contains no explicit copy of that site's present value ([SM §4.1]). The two senses are logically independent for rules of the second-order form $x(t{+}1) = F(x(t)){-}x(t{-}1)$ over $\mathbb{Z}_q$ on a periodic cubic lattice of side at least 3, with gauge acting by site-wise bijections of each site's (previous, current) pair. Such a gauge preserves which sites each update reads, so a rule that commutes with translations up to gauge reads its sites in a translation-invariant pattern. The rule $x_i(t+1) = x_i(t) + \sum_{j\sim i} x_j(t) - x_i(t-1)$ commutes with translations exactly and reads $x_i(t)$; the same rule with the $x_i(t)$ term dropped and one neighbour omitted from the sum at a single site reads no site's own present value, and its pattern of read sites is not translation-invariant. (The second sense is what the wave-equation derivation of Stage 2 uses. Its observational anchor is the homogeneity of physical law — no experiment has found a preferred location in the laws — the cosmological principle's complement to E4's isotropy; whether it is independently warranted by the observer architecture remains the question flagged in §1.)

(A5) **Linearity.** The wave equation for φ is linear. (Amplitude-scale gauge invariance — that the field-value scale is unphysical — restricts the update to an affine one via the linearity-equivalence lemma of [SM §4.1], so A5 is "partly derived" in exactly that sense: its affine part is necessary *given* that gauge principle, and the further restriction to linear is imposed. Whether the q-size gauge freedom of [SM §2.7] entails amplitude-scale gauge is an identified open step; pending it, non-affine alternatives are excluded precisely to the extent amplitude-scale gauge is assumed, and would otherwise require a separate derivation.)

(A6) **Background independence.** The dynamics is covariant under spatially-varying internal-index transformations: a site-dependent transformation $G(\mathbf{n})$ of the internal index is a symmetry of the dynamics provided the coupling data carried on the links are transformed with it, $M(\mathbf{n}, \hat{e}_j) \to G(\mathbf{n})\, M(\mathbf{n}, \hat{e}_j)\, G(\mathbf{n} + \hat{e}_j)^{-1}$ ([SM §3.1]). The content of the assumption is this covariant interface — the coupling is link-valued data on which the transformations act — and once it is in place the local transformation law imposes no further condition on the rule. A6 is not the stronger fixed-background condition that the same transformations leave the dynamics invariant with the coupling held fixed and preserved pointwise: for an invertible constant coupling that condition forces the transformation to agree across every coupled pair of sites, and at the symmetric point of the coupling matrix it fails for every transformation that differs across a coupled pair. The global commutant symmetry — the transformations constant across the lattice that leave the coupling itself unchanged, the specialization of A6 in which nothing needs transporting — is what the local gauge reading of [SM §3.1] proceeds from. The state-dependent coupling graph of [SM §3.1], under which the graph itself evolves with the state, is a separate principle that shares the name and is not A6.

The theorem's uniqueness claim holds under E1–E7 **and** A1–A6 jointly, with the nearest-neighbor premise of Stage 2(b); removing any of A3–A6 either requires new derivations or weakens the uniqueness. The set A1–A6, with that premise, is sufficient for the reconstruction but is not proved to be minimal: A4 (center independence) and A6 (background independence) overlap in physical content (A6 promotes the symmetry that A4 constrains), and A5 (linearity) is partly derived — amplitude-scale gauge invariance restricts the update to an affine one via the linearity-equivalence lemma of [SM §4.1], the further restriction to linear being imposed — with whether [SM §2.7]'s q-size gauge entails amplitude-scale gauge left as an identified open step. A tighter axiomatization may be possible but is not pursued here; the reconstruction's validity depends on the sufficiency of A1–A6 and E1–E7, not on their independence. The empirical inputs E5, E6, E7 (propagating gravity, stable matter, boundary-entropy concordance) provide explicit forward filters for $d = 3$ — see Stage 2(a) below.

Critical dependencies. Theorem 23's proof relies on the following prior theorems, whose individual correctness is assumed:

- [Main §2.3] history readback for the **memory-bearing temporal** sector used in Stage 1 (a modeling premise about the temporal data, not an inference from Bell violation)
- [Main §3.3] Bell ceiling and the adopted measurement-independent, ontically parameter-dependent Bell branch (a separate spatial-composite premise; Bell violation is not used as a witness of temporal P-indivisibility)
- [Main §3.2] Forward Stinespring dilation (bijection → CPTP channel) and reverse-direction lemma (one-step doubly stochastic matrix → uniform-prior bijection; exact for rational entries, to arbitrary precision otherwise)
- [Main §3.4] Finite-horizon process dilation (stochastic process → bijection, per finite horizon)
- [Main §3.4] Characterization theorem (accessible non-Markovianity — the framework's finite-horizon visible-law memory notion, not generic quantum non-Markovianity — $\iff$ C1/C3/C4 per finite accessible horizon; C2's slow-bath necessity conditional on the hidden-sector mixing hypothesis, ETH its physical motivation, its persistence form holding without it)
- [SM §3.2] Coupling-graph dimension → d = 3
- [SM §4.1] Center independence + isotropy + linearity → wave equation
- [SM Theorems 5–15] Gauge group, generations, hypercharges derivation chain
- [SM Theorems 17–21] T-invariance → $\theta \in \{0, \pi\}$ and reciprocal transition counts; $\bar{\theta}$ open (H-top, H-det)
- [GR §3] Gap equation from thermal self-consistency → $\hbar = c^3\epsilon^2/(4G\sigma)$

Theorem 23's confidence is bounded above by the minimum confidence in these dependencies.

3.2 Stage 1: Observed QM → deterministic embedding with C1–C4

Inputs: E1 (unitary QM), E2 (Bell violations), E3 (finite boundary entropy), A1 (finiteness), A2 (determinism), and M1, in two clauses: **M1-T**, the observed temporal quantum data used in Stage 1 lie in the accessible history-readback sector of [Main §2.3–§3.4]; and **M1-B**, Bell-violating composites take [Main §3.3]'s measurement-independent, ontically parameter-dependent branch with operational no-signaling. M1-T and M1-B are logically distinct; Bell violation is not taken to imply temporal P-indivisibility.

Output: There exists a triple (S, φ, V) with S finite, φ a bijection on S, and $V \subset S$ a distinguished subset such that: - The observed quantum dynamics is the reduced description obtained by tracing out $S \setminus V$ from the deterministic evolution under φ. - The triple satisfies C1 (non-trivial coupling), C2 (record persistence — the slow bath one realization), C3 (sufficient hidden memory capacity), and C4 (history readback) for the M1-T temporal sector: [Main §3.4]'s universal hidden-memory theorem — accessible non-Markovianity $\Longleftrightarrow$ C1/C3/C4 realization per accessible horizon, with $M_t > 0$ forcing unavoidable hidden predictive memory — makes readback-relevant hidden memory unavoidable in every faithful deterministic realization. M1-B separately requires Bell-relevant cross-wing coupling/parameter dependence and does not supply C4.

Conditional structure. C1 and C4 for the memory-bearing temporal sector are derived from E1–E3, A1–A2 together with **M1-T** — C4 via the hidden-memory theorem applied to the stipulated accessible temporal memory; conditional on M1-T, they hold without further hypothesis. C3 holds in every faithful realization by the capacity bound of [Main §3.4]. **M1-B** is separate: it maps E2 to the ontically parameter-dependent Bell composite and supplies an independent cross-wing-coupling requirement, not C4. C2 takes the two forms of [Main §1.3]. (C2-structural), retention of the record of visible history to its readback, holds by reversibility in every bijective realization. C2's necessity holds in its persistence form without further hypothesis: in the coupling-step model of [Main §3.4]'s physical-memory theorem, observed conditional memory $M_k = I(X_{<k}; X_{k+1} \mid X_k) > 0$ at an accessible order k places some realized hidden conditional at that coupling time at least $\varepsilon_{\min}(M_k, n_V)$ from its reference prior (the physical-memory theorem). Its slow-bath form is conditional on the quantitative mixing hypothesis of [Main §3.4] — uniform total-variation closeness of the hidden sector's post-relaxation conditionals to a single bath distribution — under which the same bound makes $\varepsilon_{\min}(M_k, n_V)$ a lower bound on the relaxation residue at coupling time k, hence on the bath's conditional mixing time, and a departure δ from the homogeneous Markov benchmark at step order k forces $\varepsilon \geq \delta/(k-1)$ (the bound $\|T^{(k)} - T^k\|_{\mathrm{TV}} \leq (k-1)\varepsilon$, constant optimal at $k = 2$); without it, accessible conditional memory does not bound the bath's mixing time. ETH is the physical case for the hypothesis, as discussed in [Main §3.4]'s "Remark (Status of ETH)." ETH is a well-supported conjecture for generic chaotic many-body Hamiltonians and is the standard assumption for the cosmological-horizon complement; relaxations or failures of ETH would

require a separate argument. The framework's gauge-group derivation through Stage 2 inherits this conditional.

Bell–gravity compatibility hypothesis (H-Bell). The Stage-2 reference nearest-neighbor wave-equation graph has a pointwise causal cone advancing one edge per update; whether that cone or the lift's long-wavelength speed $1/\sqrt{3}$ is the physical c is not derived [Main §3.3]. Under measurement independence, setting interventions that each read and write only their own wing and ± 1 readouts outside the remote setting's graph cone, [Main §3.3] therefore makes that **reference graph** Bell-local and caps CHSH at 2. The framework nevertheless already makes the graph state-dependent: the discrete Einstein equation of [SM §3.1] is formulated on $G(x)$, and its mod-q spatial-Markov proof explicitly assumes uniform initial conditions. M1-B may therefore be implemented by **preparation-indexed adjacency**: the vacuum/uniform reference class retains the geometric bounded-range graph used by the local SM/GR derivation, while an entangling preparation supplies bounded-degree Bell-relevant long edges (or an equivalent ontically nonlocal response map) furnishing parameter dependence without operational signaling.

H-Bell is the remaining compatibility statement for that branch: the preparation-indexed graphs must (i) supply the required ontic parameter dependence, (ii) preserve operational no-signaling, and (iii) satisfy a metric/Ollivier–Ricci stability condition strong enough that the continuum-curvature step of [SM §3.1] remains valid on the state classes to which the Einstein reconstruction is applied. The graph-boundary area law is not the primary unresolved issue: $|\partial V|$ is defined on $G(x)$ and therefore counts preparation-induced crossing edges. The open step is the identification of that graph boundary/curvature with continuum geometric area/Ricci curvature. [SM §3.1]'s sparse local-contamination estimate, $L_N m_N/N \to 0$, is a sufficient condition for typical mesoscopic neighborhoods to remain reference-like, but it does not prove global curvature convergence, control defect endpoints, or establish the integrated-curvature limit. H-Bell is therefore an explicit curvature-stability/composite-existence hypothesis, not a consequence of A1–A6 or of the nearest-neighbor wave equation.

The mixing hypothesis for the realized hidden sector remains open. The exact linear rule of [SM §4.2] carries commuting structure from translation invariance: each symmetric translation-invariant coupling B commutes with the neighbour sum and gives a conserved quadratic invariant, a sum of local densities when B is local, and at prime q not dividing the lattice size its sufficiently large energy shells split into many cycles. The rounded bijection of [SM §4.2] behaves chaotically at the sizes measured; these diagnostics concern that candidate map and do not establish mixing of the realized hidden sector. Neither statistical isotropy nor bounded coupling degree supplies the missing conditional-distribution mixing bound. The conditional attaches to the slow-bath form of C2's necessity direction: observed conditional memory gives a lower bound on the bath's conditional mixing time within the class satisfying that hypothesis.

C2's persistence form, the readback implication and the finite-horizon representation retain their own stated hypotheses and do not require ETH. ETH for few-body observables is itself insufficient for the quantitative mixing condition when conditional-distribution memory survives ([Main §3.4], Remark (role of ETH)).

Derivation. The observed quantum dynamics provides, for each finite experimental context, a stochastic law on a finite visible configuration space. By [Main]'s exact finite-horizon equivalence, this law has a finite reversible deterministic realization and an exact fixed-basis unitary/Born representation. The representation step is universal and therefore does not itself reconstruct the nontrivial OI sector. The additional modeling premise M1 separates two empirical structures. **M1-T** places the temporal data used for Stage 1 in the accessible history-readback sector; [Main §2.3] then proves that on a fixed finite representative a non-permutation rooted map, which C4 readback supplies, forces indivisibility somewhere in the recurrence cycle. **M1-B** places the Bell data on [Main §3.3]'s measurement-independent, ontically parameter-dependent branch; Bell violation therefore certifies Bell-relevant cross-wing coupling but is not used to infer temporal P-indivisibility. By [Main]'s finite-horizon equivalence ($S \Rightarrow D$, the process-dilation and response-table constructions of [Main §3.4]), the temporal law admits a realization as the visible marginal of a deterministic bijection φ on an enlarged finite space $\mathcal{C}_V \times \mathcal{C}_H$, with a hidden prior independent of the visible initial state carried as part of the realization datum; for a single one-step doubly stochastic matrix the reverse-direction lemma ([Main §3.2]) gives the uniform-prior form, exactly when its entries are rational and to arbitrary precision otherwise. This delivers the triple (S, φ, V) with $S = \mathcal{C}_V \times \mathcal{C}_H$ and $V = \mathcal{C}_V$. Non-permutation $T(t)$ in the M1-T temporal sector gives the corresponding C1 witness; independently, M1-B's ontic parameter dependence requires Bell-relevant cross-wing coupling. The necessity analysis of [Main §3.4] establishes, for every faithful realization of the resulting dynamics, C3 (the data-processing bound $\log_2 |\mathcal{C}_H| \geq I^*$, with no mixing hypothesis) and (C2-structural) (retention of the record of visible history to its readback, supplied by reversibility; record persistence is required for the non-Markovian returns); in the coupling-step model of [Main §3.4]'s physical-memory theorem, observed conditional memory $M_k > 0$ at an accessible order k keeps some realized hidden conditional at least $\varepsilon_{\min}(M_k, n_V)$ from the reference prior at that coupling time (C2's persistence form, with no mixing hypothesis), and under the hidden-sector mixing hypothesis ([Main §3.4]; ETH its physical motivation) this lower-bounds the bath's conditional mixing time (the slow-bath form).

Stage-1 gauge statement. For the **finite channel/dilation datum actually used by the temporal reconstruction**, the familiar dilation freedoms include the stated substratum-gauge directions: hidden-sector basis/relabeling freedom and observationally inert deep-sector enlargement; equal channel families need not have unitarily conjugate dilations (§4). This is not uniqueness of the full triple (S, φ, V) across all coherent composites. In particular, it does not select a preparation-indexed Bell adjacency rule, and the operational lift still

requires its separately stated finite-predictive-dimension, local-tomography and purification hypotheses.

Status: Theorem for the scoped finite dilation datum; no Bell-inclusive Stage-1 uniqueness claim.

3.3 Stage 2: Embedding + isotropy → specific lattice + gauge structure

Inputs: Stage 1 output + E4 (spatial isotropy) + E5 (propagating gravity) + E6 (stable matter) + E7 (boundary-entropy concordance) + A3 (bounded coupling) + A4 (center independence) + A5 (linearity) + A6 (background independence) + max-lattice-speed condition (for the wave-equation coupling constant; see step (b) below). Identification of this local lattice output with the Bell-violating Stage-1 completion additionally assumes H-Bell.

Output: The local propagating substratum sector is a $d = 3$ cubic lattice bijection with: - $K = 2d = 6$ internal components per site - Block multiplicities $(3, 2, 1)$ of the cubic-group decomposition of the link directions - Gauge group SU(3) × SU(2) × U(1) - Three generations of chiral fermions in the SM representations - One Higgs doublet $(\mathbf{1}, \mathbf{2}, +1/2)$ - Anomaly-free hypercharges $(Y_Q, Y_u, Y_d, Y_L, Y_e) = (1/6, 2/3, -1/3, -1/2, -1)$

Derivation.

(a) *Dimension.* The coupling graph inherited from Stage 1 has polynomial growth exponent d. Within A4's assumed periodic-lattice class and the continuum conditions of [SM §3.1–§3.2], the physical inputs select $d = 3$: propagating gravity (E5) requires $d \geq 3$ (the Weyl tensor vanishes for $d \leq 2$); stable matter (E6) requires $d \leq 3$ (Coulomb instability of atoms for $d \geq 4$, plus Tangherlini's gravitational orbit instability); boundary-entropy concordance (E7) gives $\rho_s/\rho_{\text{crit}} = 2/(d-1) = 1$ at $d = 3$ uniquely. The intersection of the first two filters already pins $d = 3$; the third is a sharpening consistency check that the framework passes exactly. Self-consistency excludes exponential growth (Lorentz-invariant dispersion fails, and Jacobson's GR derivation requires it) and fractal growth (the cubic-lattice symmetry needed for the cubic-group decomposition in step (d) below is absent), but not intermediate or irregular growth. The integer degree is an input through A4, whose lattice translations make the coupling graph a finite periodic lattice of rank d; Gromov's theorem, which concerns finitely generated groups, and statistical isotropy (E4) do not supply it for a general bounded-degree graph. Applied to the continuum description, conditional on the continuum step of [SM §3.1], the three filters then select $d = 3$. ([SM §3.2] presents this argument in full; the inputs E5, E6, E7 surfaced here make explicit the empirical content the filters require.)

(b) *Wave equation.* Center independence (A4, in its functional sense: no self-term), isotropy (E4), and linearity (A5) uniquely select, within the nearest-

neighbor class (an assumption beyond A3's bounded degree; see the scope remark at [SM §4.1]), the wave equation up to a multiplicative coupling constant α ([SM §4.1, Theorem]): the unique second-order linear dynamics on a lattice that is translation-invariant, isotropic, reversible and free of a self-term has the form $f = \alpha(x_1 + x_2 + \cdots + x_{2d}) \bmod q$ with coupling coefficient α. The coefficient is neither the propagation speed nor free to be set to 1: causal support travels at most one lattice edge per update whatever its value, and on the observer-level normalized branch it equals $1/d$ ([SM §4.1, Remark]).

(c) *Internal components.* Under H-link the link count gives $K = 2dm$, with $m = 1$ giving $K = 6$; coupling-degree minimization does not establish it (see [SM] Theorem 6). This locks the candidate-sector count at three — three physical generations only under H-spin' ([SM §4.7]) — (three spin-1/2 staggered tastes emerge from the $K = 6$ minimum).

(d) *Gauge group.* Cubic-group decomposition of the $K = 6$ components gives multiplicities $(3, 2, 1)$ (Theorem 7). The unitary group of the bicommutant of the cubic-group action on $V_K = V_3 \oplus V_2 \oplus V_1$ — equivalently, the stabilizer of any generic O-equivariant operator on V_K — is $U(3)\times U(2)\times U(1)$. (The strict pointwise commutant of the action is $U(1)^3$ by Schur's lemma; the gauge group is the unitary group of the block algebra $\mathrm{Mat}(3) \oplus \mathrm{Mat}(2) \oplus \mathrm{Mat}(1)$ generated by the action.) *Amplitude-scale invariance* (Theorem 24, Remark below) removes only the common overall phase: a global rescaling of the field value ϕ rescales every block together. (This is the field-value-scale freedom — the additive-automorphism relabellings of the alphabet — *distinct* from the alphabet-*size* freedom of generator (ii).) The reduction to $SU(3)\times SU(2)\times U(1)$, with the singlet-block direction as the hypercharge $U(1)$, additionally uses a specified charged matter action and anomaly cancellation ([SM §4.9, Theorem 15]). Background independence (A6) — covariance under site-dependent transformations, the link coupling transformed with them — then promotes the remaining global $SU(3) \times SU(2) \times U(1)$ to local gauge invariance ([SM §3.1]).

(e) *Matter content.* Staggered reduction gives three degenerate spin-1/2 sectors (Theorems 8–10). Fermion embedding is completed via the link-carrier construction ([SM §4.7.1.2]). Partition-spinor identification (Theorem 12) and trace-out (Theorem 13) give chirality. Anomaly cancellation (Theorems 14–15) fixes the hypercharges up to normalization and quark-singlet interchange, apart from the exceptional branch $Y_Q = 0$.

(f) *Discrete symmetries.* T-invariance of the bijection φ gives $\theta \in \{0, \pi\}$ and reciprocal visible transition counts at every scale (Theorems 17–21); it does not fix $\bar{\theta}$, which remains open under H-top and H-det [SM §5]. Reciprocity leaves the Jarlskog invariant free; only a mechanism making Y_u and Y_d simultaneously real would force $J = 0$, which is the falsification control on H-det proposals [SM, §5.3–5.5].

Uniqueness at Stage 2. Each step (a)–(f) is a uniqueness statement: the stated structure is the unique consistent choice given the inputs, step (b) within the nearest-neighbor class and step (e) apart from its exceptional branch. The full chain is therefore a composition of unique selections, so the output is unique within those qualifications.

Bell compatibility. Step (b)'s nearest-neighbor **reference** cone cannot itself realize M1-B, while retaining measurement independence, in a Bell protocol with one-step own-wing read/write setting interventions, a ± 1 outcome at each wing in every run, and readout regions outside the other wing's graph cone over its duration. Stage 2 therefore establishes the local lattice/gauge sector on the geometric reference class. The state-dependent $G(x)$ machinery leaves a native route for M1-B through preparation-indexed adjacency, but the Ollivier–Ricci continuum theorem presently cited by [SM] does not cover those long-edge perturbations. Nor does it cover the unperturbed reference graph, and there the situation is worse than open: the degree-6 cubic hop metric is exactly ℓ_1, with a scale-independent $\sqrt{2}$ diagonal stretch, so the hop-metric curvature route fails for the reference family and no $D = 3$ extension of the cited theorem would rescue it. The repair is to re-found curvature on the propagation/Laplacian geometry, which the substratum's isotropic leading dispersion already supplies [SM §3.2]. Promoting the local sector and M1-B to one common (S, φ) is therefore conditional on H-Bell's no-signaling and continuum-curvature requirements.

Status: Theorem for the local lattice sector; one-object Bell compatibility conditional on H-Bell.

3.4 Stage 3: Dynamics → emergent fundamental constants

Inputs: Stage 2 output.

Output: The emergent action-scale relation $\hbar = c^3\epsilon^2/(4G\sigma)$, equivalently $\epsilon = 2\sqrt{\sigma}\, l_p$, where l_p is the Planck length and σ the entropy per boundary mode ([GR §3.1]); it fixes only $\epsilon/\sqrt{\sigma}$, not the value of $\hbar$ ([GR §4]).

Derivation. The gap equation from thermal self-consistency of the boundary layer ([GR §3]) gives the stated relation. The derivation uses: existence of thermal equilibrium at the boundary layer (a consequence of the C1–C4 dynamics from Stage 1) and the infrared detailed-balance condition H-slope with H-count and the horizon and frame conditions ([GR §2, §3.2, §8.5]), which does not follow from φ being a bijection ([GR §3.2]) and is an assumption beyond Stages 1–2.

Uniqueness at Stage 3. The gap equation admits a unique solution under the stated inputs ([GR §3.2]).

Status: Conditional on H-slope with H-count and the horizon and frame conditions ([GR §3.2]).

3.5 The composite theorem

Lemma 23.0 (Local-sector uniqueness preservation). *Let (S_1, φ_1) and (S_2, φ_2) both reproduce the same inputs E1–E7 and satisfy A1–A6 and M1, with the hidden-sector mixing hypothesis where required, and both lie in the nearest-neighbor class of Stage 2(b). Then their reconstructed* ***local propagating lattice/gauge sectors*** *have the same Stage-2 invariants: $d = 3$, $K = 6$, multiplicities $(3, 2, 1)$, the SM gauge group and associated local representation data (hypercharges apart from the exceptional branch of Stage 2(e)) — $K = 6$, the multiplicities and the gauge group conditionally on H-link with its single-copy clause and H-cust [SM §4.5–§4.6] — and, conditional on H-slope with H-count and the horizon and frame conditions of [GR §3.2], the same Stage-3 action-scale relation $\hbar = c^3\epsilon^2/(4G\sigma)$.*

Proof. Stage 2 selects the same local cubic propagating sector and discrete representation-theoretic invariants from E4–E7, A3–A6 and the nearest-neighbor premise; hidden-sector relabeling and deep-sector enlargement lie in $\mathcal{G}_{\text{sub}}$, and alphabet freedom does under the alphabet-size hypothesis of [SM §2.7] (Theorem 24(ii)). Under H-slope and H-count, Stage 3 fixes the same gap-equation relation $\hbar = c^3\epsilon^2/(4G\sigma)$; an alphabet change can change the bound $\sigma \leq \ln q$, and the equality of $\hbar$ across it is part of that hypothesis. These statements compare the reconstructed local sector only. They do **not** compare preparation-indexed Bell edge rules or other Bell-nonlocal composite data, because the Stage-2 uniqueness argument does not act on that extra structure. □

Theorem 23 (Layered local reconstruction with Bell compatibility). *Let E1–E7, M1-T, M1-B, A1–A6, the hidden-sector mixing hypothesis for the slow-bath form of the C2 necessity direction, and the nearest-neighbor premise of Stage 2(b) hold. Then the reconstruction fixes the* ***local propagating lattice/gauge sector*** *with $d = 3$, $K = 6$, block multiplicities $(3, 2, 1)$, gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$, the stated local matter/representation content and discrete-symmetry outputs, and $\hbar = c^3\epsilon^2/(4G\sigma)$ conditional on H-slope with H-count and the horizon and frame conditions of [GR §3.2], with the same qualifications stated in [SM] for H-link with its single-copy clause and H-cust [SM §4.5–§4.6], H-spin', H-χ', the exceptional hypercharge branch $Y_Q = 0$, and the CKM-CP fork. A single measurement-independent deterministic completion that also realizes the Bell-violating M1-B composites exists only conditional on H-Bell. The theorem does* ***not*** *establish uniqueness of that Bell-inclusive full completion.*

Proof. Stage 1 supplies the finite reversible/dilation data and the M1-T temporal conditions used by the local reconstruction, subject to its explicit operational-lift hypotheses and the C2-mixing condition. Stage 2 uniquely fixes the listed local lattice/gauge invariants from those data and E4–E7; Stage 3, conditional on H-slope, fixes the relation $\hbar = c^3\epsilon^2/(4G\sigma)$. Lemma 23.0 preserves that uniqueness at the local-sector level. M1-B adds a separate Bell-relevant requirement. H-Bell states the compatibility/existence condition for realizing it through the state-

dependent graph or an equivalent ontically nonlocal response implementation while retaining the continuum-curvature limit. No theorem in the present chain selects a unique preparation-indexed Bell edge rule. Therefore Bell-inclusive existence is conditional and Bell-inclusive uniqueness remains open. □

3.6 Remarks

Remark (Evidential weight of the reconstruction). Within the nearest-neighbor class of Stage 2(b), the reconstruction uniquely derives, conditionally on H-link and H-cust [SM §4.5–§4.6], $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ with multiplicities $(3, 2, 1)$, three generations (physical reading conditional on H-spin', [SM §4.7]; chirality on H-χ', [SM §4.8]), one Higgs doublet $(1, 2, +1/2)$ (taken as input and accommodated in the E projection of the singlet taste, [SM] Theorem 11), and hypercharges unique up to normalization and quark-singlet interchange apart from the exceptional branch $Y_Q = 0$. Because the derivation is tightly constrained — no fitted parameters, no model-building choices, no landscape of alternatives at the level of the gauge sector — every independent experimental confirmation of the Standard Model's gauge structure is retroactive evidence for the derivation chain that produces it, with the strength of that evidence bounded by the chain's open links (the construction delivers a distinguished low-freedom representation rather than a uniquely selected one, and the family pattern's selection is not supplied; [SM] Theorem 15 Remark). The Standard Model is the most precisely tested mathematical structure in the history of science ($g - 2$ of the electron confirmed to 10^{-12}, electroweak precision observables to 10^{-5}, QCD cross-sections across decades of energy). This experimental record confirms the Standard Model structure; it transfers to the framework only to the extent that the derivation is unique, since the transitivity "data confirms structure, structure is uniquely derived, therefore data confirms the derivation" is only as strong as its middle premise. That premise is conditional on the argued — not proven — sufficiency of the A1–A6 class (Remark on hypothesis dependencies below) and is established operationally by the classified retrodiction record of [SM §7] (two parameter-free structural and five conditional structural entries among twenty-two classified observables) (Remark on finite observation sets below), not by the Standard Model precision record itself. The precision record is therefore necessary but not sufficient evidence for the *uniqueness* of the derivation — which is the framework's distinct claim, and the proper locus of skeptical scrutiny.

Remark (Substantive content of the uniqueness claim). A reasonable objection to any reconstruction theorem whose answer is an equivalence class under a large gauge group is that the result risks tautology: if $[(S, \varphi)]/\mathcal{G}_{\mathrm{sub}}$ is *defined* as the observational-equivalence class, then "observed physics uniquely determines the observational-equivalence class" is close to true by construction, and the gauge group $\mathcal{G}_{\mathrm{sub}}$ here is enormous (hidden-sector relabeling, alphabet change under the alphabet-size hypothesis, finite decoupled deep-sector enlargement of arbitrary size — see §4). The objection has force against a vacuous version

of the claim and must be answered by exhibiting the *non-tautological residue*: the specific structure the theorem fixes that does not follow merely from "is observationally equivalent to the data." That residue is a short list of discrete invariants. The theorem does not merely return "whatever is observationally equivalent to observed physics"; it returns specific integers and discrete structures — the spatial dimension $d = 3$, the internal-component count $K = 6$, the block multiplicities $(3, 2, 1)$, the gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$, the generation count exactly three, one Higgs doublet, and the anomaly-free hypercharge assignment (apart from the exceptional branch $Y_Q = 0$). None of these is a tautological consequence of observational equivalence: a priori the reconstruction could have returned any dimension, any gauge group, any generation count, or a continuum of them, and the content of the theorem is that the structural assumptions force *these specific discrete values* and no others. The large gauge group $\mathcal{G}_{\mathrm{sub}}$ is quotiented away precisely so that what remains invariant is this discrete list — the continuous and combinatorial freedoms (relabeling, alphabet under the alphabet-size hypothesis, deep-sector size) are exactly the directions along which the answer is *not* fixed and should not be, because they are unobservable. The uniqueness claim is therefore substantive in the dimensions that carry physical content (the discrete structural invariants) and silent in the dimensions that do not (the gauge directions): a tautological theorem would fix nothing specific, whereas this one fixes a particular finite list of integers and groups that the framework then matches against the observed Standard Model.

Refinement (which residue is forced, and which is selected). The defense above is correct but must be stated with one further distinction, or it over-claims. Of the discrete invariants listed, those that are *forced by representation theory given the lattice* — the multiplicities $(3, 2, 1)$ and the gauge factors, the generation count, the hypercharges (apart from the exceptional branch $Y_Q = 0$) — are non-tautological in the full sense: a priori they could have come out otherwise, and the octahedral character table fixes them. But two of the listed invariants are not forced by the construction in this way: the spatial dimension $d = 3$ is selected by the empirical filters E5–E7 (§3.3), and the simple-cubic Bravais structure is selected, within the nearest-neighbor class, by the observed $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ (Theorem 7b, [SM §4.6]) — only the cubic crystal system supplies the three-dimensional point-group irrep an SU(3) factor requires. For these, the inference runs from observed physics back to the commitment, so the theorem's "uniqueness" at these joints is uniqueness *relative to the observed inputs*, not derivation of them. This does not collapse the residue into tautology — the representation-theoretic consequences remain genuine content — but it locates the boundary precisely: the framework forces the consequences of the cubic lattice, and selects the cubic lattice from the world. The full three-way classification (forced / empirically selected / definitionally stipulated) is given in [SM §8.3, *Derivation status: the three tiers*]; the honest one-line statement of the framework's reach is that it is a maximally tight compression of observed physics onto the cubic-lattice commitment, not a derivation of physics from the bare fact of observation.

The structural assumptions A1–A6 exclude several broad classes of theory from the outset:

- Intrinsically continuum theories (violate A1)
- Intrinsically stochastic theories (violate A2)
- Theories with unbounded coupling degree (violate A3)
- Theories violating center independence, linearity, or background independence (violate A4–A6)

The framework's claim is that within the stated structural class, with the nearest-neighbor premise of Stage 2(b), the **local reconstructed residue** is unique up to gauge. The Bell-inclusive substratum is not proved unique. A separate argument would be required to rule out alternatives outside this class. The framework's defense of A1–A6 (in [SM §2] and elsewhere) is that they are natural given the empirical inputs E1–E7, but this defense is itself an argument, not a theorem.

Remark (Hypothesis dependencies). Each structural assumption can in principle be weakened or replaced:

- A1 (finiteness) is supported by E3; alternative would be an effective-finiteness argument with continuum UV completion, which the framework does not develop.
- A2 (determinism) is the key ontological commitment; stochastic alternatives would produce different reconstructions.
- A3 (bounded coupling) is conventional for physical systems; long-range couplings would give different dimensional structure.
- A4 (center independence) rules out preferred-frame theories.
- A5 (linearity) is the strongest restriction; amplitude-scale gauge invariance restricts the rule to an affine one (linearity-equivalence lemma, [SM §4.1]), so non-affine wave equations on finite lattices are the separate class one obtains exactly when that gauge principle is dropped.
- A6 (background independence), as covariance under local internal-index transformations with the link coupling transformed alongside, is the standard gauge-theoretic requirement; the stronger fixed-background invariance, with the coupling held fixed, is a separate condition and is not what A6 asserts.

Weakening any A_i does not merely enlarge the equivalence class; it potentially produces *different* reconstructions that may not agree with observation. The framework's claim is that E1–E7 + A1–A6 is a consistent set of inputs producing our observed physics; alternative input sets are not investigated.

Remark (Continuum extension). The lattice-level predictions are the framework's primary claims — the lattice is the fundamental description, not an approximation to a continuum theory. The structural results (gauge group, representations, generation count) are algebraic/topological properties of the lattice theory and are exact. Quantitative observables (scattering amplitudes, mass ratios) are compared to experiment through continuum perturbation the-

ory; the lattice-continuum discrepancy at any experimentally accessible energy E is suppressed by $(E\epsilon/\hbar c)^2 = 4\sigma(E/M_{\mathrm{Pl}})^2$, which at $\sigma = 1$ is $\sim 10^{-32}$ at LHC energies. The rigorous proof that lattice Yang-Mills defines a continuum limit with a mass gap (a Clay Millennium Prize problem) is an open problem in mathematics, but it is not required by the OI framework: the lattice theory is complete, and its predictions are lattice-exact.

The epistemic structure is as follows. A reasonable objection is that the framework treats the lattice as fundamental (so the mass-gap problem can be set aside) while also using continuum perturbation theory to extract the numbers that match data (which would make the continuum limit load-bearing). The position is consistent. The framework's claim is that the lattice theory is the complete physical content and its predictions are exact *at the lattice level*. The comparison to experiment is then a comparison between two objects: the lattice theory's prediction extracted via continuum PT (a controlled approximation with error bounded by $(E\epsilon/\hbar c)^2 = 4\sigma(E/M_{\mathrm{Pl}})^2 \sim 10^{-32}$ at $\sigma = 1$), and the continuum-QFT fit to the experimental data (the standard way the data is reported). The empirical agreement is therefore agreement between the lattice theory's continuum approximation and the continuum-QFT data fit, with the approximation error in the first object ~ 30 orders of magnitude below the experimental precision of the second, so the lattice-vs-continuum gap is not the source of any observed agreement or disagreement at accessible energies. The mass-gap problem is not load-bearing for this: it concerns whether a continuum limit *exists as a rigorous mathematical object*, whereas the framework requires only that continuum PT is an accurate *calculational approximation* to the lattice theory's predictions at low energy — the standard status of lattice-to-continuum perturbative matching, which does not require the existence of the rigorous limit. The lattice is fundamental, the continuum is a calculational tool with a quantified error, and the experimental comparison is between the tool's output and continuum-reported data.

Remark (Finite observation sets vs. idealized empirical inputs). The empirical inputs E1–E7 are stated as structural facts about observed physics — the full apparatus of QM, Bell violations attaining Tsirelson's bound, the holographic entropy bound, exact rotational invariance, propagating gravitational waves, stable matter on all observed scales, and spatial-curvature concordance. Operationally, no finite observation set fully establishes any of these; finite measurements support them only in the sense of being consistent with the data so far. The reconstruction theorem holds for the empirical inputs *as stated* — the idealized infinite-data regime — and proves uniqueness of the local lattice/gauge residue under that regime; Bell-inclusive uniqueness remains open. Reconstruction from finite observation sets is a related but distinct question: with finite data, multiple equivalence classes may all be consistent, with the data discriminating among them only weakly. The framework's operational answer to the finite-data question is the cumulative weight of the quantitative predictions of [SM §7] — twenty-two classified retrodictions of gauge couplings, CKM/PMNS angles, mass ratios, and the Koide relation (two parameter-free structural and

five conditional structural, the remainder layered, chained or fitted, the gauge couplings among the fitted), the measured comparisons each at $\lesssim 1\sigma$ — which collectively rule out reconstructions that would have produced different values. Theorem 23 establishes idealized local-sector uniqueness; the [SM §7] prediction set supplies the operational evidence for that local structural residue that a skeptical reader can check against data.

The reconstruction establishes a bidirectional correspondence for the **local reconstructed residue**:

$$\text{Observed local lattice/gauge data (E1–E7) + Structural assumptions (A1–A6)} \quad \longleftrightarrow \quad [(S,\varphi)]_{\text{loc}}/\mathcal{G}_{\text{sub}}.$$

The reconstruction theorem concerns the local lattice/gauge residue under its stated inputs and structural premises. Bell-inclusive existence remains conditional on H-Bell and Bell-inclusive uniqueness is open. Relative to a specified observation interface, two descriptions that induce identical data cannot be distinguished by a decision based only on those data. This leaves the broader metaphysical question of whether mathematics describes reality or constitutes it outside the reconstruction result; Theorem 24 supplies fixed-interface invariances, not an all-measurements undecidability theorem about that question.

Remark (ER=EPR at substratum and emergent levels). The Maldacena-Susskind ER=EPR conjecture [3] proposes that any two entangled systems are connected by an Einstein-Rosen bridge, with the strong form asserting that entanglement *is* a non-traversable wormhole — the same ontological structure under two descriptions. The framework here engages this conjecture at two distinct levels.

At the **emergent level**, the conjecture's *weak* form is reproduced: any pair of entangled subsystems in the emergent QM description shares boundary modes via the partition trace-out, and the holographic dictionary maps shared boundary modes to a connecting geometric structure in the emergent gravitational description. This recovers the ER=EPR weak form — entangled systems are connected by a wormhole-like geometric object in the emergent theory — as a structural consequence of the framework's holographic architecture rather than as an independent postulate.

At the **substratum level**, the conjecture's *strong* form (entanglement is identical to wormhole geometry, not merely correlated with it) is predicted *false* at the representative level. Two distinct partition representatives can produce identical emergent EPR correlations while having different substratum representative structures — there is no faithful bijection from EPR pairs to substratum geometric features. The strong identity holds at most up to gauge equivalence on the substratum side. Whether a gauge-orbit reformulation of strong ER=EPR — "EPR-equivalence-classes correspond to ER-equivalence-classes" — is true in the framework remains open; the test would be whether the substratum's

gauge group $\mathcal{G}_{\mathrm{sub}}$ acts transitively on the set of substratum representatives that produce a given EPR correlation pattern.

This produces a sharp empirical signature: the framework's operational predictions agree with weak ER=EPR (which would also be predicted by AdS/CFT, by quantum error correction in holography, and by other emergent-geometry approaches) but disagree with strong ER=EPR at the representative level. Distinguishing these experimentally would require probing substratum-scale features, which is currently beyond reach. The disagreement is therefore a structural prediction rather than a testable one at present, but it concretely positions the framework relative to the broader ER=EPR program.

---

4. The Substratum Gauge Group

The observational datum must specify the visible/hidden interface, the admissible preparations and the readouts. At that fixed interface, equality of the coherent reduced-channel family is stronger than equality of the rooted population transition probabilities $T_{ij}(t)$. Write $\sim$ for equality of the specified coherent-channel data at that fixed interface; population-only comparisons are stated separately. The representative freedoms denoted by $\mathcal{G}_{\mathrm{sub}}$ below are compared on their stated data: (i) and the finite normalized-product form of (iii) preserve the coherent family, while (ii) retains the alphabet-size hypothesis. These results do not classify equivalence over all observer partitions or reconstruct every physical implementation.

Theorem 24 (Generators of the substratum gauge group). *$\mathcal{G}_{sub}$ contains the independent families of transformations (i) and (iii) below, and, under the alphabet-size hypothesis of [SM, §2.7], the family (ii); (iv) is a structural graph comparison, not a family of $\mathcal{G}_{sub}$:*

(i) Hidden-sector relabeling. For any bijection $\sigma : \mathcal{C}_H \to \mathcal{C}_H$, applied identically in every visible fibre, $\pi(v,h) = (v, \sigma(h))$, the conjugate $(S, \pi \circ \varphi \circ \pi^{-1}) \sim (S, \varphi)$: with uniform hidden prior $\omega_H = \mathbb{1}/|\mathcal{C}_H|$, every visible channel $\Phi_t(\rho_V) = \mathrm{Tr}_H[\widehat{\varphi}^{\,t}\,(\rho_V \otimes \omega_H)\,\widehat{\varphi}^{-t}]$, with $\widehat{\varphi}$ the permutation unitary of φ, is unchanged. A fibre-dependent hidden relabelling $(v,h) \mapsto (v, \sigma_v(h))$ preserves the transition probabilities $T_{ij}(t)$ but not, in general, the coherent channels Φ_t.

(ii) Alphabet change (under the alphabet-size hypothesis). Replacing the local state space $\mathbb{Z}/q\mathbb{Z}$ with $\mathbb{Z}/q'\mathbb{Z}$ for any $q' \geq 2$, while preserving the coupling graph and dynamics class, leaves all observables unchanged if the alphabet-size hypothesis of [SM, §2.7] holds; [SM, §2.7] lists the results proved for every $q \geq 2$ and those that carry q. This family is parametrized by all integers $q \geq 2$.

(iii) Finite decoupled ancillary enlargement. Fix the visible/hidden interface $V \times H$, the normalized hidden density matrix ω_H, and the permutation unitary U induced by φ. Adjoin a finite factor A, prepared independently in a density matrix σ_A with $\mathrm{Tr}\,\sigma_A = 1$, and evolve it by a permutation unitary W with

product dynamics $U \otimes W$. Then, for every visible input ρ_V and discrete time t, the enlarged reduced channel equals the original one:

$$\begin{aligned}\Phi'_t(\rho_V) &= \mathrm{Tr}_{HA}\big[(U^t \otimes W^t)(\rho_V \otimes \omega_H \otimes \sigma_A)(U^{\dagger t} \otimes W^{\dagger t})\big] \\ &= \mathrm{Tr}_H\big[U^t(\rho_V \otimes \omega_H)U^{\dagger t}\big] = \Phi_t(\rho_V).\end{aligned}$$

Consequently every rooted population law $T_{ij}(t)$ is unchanged. Removing the independently prepared decoupled factor reverses this comparison. This is a finite, fixed-interface result; an infinite extension requires its own normalized measure or trace-class state and a well-defined evolution, and is not established by finite uniform counting. An enlargement coupled to the boundary layer, even slowly, falls outside this identity: [GR §3.2] supplies only $T_{ij}(t) = T_{ij}^{(B)}(t) + \mathcal{O}(t/\tau_B)$ on its stated accessible-time regime.

(iv) Graph isomorphism (up to statistical isotropy), with a qualification on the gauge sector. Two coupling graphs G_φ and $G_{\varphi'}$ that are quasi-isometric with the same polynomial growth exponent d, the same spectral mode count, and the same statistical isotropy at large scales both admit the graph-independent finite-law representation [Main §3.4] and share the growth exponent d that enters the dimensional filters [SM §3.2], but need not give the same transition probabilities $T_{ij}(t)$ at a fixed observer interface, and the continuum-curvature step of the gravitational construction [SM §3.1] requires its own geometric conditions; in this sense the regular cubic lattice $\mathbb{Z}^3$ and a bounded-degree random graph with $d = 3$ polynomial growth and statistical isotropy are structurally comparable, not observationally equivalent. The *gauge-group* content of Stage 2(d) is where this crystallography becomes physical: the multiplicities $(3, 2, 1)$ are fixed by the octahedral point group acting on the six link directions (Theorem 7; [SM §4.5–4.6]), and the local point group is *not* a quasi-isometry invariant — only the spectral mode count $2d_s = 6$ transfers to a generic isotropic graph, not the cubic decomposition. For the gauge-group content, therefore, the equivalence in (iv) is the narrower one of rigid orientation together with isomorphisms preserving the cubic (octahedral) local structure; the simple-cubic Bravais type itself is an empirically-selected input (§3, Refinement below; [SM §8.3]), not a gauge choice. A generic isotropic random graph admits the finite-law representation but need not reproduce the same observable law, and does not reproduce the Standard Model gauge group. The gauge group itself is a universality-class invariant, shared with any Standard-Model-reproducing class through different machinery ([Structure §2.3, §14.4]); within OI's class it is realized through the cubic-group commutant ([SM §4.6]).

Proof. Each family preserves the specified data under its stated hypotheses. (i): conjugation by $\mathbb{1}_V \otimes P_\sigma$ leaves ω_H invariant and cancels under the partial trace, so every Φ_t is unchanged; for a fibre-dependent σ_v the visible coordinate is fixed and each rooted fibre's uniform hidden prior is permuted bijectively, so every $T_{ij}(t)$ is unchanged. (ii): this is the alphabet-size hypothesis of [SM, §2.7], under which the family is stated; the results [SM, §2.7] proves for every $q \geq 2$ are preserved without it. (iii): the evolved state factors as

$U^t(\rho_V \otimes \omega_H)U^{\dagger t} \otimes W^t\sigma_A W^{\dagger t}$. Partial trace over HA is the original reduced state multiplied by $\mathrm{Tr}(W^t\sigma_A W^{\dagger t}) = \mathrm{Tr}\,\sigma_A = 1$. This proves the coherent-channel identity and hence its population restriction. It preserves the fixed observed data, not measurements that access A or an extension that later couples to $V \times H$. (iv): the finite-law representation [Main §3.4] holds for any finite bijection and the growth exponent is shared by hypothesis, while $T_{ij}(t)$ depends on the specific graph and the continuum-curvature step [SM §3.1] has its own conditions; the gauge-group derivation of Stage 2(d) additionally uses the octahedral point-group action on the six link directions, which the simple-cubic representative supplies and a generic isotropic graph does not, so for that content (iv) is restricted to the orientation- and octahedral-structure-preserving isomorphisms (see the qualification in the statement). □

Corollary (effective finiteness and gauge-class transfer). *(i) The bound.* E3, read as a Hilbert-space dimension cutoff, bounds the observationally relevant space by $|\mathcal{C}_V \times \mathcal{C}_B| \leq e^{S_\partial}$, with $S_\partial = A/4$ in Planck units. This reading is the interpretive premise of A1. *(ii) Accessible-time comparison.* Under [GR §3.2]'s hypotheses, $T_{ij}(t) = T_{ij}^{(B)}(t) + \mathcal{O}(t/\tau_B)$ for $t \ll \tau_B$. The leading-order description therefore uses only visible and boundary data. *(iii) Exact transfer.* For the independently prepared finite decoupled extension of Theorem 24(iii), the full fixed-interface channel family, and hence every statement expressible solely through that family or its $T_{ij}(t)$ restriction, transfers exactly at every time. This includes a recurrence-scale property of the observed law when that property is established on a finite representative; it does not imply recurrence of every unobserved sector. Infinite extensions need a separately specified normalized state or measure and dynamics before any such comparison can be made. Across slowly coupled extensions, only statements insensitive to the $\mathcal{O}(t/\tau_B)$ correction transfer. For example, $T_{ij} = 0$ does not transfer to $T_{ij} = \varepsilon > 0$ merely because $\varepsilon = \mathcal{O}(t/\tau_B)$. Finite-carrier recurrence, and the accessible-window conditional-memory conclusion of [Main §2.3], remain distinct claims. □

Remark (amplitude-scale gauge — the unobservability of absolute field scale). One gauge freedom invoked elsewhere is named here as its single authoritative home, and is best stated *operationally*: an embedded observer has no access to the absolute scale of the substratum field value — only to the emergent transition probabilities and coupling structure ([SM §2.7]). This is the discrete analogue of the field-normalization redundancy of ordinary field theory, where rescaling a field changes nothing physical because only correlators, not absolute amplitudes, are observable. It is realized as *amplitude-scale invariance*: the rescalings $\phi \mapsto \lambda\phi$ (λ a unit) and shifts $\phi \mapsto \phi + c$ — the additive-automorphism (affine) relabellings of the alphabet — leave every observable unchanged. These affine relabellings are exactly the site-wise relabellings that preserve not only the coupling graph but the field-amplitude dispersion $\omega(k)$ (the emergent Hamiltonian), so amplitude-scale gauge is one instance of the framework's master principle that *observationally indistinguishable transformations are gauge* — the same principle behind generators (i)–(iii) and the q-size

freedom of [SM §2.7], not an ad hoc addition. Two derivations invoke it: the linearity equivalence of [SM §4.1] (amplitude-independent dispersion $\iff$ affine dynamics) and the $SU(N)$ reduction of Stage 2(d) above; its status as an operational input rather than a theorem — and the sense in which those two results are conditional on it — is set out in the *Status* note immediately below.

Status (open, referee-grade). Amplitude-scale gauge is adjoined as an explicit operational principle: it is not one of (i)–(iii) — generator (i) relabels hidden states only and (ii) is alphabet-*size* only — so the results invoking it are conditional on it. Stated operationally it is *not* viciously circular: the input (absolute amplitude is unobservable) is strictly weaker than the output (affine dynamics), and a non-affine F on the same alphabet is a coherent object the principle *excludes* rather than presupposes. The one route that would make the affine restriction *unconditional* — deriving the unobservability of absolute amplitude from the observer architecture itself — is by contrast circular: what the embedded observer can resolve through the trace-out is dynamics-dependent (a non-affine F induces amplitude-dependent emergent dispersion, which *is* observable), so "absolute amplitude is unobservable" already presupposes an affine rule. Amplitude-scale gauge therefore has the status of an operational input on a par with center independence and isotropy, not a theorem; whether it is independently warranted by the observer architecture is the question flagged for external review.

Formalization status of the linearity step (resolved as a sharpened axiom, not a closure). A dedicated audit of this step settles its logical status, and the outcome is worth stating precisely because it is the prototype for how the framework's "open" foundational steps resolve. The linearity-equivalence lemma — *all first finite-differences of the update rule over $\mathbb{Z}/q\mathbb{Z}$ are base-point-independent iff the rule is affine as a function* — has been verified exhaustively for $q \leq 6$ in one neighbour-coordinate and $q \leq 3$ in two, and on 2.5×10^5 random rules at larger (q, m), with zero counterexamples; together with the telescoping proof (constant single-coordinate differences force $F(x) = F(0) + \sum_i c_i x_i$, well-defined on the torus since $qc_i \equiv 0$) this is a complete elementary result, not a genericity heuristic. The question of whether the *q-size* gauge freedom of [SM §2.7] *entails* amplitude-scale gauge resolves in the negative, and sharply: the weak reading of [SM §2.7] (size-independence of predictions) does not reach rescaling at fixed q; the strong reading (arbitrary relabellings of $\mathbb{Z}/q\mathbb{Z}$ are gauge) *overshoots* — a nonlinear permutation conjugating a linear rule preserves cycle structure (hence emergent transition probabilities up to relabelling) while being nonlinear, so it would render a nonlinear rule gauge-equivalent to a linear one and thereby *destroy* linearity-necessity rather than establish it (verified explicitly on $\mathbb{Z}/5\mathbb{Z}$). The principled object that delivers the affine restriction without overshooting is the **additive-automorphism** group $x \mapsto \lambda x + c$ (λ a unit) — a proper subgroup of all permutations that preserves the alphabet's "+", hence the dispersion. The upshot, in the framework's three-tier classification ([SM §8.3]): linearity is **not** promoted to a forced (Tier-1) consequence; it is a Tier-3 *definitional stipulation*, but a sharpened one — the more primitive and better-motivated assumption

"the alphabet's additive automorphisms are gauge" yields an affine rule as a certified theorem, and linearity is the further restriction the stipulation adds. This is the characteristic resolution shape for the framework's foundational gaps: pressing for a proof does not close the gap to a derivation but relocates the irreducible assumption, apart from the further restriction from affine to linear, to a more primitive and more defensible place, and names it exactly.

Completeness of the generators (open). Whether families (i)–(iii) generate $\mathcal{G}_{\mathrm{sub}}$, family (ii) entering under the alphabet-size hypothesis, is open. The question has two logical parts that are worth separating, because the completeness (as opposed to soundness) direction is the substantive one: soundness — that each of (i)–(iii) preserves observables — is established above; completeness — that *every* observables-preserving transformation lies in the group generated by (i)–(iii) — is not established. Equality of all time-resolved visible channels does not determine the bijection up to hidden unitary, even with trivial hidden commutants. In the Lean-checked ST3_pairC counterexample, two permutation unitaries on six states ($|V| = 2$, $|H| = 3$) induce the same visible channel Φ_t at every time, with nontrivial visible statistics and trivial hidden commutants. Their traces are 2 and 0, so no unitary conjugates them.

Specht's trace criterion and Wiegmann's simultaneous-unitary extension characterize unitary similarity through traces of words in matrices and their adjoints [5–7]. Proposition 24.2 is a fixed-carrier specialization with the additional spanning premise used by its cited Lean witnesses. The classical finite-dimensional criterion itself does not require that premise.

Proposition 24.2 (Fixed-carrier block-word rigidity). *Let V, H be finite index sets and U, U' be matrices on $\mathbb{C}^V \otimes \mathbb{C}^H$. Write $B_{vw}(U)$ for their visible blocks on $\mathbb{C}^H$, and let $\mathcal{W}(U)$ be the linear span of all words in the blocks and their adjoints, including the identity. Suppose $\mathcal{W}(U) = M_{|H|}(\mathbb{C})$ and the unnormalized trace of every corresponding block-word agrees for U and U'. Then a hidden unitary W satisfies $U' = (\mathbb{1}_V \otimes W)U(\mathbb{1}_V \otimes W)^\dagger$. Conversely, hidden-unitary conjugation preserves every block-word trace, and the hypotheses imply $\mathcal{W}(U') = M_{|H|}(\mathbb{C})$.* No unitarity assumption on U, U' is needed, but the two systems must have the same hidden index set.

Proof. Equality of all word traces yields a trace-preserving unital $*$-isomorphism of the generated word spans. When the first span is the full matrix algebra, the matrix-unit construction implements it by a unitary. Necessity follows by trace invariance. The Lean witnesses for the two directions are WT2gen_oneSided and trace_wordEval_hiddenConjugate, combined in WT3_spanning_iff; blockSpanning_of_wordTracesEqual establishes spanning for the target (OIBridge/WordTraceSufficiency.lean and OIBridge/SemigroupTransfer.lean). □

The ST3_pairC pair has unequal block-word traces (pairC_wordTraces), so it does not meet Proposition 24.2's premises.

A further candidate family is explicitly subsumed:

- *Visible-sector emergent global phase* ($U \to e^{i\theta}U$): at the substratum level S is a finite set with no complex structure; the emergent phase is trivially the identity on (S, φ).

The gauge hierarchy. Three levels of gauge symmetry appear in the framework:

Level 3 (substratum): $\mathcal{G}_{\text{sub}}$ acts on (S, φ) before the trace-out. It includes transformations with no analog in the emergent description (deep-sector enlargement, and alphabet change under the alphabet-size hypothesis).

Level 2 (emergent QFT): $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ is the gauge group acting on the emergent fields (Stage 2(d): the block-algebra unitary group $\mathrm{U}(3) \times \mathrm{U}(2) \times \mathrm{U}(1)$ with a specified charged matter action).

Level 1 (emergent Hamiltonian): The D-gauge $H \to DHD^\dagger$ with D a diagonal unitary, acting on the emergent Hamiltonian within the emergent QM. It preserves every transition probability exactly. Energy shifts and the antiunitary reflection are also invisible; completeness of these alternatives under the generic gap and overlap hypotheses is conditional on H-BG ([GR §3.3]).

The levels carry distinct equivalence notions; inclusions between them and a map from $\mathcal{G}_{\text{sub}}$ onto the Standard Model gauge group are not established.

Remark. The substratum gauge group is not a symmetry of a Lagrangian or an action — no Lagrangian exists at the substratum level. It is a symmetry of the *equivalence class of substrata*, defined by the condition that all observables are preserved. The emergent gauge symmetries (Levels 1 and 2) are Lagrangian symmetries in the standard sense, derived from the substratum through the trace-out.

4.1 Conditional reconstruction from complete word data

Multi-time records can recover more than separate one-time channels, but their observational carrier must be specified. The following classical minimal-realization result for weighted automata and rational series originates in Schützenberger, Carlyle–Paz and Fliess [12–14]; minimal-realization uniqueness up to similarity is stated in [10, Chapter 2, Theorem 2.4], with the common sequential-system and Hankel framework developed in [11]. This result and its native deterministic specialization recover predictive structure. They complement Proposition 24.2, whose input is coherent block-word traces, rather than imply its hypotheses.

Proposition 24.3 (Minimal linear word realizations). *Let a finite alphabet $\mathcal{A}$ have two finite-dimensional realizations $(E, a, b, (M_c))$ and $(E', a', b', (M'_c))$ over a field, with chronological convention $M_{c_1 \cdots c_k} = M_{c_k} \cdots M_{c_1}$ and $M_\epsilon = I$. Assume each is reachable,* $\mathrm{span}\{M_w a\} = E$, *and observable,* $\bigcap_u \ker(bM_u) = \{0\}$, *and similarly for the primed model. Equality $bM_w a = b' M'_w a'$ for every word implies a unique linear isomorphism S with $Sa = a'$, $SM_c = M'_c S$, and $b'S = b$.*

For arbitrary competing dimensions $n, m \geq 1$, equality through word length $n + m - 1$ already implies equality for all words. A nonzero series of known minimal dimension d has a scalar Hankel reconstruction using prefix and suffix lengths at most $d - 1$ and word statistics through $2d - 1$.

Proof. Define

$$S\left(\sum_i t_i M_{w_i} a\right) = \sum_i t_i M'_{w_i} a'.$$

If the source sum is zero, every target continuation evaluates the proposed image to $\sum_i t_i b' M'_u M'_{w_i} a' = \sum_i t_i b M_u M_{w_i} a = 0$; target observability makes the image zero. Thus the map is well-defined. Reversing the construction gives its inverse by reachability. The empty word gives $Sa = a'$, appending a letter gives $SM_c = M'_c S$, and equality of evaluations gives $b'S = b$. Any such map agrees on the reachable generators, proving uniqueness.

For the length bound, the reachable spaces $R_k = \mathrm{span}\{M_w a : |w| \leq k\}$ satisfy $R_{k+1} = R_k + \sum_c M_c R_k$. Equality at two consecutive stages makes the space invariant and the sequence stable; otherwise its dimension increases. In dimension N it therefore stabilizes by $N - 1$ for nonzero a, and the zero-vector case is immediate. Apply this to the $(n + m)$-dimensional discrepancy realization with operators $M_c \oplus M'_c$, vector (a, a'), and evaluation $(b, -b')$.

In a minimal d-dimensional model choose reachable-basis prefixes w_j and observable-basis suffixes u_i, each of length at most $d - 1$ by the same filtration argument. Put $R = [M_{w_j} a]_j$ and $O = [b M_{u_i}]_i$. Then

$$H_{ij} = s(w_j u_i) = (OR)_{ij}, \qquad (H_c)_{ij} = s(w_j c u_i) = (OM_c R)_{ij}.$$

Both O and R are invertible. The reconstructed coordinates are

$$\widehat{M}_c = H^{-1} H_c, \qquad \hat{a} = H^{-1}(s(u_i))_i, \qquad \hat{b} = (s(w_j))_j.$$

These equal $R^{-1} M_c R$, $R^{-1} a$, and bR. The largest required length is $(d - 1) + 1 + (d - 1) = 2d - 1$, and the invertible minor proves that full Hankel rank is d. □

The alphabet must be stationary and freely composable, with every required prefix, suffix and extension accessible; an outcome-labelled instrument uses an unnormalized linear update. A restricted protocol grammar or changing apparatus requires its own state augmentation or admissible-language theorem. The dimension/rank bound is an independent hypothesis: a finite observed Hankel rank alone does not certify it. Similarity is only a linear conclusion. Even strictly positive normalized minimal stochastic models can have a unique non-monomial similarity with an inverse having negative entries. An isomorphism mapping the entire normalized classical simplex onto itself would instead permute its extreme points, but generic linear reachability and observability do not establish that stronger condition.

Proposition 24.4 (Passive finite-permutation predictive quotient). *Let P be a permutation of a nonempty finite set Ω, with deterministic passive readout $r : \Omega \to \mathcal{A}$ and initial law μ. Identify ω, η exactly when $r(P^t\omega) = r(P^t\eta)$ for every $t \geq 0$, and let Q be the resulting future-trace quotient. It inherits a permutation $\bar{P}$, readout $\bar{r}$, and prior $\bar{\mu}$. For each model being compared, assume its quotient prior $\bar{\mu}$ has full support, and let B be a known upper bound on both quotient-state counts. For $B \geq 2$, equality of all passive word probabilities through length B, counting the time-zero readout, determines $(Q, \bar{P}, \bar{r}, \bar{\mu})$ up to a unique readout-preserving conjugacy transporting the prior. For $B = 1$, one-letter data suffice. No invariance or uniformity of μ is required. The quotient's linear Hankel rank equals $|Q|$; common-length retained-word vectors and block effects generate its full nonnegative state cone and all coordinate indicators, respectively. Minimal-realization similarity between two such quotients is a permutation.*

Proof. Future-trace equivalence is forward invariant. Since P has a finite order, every negative power is also a nonnegative power, so the equivalence is backward invariant too and $\bar{P}$ is a permutation. On an N-state deterministic labelled system let C_k be the partition by its first k readouts, with C_0 indiscrete. If $C_k = C_{k+1}$, equality of labels and successor classes makes this equivalence forward invariant and no later refinement occurs. Otherwise the number of classes increases. Thus C_{N-1} is the full future partition for $N \geq 2$.

Put $\ell = B - 1 \geq 1$. Distinct quotient states are distinguished by their length-ℓ future blocks. Full support makes the common positive-probability length-ℓ words exactly the state labels W. Each $w \in W$ has exactly one symbol x with $s(wx) > 0$; its successor label is the last ℓ symbols of wx. The first symbol of w gives its readout and $s(w)$ gives its initial mass. Hence length-B data determine the labelled quotient permutation and prior. Any readout-preserving intertwiner preserves the entire future trace, so this conjugacy is unique. For $B = 1$ the sole state has identity dynamics and its one-letter law determines its label.

For the cone and rank statements choose one common separating length $\ell \geq \max(1, |Q|-1)$. In quotient coordinates set $M_x = \bar{P}D_x$, where D_x filters readout x, $a = \bar{\mu}$, and $b = \mathbf{1}^T$. These reproduce passive cylinder probabilities because the last advance cancels under $\mathbf{1}^T\bar{P} = \mathbf{1}^T$. If w_q is the length-ℓ block of state q, then

$$bM_{w_q} = e_q^T, \qquad M_{w_q}\bar{\mu} = \bar{\mu}(q)e_{\bar{P}^\ell q}.$$

Full support gives a positive coefficient, and the common power $\bar{P}^\ell$ bijectively covers all end states. Thus the retained-word rays and block effects cover the coordinate rays and indicators. The model is reachable and observable, so its Hankel rank is $|Q|$. Matching blocks and their probabilities makes the similarity of Proposition 24.3 carry the corresponding coordinate rays with unit scale, hence be a permutation. □

Access and remaining gauge gap. This is a predictive-quotient result for passive, nondisturbing records from one fixed finite reversible carrier. It refines

[Main §3.4]'s finite-horizon canonical tail quotient; it does not infer an all-time carrier from arbitrary finite-horizon laws. Full support on the microscopic carrier is sufficient, but one rooted visible preparation $\delta_v \otimes \mu_H$ is generally not such a prior. A consistent positive mixture of every root would supply it only if that preparation family is actually accessible with the same dynamics and hidden prior. Algebraic saturation on Q does not assert externally selectable microscopic states or separate states in the same future-trace fibre. Without a known state bound, full-support uniform laws on binary de Bruijn cycles of orders k and $k+1$ agree through length k and differ later. Invisible fibres can also have different microscopic cycle structures. Neither proposition therefore supplies hidden-factor conjugacy, coherent block-word traces, a common quantum instrument theory, or completeness over all admissible partitions.

The self-contained derivation and bounded exact rational controls are retained in verification/audits/manuscript/multitime-predictive-recovery-20261028/. The native checks cover 768 four-state cases and two one-state cases, including nonstationary full-support priors, constant readouts, common-length cone/Hankel checks, and countercontrols for missing hypotheses. These are executable evidence supplementing the written proofs; neither new proposition is claimed Lean-proved.

Proposition 24.5 (Passive record fibres and state identification). *Let P be a known permutation of a nonempty finite carrier Ω, let $r : \Omega \to \mathcal{A}$ be a deterministic readout, and let μ be a known initial law with support A. Suppose the observer may retain the exact passive readings $X_j = r(P^j S_0)$. For a positive-probability word $w = (w_0, \dots, w_{k-1})$ of length $k \geq 1$, define*

$$F_w = \{s \in A : r(P^j s) = w_j \text{ for } 0 \leq j < k\}.$$

(i) A quantity $q(S_0)$ is identified by that record exactly when q is constant on F_w. In particular the initial state is identified exactly when $|F_w| = 1$. The current state $P^{k-1} S_0$ has the same singleton criterion. When $\Omega = \mathcal{C}_V \times \mathcal{C}_H$ and $r = \pi_V$, the current hidden state is identified exactly under that same criterion. (ii) Put $N = |\Omega|$ and $L = \max(1, N-1)$. Passive words of length L already induce the full future-trace equivalence on Ω, hence on A by restriction. Every positive-probability finite record remains compatible with at least two initial states if and only if every nonempty intersection of a full future-trace class with A has at least two elements.

Proof. Conditioning gives

$$\Pr(S_0 = s \mid w) = \frac{\mu(s) \mathbf{1}_{F_w}(s)}{\sum_{u \in F_w} \mu(u)}.$$

Every point of F_w has positive mass, so a quantity is known exactly when it is constant there. Invertibility maps F_w bijectively to the current compatible-state set. For a product carrier with current visible value recorded, distinct current compatible states have distinct hidden coordinates. This proves (i).

For (ii), partition Ω by its first j readings, starting with the indiscrete partition at $j = 0$. Each next reading refines it. If two consecutive partitions agree, the classes have identical labels and successors in the same class; no later refinement occurs. Otherwise the class count strictly increases. Starting with one class, there can be at most $N - 1$ strict increases, so length $N - 1$ suffices for $N \geq 2$; one reading suffices when $N = 1$. Restriction to A gives the assertion without assuming that A is invariant. Each nonempty finite-record fibre is a union of restricted full future-trace classes. Thus classes of size at least two make every such fibre non-singleton; conversely a singleton full class yields a singleton length-L fibre. □

Memory and opacity are different. Under the uniform law in the swap model, $H_1 = X_0$, $X_2 = X_0$ and $I(X_0; X_2 \mid X_1) = 1$, but $H(H_1 \mid X_0, X_1) = 0$. More generally, cyclically shifting $m + 1$ independent uniform bits and reading the first gives a hidden store of 2^m states, all identified by $m+1$ retained readings. At time m, $I(X_{<m}; X_{>m} \mid X_m) = m$ bits, saturating the hidden capacity, while $H(H_m \mid X_{\leq m}) = 0$. Indeed, conditional on the current bit, the preceding m readings specify all other bits, and the next m readings repeat them. C4's information is predictive memory relative to the present readout, not a lower bound on residual uncertainty after the full history.

Access and resource scope. The proposition assumes the dynamics, preparation support and retained records are available. It supplies no physical record-storage or intervention mechanism. A stored record with only R possible values cannot distinguish all $|A| > R$ admissible initial states: some fibre has at least $\lceil |A|/R \rceil$ members, although other fibres may be singletons. This pigeonhole limit is weaker than claiming ambiguity for every history. A decoupled spectator gives genuine non-singleton passive classes; fixing that spectator by preparation, reading it, or allowing a separating intervention can remove the ambiguity. Proposition 24.4 instead reconstructs an unknown predictive quotient from exact word laws under its own support and state-bound hypotheses. Neither result identifies the physical OI model, coherent instruments or the full gauge relation. The bounded exact controls are retained in verification/audits/manuscript/hidden-state-identifiability-20261028/.

---

5. Synthesis: Three Projections of (S, φ)

The reconstruction theorem (§3) establishes that the stated empirical and structural inputs, together with M1-T, C2-mixing and the nearest-neighbor premise of Stage 2(b), fix the **local propagating lattice/gauge residue**. M1-B and H-Bell separately condition the existence of a measurement-independent Bell-violating completion; Bell-inclusive uniqueness is open. The substratum gauge group (§4) makes the equivalence relation precise. This section makes the synthesis claim explicit: quantum mechanics and general relativity are not two independent theories but projections of the same finite deterministic construc-

tion, and a third projection bears on the arrow of time (§5.4).

5.1 The three projections

The framework's derivations apply the same trace-out machinery to the same (S, φ) along three projections: a visible-sector projection that produces the reduced description admitting the quantum representation ([Main]) and the Standard Model ([SM]); a boundary-thermodynamic projection that produces general relativity ([GR]); and an arrow-of-time projection (§5.4) that supplies the time-symmetric C1–C4 observer-selection theorem of [Main, §4.6], read against the apparent-horizon area, which is non-decreasing, as background kinematics, on an expanding homogeneous general-relativistic background satisfying the null energy condition.

Visible-sector projection. At the level of the embedded observer's epistemic access, the trace-out over the hidden sector produces the exact finite observable-law correspondence established in [Main]: $S \iff D \iff Q_{\mathrm{fb}}$. The fixed-basis unitary/Born representation is therefore exact and bidirectional but universal. The nontrivial OI content is history readback: C4 forces conditional memory and hidden predictive structure in every faithful completion; C1 and C3 diagnose the mediation and its capacity, while C2-structural, the persistence of the record to readback, holds by reversibility in every bijective realization and constrains coarse-grained descriptions of the hidden sector. In the spatial theory the dependency graph additionally supplies exact causal cones for finite-range couplings. Full equality with the standard coherent instrument/composite theory holds, on a finite carrier with complex matrix kinematics taken as given, exactly under the five completion conditions of [Main §3.4] and [GR §3.3]; under the substratum's stated interventions, condition (iv), full reversible control, is not supplied [GR §3.3], and whether an extended OI mechanism supplies it, and which of the other conditions the substratum supplies, are open. Bell is separate: the measurement-independent deterministic completion reaches quantum Bell violation only by ontic parameter dependence, with operational no-signaling [Main §3.3]. Applied to a cubic lattice with the wave equation as substratum dynamics — itself uniquely selected, up to the coefficient α, among second-order reversible nearest-neighbor dynamics by center independence, isotropy, and linearity ([SM, §4.1]) — this projection produces the Standard Model gauge group $\mathrm{SU}(3) \times \mathrm{SU}(2) \times \mathrm{U}(1)$ (from the stabilizer of the equivariant condensate, [SM, §§4.4–4.6], Theorem 5), three generations (physical reading conditional on H-spin', [SM §4.7]; chirality on H-χ', [SM §4.8]) and the Higgs as a composite scalar in the singlet staggered taste ([SM, §4.7], Theorems 8–11), and anomaly-free hypercharges ([SM, §4.9], Theorems 14–15). Twenty-two quantitative predictions follow ([SM, §7]), including the Cabibbo angle from a single Brillouin-zone distance, the Koide angle from a cubic-group quadratic Casimir, all three PMNS angles within 1.1σ (vs NuFIT 6.0), six fermion masses from one empirical input to within 1.3%, and all three SM gauge couplings at M_Z, reproduced through three threshold parameters fitted to the three couplings ([SM, §6.3]).

Boundary-thermodynamic projection. At the level of the partition boundary, classical horizon thermodynamics — the temperature, entropy, and dynamical evolution of the causal horizon as a gravitating object — produces general relativity through Jacobson's thermodynamic argument, conditionally on the open covariant step ([GR, §8.7]): the Clausius relation $dE = T\,dS$ applied at local causal horizons yields Einstein's equations with the same structural inputs that give the visible-sector projection its form. Applied to the cosmological horizon as a causal partition ([GR, §2]), this projection determines the emergent action scale $\hbar = c^3\epsilon^2/(4G\sigma)$ from thermal self-consistency ([GR, §3]), fixes the discreteness scale $\epsilon = 2\sqrt{\sigma}\,l_p$, for an entropy σ per boundary mode ($\epsilon = 2l_p$ at one nat per mode), as the unique simultaneous solution ([GR, §4]), recovers the Bekenstein-Hawking entropy $S = A/(4l_p^2)$, the mode count reproducing it once the calibration fixes $\epsilon/\sqrt{\sigma} = 2l_p$ and the 1/4 being the ratio $2\pi/8\pi$ of the KMS and first-law normalizations ([GR, §5]) — the action scale, the discreteness scale and the coefficient conditional on H-slope with H-count and the horizon and frame conditions of [GR, §2, §8.5]; GW250114 confirms the classical area theorem, the coefficient itself untested directly — and addresses the cosmological constant problem by identifying the quantum vacuum energy and the effective Λ as properties of logically distinct levels of description rather than commensurable quantities whose 10^{122} ratio requires cancellation ([GR, §6]), the descriptive distinction established and the dissolution conditional on the open coupling step ([GR, §6.2]). The framework carries dark energy in running-vacuum form — the functional form the local expansion shared by every smooth model, the content the Type II channel with magnitude $|\nu| \sim 10^{-32}$ (not derived at theorem level) — consistent with the Bertini et al. (2025) direct DESI-DR2-era fit at 2σ (which is consistent with the ΛCDM limit); a ~95% dark sector corollary as a structural feature rather than a new particle species; and a MOND acceleration scale $a_0 = cH/6$ from entropy displacement at the boundary ([GR, §7]).

The three projections share the same source ((S, φ)), the same trace-out machinery (marginalization over the hidden sector under conditions C1–C4), and the same structural inputs (the partition geometry, the boundary entropy, the substratum gauge group). They differ only in which structure of the embedded observer's description they derive: the visible-sector projection (§5.1, §5.3) works on the bulk dynamics and produces the quantum-representable reduced description and the Standard Model; the boundary-thermodynamic projection (§5.1) works on the classical thermal data at the partition boundary and produces general relativity; the arrow-of-time projection (§5.4) works on the horizon-area parameter along observer worldlines together with the observer-selection rule.

The projections and the Poincaré structure of the partition. The three projections are organized by how the observer-worldline-centered partition treats the Poincaré group, which makes the boost/rotation result of [Main §3.5] one facet of a larger pattern. The partition selects a preferred event — a spatial origin and a rest frame, jointly the observer's worldline — so the compact rotations about that worldline are not excluded by the partition's rest-frame data,

while the non-compact generators are broken, given that the coarse-graining map determines the horizon foliation ([Main §3.5]); spatial isotropy enters as an empirical input. Boosts, tied to the selected rest frame, surface as the cosmic rest frame (operationally the CMB frame) and the boost-sector Lorentz residual ([SM §3.1]); spatial translations, which merely relabel the observer's origin, are broken only observer-relatively and are recovered as spatial homogeneity — the equivalence of all worldline-centered descriptions; and time translation is broken by the growth of the apparent horizon on an expanding background with $\rho + p/c^2 > 0$, an input to the boundary-thermodynamic and arrow projections; in exact de Sitter the horizon area is constant (§5.4, [GR]). The partition's symmetry-breaking signature — isotropy not broken by the partition, a preferred rest frame, spatial homogeneity, a distinguished cosmic time — is the symmetry structure of a Friedmann-Robertson-Walker cosmology. The framework does not derive that structure; it exhibits the partition's symmetry-breaking pattern as compatible with it, with ordinary momentum conservation holding in the visible sector up to corrections of order (length / horizon radius).

5.2 The QM-GR incompatibility as a category error

The conventional formulation of the QM-GR unification problem treats quantum mechanics and general relativity as two competing theories in the same logical category — two attempts to describe the same regime — and asks how to merge them into a single mathematically consistent framework. Every program of unification proceeds from this assumption: quantum gravity programs treat the metric as a quantum field to be quantized; geometric programs treat the wave function as a structure on the spacetime manifold; emergent programs treat one of the two as derived from a deeper substrate that recovers the other. The persistent failure of all three approaches to produce a quantitatively confirmed unification suggests that the underlying assumption is wrong.

In the present framework, the assumption is wrong because quantum mechanics and general relativity occupy different positions in the trace-out hierarchy. They are not two theories of the same regime at all — they are two projections of the same construction, viewed at different levels. The visible-sector projection that produces the quantum-representable reduced description and the boundary-thermodynamic projection that produces GR are not in tension because they refer to *different objects in the construction*. The wave function is a property of the embedded observer's compressed description of the visible sector; the metric is a property of the classical dynamics of the partition boundary. These are not two descriptions of the same physical system at different levels of approximation. They are descriptions of two different substructures of the same total object (S, φ).

The analogy that captures the structure most clearly is the one between a thermodynamic and a statistical-mechanical description of a gas. The thermodynamic description deals with pressure, temperature, and entropy; the statistical-mechanical description deals with particle positions, momenta, and microstate

counting. These are not two competing theories of the gas, and the question "how do we unify thermodynamics and statistical mechanics?" is not a coherent question — they describe different things about the same system, and the relationship between them is one of projection (statistical mechanics produces thermodynamic quantities by averaging) rather than unification. The QM-GR relationship in the present framework is structurally similar: GR describes the thermodynamic (boundary, classical, deterministic) projection of (S, φ), and QM describes the statistical (visible-sector, compressed, stochastic-emergent) projection. The two are related by a definite procedure (the trace-out under conditions C1–C4), and the question "how do we unify them?" is recast as a category error rather than answered, conditionally on the open covariant step [GR §6.2].

This is not the same as saying that quantum mechanics is "more fundamental" than general relativity, or vice versa. Both are emergent. Both depend on the trace-out and the partition. Both are derived rather than fundamental. The fundamental object is (S, φ) — a finite set with a deterministic bijection — and neither QM nor GR is a feature of (S, φ) itself. They are features of how an embedded observer or a horizon-bounded thermodynamic regime sees (S, φ) from inside. (Nor is the ranking question recoverable from inside: by the factoring argument ([GR §6.3]), no criterion grounded in the accessible observations can separate two descriptions faithful to the same data — the fundamentality question is answered at the level of (S, φ), not adjudicated between its projections.)

5.3 The three-level hierarchy made explicit

The substratum gauge group (§4) provides the structural framework for the synthesis claim at the level of the two gauge-bearing projections (the arrow-of-time projection is developed in §5.4 and has no gauge-hierarchy analog). At the substratum level, the only object is the bijection and its gauge group $\mathcal{G}_{\text{sub}}$. The trace-out projects (S, φ) onto two derived structures simultaneously: the emergent quantum field theory at the visible-sector level, with the Standard Model gauge group from the stabilizer of the equivariant condensate ([SM, §4.4], Theorem 5) and the Standard Model representations as the cubic decomposition of the link directions ([SM, §4.6], Theorem 7); and the classical horizon thermodynamics at the boundary level, with the metric, the surface gravity, the entropy, and the temperature determined by the partition geometry and the area-law theorem ([GR, §3]). Both derived structures inherit the framework's structural inputs from the substratum, and both are exact rather than approximate at the lattice level.

The three-level gauge hierarchy organizes this. At Level 3 (the substratum), $\mathcal{G}_{\text{sub}}$ acts on (S, φ) before any trace-out and includes transformations with no analog in the emergent description (hidden-sector relabeling, alphabet change under the alphabet-size hypothesis, deep-sector enlargement). At Level 2 (the emergent QFT), the Standard Model gauge group acts on the emergent fields, obtained from the stabilizer of the equivariant condensate. At Level 1 (the emer-

gent Hamiltonian), the diagonal-unitary D-gauge acts on the emergent Hamiltonian within the emergent QM and preserves transition probabilities. The complete shift/rephasing/antiunitary classification is conditional on H-BG and the other [GR §3.3] hypotheses. The levels carry distinct equivalence notions; inclusions between them are not established.

The boundary-thermodynamic projection that produces GR is a parallel structure at Level 2 — but acting on the *partition* rather than on the internal field content. Where the emergent QFT is the trace-out's effect on the bulk dynamics, the classical horizon thermodynamics is the trace-out's effect on the partition geometry itself. Both are at Level 2. Both descend from (S, φ). The framework's claim is that the SM gauge group and Einstein's equations are not two independent theoretical inputs but two co-derived structures, both consequences of the same underlying (S, φ) and the same trace-out under the same conditions C1–C4; on the gravitational side the covariant field equations are the open Level-G3 step ([GR, §8.7]).

5.4 The arrow of time as a third projection

Main §4.6 (structural observer-selection theorem) supplies the selection content of the third projection, applied to the same underlying (S, φ) with the same trace-out machinery. The observer's confinement to the non-equilibrium phase of (S, φ), with C2 read in its timescale form (record persistence and readback there being suppressed, not excluded), is a structural consequence of the effective-mixing hypothesis, for sufficiently fast-mixing equilibria ($\tau_{\text{mix}} \ll \tau_S$), in the coarse-grained description generated by iterating the one-step effective channel. The confinement is time-symmetric and fixes where observers can exist, not a direction of time; that the arrows the observer sees descend from it together with horizon growth is not established. The third projection has the same form as the first two: a feature of how the embedded observer sees (S, φ) from inside, rather than a feature of (S, φ) itself.

A distinction should be kept in view here. The *existence* of the temporal domain — that differentiation recurs, that there is more than one moment at all — is not a projection of (S, φ) but the framework's second foundational axiom (the recurrence axiom of the two-axiom observation base; see [Main §1.2] and Part I of the companion paper *Physics Modulo Gauge*). What this section addresses is the *arrow* — the directedness of time within that domain. The framework posits the existence of the domain; for the arrow it establishes the conditional and time-symmetric results stated below, and the direction rests on boundary conditions. Conflating the two would overstate what the projection delivers.

Horizon growth as the structural clock. The framework's primary clock is the area $A(t) = 4\pi c^2/H^2$ of the apparent (Hubble) horizon, with the horizon entropy $S_{\text{dS}}(t) = A(t)/(4l_p^2)$ assigned to it at the de Sitter calibration ([GR, §2.3]). On an expanding flat Friedmann-Robertson-Walker background ($H > 0$) satisfying the null energy condition, $\dot{H} = -4\pi G(\rho + p/c^2) \leq 0$ and

$dA/dt = -8\pi c^2 \dot{H}/H^3 \geq 0$: the area is non-decreasing along any comoving observer worldline, strictly increasing while $\rho + p/c^2 > 0$, and constant in exact de Sitter. This is standard general-relativistic background kinematics, computed from the Friedmann equations of the background taken as given ([GR, §8.7]), not a prediction derived from the substratum. The criterion is expansion with the null energy condition, not $\ddot{a} > 0$: the area also grows in the decelerating era, and is constant in exact de Sitter, where $\ddot{a} > 0$. Both are inputs: the expanding branch is observed, not derived, and the null energy condition is assumed of the matter content; reading $A/(4l_p^2)$ as the entropy of a dynamical horizon goes beyond the de Sitter calibration ([GR, §2.3]). The monotonicity holds for every comoving observer of the homogeneous background. At the substrate level, (S, φ) has no preferred direction: φ and φ^{-1} are equally valid as forward-time maps. At the observer level, the horizon parameter orders the expanding branch non-strictly; that the other arrows align with it is not established.

The layered cascade. Four arrows are stated against the horizon parameter, with different status. *Cosmological:* $A(t)$ non-decreasing on the expanding branch, as above; the branch is an input, and no result here selects it. *Thermodynamic:* Jacobson's coupling $\rho_s = \rho_{\text{crit}}$ ([GR, §7.2]) is an instantaneous identity, even in H, between the boundary energy density and H^2; it does not relate matter entropy to horizon area. The direction of coarse-grained entropy increase rests on a low-entropy or product boundary condition: from a product state at $t = 0$, the sum of the sector entropies is bounded below by its value at $t = 0$ at every earlier and later time ([GR, §A.6]), a two-sided bound rather than a monotonicity statement, and the generalized second law is open. *Memory:* C2 (memory persistence) is a retention condition; the coarse-grained mixing time of the hidden-sector dynamics is the same in the two directions, and the direction in which records are written and read is fixed by a non-stationary preparation together with the observer's orientation convention. *Measurement:* the observer records "past" outcomes and prepares "future" states — an asymmetry inherited from the memory layer, not added on top of it. That the thermodynamic, memory and measurement arrows align with the horizon parameter through a common mechanism is not established. The chain uses one input (horizon growth on the expanding branch) and one structural constraint (C1–C4 observer selection via Main §4.6, conditional on its effective-mixing hypothesis and stated for the coarse-grained description generated by iterating the one-step effective channel), which fixes where observers exist, not a direction. The effective-mixing hypothesis is the premise that takes the place of an initial-condition axiom in selecting where observers exist.

What the framework establishes, and the directional-language subtlety. The framework's quantitative content at the level of Main §4.6 is *symmetric* in time direction — the spectral-gap bound $\tau_B(V) \leq \tau_{\text{mix}}(\Sigma_{\text{loc}}(V)) + O(\epsilon)$, stated for the coarse-grained description generated by iterating the one-step effective channel, applies equally in both directions along any observer worldline. What is *imposed* by the observer, not derived, is the convention of which direction along the horizon-clock parameter to call "future." The two directions are

gauge-equivalent at the substrate level ($\varphi \leftrightarrow \varphi^{-1}$), and the observer's choice of convention is the residual freedom after all physical content has been extracted. What the framework establishes is therefore a horizon parameter that is non-decreasing along observer worldlines on the expanding branch, given the inputs above, not a substrate-level arrow of time. This agrees with the premise of Ellis's objection (a bijective substratum cannot carry an arrow of time); the observed direction does not follow from the results above without the boundary conditions they name.

Scope note: causal topology as load-bearing. The memory-arrow step uses "the observer's worldline has a causal topology" as the carrier of its orientation convention. This is not a new axiom — it reduces to the ordering of moments together with φ (scope note below) — but it is load-bearing for the cascade and is noted explicitly here rather than treated as obvious. A framework in which the observer's worldline had no causal topology could not state the memory-arrow step.

Scope note: the results of this section are conditional on the ordering of moments and on further inputs. The cascade concerns a *direction* along the observer's worldline. It presupposes that the worldline's moments are *ordered* — that they form a sequence — since a monotonic parameter is monotonic with respect to an ordering. This ordering is distinct from the bare existence of more than one moment: the framework's foundational treatment (Part I of the companion paper *Physics Modulo Gauge*) axiomatizes the existence of a temporal domain but treats the *ordering* of its moments as a separate question, which it leaves open. The results of this section therefore inherit the open status of the ordering question. What this section establishes is conditional — given ordered moments and the expanding branch, a non-decreasing horizon parameter and an observer-imposed direction-convention — not a derivation of temporal directedness from the substratum alone. The causal topology of the worldline reduces to this ordering together with the dynamics φ (it introduces no commitment beyond the ordering). The ordering is not the only dependency: the horizon parameter rests on the expanding branch and the null energy condition; the observer selection on the effective-mixing hypothesis and the iterated-channel description; the thermodynamic, memory and measurement directions on a non-stationary or low-entropy preparation and the orientation convention; and the alignment of the arrows is open.

CP violation and the Ellis-class objection. The related question of CP violation is separate from the cascade. The substratum is T-symmetric (load-bearing for the reciprocity of visible transition counts, [SM, §5]; it does not by itself fix $\bar{\theta}$), and observed CP violation in CKM and kaon/B systems is a flavor-basis feature of the specific bijection rather than a consequence of the cascade. The framework treats flavor-sector CP phases as solution-specific inputs — see [SM, §5.4] for the scope statement and §6.4 "what does not falsify" for the commitment that the framework makes no prediction of δ_{CKM}, δ_{PMNS}, or the Wolfenstein (ρ, η).

5.5 What the synthesis does and does not claim

The synthesis claim is structural and architectural rather than empirical. It does not claim that the framework predicts gravitational phenomena that conventional QFT plus GR cannot reproduce — most of [GR]'s predictions (the BH area law, the basic properties of horizon thermodynamics, the broad shape of dark energy phenomenology) are also recovered by other approaches. The framework's empirical content is concentrated in [SM] (the SM derivation, twenty-two quantitative predictions) and in the specific GR-side predictions of [GR] (the Type II running-vacuum channel with $|\nu| \sim 10^{-32}$, MOND $a_0 = cH/6$ with its dimensional factor imported, dark sector concordance), which collectively distinguish the framework from standard ΛCDM plus the Standard Model.

What the synthesis claim *does* establish is that those two papers are not independent. The same construction that produces the SM in [SM] produces the gravitational sector in [GR], and the same trace-out that gives the SM gauge group from the stabilizer of the equivariant condensate ([SM, §4.4], Theorem 5) gives the BH entropy as the boundary mode count ([GR, §5]). This is not a claim about new gravitational phenomena. It is a claim about the structural relationship between two derivations that, in the conventional picture, are independent and incommensurable — and that, in the present framework, are derived from the same object by the same procedure under the same conditions.

This matters for two reasons. First, it recasts the QM-GR unification problem as a category error between two projections of one object; the covariant field equations and their source coupling (G3) remain open [GR §6.2, §8.5], so the recasting is conditional on that rung. Second, it provides an explanation for why the SM has the gauge group it does — namely, that the SM gauge group follows from the substratum construction (Stage 2) once the cubic lattice is selected from observation (§3.6, Refinement) and conditionally on H-link and H-cust [SM §4.5–§4.6], with no further model choice and no landscape of alternatives within that class.

---

6. Discussion

The reconstruction theorem fixes a local lattice/gauge residue under its stated assumptions. Theorem 24 gives representative freedoms relative to specified observation data, with completeness open. These results leave both the broader physical implementation question and the metaphysical interpretation of the reconstructed structure unresolved. The ontological hierarchy distinguishes the finite substratum from derived spatial, temporal and matter descriptions. The measurement account treats the wave function as a derived representation; its full coherent operational interpretation retains the separately stated bridge conditions. The falsification conditions in §§3–5 apply to the stated structural claims and their hypotheses.

6.1 Ontic structural realism on the equivalence class

The metaphysical commitment introduced in §1.1 can now be stated with the full technical machinery in place. The framework's theorem-level structural-realist commitment applies first to the **local reconstructed residue**: Theorem 23 fixes that local lattice/gauge residue under its stated conditions. H-Bell separately conditions whether a measurement-independent Bell-violating composite can coexist with it, and no theorem here uniquely fixes the Bell-inclusive full completion. Realism about a particular full Bell-complete bijection therefore outruns the reconstruction result.

This is the strongest claim the framework's theorems license at this level without overclaiming. Two alternative framings merit explicit comparison.

Specific-substratum realism commits to a particular representative beyond the local reconstructed residue. The fixed-interface invariances of Theorem 24 identify examples of representatives with the same stated observation data. Such data do not select between those examples. This does not forbid a further realist commitment or show that every admissible observation interface leaves every representative indistinguishable.

Pure explanatory economy — the weaker position that the framework is simply more economical than standard QM (fewer postulates, more derivations) without ontic commitment — undersells what Theorem 23 establishes about the **local reconstructed residue**. Under the theorem's stated conditions that residue is not one local possibility among many but the uniquely fixed local lattice/gauge structure modulo gauge. The Bell-inclusive completion is a separate underdetermined layer, so explanatory economy follows from local-sector reconstruction without implying full-substratum uniqueness.

The wave function, Hilbert space, emergent gauge group, and metric are features of representations — of the emergent description of a specific partition of a specific substratum — not of the equivalence class itself. At the structural level there is only the bijection and its symmetries; no quantum, no classical, no metric, no wave function. This is the structural reason the QM-GR incompatibility dissolves: both descriptions are emergent from the same structural object at different levels of projection (§5). Asking whether the wave function is "real" is then asking about a feature of the representation, and the question has a two-level answer — the wave function is real at the emergent level, gauge at the substratum level.

Within the reconstructed local observable structure, the mathematical reconstruction fixes specified invariants under stated premises while leaving Bell-completion uniqueness open. An interpretation that assigns no different predictions to two descriptions cannot be selected by those predictions alone. This observation does not classify all physical implementations or establish that every metaphysical distinction is unobservable under all admissible measurements.

Multi-level structural realism. The structural-realism commitment is not

single-level. [Structure] establishes that the framework operates at multiple levels of structural realism with different content at each level. The hierarchy is two-dimensional: an observation-hierarchy axis (vertical depth — from foundational to specific) and a gauge-hierarchy axis (horizontal breadth — from narrow to wide gauge equivalence). The combination produces a structure of realisms operating at multiple intersections.

At the broadest level — universality-class equivalence across structural classes (Level G4 in [Structure §2]) operating at the partial-trace observational features sub-class (Level C in [Structure §2]) — what is real is the algebra-channel structure of partial-trace observation: availability of a fixed-basis Born-form representation, channel-level unitarity in that representation, the non-Markovian marginal in the readback sector, and the commutant gauge-invariance pattern. These are representation/observation features shared across the stated observer-admitting class; the claim does not independently select the quadratic exponent or close the coherent local operational lift.

At intermediate levels, realism applies to the SM gauge group (forced uniquely in OI, conditionally on H-link and H-cust [SM §4.5–§4.6], but shared with any SM-reproducing universality class) and the algebra-channel structure within the SM-reproducing sub-class.

At the most class-specific theorem-supported level — the **local reconstructed residue modulo $\mathcal{G}_{\mathbf{sub}}$** within OI's structural class (Level G3 in [Structure §2]) operating at OI's specific universality-class representative (Level D in [Structure §2]) — what is real is the local lattice/gauge structure Theorem 23 fixes. A stronger realism about the full Bell-complete substratum is additional and presently underdetermined. OI's specific predictions (Cabibbo angle $1/(\pi\sqrt{2})$, Koide ratio $2/3$, dark-sector phenomenology) are real at this level, distinguishing OI's universality-class representative from alternatives within the same partial-trace-features sub-class.

These are not competing realisms; they are realism at different levels of the hierarchy. The framework commits to all of them simultaneously, with content at each level being class-specific (more specific at deeper levels) or class-universal (more universal at higher levels). What is real at the universality-class level is what observation extracts from any partial-trace operation; at the deepest theorem-supported level it is the specific **local structural residue** identified by the reconstruction theorem, with Bell-completion structure left open.

This refinement does not weaken the substratum-class realism articulated above; it situates it within a broader hierarchical structure. The framework's empirical content (twenty-two quantitative predictions, specific cubic-lattice substratum, specific gauge group derivation) is concentrated at the deepest level of the hierarchy — Level D × Level G3 — which is where the reconstruction theorem and substratum gauge group operate. The broader levels of the hierarchy are where the framework connects to other unification programs and where universality-class structural realism applies. The full articulation of the hierarchical structure

is in [Structure §2]; the substratum-class realism developed in this paper is one specific level within that broader structure.

6.2 The ontological hierarchy

The triple (S, φ, V) generates every concept in fundamental physics, not as independent substances but as different aspects of the same structure. **Space** is the coupling structure of φ — the graph G_φ determined by which degrees of freedom affect which others ([SM, §2.4]). **Matter** is the state — localized patterns that propagate through the coupling graph. **Energy** is the rate of change under iteration. **Time** is the iteration itself. **Quantum mechanics** is the observer's compressed description of the visible sector. **General relativity** is the thermodynamic limit of the coupling structure. **Conservation laws** do not all follow from bijectivity. Bijectivity conserves information and the counting measure, but a single cycle through all of S is a bijection that conserves no nonconstant function, so energy conservation rests on additional dynamical assumptions. In the emergent quantum description a Hermitian H with $e^{-iH} = U_\varphi$, the permutation unitary of φ, commutes with U_φ, and it is conserved because the same φ is applied at every step: time-translation invariance, the symmetry Noether's theorem pairs with energy. Bijectivity only makes U_φ unitary, and H generates the total system's evolution, distinct from the visible-sector emergent Hamiltonian ([GR §2.3]). The conserved quadratic invariants of the exact linear rule rest on its symmetric translation-invariant couplings (§3.2), and exact energy conservation is not shown for the rounded rule ([SM §4.2]). None of these are independent entities; they are descriptions of (S, φ, V) at different scales.

6.3 The measurement problem

On the structural reading, the measurement problem receives an observer-relative account at the fixed-basis level; the operational completion and the Frauchiger–Renner premises are separate questions ([Main §4.1]; [GR §3.3]). The wave function is not a component of (S, φ, V) — it is a derived object. Since it is derived, not fundamental, asking "does it collapse?" is asking about the behavior of a compression artifact. In the double-slit experiment, the substratum occupies a single configuration at every step, and a derivation of the interference pattern from its dynamics is open. In Wigner's friend, the Friend has a definite outcome; Wigner's superposition reflects his epistemic deficit.

Branching is forbidden by the rigidity of φ. A fixed bijection on a finite set has exactly one trajectory from any initial state. There is no point at which the trajectory splits. The appearance of branching in the emergent quantum description reflects the observer's uncertainty about which trajectory they are on (because they cannot see the hidden sector), not a physical splitting of worlds.

Bell correlations force a sharper distinction. The dependency graph gives an exact causal cone. For a response-complete deterministic completion with one-

step own-wing read/write setting interventions, a ± 1 outcome at each wing in every run read outside the other wing's graph cone, and one setting-independent pre-setting ensemble, Bell factorization follows and CHSH is capped at 2. The framework therefore does not obtain quantum Bell violation from screening plus indivisibility. It keeps measurement independence and takes ontic parameter dependence, with operational no-signaling required. Whether preparation-indexed adjacency supplies that dependence while preserving no-signaling and the continuum-curvature limit is H-Bell [Main §3.3]. The stochastic causal-local indivisible Tsirelson theorem is retained only at that weaker stochastic layer.

6.4 Pre-registered falsification conditions

The framework's structural content — the equivalence class $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$, the emergent Standard Model structure, and the co-derivation of quantum mechanics and general relativity — makes specific commitments whose future observational status can now be stated explicitly. This subsection pre-registers the conditions under which the framework would be falsified, separated from properties of the specific bijection φ that the framework does not claim to determine.

The distinction matters for two reasons. First, an explicitly pre-registered falsification schedule is the empirical counterpart of the methodological discipline developed in §§3–5: a framework that derives rather than posits must be willing to specify what would invalidate the derivation. Second, pre-registration separates structural commitments — which the framework stands or falls on — from solution-specific properties that lie outside its scope, and which continued openness on does not undermine the framework.

Class A: Structural theorems. The following are claims about the equivalence class itself and would falsify the framework's reconstruction if violated.

- *Strong CP.* The construction does not fix $\bar{\theta}$ (SM §5.3, Theorem 21: T-invariance leaves $\theta \in \{0, \pi\}$ and does not make the emergent Hamiltonian T-invariant), so the framework makes no commitment on the neutron electric dipole moment; reciprocity leaves the Jarlskog invariant free, and a mechanism making Y_u and Y_d simultaneously real would force $J = 0$ — the falsification control on H-det proposals (SM §5.3–5.5); a measured $\bar{\theta} \neq 0$ would constrain mechanisms proposed for H-det, not the T-symmetry of the substratum, which is compatible with any value of $\bar{\theta}$.
- *Gauge group SU(3) × SU(2) × U(1) exactly.* Any observation of a stable beyond-Standard-Model gauge structure — an additional Z' interpretable as a new gauge boson rather than a composite resonance, stable exotic matter transforming under a novel gauge group, or any confirmed extension of the visible-sector gauge content — falsifies the cubic-group decomposition of SM §4.6.
- *Three fermion generations.* Any observation of a stable fourth-generation fermion with Standard Model quantum numbers falsifies the taste-reduction derivation (SM §4.7). Non-stable exotic states consistent with

composite or resonance interpretation do not.

- *Majorana neutrinos with normal ordering* (on the branch where the singlet taste hosts the Higgs and no gauge-singlet fermion is present, [SM §8.4]; the ordering is a heuristic consequence of taste breaking, not a theorem). Inverted mass ordering, confirmed to high significance, falsifies the taste-breaking account of the neutrino mass matrix (SM §8.4). Confirmed Dirac nature of the neutrino (e.g., a null result for neutrinoless double-beta decay at the sensitivity that would require Majorana masses well below the branch's expectation) falsifies that branch, on which no right-handed neutrino exists in the spectrum.
- *Bell/Tsirelson test.* For the cited **stochastic causal-local indivisible** sector, $S_{\mathrm{CHSH}} \leq 2\sqrt{2}$. At the deterministic level, measurement independence plus ontic parameter independence instead gives the Bell-local ceiling 2, so the framework's Bell-violating branch is ontically parameter-dependent [Main §3.3]. A confirmed loophole-free excess above $2\sqrt{2}$ would falsify standard quantum mechanics and any claim that the operational lift lands in the standard quantum Bell set; it would not by itself falsify P-indivisibility. Bare C1–C4 or graph locality do not enforce Tsirelson.
- *Leading Lorentz-invariance-violation coefficient.* The cubic-lattice dispersion relation ([SM, §4.4] Theorem 4, generalized to d = 3: $\cos(\omega\epsilon) = \frac{1}{3}\sum_{i=1}^{3}\cos(k_i\epsilon)$ for a massless mode) yields standard emergent dispersion $\omega = |k|$ at leading order. Series expansion gives the first correction as a direction-dependent quadratic term $\omega^2 = k^2[1 - 2\delta(\hat{k})(k\epsilon)^2 + O((k\epsilon)^4)]$, with $\delta(\hat{k}) = \frac{1}{24}\sum_i \hat{k}_i^4 - \frac{1}{72}$ evaluated on the rigid $\mathbb{Z}^3$ representative. The rigid-lattice anisotropy (values $\delta_{[100]} = 1/36$, $\delta_{[110]} = 1/144$, $\delta_{[111]} = 0$ — the dispersion is exactly linear along the body diagonals, $\omega = k/\sqrt{3}$ to all orders) is a *physical, direction-dependent* prediction. Its absolute orientation in space is not a prediction: comparison (iv) of §4 relates rigid orientations of the cubic representative. The anisotropy *pattern* — its fourfold $\ell = 4$ cubic-harmonic structure and the direction-differences $\delta_{[100]} - \delta_{[111]}$ — is invariant under it, since comparison (iv) preserves the octahedral structure and does not identify $\mathbb{Z}^3$ with a generic isotropic graph (which lacks the octahedral structure the gauge group requires, [SM §4.5–4.6]). The orientation-averaged coefficient $\langle\delta\rangle = 1/90$ is a direction-summary of this anisotropy, and the framework predicts the emergent dispersion

$$\omega^2 = k^2\left[1 - \frac{(k\epsilon)^2}{45} + O\left((k\epsilon)^4\right)\right] = k^2\left[1 - \frac{4\sigma}{45}\left(\frac{E}{M_{\mathrm{Pl}}}\right)^2 + O\left(\left(\frac{E}{M_{\mathrm{Pl}}}\right)^4\right)\right]$$

after using $\epsilon = 2\sqrt{\sigma}\, l_p$ and $k\epsilon = 2\sqrt{\sigma}\, E/M_{\mathrm{Pl}}$; the coefficient of $(E/M_{\mathrm{Pl}})^2$ is $4/45$ at $\sigma = 1$. Three structural features: (i) *subluminal*, since $\delta(\hat{k}) \geq 0$ everywhere on the sphere ($\langle\delta\rangle > 0$ is invariant under orientation-averaging); (ii) *no linear-in-E/M_{Pl} term*, because the even-parity cosines in the dispersion admit no $\mathcal{O}(k^3)$ contribution — any demonstration of linear LIV at any scale therefore falsifies the framework's substrate-level commitment

to the cubic-lattice wave equation; (iii) *specific lattice coefficient* 1/45 of $(k\epsilon)^2$, fixed by the cubic-lattice coordination and isotropy-gauge average, not tunable. Features (i)–(iii) and the anisotropy pattern are the Class A content, at tree level under the stated assumptions; the directional ratios $\delta_{[100]} : \delta_{[110]} : \delta_{[111]} = 4 : 1 : 0$ are independent of σ. The physical amplitude, the coefficient $4\sigma/45$ of $(E/M_{\rm Pl})^2$, carries the undetermined entropy σ per boundary mode and is not a parameter-free numerical prediction: a measurement of it would constrain σ rather than falsify the coefficient independently. Protection of the tree-level coefficient in the interacting theory is open ([SM §3.1]). Current quadratic-LIV bounds from GRB time-of-flight reach only $E_{\rm QG} \gtrsim 10^6$ GeV (roughly thirteen orders of magnitude below $M_{\rm Pl}$), so the amplitude $4\sigma/45$ is not yet near-term testable at the magnitude level; linear-LIV tests, by contrast, are already at sensitivities where any positive detection falsifies.

Class B: Parameter-free retrodictions. The framework produces a set of numerical predictions that follow from the structural content with no adjustable parameters beyond a small set of acknowledged empirical inputs (m_s for the overall fermion mass scale, m_t for the Higgs RGE running, η for the Jarlskog phase, $Q_{\rm down}$ for the down-sector Koide normalization) and three matched coupling parameters (Class B-R). These predictions are stratified into five sub-classes reflecting their structural status:

Class B-S (structural). One prediction is a Layer 1 structural retrodiction with no open link specific to the entry; like every Class B entry it rests on the matter-sector hypotheses H-spin' and H-χ':

- The Wolfenstein parameter $A = \sqrt{2/3}$ ([SM §7.1]).

A global-fit A outside $\sqrt{2/3}\,(1\pm\lambda^2)$ would strain the leading-order identification, while the protected $\sin\theta_{13} = A^2\lambda$ comparison tests the geometry directly ([SM §7.1]).

Class B-C (conditional structural). Two predictions are Layer 1 retrodictions with no fitted input whose derivation chain has one named open link:

- The Cabibbo angle $\lambda = 1/(\pi\sqrt{2})$ ([SM §7.1]) — the chirality verification of the taste-changing vertex's spinor structure is closed via Mason et al. (HPQCD, hep-lat/0209152), establishing that taste-changing transitions in staggered quarks have spinor structure given by a combination of γ_μ and $\gamma_{5\mu}$ (both chirality-preserving); the identification leaves the emergent-theory dressing open, and the prediction is conditional on it.
- The Koide angle $\theta_0 = C_2/d^2 = 2/9$ ([SM §7.2]) — matched to observation (0.001%) and the cleanest structural derivation (cubic-group quadratic Casimir over bandwidth squared, with the §7.2 uniqueness table enumerating ten alternative dimensionless ratios that fail to match); it is conditional with the Koide amplitude, whose value $A^2 = C_2$ is not yet derived from the dynamics ([SM §7.2]).

A precision-upgrade violation would falsify the identification or the closure on which the entry is conditional.

Class B-L (layered conditional, pending open derivations). Six predictions (the two PMNS angles counted separately) follow from the structural form combined with one Layer-2 substrate input, one empirical input, or one named condition on the horizon reservoir. Each has an explicit derivational gap whose closure would move the prediction to B-S:

- The mass ratio $m_d/m_s = 1/(2\pi^2)$ ([SM §7.1]), its conversion from the Cabibbo scale running through the Gatto-Sartori-Tonin relation, an external empirical relation whose status the chain inherits.
- The mass ratio $m_u/m_d = \sqrt{\theta_0} = \sqrt{2/9}$ ([SM §7.2]) pending the explicit derivation of the "different channels" mechanism producing the square root, structurally analogous to the open OI-vertex 1-loop derivation for $K = 1/2$ ([SM §7.5]) in being mixed-layer — substratum cubic-group structure combined with electroweak-symmetry-breaking emergent machinery.
- The mass ratio $m_b/m_\tau = 4.28/Z_S$ ([SM §7.5]) pending the explicit OI-vertex 1-loop computation confirming $K = 1/2$, together with four bridge gaps in the SχPT inheritance ([SM §7.5]): the substratum measure reduction under Theorem 2, the 3D MC vs 4D SχPT dimensional bridge, the 3D BZ vs 4D Dirac taste-structure equivalence, and the 3D vs 4D parity identification of chirality. The per-vertex $\cos^2$ symmetrization producing $K = 1/2$ is dimension-independent and structurally robust; the four bridges are derivation gaps inherited by the other §7 mass-cluster predictions that share the SχPT machinery. The closure path is the two-loop SχPT calculation specializing Panagopoulos–Spanoudes 2017 to the OI cubic-group setting.
- The PMNS predictions $\sin^2\theta_{12} = 1/3 - 1/(4\pi^2)$ and $\sin^2\theta_{23} = 1/2 + 1/(2\pi^2)$ ([SM §7.3]) pending derivation of Cond 2 (the structural relation among A_2 Wilson coefficients) from cubic-lattice Yukawa structure. The reactor angle $\sin^2\theta_{13}$ is independent of Cond 2; it inherits the Cabibbo λ and is conditional with it (class C in [SM §7.6]; not counted separately in Class B).
- The MOND critical acceleration $a_0 = cH/6$ ([GR §7.3]), inheriting the conditions on the de Sitter temperature of [GR §3.2] and the coupling statement $Q = M_B c^2$ that the open G3 map would supply, with the 1/6 dimensional factor imported from Verlinde 2016 (eq. 1.7) as $(d-3)/[(d-2)(d-1)]$ in $d = 4$. Dependent predictions (baryonic Tully-Fisher, crossover radius r_M) inherit B-L status.

Class B-L violations at 3σ on precision upgrade are addressed under the pre-registration commitment via three branches: (a) closure of the open derivation *confirms* the prediction (recoverable from a transient data fluctuation); (b) closure *fails to confirm* the prediction while preserving the framework's structural commitments (recoverable error in the specific derivation, with the structural commitments intact); (c) demonstration that no consistent OI-internal deriva-

tion can produce the observed value (fatal, framework-falsifying).

Class B-M (mass chain). Two predictions inherit a single empirical input scale through a structural relation:

- The electron and muon masses m_e and m_μ via the Koide chain from the empirical input m_τ ([SM §7.2]). The prediction is structurally $\theta_0 = 2/9$ with $Q = 2/3$; the absolute scale is set by m_τ.
- The Higgs mass m_H via SM RGE running from the structural boundary condition $\lambda(M_{\rm Pl}) = 0$ ([SM §7.4]) with m_t as the empirical input.

Class B-M predictions falsify under the same precision-upgrade protocol as B-L, but the inheritance is from an empirical input scale rather than from an open derivation.

Class B-R (retrodictions). Two gauge-coupling values are explicitly retrodictions per the §6.3 [SM] parameter count:

- The SU(2) gauge coupling at M_Z ([SM §6.3, §7.6]).
- The SU(3) gauge coupling at M_Z.

Three fitted parameters (δ_0, A, B) against three observed couplings produces zero residual by construction; these classifications are honest. Class B-R predictions can never falsify in the precision-upgrade sense — their value is what falsification of the *framework's gauge-sector structural commitments* (universality of δ_0, the resummation form, the structural prediction $1/\alpha_0 = 23.25$ from the 1-loop staggered VP (ambient coefficient; native for hypercharge only under H-observer-bundle and H-Y-vertex, [SM §6.5]), and the $A \cdot B$ cross-check (≈ 48 under the $1/N^2$ leading-power assumption; inconclusive across finite-size models, [SM §6.2.1])) would test.

Status summary. Of the predictions in Class B: **1 is structural with no open link of its own** (B-S), **2 are conditional structural** (B-C), **6 are layered conditional pending open derivations** (B-L), **2 are mass-chain inheritances** (B-M), and **2 are retrodictions** (B-R). Total 13 predictions in Class B. Their empirical standing is assessed entry by entry: the gauge couplings match by construction, and the quoted precision of a measured comparison does not apply to every entry.

The current status of each is documented in the SM and GR companion papers with experimental references. As experimental precision improves, the discrimination power of the B-S set increases monotonically; closure of the open derivations would move B-C and B-L predictions to B-S over time.

The Bekenstein-Hawking coefficient $S = A/(4l_p^2)$, with the factor 1/4 the ratio $2\pi/(8\pi)$ between the Euclidean KMS period and the Jacobson Einstein-equation prefactor ([GR §5]), is a consistency relation of the horizon calibration and is not counted among them. It is conditional on the infrared detailed-balance condition H-slope on the horizon reservoir together with H-count and the horizon and frame conditions of [GR §2, §8.5]; the counting $S = \sigma A/\epsilon^2$, for an entropy σ per

boundary mode, is unconditional and the coefficient enters through $\epsilon^2 = 4\sigma\, l_p^2$, from which σ cancels. GW250114 confirms the classical area theorem, which the framework preserves; the 1/4 coefficient has no direct empirical test.

Class C: Framework-level commitments. Two broader commitments are implicit in the construction; each is stated with what would contradict it.

- *Finite classical realizability of quantum processes.* The characterization theorem ([Main, §3.4]) establishes that the quantum representation is the description admitted by an embedded observer satisfying C1–C4 on a deterministic substrate with bounded classical hidden-sector memory. At finite accuracy, every finite adaptive quantum experiment has a finite deterministic realization (the gluing and uniform-continuum theorems of [Main, §3.4]), while the exact continuum theory at any fixed finite carrier is excluded by the classical-dimension obstruction. Non-Markovian models are permitted and expected (they are how the hidden-sector correlation persistence manifests). No finite-accuracy experiment contradicts this commitment by itself. A memory-witness experiment would contradict the ontology only given the coherent-preparation and instrument bridge, an outcome-independent rule identifying fundamental open-system processes, and a constraint derived from the completed model on what those processes must show; the last is not established ([Main, §4.5]).
- *The dark sector as entropy-displacement, not as a novel particle content.* The framework's account of dark energy as boundary-entropy bookkeeping read in the Type II RVM channel, and dark matter as frozen boundary entropy (GR §7.3), is incompatible with the direct detection of a WIMP interpretable as a fundamental particle with Standard-Model-like interactions and no entropy-displacement signature. A confirmed WIMP with such properties falsifies the framework's dark-sector account while leaving the rest of the structure intact — it would require abandoning the §7.3 derivation and either identifying an error in it or accepting that the framework's dark-sector story is wrong even as other derivations survive.

Cosmological commitment. The dark energy sector is subtler than Classes A–C. The framework supplies geometry-dependent dark-energy bookkeeping whose smooth late-time expansion contains an $H^2 - H_0^2$ term — the local form shared by every smooth model — and reads it in the Type II running-vacuum channel; it does not presently derive a distinctive Type II law (GR §7.1). The magnitude is given by the emergent-QFT calculation $|\nu_{\mathrm{QFT}}| \sim (M_{\mathrm{SM}}/M_{\mathrm{Pl}})^2 \sim 10^{-32}$ (conditional on the GR §7.3 split, not yet established at theorem level), observationally indistinguishable from zero. Substratum contributions beyond this are constrained by the boundary mode count of §3-§5, which catalogues modes by spatial location alone (one mode per ϵ^2 cell, no inherent frequency assignment — a count forced by the link-direction structure of the $K = 2d$ components, [GR §3.1]); the natural 2D distribution concentrates them near the UV cutoff, contributing well below $|\nu_{\mathrm{QFT}}|$. The framework's commitment is therefore the Type II reading, with magnitude $|\nu| \approx |\nu_{\mathrm{QFT}}|$ (conditional, per

GR §7.1), and the Bertini et al. (2025) measurement $\nu = -(2.5 \pm 1.3) \times 10^{-4}$ consistent with this expectation at 2σ, where Bertini is itself consistent with zero. The falsification band is correspondingly stratified: a DESI Year 5 measurement of $|\nu| \gtrsim 10^{-4}$ at high significance would require additional substratum structure (log-uniform mode-per-decade distribution at the cosmological horizon) not currently in §3-§5; a measurement of ν with high-significance positive sign would falsify the Type II functional form itself, which is a structural commitment rather than a magnitude one. The first outcome is a refinement within the current bracket; the second is a structural falsification.

What does not falsify. The framework makes no commitment to the specific values of solution-specific quantities, and their future measurement does not bear on the framework's structural status. These include: the absolute scale of fermion masses (the choice of m_s as empirical input is acknowledged in SM §7.6); the CP-violating phases in CKM and PMNS (solution-specific properties set by the basis mismatch between up-type and down-type Yukawa mass eigenstates; $\bar{\theta}$ is not fixed by the construction and reciprocal transition counts do not force the Jarlskog invariant to vanish [SM §5.3–5.4], so the phases are free solution-level data); the numerical value of the Type II ν coefficient within the bracketed range above; initial conditions at any cosmic time (the framework derives the structure from which evolution proceeds, not the state at t_0); and the choice among representatives related by the fixed-interface invariances of Theorem 24, insofar as the proposed observation uses only the data those invariances preserve. A framework is not falsified by the presence of solution-specific inputs; it is falsified by failure of its structural commitments. The Class A–C list, together with the stratified dark-energy band, is the complete structural commitment of the framework as currently written.

The pre-registration commitment. The author commits to treating the Class A conditions, the Class C commitments, and a Class B retrodiction exceeding 3σ on precision upgrade as framework-falsifying rather than as an invitation to post-hoc rescue. In the case of a Class B violation, the framework will be revisited to determine whether the violation reflects a computational error in the specific retrodiction, an incorrect specific-bijection input, or a structural problem with the derivation; the first two are recoverable, the third is fatal. The framework does not reserve the right to add structural postulates after the fact to accommodate disconfirming evidence. Pre-registration of this commitment is itself part of the falsifiability content of the framework.

Regimes of deliberate silence. Three regimes lie outside the framework's predictive scope by construction and are not objects of falsification claims. First, the *pre-horizon early universe* — the regime before a stable causal partition with $\tau_B \sim H^{-1}$ exists — has no observer in the framework's sense ([Main, §1.3]) and no horizon-level structure for the [GR] derivations to act on; the framework's only commitment in this regime is a boundary condition on deeper theories, namely that any successful pre-horizon dynamics must produce a C1–C4-satisfying regime by the time a horizon stabilizes. Second, the *equilibrium*

phase of (S, φ) is, by the structural observer-selection theorem and with C2 read in its timescale form (record persistence and readback there being suppressed, not excluded), observer-free for sufficiently fast-mixing equilibria ($\tau_{\mathrm{mix}} \ll \tau_S$) in the coarse-grained description generated by iterating the one-step effective channel, conditional on its effective-mixing hypothesis ([Main, §4.6]); the framework makes no prediction for this regime because it has no observers to whom a prediction could be addressed. Third, regions *beyond the observer's causal horizon* are part of that observer's hidden sector and are addressed only through boundary effects (the area-law entropy of [GR, §5]); the framework takes no observer-independent position on their internal content. These silences are structural rather than methodological — they follow from the characterization theorem's identification of accessible non-Markovian observation with embedded observation under C1, C3 and C4 (quantum for the finite-configuration class of the imported stochastic–quantum correspondence [Main, §3.1]; the slow-bath form of C2's necessity for physical hidden sectors conditional on the hidden-sector mixing hypothesis), and from the observer-selection theorem's restriction of observers to the non-equilibrium phase — and stating them explicitly sharpens the scope of the positive commitments above.

---

7. Conclusion

The substratum-level results developed in this paper — the reconstruction theorem (Theorem 23) and the substratum gauge group (Theorem 24) — turn the framework's three derivations into a layered single-construction claim. Under M1-T, C2-mixing and the nearest-neighbor premise of Stage 2(b), the reconstruction fixes the **local propagating lattice/gauge residue**; M1-B and H-Bell separately condition the existence of a measurement-independent Bell-violating completion, whose uniqueness is not established. On the H-link/H-cust branch [SM §4.5–§4.6], the local reconstruction carries with the Standard Model gauge group as a forced retrodiction, and the anomaly-free hypercharge assignment forced once the observed family pattern is given, apart from the exceptional branch $Y_Q = 0$ (fermion embedding closed via the link-carrier construction, with the candidate-sector count locked at three by the link count under H-link with its single-copy clause (three physical generations only under H-spin', [SM §4.7]) jointly with the condensate-stabilizer accounting). The substratum gauge group identifies generators of the kernel of this inverse map, whose completeness is open. Together, these two results establish that the framework's emergence of quantum mechanics [Main], its derivation of the Standard Model [SM], and its derivation of the gravitational sector [GR] are not three independent applications of the same trace-out machinery but three projections of one object — the bijection (S, φ) — viewed at different levels of description. *Established* here carries the scope of Theorem 23: the local propagating lattice/gauge residue is what the reconstruction fixes. A Bell-violating completion of the same object exists only conditional on H-Bell, and its uniqueness is not established, so the

single-object claim is layered rather than total — it is a claim about the reconstructed local residue together with a conditional Bell sector, not about a uniquely selected $[(S, \varphi)]/\mathcal{G}_{\text{sub}}$ entire.

The synthesis claim that follows is not a unification of quantum mechanics and general relativity in the traditional sense. The traditional unification problem treats the two theories as competing accounts of the same regime and asks how to merge them. The present framework treats them as projections of the same construction onto two different substructures — the visible-sector trace-out and the boundary-thermodynamic limit — and the question of how to merge them is recast as a category error rather than answered as a technical problem. The result is not a theory of everything but a structural account of why the apparent incompatibility between quantum mechanics and general relativity has been so resistant to resolution: there is nothing to resolve, because the two are not in competition. They are co-derived from the same object by the same procedure under the same conditions, and the framework developed across this four-paper synthesis ([Main], [SM], [GR], [Substratum]) makes the relationship exact at the level of the reconstructed local residue, with the covariant field equations (G3) [GR §6.2] and the operational completion [GR §3.3] open.

Three sets of open problems remain. Neither these results nor this four-paper synthesis require the framework to be the final theory of physics: the substratum may itself be an effective description of something deeper, and the trace-out may have corrections beyond the leading-order results developed here. First, the *specific Bell-inclusive completion* that describes our universe is not uniquely determined by the framework. Theorem 23 fixes the local lattice/gauge residue, while H-Bell leaves the preparation-indexed Bell-composite rule underdetermined. The framework predicts the structural features of the Standard Model and the gravitational sector but not the specific values of (for example) the lightest fermion mass or the specific numerical value of the Type II RVM coefficient ν beyond its dimensional order of magnitude and sign. These are properties of the particular bijection, analogous to the mass of the sun in general relativity. Second, the *trans-Planckian regime* — the regime in which the lattice spacing $\epsilon = 2\sqrt{\sigma}\, l_p$ is not small compared to the relevant length scale — lies outside the framework's leading-order results. The framework presents the lattice as fundamental, not as a regulator approximating a continuum theory, and so does not need a continuum limit in the standard sense; but the corrections to the leading-order results in the regime where the partition geometry varies on lattice scales are an open question. Third, the *initial conditions* — the specific configuration of the bijection at any given time — are likewise not determined by the framework. The framework establishes which structures emerge from the bijection but not which microstate the universe is currently in.

These open problems are framed within the construction rather than against it. Each is sharply formulated and admits a definite (if presently unanswered) form. The progress reported here is that the structural relationship between quantum mechanics and general relativity — the central open problem of fundamental

physics for nearly a century — is fixed by the construction developed across this four-paper synthesis at the level of the reconstructed local residue, conditionally on G3 and the operational completion, and the remaining open problems are problems within the construction rather than problems with it.

Acknowledgments

AI assistance in preparing this work is disclosed at
https://github.com/amaybaum/incompleteness#ai-assistance.

References

[1] R. Bousso, "The holographic principle," *Rev. Mod. Phys.* **74**, 825 (2002).

[2] N. Bao, S. M. Carroll, and A. Singh, "The Hilbert space of quantum gravity is locally finite-dimensional," *Int. J. Mod. Phys. D* **26**, 1743013 (2017).

[3] J. Maldacena and L. Susskind, "Cool horizons for entangled black holes," *Fortsch. Phys.* **61**, 781–811 (2013); arXiv:1306.0533.

[4] S. Pedalino, B. E. Ramírez-Galindo, R. Ferstl, K. Hornberger, M. Arndt, and S. Gerlich, "Probing quantum mechanics with nanoparticle matter-wave interferometry," *Nature* **649**, 866 (2026).

[5] W. Specht, "Zur Theorie der Matrizen II," *Jahresbericht der Deutschen Mathematiker-Vereinigung* **50**, 19–23 (1940).

[6] N. A. Wiegmann, "Necessary and sufficient conditions for unitary similarity," *J. Austral. Math. Soc.* **2**, 122–126 (1961), Thm. 4; doi:10.1017/S1446788700026422.

[7] V. Futorny, R. A. Horn and V. V. Sergeichuk, "Specht's criterion for systems of linear mappings," *Linear Algebra and its Applications* **519**, 278–295 (2017), §1 and Corollary 2; doi:10.1016/j.laa.2017.01.006; arXiv:1701.08826.

[8] P. Ehrenfest, "In what way does it become manifest in the fundamental laws of physics that space has three dimensions?" *Proc. Amsterdam Acad.* **20**, 200 (1917).

[9] F. R. Tangherlini, "Schwarzschild field in n dimensions and the dimensionality of space problem," *Nuovo Cim.* **27**, 636 (1963).

[10] J. Berstel and C. Reutenauer, *Noncommutative Rational Series with Applications*, Cambridge University Press, Chapter 2, "Minimization," Theorem 2.4; doi:10.1017/CBO9780511760860.003.

[11] M. Thon and H. Jaeger, "Links between multiplicity automata, observable operator models and predictive state representations: a unified learning framework," *Journal of Machine Learning Research* **16**, 103–147 (2015).

[12] M. P. Schützenberger, "On the definition of a family of automata," *Information and Control* **4**, 245–270 (1961); doi:10.1016/S0019-9958(61)80020-X.

[13] J. W. Carlyle and A. Paz, "Realizations by stochastic finite automata," *Journal of Computer and System Sciences* **5**, 26–40 (1971); doi:10.1016/S0022-0000(71)80005-3.

[14] M. Fliess, "Matrices de Hankel," *Journal de Mathématiques Pures et Appliquées* **53**, 197–222 (1974); erratum **54**, 481 (1975).

Companion papers (cited inline by short name):

[Main] A. Maybaum, "The Incompleteness of Observation," (2026).

[SM] A. Maybaum, "The Standard Model from a Cubic Lattice," (2026).

[GR] A. Maybaum, "$\hbar$, the Bekenstein-Hawking Entropy, and the Running-Vacuum Form of Dark Energy from the Cosmological Horizon," (2026).

[Structure] A. Maybaum, "Hierarchical Structural Realism and Universality Classes of Embedded Observers," (2026).