Symmetric graphs and their automorphisms

A.A. Makhnev

Institute of Mathematics and Mechanics UB RAS
Ekaterinburg, Russia
makhnev@imm.uran.ru

Berlin, September 2009

Amply regular graph

We consider undirected graphs without loops or multiple edges. Let Γ be a graph. For vertex a of Γ the subgraph $\Gamma_i(a) = \{b \,|d(a,b) = i\}$ is called i-neighboorhod of a in Γ. We set $[a] = \Gamma_1(a)$, $a^\perp = \{a\} \cup [a]$.

The degree of a vertex a of Γ is the number of vertices in $[a]$. Γ is called regular of degree k, if the degree of any its vertex is equal k. Γ is called amply regular with parameters (v, k, λ, μ) if Γ is regular of degree k on v vertices, and $|[u] \cap [w]|$ is equal λ, if u is adjacent to w, is equal μ, if $d(u, w) = 2$. Amply regular graph of diameter 2 is called strongly regular.

Some strongly regular graphs

Let X be a set of power m. Triangular graph $T(m)$ has the vertex set $\binom{X}{2} = \{\{u, w\} \mid u, w \in X, u \neq w\}$ and $\{u, w\}$ is adjacent to $\{x, y\}$, if and only if $|\{u, w\} \cap \{x, y\}| = 1$.

Let Y be a set of power n. Grid graph $m \times n$ has the vertex set $X \times Y$ and (x_1, y_1) is adjacent to (x_2, y_2), if and only if $x_1 = x_2$ or $y_1 = y_2$.

Distance-regular graph

Let $u, w \in \Gamma$ such that $d(u, w) = i$. By $b_i(u, w)$ (by $c_i(u, w)$) we denote the number of vertices in $\Gamma_{i+1}(u) \cap [b]$ (in $\Gamma_{i-1}(u) \cap [b]$). The graph Γ with diameter d is called distance-regular with intersection array $\{b_0, b_1, ..., b_{d-1}; c_1, ..., c_d\}$ if for every $i \in \{0, ..., d\}$ the values $b_i = b_i(u, w)$ and $c_i = c_i(u, w)$ do not depend on the choice of vertices u, w at distance i. A distance-regular graph with diameter 2 is strongly regular with parameters (v, k, λ, μ), where v is the number of vertices of the graph, $k = b_0$, $\lambda = k - b_1 - 1$ and $\mu = c_2$.

Higman method

Automorphism

Let g be an automorphism of a graph Γ. Denote by $\alpha_i(g)$ the number of vertices u such that $d(u, u^g) = i$ and by $\mathrm{Fix}(g)$ the subgraph $\{a \in \Gamma \mid a^g = a\}$.

Intersection numbers

Let Γ be a distance-regular graph of diameter d on v vertices. Let us consider the symmetric association scheme $(X, \mathcal{R})$ with d classes, where X is the set of vertices of Γ and $R_i = \{(u, w) \in X^2 \mid d(u, w) = i\}$. For vertex $u \in X$ set $k_i = |\Gamma_i(u)|$. Let A_i be the adjacency matrix of the graph Γ_i, wich is corresponding to the relation R_i. Then $A_i A_j = \sum p_{ij}^l A_l$ for some integer numbers $p_{ij}^l \geq 0$, which are called the intersection numbers. Note that $p_{ij}^l = |\Gamma_i(u) \cap \Gamma_j(w)|$ for any vertices u, w with $d(u, w) = l$.

Eigenmatrices

Let P_i be the matrix in which in the (j,l) entry there is p_{ij}^l. Then the eigenvalues $k = p_1(0), ..., p_1(d)$ of the matrix P_1 are eigenvalues of Γ with multiplicities $m_0 = 1, ..., m_d$. Note that the matrix P_i is the value of some integer polynom of P_1, so the ordering of eigenvalues of the matrix P_1 gives the ordering of eigenvalues of P_i. The matrices P and Q with (i,j) entry $p_j(i)$ and $Q_{ji} = m_j p_i(j)/k_i$ are called the first and the second eigenmatrix of Γ and $PQ = QP = vI$ holds, where I is the identity matrix of order $d+1$.

Proposition 1

▸ [1, Theorem 17.12]

Let u_j and w_j be the left and the right eigenvectors of matrix P_1 affording eigenvalue $p_1(j)$ and having the first coordinate 1. Then the multiplicity m_j of the eigenvalue $p_1(j)$ is equal $v/\langle u_j, w_j \rangle$.

Character

In fact, from the proof of the Theorem 17.12 we have that w_j are the columns of the matrix P and $m_j u_j$ are the rows of the matrix Q.

The permutation representation of the group $G \leq \mathrm{Aut}(\Gamma)$ on the vertex set of Γ naturally gives the matrix representation ψ of G in $GL(v, \mathbb{C})$. The space $\mathbb{C}^v$ is the orthogonal direct sum of the eigenspaces $W_0, W_1, ..., W_d$ of the adjacency matrix A_1 of Γ. For every $g \in G$ we have $\psi(g)A_1 = A_1\psi(g)$, so the subspace W_i is $\psi(G)$-invariant. Let χ_i be a character of the representation ψ_{W_i}. Then for $g \in G$ we obtain

▸ [2,§3.7]

$\chi_i(g) = v^{-1} \sum_{j=0}^{d} Q_{ij}\alpha_j(g)$, where $\alpha_j(g)$ is the numbers of vertices x of X such that $d(x, x^g) = j$.

Note that the value of character is an integer algebraic number, and if the numbers Q_{ij} are rational then $\chi_i(g)$ is integer.
The Higman method was published in [2]. And in [2] this method was applied only to involutive automorphisms of strongly regular graph with parameters (3250,57,0,1).

Automorphisms of srg

If g is an automorphism of a strongly regular graph Γ, then either Γ has parameters $k = 2\mu$ and $\lambda = \mu - 1$ (a conference graph), or $\chi_i(g)$ is integer.
P. Cameron noted that the main difficulte in the Higman method is the calculation of parameters $\alpha_j(g)$. But in the class of graphs without triangles, for every automorphism f of ofder 3 we obtaine $\alpha_1(f) = 0$. Apart from, the structure of subgraphs of fixed points of automorphism is strongly restricted in graphs with small λ and μ. For example, M. Aschbacher noted that nonempty subgraph of fixed points of automorphism of strongly regular Moore graph (graph with $\lambda = 0$ and $\mu = 1$) is Moore graph or star.
The common properties of strongly regular graphs are in

Lemma 1

Let Γ be a strongly regular graph with parameters (v, k, λ, μ). Then $v - k - 1 = k(k - \lambda - 1)/\mu$ and one of the following holds:

1. $k = 2\mu$, $\lambda = \mu - 1$ and $v = 4\mu + 1$ is the sum of two squares of some integers;
2. $(\lambda - \mu)^2 + 4(k - \mu)$ is the square of some positive integer n, and Γ has spectrum k^1, r^f, s^{v-f-1}, where $r = (\lambda - \mu + n)/2$, $s = (\lambda - \mu - n)/2$ and $f = (s + 1)k(s - k)/(n\mu)$.

J. Seidel poset the following

Seidel Problem

Does a strongly regular graph with parameters $(99, 14, 1, 2)$ exist?

Let Γ be a strongly regular graph with parameters $(99, 14, 1, 2)$, g be an automorphism of Γ. Then for every vertex $a \in \Gamma$ the subgraph $[a]$ is the union of 7 isolated edges, Γ has spectrum $14^1, 3^{54}, -4^{44}$, and

$$Q = \begin{pmatrix} 1 & 1 & 1 \\ 44 & -88/7 & 11/7 \\ 54 & 81/7 & -18/7 \end{pmatrix}.$$

So the value of character, obtaining by projection of monomial representation on the eigenspace W_2 of the dimension 54 is equal

$$\chi_2(g) = \frac{1}{99}(54\alpha_0(g) + \frac{81}{7}\alpha_1(g) - \frac{18}{7}\alpha_2(g) =$$

$$\frac{1}{77}(42\alpha_0(g) + 9\alpha_1(g) - 2\alpha_2(g).$$

By substitution $\alpha_2(g) = 99 - \alpha_0(g) - \alpha_1(g)$ we have

$$\chi_2(g) = (4\alpha_0(g) + \alpha_1(g) - 18)/7.$$

Automorphisms of Seidel graph

Involutive automorphisms

A. Makhnev and I. Minakova investigated involutive automorphisms of strongly regular graph with parameters $(99, 14, 1, 2)$.

Let t be an involutive automorphisms of strongly regular graph with parameters $(v, k, 1, 2)$, $\Omega = \mathrm{Fix}(t)$. If $a \in \Omega, b \in \Omega - a^{\perp}$, then $[a] \cap [b] = \{u, u^t\}$ is a 2-coclique. If $u \in [a] - \Omega$ and $\{u, u^t\}$ is a 2-coclique, then $[u] \cap [u^t] = \{a, b\}$ and $b \in \Omega - a^{\perp}$. So $|\Omega| = 1 + |\Omega(a)| + 2l$, where l — is the number of t-orbits of size 2 on the edges in $[a]$.

Proposition 2

Let Γ be a strongly regular graph with parameters $(99, 14, 1, 2)$, having involutive automorphism t. Then $\mathrm{Fix}(t)$ is one of the following graphs:

1. one-vertex graph;
2. triangle;
3. three isolated triangles;
4. vertex and two isolated triangles;
5. four isolated vertices and triangle;
6. n-coclique, $n = 3, 5$ or 7;
7. 3×3-grid.

Only in the case (1) the value $\chi_2(t)$ is integer. Consider, for example, the case, when $\mathrm{Fix}(t)$ is the union of 4-coclique and triangle. Then t has 3 orbits of size 2 on the edges in $[a]$, if a is isolated in $\mathrm{Fix}(t)$ and t has 2 orbits of size 2 on the edges in $[b]$, if b belongs to triangle of $\mathrm{Fix}(t)$, so $\alpha_1(t) = 4 \cdot 2 + 3 \cdot 4 = 20$ and $\chi_2(t) = (28 + 20 - 18)/7$, a contradiction.

Theorem 1 ▸ Makhnev A., Minakova I. [3]

Let Γ be a strongly regular graph with parameters $(99, 14, 1, 2)$, g an element of prime order p from $\mathrm{Aut}(\Gamma)$ and $\Delta = \mathrm{Fix}(g)$. Then one of the following holds:

1. Δ is one vertex graph and $p = 2$ or 7;
2. Δ is empty graph and $p = 3$ or 11;
3. Δ is triangle and $p = 3$.

H. Wilbrink [4] proved that Γ does not admit an automorphism of order 11. In particular, the order of the automorphism group of strongly regular graph with parameters $(99, 14, 1, 2)$ divides $2 \cdot 3^3 \cdot 7$.

Corollary 1

Let Γ be the strongly regular graph with parameters $(99, 14, 1, 2)$, $G = \mathrm{Aut}(\Gamma)$. If G contains an involution t, then one of the following holds:

1. $|G|$ is divided by 7 and divides 42, $[O(G), t] = 1$ and in the case $|G| = 42$ the subgroup $O(G)$ is nonabelian;
2. $|G|$ divides 6.

Graphs with $\lambda = 0$ and $\mu = 2$

Known srg without triangles

The known strongly regular graph without triangles are in 3 classes:

1. Bipartite graphs $K_{k,k}$, $k \geq 2$.
2. Moore graphs (pentagon, Petersen graph with parameters (10,3,0,1) and Hoffman-Singleton graph with parameters (50,7,0,1)),
3. Clebsh graph with parameters (16,5,0,2), Gewirtz graph with parameters (56,10,0,2), Higman-Sims graph with parameters (100,22,0,6), graph with parameters (77,16,0,4) (the second neighborhood of vertex in the Higman-Sims graph).

It is old problem in the theory of symmetric graphs.

Problem 2

Is a strongly regular graph with $\lambda = 0$ and $\mu = 2$ the quadrangle, Clebsh graph or Gewirtz graph?

Let Γ be a strongly regular graph with parameters $(v, k, 0, 2)$. Then $k = w^2 + 1$ for some integer w, Γ has spectrum $k^1, (w-1)^f, -(w+1)^{v-f-1}$, where $f = w(w^2+1)(w^2+w+2)/(4w)$, so w is not divided by 4.

A. Makhnev and V. Nosov investigated automorphisms of strongly regular graph Γ with parameters $(v, k, 0, 2)$, where $v = (w^4 + 3w^2 + 4)/2$, $k = w^2 + 1$. Then

$$Q = \begin{pmatrix} 1 & 1 & 1 \\ \frac{(w^2+1)(w^2-w+2)}{4} & -\frac{w^3+w+2}{4} & \frac{w^2-w+2}{2w} \\ \frac{(w^2+1)(w^2+w+2)}{4} & \frac{w^3+w-2}{4} & -\frac{w^2+w+2)}{2w} \end{pmatrix}.$$

So the value of character, obtaining by projection of monomial representation on the eigenspace W_1 of the dimension $(w^2+1)(w^2-w+2)/4$ is equal

$$\chi_1(g) = \frac{1}{v}((w^2+1)(w^2-w+2)\alpha_0(g)/4-$$

$$(w^2+w+2)\alpha_1(g)/4+(w^2-w+2)\alpha_2(g)/(2w)).$$

By substitution $\alpha_2(g) = v-\alpha_0(g)-\alpha_1(g)$ we have

$$\chi_1(g) = \frac{1}{2w}((w-1)\alpha_0(g)-\alpha_1(g)+(w^2-w+2)).$$

Proposition 3

Let Γ be a strongly regular graph with parameters $((w^4+3w^2+4)/2, w^2+1, 0, 2)$, f — an element of order 3 from $\mathrm{Aut}(\Gamma)$ and $\Omega = \mathrm{Fix}(f)$. Then one of the following holds:

1. Ω is an edge, w is odd and divided by 3 (this case is in Gewirtz graph);
2. Ω is strongly regular graph with parameters $((u^4+3u^2+4)/2, u^2+1, 0, 2)$, $u^2 \equiv w^2 \pmod 3$ and w divides $(u^4+3u^2)/2$.

N Nakagawa [5] proved that if Γ is a strongly regular graph with parameters $(v, q^2+1, 0, 2)$, $q = p^n$, p is odd prime number, and $G = \mathrm{Aut}(\Gamma)$ contains the subgroup $H = PGL_2(q^2)$, fixing the vertex a and is transitive on $\Gamma_2(a)$, then $q = 3$ and Γ is Gewirtz graph.

Proposition 4

Let Γ be a strongly regular graph with parameters $(v, q^2+1, 0, 2)$, t — an involutive automorphism of Γ, $a \in \mathrm{Fix}(t)$ and t acts on $\Gamma_2(a)$ as an involution of $G = PGL_2(q^2)$ by conjugating on z^G, where z is the involution of G', then $q = 3$ and Γ is Gewirtz graph.

Theorem 2 ▸ A. Makhnev, V. Nosov [6]

Let Γ be a strongly regular graph with parameters $(352, 26, 0, 2)$, g be an element of prime order p from $\mathrm{Aut}(\Gamma)$ and $\Omega = \mathrm{Fix}(g)$. Then one of the following holds:

1. $p = 2$, either Ω is the empty graph, 14-coclique, connected graph of degree 6 on 32 vertices, or Ω has four connected components, that are isomorphic to a quadrangle;
2. $p = 5$ and Ω is the edge;
3. $p = 11$ and Ω is an empty graph;
4. $p = 13$ and Ω is one vertex graph.

Theorem 2 implies that the order of the automorphism group G of strongly regular graph with parameters $(352, 26, 0, 2)$ divides $2^l \cdot 5^2 \cdot 11 \cdot 13$ and G is the solvable group.

Krein graphs

Krein conditions

Let strongly regular graph Γ with parameters (v, k, λ, μ) have eigenvalues k, r, s. If graphs Γ and $\bar{\Gamma}$ are connected then the next unequalities (Krein conditions) hold:
(1) $(r+1)(k+r+2rs) \leq (k+r)(s+1)^2$ and
(2) $(s+1)(k+s+2rs) \leq (k+s)(r+1)^2$.

Γ is called Krein graph, if we have equality in (1) or in (2). It is interesting that in Krein graph the first and the second neighboorhods of any vertex are strongly regular. Krein graph without triangles has parameters $((r^2+3r)^2, r^3+3r^2+r, 0, r^2+r)$ and we denote this graph as $Kre(r)$. For each $r = 1, 2$ there is the unique graph $Kre(r)$ — the Clebsh graph and the Higman-Sims graph respectively. By [7] graph $Kre(3)$ does not exist.

Character of $Kre(r)$

Let Γ be a Krein graph $Kre(r)$. Then

$$P = \begin{pmatrix} 1 & 1 & 1 \\ r^3+3r^2+r & -(r^2+2r) & r \\ r^4+5r^3+6r^2-r-1 & r^2+2r-1 & -r-1 \end{pmatrix}$$

and $P = Q$. So the value of character, obtaining by projection of monomial representation on the eigenspace W_1 of the dimension r^3+3r^2+r is equal
$\chi_1(g) = (r^2+3r)^{-2}((r^3+3r^2+r)\alpha_0(g)-(r^2+2r)\alpha_1(g)+r\alpha_2(g))$.
By substitution $\alpha_2(g) = v - \alpha_0(g) - \alpha_1(g)$ we have

$$\chi_1(g) = (r\alpha_0(g) - \alpha_1(g) + r^2(r+3))/(r^2+3r).$$

Proposition 5

Let Γ be a strongly regular graph with parameters $((r^2+3r)^2, r^3+3r^2+r, 0, r^2+r)$, g be an automorphism of order 3 of Γ and $\Omega = \mathrm{Fix}(g)$. Then $|\Omega|$ is divided by $r+3$ and if r is not divided by 3, then the following hold:

1. either Ω is a quadrangle and Γ is Clebsh graph, or Ω contains 3-coclique and every 3-coclique of Ω is subset of $[a]$ for some vertex $a \in \Omega$;
2. either Γ is Clebsh graph, or degree of every vertex in Ω is greater 2;
3. if Ω contains two nonadjacent vertices, not belonging any 3-coclique of Ω, then either Ω is quadrangle or Ω is $K_{r+3,r+3}$ without matching and $r \equiv 0, 2 \pmod 3$;

Example 1

Consider automorphisms of the Higman-Sims graph Γ (Krein graph with $r = 2$). Let g be an automorphism of prime order p of Γ. Then 10 divides $2\alpha_0(g) - \alpha_1(g)$ and for the graph $\Omega = \mathrm{Fix}(g)$ the following holds:

a) Ω is an empty graph, $|g| = 2$ or 5 and $\alpha_1(g)$ is divided by 10;

b) $\Omega = \{a\}$ is one vertex graph, $|g| = 11$ and $\alpha_1(g) - 2$ is divided by 10;

c) Ω is an edge, $|g| = 7$ and $\alpha_1(g) = 14$;

d) $|g| = 2$ and Ω is a 2-coclique extension of Petersen graph, 6-coclique or amply regular graph with parameters $(30, 8, 0, 4)$;

e) $|g| = 3$, Ω is $K_{5,5}$ without matching and $\alpha_1(g) = 0$;

f) $|g| = 5$, g fixes pentagon and $\alpha_1(g) = 20$.

Thus, for $G = \mathrm{Aut}(Kre(2))$ we have $\pi(G) \subseteq \{2, 3, 5, 7, 11\}$. In fact, G is the automorphism group of the Higman-Sims group and $|G| = 2^{10}3^2 5^3 7 \cdot 11$.

Theorem 3 ▸ A. Gavrilyuk [9]

Let Γ be a graph $Kre(4)$, g be an element of prime order p of $\mathrm{Aut}(\Gamma)$ and $\Omega = \mathrm{Fix}(g)$. Then either $p = 2$ or $p = 7$ and Ω is empty graph.

Theorem 4 ▸ A. Makhnev, V. Nosov [8]

Let Γ be a graph $Kre(5)$, g be an element of odd prime order p of $\mathrm{Aut}(\Gamma)$ and $\Omega = \mathrm{Fix}(g)$. Then one of the following holds:

1. Ω is empty graph and $p = 5$;
2. either $|\Omega| = 1$ and $p = 41$, or Ω is 2-clique and $p = 17$;
3. Ω is $K_{8,8}$ without matching and $p = 3$.

Generalized poligons

The incidence system $(X, \mathcal{L})$, where X is a point set and $\mathcal{L}$ is a line set, is called near $2n$-gon of order (s, t) if every line contains $s + 1$ points, every point belongs to $t + 1$ lines (distinct lines intersect in at most one point), diameter of collinearity graph Γ is n and for any pair $(a, L) \in (X, \mathcal{L})$ the line L contains a unique point with minimal distance of a in Γ. Near $2n$-gon is called generalized $2n$-gon, if every two point u, w with $d_\Gamma(u, w) < n$ belongs to the unique geodesic way in Γ from u to w.
Generalized $2n$-gon of order (s, t) is called thick, if $s > 1$ and $t > 1$.

Generalized octagon

It is known the existence of thick generalized octagon only for $\{s, t\} = \{q, q^2\}$, where $q = 2^{2l-1}$. Apart from, known $GO(q, q^2)$ corresponds to the building of the group ${}^2F_4(q)$.

I. Belousov and A. Makhnev investigated automorphisms of distance-regular graph with intersection array $\{10, 8, 8, 8; 1, 1, 1, 5\}$ on 1755 vertices. This graph have spectrum $10^1, 5^{351}, 1^{650}, -3^{675}, -5^{78}$, and is the collinearity graph of generalized octagon $GO(2, 4)$ (see [10,chapter 6]).

An involution t of the finite group G is said to be central, if $|G : C_G(t)|$ is odd.

Example 2

In the case $q = 2$ vertices of the graph Γ are central involutions of the Tits group ${}^2F_4(2)'$, and two central involutions u, w are adjacent if and only if uw also is central involution in ${}^2F_4(2)'$. Further, $d(u, w) = 2$ if and only if uw is noncentral involution in ${}^2F_4(2)'$, $d(u, w) = 3$ if and only if $|uw| = 4$ (and in this case $(uw)^2$ is noncentral involution in ${}^2F_4(2)'$), $d(u, w) = 4$ if and only if $|uw| = 5$.

Let Γ be a distance-regular graph with intersection array $\{10, 8, 8, 8; 1, 1, 1, 5\}$ on 1755 vertices. Then

$$Q = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 351 & 351/2 & 351/8 & 0 & -351/64 \\ 650 & 65 & -325/4 & -65/8 & 325/32 \\ 675 & -405/2 & 135/8 & 135/8 & -675/64 \\ 78 & -39 & 39/2 & -39/4 & 39/8 \end{pmatrix}.$$

So $\chi_1(g) = 351/1755(\alpha_0(g) + \alpha_1(g)/2 + \alpha_2(g)/8 - \alpha_4(g)/64)$ and $\chi_1(g) = (64\alpha_0(g) + 32\alpha_1(g) + 8\alpha_2(g) - \alpha_4(g))/(5 \cdot 64)$.
Further, $\chi_2(g) =$
$65/1755(10\alpha_0(g) + \alpha_1(g) + 5\alpha_2(g)/4 - \alpha_3(g)/8 + 5\alpha_4(g)/32)$. By substitution $\alpha_4(g) = 1755 - \alpha_0(g) - \alpha_1(g) - \alpha_2(g) - \alpha_3(g)$, we have
$\chi_2(g) = (35\alpha_0(g) + 3\alpha_1(g) - 5\alpha_2(g) - \alpha_3(g))/96 + 325/32$.

Also $\chi_3(g) =$
$135/1755(5\alpha_0(g) - 3\alpha_1(g)/2 + \alpha_2(g)/8 + \alpha_3(g)/8 - 5\alpha_4(g)/64)$.
By substitution $\alpha_2(g) + \alpha_3(g) = 1755 - \alpha_0(g) - \alpha_1(g) - \alpha_4(g)$,
we have

$$\chi_3(g) = (24\alpha_0(g) - 8\alpha_1(g) - \alpha_4(g))/64 + 135/8.$$

Finally,
$\chi_4(g) = 39/1755(2\alpha_0(g) - \alpha_1(g) + \alpha_2(g)/2 - \alpha_3(g)/4 + \alpha_4(g)/8)$.
By substitution $\alpha_4(g) = 1755 - \alpha_0(g) - \alpha_1(g) - \alpha_2(g) - \alpha_3(g)$,
we have

$$\chi_4(g) = (5\alpha_0(g) - 3\alpha_1(g) + \alpha_2(g) - \alpha_3(g))/120 + 39/8.$$

Theorem 5

▸ I. Belousov, A. Makhnev [11]

Let Γ be a distance-regular graph with intersection array $\{10, 8, 8, 8; 1, 1, 1, 5\}$, g be an element of prime order p of $\mathrm{Aut}(\Gamma)$ and $\Omega = \mathrm{Fix}(g)$. Then one of the following holds:

1. $p = 3$ or 13 and Ω is an empty graph;
2. $p = 5$ and Ω is the union of 5ω isolated vertices, and $d(u, w) = 4$ for every two vertices $u, w \in \Omega$;
3. $p = 2$, $|\Omega|$ is odd and either

   (i) Ω contains triangle $\{x, y, z\}$ such that Ω is the union of balls with radius 1 and centers in $\{x, y, z\}$;

   (ii) Ω contains the vertex b such that $[b] \subset \Omega$, Ω is a subset of the ball with radius 2 and center b, and there are two nonadjacent vertices $a, c \in [b]$, such that $b^{\perp}$ does not contain $\Omega(a)$ and $\Omega(c)$.

Let us prove, for example, that if Fix(g) is an empty graph, then $p \neq 5$.

Let $p = 5$. Then $\alpha_i(g)$ is divided by 5 and $\alpha_0(g) = \alpha_1(g) = 0$. As $\chi_3(g) = (24\alpha_0(g) - 8\alpha_1(g) - \alpha_4(g))/64 + 135/8$ is integer, then $\alpha_4(g) = 40r$ and $5(r-3)$ is divided by 8. So $r = 8r' + 3$ is odd.

As $\chi_4(g) = (5\alpha_0(g) - 3\alpha_1(g) + \alpha_2(g) - \alpha_3(g))/120 + 39/8$ is integer, then $\alpha_2(g) = 5s$, $\alpha_3(g) = 5u$, $s - u$ is divided by 3 and $(s-u)/3 - 1$ is divided by 8. So $s - u = 3(8y+1)$. Further, $\sum \alpha_i(g) = 5s + 5u + 40r = 1755$, so

$s + u = 351 - 8r = 2u + 24y + 3$ and $u = 174 - 12y - 4r$.

As $\chi_2(g) = (35\alpha_0(g) + 3\alpha_1(g) - 5\alpha_2(g) - \alpha_3(g))/96 + 325/32$ is integer, then $(5s+u)/3 - 1 = 2u + 40y + 4$ is divided by 32.

Thus, $u + 20y + 2 = 176 + 8y - 4r$ is divided by 16, a contradiction with r is odd.

Corollary 2

Let Γ be a distance-regular graph with intersection array $\{10, 8, 8, 8; 1, 1, 1, 5\}$ and a group $G = \mathrm{Aut}(\Gamma)$ acts transitively on the set of vertices of Γ. Then Γ is the collinearity graph of the generalized octagon of Tits group ${}^2F_4(2)'$.

Generalized hexagon

I. Belousov and A. Makhnev investigated automorphisms of distance-regular graph with intersection array $\{84, 81, 81; 1, 1, 28\}$ on 26572 vertices. This graph is the collinearity graph of generalized hexagon $GH(3, 27)$ and does not contain m-circles for $4 \leq m \leq 5$. It is known the existence of thick generalized hexagon only for $\{s, t\} = \{q, q\}$ or $\{s, t\} = \{q, q^3\}$, where $q = p^l$, p is a prime number. Apart from, $GH(q, q)$ corresponds to the building of the group $G_2(q)$ and $GH(q, q^3)$ corresponds to the building of the group ${}^3D_4(q)$.

It is known (see [12]), that there is a unique generalized hexagon $GH(2, 8)$. Let Γ be a distance-regular graph with intersection array $\{q(q^3 + 1), q^4, q^4; 1, 1, q^3 + 1\}$ on $(q^3 + 1)(q^2 + q + 1)(q^4 - q^2 + 1)$ vertices.

This graph has eigenvalues $q(q^3+1), q^2+q-1, -(q^2-q+1), -(q^3+1)$ with multiplicities $1, \frac{1}{2}q^3(q^3+1)^2, \frac{1}{2}q^3(q+1)^2(q^4-q^2+1), q^5-q^3+q$ respectively, and is the collinearity graph of $GH(q,q^3)$. Further

$$\chi_1(g) = \frac{(q^2+1)(q^3+1)\alpha_0(g) + (q^3+q^2+1)\alpha_1(g) + \alpha_2(g)}{2q^3(q^2+q+1)} - \frac{(q^3+1)^2}{2q^3}.$$

$$\chi_2(g) = \frac{(q^2-1)(q^3+1)\alpha_0(g) - (q^3-q^2+1)\alpha_1(g) - \alpha_2(g)}{2q^3(q^2-q+1)} + \frac{(q+1)^2(q^4-q^2+1)}{2q^3}.$$

$$\chi_3(g) = \frac{(q^2-q+1)\alpha_0(g) - (q-1)\alpha_1(g) + \alpha_2(g)}{q^2(q^2-q+1)(q^2+q+1)} - \frac{q^4-q^2+1}{q^2}.$$

Next two lemmas have independent interest.

Lemma 2

Let Γ be a distance-regular graph with intersection array $\{q(q^3+1), q^4, q^4; 1, 1, q^3+1\}$. If Γ contains proper subgraph Δ such that Δ is the collinearity graph of generalized hexagon of order (q,t), then $t \leq q$.

Proof

Let Γ contain proper subgraph Δ such that Δ is the collinearity graph of generalized hexagon of order (q,t). Then the number vertices of Δ is equal $v' = (q+1)(q^2t^2+qt+1)$, the degree of the graph Δ is $k' = q(t+1)$ and the number of edges between Δ and $\Gamma - \Delta$ is equal $v'(k-k') = (q+1)(q^2t^2+qt+1)q(q^3-t)$. As every vertex of $\Gamma - \Delta$ is adjacent to at most one vertex of Δ, then $v = (q+1)(q^8+q^4+1)$ and $v \geq v' + v'(k-k')$, so $t^3 - q^3t^2 - q^2t + q^5 = (t-q)(t+q)(t-q^3) \geq 0$. Hence $t \leq q$.

Lemma 3

Let Γ be a distance-regular graph with intersection array $\{q(q^3+1), q^4, q^4; 1, 1, q^3+1\}$, g be an element of prime order p of $\mathrm{Aut}(\Gamma)$ and $\Omega = \mathrm{Fix}(g)$ is nonempty subgraph. Then:

1. if Ω is nonconnected, then Ω is coclique, and if $p > q$, then p divides $q^3 + 1$;
2. if Ω is connected and $p > q$, then Ω is the collinearity graph of generalized hexagon of order (q, t), and either $t = 1$, or the number qt is the square, $q \leq t^3$ and $t \leq q$.

Proof

Let Ω be a nonconnected subgraph. Then vertices from distinct connected components of Ω are at the distance 3 in Γ. Let a, b are adjacent in Ω and c belongs to other connected component of Ω. By definition of generalized hexagon the clique $a^\perp \cap b^\perp$ contains the unique vertex e with $d(c, e) = 2$. The contradiction with $d \in \Omega$.

Hence, Ω is a coclique.
Let $p > q$. Then $k = q(q^3 + 1)$, so p devides $q^3 + 1$. The statement (1) is proved.
Let Ω be connected and $p > q$. Consider two adjacent vertices a, b. Then the clique $L = a^\perp \cap b^\perp$ is a subset of Ω. Further g acts on the set of q^3 maximal cliques from $b^\perp$ different of L and fixes one more clique L_1 in this set. For $c \in L_1 - \{b\}$ we get the maximal clique L_2 from $c^\perp$, fixed by g.
Note that every two vertices $a, e \in \Omega$, that are antipodal in Γ, have the same degree in Ω. By connectivity Ω there is the vertex $b \in \Omega(a) \cap \Gamma_2(e)$ such that the clique $a^\perp \cap b^\perp$ is a subset in Ω and contains the unique vertex c with $d(c, e) = 2$. Further for every vertex $b \in \Omega(a) \cap \Gamma_2(e)$ there is a unique vertex c adjacent with b from $\Omega(d) \cap \Gamma_2(a)$ and $c^\perp \cap e^\perp \subset \Omega$. As $\mu = 1$, so distinct vertices $b, b' \in \Omega(a) \cap \Gamma_2(e)$ correspond to distinct vertices c, c' and $|\Omega(a)| = |\Omega(e)|$.

We have $p_{33}^1 = q^8(q-1)$, so for adjacent vertices a, b of Ω there is a vertex $e \in \Omega$ such that $d(a,e) = d(b,e) = 3$. Hence Ω is regular graph. Thus Ω is the collinearity graph of near $2d$-gon with $d = 3$, and $c_2(\Omega) = 1$. So Ω is the collinearity graph of generalized hexagon of order (q,t), and by theorem 6.5.1 of [10] either $t = 1$, or the number qt is the square, $q \leq t^3$ and $t < q^3$. By Lemma 2 we have $t \leq q$.

Theorem 6 ▸ I. Belousov, A. Makhnev [11]

Let Γ be a distance-regular graph with intersection array $\{84, 81, 81; 1, 1, 28\}$, g be an element of prime order p of $\mathrm{Aut}(\Gamma)$ and $\Omega = \mathrm{Fix}(g)$. Then one of the following holds:

1. $p = 7, 13$ or 73 and Ω is an empty graph;
2. $p = 7$ and Ω consists of 7ω vertices, with $d(a, b) = 3$ for every two vertices $a, b \in \Omega$;
3. $p = 13$ and Ω is the collinearity graph of $GH(3, 1)$;
4. $p = 3$, $|\Omega| \equiv 1 \pmod 3$ and either
   (i) Ω is 1-clique or 4-clique, or
   (ii) $\Omega \subseteq a^{\perp}$ for some vertex a, or
   (iii) Ω is the collinearity graph of $GH(3, 3)$, or
   (iv) Ω contains 4-clique L, such that Ω is the subset of the union of balls with radius 1 and centers in L;
5. $p = 2$, $|\Omega|$ is even and either
   (i) Ω is $54m + 28$-coclique, where $0 \leq m \leq 5$, or
   (ii) Ω is the collinearity graph of $GH(1, 9)$, or
   (iii) Ω is the collinearity graph of $GH(3, 3)$, or
   (iv) $|\Omega| = 116$, Ω contains four vertices of degree 28 and twenty eight 4-cliques, whose vertices have degree 4 in Ω.

Let Γ — be a collinearity graph of generalized $2n$-gon, $G = \mathrm{Aut}(\Gamma)$. If G acts transitively on the set of $2n$-gons in each of graphs: collinearity graph, line graph and flag graph, then generalized $2n$-gon is classic [12].

Corollary 3

Let Γ be a vertex-symmetric distance-regular graph with intersection array $\{84, 81, 81; 1, 1, 28\}$. Then Γ is the collinearity graph of classic generalized hexagon $GH(3, 27)$, affording to the Steinberg group ${}^3D_4(3)$.

Lemma 4

Let Γ be a distance-regular graph with intersection array $\{84, 81, 81; 1, 1, 28\}$ and $G = \mathrm{Aut}(\Gamma)$. Then the following statments hold:

1. G does not contain an element of order 49, and 7^3 does not divided $|G|$;
2. G does not does not contain an element of order 169, and 13^3 does not divided $|G|$;
3. if G contains an element f of order 73, then $C_G(f) = \langle f \rangle$.

Proof of the corollary 3

Let Γ be a distance-regular graph with intersection array $\{84, 81, 81; 1, 1, 28\}$ and $G = \mathrm{Aut}(\Gamma)$. Then $|G|$ divides $2^\beta 3^\gamma 7^2 13^2 73$. Let G act transitively on the set of vertices of Γ. Fix the vertex a of Γ and let G_a be a stabilizer a in G. Then $|G_a|$ divides $2^{\beta-2} 3^\gamma \cdot 7 \cdot 13$ and $|G : G_a| = 4 \cdot 7 \cdot 13 \cdot 73$.

Let $P = \langle g \rangle$ be a Sylow 73-subgroup of G. We have following statements.

1. The factorgroup $\bar{G} = G/O_2(G)$ has a simple socle T and $\bar{G}/T$ is the subgroup of the cyclic group of order 72.
2. Let T be a simple group, $|T|$ devides $2^\beta 3^\gamma 7^2 13^2 73$ and devided by 73. If T does not contain elements of order $73p$ for any $p \in \{2, 3, 7, 13\}$, then T is isomorphic to the Steinberg group ${}^3D_4(3)$.
3. $O_2(G) = 1$.

From (1–3) it is follows that Γ is the collinearity graph of classic generalized hexagon $GH(3, 27)$, affording to the Steinberg group ${}^3D_4(3)$.

Graphs with block-designs

A. Gavrilyuk and A. Makhnev investigated amply regular graphs with diameter $d \geq 3$, such that for some vertex $a \in \Gamma$ the pair $(\Gamma_d(a), \Gamma_{d-1}(a))$ is a 2-(V, K, Λ) design [14]. In this case $\Gamma_d(a)$ is clique, coclique or strongly regular graph with parameters $(v' = V, k' = k - R, \lambda' = \lambda - \Lambda, \mu' = \mu - \Lambda)$.
For distance-regular graphs with diameter $d \geq 3$, such that for every vertex $a \in \Gamma$ the pair $(\Gamma_d(a), \Gamma_{d-1}(a))$ is a 2-design can be wright the following.

Conjecture Σ

Let Γ be a distance-regular graphs with diameter d, such that for every vertex $a \in \Gamma$ the pair $(\Gamma_d(a), \Gamma_{d-1}(a))$ is a 2-design, then the subgraph $\Sigma = \Gamma_d(a)$ is clique or coclique.

This Conjecture is based on the fact that in the list of feasible arrays in [10] only two arrays $\{60, 45, 8; 1, 12, 50\}$ and $\{49, 36, 8; 1, 6, 42\}$ correspond to graphs satisfying conditions of the Conjecture Σ. In the first case $\Gamma_3(a)$ is 6×6-grid, and in the second case $\Gamma_3(a)$ is the union of seven isolated 8-cliques. It is proved in [14] that a distance-regular graph with intersection array $\{60, 45, 8; 1, 12, 50\}$ contains a vertex x such that $\Gamma_3(x)$ is not 6×6-grid. In the second case the pair $(\Gamma_3(a), \Gamma_2(a))$ is a 2-(V, K, Λ) design, where $V = 56$, $K = 8$ and $\Lambda = 42 \cdot 7/55$.

In [15] it is investigated automorphisms of distance-regular graph with intersection array $\{60, 45, 8; 1, 12, 50\}$. This graph have the spectrum $60^1, 14^{45}, 0^{207}, -10^{69}$ and is Q-polinomial. Further

$$Q = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 45 & 21/2 & -1 & -25/2 \\ 207 & 0 & -23/5 & 23 \\ 69 & -23/2 & 23/5 & -23/2 \end{pmatrix}.$$

So $\chi_2(g) = 1/322(207\alpha_0(g) - 23\alpha_2(g)/5 + 23\alpha_3(g))$ and

$$\chi_2(g) = (45\alpha_0(g) - \alpha_2(g) + 5\alpha_3(g))/70.$$

Simirlaly,
$\chi_3(g) = 1/322(69\alpha_0(g) - 23\alpha_1(g)/2 + 23\alpha_2(g)/5 - 23\alpha_3(g)/2)$.
By substitution $\alpha_1(g) + \alpha_3(g) = 322 - \alpha_0(g) - \alpha_2(g)$, we have

$$\chi_3(g) = (5\alpha_0(g) + \alpha_2(g))/20 - 23/2.$$

Theorem 7 ▸ A. Gavrilyuk, A. Makhnev [15]

Let Γ be a distance-regular graph with intersection array $\{60, 45, 8; 1, 12, 50\}$, g be an element of prime order $p \geq 5$ of $\mathrm{Aut}(\Gamma)$ and $\Omega = \mathrm{Fix}(g)$. Then one of the following holds:

1. $p = 7$ or 23 and Ω is the empty graph;
2. $p = 5$ and either
   (i) $\Omega = \{a, b\}$ and $d(a, b) = 3$, or
   (ii) $|\Omega| = 7$ and $\Gamma_3(a) \cap \Omega$ is 6-clique for some vertex $a \in \Omega$.

Covers of srg

Antipodality

Let Γ be a graph of diameter d. Γ is called antipodal graph, if $\{(u,w) \mid d(u,w) \in \{0,d\}\}$ is equivalence relation on the vertex set of Γ. The graph Γ' with the set of antipodal classes of Γ as the vertex set and u' is adjacent to w' iff there exist $u \in u'$, that is adjacent to some $w \in w'$ in Γ, is called the antipodal quotient (folded graph) of Γ. Then Γ is called antipodal cover of Γ', Γ is called r-cover if $|u'| = r$ for each antipodal class u' of Γ.

Covers of HiS

The Higman-Sims graph is the unique strongly regular graph with parameters (100,22,0,6). The existence of cover with diameter 4 of the Higman-Sims graph is unknown. A.Makhnev, V. Nosov and D. Paduchikh investigated automorphisms of a distance-regular covers of the Higman-Sims graph.

Covers of diameter 5

Let Γ be an antipodal distance-regular graph with diameter d such that folded graph Γ' is the Higman-Sims graph. If $d = 5$ then Γ has the intersection array $\{22, 21, t(r-1), 6, 1; 1, 6, t, 21, 22\}$ and $r = 2, 3$. In the case $r = 3$ we have $t \in \{6, 7, 8\}$ and some eigenvalue of Γ has nonintegral multiplicity. In the case $r = 2$ by Theorem 4.2.11 [10] the graph Γ is bipartite double of the Higman-Sims graph.

Problem 3

Does a distance-regular graph of diameter 4, which is r-cover of the Higman-Sims graph ($r \in \{2, 3, 6\}$), exist?

Let Γ be a distance-regular graph with diameter 4, which is 2-cover of the Higman-Sims graph. Then

$$Q = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 50 & 50\sqrt{22}/22 & 0 & -50\sqrt{22}/22 & -50 \\ 77 & 7 & -3 & 7 & 77 \\ 50 & -50\sqrt{22}/22 & 0 & 50\sqrt{22}/22 & -50 \\ 22 & -8 & 2 & -8 & 22 \end{pmatrix}.$$

So
$\chi_4(g) = 1/200(22\alpha_0(g) - 8\alpha_1(g) + 2\alpha_2(g) - 8\alpha_3(g) + 22\alpha_4(g))$.
By substitution $\alpha_1(g) + \alpha_3(g) = 200 - \alpha_0(g) - \alpha_2(g) - \alpha_4(g)$ we have
$\chi_4(g) = 1/20(3\alpha_0(g) + \alpha_2(g) + 3\alpha_4(g)) - 8$.
By substitution $\alpha_0(g) + \alpha_4(g) = 200 - \alpha_1(g) - \alpha_2(g) - \alpha_3(g)$ we have $\chi_4(g) = 1/20(-3\alpha_1(g) - 2\alpha_2(g) - 3\alpha_3(g)) + 22$.
Similarly
$\chi_1(g) = (\alpha_0(g) - \alpha_4(g))/4 + \sqrt{22}(\alpha_1(g)/22 - \alpha_3(g))/88$.

Theorem 8

▸ A. Makhnev, D. Paduchikh [16]

Let Γ be a distance-regular graph with intersection array $\{22, 21, 3, 1; 1, 3, 21, 22\}$, $G = \mathrm{Aut}(\Gamma)$, g be an element of prime order p of G and $\Omega = \mathrm{Fix}(g)$. Then one of the following holds:

1. $p = 11$, Ω is an antipodal class, $\alpha_2(g) = 154$ and $\alpha_1(g) = \alpha_3(g) = 22$;
2. $p = 7$, Ω is the union of two antipodal classes having degree 1, $\alpha_1(g) = \alpha_3(g) = 14$ and $\alpha_2(g) = 168$;
3. $p = 5$, Ω is an empty graph, $\alpha_4(g) = 0$ and either $\alpha_1(g) = \alpha_3(g) = 0$, or $\alpha_1(g) = \alpha_3(g) = 50$;
4. $p = 3$, Ω is the union of two isolated $K_{5,5}$-subgraphs with deleted matching and $\alpha_2(g) = 180$;
5. $p = 2$, Ω is an empty graph and $\alpha_4(g) = 200$.

Theorem 9

▸ A. Makhnev, D. Paduchikh [16]

Let Γ be a distance-regular graph with intersection array $\{22, 21, 4, 1; 1, 2, 21, 22\}$, $G = \mathrm{Aut}(\Gamma)$, g be an element of prime order p of G and $\Omega = \mathrm{Fix}(g)$. Then one of the following holds:

1. $p = 11$, Ω is an antipodal class, $\alpha_2(g) = 231$ and $\alpha_1(g) = 66$;
2. $p = 7$, Ω is the union of two antipodal classes having degree 1, $\alpha_1(g) = \alpha_3(g)/2 = 14$ and $\alpha_2(g) = 252$;
3. $p = 5$, Ω is an empty graph, $\alpha_4(g) = 0$ and either $\alpha_2(g) = 300$, or $\alpha_1(g) = \alpha_3(g)/2 = 50$ and $\alpha_2(g) = 150$;
4. $p = 3$, Ω is an empty graph, and either $\alpha_1(g) = \alpha_3(g) = 0$, $\alpha_4(g) = 300$ or $\alpha_4(g) = 30$, $\alpha_2(g) = 270$;
5. $p = 2$, Ω is an empty graph, $\alpha_4(g) = 0$, $\alpha_2(g) = 180$ and $\alpha_1(g) = \alpha_3(g)/2 = 40$.

Theorem 10

▸ A. Makhnev, V. Nosov [17]

Let Γ be a distance-regular graph with intersection array $\{22, 21, 5, 1; 1, 1, 21, 22\}$, $G = \mathrm{Aut}(\Gamma)$, g be an element of prime order p of G and $\Omega = \mathrm{Fix}(g)$. Then $\pi(G) \subseteq \{2, 3, 5, 7\}$ and one of the following holds:

1. $p = 7$, Ω — a graph of degree 1, which is the union of two antipodal classes, $5\alpha_1(g) = \alpha_3(g) = 70$ and $\alpha_2(g) = 504$;
2. $p = 5$, either Ω is an empty graph, $\alpha_4(g) = 0$ and $\alpha_1(g) = \alpha_3(g) = 0$, or $5\alpha_1(g) = \alpha_3(g) = 250$, or Ω is pentagon, $5\alpha_1(g) = \alpha_3(g) = 50$, $\alpha_2(g) = 510$ and $\alpha_4(g) = 25$;
3. $p = 3$, Ω is an empty graph, either $\alpha_4(g) = 600$, or $\alpha_4(g) = 60$ and $\alpha_2(g) = 540$;
4. $p = 2$, Ω is an empty graph and $\alpha_4(g) = 600$.

Corollary 4

Let Γ be a distance-regular graph with diameter 4 such that Γ is r-cover ($r = 2, 3, 6$) of the Higman-Sims graph. Then the group $G = \mathrm{Aut}(\Gamma)$ acts intransitively on the set of antipodal classes of Γ.

For the proof of theorems 6.1–6.3 we have very useful

Lemma 5

Let Γ be a Higman-Sims graph, $G = \mathrm{Aut}(\Gamma)$, g an element of prime order p of G and $\Omega = \mathrm{Fix}(g)$. Then the following statements hold:

1. $p = 11$, $|\Omega| = 1$, $\alpha_1(g) = 22$, five $\langle g \rangle$-orbits are regular of degree 4, and four — cocliques;
2. $p = 7$, $|\Omega| = 2$, $\alpha_1(g) = 14$, six $\langle g \rangle$-orbits are 7-gon;
3. $p = 5$, either Ω is pentagon, or Ω is an empty graph;
4. $p = 3$, $\alpha_1(g) = 0$, Ω is $K_{5,5}$-subgraph without matching;
5. $p = 2$, and either

   (i) Ω is an empty graph, $\alpha_1(g) = 40$, or

   (ii) Ω is a 2-coclique extension of Petersen graph, $\alpha_2(g) = 80$ and each vertex of $\Gamma - \Omega$ is adjacent with 4 vertices of Ω, or

   (iii) Ω is 6-coclique (this is μ-subgraph of two α_2-vertices), each α_2-vertex is adjacent with 2 or 6 vertices of Ω, or

   (iv) Ω is an amply regular graph with parameters $(30, 8, 0, 4)$, $\alpha_2(g) = 70$ and each α_2-vertex is adjacent with 6 vertices of Ω.

Covers of srg(81,20,1,6)

Wenbin Guo, A. Makhnev and D. Paduchikh investigated automorphisms of a distance-regular r-covers Γ of the strongly regular graph with parameters (81,20,1,6). It is known that there exists a unique strongly regular graph with this parameters. If $d(\Gamma) = 5$ then Γ has the intersection array $\{20, 18, t(r-1), 6, 1; 1, 6, t, 18, 20\}$ and $c_2 \leq t \leq \min\{b_1, a_2\}/(r-1)$ (parameters a_2, b_1, c_2 correspond to the strongly regular graph), so $r \leq 3$. If $r = 3$, then $t = 6, 7$. If $r = 2$, then $6 \leq t \leq 14$. In any case some eigenvalue of Γ has nonintegral multiplicity.

Covers with diameter 4

In the case $d(\Gamma) = 4$ by Theorem 4.2.11 [10] the graph Γ has feasible intersection array for $r \in \{2, 3, 6\}$. If $r = 3$ then cover exist (it is the graph on the cosets of ternary Goley code).

Let Γ be a distance-regular 2-cover of strongly regular graph with parameters (81,20,1,6). Then

$$Q = \begin{pmatrix} 1 & 1 & 1 & 1 & 1 \\ 36 & 9 & 0 & -9 & -36 \\ 60 & 6 & -3 & -6 & 60 \\ 45 & -9 & 0 & 9 & -45 \\ 20 & -7 & 2 & -7 & 20 \end{pmatrix}.$$

So

$$\chi_1(g) = 1/18(4\alpha_0(g) + \alpha_1(g) - \alpha_3(g) - 4\alpha_4(g)).$$

$$\chi_2(g) = 1/54(20\alpha_0(g) + 3\alpha_1(g) - \alpha_2(g) - 3\alpha_3(g) + 20\alpha_4(g)).$$

$$\chi_3(g) = 1/18(5\alpha_0(g) - \alpha_1(g)/2 + \alpha_3(g) - 5\alpha_4(g)).$$

Theorem 11 ‣ Wenbin Guo, A. Makhnev, D. Paduchikh [18]

Let Γ be a distance-regular graph with intersection array $\{20, 18, 3, 1; 1, 3, 18, 20\}$, $G = \mathrm{Aut}(\Gamma)$, g be an element of prime order p of G and $\Omega = \mathrm{Fix}(g)$. Then $|G|$ divides 18 and one of the following holds:

1. $p = 2$, $|\Omega| = 0$ and $d(u, u^g) = 4$ for every vertex $u \in \Gamma$;
2. $p = 3$, Ω is either an empty graph or the union of two triangles, distance between them in Γ is equal 3 and $\alpha_3(g) = \alpha_4(g) = 0$.

Theorem 12 ▸ Wenbin Guo, Makhnev A., Paduchikh D. [18]

Let Γ be a distance-regular graph with intersection array $\{20, 18, 4, 1; 1, 2, 18, 20\}$, $G = \mathrm{Aut}(\Gamma)$, g be an element of prime order p of G and $\Omega = \mathrm{Fix}(g)$. Then one of the following holds:

1. $p = 5$, Ω is an antipodal class, $\alpha_4(g) = 0$, $\alpha_2(g) = 180$ and $\alpha_1(g) = \alpha_3(g) = 30$;
2. $p = 2$, $\bar{\Omega}$ is m-coclique, $m \in \{1, 3, 9\}$ and Ω is n-coclique, $n \in \{1, 9, 9\}$ respectively;
3. $p = 3$, $\alpha_3(g) = \alpha_4(g) = 0$ and either
   (i) $\alpha_4(g) = 243$,
   (ii) $\alpha_1(g) + \alpha_3(g) = 243$ and $\alpha_1(g)$ is divided by 9,
   (iii) $\alpha_2(g) = 243$,
   (iv) $\alpha_0(g) = 0$, $\alpha_1(g) + \alpha_3(g) = 54$, $\alpha_2(g) = 162$,
   $\alpha_4(g) = 27$ and $\alpha_1(g) \in \{0, 27, 54\}$, or
   (v) $\alpha_0(g) = 9$, $\alpha_1(g) + \alpha_3(g) = 54$, $\alpha_2(g) = 162$,
   $\alpha_4(g) = 18$ and $\alpha_1(g) \in \{0, 27, 54\}$.

Theorem 13 ‣ Wenbin Guo, A. Makhnev A, D. Paduchikh [18]

Let Γ be a distance-regular graph with intersection array $\{20, 18, 5, 1; 1, 1, 18, 20\}$, $G = \mathrm{Aut}(\Gamma)$, g be an element of prime order p of G and $\Omega = \mathrm{Fix}(g)$. Then one of the following holds:

1. $p = 5$, Ω is an antipodal class, $\alpha_1(g) + \alpha_3(g) = 120$, $\alpha_2(g) = 360$;
2. $p = 3$ and either
   (i) $\alpha_4(g) = 486$,
   (ii) $\alpha_1(g) + \alpha_3(g) = 486$ and $\alpha_1(g)$ is divided by 9,
   (iii) $\alpha_2(g) = 486$,
   (iv) $\alpha_0(g) = 0$, $\alpha_1(g) + \alpha_3(g) = 108$, $\alpha_2(g) = 324$, $\alpha_4(g) = 54$ and $\alpha_1(g)$ is divided by 9, or
   (v) $\alpha_0(g) = 18$, $\alpha_1(g) + \alpha_3(g) = 108$, $\alpha_2(g) = 324$, $\alpha_4(g) = 36$ and $\alpha_1(g) - 3$ is divided by 9.

Corollary 5

Let Γ be a distance-regular graph with diameter 4 such that Γ is r-cover ($r = 2, 3, 6$) of the strongly regular graph wirh parameters $(81, 20, 1, 6)$. If $\mathrm{Aut}(\Gamma)$ acts transitively on the set vertices of Γ, then $r = 3$.

Terwilliger graphs

Known graphs

Noncomplete connected graph is called Terwilliger graph, if for every two vertices u, w at distance 2 the subgraph $[u] \cap [w]$ is μ-clique for some constant μ. All of known distance-regular Terwilliger graph with $\mu > 1$ are locally Moore graphs. A connected locally pentagon graph is isomorphic to the icosahedron. There are exactly three connected locally Petersen graphs [10, theorem 1.16.5]: $\bar{T}(7)$, Conway-Smith graph (the unique distance-regular graph with intersection array $\{10, 6, 4, 1; 1, 2, 6, 10\}$) and Doro graph (the unique distance-regular graph with intersection array $\{10, 6, 4; 1, 2, 5\}$). The existence of locally Hoffman-Singleton graphs is unknown.

A. Gavrilyuk and A. Makhnev [19] classified connected Terwilliger graphs, containing a some vertex u such that $[u]$ is Petersen graph.

Proposition 6

Let Γ be a Terwilliger graph with vertex u such that $[u]$ is Petersen graph. Then one of the following holds:

1. $\Gamma = u^{\perp}$;
2. Γ is Conway-Smith graph;
3. Γ is Doro graph.

Locally Ho-Si graphs

A. Gavrilyuk and A. Makhnev [20] proved that a distance-regular locally Hoffman-Singleton graph has intersection array $\{50, 42, 1; 1, 2, 50\}$ or $\{50, 42, 9; 1, 2, 42\}$.
A. Gavrilyuk, Wenbin Guo and A. Makhnev investigated automorphisms of distance-regular Terwilliger graphs with intersection arrays $\{50, 42, 1; 1, 2, 50\}$ and $\{50, 42, 9; 1, 2, 42\}$.

Let Γ be a distance-regular graph with intersection array $\{50, 42, 1; 1, 2, 50\}$. Then

$$Q = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 357 & 357/5 & -17/5 & -17 \\ 50 & -1 & -1 & 50 \\ 714 & -357/5 & 17/5 & -34 \end{pmatrix}.$$

So
$\chi_1(g) = 1/1122(357\alpha_0(g) + 357\alpha_1(g)/5 - 17\alpha_2(g)/5 - 17\alpha_3(g))$.
As $\alpha_2(g) = 1122 - \alpha_0(g) - \alpha_1 - \alpha_3(g)$, then

$$\chi_1(g) = (53\alpha_0(g) + 11\alpha_1(g) - 2\alpha_3(g))/165 - 17/5.$$

Further, $\chi_2(g) = 1/1122(50\alpha_0(g) - \alpha_1(g) - \alpha_2(g) + 50\alpha_3(g)$. By substitution $\alpha_1(g) + \alpha_2(g) = 1122 - \alpha_0(g) - \alpha_3(g)$ we have

$$\chi_2(g) = (\alpha_0(g) + \alpha_3(g))/22 - 1.$$

Theorem 14 ‣ A. Gavrilyuk, Wenbin Guo, A. Makhnev [21]

Let Γ be a distance-regular Terwilliger graph with intersection array $\{50, 42, 1; 1, 2, 50\}$, g be an element of prime order p of $\mathrm{Aut}(\Gamma)$ and $\Omega = \mathrm{Fix}(g)$. Then one of the following holds:

1. $p = 3, 11$ or 17 and Ω is an empty graph;
2. $p = 5$, $d(a, b) = 3$ for any two vertices $a, b \in \Omega$ and $|\Omega| \in \{2, 7, 12, 17\}$.

Let Γ be a distance-regular graph with intersection array $\{50, 42, 9; 1, 2, 42\}$. Then

$$Q = \begin{pmatrix} 1 & 1 & 1 & 1 \\ 350 & 77 & -1 & -14 \\ 390 & -39/5 & -39/5 & 182/5 \\ 585 & -351/5 & 117/15 & -117/5 \end{pmatrix}.$$

So $\chi_1(g) = 1/1326(350\alpha_0(g) + 77\alpha_1(g) - \alpha_2(g) - 14\alpha_3(g))$,
$\alpha_2(g) = 1326 - \alpha_0(g) - \alpha_1(g) - \alpha_3(g)$ and
$\chi_1(g) = (27\alpha_0(g) + 6\alpha_1(g) - \alpha_3(g))/102 - 1$.
Further,
$\chi_2(g) = 1/1326(390\alpha_0(g) - 39\alpha_1(g)/5 - 39\alpha_2(g)/5 + 182\alpha_3(g)/5)$.
By substitution $\alpha_1(g) + \alpha_2(g) = 1326 - \alpha_0(g) - \alpha_3(g)$, we have
$\chi_2(g) = (9\alpha_0(g) + \alpha_3(g))/30 - 39/5$.

Theorem 15 ‣ A. Gavrilyuk, Wenbin Guo, A. Makhnev [21]

Let Γ be a distance-regular Terwilliger graph with intersection array $\{50, 42, 9; 1, 2, 42\}$, g be an element of prime order p of $\mathrm{Aut}(\Gamma)$ and $\Omega = \mathrm{Fix}(g)$. Then one of the following holds:

1. $p = 3, 13$ or 17 and Ω is an empty graph;
2. $p = 7$ and Ω is the union of five isolated 2-cliques;
3. $p = 5$ and Ω is the one vertex graph.

Corollary 6

Distance-regular Terwilliger graphs with intersection arrays $\{50, 42, 1; 1, 2, 50\}$ and $\{50, 42, 9; 1, 2, 42\}$ are not vertex-symmetric.

Earlier John van Bon has proved that a locally Hoffman-Singleton graph is not distance-transitive.

The following proposition has independent interest.

Proposition 7

Let Γ be a Terwilliger locally Hoffman-Singleton graph, t be an automorphism of prime order p of Γ, having fixed point and $\Omega = \mathrm{Fix}(t)$. Then one of the following holds:

1. $p = 2$, any component of Ω is isomorphic to Conway-Smith graph or Doro graph, for any vertex $w \in \Gamma - \Omega$ we have $d(w, w^t) \geq 2$ and w is adjacent to 0 or 2 vertices of Ω;
2. $p = 5$, any component of Ω is the one vertex graph or the icosahedron graph and the distance between two vertices from different components is at least 3;
3. $p = 7$, any component of Ω is an edge and the distance between two vertices from different components is at least 3.

References

1. Cameron P., van Lint J. Graphs, Codes and Designs, Cambr. Univ. Press 1991. ▸ ->Proposition 1
2. Cameron P.J. Permutation Groups, Cambridge Univ. Press, 1999. ▸ ->Character
3. Makhnev, Minakova I.M. On automorphisms of graphs with $\lambda = 1, \mu = 2$, Discret. Matem. 2004, v. 16, 95-104. ▸ ->Theorem 1
4. Wilbrink H.A. On the (99,14,1,2) strongly regular graph, In: P.J. de Doelder, J. de Graaf, J.H. van Lint, Papers dedicated to J.J. Seidel, Eindhoven: Technische Hogeschool Eindhoven, 1984, 342-355.
5. Makhnev A.A., Nosov V.V. On automorphisms of strongly regular graphs with $\lambda = 0$, $\mu = 2$, Matem. Sbornik 2004, v. 185, N 3, 47-68. ▸ ->Theorem 2
6. Nakagawa N. On strongly regular graphs with parameters $(k, 0, 2)$ and their antipodal double cover, Hokkaido Math.

References

7. Gavrilyuk A.L., Makhnev A.A. On Krein graphs without triangles, Doklady RAN, 2006, v. 403, N 6, 727-730.
8. Makhnev A.A., Nosov V.V. On automorphisms of strongly regular Krein graphs without triangles, Algebra i Logika 2005, v. 44, N 3, 335-354. ▸ ->Theorem 4
9. Gavrilyuk A.L. On automorphisms of strongly regular graphs with parameters $(784, 116, 0, 20)$, Sibirean Electron Math. Izv. 2008, v. 5, 80-87. ▸ ->Theorem 3
10. Brouwer A.E., Cohen A.M., Neumaier A. Distance-Regular Graphs, Springer-Verlag, 1989.
11. Belousov I.N., Makhnev A.A. On automorphisms of generalized octagon of order (2,4), Doklady RAN 2008, v. 423, N 2, 151-154. ▸ ->Theorem 5
12. Cohen A., Tits J., On generalized hexagons and a near octagon whose lines have three points, Europ. J. Comb. 1985, v. 6, 13–27.

References

13. Belousov I.N., Makhnev A.A. On automorphisms of generalized gexagon of order (3,27), Proc. Inst. Math. and Mechanics UB RAS 2009, v. 15, N 2, 34-44. ▸ ->Theorem 6
14. Gavrilyuk A.L., Makhnev A.A. Amply regular graphs and block-designs, Sibirskii Math. J. 2006, v. 47, N 4, 609-619.
15. Gavrilyuk A.L., Makhnev A.A. On automorphisms of distance-regular graph with the intersection array $\{60, 45, 8; 1, 12, 50\}$, Proc. Inst. Math. and Mechanics UB RAS 2007, v. 13, N 2, 41-53. ▸ ->Theorem 7
16. Makhnev A.A., Paduchikh D.V. Covers of the Higman-Sims graph and their automorphisms, Doklady RAN 2008, v. 422, N 1, 26-29. ▸ ->Theorem 8
17. Makhnev A.A., Nosov V.V. On automorphisms of 6-cover of the Higman-Sims graph, Doklady RAN 2009, v. 425, N 3, 584-600.

References

18. Guo Wenbin, Makhnev A.A., Paduchikh D.V. On automorphisms of covers of strongly regular graph with parameters (81,20,1,6), Mathem. notes 2009, v. 86, N 1, 22-36. ->Theorem 11
19. Gavrilyuk A.L., Makhnev A.A. Terwilliger graphs, in which neighborhood of some vertex is isomorphic to Petersen graph, Doklady RAN 2008, v. 421, N 4, 445-448.
20. Gavrilyuk A.L., Makhnev A.A. Distance-regular graph, in which neighborhoods of vertices are isomorphic to Hoffman-Singleton graph, Doklady RAN 2009, v. 428, N 2, 445-449.
21. Gavrilyuk A.L., Guo Wenbin, Makhnev A.A. On automorphisms of Terwilliger graphs with $\mu = 2$, Algebra i Logika, 2008, v. 47, N 5, 584-600. ->Theorem 14