



Draft Genome Sequence of *Pseudomonas aeruginosa* ATCC 9027, Originally Isolated from an Outer Ear Infection

Ambikesh Jayal,^a Benjamin E. Johns,^b Kevin J. Purdy,^c Sarah E. Maddocks^b

Department of Computing and Information Systems, School of Management, Cardiff Metropolitan University, Cardiff, United Kingdom^a; Department of Biomedical Sciences, Cardiff School of Sport and Health Sciences, Cardiff Metropolitan University, Cardiff, United Kingdom^b; School of Life Sciences, University of Warwick, Coventry, United Kingdom^c

ABSTRACT *Pseudomonas aeruginosa* ATCC 9027 was isolated in 1943 from a case of otitis externa and is commonly employed as a quality control strain for sterility, assessment of antibiofilm agents, and *in vitro* study of wound infection. Here, we present the 6.34-Mb draft genome sequence and highlight some pertinent genes that are associated with virulence.

Pseudomonas aeruginosa ATCC 9027 was originally isolated from a case of otitis externa and is used as a quality control organism (https://www.lgcstandards-atcc.org/Products/Cells_and_Microorganisms/Bacteria.aspx). Despite its origin as a skin pathogen, this strain has been regarded as avirulent and has been utilized to produce rhamnolipid surfactants (1, 2). Nonetheless, *P. aeruginosa* ATCC 9027 has been used extensively to study wound infection *in vitro*, because it is an excellent biofilm former (3). *P. aeruginosa* ATCC 9027 is closely related to PA7, a taxonomic outlier which exhibits numerous multidrug resistance mechanisms (4). PA7 lacks several factors associated with virulence, including a 36-gene cluster encoding the type III secretory system (5).

Genomic DNA derived from a biofilm of *P. aeruginosa* ATCC 9027 was purified using the GenElute bacterial genomic DNA extraction kit (Sigma-Aldrich). The sample library was prepared using an Illumina Nextera XT before DNA sequencing using an Illumina MiSeq system. The genomic data were transferred to and stored in Illumina BaseSpace. The raw reads were processed for quality using Trimmomatic version 0.36 (6) and quality reports generated using FastQC (7). A 6,340,907-bp assembly with 126-fold coverage was constructed using SPAdes version 3.8 (8) and run remotely using Cloud Infrastructure for Microbial Bioinformatics (CLIMB) (9, 10). Quality assessment of assembly was done using QUAST (11). The assembly consists of 53 contigs (500 bp or more) with a mean G+C content of 66.65%. The N_{50} and N_{75} values of the assembly were 290,774 bp and 197,962 bp, respectively. The L_{50} and L_{75} values of the assembly were 8 and 15, respectively. The annotation was performed with the NCBI Prokaryotic Genome Annotation Pipeline (http://www.ncbi.nlm.nih.gov/genome/annotation_prok), which predicted a total of 5,841 protein-coding genes and 58 tRNAs.

The draft genome contained several virulence-associated genes, including *flg*, *pil*, and *gld* (flagellar and fimbrial biosynthesis and motility [12, 13]), *mviM* (putative virulence factor [<http://www.uniprot.org/uniprot/Q1RD89>]), *hudR* (virulence-associated transcriptional regulator [14]), the *pvd* operon (pyoverdine synthesis [15]), *bvgS* (virulence sensor protein [16]), *mvfR-pqsR* (multiple virulence factor regulator [17]), *vrrR* (sigma factor regulator associated with virulence [18]), and *hemO* (heme oxygenase [19]), to name a few. The presence of these genes demonstrates that *P. aeruginosa* ATCC 9027 should be classified as pathogenic, rather than avirulent, and could have implications for its use as a workhorse to produce surfactants.

Received 6 November 2017 Accepted 6 November 2017 Published 30 November 2017

Citation Jayal A, Johns BE, Purdy KJ, Maddocks SE. 2017. Draft genome sequence of *Pseudomonas aeruginosa* ATCC 9027, originally isolated from an outer ear infection. Genome Announc 5:e01397-17. <https://doi.org/10.1128/genomeA.01397-17>.

Copyright © 2017 Jayal et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

Address correspondence to Ambikesh Jayal, ajayal@cardiffmet.ac.uk, or Sarah E. Maddocks, smaddocks@cardiffmet.ac.uk.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [PDLX00000000](#). The version described in this paper is version PDLX01000000. Unassembled reads are also available from the NCBI Sequence Read Archive (SRA) under the accession number SRP120030.

ACKNOWLEDGMENTS

This study was funded by a Cardiff Metropolitan University Research Innovation Award to S.E.M. This study has also used many resources from the Cloud Infrastructure for Microbial Bioinformatics (CLIMB) project, which is funded by UK's Medical Research Council and provides free cloud-based computing resources for analyzing microbial genomics data.

We gratefully acknowledge Lesley Ward and University of Warwick for performing Illumina MiSeq sequencing. We gratefully acknowledge the support provided by the CLIMB project and would like to thank the team managing and funding the CLIMB project.

REFERENCES

- Setoodeh P, Jahanmiri A, Eslamloueyan R, Niazi A, Ayatollahi SS, Aram F, Mahmoodi M, Hortamani A. 2014. Statistical screening of medium components for recombinant production of *Pseudomonas aeruginosa* ATCC 9027 rhamnolipids by non-pathogenic cell factory *Pseudomonas putida* KT2440. *Mol Biotechnol* 56:175–191. <https://doi.org/10.1007/s12033-013-9693-1>.
- Grosso-Becerra MV, González-Valdez A, Granados-Martínez MJ, Morales E, Servín-González L, Méndez JL, Delgado G, Morales-Espinosa R, Ponce-Soto GY, Cocotl-Yañez M, Soberón-Chávez G. 2016. *Pseudomonas aeruginosa* ATCC 9027 in a non-virulent strain suitable for mono-rhamnolipids production. *Appl Microbiol Biotechnol* 100:9995–10004. <https://doi.org/10.1007/s00253-016-7789-9>.
- Camplin AL, Maddocks SE. 2014. Manuka honey treatment of biofilms of *Pseudomonas aeruginosa* results in the emergence of isolates with increased honey resistance. *Ann Clin Microbiol Antimicrob* 13:19. <https://doi.org/10.1186/1476-0711-13-19>.
- Mai-Prochnow A, Bradbury M, Murphy AB. 2015. Draft genome sequence of *Pseudomonas aeruginosa* ATCC 9027 (DSM 1128), an important rhamnolipid surfactant producer and sterility testing strain. *Genome Announc* 3:e01259-15. <https://doi.org/10.1128/genomeA.01259-15>.
- Roy PH, Tetu SG, Larouche A, Elbourne L, Tremblay S, Ren Q, Dodson R, Harkins D, Shay R, Watkins K, Mahamoud Y, Paulsen IT. 2010. Complete genome sequence of the multidrug-resistant taxonomic outlier *Pseudomonas aeruginosa* PA7. *PLoS One* 5:e8842. <https://doi.org/10.1371/journal.pone.0008842>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data. <http://www.bioinformatics.babraham.ac.uk/projects/fastqc>.
- Nurk S, Bankevich A, Antipov D, Gurevich A, Korobeynikov A, Lapidus A, Pribelsky A, Pyshkin A, Sirotkin A, Sirotkin Y, Stepanauskas R, McLean J, Lasken R, Clingenpeel SR, Woyke T, Tesler G, Alekseyev MA, Pevzner PA. 2013. Assembling genomes and mini-metagenomes from highly chimeric reads. In Deng M, Jiang R, Sun F, Zhang X (ed), *Research in Computational Molecular Biology (RECOMB)* 2013. Lecture notes in computer science, vol 7821. Springer, Berlin, Germany.
- Afgan E, Sloggett C, Goonasekera N, Makunin I, Benson D, Crowe M, Gladman S, Kowsar Y, Pheasant M, Horst R, Lonie A, Genomics Virtual Laboratory. 2015. A practical bioinformatics workbench for the cloud. *PLoS One* 10:e0140829. <https://doi.org/10.1371/journal.pone.0140829>.
- Connor TR, Loman NJ, Thompson S, Smith A, Southgate J, Poplawski R, Bull MJ, Richardson E, Ismail M, Thompson SE, Kitchen C, Guest M, Bakke M, Sheppard SK, Pallen MJ. 2016. CLIMB (the Cloud Infrastructure for Microbial Bioinformatics): an online resource for the medical microbiology community. *Microb Genom* 2:e000086. <https://doi.org/10.1099/mgen.0.000086>.
- Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
- Drake D, Montie TC. 1988. Flagella, motility and invasive virulence of *Pseudomonas aeruginosa*. *J Gen Microbiol* 134:43–52. <https://doi.org/10.1099/00221287-134-1-43>.
- Persat A, Inclan YF, Engel JN, Stone HA, Gitai Z. 2015. Type IV pili mechanistically regulate virulence factors in *Pseudomonas aeruginosa*. *Proc Natl Acad Sci U S A* 112:7563–7568. <https://doi.org/10.1073/pnas.1502025112>.
- Kim SH, Park SY, Heo YJ, Cho YH. 2008. *Drosophila melanogaster*-based screening for multihost virulence factors of *Pseudomonas aeruginosa* PA14 and identification of a virulence attenuating factor, HudA. *Infect Immun* 76:4152–4162. <https://doi.org/10.1128/IAI.01637-07>.
- Meyer JM, Neely A, Stintzi A, Georges C, Holder IA. 1996. Pyoverdine is essential for virulence of *Pseudomonas aeruginosa*. *Infect Immun* 64:518–523.
- Linz B, Ivanov YV, Preston A, Brinkac L, Parkhill J, Kim M, Harris SR, Goodfield LL, Fry NK, Gorringer AR, Nicholson TL, Register KB, Losada L, Harvill ET. 2016. Acquisition and loss of virulence-associated factors during genome evolution and speciation in three clades of *Bordetella* species. *BMC Genomics* 17:767. <https://doi.org/10.1186/s12864-016-3112-5>.
- Déziel E, Gopalan S, Tampakaki AP, Lépine F, Padfield KE, Saucier M, Xiao G, Rahme LG. 2005. The contribution of MvfR to *Pseudomonas aeruginosa* pathogenesis and quorum sensing-regulated genes are modulated without affecting *lasRI*, *rhlRI* or the production of *N*-acyl-homoserine-lactones. *Mol Microbiol* 55:998–1014. <https://doi.org/10.1111/j.1365-2958.2004.04448.x>.
- Quesada JM, Otero-Asman JR, Bastiaansen KC, Civantos C, Llamas MA. 2016. The activity of the *Pseudomonas* virulence regulator σ^{VreI} is modulated by the anti- σ factor VreR and the transcription factor PhoB. *Front Microbiol* 7:1159. <https://doi.org/10.3389/fmicb.2016.01159>.
- Zhu W, Hunt DJ, Richardson AR, Stojiljkovic I. 2000. Use of heme compounds as iron sources by pathogenic neisseriae requires the product of the *hemO* gene. *J Bacteriol* 182:439–447. <https://doi.org/10.1128/JB.182.2.439-447.2000>.