

# Assignment 3: Data Exploration

Amanda Braun

## OVERVIEW

This exercise accompanies the lessons in Environmental Data Analytics on Data Exploration.

## Directions

Change “Student Name” on line 3 (above) with your name.

Work through the steps, **creating code and output** that fulfill each instruction.

Be sure to **answer the questions** in this assignment document.

When you have completed the assignment, **Knit** the text and code into a single PDF file.

After Knitting, submit the completed exercise (PDF file) to the dropbox in Sakai. Add your last name into the file name (e.g., “Salk\_A03\_DataExploration.Rmd”) prior to submission.

The completed exercise is due on Tuesday, January 28 at 1:00 pm.

## Set up your R session

Check your working directory, load necessary packages (tidyverse), and upload two datasets: the ECOTOX neonicotinoid dataset (ECOTOX\_Neonicotinoids\_Insects\_raw.csv) and the Niwot Ridge NEON dataset for litter and woody debris (NEON\_NIWO\_Litter\_massdata\_2018-08\_raw.csv). Name these datasets “Neonics” and “Litter”, respectively.

```
getwd()
```

```
## [1] "/Users/amandabraun/Documents/Classes Spring 2020/Data  
Analytics/Environmental_Data_Analytics_2020"
```

```
library(tidyverse)
```

```
Neonics <- read.csv("/Data/Raw/ECOTOX_Neonicotinoids_Insects_raw.csv")
```

```
view(Neonics)
```

```
Litter<-read.csv("/Data/Raw/NEON_NIWO_Litter_massdata_2018-08_raw.csv")
```

*#had to change object name from file name to not include a dash -*

```
view(Litter)
```

## Learn about your system

The neonicotinoid dataset was collected from the Environmental Protection Agency’s ECOTOX Knowledgebase, a database for ecotoxicology research. Neonicotinoids are a class of insecticides used widely in agriculture. The dataset that has been pulled includes all studies published on insects. Why might we be interested in the ecotoxicology of neonicotinoids on insects? Feel free to do a brief internet search if you feel you need more background information.

Answer: We could use the data to determine the efficacy of using neonicotinoids in eradicating different classes of insects. We could look at the amount of neonicotinoids used, and the amount of insects that existed after the application. We could look at the amount of time between application of neonicotinoids and decline in insect population.

The Niwot Ridge litter and woody debris dataset was collected from the National Ecological Observatory Network, which collectively includes 81 aquatic and terrestrial sites across 20 ecoclimatic domains. 32 of these sites sample forest litter and woody debris, and we will focus on the Niwot Ridge long-term ecological research (LTER) station in Colorado. Why might we be interested in studying litter and woody debris that falls to the ground in forests? Feel free to do a brief internet search if you feel you need more background information.

Answer: The amount and condition of woody debris and litter is emerging as a major factor that influences forest ecosystem processes such as carbon cycling, fire behavior, and tree regeneration

How is litter and woody debris sampled as part of the NEON network? Read the NEON\_Litterfall\_UserGuide.pdf document to learn more. List three pieces of salient information about the sampling methods here:

Answer: Trap placement within plots may be either targeted or randomized, depending on the vegetation. In sites with > 50% aerial cover of woody vegetation >2m in height, placement of litter traps is random. In sites

with < 50% cover of woody vegetation that is heterogeneously distributed, trap placement is targeted beneath vegetation. The finest resolution at which spatial data are reported is a single trap. \*Ground traps are sampled once per year. Elevated traps are sampled more frequently. Deciduous forest sites are sampled once every two weeks, and evergreen sites are sampled once every one to two months.

## Obtain basic summaries of your data (Neonics)

What are the dimensions of the dataset?

**dim**(Neonics)

```
## [1] 4623 30
```

5. There are 30 columns of variables and 4623 rows of observed data in the Neonictinoids dataset.

Using the summary function, determine the most common effects that are studied. Why might these effects specifically be of interest?

**summary**(Neonics)

```
## CAS.Number
## Min. : 58842209
## 1st Qu.: 138261413
## Median : 138261413
## Mean : 147651982
## 3rd Qu.: 153719234
## Max. : 210880925
##
##
## Chemical.Name
## (2E)-1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine : 2658
## 3-[(2-Chloro-5-thiazolyl)methyl]tetrahydro-5-methyl-N-nitro-4H-1,3,5-oxadiazin-4-imine: 686
##
## [C(E)]-N-[(2-Chloro-5-thiazolyl)methyl]-N'-methyl-N"-nitroguanidine : 452
## (1E)-N-[(6-Chloro-3-pyridinyl)methyl]-N'-cyano-N-methylethanimidamide : 420
## N"-Methyl-N-nitro-N'-[(tetrahydro-3-furanyl)methyl]guanidine : 218
## [N(Z)]-N-[3-[(6-Chloro-3-pyridinyl)methyl]-2-thiazolidinylidene]cyanamide : 128
## (Other) : 61
##
## Chemical.Grade
## Not reported : 3989
## Technical grade, technical product, technical formulation: 422
## Pestanal grade : 93
## Not coded : 53
## Commercial grade : 27
## Analytical grade : 15
## (Other) : 24
##
## Chemical.Analysis.Method
## Measured : 230
## Not coded : 51
## Not reported : 5
## Unmeasured : 4321
## Unmeasured values (some measured values reported in article): 16
##
##
## Chemical.Purity Species.Scientific.Name
## NR : 2502 Apis mellifera : 667
## 25 : 244 Bombus terrestris : 183
## 50 : 200 Apis mellifera ssp. carnica : 152
## 20 : 189 Bombus impatiens : 140
## 70 : 112 Apis mellifera ssp. ligustica: 113
## 75 : 89 Popillia japonica : 94
## (Other): 1287 (Other) : 3274
##
## Species.Common.Name
## Honey Bee : 667
## Parasitic Wasp : 285
## Buff Tailed Bumblebee: 183
## Carniolan Honey Bee : 152
## Bumble Bee : 140
## Italian Honeybee : 113
## (Other) : 3083
##
## Species.Group
## Insects/Spiders : 3569
## Insects/Spiders; Standard Test Species : 27
## Insects/Spiders; Standard Test Species; U.S. Invasive Species: 667
## Insects/Spiders; U.S. Invasive Species : 360
##
##
##
## Organism.Lifestage Organism.Age Organism.Age.Units
## Not reported: 2271 NR : 3851 Not reported : 3515
## Adult : 1222 2 : 111 Day(s) : 327
## Larva : 437 3 : 105 Instar : 255
## Multiple : 285 < 24 : 81 Hour(s) : 241
```

```

## Multiple : 285 <24 : 81 Hour(s) : 241
## Egg : 128 4 : 81 Hours post-emergence: 99
## Pupa : 69 1 : 59 Year(s) : 64
## (Other) : 211 (Other): 335 (Other) : 122
## Exposure.Type Media.Type
## Environmental, unspecified:1599 No substrate:2934
## Food :1124 Not reported: 663
## Spray : 393 Natural soil: 393
## Topical, general : 254 Litter : 264
## Ground granular : 249 Filter paper: 230
## Hand spray : 210 Not coded : 51
## (Other) : 794 (Other) : 88
## Test.Location Number.of.Doses Conc.1.Type..Author.
## Field artificial : 96 2 :2441 Active ingredient:3161
## Field natural :1663 3 : 499 Formulation :1420
## Field undeterminable: 4 5 : 314 Not coded : 42
## Lab :2860 6 : 230
## 4 : 221
## NR : 217
## (Other): 701
## Conc.1..Author. Conc.1.Units..Author. Effect
## 0.37/ : 208 AI kg/ha : 575 Population :1803
## 10/ : 127 AI mg/L : 298 Mortality :1493
## NR/ : 108 AI lb/acre: 277 Behavior : 360
## NR : 94 AI g/ha : 241 Feeding behavior: 255
## 1 : 82 ng/org : 231 Reproduction : 197
## 1023 : 80 ppm : 180 Development : 136
## (Other):3924 (Other) :2821 (Other) : 379
## Effect.Measurement Endpoint Response.Site
## Abundance :1699 NOEL :1816 Not reported :4349
## Mortality :1294 LOEL :1664 Midgut or midgut gland: 63
## Survival : 133 LC50 : 327 Not coded : 51
## Progeny counts/numbers: 120 LD50 : 274 Whole organism : 41
## Food consumption : 103 NR : 167 Hypopharyngeal gland : 27
## Emergence : 98 NR-LETH: 86 Head : 23
## (Other) :1176 (Other): 289 (Other) : 69
## Observed.Duration..Days. Observed.Duration.Units..Days.
## 1 : 713 Day(s) :4394
## 2 : 383 Emergence : 70
## NR : 355 Growing season : 48
## 7 : 207 Day(s) post-hatch : 20
## 3 : 183 Day(s) post-emergence: 17
## 0.0417 : 133 Tiller stage : 15
## (Other):2649 (Other) : 59
## Author
## Peck,D.C. : 208
## Frank,S.D. : 100
## El Hassani,A.K., M. Dacher, V. Gary, M. Lambin, M. Gauthier, and C. Armengaud: 96
## Williamson,S.M., S.J. Willis, and G.A. Wright : 93
## Laurino,D., A. Manino, A. Patetta, and M. Porporato : 88
## Scholer,J., and V. Krischik : 82
## (Other) :3956
## Reference.Number
## Min. : 344
## 1st Qu.:108459
## Median :165559
## Mean :142189
## 3rd Qu.:168998
## Max. :180410
##
## Title
## Long-Term Effects of Imidacloprid on the Abundance of Surface- and Soil-Active Nontarget
Fauna in Turf : 200
## Reduced Risk Insecticides to Control Scale Insects and Protect Natural Enemies in the
Production and Maintenance of Urban Landscape Plants: 100
## Effects of Sublethal Doses of Acetamiprid and Thiamethoxam on the Behavior of the Honeybee
(Apis mellifera) : 96
## Exposure to Neonicotinoids Influences the Motor Function of Adult Worker Honeybees
: 93
## Toxicity of Neonicotinoid Insecticides on Different Honey Bee Genotypes
: 88
## Chronic Exposure of Imidacloprid and Clothianidin Reduce Queen Survival, Foraging, and
Nectar Storing in Colonies of Bombus impatiens : 82
## (Other) :3964
## Source Publication.Year
## Agric. For. Entomol.11(4): 405-419 : 200 Min. :1982

```

```

## Environ. Entomol.41(2): 377-386 : 100 1st Qu.:2005
## Arch. Environ. Contam. Toxicol.54(4): 653-661: 96 Median :2010
## Ecotoxicology23:1409-1418 : 93 Mean :2008
## Bull. Insectol.66(1): 119-126 : 88 3rd Qu.:2013
## PLoS One9(3): 14 p. : 82 Max. :2019
## (Other) :3964
## Summary.of.Additional.Parameters

## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Active ingredient NR/ - NR/ AI kg/ha | Duration (Days):\xca NR - NR NR | Conc 2 (Author):\xca
NR (NR - NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR : 389
## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Active ingredient NR - NR AI lb/acre | Duration (Days):\xca NR - NR NR | Conc 2 (Author):\xca
NR (NR - NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR : 138
## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Active ingredient NR - NR AI kg/ha | Duration (Days):\xca NR - NR NR | Conc 2 (Author):\xca
NR (NR - NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR : 136
## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Active ingredient NR/ - NR/ AI lb/acre | Duration (Days):\xca NR - NR NR | Conc 2 (Author):
\xca NR (NR - NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR: 124
## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Active ingredient NR - NR AI ng/org | Duration (Days):\xca NR - NR NR | Conc 2 (Author):\xca
NR (NR - NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR : 94
## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Formulation NR - NR ml/ha | Duration (Days):\xca NR - NR NR | Conc 2 (Author):\xca NR (NR -
NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR : 80
## (Other)

```

:3662

Answer: The effects that are studied are abundance, mortality, survival, food consumption, progeny count and emergence. These effects might be of interest because we can determine the number of insects that Neonictinoids kill, the percentage of an insect population that is killed or effected by the Neonictinoid and how Neonictinoids affect reproduction.

Using the summary function, determine the six most commonly studied species in the dataset (common name). What do these species have in common, and why might they be of interest over other insects? Feel free to do a brief internet search for more information if needed.

[summary](#)(Neonics)

```

## CAS.Number
## Min. : 58842209
## 1st Qu.:138261413
## Median :138261413
## Mean :147651982
## 3rd Qu.:153719234
## Max. :210880925
##
##
## Chemical.Name
## (2E)-1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine :2658
## 3-[(2-Chloro-5-thiazolyl)methyl]tetrahydro-5-methyl-N-nitro-4H-1,3,5-oxadiazin-4-imine: 686

## [C(E)]-N-[(2-Chloro-5-thiazolyl)methyl]-N'-methyl-N"-nitroguanidine : 452
## (1E)-N-[(6-Chloro-3-pyridinyl)methyl]-N'-cyano-N-methylethanimidamide : 420
## N"-Methyl-N-nitro-N'-[(tetrahydro-3-furanyl)methyl]guanidine : 218
## [N(Z)]-N-[3-[(6-Chloro-3-pyridinyl)methyl]-2-thiazolidinylidene]cyanamide : 128
## (Other) : 61
##
## Chemical.Grade
## Not reported :3989
## Technical grade, technical product, technical formulation: 422
## Pestanal grade : 93
## Not coded : 53
## Commercial grade : 27
## Analytical grade : 15
## (Other) : 24
##
## Chemical.Analysis.Method
## Measured : 230
## Not coded : 51
## Not reported : 5
## Unmeasured :4321
## Unmeasured values (some measured values reported in article): 16
##
##
## Chemical.Purity Species.Scientific.Name
## NR :2502 Apis mellifera : 667
## 25 : 244 Bombus terrestris : 183
## 50 : 200 Apis mellifera ssp. carnica : 152
## 20 : 180 Bombus impatiens : 140

```

```

## 20 : 189 Bombus impatiens : 140
## 70 : 112 Apis mellifera ssp. ligustica: 113
## 75 : 89 Popillia japonica : 94
## (Other):1287 (Other) :3274
## Species.Common.Name
## Honey Bee : 667
## Parasitic Wasp : 285
## Buff Tailed Bumblebee: 183
## Carniolan Honey Bee : 152
## Bumble Bee : 140
## Italian Honeybee : 113
## (Other) :3083
## Species.Group
## Insects/Spiders :3569
## Insects/Spiders; Standard Test Species : 27
## Insects/Spiders; Standard Test Species; U.S. Invasive Species: 667
## Insects/Spiders; U.S. Invasive Species : 360
##
##
## Organism.Lifestage Organism.Age Organism.Age.Units
## Not reported:2271 NR :3851 Not reported :3515
## Adult :1222 2 : 111 Day(s) : 327
## Larva : 437 3 : 105 Instar : 255
## Multiple : 285 <24 : 81 Hour(s) : 241
## Egg : 128 4 : 81 Hours post-emergence: 99
## Pupa : 69 1 : 59 Year(s) : 64
## (Other) : 211 (Other): 335 (Other) : 122
## Exposure.Type Media.Type
## Environmental, unspecified:1599 No substrate:2934
## Food :1124 Not reported: 663
## Spray : 393 Natural soil: 393
## Topical, general : 254 Litter : 264
## Ground granular : 249 Filter paper: 230
## Hand spray : 210 Not coded : 51
## (Other) : 794 (Other) : 88
## Test.Location Number.of.Doses Conc.1.Type..Author.
## Field artificial : 96 2 :2441 Active ingredient:3161
## Field natural :1663 3 : 499 Formulation :1420
## Field undeterminable: 4 5 : 314 Not coded : 42
## Lab :2860 6 : 230
## 4 : 221
## NR : 217
## (Other): 701
## Conc.1..Author. Conc.1.Units..Author. Effect
## 0.37/ : 208 AI kg/ha : 575 Population :1803
## 10/ : 127 AI mg/L : 298 Mortality :1493
## NR/ : 108 AI lb/acre: 277 Behavior : 360
## NR : 94 AI g/ha : 241 Feeding behavior: 255
## 1 : 82 ng/org : 231 Reproduction : 197
## 1023 : 80 ppm : 180 Development : 136
## (Other):3924 (Other) :2821 (Other) : 379
## Effect.Measurement Endpoint Response.Site
## Abundance :1699 NOEL :1816 Not reported :4349
## Mortality :1294 LOEL :1664 Midgut or midgut gland: 63
## Survival : 133 LC50 : 327 Not coded : 51
## Progeny counts/numbers: 120 LD50 : 274 Whole organism : 41
## Food consumption : 103 NR : 167 Hypopharyngeal gland : 27
## Emergence : 98 NR-LETH: 86 Head : 23
## (Other) :1176 (Other): 289 (Other) : 69
## Observed.Duration..Days. Observed.Duration.Units..Days.
## 1 : 713 Day(s) :4394
## 2 : 383 Emergence : 70
## NR : 355 Growing season : 48
## 7 : 207 Day(s) post-hatch : 20
## 3 : 183 Day(s) post-emergence: 17
## 0.0417 : 133 Tiller stage : 15
## (Other):2649 (Other) : 59
## Author
## Peck,D.C. : 208
## Frank,S.D. : 100
## El Hassani,A.K., M. Dacher, V. Gary, M. Lambin, M. Gauthier, and C. Armengaud: 96
## Williamson,S.M., S.J. Willis, and G.A. Wright : 93
## Laurino,D., A. Manino, A. Patetta, and M. Porporato : 88
## Scholer,J., and V. Krischik : 82
## (Other) :3956

```

```
## Reference.Number
## Min. : 344
## 1st Qu.:108459
## Median :165559
## Mean :142189
## 3rd Qu.:168998
## Max. :180410
##
##
## Title
## Long-Term Effects of Imidacloprid on the Abundance of Surface- and Soil-Active Nontarget
Fauna in Turf : 200
## Reduced Risk Insecticides to Control Scale Insects and Protect Natural Enemies in the
Production and Maintenance of Urban Landscape Plants: 100
## Effects of Sublethal Doses of Acetamiprid and Thiamethoxam on the Behavior of the Honeybee
(Apis mellifera) : 96
## Exposure to Neonicotinoids Influences the Motor Function of Adult Worker Honeybees
: 93
## Toxicity of Neonicotinoid Insecticides on Different Honey Bee Genotypes
: 88
## Chronic Exposure of Imidacloprid and Clothianidin Reduce Queen Survival, Foraging, and
Nectar Storing in Colonies of Bombus impatiens : 82
## (Other) :3964
## Source Publication.Year
## Agric. For. Entomol.11(4): 405-419 : 200 Min. :1982
## Environ. Entomol.41(2): 377-386 : 100 1st Qu.:2005
## Arch. Environ. Contam. Toxicol.54(4): 653-661: 96 Median :2010
## Ecotoxicology23:1409-1418 : 93 Mean :2008
## Bull. Insectol.66(1): 119-126 : 88 3rd Qu.:2013
## PLoS One9(3): 14 p. : 82 Max. :2019
## (Other) :3964
## Summary.of.Additional.Parameters
```

```
## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Active ingredient NR/ - NR/ AI kg/ha | Duration (Days):\xca NR - NR NR | Conc 2 (Author):\xca
NR (NR - NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR : 389
## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Active ingredient NR - NR AI lb/acre | Duration (Days):\xca NR - NR NR | Conc 2 (Author):\xca
NR (NR - NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR : 138
## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Active ingredient NR - NR AI kg/ha | Duration (Days):\xca NR - NR NR | Conc 2 (Author):\xca
NR (NR - NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR : 136
## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Active ingredient NR/ - NR/ AI lb/acre | Duration (Days):\xca NR - NR NR | Conc 2 (Author):
\xca NR (NR - NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR: 124
## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Active ingredient NR - NR AI ng/org | Duration (Days):\xca NR - NR NR | Conc 2 (Author):\xca
NR (NR - NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR : 94
## Purity:\xca NR - NR | Organism Age:\xca NR - NR Not reported | Conc 1 (Author):\xca
Formulation NR - NR ml/ha | Duration (Days):\xca NR - NR NR | Conc 2 (Author):\xca NR (NR -
NR) NR | Conc 3 (Author):\xca NR (NR - NR) NR : 80
## (Other)
```

:3662

Answer: The six most commonly studied species in the dataset are the Honey Bee, Parasitic Wasp, Buff Tailed Bumblebee, Carniolan Honey Bee, Bumble Bee and Italian Honeybee. They are all part of the bee family and neonicotinoids are believed to harm them.

Concentrations are always a numeric value. What is the class of Conc.1..Author. in the dataset, and why is it not numeric?


```
class(Neonics$Conc.1..Author)
## [1] "factor"
```


Answer:Conc.1..Author concentration class is a factor in this dataset. It is not numeric because some of the results in this column are NR and this wouldn't satisfy a numeric value.

## Explore your data graphically (Neonics)

Using geom\_freqpoly, generate a plot of the number of studies conducted by publication year.


```
ggplot(Neonics) +
  geom_freqpoly(aes(x = Publication.Year), bins=50)
```





Reproduce the same graph but now add a color aesthetic so that different Test.Location are displayed as different colors.

```
ggplot(Neonics) +  
  geom_freqpoly(aes(x = Publication.Year, colour = Test.Location))  
## `stat_bin()` using `bins = 30`. Pick better value with `binwidth`.
```





Interpret this graph. What are the most common test locations, and do they differ over time?

Answer: Field natural and lab are the most common test locations. Between 1990 and 2000 field natural and lab had about the same count. Between 2010 and 2020, lab tests became much more prevalent (6 to 10 times more prevalent) than field natural test locations.


Create a bar graph of Endpoint counts. What are the two most common endpoints, and how are they defined? Consult the ECOTOX\_CodeAppendix for more information.

```
ggplot(Neonics) + (aes(x=Endpoint)) + geom_bar()
```










```
ggplot(Litter) +  
  geom_violin(aes(x = functionalGroup, y = dryMass))
```



Why is the boxplot a more effective visualization option than the violin plot in this case?

Answer: The boxplot is more effective. The violin plot is not showing the median, interquartile spread, or outliers like the boxplot.

What type(s) of litter tend to have the highest biomass at these sites?

Answer: Needles and Twigs/Branches