

Improving BiqMac

Angelika Wiegele

Institut für Mathematik, Alpen-Adria-Universität Klagenfurt, Austria

July 2016

BiqMac...

- solver for Binary quadratic and Max-cut problems

BiqMac...

- solver for Binary quadratic and Max-cut problems
- C implementation of a branch-and-bound method

BiqMac...

- solver for Binary quadratic and Max-cut problems
- C implementation of a branch-and-bound method
- using a relaxation based on semidefinite optimization (SDO)

BiqMac...

- solver for Binary quadratic and Max-cut problems
- C implementation of a branch-and-bound method
- using a relaxation based on semidefinite optimization (SDO)
- solved by combination of bundle and interior point methods

BiqMac...

- solver for Binary quadratic and Max-cut problems
- C implementation of a branch-and-bound method
- using a relaxation based on semidefinite optimization (SDO)
- solved by combination of bundle and interior point methods
- for several problem classes best performing solver

BiqMac...

- solver for Binary quadratic and Max-cut problems
- C implementation of a branch-and-bound method
- using a relaxation based on semidefinite optimization (SDO)
- solved by combination of bundle and interior point methods
- for several problem classes best performing solver
- available via biqmac.aau.at or NEOS-server

Overview

1 Introduction

- BiqMac
- Semidefinite Optimization
- Max Cut and Unconstrained Binary Quadratic Problems

2 SDO Relaxation

- Solving the SDO Relaxation
- Strengthening the SDO Relaxation

3 Improving BiqMac

Semidefinite Optimization

$$\begin{array}{ll} \text{(LO)} & \min \quad c^\top x \\ & \text{s.t.} \quad Ax = b \\ & \quad \quad x \geq 0, x \in \mathbb{R}^n \end{array}$$

Semidefinite Optimization

$$\begin{array}{ll} \text{(LO)} & \min \quad c^\top x \\ & \text{s.t.} \quad Ax = b \\ & \quad \quad x \geq 0, x \in \mathbb{R}^n \end{array}$$

$$\begin{array}{ll} \text{(SDO)} & \min \quad \langle C, X \rangle \\ & \text{s.t.} \quad \mathcal{A}(X) = b \\ & \quad \quad X \succeq 0, X \in \mathcal{S}_n \end{array}$$

Semidefinite Optimization

$$\begin{array}{ll} \text{(LO)} & \min \quad c^\top x \\ & \text{s.t.} \quad Ax = b \\ & \quad \quad x \succeq 0, x \in \mathbb{R}^n \end{array}$$

$$\begin{array}{ll} \text{(SDO)} & \min \quad \langle C, X \rangle \\ & \text{s.t.} \quad \mathcal{A}(X) = b \\ & \quad \quad X \succeq 0, X \in \mathcal{S}_n \end{array}$$

$$\langle C, X \rangle = \text{trace}(CX) = (\text{vec}(C))^\top \text{vec}(X)$$

Semidefinite Optimization

$$\begin{array}{ll} \text{(LO)} & \min \quad c^\top x \\ & \text{s.t.} \quad Ax = b \\ & \quad x \geq 0, x \in \mathbb{R}^n \end{array}$$

$$\begin{array}{ll} \text{(SDO)} & \min \quad \langle C, X \rangle \\ & \text{s.t.} \quad \mathcal{A}(X) = b \\ & \quad X \succeq 0, X \in \mathcal{S}_n \end{array}$$

$$\mathcal{A}(X) = \begin{pmatrix} \langle A_1, X \rangle \\ \langle A_2, X \rangle \\ \vdots \\ \langle A_m, X \rangle \end{pmatrix}, \text{ thus } \begin{pmatrix} \text{vec}(A_1)^\top \\ \text{vec}(A_2)^\top \\ \vdots \\ \text{vec}(A_m)^\top \end{pmatrix} \text{vec}(X) = b$$

Semidefinite Optimization

$$\begin{array}{ll} \text{(LO)} & \min \quad c^\top x \\ & \text{s.t.} \quad Ax = b \\ & \quad \quad x \succeq 0, x \in \mathbb{R}^n \end{array}$$

$$\begin{array}{ll} \text{(SDO)} & \min \quad \langle C, X \rangle \\ & \text{s.t.} \quad \mathcal{A}(X) = b \\ & \quad \quad X \succeq 0, X \in \mathcal{S}_n \end{array}$$

SDO as LO, except “ \succeq ”

$$(X \succeq 0 \Leftrightarrow y^\top X y \geq 0 \quad \forall y \in \mathbb{R}^n)$$

Semidefinite Optimization - Duality

- $S_n^+ = \{X \in \mathcal{S}_n: X \succeq 0\}$ is a **closed convex cone** in $\mathbb{R}^{\binom{n+1}{2}}$.
- **Lemma:** $\mathcal{S}_n^{+*} = S_n^+$,
i.e., the cone of semidefinite matrices is **self-dual**.

$$\begin{array}{ll} \textbf{(SDO)} & \min \quad \langle C, X \rangle \\ & \text{s.t.} \quad \mathcal{A}(X) = b \\ & \quad \quad X \succeq 0 \end{array}$$

$$\begin{array}{ll} \textbf{(DSDO)} & \max \quad b^\top y \\ & \text{s.t.} \quad \mathcal{A}^t(y) + Z = C \\ & \quad \quad y \in \mathbb{R}^m, Z \succeq 0 \end{array}$$

Semidefinite Optimization - Duality

- $S_n^+ = \{X \in \mathcal{S}_n: X \succeq 0\}$ is a **closed convex cone** in $\mathbb{R}^{\binom{n+1}{2}}$.
- **Lemma:** $\mathcal{S}_n^{+*} = S_n^+$,
i.e., the cone of semidefinite matrices is **self-dual**.

$$\begin{array}{ll} \textbf{(SDO)} & \min \quad \langle C, X \rangle \\ & \text{s.t.} \quad \mathcal{A}(X) = b \\ & \quad \quad X \succeq 0 \end{array}$$

$$\begin{array}{ll} \textbf{(DSDO)} & \max \quad b^\top y \\ & \text{s.t.} \quad \mathcal{A}^t(y) + Z = C \\ & \quad \quad y \in \mathbb{R}^m, Z \succeq 0 \end{array}$$

- Slater-constraint qualification guarantees strong duality,
- solvable in polynomial time.

Max-Cut Problem

given: graph $G = (V, E)$, $V(G) = \{1, \dots, n\}$, $|E(G)| = m$, w_{ij}

Max-Cut Problem

given: graph $G = (V, E)$, $V(G) = \{1, \dots, n\}$, $|E(G)| = m$, w_{ij}

find: a partition of the vertex set into two parts such that the sum of the weights on the edges having one endpoint in each set is maximized.

Max-Cut Problem

given: graph $G = (V, E)$, $V(G) = \{1, \dots, n\}$, $|E(G)| = m$, w_{ij}

find: a partition of the vertex set into two parts such that the sum of the weights on the edges having one endpoint in each set is maximized.

i.e., find $x \in \{\pm 1\}^n$ such that $\sum w_{ij}(1 - x_i x_j) = x^t L x$ is maximized.

Max-Cut Problem

given: graph $G = (V, E)$, $V(G) = \{1, \dots, n\}$, $|E(G)| = m$, w_{ij}

find: a partition of the vertex set into two parts such that the sum of the weights on the edges having one endpoint in each set is maximized.

i.e., find $\mathbf{x} \in \{\pm 1\}^n$ such that $\sum w_{ij}(1 - x_i x_j) = \mathbf{x}^t \mathbf{L} \mathbf{x}$ is maximized.

Max-Cut Problem

$$\begin{aligned} \max \quad & x^t L x \\ \text{s.t.} \quad & x \in \{\pm 1\}^n \end{aligned}$$

where $L \in \mathcal{S}_n$ is the Laplacian of the graph.

Max-Cut Problem

$$\begin{aligned} \max \quad & x^t L x \\ \text{s.t.} \quad & x \in \{\pm 1\}^n \end{aligned}$$

where $L \in \mathcal{S}_n$ is the Laplacian of the graph.

Unconstrained Binary Quadratic Problem

given: symmetric matrix Q , vector c

$$\begin{aligned} \min \quad & x^t Q x + c^t x \\ \text{s.t.} \quad & x \in \{0, 1\}^n \end{aligned}$$

Max-Cut Problem

$$\begin{aligned} \max \quad & x^t L x \\ \text{s.t.} \quad & x \in \{\pm 1\}^n \end{aligned}$$

where $L \in \mathcal{S}_n$ is the Laplacian of the graph.

Unconstrained Binary Quadratic Problem

given: symmetric matrix Q , vector c

$$\begin{aligned} \min \quad & x^t Q x + c^t x \\ \text{s.t.} \quad & x \in \{0, 1\}^n \end{aligned}$$

Solving MC and solving UBQP is essentially the same.

SDO Relaxation

Max-Cut Problem

$$\begin{aligned} \max \quad & x^t L x \\ \text{s.t.} \quad & x \in \{\pm 1\}^n \end{aligned}$$

where $L \in \mathcal{S}_n$ is the Laplacian of the graph.

SDO Relaxation

Max-Cut Problem

$$\begin{aligned} \max \quad & x^T L x \\ \text{s.t.} \quad & x \in \{\pm 1\}^n \end{aligned}$$

where $L \in \mathcal{S}_n$ is the Laplacian of the graph.

linearize and simplify:

$$x^T L x = \langle L, x x^T \rangle = \langle L, X \rangle \text{ where } X = x x^T$$

SDO Relaxation

Max-Cut Problem

$$\begin{aligned} \max \quad & x^T L x \\ \text{s.t.} \quad & x \in \{\pm 1\}^n \end{aligned}$$

where $L \in \mathcal{S}_n$ is the Laplacian of the graph.

linearize and simplify:

$$x^T L x = \langle L, x x^T \rangle = \langle L, X \rangle \text{ where } X = x x^T \rightsquigarrow$$

$$\begin{aligned} \max \quad & \langle L, X \rangle \\ \text{s.t.} \quad & \text{diag}(X) = e \\ & \text{rank}(X) = 1 \\ & X \succeq 0 \end{aligned}$$

SDO Relaxation

Max-Cut Problem

$$\begin{aligned} \max \quad & x^T L x \\ \text{s.t.} \quad & x \in \{\pm 1\}^n \end{aligned}$$

where $L \in \mathcal{S}_n$ is the Laplacian of the graph.

linearize and simplify:

$$x^T L x = \langle L, x x^T \rangle = \langle L, X \rangle \text{ where } X = x x^T \rightsquigarrow$$

$$\begin{aligned} \max \quad & \langle L, X \rangle \\ \text{s.t.} \quad & \text{diag}(X) = e \\ & \text{rank}(X) = 1 \\ & X \succeq 0 \end{aligned}$$

SDO Relaxation

Max-Cut Problem

$$\begin{aligned} \max \quad & x^T L x \\ \text{s.t.} \quad & x \in \{\pm 1\}^n \end{aligned}$$

where $L \in \mathcal{S}_n$ is the Laplacian of the graph.

linearize and simplify:

$$x^T L x = \langle L, x x^T \rangle = \langle L, X \rangle \text{ where } X = x x^T \leadsto$$

$$\begin{aligned} \max \quad & \langle L, X \rangle \\ \text{s.t.} \quad & \text{diag}(X) = e \\ & \text{rank}(X) = 1 \\ & X \succeq 0 \end{aligned}$$

SDO Relaxation

$$\begin{aligned} \max \quad & \langle L, X \rangle \\ \text{s.t.} \quad & \text{diag}(X) = e \\ & X \succeq 0 \end{aligned}$$

$$\begin{aligned} \min \quad & e^t y \\ \text{s.t.} \quad & \text{Diag}(y) - Z = L \\ & Z \succeq 0 \end{aligned}$$

SDO Relaxation

$$\max \langle L, X \rangle$$

$$\text{s.t. } \text{diag}(X) = e$$

$$X \succeq 0$$

$$\min e^t y$$

$$\text{s.t. } \text{Diag}(y) - Z = L$$

$$Z \succeq 0$$

solving the SDO relaxation: primal-dual interior-point method

SDO Relaxation

$$\max \langle L, X \rangle$$

$$\text{s.t. } \text{diag}(X) = e$$

$$X \succeq 0$$

$$\min e^t y$$

$$\text{s.t. } \text{Diag}(y) - Z = L$$

$$Z \succeq 0$$

solving the SDO relaxation: primal-dual interior-point method

- based on Newton's method

SDO Relaxation

$$\max \langle L, X \rangle$$

$$\text{s.t. } \text{diag}(X) = e$$

$$X \succeq 0$$

$$\min e^t y$$

$$\text{s.t. } \text{Diag}(y) - Z = L$$

$$Z \succeq 0$$

solving the SDO relaxation: primal-dual interior-point method

- based on Newton's method
- let the constraints be given as $\mathcal{A}(X) = (\cdot)$ with

$$\mathcal{A}(X) = \begin{pmatrix} \langle A_1, X \rangle \\ \langle A_2, X \rangle \\ \vdots \\ \langle A_m, X \rangle \end{pmatrix}, \text{ in each iteration solve } \boxed{M \Delta y = rhs}$$

where

$$m_{ij} = \text{trace}(X A_i Z^{-1} A_j)$$

SDO Relaxation

$$\max \langle L, X \rangle$$

$$\text{s.t. } \text{diag}(X) = e$$

$$X \succeq 0$$

$$\min e^t y$$

$$\text{s.t. } \text{Diag}(y) - Z = L$$

$$Z \succeq 0$$

solving the SDO relaxation: primal-dual interior-point method

- based on Newton's method
- let the constraints be given as $\mathcal{A}(X) = (\cdot)$ with

$$\mathcal{A}(X) = \begin{pmatrix} \langle A_1, X \rangle \\ \langle A_2, X \rangle \\ \vdots \\ \langle A_m, X \rangle \end{pmatrix}, \text{ in each iteration solve } M \Delta y = rhs$$

where

$$m_{ij} = \text{trace}(X A_i Z^{-1} A_j)$$

- forming the final system matrix: $O(mn^3 + m^2 n^2)$

SDO Relaxation of k -QKP

for max-cut we have the constraints

$$\text{diag}(X) = \begin{pmatrix} \left\langle \begin{pmatrix} 1 & 0 & \dots \\ 0 & 0 & \dots \\ \vdots & \vdots & \ddots \end{pmatrix}, X \right\rangle \\ \left\langle \begin{pmatrix} 0 & 0 & \dots \\ 0 & 1 & \dots \\ \vdots & \vdots & \ddots \end{pmatrix}, X \right\rangle \\ \vdots \\ \left\langle \begin{pmatrix} \vdots & \vdots & \vdots \\ \dots & 0 & 0 \\ \dots & 0 & 1 \end{pmatrix}, X \right\rangle \end{pmatrix} = e$$

SDO Relaxation of k -QKP

for max-cut we have the constraints

$$\text{diag}(X) = \begin{pmatrix} \left\langle \begin{pmatrix} 1 & 0 & \dots \\ 0 & 0 & \dots \\ \vdots & \vdots & \ddots \end{pmatrix}, X \right\rangle \\ \left\langle \begin{pmatrix} 0 & 0 & \dots \\ 0 & 1 & \dots \\ \vdots & \vdots & \ddots \end{pmatrix}, X \right\rangle \\ \vdots \\ \left\langle \begin{pmatrix} \vdots & \vdots & \vdots \\ \dots & 0 & 0 \\ \dots & 0 & 1 \end{pmatrix}, X \right\rangle \end{pmatrix} = e$$

this yields the system

$$(Z^{-1} \circ X) \Delta y = rhs$$

SDO Relaxation of k -QKP

for max-cut we have the constraints

$$\text{diag}(X) = \begin{pmatrix} \left\langle \begin{pmatrix} 1 & 0 & \dots \\ 0 & 0 & \dots \\ \vdots & \vdots & \ddots \end{pmatrix}, X \right\rangle \\ \left\langle \begin{pmatrix} 0 & 0 & \dots \\ 0 & 1 & \dots \\ \vdots & \vdots & \ddots \end{pmatrix}, X \right\rangle \\ \vdots \\ \left\langle \begin{pmatrix} \vdots & \vdots & \vdots \\ \dots & 0 & 0 \\ \dots & 0 & 1 \end{pmatrix}, X \right\rangle \end{pmatrix} = e$$

this yields the system

$$(Z^{-1} \circ X) \Delta y = rhs$$

\Rightarrow cheap construction of system matrix.

SDO Relaxation of k -QKP

Hence,

$$\begin{aligned} \max \quad & \langle L, X \rangle \\ \text{s.t.} \quad & \text{diag}(X) = e \\ & X \succeq 0 \end{aligned}$$

can be solved efficiently by interior-point methods.

SDO Relaxation of k -QKP

Hence,

$$\begin{aligned} \max \quad & \langle L, X \rangle \\ \text{s.t.} \quad & \text{diag}(X) = e \\ & X \succeq 0 \end{aligned}$$

can be solved efficiently by interior-point methods.

SDO relaxation + primal heuristic Goemans-Williamson rounding

SDO Relaxation of k -QKP

Hence,

$$\begin{aligned} \max \quad & \langle L, X \rangle \\ \text{s.t.} \quad & \text{diag}(X) = e \\ & X \succeq 0 \end{aligned}$$

can be solved efficiently by interior-point methods.

SDO relaxation + primal heuristic Goemans-Williamson rounding

SDO Relaxation of k -QKP

Hence,

$$\begin{aligned} \max \quad & \langle L, X \rangle \\ \text{s.t.} \quad & \text{diag}(X) = e \\ & X \succeq 0 \end{aligned}$$

can be solved efficiently by interior-point methods.

SDO relaxation + primal heuristic Goemans-Williamson rounding
 \leadsto **branch-and-bound algorithm**

SDO Relaxation of k -QKP

Hence,

$$\begin{aligned} \max \quad & \langle L, X \rangle \\ \text{s.t.} \quad & \text{diag}(X) = e \\ & X \succeq 0 \end{aligned}$$

can be solved efficiently by interior-point methods.

SDO relaxation + primal heuristic Goemans-Williamson rounding
 \leadsto branch-and-bound algorithm

next step: stronger bounds by including triangle inequalities

strengthen SDO relaxation

observation: $x_i, x_j, x_k \in \{\pm 1\}$ and $\mathbf{X} = \mathbf{x}\mathbf{x}^t$ then the triangle inequalities hold, i.e.,

$$-x_{ij} - x_{ik} - x_{jk} \leq 1$$

$$-x_{ij} + x_{ik} + x_{jk} \leq 1$$

$$x_{ij} - x_{ik} + x_{jk} \leq 1$$

$$x_{ij} + x_{ik} - x_{jk} \leq 1$$

$$1 \leq i < j < k \leq n$$

... $\mathbf{X} \in MET$

- matrix \mathbf{X} dimension n
- $4\binom{n}{3}$ additional constraints

Strengthen SDO Relaxation

$$\begin{aligned} \max \quad & \langle L, X \rangle \\ \text{s.t.} \quad & \text{diag}(X) = e \\ & X \in MET \\ & X \succeq 0 \end{aligned}$$

- matrix X dimension n
- number of linear constraints: $n + 4\binom{n}{3}$

Strengthen SDO Relaxation

$$\begin{aligned} \max \quad & \langle L, X \rangle \\ \text{s.t.} \quad & \text{diag}(X) = e \\ & X \in MET \\ & X \succeq 0 \end{aligned}$$

- matrix X dimension n
- number of linear constraints: $n + 4\binom{n}{3}$
- serious computational effort
- use [bundle method](#)

Bundle Method

We would like to compute

$$z^* = \max \{ \langle L, X \rangle : \text{diag}(X) = e, \mathcal{M}(X) \leq b, X \succeq 0 \}$$

Optimizing over $\text{diag}(X) = e, X \succeq 0$ without $\mathcal{M}(X) \leq b$ is “easy”, but inclusion of $\mathcal{M}(X) \leq b$ makes SDO difficult.

Bundle Method

We would like to compute

$$z^* = \max \{ \langle L, X \rangle : \text{diag}(X) = e, \mathcal{M}(X) \leq b, X \succeq 0 \}$$

Optimizing over $\text{diag}(X) = e, X \succeq 0$ without $\mathcal{M}(X) \leq b$ is “easy”, but inclusion of $\mathcal{M}(X) \leq b$ makes SDO difficult.

partial Lagrangian dual ($\gamma \geq 0$ dual to $\mathcal{M}(X) \leq b$):

$$\mathcal{L}(X; \gamma) = \langle L, X \rangle + \gamma^t(b - \mathcal{M}(X))$$

dual functional:

$$f(\gamma) = \max_{\text{diag}(X)=e, X \succeq 0} \mathcal{L}(X; \gamma)$$

Bundle Method

We would like to compute

$$z^* = \max \{ \langle L, X \rangle : \text{diag}(X) = e, \mathcal{M}(X) \leq b, X \succeq 0 \}$$

Optimizing over $\text{diag}(X) = e, X \succeq 0$ without $\mathcal{M}(X) \leq b$ is “easy”, but inclusion of $\mathcal{M}(X) \leq b$ makes SDO difficult.

partial Lagrangian dual ($\gamma \geq 0$ dual to $\mathcal{M}(X) \leq b$):

$$\mathcal{L}(X; \gamma) = \langle L, X \rangle + \gamma^t(b - \mathcal{M}(X))$$

dual functional:

$$f(\gamma) = \max_{\text{diag}(X)=e, X \succeq 0} \mathcal{L}(X; \gamma)$$

and thus

$$z^* = \min_{\gamma \in \mathbb{R}_{\geq 0}^m} f(\gamma)$$

Bundle Method

since

$$z^* = \min_{\gamma \in \mathbb{R}_{\geq 0}^m} f(\gamma)$$

Bundle Method

since

$$z^* = \min_{\gamma \in \mathbb{R}_{\geq 0}^m} f(\gamma) \leq f(\tilde{\gamma}) \quad \forall \tilde{\gamma} \in \mathbb{R}_{\geq 0}^m$$

any $\tilde{\gamma} \in \mathbb{R}_{\geq 0}^m$ provides upper bound on z^*

Bundle Method

since

$$z^* = \min_{\gamma \in \mathbb{R}_{\geq 0}^m} f(\gamma) \leq f(\tilde{\gamma}) \quad \forall \tilde{\gamma} \in \mathbb{R}_{\geq 0}^m$$

any $\tilde{\gamma} \in \mathbb{R}_{\geq 0}^m$ provides upper bound on z^*

→ try to find tight upper bound (i.e., approximate minimizer of $f(\gamma)$) by using **bundle methods**.

Bundle Method

since

$$z^* = \min_{\gamma \in \mathbb{R}_{\geq 0}^m} f(\gamma) \leq f(\tilde{\gamma}) \quad \forall \tilde{\gamma} \in \mathbb{R}_{\geq 0}^m$$

any $\tilde{\gamma} \in \mathbb{R}_{\geq 0}^m$ provides upper bound on z^*

→ try to find tight upper bound (i.e., approximate minimizer of $f(\gamma)$) by using **bundle methods**.

iterative procedure,

Bundle Method

since

$$z^* = \min_{\gamma \in \mathbb{R}_{\geq 0}^m} f(\gamma) \leq f(\tilde{\gamma}) \quad \forall \tilde{\gamma} \in \mathbb{R}_{\geq 0}^m$$

any $\tilde{\gamma} \in \mathbb{R}_{\geq 0}^m$ provides upper bound on z^*

→ try to find tight upper bound (i.e., approximate minimizer of $f(\gamma)$) by using **bundle methods**.

iterative procedure, main computational effort in each iteration:

- evaluate $f(\hat{\gamma})$ to yield new \hat{x} and a subgradient \hat{g} ,

Bundle Method

since

$$z^* = \min_{\gamma \in \mathbb{R}_{\geq 0}^m} f(\gamma) \leq f(\tilde{\gamma}) \quad \forall \tilde{\gamma} \in \mathbb{R}_{\geq 0}^m$$

any $\tilde{\gamma} \in \mathbb{R}_{\geq 0}^m$ provides upper bound on z^*

→ try to find tight upper bound (i.e., approximate minimizer of $f(\gamma)$) by using **bundle methods**.

iterative procedure, main computational effort in each iteration:

- evaluate $f(\hat{\gamma})$ to yield new \hat{x} and a subgradient \hat{g} , i.e., solve

$$\max_{\text{diag}(\hat{x})=e, \hat{x} \succeq 0} \langle L - \mathcal{M}^t(\hat{\gamma}), \hat{x} \rangle$$

Experience using bundle method

- in combination with interior point methods is a good tool to approximate SDOs with a huge number of constraints

Experience using bundle method

- in combination with interior point methods is a good tool to approximate SDOs with a huge number of constraints
- the number of function evaluations to reach good approximations is surprisingly small

Experience using bundle method

- in combination with interior point methods is a good tool to approximate SDOs with a huge number of constraints
- the number of function evaluations to reach good approximations is surprisingly small
- getting to the “real” optimum is hard

Overview

1 Introduction

- BiqMac
- Semidefinite Optimization
- Max Cut and Unconstrained Binary Quadratic Problems

2 SDO Relaxation

- Solving the SDO Relaxation
- Strengthening the SDO Relaxation

3 Improving BiqMac

Bundle Method

Drawback of interior point methods: warm start not possible!

Bundle Method

Drawback of interior point methods: warm start not possible!

How to improve Biq Mac?

- replace interior point by method that can make use of information from previous iteration and previously parent nodes in branch-and-bound tree

Drawback of interior point methods: warm start not possible!

How to improve Biq Mac?

- replace interior point by method that can make use of information from previous iteration and previously parent nodes in branch-and-bound tree
- use stronger SDO relaxation

Drawback of interior point methods: warm start not possible!

How to improve Biq Mac?

- replace interior point by method that can make use of information from previous iteration and previously parent nodes in branch-and-bound tree
- use stronger SDO relaxation

Bundle Method

Drawback of interior point methods: warm start not possible!

How to improve Biq Mac?

- replace interior point by method that can make use of information from previous iteration and previously parent nodes in branch-and-bound tree
- use stronger SDO relaxation

stronger relaxation: “exact subgraphs” [Malwina Duda, PhD Thesis, forthcoming]

Bundle Method

Drawback of interior point methods: warm start not possible!

How to improve Biq Mac?

- **replace interior point** by method that can make use of information from previous iteration and previously parent nodes in branch-and-bound tree
- use stronger SDO relaxation

stronger relaxation: “**exact subgraphs**” [Malwina Duda, PhD Thesis, forthcoming]

Augmented Lagrangian and Low Rank

$$\begin{array}{ll}\min & b^t y \\ \text{s.t.} & \mathcal{A}^t y - Z = C \\ & Z \in S_n^+\end{array}$$

Augmented Lagrangian and Low Rank

$$\begin{array}{ll}\min & b^t y \\ \text{s.t.} & \mathcal{A}^t y - Z = C \\ & Z \in S_n^+\end{array}$$

$$Z \in S_n^+ \Rightarrow \exists V \in \mathbb{R}^{(m+1) \times r}: Z = VV^t$$

Augmented Lagrangian and Low Rank

$$\begin{array}{ll}\min & b^t y \\ \text{s.t.} & \mathcal{A}^t y - Z = C \\ & Z \in S_n^+\end{array}$$

$$Z \in S_n^+ \Rightarrow \exists V \in \mathbb{R}^{(m+1) \times r}: Z = VV^t$$

$$\begin{array}{ll}\min & b^t y \\ \text{s.t.} & \mathcal{A}^t y - VV^t = C\end{array}$$

...non-convex problem

Augmented Lagrangian and Low Rank

$$\begin{array}{ll}\min & b^t y \\ \text{s.t.} & \mathcal{A}^t y - Z = C \\ & Z \in S_n^+\end{array}$$

$$Z \in S_n^+ \Rightarrow \exists V \in \mathbb{R}^{(m+1) \times r}: Z = VV^t$$

$$\begin{array}{ll}\min & b^t y \\ \text{s.t.} & \mathcal{A}^t y - VV^t = C\end{array}$$

...non-convex problem

$$\mathcal{L}_\sigma(y, V; Y) = b^t y + \langle Y, C - \mathcal{A}^t y + VV^t \rangle + \frac{\sigma}{2} \|C - \mathcal{A}^t y + VV^t\|^2$$

Augmented Lagrangian

$$\min_{y, V} \mathcal{L}_\sigma(y, V; Y)$$

Augmented Lagrangian

$$\min_{y, V} \mathcal{L}_\sigma(y, V; Y)$$

First-order optimality conditions:

$$\nabla_y \mathcal{L}_\sigma(y, V; Y) = 0$$

$$\nabla_V \mathcal{L}_\sigma(y, V; Y) = 0$$

Augmented Lagrangian

$$\min_{y, V} \mathcal{L}_\sigma(y, V; Y)$$

First-order optimality conditions:

$$\nabla_y \mathcal{L}_\sigma(y, V; Y) = 0$$

$$\nabla_V \mathcal{L}_\sigma(y, V; Y) = 0$$

$$\nabla_y \mathcal{L}_\sigma(y, V; Y) = b - AY - \sigma(A(C - A^t y + VV^t)) = 0$$

$$\Rightarrow y = (AA^t)^{-1} \left[A \left(VV^t + C + \frac{1}{\sigma} Y \right) - \frac{1}{\sigma} b \right]$$

Augmented Lagrangian

$$\min_{y, V} \mathcal{L}_\sigma(y, V; Y)$$

First-order optimality conditions:

$$\nabla_y \mathcal{L}_\sigma(y, V; Y) = 0$$

$$\nabla_V \mathcal{L}_\sigma(y, V; Y) = 0$$

$$\nabla_y \mathcal{L}_\sigma(y, V; Y) = b - AY - \sigma(A(C - \mathcal{A}^t y + VV^t)) = 0$$

$$\implies y = (AA^t)^{-1} \left[A \left(VV^t + C + \frac{1}{\sigma} Y \right) - \frac{1}{\sigma} b \right]$$

→ express \mathcal{L}_σ as a function of V only.

augmented Lagrangian

$$\begin{aligned} \min \quad & f(\mathbf{V}) \\ \text{s.t.} \quad & h(\mathbf{V}) = 0 \end{aligned}$$

$$\mathcal{L}_\sigma(\mathbf{V}; \mathbf{Y}) := f(\mathbf{V}) + \langle \mathbf{Y}, h(\mathbf{V}) \rangle + \frac{\sigma}{2} \|h(\mathbf{V})\|^2$$

f and h are sufficiently smooth functions

augmented Lagrangian

Data: f, h

Output: V, Y

augmented Lagrangian

Data: f, h

Output: V, Y

Initialization: $k = 0$, select $\sigma_k > 0$, $V_k \in \mathbb{R}^{(m+1) \times r}$

augmented Lagrangian

Data: f, h

Output: V, Y

Initialization: $k = 0$, select $\sigma_k > 0$, $V_k \in \mathbb{R}^{(m+1) \times r}$

while $\|h(V_k)\| > \varepsilon$

augmented Lagrangian

Data: f, h

Output: V, Y

Initialization: $k = 0$, select $\sigma_k > 0$, $V_k \in \mathbb{R}^{(m+1) \times r}$

while $\|h(V_k)\| > \varepsilon$

(a) Solve $\min_V \mathcal{L}_{\sigma_k}(V; Y_k)$ approximately, giving V_k

augmented Lagrangian

Data: f, h

Output: V, Y

Initialization: $k = 0$, select $\sigma_k > 0$, $V_k \in \mathbb{R}^{(m+1) \times r}$

while $\|h(V_k)\| > \varepsilon$

(a) Solve $\min_V \mathcal{L}_{\sigma_k}(V; Y_k)$ approximately, giving V_k

(b) Update Y : $Y_{k+1} = Y_k + \sigma_k h(V_k)$

augmented Lagrangian

Data: f, h

Output: V, Y

Initialization: $k = 0$, select $\sigma_k > 0$, $V_k \in \mathbb{R}^{(m+1) \times r}$

while $\|h(V_k)\| > \varepsilon$

(a) Solve $\min_V \mathcal{L}_{\sigma_k}(V; Y_k)$ approximately, giving V_k

(b) Update Y : $Y_{k+1} = Y_k + \sigma_k h(V_k)$

(c) Select $\sigma_{k+1} \geq \sigma_k$

augmented Lagrangian

Data: f, h

Output: V, Y

Initialization: $k = 0$, select $\sigma_k > 0$, $V_k \in \mathbb{R}^{(m+1) \times r}$

while $\|h(V_k)\| > \varepsilon$

(a) Solve $\min_V \mathcal{L}_{\sigma_k}(V; Y_k)$ approximately, giving V_k

(b) Update Y : $Y_{k+1} = Y_k + \sigma_k h(V_k)$

(c) Select $\sigma_{k+1} \geq \sigma_k$

(d) Check the stopping condition and increase k

augmented Lagrangian

Data: f, h

Output: V, Y

Initialization: $k = 0$, select $\sigma_k > 0$, $V_k \in \mathbb{R}^{(m+1) \times r}$

while $\|h(V_k)\| > \varepsilon$

(a) Solve $\min_V \mathcal{L}_{\sigma_k}(V; Y_k)$ approximately, giving V_k

(b) Update Y : $Y_{k+1} = Y_k + \sigma_k h(V_k)$

(c) Select $\sigma_{k+1} \geq \sigma_k$

(d) Check the stopping condition and increase k

How to solve the inner minimization?

→ descent method (e.g., Fletcher-Reeves)

Computational effort: Depends on cost of function evaluation and computation of the gradient.

↪ relatively cheap in our case.

Optimality Conditions

Optimality conditions of the SDO:

$$\mathcal{A}Y = b$$

$$Y \succeq 0$$

$$\mathcal{A}^t y - C = Z$$

$$Z \succeq 0$$

$$ZY = 0$$

Optimality Conditions

Optimality conditions of the SDO:

$$\mathcal{A}Y = b \quad \text{by construction}$$

$$Y \succeq 0$$

$$\mathcal{A}^t y - C = Z$$

$$Z \succeq 0 \quad \text{by construction}$$

$$ZY = 0 \quad \text{by construction}$$

Optimality Conditions

Optimality conditions of the SDO:

$$\begin{array}{lll} \mathcal{A}Y & = & b \quad \text{by construction} \\ Y & \succeq & 0 \quad \text{to be maintained} \\ \mathcal{A}^t y - C & = & Z \quad \text{stopping criterion} \\ Z & \succeq & 0 \quad \text{by construction} \\ ZY & = & 0 \quad \text{by construction} \end{array}$$

Summary

- Exact solution method for max cut and unconstrained binary quadratic problems

Summary

- Exact solution method for max cut and unconstrained binary quadratic problems
- Relaxation using SDO

Summary

- Exact solution method for max cut and unconstrained binary quadratic problems
- Relaxation using SDO
- High quality bounds

Summary

- Exact solution method for max cut and unconstrained binary quadratic problems
- Relaxation using SDO
- High quality bounds
- Improvement 1: stronger SDO relaxation (exact subgraph)

Summary

- Exact solution method for max cut and unconstrained binary quadratic problems
- Relaxation using SDO
- High quality bounds
- Improvement 1: stronger SDO relaxation (exact subgraph)
- Improvement 2: use method that allows warm start

Summary

- Exact solution method for max cut and unconstrained binary quadratic problems
- Relaxation using SDO
- High quality bounds
- Improvement 1: stronger SDO relaxation (exact subgraph)
- Improvement 2: use method that allows warm start
- Todo: efficient implementation

Summary

- Exact solution method for max cut and unconstrained binary quadratic problems
- Relaxation using SDO
- High quality bounds
- Improvement 1: stronger SDO relaxation (exact subgraph)
- Improvement 2: use method that allows warm start
- Todo: efficient implementation

Summary

- Exact solution method for max cut and unconstrained binary quadratic problems
- Relaxation using SDO
- High quality bounds
- Improvement 1: stronger SDO relaxation (exact subgraph)
- Improvement 2: use method that allows warm start
- Todo: efficient implementation

Thank you!