

FERMIONIC MATRICES AND SUPER CAYLEY–HAMILTON ALGEBRAS.

CLAUDIO PROCESI

CONTENTS

1. Introduction	1
1.1. 1 Fermionic matrix	2
2. Grassmann matrices	3
2.1. Superalgebras	3
3. Superalgebras with trace	4
3.4. Free algebras	6
3.6. Strings and encoding map	6
3.13. Trace identities	9
4. Matrices	11
4.1. Generic matrices	11
5. Relations	18
5.2. The Cayley–Hamilton identities	18
5.7. The second fundamental Theorem	20
6. Super Cayley–Hamilton algebras	24
6.3. n -dimensional superrepresentations	25
References	27

1. INTRODUCTION

In a recent preprint:

Fermionic trace relations and supersymmetric indices at finite N ,
the authors Giorgos Eleftheriou, Ziming Ji, Sameer Murthy start in their words *to study invariants of bosonic and fermionic (Grassmann-valued) matrices under the adjoint action of $U(N)$, weighted by the fermion number* [8].

In this paper they formulate two natural conjectures about trace relations, which can be viewed as the content of the *second fundamental theorem of matrix invariants*. We prove both in this paper.

Conjecture 1 is our Theorem 5.5 while Conjecture 2 is our Theorem 5.13.

The proof of these theorems and also of the First Fundamental Theorem 4.3 follows the lines of my 1976 paper [15].

In fact the first fundamental theorem of matrix invariants was already proved by Sibirskii[20], the second fundamental theorem of matrix invariants was independently proved by Razmyslov [19].

Some literature of 3-dimensional invariant theory appears in papers on continuous mechanics, [22], [23], [24], [25], [26],

As an application I define super Cayley–Hamilton algebras and prove the strong embedding theorem Theorem 6.7 for this class of graded algebras, following the lines of my 1987 paper [17].

The main novelty in this paper is an appropriate generalization of the symbolic method and a very careful bookkeeping of the signs.

1.1. 1 Fermionic matrix. This case has been completely described in [16] and [3].

The problem is to describe the invariant algebra $(\bigwedge M_n(\mathbb{C})^* \otimes M_n(\mathbb{C}))^G$ under conjugation by $G = GL(n, \mathbb{C})$.

The matrix algebra $M_n(\mathbb{C})$ has as basis the matrix units $e_i \otimes e^j$ and its dual $M_n(\mathbb{C})^*$ the duals $\xi_{i,j} := e^i \otimes e_j$ so that defining $\xi := \sum_{i,j} \xi_{i,j} e_i \otimes e^j$ we have a conjugation invariant *generic matrix* with entries $\xi_{i,j}$ in $\bigwedge M_n(\mathbb{C})^*$, we write $\xi = (\xi_{i,j})$.

One has $\xi^{2n} = 0$ see [16], which is an interpretation of the classical Theorem of Amitsur–Levitzi on the vanishing of the standard identity in $2n$ variables [2].

The computation of $(\bigwedge M_n(\mathbb{C})^*)^G$ is a classical Theorem of Dynkin, see [7]. One has $tr(\xi^{2k}) = 0, \forall k$.

Then $(\bigwedge M_n(\mathbb{C})^*)^G = \bigwedge(t_1, \dots, t_n)$, $t_i = tr(\xi^{2n-i})$ is the exterior algebra in the n Grassmann variables t_i of degree $2i - 1$.

Finally $(\bigwedge M_n(\mathbb{C})^* \otimes M_n(\mathbb{C}))^G$ has been described in a paper by Brešar; Procesi; Špenko [3].

It is a free module with basis the $2n$ elements ξ^i , $i = 0, \dots, 2n - 1$ over $\bigwedge(t_1, \dots, t_{n-1})$ and, by Lemma 4.3 of [3]

$$0 = n\xi^{2n-1} - \sum_{i=0}^{n-1} \xi^{2i} \wedge t_{n-i-1}, \quad \text{or} \quad t_n = -n\xi^{2n-1} + \sum_{i=0}^{n-2} \xi^{2i} \wedge t_{n-i-1}.$$

In particular the graded dimension polynomial is

$$\prod_{i=0}^{n-1} (1 + q^{2^{i-1}}) \left(\sum_{i=0}^{2n-1} (1 + q^i) \right).$$

For several variables the results are not as precise, since they are not so precise even in the classical case. Nevertheless most of the classical results extend to this more general case.

The previous Theorem has then a vast generalization to simple Lie algebras by De Concini, Papi, Procesi [4].

2. GRASSMANN MATRICES

2.1. Superalgebras. We use the language of superspaces and $\mathbb{Z}/(2)$ graded algebras or superalgebras.

Definition 2.2.

- (1) A *superspace* is a $\mathbb{Z}/(2)$ graded space $W = W_0 \oplus W_1$.
- (2) A *superalgebra* is an associative algebra A with a $\mathbb{Z}/(2)$ grading $A = A_0 \oplus A_1$ so that $A_j A_i \subset A_{i+j}$, $i + j \bmod 2$.
- (3) We denote by $d(a) \in \{0, 1\}$ the degree of an homogeneous element of A .
- (4) The supercommutator of two homogeneous elements a, b is

$$[a, b] := ab - (-1)^{d(a)d(b)} ba.$$

- (5) An algebra is supercommutative if the supercommutator of two homogeneous elements is always zero,
- (6) The supercenter of A is the set of elements of A which super commute with all elements of A . It is a supercommutative subalgebra.

One has the notion of *graded tensor product* $A \hat{\otimes} B$ of superspaces, when they are both superalgebras this is also a superalgebra which as vector space is $A \otimes B$ but for multiplication of graded elements one defines:

$$a_1 \otimes b_1 \cdot a_2 \otimes b_2 = (-1)^{d(a_2)d(b_1)} a_1 a_2 \otimes b_1 b_2.$$

If $A = A_0$ or $B = B_0$ the multiplication in the graded tensor product equals that of usual tensor product.

Proposition 2.3. *The graded tensor product $A \hat{\otimes} B$ of two supercommutative superalgebras is supercommutative.*

Given a vector space V we consider the Grassmann algebra $\bigwedge V$ as $\mathbb{Z}/(2)$ graded and then one has an isomorphism as $\mathbb{Z}/(2)$ graded algebras

$$(1) \quad \bigwedge (V_1 \oplus V_2) = \bigwedge V_1 \hat{\otimes} \bigwedge V_2.$$

In grater generality given a supervector space $V_0 \oplus V_1$ one may define

$$(2) \quad \mathbb{K}(V) := S(V_0) \otimes \bigwedge V_1$$

where for a vector space W the symbol $S(W)$ is the symmetric algebra (in practice the polynomials in the elements of a basis of W).

As in Formula (1) one has for a direct sum $U \oplus V$ of graded vector spaces:

$$(3) \quad \mathbb{K}(U \oplus V) = \mathbb{K}(U) \hat{\otimes} \mathbb{K}(V).$$

One has the simple

Theorem 2.4. $\mathbb{K}(V)$ is a free algebra in the category of supercommutative superalgebras.

This means that, given a supercommutative superalgebra A and a graded linear map $V \rightarrow A$ this extends to a unique homomorphism of $\mathbb{K}(V)$ to A .

If A is $\mathbb{Z}/(2)$ graded we have that the space $M_n(A) = M_n(\mathbb{C}) \otimes A$ of $n \times n$ matrices over A is also $\mathbb{Z}/(2)$ graded considering $M_n(\mathbb{C})$ as totally in degree 0. The supercenter of $M_n(A)$ is $A = 1 \otimes A$.

Definition 2.5. If A is supercommutative, an element of degree 0, resp. 1, of $M_n(A)$ is called a *Bosonic*, resp. *Fermionic matrix*.

An element of degree 1, of $M_n(\mathbb{K}(V))$ is called a *Grassmann matrix*.

3. SUPERALGEBRAS WITH TRACE

In order to explain the SFT for Grassmann matrices we develop a language which is a natural evolution of the one developed in the previous paragraphs, at least in characteristic 0, the language of *superalgebras with trace* and their identities (cf. [1]).

Definition 3.1. An associative superalgebra with trace, also called SAT over a field F is an associative F -superalgebra R with a 1-ary operation

$$t : R \rightarrow R$$

which is assumed to satisfy the following axioms:

- (1) t is F -linear and compatible with the $\mathbb{Z}/(2)$ grading.
- (2) $t(a)b = (-1)^{d(a)d(b)}b \cdot t(a)$, $\forall a, b \in R$ homogeneous.
- (3) $t(ab) = (-1)^{d(a)d(b)}t(ba)$, $\forall a, b \in R$.
- (4) $t(t(a)b) = t(a)t(b)$, $\forall a, b \in R$.

This operation is called a *formal trace*.

We denote $t(R) := \{t(a), a \in R\}$ the image of t .

Remark 3.2. We have the following implications:

- Axiom 1) implies that $t(R)$ is an F –submodule.
- Axiom 2) implies that $t(R)$ is contained in the supercenter of R .
- Axiom 3) implies that t is 0 on the space of supercommutators $[R, R]$.
- Axiom 4) implies that $t(R)$ is an F –subalgebra and that t is $t(R)$ –linear.

We call $t(R)$ the *trace algebra* of R .

The basic example of algebra with trace is of course the algebra of $m \times m$ matrices over a supercommutative algebra B with the usual trace.

We have made no special requirements on the value of $t(1)$. Of course in many important examples this is a positive integer.

SAT form a category where objects are SAT and morphisms algebra homomorphisms φ which commute with the trace and grading.

$$d(\varphi(a)) = d(a), \quad t(\varphi(a)) = \varphi(t(a)).$$

Recall that in an algebra R with some extra operations, an ideal I must be stable under these operations so that R/I can inherit the structure.

In a SAT a superideal is an $\mathbb{Z}/(2)$ graded ideal I stable under the trace, that is if $a \in I$ then $t(a) \in I$. The usual homomorphism theorems are then valid for SAT.

Given a SAT R , if we *forget the trace* we just have an associative superalgebra.

On the other hand, there is a simple formal construction which associates to any F superalgebra R a superalgebra with trace R_t .

Denote by $[R, R]$ the linear span of supercommutators, a graded subspace of R .

We then define $R_t := R \otimes \mathbb{K}[R/[R, R]]$, with $\mathbb{K}[R/[R, R]]$ the superalgebra of Formula (2).

Given an element $r \in R$, its trace $t(r)$ is the class of r in $R/[R, R]$ (this is a functor adjoint to the previous forgetful functor). This extends to a trace on R_t with trace algebra $\mathbb{K}[R/[R, R]]$.

Proposition 3.3. *A graded homomorphism $f : R \rightarrow A$ with A a trace superalgebra extends uniquely to a trace preserving map $f : R_t \rightarrow A$.*

3.4. Free algebras. In particular for a given a finite or countable alphabet $X = \{x_1, x_2, \dots, x_r, \dots\}$, $Y = \{y_1, y_2, \dots, y_r, \dots\}$, one gets a *free superalgebra with trace* in X, Y applying the previous construction to the free superalgebra $F\langle X, Y \rangle$ where we give degree 0 the variables Y and 1 to X .

Each monomial M in X, Y has a degree in X and in Y and as $\mathbb{Z}/(2)$ degree we take the degree in X modulo 2 and denote it by $d(M) \in \{0, 1\}$.

Lemma 3.5. *The space $F\langle X, Y \rangle / [F\langle X, Y \rangle, F\langle X, Y \rangle]$ has as basis over F the monomials up to signed cyclic equivalence.*

The class of a monomial M is denoted by $t(M)$.

Proof. Given a monomial $M = AB$ in X, Y one has, by definition 2.2 (5) $[A, B] := AB - (-1)^{d(A)d(B)}BA$, so

$$t(AB) = (-1)^{d(A)d(B)}t(BA).$$

This is the meaning of signed cyclic equivalence □

Having denoted by $t(M)$ the cyclic equivalence class of a monomial we have that $\mathbb{K}[F\langle X, Y \rangle / [F\langle X, Y \rangle, F\langle X, Y \rangle]]$ can be thought of as superpolynomial algebra $\mathcal{T} = \mathbb{K}[t(M)]$, that is the algebra of polynomials in the degree 0 elements $t(M)$ tensor the exterior algebra in the degree 1 elements $t(M)$.

We denote by $\mathcal{F}_T = F\langle X, Y \rangle_t = F\langle X, Y \rangle \otimes_A \mathcal{T}$, the free superalgebra with trace. The trace of a monomial M is $t(M)$.

3.6. Strings and encoding map. *This paragraph establishes the notations needed to discuss the invariant theory.*

We now discuss the *encoding map* which describes the subspace $\mathcal{F}_T(e, f)$ of the free algebra $\mathcal{F}_T$, of the elements which are multilinear in the variables $x_1, \dots, x_e; y_1, \dots, y_f$ in terms of the symmetric group S_{e+f+1} .

Respectively the subspace $\mathcal{T}(e, f)$ of its trace algebra $\mathcal{T}$, which are multilinear in $x_1, \dots, x_e; y_1, \dots, y_f$ in terms of the symmetric group S_{e+f} .

Consider a set $A := \{a_1 < a_2 < \dots < a_m\} \subset \mathbb{N}^+$ of positive integers with $|A| = m$.

Definition 3.7. A coloring of A is a map $C : A \rightarrow \{0, 1\}$. The elements with $C(a) = 0$ are called *bosonic* while those with $C(a) = 1$ *fermionic*.

A string with support in A is a bijection from $\{1, 2, \dots, m\}$ to A and it is displayed as a word $w = a_{\sigma(1)}, a_{\sigma(2)}, \dots, a_{\sigma(m)}$. This word can also be interpreted as a permutation σ of A .

For such a word and coloring C set w_C^f to be the substring formed by the fermionic elements.

We set $d_f^C(w) := |w_C^f|$ its cardinality.

The sign $\epsilon_C(w)$ of w is the sign of the permutation which reorders the fermionic string w_C^f .

Given a permutation σ of A this can also be displayed in its cycle form.

This second display is not unique since $(A, B) = (B, A)$.

The canonical way of displaying is to start from the first element of A write the string of its cycle close it with a parenthesis, open a parenthesis write the first of the elements of A which have not appeared and iterate the procedure.

In this display the first element of each cycle is minimal in the cycle and the first elements of the various cycles are increasing from left to right.

We define the corresponding canonical parenthesized string or p-string as $\lambda(\sigma)$.

We also need a second string.

Start from $\lambda(\sigma)$ single out the cycle which contains the last element a_m of A displayed as (Ba_m) then remove this cycle and place the string B at the end denote this $\mu(\sigma)$.

Example 3.8. $A = \{1, 2, \dots, 8, 9\}$; $\sigma = 4, 1, 8, 2, 3, 7, 6, 9, 5$

$$\lambda(\sigma) = (1, 4, 2)(3, 8, 9, 5)(6, 7) \implies \mu(\sigma) = (1, 4, 2)(6, 7)5, 3, 8.$$

$$\lambda(\tau) = (1, 4, 2)(3, 9, 5)(6, 8, 7) \implies \mu(\tau) = (1, 4, 2)(6, 8, 7)5, 3.$$

3.8.1. Colors. Let $\mathcal{F}_T$ be as before the free superalgebra with trace in infinitely many fermionic variables x_i and bosonic variables y_i .

Given a subset $A \subset \mathbb{N}$ with a coloring C and a string $w = a_1 a_2 \dots a_m$ we evaluate this string in $\mathcal{F}_T$ by setting

$$(4) \quad \tau_C(j) = \begin{cases} y_j & \text{if } j \text{ is bosonic} \\ x_j & \text{if } j \text{ is fermionic} \end{cases}$$

$$\tau_C(w) = \epsilon_C(w) \tau_C(a_1) \tau_C(a_2) \dots \tau_C(a_m)$$

For a p-string $w = (w_1)(w_2) \dots w(r)$ we set, use Definition 3.7

$$\tau_C(w) = \epsilon_C(w) \text{tr}(\tau_C(w_1)) \text{tr}(\tau_C(w_2)) \dots \text{tr}(\tau_C(w_r)) \in \text{tr}(\mathcal{F}_T).$$

We also consider the partial p -strings $w = (w_1)(w_2) \cdots w(r)w_{r+1}$ and their colored evaluation

$$\tau_C(w) = \epsilon_C(w) \text{tr}(\tau_C(w_1)) \text{tr}(\tau_C(w_2)) \cdots \text{tr}(\tau_C(w_r)) \tau_C(w_{r+1}) \in \mathcal{F}_T.$$

To a p -strings $w = (w_1)(w_2) \cdots w(r)(w_{r+1}a_m)$ we associate the partial p -strings $\bar{w} = (w_1)(w_2) \cdots w(r)w_{r+1}$.

Now let $A = \{1, 2, \dots, e, e+1, \dots, e+f\}$ and color

$$C_{e,f}(i) = \begin{cases} 0 & \text{if } i \leq e \\ 1 & \text{if } i > e \end{cases}$$

It is then convenient to redefine $\tau_{C_{e,f}}$ on the Fermionic indices $e+j$ by setting $\tau_{C_{e,f}}(e+j) := x_j$.

In this way p -strings in A correspond to trace monomials in the variables $\{y_1, y_2, \dots, y_e, x_1, \dots, x_f\}$.

Definition 3.9. We define $\mathcal{T}(e, f) \subset \mathcal{T}$ as the space of elements of $\mathcal{T}$ which are multilinear in the variables $y_1, \dots, y_e; x_1, \dots, x_f$.

We define $\mathcal{F}_T(e, f) \subset \mathcal{F}_T$ as the space of elements of $\mathcal{F}_T$ which are multilinear in the variables $y_1, \dots, y_e; x_1, \dots, x_f$.

Lemma 3.10. *The map $G \mapsto \text{tr}(Gy_{e+1})$ is a linear isomorphism between $\mathcal{F}_T(e, f)$ and $\mathcal{T}(e+1, f)$.*

The map $G \mapsto \text{tr}(Ax_{f+1})$ is a linear isomorphism between $\mathcal{F}_T(e, f)$ and $\mathcal{T}(e, f+1)$.

Proposition 3.11. *[Encoding map]*

Take a permutation $\sigma \in S_{e+f}$ written as a p -string,

$\tau_{C_{e,f}}(\sigma)$ depends only on $\sigma \in S_m$ and not on its display as p -string. We set $\Phi_\sigma := \tau_{C_{e,f}}(\sigma)$.

For a permutation $\delta \in S_{e+f+1}$ written as a parenthesized string

$$(w_1)(w_2) \cdots w(r)(w_{r+1}, m+1)$$

in which the last cycle is a string ending with $m+1$, we set $\bar{\delta}$ the partial p -string $w = (w_1)(w_2) \cdots w(r)w_{r+1}$.

$\tau_{C_{e,f}}(\bar{\delta})$ depends only on $\delta \in S_{e+f+1}$ and not on its display as p -string. We set $\Psi_\sigma := \tau_{C_{e,f}}(\delta)$.

- (1) The map $\sigma \mapsto \Phi_\sigma = \tau_{C_{e,f}}(\sigma)$ is a linear isomorphism with the subspace $\mathcal{T}(e, f)$ of its trace algebra $\mathcal{T}$ which are multilinear in the variables $y_1, \dots, y_e; x_1, \dots, x_f$.
- (2) The map $\sigma \mapsto \Psi_\sigma = \tau_{C_{e,f}}(\bar{\sigma})$ is a linear isomorphism with the space $\mathcal{F}_T(e, f)$ of the free algebra $\mathcal{F}_T$ which are multilinear in the variables $y_1, \dots, y_e; x_1, \dots, x_f$.
- (3) $\sigma \in S_{e+f+1}$ then $\Phi_\sigma = \text{tr}(\Psi_\sigma x_{f+1})$.

Proof. The fact that the two maps depend only on σ is a consequence of the axioms on the trace in $\mathcal{F}_T$. The fact that they are linear isomorphisms depends on the definition of multilinear. $\square$

Definition 3.12. $\tau_{C_{e,f}}$ is called the *encoding map*.

By definition of free superalgebra given any F –superalgebra with trace $R = R_0 \oplus R_1$ the homomorphisms from $\mathcal{F}_T = F\langle X, Y \rangle[t(M)]$ to R (trace preserving) are given by *evaluations* of polynomials associated to maps $X \rightarrow R_1, Y \rightarrow R_0$.

3.13. Trace identities. Using evaluations allows us to introduce the notion of *trace identity* for a trace superalgebra R .

Definition 3.14. A trace identity for a trace superalgebra R is a element $\mathbf{f} \in \mathcal{F}_T = F\langle X, Y \rangle[t(M)]$ of the free algebra such that $\mathbf{f}$ vanishes under all evaluations in R .

Let us now remark that the set of all trace identities of a trace algebra R is a special ideal of $\mathcal{F}_T$, called a T –ideal, that is an ideal in $\mathcal{F}_T$ closed under trace and under all substitutions of variables.

By definition a substitution of a variable x must be an element of degree 1 in $\mathcal{F}_T$, while a substitution of a variable y must be an element of degree 0 in $\mathcal{F}_T$.

A basic question for any algebra R is to describe a set of generators of its trace identities as T –ideal,

The theory is really effective only when F is a field of characteristic 0, so from now on, in this section, we assume that this is the case (the reader can assume $F = \mathbb{C}$).

3.14.1. *Varieties.* A subset $S \subset F\langle X, Y \rangle \otimes_A \mathcal{T}$ generates a T -ideal I_S and the *variety* $\mathcal{V}_S$ of all trace superalgebras satisfying all the trace identities S .

Of course the superalgebras in $\mathcal{V}_S$ satisfy also all the trace identities in the T -ideal I_S .

The quotient $F_S := F\langle X, Y \rangle \otimes_A \mathcal{T} / I_S$ satisfies I_S and any identity of F_S is also in I_S .

One easily sees that F_S is a free algebra, in the classes of the elements x, y for the variety $\mathcal{V}_S$.

Let $I_S \neq \{0\}$ be a T -ideal. Given a trace superalgebra $R \in \mathcal{V}_S$, consider the trace superalgebra $M_n(R)$. By a Theorem of Regev the T -ideal $I_{n,R}$ of trace identities of $M_n(R)$ is also $\neq \{0\}$.

Let $F_{S,n}$ denote the free superalgebra for the variety associated to $I_{n,R}$, then the corresponding free algebra $F_{n,R}\langle X, Y \rangle := F\langle X, Y \rangle \otimes_A \mathcal{T} / I_{n,R}$ can be described by the method of *generic matrices*.¹

Consider the free algebra $F\langle X(n), Y(n) \rangle \otimes_A \mathcal{T}$ in the variables

$$(5) \quad X(n) := \{x_{h,k}^{(i)}\}, \quad Y(n) := \{y_{h,k}^{(i)}\}. \quad h, k = 1, \dots, n, \quad i = 1, 2, \dots$$

Define the *generic matrices*

$$(6) \quad X_i := (x_{h,k}^{(i)}), \quad Y_j := (y_{h,k}^{(j)}), \quad i, j = 1, 2, \dots$$

We then have an injective map

$$\mathbf{i} : F\langle X, Y \rangle \otimes_F \mathcal{T} \mapsto M_n(F\langle X(n), Y(n) \rangle \otimes_F \mathcal{T}), \quad \mathbf{i} : x_i \mapsto X_i, \quad \mathbf{i} : y_j \mapsto Y_j.$$

Given any evaluation $\pi : F\langle X, Y \rangle \otimes_A \mathcal{T} \rightarrow M_n(R)$, let $\pi(x_i) = (a_{h,k}^{(i)})$, $\pi(y_j) = (b_{h,k}^{(j)})$.

Consider the evaluation $\bar{\pi} : F\langle X(n), Y(n) \rangle \otimes_A \mathcal{T} \rightarrow R$ where $\bar{\pi} : x_{h,k}^{(i)} \mapsto a_{h,k}^{(i)}$, $\bar{\pi} : y_{h,k}^{(j)} \mapsto b_{h,k}^{(j)}$ we have the factorization

$$(7) \quad \begin{array}{ccc} F\langle X, Y \rangle \otimes_A \mathcal{T} & \xrightarrow{\mathbf{i}} & M_n(F\langle X(n), Y(n) \rangle \otimes_F \mathcal{T}) \\ & \searrow \pi & \downarrow M_n(\bar{\pi}) \\ & & M_n(R) \end{array}$$

and clearly the evaluation π vanishes if and only if $\bar{\pi}$ vanishes.

¹Although I introduced generic matrices in my 1966 Ph thesis I am not aware if this remark has been made before in this generality

Both maps factor through the free algebras in the varieties of $M_n(R)$ and of R giving:

$$(8) \quad \begin{array}{ccc} F\langle X, Y \rangle \otimes_F \mathcal{T} & \xrightarrow{\mathbf{i}} & M_n(F\langle X(n), Y(n) \rangle \otimes_F \mathcal{T}) \\ \rho \downarrow & & \downarrow M_n(\bar{\rho}) \\ F\langle X, Y \rangle \otimes_F \mathcal{T} / I_{M_n(R)} & \xrightarrow{\bar{\mathbf{i}}} & M_n(F\langle X(n), Y(n) \rangle \otimes_F \mathcal{T} / I_R) \\ & \searrow \pi' & \downarrow M_n(\bar{\pi}') \\ & & M_n(R) \end{array}$$

With ρ the quotient map and $\pi = \pi' \circ \rho$. One can apply this construction to the superalgebra $R = F\langle X(n), Y(n) \rangle \otimes_F \mathcal{T} / I_R$ and $\pi = \mathbf{i} \circ \rho$, $\pi' = \mathbf{i}$.

We obtain that $\bar{\mathbf{i}}$ is injective so it is an isomorphism with the trace subalgebra of $M_n(F\langle X(n), Y(n) \rangle \otimes_F \mathcal{T} / I_R)$ generated by the images $\bar{\mathbf{i}}$ of the elements $\rho(x_i)$, $\rho(y_i)$ which are the images under $M_n(\bar{\rho})$ of the *matrix variables* X_i , Y_j .

We have proved

Theorem 3.15. *The trace subalgebra of $M_n(F\langle X(n), Y(n) \rangle \otimes_F \mathcal{T} / I_R)$ generated by the images under $M_n(\bar{\rho})$ of the matrix variables X_i , Y_j is a free algebra in the variety generated by $M_n(R)$.*

4. MATRICES

4.1. Generic matrices. We want to apply Theorem 6 to $R = \mathbb{K}(V)$ and show that the free algebra in the variety generated by $M_n(R)$ is described as an algebra of invariants, Theorem 4.3.

Let $G = GL(n, \mathbb{C})$ be the linear group acting by conjugation on matrices.

This action factors through the projective linear group $P_n = GL(n, \mathbb{C}) / \mathbb{C}^*$.

This action induces an action of G on the vector space $V = (M_n)^{\oplus m}$ and on its dual $V^* = (M_n^*)^{\oplus m}$ of linear functions on m -tuples of matrices.

Decompose $m = e + f$ and give a structure to V , and so also to V^* , of graded vector space, by considering the first h summands in degree 0, the *bosonic matrices* and the remaining summands of degree 1, the *fermionic matrices*.

We then have that the action of G extends as automorphisms to $\mathbb{K}(V^*)$ and we may then consider the invariant subalgebra $\mathbb{K}(V^*)^G$ of invariants for this action.

We also take $\mathbb{K}(V^*) \otimes M_n(\mathbb{C})$, with the diagonal action of G .

We then think of vectors $v \in V_0$ as elements with a basis $\{\eta_{h,k}^{(i)}, h, k = 1, \dots, n, i = 1, 2, \dots\}$, the $\eta_{h,k}^{(i)}$ are polynomial variables, and of $v \in V_1$ as Grassmann (fermionic) elements with a basis $\{\xi_{h,k}^{(i)}, h, k = 1, \dots, n, i = 1, 2, \dots\}$ as Grassmann (fermionic) variables.

They give rise to $e + f$ generic Grassmann (bosonic and fermionic) matrices

$$(9) \quad \eta_1, \dots, \eta_e; \quad \eta_i = (\eta_{h,k}^{(i)}) = \sum_{h,k} e_{h,k} \otimes \eta_{h,k}^{(i)} \in M_n(\mathbb{C}) \otimes S(V_0)$$

$$(10) \quad \xi_1, \dots, \xi_f; \quad \xi_i = (\xi_{h,k}^{(i)}) = \sum_{h,k} e_{h,k} \otimes \xi_{h,k}^{(i)} \in M_n(\mathbb{C}) \otimes \bigwedge V_1$$

of size n .

Remark 4.2. The action of G is defined so that the matrices

$$g \cdot \eta_i = \sum_{h,k} g \cdot e_{h,k} \otimes g \cdot \eta_{h,k}^{(i)}, \quad g \cdot \xi_i = \sum_{h,k} g \cdot e_{h,k} \otimes g \cdot \xi_{h,k}^{(i)}$$

are G -invariant. They are *generic matrices* as in Theorem 3.15 or in page 1.1 for 1 variable.

We set $S = S_{e,f,n} = \mathbb{K}(V^*) = S[\eta_{h,k}^{(i)}] \otimes \bigwedge[\xi_{h,k}^{(i)}]$.

We want to study the G invariants of the algebra $S_{e,f,n}$ and of $n \times n$ matrices $M_n(S_{e,f,n}) = M_n(\mathbb{C}) \otimes S_{e,f,n}$ under the diagonal tensor product action.

For this consider the evaluation of the free algebra $\mathcal{F}_T = F\langle X, Y \rangle \otimes \mathcal{T}$, with $X = x_1, x_2, \dots, x_f$; $Y = y_1, y_2, \dots, y_e$ in $M_n(S_{e,f,n}) = M_n(\mathbb{C}) \otimes S_{e,f,n}$ given by $ev : y_i \mapsto \eta_i$ and $ev : x_i \mapsto \xi_i$. We want to prove:

Theorem 4.3. *The previous evaluation map surjects to the G -invariants. That is:*

The algebra $\mathcal{T}_n$ of invariants of both bosonic and fermionic matrices is generated by the elements $tr(M)$ for all monomials M .

The algebra $\mathcal{E}_n$ of equivariant maps of both bosonic and fermionic matrices is generated by the elements $tr(M)$ and by the generic matrices ξ_i, η_j .

Proof. We start by observing that ev maps to the invariants.

Lemma 4.4. *The elements η_i, ξ_j are invariant.*

The map trace $tr : M_n(\mathbb{C}) \otimes S_{e,f,n} \rightarrow S_{e,f,n}$, $tr(A \otimes s) = tr(A)s$ is equivariant.

Using the classical method of polarization of Aronhold we can reduce to prove that the map ev is surjective for multilinear invariants.

For this we use a variation of the classical *symbolic method* suggested by the Formulas of Kostant [11], Lemma 4.9.

Multilinear elements are defined as follows:

we have the diagonal $m \times m$ matrices $d = (d_1, \dots, d_m)$ acting on $\xi_{h,k}^{(i)}$ by $d(A_1, \dots, A_m) = (d_1 A_1, \dots, d_m A_m)$ this action commutes with that of G .

The multilinear elements of $M_n(\mathbb{C}) \otimes S_{e,f,n}$, $S_{e,f,n}$ are the ones for which $d \cdot a = \prod_i d_i a$.

We have

$$S = \mathbb{K}(V^*) = \underbrace{S(M_n^*) \otimes \dots \otimes S(M_n^*)}_{e \text{ times}} \otimes \underbrace{\bigwedge M_n^* \hat{\otimes} \dots \hat{\otimes} \bigwedge M_n^*}_{f \text{ times}}$$

and the multilinear elements are

$$\mathbb{K}(V^*)_{mult} = ((M_n^*)^{\otimes m}) = \underbrace{M_n^* \otimes \dots \otimes M_n^*}_{m \text{ times}},$$

we thus want to understand a symbolic formula for $((M_n^*)^{\otimes m})^G \subset \mathbb{K}(V^*)$.

Similarly for the multilinear elements of $M_n(\mathbb{K}(V^*))$ which are identified to $M_n(\mathbb{C}) \otimes \underbrace{M_n^* \otimes \dots \otimes M_n^*}_{m \text{ times}}$.

We have

$$(M_n^{\otimes m}) = \text{End}(\mathbb{C}^{\otimes m})$$

and the invariants are linearly generated by the symmetric group S_m acting on $\mathbb{C}^{\otimes m}$ by

$$\sigma(v_1 \otimes v_2 \otimes \dots \otimes v_m) = v_{\sigma^{-1}(1)} \otimes v_{\sigma^{-1}(2)} \otimes \dots \otimes v_{\sigma^{-1}(m)}.$$

We have a non degenerate duality pairing between $\text{End}(\mathbb{C}^{\otimes m})$ and its dual $\text{End}(\mathbb{C}^{\otimes m})^*$ given by $\text{tr}(AB)$.

Since the invariants $(M_n^{\otimes m})^G$ are encoded by elements of the symmetric group so we have to understand a permutation σ as element of $\mathbb{K}(V^*)^G$.

Lemma 4.5. *Consider the map $\varphi : M_n(\mathbb{C})^{\otimes m} \rightarrow \mathbb{K}(V^*)$ defined by*

$$(11) \quad \varphi : A_1 \otimes \dots \otimes A_m \mapsto \text{tr}(A_1 \eta_1 \otimes \dots \otimes A_e \eta_e \otimes A_{e+1} \xi_1 \otimes \dots \otimes A_m \xi_f)$$

$$(12) \quad = \text{tr}(A_1 \eta_1) \dots \text{tr}(A_e \eta_e) \text{tr}(A_{e+1} \xi_1) \dots \text{tr}(A_m \xi_f).$$

this map is injective G equivariant and its image is the space of elements of $\mathbb{K}(V^)$ which are multilinear in the η_i, ξ_j ,*

Proof. Given $A = (a_{i,j})$ and $\eta = (\eta_{i,j})$ bosonic, $\xi = (\xi_{i,j})$ fermionic matrix we have

$$\text{tr}(e_{h,k} \cdot \eta) = \eta_{k,h}, \quad \text{tr}(e_{h,k} \cdot \xi) = \xi_{k,h}.$$

□

At this point the map φ identifies the invariants in $M_n(\mathbb{C})^{\otimes m}$, which are the linear span of S_m , to the multilinear invariants in $\mathbb{K}(V^*)$.

For multilinear invariants in $M_n(\mathbb{K}(V^*))$ we use the usual trick, we use Lemma 3.10, we add another generic matrix, which can be either bosonic or fermionic, let us say it is η_{e+1} then we have a bijective map between multilinear G invariants in $\eta_1, \dots, \eta_e; \xi_1, \dots, \xi_f$ of $M_n(\mathbb{K}(V^*))$ and multilinear G invariants in $\eta_1, \dots, \eta_e, \eta_{e+1}; \xi_1, \dots, \xi_f$ of $M_n(\mathbb{K}(V^*))$ by the formula

$$(13) \quad f(\eta_1, \dots, \eta_e; \xi_1, \dots, \xi_f) \mapsto \text{tr}(f(\eta_1, \dots, \eta_e; \xi_1, \dots, \xi_f) \eta_{e+1}).$$

We will see how, knowing a Formula for $\text{tr}(f(\eta_1, \dots, \eta_e; \xi_1, \dots, \xi_f) \eta_{e+1})$ we can decode it as a Formula for $f(\eta_1, \dots, \eta_e; \xi_1, \dots, \xi_f)$.

Definition 4.6.

$$(14) \quad \varphi_\sigma := \text{tr}(\sigma^{-1} \circ \eta_1 \otimes \dots \otimes \eta_e \otimes \xi_1 \otimes \dots \otimes \xi_f), \quad \sigma \in S_m.$$

$$(15) \quad \psi_\sigma : \quad \varphi_\sigma = \text{tr}(\psi_\sigma \eta_{e+1}), \text{ or } \text{tr}(\psi_\sigma \xi_{f+1}). \quad \sigma \in S_{m+1}.$$

Proposition 4.7. *The multilinear G invariants in $\mathbb{K}(V^*)$, by Lemma 4.5, are linearly generated by the elements φ_σ .*

The multilinear G invariants in $M_n(\mathbb{K}(V^))$, by Lemma 4.5, are linearly generated by the elements ψ_σ .*

4.7.1. *Symbolic method.*

We want to prove

Theorem 4.8. *We have, with the notations of Proposition 3.11 and ev the evaluation of Theorem 4.3*

$$(16) \quad \varphi_\sigma = ev(\Phi_\sigma), \quad \psi_\sigma = ev(\Psi_\sigma).$$

In order to prove this Theorem we develop an analogue of the argument of Kostant in the usual case [11] Lemma 4.9.

Consider

$$\{u_{i,1}, \dots, u_{i,n}, \quad \varphi_{i,1}, \dots, \varphi_{i,n}\}, \quad i = 1, \dots, m$$

with $u_{i,j}$ ordinary variables for $i \leq e$ and Grassmann variables for $i > e$ and $\varphi_{i,j}$ usual variables which generate the algebra

$$U = \mathbb{K}(u_{i,1}, \dots, u_{i,n}) \otimes S(\varphi_{i,1}, \dots, \varphi_{i,n}); \quad i = 1, \dots, m.$$

Consider $u_i = (u_{i,1}, \dots, u_{i,n})$ Grassmann vectors in $U^n = \mathbb{C}^n \otimes U$ and $\varphi_i = (\varphi_{i,1}, \dots, \varphi_{i,n})$ ordinary forms in U^n , we act by G on vectors and forms in the usual way.

We consider U^n as right U module and φ as an U linear map

$$\varphi_i(s_1, \dots, s_m) := \sum_j \varphi_{i,j} s_j = \langle \varphi_i, (s_1, \dots, s_m) \rangle.$$

On the algebra U acts the group $G = GL(n)$. This is the usual action, we think of φ as a row vector and then, for $g \in G$ we set $g \cdot \varphi := \varphi g^{-1}$ and u as column vector and $g \cdot u$ is matrix multiplication

We identify $u \otimes \varphi$ with a (rank 1) matrix with coefficients in U by the usual Formula $v \in U^n$, $v \mapsto u \langle \varphi, v \rangle$. We then have

$$(17) \quad u_i \otimes \varphi_i \circ u_j \otimes \varphi_j(s) = u_i \otimes \varphi_i(u_j \langle \varphi_j, s \rangle) = u_i \langle \varphi_i, u_j \rangle \langle \varphi_j, s \rangle$$

$$\implies u_i \otimes \varphi_i \circ u_j \otimes \varphi_j = u_i \langle \varphi_i, u_j \rangle \otimes \varphi_j; \quad \text{tr}(u \otimes \varphi) = \langle \varphi, u \rangle$$

This map is G equivariant in the sense that $(gu) \otimes (\varphi g^{-1}) = g(u \otimes \varphi) g^{-1}$.

We have a G equivariant map $\pi : \mathbb{K}(V^*) \rightarrow U$ of $\mathbb{Z}/(2)$ graded algebras.

$$\pi : \eta_{h,k}^{(i)} \mapsto u_{i,h} \otimes \varphi_{i,k}, \quad \xi_{h,k}^{(i)} \mapsto u_{i+e,h} \otimes \varphi_{i+e,k}.$$

We set

$$(18) \quad \alpha_i := u_i \otimes \varphi_i, \forall i, \quad \bar{\eta}_i := u_i \otimes \varphi_i, \quad i \leq e; \quad \bar{\xi}_i := u_{e+i} \otimes \varphi_{e+i}, \quad i \leq f.$$

This map induces a homomorphism $\bar{\pi} : M_n(\mathbb{C}) \otimes \mathbb{K}(V^*) \rightarrow M_n(\mathbb{C}) \otimes U$ of $\mathbb{Z}/(2)$ graded algebras.

Under $\bar{\pi}$ we have $\bar{\pi}(\eta_i) = \bar{\eta}_i$, $\bar{\pi}(\xi_i) = \bar{\xi}_i$.

The map $\bar{\pi}$ is not injective but, since rank 1 matrices linearly span the space of matrices we have

Lemma 4.9. *The map $\bar{\pi}$ restricted to the multilinear elements is an isomorphism to the space of elements which are multilinear in both the u_i and φ_j .*

Proof. It is clearly enough to do this for linear elements that is $m = 1$ with z a generic matrix out of the list ξ, η mapping to $\alpha = u \otimes \varphi$.

For $m = 1$ the elements which are linear both in u and in φ are the n^2 linearly independent elements $u_j \varphi_i$.

For $A \in M_n(\mathbb{C})$ we have $A \xrightarrow{\tau} A \cdot z \xrightarrow{\bar{\pi}} A \cdot u \otimes \varphi$. For a matrix unit $e_{i,j} = e_i \otimes e^j$ we have $e_i \otimes e^j \cdot u \otimes \varphi = u_j e_i \otimes \varphi = u_j \varphi_i e_i \otimes e^j$. $\square$

As a consequence for computing Formula (16) we can compose this with $\bar{\pi}$ and compute

$$(19) \quad \bar{\varphi}_\sigma = \bar{\pi}(ev(\Phi_\sigma)), \quad \bar{\psi}_\sigma = \bar{\pi}(ev(\Psi_\sigma)).$$

That is we evaluate x_i in $\bar{\xi}_i$ and y_j in $\bar{\eta}_j$.

We have the usual formula

$$u_1 \otimes \varphi_1 \otimes \cdots \otimes u_m \otimes \varphi_m \circ \sigma = u_1 \otimes \varphi_{\sigma(1)} \otimes \cdots \otimes u_m \otimes \varphi_{\sigma(m)}$$

hence

$$(20) \quad \bar{\varphi}_\sigma = tr(u_1 \otimes \varphi_1 \otimes \cdots \otimes u_m \otimes \varphi_m \circ \sigma^{-1}) = \langle \varphi_{\sigma^{-1}(1)}, u_1 \rangle \cdots \langle \varphi_{\sigma^{-1}(m)}, u_m \rangle.$$

We have to *decode* Formula (20).

Lemma 4.10.

$$(21) \quad \alpha_{i_1} \circ \alpha_{i_2} \circ \cdots \circ \alpha_{i_k} = u_{i_1} \langle \varphi_{i_1}, u_{i_2} \rangle \langle \varphi_{i_2}, u_{i_3} \rangle \cdots \langle \varphi_{i_{k-1}}, u_{i_k} \rangle \otimes \varphi_{i_k}.$$

$$(22) \quad tr(\alpha_{i_1} \circ \alpha_{i_2} \circ \cdots \circ \alpha_{i_k}) = \langle \varphi_{i_k}, u_{i_1} \rangle \langle \varphi_{i_1}, u_{i_2} \rangle \langle \varphi_{i_2}, u_{i_3} \rangle \cdots \langle \varphi_{i_{k-1}}, u_{i_k} \rangle.$$

Proof. By induction on k , we have

$$\alpha_{i_2} \circ \cdots \circ \alpha_{i_k} = u_{i_2} \langle \varphi_{i_2}, u_{i_3} \rangle \cdots \langle \varphi_{i_{k-1}}, u_{i_k} \rangle \otimes \varphi_{i_k}$$

by Formula (17)

$$\alpha_{i_1} \circ u_{i_2} \langle \varphi_{i_2}, u_{i_3} \rangle \cdots \langle \varphi_{i_{k-1}}, u_{i_k} \rangle \otimes \varphi_{i_k}$$

gives Formula (21) since

$$\langle \varphi_{i_1}, u_{i_2} \langle \varphi_{i_2}, u_{i_3} \rangle \cdots \langle \varphi_{i_{k-1}}, u_{i_k} \rangle \rangle = \langle \varphi_{i_1}, u_{i_2} \rangle \langle \varphi_{i_2}, u_{i_3} \rangle \cdots \langle \varphi_{i_{k-1}}, u_{i_k} \rangle$$

□

We start by computing for $\sigma = (i_1, i_2, \dots, i_k)$.

Then we have $\langle \varphi_{i_{h-1}}, u_{i_h} \rangle = \langle \varphi_{\sigma^{-1}(i_h)}, u_{i_h} \rangle$, $\forall h$ so the product of Formula (22) is a rearrangement of the product of Formula (20).

We can then reorder the formula (20)

$$(23) \quad \bar{\pi}(\varphi_\sigma) = \langle \varphi_{\sigma^{-1}(1)}, u_1 \rangle \cdots \langle \varphi_{\sigma^{-1}(k)}, u_k \rangle = \epsilon_f(\sigma) \langle \varphi_{i_1}, u_{i_2} \rangle \langle \varphi_{i_2}, u_{i_3} \rangle \cdots \langle \varphi_{i_{k-1}}, u_{i_k} \rangle.$$

where $\epsilon_f(\sigma) = \epsilon_{C_{e,f}}(\sigma)$ is the sign of the permutation which rearranges the Fermionic indices of the vectors u_i which in the left are increasing to the list $i_1, i_2, \dots, i_k$.

Denote by $z_i := y_i$, $i \leq e$, $z_{e+i} := x_i$ then:

In general by decomposing

$$\sigma^{-1} = (i_1, i_2, \dots, i_{k_1}) \cdots (i_{k+a}, i_{k+a+1}, \dots, i_{k+a+r})$$

into cycles $c_1 c_2 \cdots c_r$ we claim that

Proposition 4.11. *We have*

$$\varphi_\sigma = ev(\tau_{C_{e,f}}(\sigma)) = ev(\Phi_\sigma), \quad \psi_\sigma = ev(\tau_{C_{e,f}}(\bar{\sigma})) = ev(\Psi_\sigma).$$

Proof.

$$(24) \quad \tau_{C_{e,f}}(\sigma) \stackrel{(4)}{=} \epsilon_f(\sigma) \cdot tr(z_{i_1}, z_{i_2} \cdots z_{i_{k_1}}) \cdots tr(z_{i_{k+a}}, z_{i_{k+a+1}} \cdots z_{i_{k+a+r}})$$

with $\epsilon_f(\sigma) = \epsilon_{C_{e,f}}(\sigma)$ the sign of the permutation which rearranges the fermionic indices of the p-string of σ .

To prove that $\varphi_\sigma = ev(\tau_{C_{e,f}}(\sigma))$ we compute $\bar{\pi}(\varphi_\sigma) = \bar{\varphi}_\sigma$, which by Formula (23) and Lemma 4.10 is

$$(25) \quad \epsilon_f(\sigma) \cdot tr(\alpha_{i_1}, \alpha_{i_2} \cdots \alpha_{i_{k_1}}) \cdots tr(\alpha_{i_{k+a}}, \alpha_{i_{k+a+1}} \cdots \alpha_{i_{k+a+r}}).$$

By Lemma 4.9 and the definition of ψ_σ by Formula (15) the claim follows. $\square$

We have completed the proof of Theorem 4.3. $\square$

Example 4.12. $m = 3 + 4 = 7$, $\sigma = 3, 1, 5, 7, 2, 6, 4 = (1, 3, 5, 2)(4, 7)(6)$

$$\Phi_\sigma = tr(y_1 y_3 x_2 y_2) tr(x_1 x_4) tr(x_3)$$

$$\Psi_\sigma = tr(y_1 y_3 x_2 y_2) tr(x_3) x_1$$

$m = 4 + 3 = 7$, $\sigma = 3, 1, 5, 7, 2, 6, 4 = (1, 3, 5, 2)(4, 7)(6)$

$$\Phi_\sigma = -tr(y_1 y_3 x_1 y_2) tr(y_4 x_3) tr(x_2)$$

$$\Psi_\sigma = tr(y_1 y_3 x_1 y_2) tr(x_2) y_4$$

Remark 4.13. Beware of the fact that for an even cycle c the formula for φ_c may depend also from the way in which c is displayed, since $tr(\eta_1 \circ \eta_2) = \pm tr(\eta_2 \circ \eta_1)$ while $(1, 2) = (2, 1)$.

For a permutation it also depends on the order in which we write the cycles.

One can use the convention $\lambda(\sigma)$ of page 7.

Example 4.14. $m = 0 + 5 = 5$, $\sigma = (2, 4, 1)(5, 3) = (1, 2, 4)(3, 5)$ then

$$\Phi_\sigma = tr(x_2 x_4 x_1) tr(x_5 x_3) = -tr(x_1 x_2 x_4) tr(x_3 x_5)$$

$$\Psi_\sigma = tr(x_2 x_4 x_1) tr(x_5 x_3) = -tr(x_1 x_2 x_4) x_3$$

the sign is that of the permutations $2, 4, 1, 5, 3$, $1, 2, 4, 3, 5$.

5. RELATIONS

By Theorem 3.15 the T -ideal of trace relations for the algebra $M_n(\mathbb{K}(V))$ is the kernel of the evaluation of the free variables to the generic matrices η_i, ξ_j .

By Proposition 3.11 (4) Lemma 4.5 and the definition of the evaluation in matrices, the multilinear trace identities of degrees a, b in the space $\mathcal{T}(a, b)$ of the trace algebra $\mathcal{T}$ are the image under the encoding map $\tau_{C_{a,b}}$ of the kernel $K_{m,n}$ of the map of the group algebra $\mathbb{C}[S_m]$ to the space of endomorphisms of $(\mathbb{C}^n)^{\otimes m}$.

As a vector space this is thus independent of $e, f \mid e + f = m$. As in the Theory of polynomial identities rather than discussing the dimension of $K_{m,n}$, which grows as $m!$ it is better to study its codimension $c_m(M_n(F))$ in $\mathbb{C}[S_m]$ which equals the dimension of its image.

By standard representation theory of the symmetric group this codimension is $c_m(M_n(F)) = \sum_{\lambda \vdash m \mid ht(\lambda) \leq n} \chi_\lambda(1)^2$ where $\chi_\lambda(1)$ is the dimension of the irreducible representation of S_m associated to the partition λ of m which must be restricted to have height $\leq n$.

This sum is hard to compute, for $n = 2$ it is given by $C_n := \frac{1}{n+1} \binom{2n}{n}$ the n^{th} Catalan number. The first are

$$1, 1, 2, 5, 14, 42, 132, 429, 1430, 4862, 16796, 58786, \dots$$

In general we have an asymptotic Formula due to Regev, see [1] Theorem 21.1.2.

Theorem 5.1. (*Regev*)

$$(26) \quad c_m(M_n(F)) \sim A_n m^{\frac{1-n^2}{2}} n^{2(m+1)}.$$

where the constant

$$(27) \quad A_n = \left[\left(\frac{1}{\sqrt{2\pi}} \right)^{n-1} \left(\frac{1}{2} \right)^{\frac{(n^2-1)}{2}} \cdot 1!2! \cdots (n-1)! \cdot n^{\frac{n^2}{2}} \right].$$

5.2. The Cayley–Hamilton identities. The theory follows in a way similar to the classical case, reducing to multilinear identities.

In the classical case the space of multilinear relations of minimal dimension is in dimension n and it is formed by the multiples of the polarized

Cayley–Hamilton identity defined:

(28)

$$CH_n(y_1, \dots, y_n) = \sum_{\sigma \in S_{n+1}} \epsilon_\sigma \Psi_\sigma, \quad \text{tr}(CH_n(y_1, \dots, y_n) y_{n+1}) = \sum_{\sigma \in S_{n+1}} \epsilon_\sigma \Phi_\sigma.$$

This follows from the vanishing of the antisymmetrizer $A_{n+1} = \sum_{\sigma \in S_{n+1}} \epsilon_\sigma \sigma$ as operator on $(\mathbb{C}^n)^{\otimes n+1}$.

Then its interpretation by Formula (14). Therefore the same Formula holds, provided we remember that the formula for φ_σ in the presence of Grassmann variables may contain a sign.

Definition 5.3. For any $e + f = n + 1$ we define:

$$(29) \quad CH_{e,f}(y_1, \dots, y_e, x_1, \dots, x_f) := \tau_{C_{e,f}} \left(\sum_{\sigma \in S_{n+1}} \epsilon_\sigma \bar{\sigma} \right) = \sum_{\sigma \in S_{n+1}} \epsilon_\sigma \Psi_\sigma$$

$$(30) \quad T_{e,f}(y_1, \dots, y_e, x_1, \dots, x_{f+1}) := \tau_{C_{e,f}} \left(\sum_{\sigma \in S_{n+1}} \epsilon_\sigma \sigma \right) = \sum_{\sigma \in S_{n+1}} \epsilon_\sigma \Phi_\sigma,$$

for $C_{e,f}$ the coloring $C_{e,f}(i) = 0$, $i \leq e$ and $C_{e,f}(i) = 1$, $i > e$.

Remark 5.4. By Lemma 3.10 we have two possible identities:

(31)

$$T_{e,f}(y_1, \dots, y_e, x_1, \dots, x_f, x_{f+1}) = \text{tr}(CH_{e,f}(y_1, \dots, y_e, x_1, \dots, x_f) x_{f+1})$$

(32)

$$T_{e,f}(y_1, \dots, y_e, x_1, \dots, x_{f+1}) := \text{tr}(CH_{e-1,f}(y_1, \dots, y_{e-1}, x_1, \dots, x_f, x_{f+1}) y_e)$$

Theorem 5.5. When evaluated in $n \times n$ matrices the two expressions of Formula (29) vanish.

Proof. For the second Formula this follows from the fact that the operator $\sum_{\sigma \in S_{n+1}} \epsilon_\sigma \sigma = 0$ on $(\mathbb{C}^n)^{\otimes n+1}$. For the first one uses the fact that

$$T_{e,f}(y_1, \dots, y_e, x_1, \dots, x_f, x_{f+1}) = \text{tr}(CH_{e,f}(y_1, \dots, y_e, x_1, \dots, x_f) x_{f+1})$$

since x_{f+1} is a disjoint variable this implies the claim. $\square$

For $m = 0 + 3 = 3$ and all 3 matrices fermionic

$$\Phi_1 = \text{tr}(x_1) \text{tr}(x_2) \text{tr}(x_3), \quad \Phi_{(1,2)} = \text{tr}(x_1 x_2) \text{tr}(x_3), \quad \Phi_{(2,3)} = \text{tr}(x_1) \text{tr}(x_2 x_3),$$

$$\Phi_{(1,3)} = -\text{tr}(x_1 x_3) \text{tr}(x_2), \quad \Phi_{(1,2,3)} = \text{tr}(x_1 x_2 x_3), \quad \Phi_{(1,3,2)} = -\text{tr}(x_1 x_3 x_2)$$

Then $\sum_{\sigma \in S_3} \epsilon_\sigma \Phi_\sigma$ is

$$\begin{aligned} & \text{tr}(x_1) \text{tr}(x_2) \text{tr}(x_3) - \text{tr}(x_1 x_2) \text{tr}(x_3) - \text{tr}(x_1) \text{tr}(x_2 x_3) \\ & + \text{tr}(x_1 x_3) \text{tr}(x_2) + \text{tr}(x_1 x_2 x_3) - \text{tr}(x_1 x_3 x_2) \end{aligned}$$

Then $\sum_{\sigma \in S_3} \epsilon_\sigma \Psi_\sigma$ is

$$\text{tr}(x_1)\text{tr}(x_2) - \text{tr}(x_1x_2) - \text{tr}(x_1)x_2 + \text{tr}(x_2)x_1 + x_1x_2 - x_2x_1$$

notice that in this case the expression for CH is antisymmetric while for ordinary matrices is symmetric and the polarization of the 2-variable Cayley–Hamilton identity.

If x_2 is fermionic and $x_1 = y_1$ bosonic then CH equals the same formula of the ordinary case,

$$\text{tr}(y_1)\text{tr}(x_2) - \text{tr}(y_1x_2) - \text{tr}(y_1)x_2 - \text{tr}(x_2)y_1 + y_1x_2 + x_2y_1.$$

This means that the same Formula which expresses the relation in degree $n+1$ has different interpretations for $n+1 = e+f$ so we write these $n+1$ formulas as $CH_{e,f}(y_1, \dots, y_e, x_1, \dots, x_f)$.

Remark 5.6. $CH_{e,f}$ is symmetric in the bosonic and antisymmetric in the fermionic variables. If we set equal to a single y all the bosonic variables we have a non zero element, which for $f=0$ is $n!$ times the usual 1-variable Cayley–Hamilton identity.

5.7. The second fundamental Theorem. We start with some combinatorics.

Decompose $\underline{m} := \{1, 2, \dots, m\} = A \cup B$ into two disjoint sets.

A coloring C of $\underline{m}$ induces colorings of A, B so determines fermionic indices A_f, B_f .

Lemma 5.8. *Let D, E two words in A and B respectively. Then*

$$(33) \quad \epsilon_C(DE) = \epsilon_C(AB)\epsilon_C(D)\epsilon_C(E).$$

Proof. When we reorder the word DE we can first reorder D, E separately, getting the sign $\epsilon_C(D)\epsilon_C(E)$ and then AB getting the sign $\epsilon_C(AB)$.

Thus Formula (33). $\square$

Remark 5.9. For each $j \in B_f$ let $|j|$ the number of $i \in A_f$ with $i > j$. For each $i \in A_f$ let $|i|$ be the number of $j \in B_f$ with $i > j$.

$$(34) \quad \epsilon_C(AB) = \prod_{j \in B_f} (-1)^{|j|} = \prod_{i \in A_f} (-1)^{|i|}.$$

Lemma 5.10. *Let A, B and C as before. Consider a permutation σ of A , an index $i \in A$ and a string E permutation of B .*

Let $\mathbf{c} := (i, E)$ a cycle. Then if $C(i) \neq C(i, E)$ change the color of i and call the new coloring C' . Otherwise $C = C'$. We have:

$$(35) \quad \tau_C(\sigma \circ \mathbf{c}) = \gamma \tau_{C'}(\sigma)|_{z_i \mapsto \tau_C((i, E))}$$

either $\gamma = \pm 1$ a sign dependent only on A, B, i, C and NOT on σ .

Proof. Let us denote as in page 16, for all h , $z_h = y_h$ if $C(h) = 0$ and $z_h = x_h$ if $C(h) = 1$.

Write $\sigma = c_1 \circ \dots \circ c_r \circ (\dots, i)$ a p -string with support a string F, i and then

$$\sigma \circ c = c_1 \circ \dots \circ c_r \circ (\dots, i, E)$$

By Formula (4) $\tau_C(\sigma \circ c)$ is the trace monomial obtained by the substitutions $h \mapsto z_h$ in the p -string of support F, i, E obtaining a trace monomial M times the sign $\epsilon_C(F, i, E)$.

The support of F is $A \setminus \{i\}$.

Now $\tau_{C'}(\sigma)$ is the trace monomial obtained by the substitutions $h \mapsto z_h$ in the p -string of support F, i times the sign $\epsilon_{C'}(F, i)$.

By the hypotheses on C' in $\tau_{C'}(\sigma)$ we can perform the substitution $\tau_{C'}(\sigma)|_{z_i \mapsto \tau_C((i, E))}$ obtaining the same trace monomial M of $\tau_C(\sigma \circ c)$ times the sign $\epsilon_{C'}(F, i)\epsilon_C(i, E)$ hence we have that

$$\gamma = \epsilon_C(F, i, E)\epsilon_{C'}(F, i)\epsilon_C(i, E)$$

By Lemma 5.8 and Formula (33) for $D = F, i$.

$$(36) \quad \epsilon_C(F, i; E) = \epsilon_C(AB)\epsilon_C(F, i)\epsilon_C(E)$$

Hence

$$\gamma = \epsilon_C(AB)\epsilon_C(E)\epsilon_C(i, E) \boxed{\epsilon_C(F, i)\epsilon_{C'}(F, i)}.$$

If $C = C'$ we have $\epsilon_C(F, i)\epsilon_{C'}(F, i)$ otherwise in one of these two factors the index i is bosonic (say for instance for the color C) while in the other it is fermionic.

Then $\epsilon_C(F, i) = \epsilon_C(F)$ while $\epsilon_{C'}(F, i) = (-1)^{|i|}\epsilon_{C'}(F)$ with $|i|$ the number of indices j in $A \setminus \{i\}$ with $j > i$.

Therefore, since on the support $A \setminus \{i\}$ of F the two signs coincide, we have:

$$\gamma = \begin{cases} \epsilon_C(AB)\epsilon_C(E)\epsilon_C(i, E) & \text{if } C = C' \\ (-1)^{|i|}\epsilon_C(AB)\epsilon_C(E)\epsilon_C(i, E) & \text{if } C \neq C' \end{cases}$$

□

In order to proceed we need, for $p \geq m$:

Lemma 5.11. *Every permutation γ in S_p can be written as a product $\gamma = \alpha \circ \beta$ where $\alpha \in S_m$ and, in each cycle of β , there is at most 1 element in $1, 2, \dots, m$.*

Proof. It is enough to do it when γ is a cycle.

Observe that, for $a, b, c, \dots$ numbers $\leq m$ and $A, B, C, \dots$ strings of numbers $> m$, we have:

$$(37) \quad \gamma = (a \ A \ b \ B \ c \ C \ \dots \ e \ E) = (a \ b \ c \ \dots)(A \ a)(B \ b)(C \ c) \dots (E \ e).$$

□

Example 5.12. If $A = 7, 5, 4$; $B = 6, 3$; $a = 1$, $b = 2$,

$$(1, 7, 5, 4, 2, 6, 3) = (1, 2)(7, 5, 4, 1)(6, 3, 2).$$

Recall on page 6 we have described the free superalgebra with trace $\mathcal{F}_T = F\langle X, Y \rangle \otimes_A \mathcal{T}$.

We are now ready to prove the second fundamental theorem, we use the notations $\mathcal{T}_n$ and $\mathcal{E}_n$ as in Theorem 4.3 for the algebra of invariants of both bosonic and fermionic and the algebra of equivariant maps of both bosonic and fermionic matrices.

Theorem 5.13. *[SFT] i) The T -ideal of trace relations, kernel of the evaluation map of $\mathcal{T}$ into $\mathcal{T}_n$ is generated (as a T -ideal) by the trace $n+1$ trace relations $T_{e,f}$ with $e+f = n+1$*

ii) The T -ideal of (trace) identities of $n \times n$ matrices, is generated (as a T -ideal) by the $n+1$ Cayley Hamilton identities $CH_{e,f}(x, y)$, $e+f = n+1$, and $t(1) = n$.

Proof. i) From the Aronhold method every relation is a sum of multihomogeneous relations which can be fully polarized to multilinear trace relations.

The original relation is then obtained up to a multiplicative integer factor by the process of restitution that is making the groups of variables all equal.

It is thus sufficient to prove that a multilinear trace relation (resp. identity) (in the first n variables) is in the T -ideal generated by $T_{e,f}$ (resp. $CH_{e,f}$).

Let us first look at trace relations. By Proposition 3.11 (4) Lemma 4.5 and the definition of the evaluation in matrices, the multilinear trace identities of degrees a, b in the space $\mathcal{T}(a, b)$ of the trace algebra $\mathcal{T}$ are the image under the encoding map $\tau_{C_{a,b}}$ of the kernel $K_{m,n}$ of the map of the group algebra $\mathbb{C}[S_m]$ in the space of endomorphisms of $(\mathbb{C}^n)^{\otimes m}$.

By the classical theory this kernel is the two sided ideal of $\mathbb{C}[S_m]$ generated by the antisymmetrizer $A_{n+1} \in \mathbb{C}[S_{n+1}]$.

We thus have to decode this Kernel as elements of $\mathcal{T}(a, b)$ under the encoding map.

The space $K_{m,n}$ is linearly spanned by elements $\tau(\sum_{\sigma \in S_{n+1}} \epsilon_\sigma \sigma)\beta$, $\tau, \beta \in S_{m+1}$. Hence it is enough to look at the relations of the type

$$R = \tau_{C_{a,b}}(\tau(\sum_{\sigma \in S_{n+1}} \epsilon_\sigma \sigma)\beta), \quad \tau, \beta \in S_m.$$

We write such a relation as $\tau_{C_{a,b}}(\tau(\sum_{\sigma \in S_{n+1}} \epsilon_\sigma \sigma)\tau^{-1}\tau\beta)$.

We set $A := \tau(\{1, 2, \dots, n+1\})$ and B its complement in $\underline{m}$.

By Lemma 5.11 we can write $\tau\beta = \alpha\theta$ with $\alpha \in S_A$ and in each cycle of θ there is at most 1 element in A . We have

$$\tau(\sum_{\sigma \in S_{n+1}} \epsilon_\sigma \sigma)\tau^{-1}\alpha = \pm \tau(\sum_{\sigma \in S_{n+1}} \epsilon_\sigma \sigma)\tau^{-1}$$

so we are reduced to study a relation

$$\tau_{C_{e,f}}(\tau(\sum_{\sigma \in S_{n+1}} \epsilon_\sigma \sigma)\tau^{-1}\theta)$$

in each cycle of θ there is at most 1 element in A .

We apply in iteration Lemma 5.10 and see that R is deduced from the relation $\tau_C(\tau(\sum_{\sigma \in S_{n+1}} \epsilon_\sigma \sigma)\tau^{-1})$, for some color computed recursively from the color $C_{a,b}$ restricted to A , by a sequence of substitutions of the variables corresponding to the indices in A of the cycles of θ .

Since $\tau_C(\tau(\sum_{\sigma \in S_{n+1}} \epsilon_\sigma \sigma)\tau^{-1})$ is obtained from one of the trace relations of Formula (29) by a substitution of variables hence i) is proved.

ii) Consider now a multilinear relation R in $\mathcal{F}_T$ which we may assume to be in the variables $y_1, \dots, y_a, x_1, \dots, x_b$.

Then the formula $tr(Rx_{b+1})$ is a multilinear trace relation R in $\mathcal{T}$ in the variables $y_1, \dots, y_a, x_1, \dots, x_b, x_{b+1}$ so, from part i), it is a linear combination of terms of types

$$(38) \quad T = T_{e,f}(M_1, \dots, M_e, N_1, \dots, N_f, N_{f+1})tr(P_1) \cdots tr(P_r)$$

with the M_i monomials of degree 0, the N_i monomials of degree 1 and the P_i monomials of arbitrary degree.

We can write each $T = tr(Ux_{b+1})$ for an element $U \in \mathcal{F}_T$ vanishing on $n \times n$ matrices and R is then a linear combination of these elements U .

The Formula for U depends upon where x_{b+1} appears in the Formula (38).

If it appears in one of the $P_i = Ax_{b+1}B$ then up to a sign $U = BA$ times the element T with P_i omitted.

Otherwise up to some reordering, cf. 5.6 and sign we may assume that we have $N_{f+1} = Ax_{b+1}B$ with A, B monomials or $M_e = Ax_{b+1}B$ with A, B monomials. Then we see that U is up to sign

$$U = B \cdot C_{e,f}(M_1, \dots, M_e, N_1, \dots, N_f)A \cdot \text{tr}(P_1) \cdots \text{tr}(P_r)$$

or

$$U = B \cdot C_{e-1,f+1}(M_1, \dots, M_{e-1}, N_1, \dots, N_f, N_{f+1})A \cdot \text{tr}(P_1) \cdots \text{tr}(P_r).$$

The proof is complete. $\square$

Both the first and second fundamental theorem for matrices are not as precise as the corresponding theorems for vectors and forms.

Several things are lacking; a description of a minimal set of generators for the trace invariants and a description of a minimal set of generators of the algebra $F_{T,n}\langle X, Y \rangle$ of equivariant maps from m -tuples of matrices to matrices is spanned as a module over the ring on invariants.

The results are identical to the classical results for which at the moment the best result is given by the following:

Theorem 5.14 (Razmyslov). *The invariants of $n \times n$ matrices are generated by the elements $\text{tr}(M)$ where M is a monomial of degree $\leq n^2$.*

The algebras $F_{T,n}\langle X, Y \rangle$ of equivariant maps from m -tuples of matrices to matrices is spanned as a module over the ring on invariants, by monomials of degree $< n^2$.

It is conjectured that the best estimate should be $< \binom{n}{2}$. Kuzmin has proved that this is a lower bound, [12].

6. SUPER CAYLEY–HAMILTON ALGEBRAS

The theory follows verbatim the classical theory but we repeat it for convenience of the reader.

Definition 6.1. A super n –Cayley–Hamilton algebra R is a trace super-algebra R satisfying $t(1) = n$ and all the identities $CH_{e,f}(x, y)$ of Formula (29).

By Lemma 3.10 its trace algebra satisfies the identities $T_{e,f}(x, y)$ of Formula (29).

The category of super n –Cayley–Hamilton algebras admits free algebras and an interpretation of the SFT is

Theorem 6.2. *The superalgebra $F_{T,n}\langle X, Y \rangle$ of equivariant polynomial maps from I -tuples of $n \times n$ bosonic and J -tuples of $n \times n$ fermionic matrices to $n \times n$ matrices, is the free algebra with trace modulo the T -ideal generated by the n^{th} Cayley Hamilton polynomials of Formula (5.3) and $t(1) = n$.*

$$(39) \quad \boxed{F_{T,n}\langle X, Y \rangle := \mathcal{F}_T\langle X, Y \rangle / \langle CH_{e,f}(x, y), e + f = n + 1; t(1) = n \rangle}.$$

6.3. n -dimensional superrepresentations. For a given $n \in \mathbb{N}$ and a supercommutative ring A by $M_n(A)$ we denote the ring of $n \times n$ matrices with coefficients in A . By a symbol $(a_{i,j})$ we denote a matrix with entries $a_{i,j} \in A$, for $i, j = 1, \dots, n$. We may distinguish matrices by the $\mathbb{Z}/(2)$ degree.

The construction $A \mapsto M_n(A)$ is a functor from the category $\mathcal{C}$ of commutative superalgebras to the category $\mathcal{R}$ of associative superalgebras.

To a map $f : A \rightarrow B$ is associated a map $M_n(f) : M_n(A) \rightarrow M_n(B)$ in the obvious way $M_n(f)((a_{i,j})) := (f(a_{i,j}))$.

Definition 6.4. By an n -dimensional superrepresentation of a superalgebra R we mean a homomorphism $f : R \rightarrow M_n(A)$ of superalgebras with A supercommutative.

It is easily seen that the set valued functor $A \mapsto \text{hom}_{\mathcal{R}}(R, M_n(A))$ is representable. That is:

Proposition 6.5. *There is a commutative superalgebra $T_n(R)$ and a natural isomorphism j_A of functors:*

$$\text{hom}_{\mathcal{R}}(R, M_n(A)) \xrightarrow{j_A} \text{hom}_{\mathcal{C}}(T_n(R), A), \quad j_A : f \mapsto \bar{f}$$

given by the commutative diagram $f = M_n(\bar{f}) \circ j_R$:

$$(40) \quad \begin{array}{ccc} R & \xrightarrow{j_R} & M_n(T_n(R)) \\ & \searrow f & \downarrow M_n(\bar{f}) \\ & & M_n(A) \end{array}$$

The map $j_R : R \rightarrow M_n(T_n(R))$ is called the *universal n -dimensional superrepresentation of R* or *the universal map into $n \times n$ matrices*.

Of course it is possible that R has no n -dimensional superrepresentations, in which case $T_n(R) = \{0\}$.

Remark 6.6. A completely tautological statement is that R can be embedded in the ring $M_n(A)$ of $n \times n$ matrices over a supercommutative ring A if and only if j_R is an embedding.

The projective linear group $P_n = GL(n, \mathbb{C})/\mathbb{C}^*$ acts by conjugation of superrepresentations and as in the classical case it acts on the universal object $M_n(T_n(R))$ and R maps to the P_n invariants.

Theorem 6.7. *If R is an n -Cayley-Hamilton algebra, then the universal map $j_R : R \rightarrow M_n(T_n(R))^{P_n}$ is an isomorphism inducing an isomorphism of the trace superalgebra of R with $T_n(R)^{P_n}$.*

Proof. For the construction of the universal map we present the superalgebra R as a quotient of the free algebra $R = F_{T,n}\langle X, Y \rangle / I$

$$(41) \quad \begin{array}{ccc} F_{T,n}\langle X, Y \rangle & \xrightarrow{j} & M_n(\mathbb{K}[t(M)]) \\ \pi \downarrow & & \downarrow \bar{\pi} \\ R & \xrightarrow{\bar{j}} & M_n(\mathbb{K}[t(M)]/\bar{U}) \end{array}$$

where $M_n(\bar{U})$ is the two sided ideal of $M_n(\mathbb{K}[t(M)])$ generated by I , then $T_n(R) = \mathbb{K}[t(M)]/\bar{U}$.

So the question is to prove that $F_{T,n}\langle X, Y \rangle \cap M_n(\bar{U}) = M_n(\bar{U})^{P_n} = I$.

In order to simplify notations let us denote

$$B := M_n(\mathbb{K}[t(M)]), \quad J := M_n(\bar{U})$$

we may also assume that the set of variables in the free algebra is infinite.

First $J = \sum_j Bu_jB$, $u_j \in I$. Each Bu_jB is a P_n submodule and since the group is linearly reductive we have $J^{P_n} = \sum_j (Bu_jB)^{P_n}$, $u_j \in I$.

So it is enough to prove that for every $u \in I$ of some degree we have $(BuB)^{P_n} \subset I$.

Take an element $c = \sum_h a_h u b_h \in (BuB)^{P_n}$ with a_h, b_h homogeneous.

As in [17] we then use a matrix variable x with entries not appearing in a_h, b_h, u and consider the element

$$tr(cx) = tr\left(\sum_h a_h u b_h x\right) = tr\left(\sum_h \pm b_h x a_h u\right) \in \mathbb{K}[t(M)]^{P_n}.$$

Since P_n is linearly reductive we have the Reynolds operator $\Re$ which projects to the invariants, commutes with trace and satisfies the identity $\Re(ru) = \Re(r)u$, $\forall r \in B$, $u \in B^{P_n}$ so

$$(42) \quad tr(cx) = tr\left(\sum_h \Re(a_h u b_h x)\right) = tr\left(\sum_h \pm \Re(b_h x a_h)u\right).$$

From this one can easily prove the main Theorem.

In fact now we have the invariant $\sum_h \pm \Re(b_h x a_h)$ which is linear in x so by the FFT 4.3 it is of the form $\sum_h A_h x B_h + \sum_k C_k \text{tr}(d_k x)$, with A_h, B_h, C_k, d_k invariants independent of x .

$$\begin{aligned}
 \text{tr}(cx) &= \text{tr}\left(\sum_h A_h x B_h u + \sum_k C_k \text{tr}(d_k x) u\right) = \\
 &= \text{tr}\left(\sum_h B_h u A_h x\right) + \text{tr}\left(\sum_k u C_k \text{tr}(d_k x)\right) \\
 &= \text{tr}\left(\sum_h B_h u A_h x + \sum_k \text{tr}(u C_k) d_k x\right) \\
 (43) \quad &\implies c = \sum_h B_h u A_h + \sum_k \text{tr}(u C_k) d_k \in I
 \end{aligned}$$

since I is an ideal closed under trace. □

REFERENCES

- [1] E. Aljadeff, A. Giambruno, C. Procesi, A. Regev. *Rings with polynomial identities and finite dimensional representations of algebras*, Amer. Math. Soc. Colloq. Publ., 66 American Mathematical Society, [Providence], RI, 2020, xii+630 pp. 3, 5
- [2] Amitsur, A. S.; Levitzki, J. Minimal identities for algebras. Proc. Amer. Math. Soc. 1 (1950), 449?463
- [3] Brešar, Matej; Procesi, Claudio; Špenko, Špela, Quasi-identities on matrices and the Cayley-Hamilton polynomial. Adv. Math. 280 (2015), 439?471. 1.1
- [4] De Concini, Corrado; Papi, Paolo; Procesi, Claudio, The adjoint representation inside the exterior algebra of a simple Lie algebra Adv. Math. 280 (2015), 21?46.
- [5] C. De Concini, C. Procesi, *The invariant theory of matrices* A.M.S. University Lecture Series v. 69, 151 pp. (2017).
- [6] Drensky, Vesselin, Computing with matrix invariants. Math. Balkanica (N.S.) 21 (2007), no. 1-2, 141–172. 1.1
- [7] Dynkin, E. B. Topological characteristics of homomorphisms of compact Lie groups.(Russian) Mat. Sb. (N.S.) 35(77) (1954), 129?173. 1.1
- [8] Giorgos Eleftheriou, Ziming Ji, Sameer Murthy Fermionic trace relations and supersymmetric indices at finite N, arXiv:2605.08085
- [9] Formanek, E.: The invariants of $n \times n$ matrices. Lect. Notes Math. 1278, 18–43 (1987)
- [10] Formanek, E.: The polynomial identities and invariants of $n \times n$ matrices. In: Regional Conference Series in Mathematics, vol. 78. American Math. Soc., Providence (1991) 1.1
- [11] B. Kostant, A theorem of Frobenius, a theorem of Amitsur-Levitzki and cohomology theory, J. Mathematics and Mechanics 7 (1958), no. 2, 237-264.

- [12] E.N. Kuzmin, On the Nagata-Higman theorem, in: Mathematical Structures-Computational Mathematics-Mathematical Modeling, Sofia, 1975, pp. 101–107, in Russian. [4.1](#), [4.7.1](#)
- [13] Lew, John S. The generalized Cayley-Hamilton theorem in n dimensions. Z. Angew. Math. Phys. 17 1966 pp. 650–653. [5.7](#)
- [14] C. Procesi, The invariants of $n \times n$ matrices, Bull. Amer. Math. Soc. 82 (1976), no. 6, pp. 891–892.
- [15] C. Procesi, The invariant theory of $n \times n$ matrices, Advances in Math. 19 (1976), pp. 306–381. [1](#)
- [16] Procesi, Claudio On the theorem of Amitsur-Levitzki.(English summary) Israel J. Math. 207 (2015), no. 1, 151–154. [1.1](#)
- [17] C. Procesi, A formal inverse to the Cayley Hamilton theorem. Journal of Algebra 107 1 (1987), pp. 63–74. [1](#), [6.3](#)
- [18] C. Procesi, *Lie Groups*. An approach through invariants and representations, pp. xxiv+596, Springer, Universitext, (2007) .
- [19] J. Razmyslov, Trace identities of matrix algebras via a field of characteristic zero, Math. USSR Izvestia (translation). 8 (1974), pp. 727–760. [1](#)
- [20] K. S. Sibirskii, Algebraic invariants for a set of matrices, Sib. Mat. Zh. 9 (1968), pp. 152–164.
- [21] Spencer, A. J. M, The theory of matrix polynomials and its application to the mechanics of isotropic continua. Arch. Rational Mech. Anal. 2 1958/1959 309–336. [1](#)
- [22] Spencer, A. J. M.; Rivlin, R. S. Finite integrity bases for five or fewer symmetric 3×3 matrices. Arch. Rational Mech. Anal. 2 1958/1959 435–446.
- [23] Spencer, A. J. M. The invariants of six symmetric 3×3 matrices. Arch. Rational Mech. Anal. 7 1961 64–77. Review PDF [1](#)
- [24] Spencer, A. J. M.; Rivlin, R. S. Further results in the theory of matrix polynomials. Arch. Rational Mech. Anal. 4 1960 214–230 (1960). [1](#)
- [25] Spencer, A.J.M., Theory of invariants A.C. Eringen (Ed.), Continuum Physics, vol. I, Academic Press, San Diego (1971), pp. 240–353 [1](#)
- [26] Spencer, A. J. M., Ronald Rivlin and invariant theory. Internat. J. Engrg. Sci. 47 (2009), no. 11-12, 1066–1078. [1](#)

DIPARTIMENTO DI MATEMATICA, G. CASTELNUOVO, UNIVERSITÀ DI ROMA LA SAPIENZA,
PIAZZALE A. MORO, 00185, ROMA, ITALIA

Email address: `procesi@mat.uniroma1.it`