

Adiabatic Quantum Phase Estimation

Alexander Schmidhuber^{1,*} and Seth Lloyd^{2,†}

¹*Center for Theoretical Physics, Massachusetts Institute of Technology*

²*Department of Mechanical Engineering, Massachusetts Institute of Technology*

(Dated: May 22, 2026)

Quantum phase estimation (QPE) is a central algorithmic primitive which estimates eigenvalues of a Hamiltonian up to precision ε in Heisenberg-limited time $T = \Theta(1/\varepsilon)$. Standard gate-based implementations of QPE require deep controlled time-evolution circuits and are not native to analog hardware. Here, we present a simple adiabatic protocol for QPE that achieves (up to logarithmic factors) the optimal Heisenberg-limited scaling $T = O(\frac{1}{\varepsilon} \log(\delta^{-1}))$ in both the precision ε and failure probability δ . By encoding eigenvalues in the populations of computational basis states rather than complex phases, our approach is naturally robust against certain dephasing errors. The adiabatic protocol only requires the ability to couple a single ancilla qubit to the system Hamiltonian as well as pairwise couplings within the ancilla.

INTRODUCTION

Quantum Phase Estimation (QPE) is a fundamental quantum primitive that underlies the majority of existing quantum applications, including quantum simulation, integer factorization, quantum chemistry, and linear-systems methods [1–7]. For a Hamiltonian H_S with eigenpairs $H_S|\alpha\rangle = \lambda_\alpha|\alpha\rangle$ and an arbitrary initial state $|\psi\rangle_S = \sum_\alpha c_\alpha|\alpha\rangle$, idealized QPE implements the operation

$$\sum_\alpha c_\alpha|\alpha\rangle|0\rangle \mapsto \sum_\alpha c_\alpha|\alpha\rangle|\tilde{\lambda}_\alpha\rangle, \quad (1)$$

where $\tilde{\lambda}_\alpha$ is an m -bit estimate of λ_α . Gate-based QPE achieves additive accuracy $\varepsilon \simeq 2^{-m}$ in time $T = \Theta(2^m) = \Theta(1/\varepsilon)$, saturating the optimal tradeoff between time, energy, and precision known as the Heisenberg limit [8].

Standard gate-based implementations of QPE require a Quantum Fourier Transform and deep controlled powers of $e^{-iH_S t}$, typically through product formulas, block-encodings, or qubitization methods [9–11]. Such digital constructions are powerful but expensive and not native to analog platforms in which H_S is available as a physical interaction. This motivates an adiabatic implementation of QPE, which avoids Trotterization and is naturally robust to certain control and dephasing errors.

In this paper, we present a conceptually simple adiabatic protocol for QPE. Its time complexity is $T = \tilde{O}(\frac{1}{\varepsilon} \log \frac{1}{\delta})$, which is optimal (up to logarithmic factors) in both the accuracy ε and the failure probability δ [12]. Our protocol requires the ability to couple the system Hamiltonian to a single ancilla qubit at a time via the interaction $H_S \otimes Z_j$, as well as pairwise $Z_i Z_j$ interactions between ancilla qubits. To obtain logarithmic scaling in the failure probability δ , we devise an improved two-level adiabatic schedule that exponentially suppresses diabatic leakage. This schedule also naturally improves the adiabatic version of Grover’s algorithm [13], and we expect it to apply to a wider class of adiabatic applications.

Challenges. A core difficulty when implementing QPE adiabatically is retaining the Heisenberg limit. While an adiabatic implementation follows immediately from the universality proof of adiabatic quantum computation [14], such mappings are far from optimal due to unnatural history-state constructions. More broadly, adiabatic critical metrology is known to fall short of Heisenberg scaling in certain settings [15].

Standard adiabatic guarantees yield suboptimal scaling in both the accuracy ε and the failure probability δ . Consider a time-varying two-level Hamiltonian $h(t)$ with instantaneous ground and excited state $|g(t)\rangle$ and $|e(t)\rangle$, and associated energies $E_g(t)$ and $E_e(t)$. A commonly employed¹ version of the “adiabatic approximation theorem” states that diabatic leakage is suppressed provided the Hamiltonian evolves slowly relative to the square of its spectral gap [20]. More precisely, the final state $|\psi(T)\rangle$ after adiabatic evolution is said to satisfy $|\langle\psi(T)|g(T)\rangle|^2 \geq 1 - \delta$ provided

$$\forall t : \frac{|\langle e(t)|\partial_t h(t)|g(t)\rangle|}{(E_e(t) - E_g(t))^2} \leq \sqrt{\delta}. \quad (2)$$

This condition results in two immediate issues. First, applying the bound (2) to a Hamiltonian resolving energies up to precision ε —which must necessarily have a spectral gap of order $O(\varepsilon)$ —yields a suboptimal runtime of $T = \Omega(\frac{1}{\varepsilon^2})$ for fixed evolution speed $\partial_t h(t)$, which is quadratically worse than the Heisenberg limit. Generic superadiabatic guarantees require even longer runtimes scaling with the cube of the inverse gap [16]. Second, the diabatic leakage (and thus the overall failure probability of the algorithm) decays only polynomially in the total interrogation time T_{tot} according to (2): Lowering δ by a factor of four requires halving the evolution speed

¹ While this simple criterion is frequently used and often sets the right scale, it is neither necessary nor sufficient (see e.g. [13, 16–19] and our later discussion).

and thus doubling the total interrogation time. In contrast, the failure probability of gate-based QPE decays exponentially in the number of gates.

Our algorithm resolves the above two obstacles by (i) using a locally varying schedule in the spirit of [13] to recover the Heisenberg-optimal scaling in ε and (ii) reducing the analysis of adiabatic QPE to a two-level system, for which we design an adiabatic schedule based on the theory of superadiabaticity [16, 23, 38, 39]. The schedule is obtained by reparametrizing the Roland–Cerf local-gap schedule with an endpoint-flat Denjoy–Carleman clock [13, 21, 22]. The use of ultradifferentiable regularity in adiabatic theory has important precedents: Nenciu proved exponential adiabatic estimates under Gevrey-type regularity, and Jung and Elgart–Hagedorn developed Gevrey switching theorems for compactly supported smooth interpolations [23–25]. Compared to these Gevrey-class clock approaches that achieve a δ -dependent overhead of $\tilde{O}(\log^q(\delta^{-1}))$ for $q > 1$, the Denjoy–Carleman clock we use achieves the near-optimal scaling $\tilde{O}(\log(\delta^{-1}))$.

ADIABATIC QUANTUM PHASE ESTIMATION

Let H_S be the Hamiltonian of interest acting on an n -qubit system Hilbert space $\mathcal{H}_S$ and assume, after shifting and rescaling, that $\text{spec}(H_S) \subset [0, 1)$. The runtimes throughout are measured in physical time defined by this normalization. Fix a target additive accuracy $\varepsilon > 0$ and set

$$m = \left\lceil \log_2 \frac{1}{\varepsilon} \right\rceil, \quad \nu = 2^{-m-1}. \quad (3)$$

Let $\mathcal{A}$ be an m -qubit ancilla register with qubits $a_1, \dots, a_m$, initialized in $|+\rangle^{\otimes m}$ with $|+\rangle = (|0\rangle + |1\rangle)/\sqrt{2}$. Define the counting operators

$$B_{j-1} \equiv \sum_{\ell=1}^{j-1} 2^{-\ell} n_\ell, \quad n_\ell = \frac{\mathbb{I} - Z_{a_\ell}}{2}, \quad (4)$$

and the *midpoint operator* at stage j as

$$\mu_j \equiv B_{j-1} + 2^{-j} \quad (5)$$

acting on the ancilla register. Any computational basis string $b \in \{0, 1\}^m$ is an eigenstate of B_m with eigenvalue $\text{val}(b) = \sum_{\ell=1}^m 2^{-\ell} b_\ell$. We use the center-valued estimate

$$\hat{\lambda}(b) = \text{val}(b) + 2^{-m-1} \quad (6)$$

to associate a number to a computational basis state.

Algorithm. Our adiabatic algorithm consists of m sequential stages, each resolving one bit of the eigenvalues of H_S . At stage j we evolve under the Hamiltonian

$$H^{(j)}(s) = -(1-s)X_{a_j} + s(H_S - \mu_j)Z_{a_j}. \quad (7)$$

The protocol is:

1. Initialize the system in an arbitrary state $|\psi\rangle_S = \sum_i c_i |i\rangle$ and the ancilla in $|+\rangle_{\mathcal{A}}^{\otimes m}$.
2. For $j = 1, 2, \dots, m$: evolve $s : 0 \rightarrow 1$ under $H^{(j)}(s)$ using the schedule $s(t)$ specified in Eq. (17).
3. (Optional) Measure $\mathcal{A}$ in the computational basis to obtain a bit string b and output the center estimate $\hat{\lambda}(b)$ from Eq. (6).

Before proving correctness and efficiency of this protocol, we highlight several convenient properties. First, note that for all j and s , $[H^{(j)}(s), H_S \otimes \mathbb{I}_{\mathcal{A}}] = 0$, so the adiabatic evolution is block-diagonal in the eigenbasis of H_S . In particular, degenerate eigensectors of H_S are preserved. Moreover, for $\ell < j$, the ancilla qubit a_ℓ appears in $H^{(j)}(s)$ only through Z_{a_ℓ} terms, hence if it is set to a bit $\in \{0, 1\}$ it remains conserved for all $j > \ell$. Thus the entire adiabatic evolution is block-diagonal across superpositions of energy eigenstates and across already-decided prefixes on the ancilla qubits $a_1, \dots, a_m$, and is thus described by an effective two-level Hamiltonian.

Two-level reduction. Fix a system eigenstate $|\alpha\rangle$ with eigenvalue $\lambda \equiv \lambda_\alpha$, and fix a computational-basis prefix $x \in \{0, 1\}^{j-1}$. On the joint subspace $|\alpha\rangle_S \otimes |x\rangle_{a_{<j}} \otimes \mathbb{C}_{a_j}^2$, the midpoint μ_j becomes a scalar $\mu_j(x)$, and $H^{(j)}(s)$ acts nontrivially only on a_j as

$$h_\Delta(s) = -(1-s)X + s\Delta Z, \quad \Delta \equiv \lambda - \mu_j(x). \quad (8)$$

At $s = 0$ the unique ground state is $|+\rangle$. At $s = 1$, the ground state is $|1\rangle$ if $\Delta > 0$ and $|0\rangle$ if $\Delta < 0$. Thus adiabatic evolution of the ancilla from $s = 0$ to $s = 1$ implements a threshold test: the sign of Δ determines whether the ground-state branch ends in $|1\rangle$ or $|0\rangle$. Repeating this threshold test for all m stages coherently entangles the eigenvector $|\alpha\rangle$ with an ε -approximation to the eigenvalue λ_α . The above two-level reduction greatly simplifies the analysis of our algorithm and allows us to devise an explicit schedule that near-exponentially suppresses diabatic leakage.

For small Δ , there seems to be an apparent gap closing in Eq. (8). This is avoided by the fact that our algorithm only has to resolve eigenvalues up to accuracy $\varepsilon \approx 2\nu$. More specifically, our analysis distinguishes between (i) *resolved branches* $|\Delta| > \nu$, where the sign of Δ must be determined reliably, and (ii) *unresolved branches* $|\Delta| \leq \nu$, where the eigenvalue is effectively indistinguishable from the midpoint at this resolution. We only need to bound diabatic leakage in the resolved case: in the unresolved regime, both bit-choices remain valid ε -estimates of the true eigenvalue. The following two sections describe our schedule $s(t)$ and make the above error analysis in the resolved and unresolved case concrete.

ADIABATIC SCHEDULE

To guarantee adiabatic evolution, we require a schedule $s(t)$ that evolves slowly relative to the spectral gap of the two-level Hamiltonian (8). This gap is

$$E_e(s) - E_g(s) = 2r_\Delta(s), \quad (9)$$

where $r_x(s) = \sqrt{(1-s)^2 + s^2 x^2}$ for a general placeholder variable x . To get around the suboptimal scaling in ε implied by (2), we employ a locally varying adiabatic schedule. The simplest candidate is the Roland-Cerf schedule, originally described in [13] to recover Grover's quadratic speedup adiabatically.

Roland-Cerf schedule. The Roland-Cerf schedule is given by (visualized in Fig. 1)

$$\dot{s}(t) = 4\eta r_\nu(s(t))^2 = 4\eta ((1-s)^2 + \nu^2 s^2), \quad (10)$$

chosen such that the velocity approximately matches the square of the spectral gap of h_Δ . The free parameter η is chosen later to guarantee adiabaticity and will only depend on the overall failure probability δ . The duration of a single stage is

$$T_{\text{stage}} = \int_0^1 \frac{ds}{4\eta r_\nu(s)^2} = \frac{\pi}{8\eta\nu} = O(\varepsilon^{-1}). \quad (11)$$

This yields an overall runtime of $T_{\text{tot}} = mT_{\text{stage}} = \Theta(\varepsilon^{-1} \log \varepsilon^{-1})$, which is already Heisenberg-optimal in ε up to a logarithmic factor. However, the Roland-Cerf schedule suppresses diabatic leakage only polynomially (cf. Figure 2): it is easy to verify that the total runtime scales with the failure probability δ as $T_{\text{tot}} = \Theta(\delta^{-1/2})$, exponentially worse than the optimal bound of $T_{\text{tot}} = \Theta(\log \delta^{-1})$ [12] achieved by gate-based QPE. This is because the speed (10) and its derivatives are large at the endpoints of the schedule (cf. Fig. 1), causing amplitude leakages into the excited state of order $O(1/T)$.

Superadiabatic Roland-Cerf schedule. To obtain exponentially stronger bounds on the diabatic leakage, we instead reparametrize the Roland-Cerf schedule [13] with a non-quasianalytic Denjoy–Carleman function class clock [21, 22]. Define the Roland–Cerf action coordinate

$$\Theta_\nu(s) \equiv \int_0^s \frac{dz}{r_\nu(z)^2}, \quad r_\nu(s) = \sqrt{(1-s)^2 + \nu^2 s^2}. \quad (12)$$

A direct evaluation gives

$$\Theta_\nu(s) = \frac{1}{\nu} \left[\arctan\left(\frac{(1+\nu^2)s-1}{\nu}\right) + \arctan\left(\frac{1}{\nu}\right) \right]. \quad (13)$$

The ordinary Roland–Cerf schedule is constant-speed motion in the coordinate $\Theta_\nu(s)$. Let $q > 1$ be fixed and




FIG. 1. A visualization of three different schedule profiles: (a) The linear ramp profile $s(t) = t/T$. (b) The Roland-Cerf schedule (10) mirroring the behavior of the spectral gap. (c) The Roland-Cerf schedule reparametrized by a DC clock, which we call superadiabatic RC. Also shown is the behavior of the DC clock function, whose endpoint derivatives vanish and which thereby near-exponentially suppresses diabatic transitions.

let $F_q : [0, 1] \rightarrow [0, 1]$ be monotone, with $F_q(u) = 0$ near 0, $F_q(u) = 1$ near 1, and

$$\|F_q^{(n)}\|_\infty \leq C_F R_F^n n! [\log(n+e)]^{qn} \quad (n \geq 1). \quad (14)$$

The existence of such Denjoy–Carleman clock functions is proved constructively in Lemma 2, and visualized in Fig. 1. For a dimensionless adiabatic parameter $A = \eta^{-1} > 0$, define the duration of one stage by

$$T_{\text{stage}}(A) = \frac{A}{4} \Theta_\nu(1) = \frac{\pi A}{8\nu}, \quad (15)$$

and define the schedule implicitly by

$$\Theta_\nu(s(t)) = \Theta_\nu(1) F_q\left(\frac{t}{T_{\text{stage}}(A)}\right) \quad (16)$$

for $0 \leq t \leq T_{\text{stage}}(A)$. Differentiating (16) gives

$$\dot{s}(t) = \frac{4}{A} r_\nu(s(t))^2 F_q'\left(\frac{t}{T_{\text{stage}}(A)}\right). \quad (17)$$

Thus the schedule retains the Roland–Cerf local-gap factor $r_\nu(s)^2$, but the clock is constant near both endpoints. Equivalently, all positive time derivatives of the Hamiltonian vanish at the beginning and end of each stage, which removes the boundary terms that would otherwise appear in the high-order adiabatic expansion. These boundary cancellations allow us to prove near-exponentially small bounds on the diabatic leakage, significantly stronger than generically available bounds [16], as we show now.

Diabatic leakage for superadiabatic RC. Consider the two-level Hamiltonian (8) for a resolved branch $|\Delta| > \nu$. Let $P_\Delta(s)$ denote the instantaneous ground-state projector of $h_\Delta(s)$. The following theorem bounds transitions from the initial ground-state sector to the final excited sector.

Theorem 1 (Near-exponentially small diabatic leakage). *Fix $q > 1$ and let $\mathcal{V}_{\Delta,A}^{\text{stage}}$ be the single-stage two-level adiabatic evolution generated by the Hamiltonian (8) together with the superadiabatic RC schedule (16). There exist constants $C, c > 0$, depending only on q , such that for every $\nu \in (0, 1/2]$, every resolved detuning $1 \geq |\Delta| > \nu$, and every $A \geq 1$,*

$$\|(I - P_\Delta(1))\mathcal{V}_{\Delta,A}^{\text{stage}}P_\Delta(0)\| \leq C \exp\left[-c \frac{A}{[\log(e + A)]^q}\right]. \quad (18)$$

The proof is given in Appendix A.4. The main technical input is the weighted superadiabatic estimate, Theorem 3, applied to the normalized two-level Hamiltonian in the Roland–Cerf action coordinate.




FIG. 2. Numerical computation of the diabatic leakage via exact diagonalization of the two-level Hamiltonian (8). Four different schedules are considered: linear, Roland–Cerf, superadiabatic Roland–Cerf, and the DC clock function, which can be interpreted as the linear schedule reparametrized by the DC clock. While the linear and Roland–Cerf schedules yield a power-law decay, the reparametrized schedules suppress diabatic transitions near-exponentially. Both RC-type schedules display Stückelberg oscillations arising from coherent interference between nonadiabatic transitions.

We note that the same proof gives a high-confidence version of local adiabatic Grover search [13], as we show in Appendix A.7. Indeed, after symmetry reduction the Grover Hamiltonian is an affine two-level avoided crossing with minimum gap $\sqrt{M/N}$, and the same superadiabatic Roland–Cerf schedule (16) applies.

ERROR ANALYSIS

We now show that the protocol specified by the Hamiltonians (8) and the superadiabatic RC schedule (16) evolves adiabatically and accurately implements QPE. To formalize this claim, let

$$H_S = \sum_{\alpha} \lambda_{\alpha} \Pi_{\alpha}$$

be the spectral decomposition of H_S , with $\lambda_{\alpha} \in [0, 1)$. Because all stage Hamiltonians (7) commute with H_S , degenerate eigenspaces of H_S are preserved. For each eigenvalue λ_{α} , let

$$G^{\lambda_{\alpha}} = \sum_{\substack{b \in \{0,1\}^m \\ |\hat{\lambda}(b) - \lambda_{\alpha}| \leq \varepsilon}} |b\rangle\langle b|_{\mathcal{A}}$$

be the projector onto valid m -bit estimates of λ_{α} . Then the probability of failure for our algorithm is specified by the block diagonal spectral projector

$$\Pi_{\text{fail}} := I_{S \otimes \mathcal{A}} - \sum_{\alpha} \Pi_{\alpha} \otimes G^{\lambda_{\alpha}} \quad (19)$$

applied to the output state of our protocol

$$\mathcal{V}_A(|\psi\rangle_S |+\rangle_{\mathcal{A}}^{\otimes m}).$$

Here, $\mathcal{V}_A$ is the full adiabatic QPE algorithm, that is, sequential adiabatic evolution under Hamiltonians (7) and the single schedule (16) (with parameter A) for all stages $j = 1, 2, \dots, m$.

Theorem 2 (Correctness and efficiency). *Fix $q > 1$, target accuracy $\varepsilon \in (0, 1/2]$, and error probability $\delta \in (0, 1)$. Let $\mathcal{V}_A$ be the unitary adiabatic evolution under Hamiltonians (7) and the schedule (16) with parameter A . Then there exists a choice of A given by*

$$A = C_q L [\log(e + L)]^q, \quad L = 1 + \log \frac{C_0 m}{\sqrt{\delta}}, \quad (20)$$

with constants C_0, C_q sufficiently large depending only on q such that:

- For every n -qubit input state $|\psi\rangle_S$,

$$\|\Pi_{\text{fail}} \mathcal{V}_A(|\psi\rangle_S |+\rangle_{\mathcal{A}}^{\otimes m})\| \leq \sqrt{\delta}.$$

- The total interrogation time of $\mathcal{V}_A$ is

$$\begin{aligned} T_{\text{tot}} &= O\left(\frac{\log(1/\varepsilon)}{\varepsilon} \left[1 + \log \frac{1}{\delta} + \log \log \frac{e}{\varepsilon}\right] \times \left[\log\left(e + \log \frac{1}{\delta} + \log \log \frac{e}{\varepsilon}\right)\right]^q\right) \\ &= \tilde{O}\left(\frac{1}{\varepsilon} \log \frac{1}{\delta}\right). \end{aligned}$$

The proof is given in Appendix A.6. It combines Theorem 1 with a good-prefix invariant over the $m = \lceil \log_2(1/\varepsilon) \rceil$ midpoint comparisons and then sums the stage times.

DISCUSSION

This section compares adiabatic QPE to alternative implementations and discusses its practicality, robustness, and coherence. In general, adiabatic QPE is naturally applicable if the Hamiltonian H_S is available as a physical Hamiltonian in the lab via the interaction $H_S \otimes Z$, in which case adiabatic QPE avoids expensive compiling of controlled powers of $e^{-iH_S t}$ through product formulas, block encodings, or qubitization [9–11].

Robustness. Adiabatic QPE enjoys useful robustness properties. First, it is well-known that adiabatic quantum computation is naturally resistant to dephasing errors in the energy eigenbasis [26–30], which is an important source of noise in systems weakly coupled to the environment [31]. Second, unlike gate-based QPE which writes eigenvalues into relative Fourier phases before the final QFT, our protocol stores eigenvalue estimates in quantum state populations of the ancilla register. Since later stages interact with the ancilla register only through Z -diagonal terms, Z -basis dephasing of already written eigenvalue bits does not affect the final measurement statistics and commutes with all subsequent stages. Naturally, this observation does not guarantee robustness to other errors, such as relaxation, bit flips, dephasing of the active qubit in the wrong basis, and bath-induced transitions between instantaneous eigenstates.

Hardware assumptions. From a hardware perspective, the Hamiltonians (7) require only (i) a transverse field on the register, (ii) pairwise $Z_i Z_j$ interactions between register qubits, and (iii) a coupling $H_S Z_{a_j}$ between the system energy and a single ancilla qubit. This distinguishes AQPE from a non-adiabatic but analog von-Neumann pointer measurement of energy, which would couple H_S to a many-level pointer or clock degree of freedom through an interaction such as $H_S \otimes P$ [32]. Our protocol is closer in spirit to iterative or semiclassical forms of phase estimation, which can also use a single control qubit at a time [1, 33, 34]. The difference is that adiabatic QPE runs without intermediate measurements or classical feed-forward, and can be implemented coherently.

Previous adiabatic approaches. To the best of our knowledge, no prior adiabatic implementation of QPE achieves the near-optimal scaling $T = \tilde{O}(\frac{1}{\varepsilon} \log \frac{1}{\delta})$. Ref. [35] constructs an adiabatic implementation of the Quantum Fourier Transform (QFT) via circulant Hamiltonians and combines it with gate-based circuits to construct the QPE. This approach is hybrid and the precise runtime is not analyzed. Hen [36] constructs adiabatic implementations of several two-qubit gates and uses them to assemble an adiabatic gate-by-gate realization of the QFT and thus QPE, but this approach inherits the circuit structure of gate-based QPE, and more-

over requires relative phases to be maintained throughout the adiabatic gates. Ref. [37] gives an adiabatic phase-estimation-like construction for quantum counting based on Berry phases, achieving $O(\varepsilon^{-3/2})$ scaling, which is suboptimal relative to the $O(\varepsilon^{-1})$ scaling of gate-based quantum counting and highlights the difficulty of recovering optimal scaling adiabatically.

Coherence. Our adiabatic evolution is block diagonal in the eigenspaces of H_S , but as written it need not implement the ideal coherent map (1) with the same phases: different eigenvalue and prefix branches can accumulate different dynamical phases during the adiabatic evolution. (Geometric phases do not play a role because each two-level Hamiltonian (8) is real.) This does not affect applications of QPE if they rely only on the measurement statistics (such as Shor’s algorithm [4]), or if they only use QPE to apply classical functions $f(\lambda)$ to the eigenvalues where f does not vary strongly over ε —valid estimates (such as $f(\lambda) = \lambda^{-1}$ for linear system solvers [7]). In the latter case, simply apply the adiabatic protocol $\mathcal{V}$, apply a diagonal operation f on the eigenvalue register, and subsequently reverse the adiabatic evolution which uncomputes the dynamical phases.

If desired, a genuinely coherent implementation of QPE can be achieved by applying the adiabatic evolution $\mathcal{V}$, copying the eigenvalues from the ancilla register $\mathcal{A}$ to a separate register $\mathcal{R}$ with CNOT gates, and subsequently implementing the inverse adiabatic evolution $\mathcal{V}^\dagger$ on only the system $\mathcal{S}$ and the ancilla $\mathcal{A}$. This coherently uncomputes all dynamical phases while leaving eigenvectors entangled with corresponding eigenvalue estimates, *provided* each eigenvector $|\alpha\rangle$ is associated with a single eigenvalue estimate $\tilde{\lambda}_\alpha$. This is the case if no eigenvalue lies too close to a dyadic m -bit boundary, such that all eigenvalues are resolved. Alternatively, one can overresolve to $m + r$ bits and copy only an m -bit coarse label, thereby reducing the probability that the copied label distinguishes two valid boundary branches.

Imperfect implementations of the schedule. The superadiabatic schedule in (16) is a mathematically convenient schedule that should be viewed as a compact asymptotic way to package a family of finite-order endpoint-flat controls. For any fixed target δ , the leakage bound proof only uses integration-by-parts/superadiabatic orders up to $N = O(\log(1/\delta) + \log \log(1/\varepsilon))$. Thus an implementation need not realize an infinitely differentiable clock with infinite precision; it suffices to approximate the switch and its first N derivatives to the accuracy required by the target error budget.

In practice, one might want to replace the superadiabatic schedule with a schedule that is easier to implement with noisy waveform generators available in the lab. Many different simpler schedule choices exist that achieve nearly the same scaling: One approach is to re-parametrize the Roland-Cerf schedule by a smooth

Gevrey-class clock [23], which achieves total interrogation time $T_{\text{tot}} = \tilde{O}(\frac{1}{\varepsilon} \log^q \frac{1}{\delta})$ for any $q > 1$. Indeed, we have found that the simple Gevrey-class clock at $q = 2$, while asymptotically worse, performs better than the DC-clock for realistic choices of δ . If the initial state $|\psi\rangle_S$ is already close to an eigenstate, one could also directly use the simple Roland-Cerf schedule and exponentially amplify the success probability by repetition and majority vote.

CONCLUSION

We have presented an adiabatic analogue of Quantum Phase Estimation: instead of implementing controlled powers of $e^{-iH_S t}$ and a quantum Fourier transform, we implement m adiabatic midpoint comparisons that write an additive estimate directly into a register. Our protocol requires only (i) a transverse field on the register, (ii) pairwise $Z_i Z_j$ interactions between register qubits, and (iii) a coupling $H_S Z_j$ between the system energy and a single ancilla qubit at a time. Our algorithm is further robust against certain dephasing errors.

By reducing the analysis of adiabatic QPE to a two-level system, in which we reparametrize the Roland-Cerf schedule with a Denjoy-Carleman clock, we prove a total time complexity of $T = \tilde{O}(\varepsilon^{-1} \log(1/\delta))$, which is optimal (up to logarithmic factors) in all parameters.

This scheduling idea is not specific to quantum phase estimation. We expect it to apply to general adiabatic algorithms whose relevant dynamics reduce, exactly or perturbatively, to a family of uniformly controlled avoided crossings. The clearest example is local adiabatic Grover search: the usual Roland-Cerf schedule already gives the optimal $\Theta(\sqrt{N/M})$ search time, while reparameterizing the linear Roland-Cerf clock by a Denjoy-Carleman clock suppresses the single-run diabatic failure probability exponentially in the dimensionless adiabatic parameter. Similar remarks apply to fixed-point adiabatic search and adiabatic amplitude-amplification/state-conversion problems. Our result thus suggests a general route to recovering the complexity of optimal gate-based quantum algorithms adiabatically.

We thank Eddie Farhi, Alioscia Hama, Aram Harrow, and Rolando Somma for helpful discussions. A.S. is supported by a Google PhD Fellowship and the Simons Foundation (MP-SIP-00001553, AWH). S.L. is supported by, or in part by, the U. S. Army Research Laboratory and the U. S. Army Research Office under contract/grant number W911NF2310255, and by DoE under contract DE-SC0012704.

[†] slloyd@mit.edu

- [1] A. Y. Kitaev, Quantum measurements and the Abelian stabilizer problem (1995), [arXiv:quant-ph/9511026](https://arxiv.org/abs/quant-ph/9511026).
- [2] R. Cleve, A. Ekert, C. Macchiavello, and M. Mosca, Quantum algorithms revisited, *Proc. R. Soc. Lond. A* **454**, 339 (1998).
- [3] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
- [4] P. W. Shor, Algorithms for quantum computation: Discrete logarithms and factoring, in *Proceedings of the 35th Annual Symposium on Foundations of Computer Science* (IEEE, 1994) pp. 124–134.
- [5] D. S. Abrams and S. Lloyd, Quantum algorithm providing exponential speed increase for finding eigenvalues and eigenvectors, *Phys. Rev. Lett.* **83**, 5162 (1999).
- [6] A. Aspuru-Guzik, A. D. Dutoi, P. J. Love, and M. Head-Gordon, Simulated quantum computation of molecular energies, *Science* **309**, 1704 (2005).
- [7] A. W. Harrow, A. Hassidim, and S. Lloyd, Quantum algorithm for linear systems of equations, *Phys. Rev. Lett.* **103**, 150502 (2009).
- [8] V. Giovannetti, S. Lloyd, and L. Maccone, Quantum metrology, *Phys. Rev. Lett.* **96**, 010401 (2006).
- [9] S. Lloyd, Universal quantum simulators, *Science* **273**, 1073 (1996).
- [10] D. W. Berry, A. M. Childs, and R. Kothari, Hamiltonian simulation with nearly optimal dependence on all parameters, in *Proceedings of the 56th IEEE Symposium on Foundations of Computer Science* (IEEE, 2015) pp. 792–809.
- [11] G. H. Low and I. L. Chuang, Hamiltonian simulation by qubitization, *Quantum* **3**, 163 (2019).
- [12] N. S. Mande and R. de Wolf, Tight bounds for quantum phase estimation and related problems, in *31st Annual European Symposium on Algorithms (ESA 2023)*, LIPIcs, Vol. 274 (Schloss Dagstuhl, 2023) pp. 81:1–81:16.
- [13] J. Roland and N. J. Cerf, Quantum search by local adiabatic evolution, *Phys. Rev. A* **65**, 042308 (2002).
- [14] D. Aharonov, W. van Dam, J. Kempe, Z. Landau, S. Lloyd, and O. Regev, Adiabatic quantum computation is equivalent to standard quantum computation, *SIAM J. Comput.* **37**, 166 (2007).
- [15] K. Gietka, F. Metz, T. Keller, and J. Li, Adiabatic critical quantum metrology cannot reach the Heisenberg limit even when shortcuts to adiabaticity are applied, *Quantum* **5**, 489 (2021).
- [16] D. A. Lidar, A. T. Rezakhani, and A. Hama, Adiabatic approximation with exponential accuracy for many-body systems and quantum computation, *J. Math. Phys.* **50**, 102106 (2009).
- [17] E. Farhi, J. Goldstone, S. Gutmann, and M. Sipser, Quantum computation by adiabatic evolution (2000), [arXiv:quant-ph/0001106](https://arxiv.org/abs/quant-ph/0001106).
- [18] E. Farhi, J. Goldstone, S. Gutmann, J. Lapan, A. Lundgren, and D. Preda, A quantum adiabatic evolution algorithm applied to random instances of an NP-complete problem, *Science* **292**, 472 (2001).
- [19] J. Du, L. Hu, Y. Wang, J. Wu, M. Zhao, and D. Suter, Experimental study of the validity of quantitative conditions in the quantum adiabatic theorem, *Phys. Rev. Lett.* **101**, 060403 (2008).
- [20] A. Messiah, *Quantum Mechanics, Vol. II* (North-Holland Publishing Company, Amsterdam, 1962).

* alexsc@mit.edu

- [21] A. Rainer, Ultradifferentiable extension theorems: A survey, *Expo. Math.* **40**, 679 (2022).
- [22] A. Kriegl, P. W. Michor, and A. Rainer, The convenient setting for non-quasianalytic superadiabatic differentiable mappings, *J. Funct. Anal.* **256**, 3510 (2009).
- [23] G. Nenciu, Linear adiabatic theory. exponential estimates, *Commun. Math. Phys.* **152**, 479 (1993).
- [24] K. Jung, *The Adiabatic Theorem for Switching Processes with Gevrey Class Regularity*, Tech. Rep. Preprint 442 (Sonderforschungsbereich 288, Technische Universität Berlin, 2000) cERN-EXT-2000-159.
- [25] A. Elgart and G. A. Hagedorn, A note on the switching adiabatic theorem, *Journal of Mathematical Physics* **53**, 102202 (2012), [arXiv:1204.2318 \[math-ph\]](#).
- [26] A. M. Childs, E. Farhi, and J. Preskill, Robustness of adiabatic quantum computation, *Physical Review A* **65**, 012322 (2002).
- [27] M. S. Sarandy and D. A. Lidar, Adiabatic quantum computation in open systems, *Physical Review Letters* **95**, 250503 (2005).
- [28] T. Albash, S. Boixo, D. A. Lidar, and P. Zanardi, Quantum adiabatic markovian master equations, *New Journal of Physics* **14**, 123016 (2012).
- [29] T. Albash and D. A. Lidar, Decoherence in adiabatic quantum computation, *Physical Review A* **91**, 062320 (2015).
- [30] T. Albash and D. A. Lidar, Adiabatic quantum computation, *Rev. Mod. Phys.* **90**, 015002 (2018).
- [31] J. P. Paz and W. H. Zurek, Quantum limit of decoherence: Environment induced superselection of energy eigenstates, *Physical Review Letters* **82**, 5181 (1999).
- [32] J. von Neumann, *Mathematical Foundations of Quantum Mechanics* (Princeton University Press, 1955).
- [33] R. B. Griffiths and C.-S. Niu, Semiclassical fourier transform for quantum computation, *Physical Review Letters* **76**, 3228 (1996).
- [34] W.-C. Lin and C.-H. Wang, Analog quantum phase estimation with single-mode readout, *arXiv preprint arXiv:2506.15668* (2025).
- [35] B. T. Torosov and N. V. Vitanov, Design of quantum fourier transforms and quantum algorithms by using circulant hamiltonians, *Physical Review A—Atomic, Molecular, and Optical Physics* **80**, 022329 (2009).
- [36] I. Hen, Fourier-transforming with quantum annealers, *Frontiers in Physics* **2**, 44 (2014).
- [37] C. Zhang, Z. Wei, and A. Papageorgiou, Adiabatic quantum counting by geometric phase estimation, *Quantum Inf. Process.* **9**, 369 (2010).
- [38] G. A. Hagedorn and A. Joye, Elementary exponential error estimates for the adiabatic approximation, *J. Math. Anal. Appl.* **267**, 235 (2002).
- [39] S. Jansen, M.-B. Ruskai, and R. Seiler, Bounds for the adiabatic approximation with applications to quantum computation, *J. Math. Phys.* **48**, 102111 (2007).

Proof overview and conventions

This appendix proves the two estimates used in the main text. It is deliberately explicit to provide a detailed and self-contained proof of our main results. The argument has four ingredients. First, we construct a compactly supported flat clock in a non-quasianalytic Denjoy–Carleman class. Second, we prove a finite-dimensional superadiabatic estimate with endpoint cancellation. This is a standard superadiabatic construction in the spirit of Refs. [16, 23, 38, 39], with one modification: the residual is estimated in L^1 so that the constants do not depend on the length of the rescaled gap-time interval. Third, we apply these constructions to the normalized two-level comparator obtained from the Roland–Cerf action coordinate [13]. Finally, we compose the one-stage leakage estimate over the m midpoint-comparison stages.

Throughout the appendix, $q > 1$ is fixed and

$$M_n = n! [\log(n + e)]^{qn}, \quad n \geq 0. \quad (21)$$

Constants denoted by $C, R, c, C_0, C_1, \dots$ may change from line to line. They may depend on q and on the fixed clock, but never on $\nu, \Delta, A, m, \varepsilon$, or δ . All operator norms are spectral norms, and $\|X\|_1$ denotes the integral of $\|X(z)\|$ over the relevant z -interval. We use the usual big-O convention with $\tilde{O}(f(n))$ suppressing polylogarithmic factors in $f(n)$.

The flat Denjoy–Carleman clock

We first record two elementary estimates for the weights M_n , and then construct the endpoint-flat clock. The following construction is known as the infinite-convolution construction of a non-quasianalytic Denjoy–Carleman cutoff; see, e.g., Refs. [21, 22].

Lemma 1 (Product stability). *There is a constant $C_M \geq 1$, depending only on q , such that for all nonnegative integers a, b, r and $0 \leq \ell \leq r$,*

$$\binom{r}{\ell} M_{a+\ell} M_{b+r-\ell} \leq C_M^{a+b+r+1} M_{a+b+r}. \quad (22)$$

More generally, if $a_j, r_j \geq 0$, $\sum_{j=1}^p a_j = a$, and $\sum_{j=1}^p r_j = r$, then

$$\binom{r}{r_1, \dots, r_p} \prod_{j=1}^p M_{a_j+r_j} \leq C_M^{a+r+p} M_{a+r}. \quad (23)$$

After increasing C_M , one also has

$$M_{N+1} \leq C_M^N (N[\log(N+e)]^q)^N, \quad N \geq 1. \quad (24)$$

Proof. For the logarithmic factors we use

$$\log(a_j + r_j + e) \leq \log(a + r + e).$$

For the factorials,

$$\binom{r}{r_1, \dots, r_p} \prod_{j=1}^p (a_j + r_j)! = r! \prod_{j=1}^p \frac{(a_j + r_j)!}{r_j!}.$$

Since

$$\prod_{j=1}^p \frac{(a_j + r_j)!}{r_j!} \leq (a + r)^a \leq e^a \frac{(a + r)!}{r!},$$

we obtain the multinomial estimate up to an exponential factor in $a + r + p$. The two-factor estimate is the case $p = 2$. Finally, Stirling's bound and the elementary absorption of the extra factor $(N + 1)[\log(N + e)]^q$ into C^N give (24). $\square$

Lemma 2 (Flat Denjoy–Carleman switch). *For every $q > 1$, there exists a monotone $F_q \in C^\infty([0, 1])$ such that*

$$F_q(u) = 0 \quad (u \leq 1/4), \quad F_q(u) = 1 \quad (u \geq 3/4), \quad F_q'(u) \geq 0, \quad (25)$$

and constants $C_F, R_F > 0$ such that

$$\|F_q^{(n)}\|_\infty \leq C_F R_F^n M_n, \quad n \geq 1. \quad (26)$$

For $f_q = F_q'$, there are constants $C_f, R_f > 0$ such that

$$\|f_q^{(n)}\|_\infty \leq C_f R_f^n M_n, \quad n \geq 0. \quad (27)$$

Proof. Choose

$$a_j = \frac{a}{(j+2)[\log(j+e)]^q}, \quad j \geq 0, \quad (28)$$

with $a > 0$ small enough that $\sum_{j \geq 0} a_j \leq 1/2$. This is possible because $q > 1$. Let

$$\chi_j = a_j^{-1} \mathbf{1}_{[-a_j/2, a_j/2]}, \quad \rho_N = \chi_0 * \dots * \chi_N.$$

The sequence ρ_N converges, with all derivatives, to a compactly supported C^∞ probability density ρ , supported in an interval of length at most $1/2$. For fixed n and $N \geq n$, differentiating in the sense of distributions and placing the n derivatives on the first n box functions gives

$$\rho_N^{(n)} = \chi_0' * \dots * \chi_{n-1}' * \chi_n * \dots * \chi_N.$$

Since $\|\chi_j'\|_{\text{TV}} = 2/a_j$, $\|\chi_n\|_\infty = 1/a_n$, and convolution with probability densities does not increase the L^∞ norm,

$$\|\rho_N^{(n)}\|_\infty \leq \frac{2^n}{a_0 a_1 \dots a_n}. \quad (29)$$

Passing to the C^n -limit yields the same estimate for ρ . From the definition of a_j ,

$$\frac{2^n}{a_0 a_1 \cdots a_n} \leq C R^n (n+1)! [\log(n+e)]^{q(n+1)} \leq C' (R')^n M_n,$$

where the final extra factor is absorbed into the exponential constant.

Define

$$F_q(u) = \int_{-\infty}^{u-1/2} \rho(x) dx. \quad (30)$$

The support of ρ gives the plateaus in (25), and monotonicity follows from $\rho \geq 0$. Since $F_q^{(n)} = \rho^{(n-1)}(\cdot - 1/2)$ for $n \geq 1$, (26) follows. Finally, $f_q^{(n)} = F_q^{(n+1)}$, and $M_{n+1} \leq C^n M_n$ after increasing the exponential constant, giving (27). $\square$

A weighted superadiabatic estimate

The rescaled gap-time interval used below may have length of order ν^{-1} . The following product class and superadiabatic theorem are arranged so that no constant grows with that interval length.

Lemma 3 (Weighted L^∞/L^1 product calculus). *Let $I = [0, Z]$. A smooth matrix-valued function $X : I \rightarrow \mathbb{C}^{2 \times 2}$ has weighted order $\mu \geq 0$, with constants B, R , if*

$$\|X^{(r)}\|_\infty \leq B R^r M_{\mu+r} \quad (r \geq 0), \quad (31)$$

and, whenever $\mu + r \geq 1$,

$$\|X^{(r)}\|_1 \leq B R^r M_{\mu+r}. \quad (32)$$

No L^1 bound is required in the single case $\mu = r = 0$.

If X has weighted order μ and Y has weighted order η , then XY has weighted order $\mu + \eta$, with constants depending only on the previous constants and on q . The same holds for any fixed finite product.

Proof. Leibniz's rule gives

$$(XY)^{(r)} = \sum_{\ell=0}^r \binom{r}{\ell} X^{(\ell)} Y^{(r-\ell)}.$$

The L^∞ estimate follows from Lemma 1. For the L^1 estimate, assume $\mu + \eta + r \geq 1$. In each Leibniz term at least one of $\mu + \ell$ and $\eta + r - \ell$ is positive; put the L^1 norm on such a factor and the L^∞ norm on the other. Lemma 1 again controls the weights. Iteration gives the finite-product claim. $\square$

Theorem 3 (Weighted global superadiabatic estimate). *Let $K : [0, Z] \rightarrow \mathbb{C}^{2 \times 2}$ be a smooth self-adjoint Hamiltonian with two simple eigenvalues separated by a gap at least 2. Let $P(z)$ be the instantaneous ground-state projector and $Q(z) = \mathbb{I} - P(z)$. Assume*

$$K^{(n)}(0) = K^{(n)}(Z) = 0 \quad (n \geq 1), \quad (33)$$

and, for constants C_0, R_0 ,

$$\|K^{(n)}\|_\infty \leq C_0 R_0^n M_n, \quad \|K^{(n)}\|_1 \leq C_0 R_0^n M_n \quad (n \geq 1). \quad (34)$$

Let $U_A(z)$ solve

$$i \frac{d}{dz} U_A(z) = A K(z) U_A(z), \quad U_A(0) = \mathbb{I}. \quad (35)$$

Then there exist constants $C, c > 0$, depending only on C_0, R_0, q , such that for all $A \geq 1$,

$$\|(\mathbb{I} - P(Z)) U_A(Z) P(0)\| \leq C \exp \left[-c \frac{A}{[\log(e+A)]^q} \right]. \quad (36)$$

Proof. For an operator X , write

$$X^D = PXP + QXQ, \quad X^{OD} = PXQ + QXP.$$

On off-diagonal operators, $\text{ad}_K : X \mapsto [K, X]$ is invertible with norm at most $1/2$: if $K = E_g P + E_e Q$ and $E_e - E_g \geq 2$, then

$$[K, PXP] = -(E_e - E_g)PXP, \quad [K, QXP] = (E_e - E_g)QXP.$$

Bounds for P . We first show that P has weighted order 0:

$$\|P^{(n)}\|_\infty + \|P^{(n)}\|_1 \leq CR^n M_n, \quad n \geq 1, \quad (37)$$

and $\|P\|_\infty = 1$. Differentiating $P^2 = P$ and $[K, P] = 0$ n times gives

$$PP^{(n)} + P^{(n)}P - P^{(n)} = - \sum_{k=1}^{n-1} \binom{n}{k} P^{(k)} P^{(n-k)}, \quad (38)$$

$$[K, P^{(n)}] = - \sum_{k=1}^n \binom{n}{k} [K^{(k)}, P^{(n-k)}]. \quad (39)$$

The left side of (38) determines the diagonal part of $P^{(n)}$, while (39) and the gap determine the off-diagonal part. Induction on n , using Lemma 3, proves (37); in every L^1 estimate, at least one factor has positive weighted index, so $\|P\|_1$ is never used.

The same equations imply endpoint flatness. If $P^{(k)} = 0$ at an endpoint for $1 \leq k < n$, then (33) forces both the diagonal and off-diagonal parts of $P^{(n)}$ to vanish there. Hence

$$P^{(n)}(0) = P^{(n)}(Z) = 0, \quad n \geq 1. \quad (40)$$

The homological inverse. Let $Y = Y^{OD}$ have weighted order $\mu \geq 1$, and let $X = \mathfrak{S}(Y)$ be the unique solution of

$$i[K, X] = -Y, \quad X^D = 0. \quad (41)$$

Then X has the same weighted order:

$$\|X^{(r)}\|_\infty + \|X^{(r)}\|_1 \leq CR^r M_{\mu+r}, \quad r \geq 0. \quad (42)$$

For $r = 0$, this is the inverse-gap estimate. For $r \geq 1$, differentiating (41) gives

$$i[K, X^{(r)}] = -Y^{(r)} - i \sum_{\ell=1}^r \binom{r}{\ell} [K^{(\ell)}, X^{(r-\ell)}],$$

which controls the off-diagonal part of $X^{(r)}$. The diagonal part is controlled by differentiating $PXP + QXQ = 0$. All remaining terms contain either a positive derivative of P , a positive derivative of K , Y , or a lower derivative of X , so Lemma 3 closes the induction without any factor of Z . If all derivatives of Y vanish at the endpoints, then the same differentiated equations imply that all derivatives of X vanish at the endpoints.

Superadiabatic coefficients. Set $\Pi_0 = P$ and define

$$D_A(X) = A^{-1}X' + i[K, X].$$

We construct Π_n recursively. Suppose $\Pi_0, \dots, \Pi_{n-1}$ have been chosen so that, with

$$\Sigma_{n-1} = \sum_{j=0}^{n-1} A^{-j} \Pi_j,$$

the projection defect $\Sigma_{n-1}^2 - \Sigma_{n-1}$ and invariance defect $D_A(\Sigma_{n-1})$ vanish through order $A^{-(n-1)}$. Let G_n and F_n be the leading coefficients at order A^{-n} :

$$G_n = \sum_{a=1}^{n-1} \Pi_a \Pi_{n-a}, \quad F_n = \Pi'_{n-1}, \quad (43)$$

with the empty sum $G_1 = 0$.

The coefficient G_n is diagonal. Indeed, $\Sigma_{n-1}^2 - \Sigma_{n-1}$ commutes with Σ_{n-1} , and the coefficient of A^{-n} in this commutator is $[P, G_n]$. The coefficient F_n is off-diagonal. Since D_A is a derivation,

$$D_A(\Sigma_{n-1}^2 - \Sigma_{n-1}) = D_A(\Sigma_{n-1})\Sigma_{n-1} + \Sigma_{n-1}D_A(\Sigma_{n-1}) - D_A(\Sigma_{n-1}).$$

Taking the coefficient of A^{-n} gives

$$i[K, G_n] = F_n P + P F_n - F_n.$$

Because G_n is diagonal, it commutes with K , and therefore $F_n P + P F_n - F_n = 0$, which is equivalent to $F_n = F_n^{\text{OD}}$.

Define

$$D_n = -P G_n P + Q G_n Q, \quad O_n = \mathfrak{S}(F_n), \quad \Pi_n = D_n + O_n. \quad (44)$$

Then

$$P \Pi_n + \Pi_n P - \Pi_n = -G_n, \quad F_n + i[K, \Pi_n] = 0,$$

so both defects are cancelled through order A^{-n} .

We now record the bounds. The induction gives

$$\|\Pi_j^{(r)}\|_\infty + \|\Pi_j^{(r)}\|_1 \leq C^{j+r+1} R^{j+r} M_{j+r}, \quad j+r \geq 1. \quad (45)$$

For $j = 0$, this is (37). If the estimate holds up to $j = n-1$, then G_n has weighted order n by Lemma 3, $F_n = \Pi'_{n-1}$ has weighted order n , D_n has weighted order n , and the homological inverse gives the same order for O_n . Finite sums and the number of Leibniz terms are absorbed into the exponential constants.

Endpoint flatness is inherited by the same recursion. For $j = 0$, it is (40). For $j \geq 1$, the terms defining G_j contain lower positive-order coefficients, and $F_j = \Pi'_{j-1}$. By induction, these are flat at the endpoints; hence so are D_j , O_j , and Π_j . Therefore

$$\Pi_j^{(r)}(0) = \Pi_j^{(r)}(Z) = 0 \quad (j \geq 1, r \geq 0). \quad (46)$$

For $N \geq 1$, set

$$\Pi^{[N]} = \sum_{j=0}^N A^{-j} \Pi_j.$$

By (46),

$$\Pi^{[N]}(0) = P(0), \quad \Pi^{[N]}(Z) = P(Z). \quad (47)$$

The construction also gives the exact residual identity

$$\frac{d}{dz} \Pi^{[N]} + iA[K, \Pi^{[N]}] = A^{-N} \Pi'_N. \quad (48)$$

Thus, by (45) and (24),

$$\begin{aligned} \|A^{-N} \Pi'_N\|_1 &\leq A^{-N} C^{N+2} R^{N+1} M_{N+1} \\ &\leq C \left(\frac{C_1 N [\log(N+e)]^q}{A} \right)^N. \end{aligned} \quad (49)$$

Finally,

$$\frac{d}{dz} \left(U_A(z)^\dagger \Pi^{[N]}(z) U_A(z) \right) = U_A(z)^\dagger (A^{-N} \Pi'_N(z)) U_A(z).$$

Using (47) and integrating,

$$\|P(Z) U_A(Z) - U_A(Z) P(0)\| \leq \|A^{-N} \Pi'_N\|_1.$$

Multiplying by $\mathbb{I} - P(Z)$ on the left and $P(0)$ on the right gives the same bound for the leakage.

Choose

$$N = \left\lfloor a \frac{A}{[\log(e + A)]^q} \right\rfloor$$

with $a > 0$ small enough. For A larger than a q -dependent constant, (49) is bounded by

$$C \exp \left[-c \frac{A}{[\log(e + A)]^q} \right].$$

For bounded A , the leakage norm is at most one, and increasing C extends the estimate to every $A \geq 1$. $\square$

Proof of Theorem 1: resolved two-level leakage

We now apply Theorem 3 to the two-level comparator

$$h_\Delta(s) = -(1 - s)X + s\Delta Z$$

under the schedule (16). Fix a resolved detuning $1 \geq |\Delta| > \nu$, and write

$$d = |\Delta|, \quad \sigma = \text{sgn}(\Delta), \quad r_d(s) = \sqrt{(1 - s)^2 + d^2 s^2}.$$

Define the rescaled gap time

$$z(t) = \frac{1}{A} \int_0^t r_d(s(t')) dt'. \quad (50)$$

Then the two-level Schrödinger equation is

$$i \frac{d}{dz} U(z) = A K_\Delta(z) U(z), \quad K_\Delta(z) = \frac{h_\Delta(s(t(z)))}{r_d(s(t(z)))}. \quad (51)$$

The normalized Hamiltonian K_Δ has eigenvalues ± 1 , hence gap 2, and has the same instantaneous projectors as h_Δ .

Introduce $\phi \in [0, \pi/2]$ by

$$\cos \phi = \frac{1 - s}{r_d(s)}, \quad \sin \phi = \frac{ds}{r_d(s)}. \quad (52)$$

Then

$$K_\Delta = -\cos \phi X + \sigma \sin \phi Z \equiv G_\sigma(\phi). \quad (53)$$

Let

$$x = \frac{t}{T_{\text{stage}}(A)}, \quad f(x) = F'_q(x).$$

Using (17), (15), and (50), one obtains

$$\frac{d}{dz} = \mathcal{L}, \quad \mathcal{L} = a_d(\phi) \partial_x + b_{d,\nu}(\phi) f(x) \partial_\phi, \quad (54)$$

where

$$a_d(\phi) = \frac{8}{\pi} \left(\nu \cos \phi + \frac{\nu}{d} \sin \phi \right), \quad (55)$$

$$b_{d,\nu}(\phi) = 4(d \cos \phi + \sin \phi) \left(\cos^2 \phi + \frac{\nu^2}{d^2} \sin^2 \phi \right). \quad (56)$$

The inequalities $d \in [\nu, 1]$ and $\nu \in (0, 1/2]$ imply uniform analytic bounds

$$\|\partial_\phi^r a_d\|_\infty + \|\partial_\phi^r b_{d,\nu}\|_\infty + \|\partial_\phi^r G_\sigma\|_\infty \leq C R^r r! \leq C R^r M_r. \quad (57)$$

Also $a_d(\phi) > 0$.

Lemma 4 (Uniform derivative bounds for the normalized comparator). *There are constants $C, R > 0$, depending only on q and on the fixed clock F_q , such that for every $\nu \in (0, 1/2]$, every resolved $d \in [\nu, 1]$, and every $n \geq 1$,*

$$\|K_\Delta^{(n)}\|_\infty \leq CR^n M_n, \quad \|K_\Delta^{(n)}\|_1 \leq CR^n M_n, \quad (58)$$

where derivatives and the L^1 norm are taken with respect to z . Moreover,

$$K_\Delta^{(n)}(0) = K_\Delta^{(n)}(Z_\Delta) = 0, \quad n \geq 1, \quad (59)$$

where Z_Δ is the final value of $z(t)$.

Proof. Write

$$f(x) = F'_q(x), \quad \mathcal{L} = a_d(\phi)\partial_x + b_{d,\nu}(\phi)f(x)\partial_\phi,$$

and define

$$W_n = \mathcal{L}^n G_\sigma$$

as a function of the independent variables $(x, \phi) \in [0, 1] \times [0, \pi/2]$. Since $\mathcal{L}$ is the z -derivative along the trajectory, we have

$$K_\Delta^{(n)}(z) = W_n(x(z), \phi(z)).$$

We first prove the mixed derivative estimate

$$\|\partial_x^r \partial_\phi^s W_n\|_\infty \leq C^{n+r+s+1} R^{n+r+s} M_{n+r+s} \quad (60)$$

for all $n, r, s \geq 0$, uniformly in $\nu \in (0, 1/2]$, $d \in [\nu, 1]$, and $\sigma \in \{\pm 1\}$. Here the supremum is taken over $(x, \phi) \in [0, 1] \times [0, \pi/2]$.

The case $n = 0$ follows from the analyticity of

$$G_\sigma(\phi) = -\cos \phi X + \sigma \sin \phi Z.$$

Moreover, since $0 < \nu/d \leq 1$ and $0 < d \leq 1$, the coefficients a_d , $b_{d,\nu}$, and G_σ satisfy uniform analytic bounds: for suitable constants C_a, R_a ,

$$\|\partial_\phi^\ell a_d\|_\infty + \|\partial_\phi^\ell b_{d,\nu}\|_\infty + \|\partial_\phi^\ell G_\sigma\|_\infty \leq C_a R_a^\ell \ell! \leq C_a R_a^\ell M_\ell. \quad (61)$$

The clock derivative satisfies

$$\|f^{(\ell)}\|_\infty \leq C_f R_f^\ell M_\ell.$$

Assume (60) holds for W_n . Since

$$W_{n+1} = a_d(\phi)\partial_x W_n + b_{d,\nu}(\phi)f(x)\partial_\phi W_n,$$

Leibniz's rule gives two types of terms. For the first term,

$$\partial_x^r \partial_\phi^s (a_d(\phi)\partial_x W_n) = \sum_{\ell=0}^s \binom{s}{\ell} (\partial_\phi^\ell a_d) \partial_x^{r+1} \partial_\phi^{s-\ell} W_n.$$

Using (61), the induction hypothesis, and Lemma 1, this is bounded by

$$C^{n+r+s+2} R^{n+r+s+1} M_{n+r+s+1}.$$

For the second term,

$$\partial_x^r \partial_\phi^s (b_{d,\nu}(\phi)f(x)\partial_\phi W_n) = \sum_{\alpha=0}^r \sum_{\beta=0}^s \binom{r}{\alpha} \binom{s}{\beta} f^{(\alpha)}(x) (\partial_\phi^\beta b_{d,\nu})(\phi) \partial_x^{r-\alpha} \partial_\phi^{s-\beta+1} W_n.$$

The derivative bounds for f , the analytic bounds for $b_{d,\nu}$, the induction hypothesis, and Lemma 1 again give

$$C^{n+r+s+2} R^{n+r+s+1} M_{n+r+s+1}.$$

After increasing C and R , the induction closes. Taking $r = s = 0$ in (60) gives

$$\|K_\Delta^{(n)}\|_\infty \leq CR^n M_n.$$

We now prove the L^1 estimate. Parameterize the trajectory by $x \mapsto \phi(x)$. Since F_q is monotone, $s(x)$ is monotone, and hence $\phi(x)$ is monotone increasing from 0 to $\pi/2$. Thus

$$\int_0^1 \phi'(x) dx = \frac{\pi}{2}. \quad (62)$$

Along the trajectory,

$$\mathcal{L} = a_d(\phi(x))(\partial_x + \phi'(x)\partial_\phi) = a_d(\phi(x))\frac{d}{dx}, \quad dz = \frac{dx}{a_d(\phi(x))}.$$

Therefore, for $n \geq 1$,

$$\begin{aligned} \|K_\Delta^{(n)}\|_1 &= \int_0^{Z_\Delta} \|W_n(x(z), \phi(z))\| dz \\ &= \int_0^1 \left\| a_d(\phi(x)) \frac{d}{dx} W_{n-1}(x, \phi(x)) \right\| \frac{dx}{a_d(\phi(x))} \\ &= \int_0^1 \left\| \frac{d}{dx} W_{n-1}(x, \phi(x)) \right\| dx \\ &\leq \int_0^1 \|\partial_x W_{n-1}(x, \phi(x))\| dx + \int_0^1 \phi'(x) \|\partial_\phi W_{n-1}(x, \phi(x))\| dx \\ &\leq \|\partial_x W_{n-1}\|_\infty + \frac{\pi}{2} \|\partial_\phi W_{n-1}\|_\infty \\ &\leq CR^n M_n. \end{aligned} \quad (63)$$

In the last step we used (60) with $(n-1, r, s) = (n-1, 1, 0)$ and $(n-1, 0, 1)$. This estimate is uniform in ν and d ; in particular, the length Z_Δ of the z -interval never appears.

Finally, F_q is constant near $x = 0$ and near $x = 1$. Hence $s(x)$, $\phi(x)$, and therefore $K_\Delta(z)$, are constant on neighborhoods of both endpoints. All positive z -derivatives vanish there, so

$$K_\Delta^{(n)}(0) = K_\Delta^{(n)}(Z_\Delta) = 0, \quad n \geq 1.$$

□

Proof of Theorem 1. Lemma 4 verifies the hypotheses of Theorem 3 for $K_\Delta(z)$, uniformly over all $\nu \in (0, 1/2]$ and all resolved detunings $1 \geq |\Delta| > \nu$. Since K_Δ is obtained from h_Δ by division by the positive scalar $r_d(s(t(z)))$, their instantaneous spectral projectors agree. Reparameterizing time does not change the physical unitary. Therefore

$$\|(\mathbb{I} - P_\Delta(1))\mathcal{V}_{\Delta,A}^{\text{stage}} P_\Delta(0)\| \leq C \exp\left[-c \frac{A}{[\log(e+A)]^q}\right],$$

which is (18). □

Choosing the adiabatic parameter

Lemma 5 (Inverting the near-exponential bound). *Fix $q > 1$ and $c > 0$. There is a constant C_q such that, for every $L \geq 1$, the choice*

$$A = C_q L [\log(e+L)]^q \quad (64)$$

implies

$$\exp\left[-c \frac{A}{[\log(e+A)]^q}\right] \leq e^{-L}. \quad (65)$$

Proof. For a parameter $\kappa \geq 1$, set $A_\kappa = \kappa L[\log(e + L)]^q$. For all $L \geq 1$,

$$\log(e + A_\kappa) \leq B_q(1 + \log \kappa) \log(e + L)$$

with a constant B_q independent of L and κ . Hence

$$\frac{A_\kappa}{[\log(e + A_\kappa)]^q} \geq \frac{\kappa}{B_q^q(1 + \log \kappa)^q} L.$$

Since $\kappa/(1 + \log \kappa)^q \rightarrow \infty$, choose $\kappa = C_q$ large enough that the prefactor is at least $1/c$. $\square$

Let C_*, c_* be the constants from Theorem 1. Choose

$$L = 1 + \log \frac{C_0 m}{\sqrt{\delta}}, \quad A = C_q L[\log(e + L)]^q, \quad (66)$$

where C_q is chosen using Lemma 5 with $c = c_*$, and C_0 is large enough to absorb the prefactor C_* . Then the resolved-branch leakage amplitude of one stage obeys

$$\beta_{\text{stage}} \leq \frac{\sqrt{\delta}}{m}. \quad (67)$$

The corresponding stage time is

$$T_{\text{stage}}(A) = \frac{\pi A}{8\nu} = O\left(\frac{1}{\nu} L[\log(e + L)]^q\right). \quad (68)$$

Proof of Theorem 2: composition over QPE stages

We prove the measurement-success guarantee first for one fixed eigenvalue λ , and then sum over eigenspaces.

For a length- j prefix $b \in \{0, 1\}^j$, let

$$\text{val}_j(b) = \sum_{\ell=1}^j 2^{-\ell} b_\ell, \quad I_b = [\text{val}_j(b), \text{val}_j(b) + 2^{-j}]$$

be the associated dyadic interval. We call b good for λ if

$$\text{dist}(\lambda, I_b) \leq \nu. \quad (69)$$

Let G_j^λ be the projector onto the span of good length- j prefixes, tensored with the identity on unused qubits. The empty prefix is good, so $G_0^\lambda = \mathbb{I}$.

Consider stage j , conditioned on a good parent prefix $b \in \{0, 1\}^{j-1}$. Its midpoint is

$$\mu_j(b) = \text{val}_{j-1}(b) + 2^{-j}.$$

If $|\lambda - \mu_j(b)| \leq \nu$, then both children are good, since the midpoint is an endpoint of both child intervals. If $|\lambda - \mu_j(b)| > \nu$, then the child on the same side of $\mu_j(b)$ as λ is good. The opposite child is exactly the excited final state of the two-level comparator h_Δ , with $\Delta = \lambda - \mu_j(b)$, and therefore has transition amplitude at most $\sqrt{\delta}/m$ by (67).

This blockwise statement gives an operator bound for the full stage. Let U_j^λ be the stage- j unitary restricted to the eigenspace of H_S with eigenvalue λ , and define

$$\Gamma_{j-1}^\lambda = G_{j-1}^\lambda \otimes |+\rangle\langle +|_{a_j} \otimes \mathbb{I}_{a_{>j}}. \quad (70)$$

The fresh qubit a_j has not appeared in earlier stages, so the good component before stage j lies in the range of Γ_{j-1}^λ . Since the prefix blocks are orthogonal and the unresolved blocks have both children good,

$$\|(\mathbb{I} - G_j^\lambda)U_j^\lambda\Gamma_{j-1}^\lambda\| \leq \frac{\sqrt{\delta}}{m}. \quad (71)$$

Let

$$\psi_j^\lambda = U_j^\lambda \cdots U_1^\lambda |+\rangle^{\otimes m}, \quad e_j^\lambda = \|(\mathbb{I} - G_j^\lambda)\psi_j^\lambda\|.$$

Decompose ψ_{j-1}^λ into its good and bad parts. The good part is controlled by (71); the bad part has norm e_{j-1}^λ , and unitarity cannot increase it. Hence

$$e_j^\lambda \leq e_{j-1}^\lambda + \frac{\sqrt{\delta}}{m}.$$

Since $e_0^\lambda = 0$, iteration gives

$$e_m^\lambda \leq \sqrt{\delta}. \quad (72)$$

If a final prefix b is good, then $\text{dist}(\lambda, I_b) \leq \nu$ and $|I_b| = 2^{-m}$. Therefore the center estimate satisfies

$$|\hat{\lambda}(b) - \lambda| \leq \nu + \frac{2^{-m}}{2} = 2^{-m} \leq \varepsilon, \quad (73)$$

using $\nu = 2^{-m-1}$ and $m = \lceil \log_2(1/\varepsilon) \rceil$. Thus G_m^λ is supported on the valid-estimate projector G^λ from Eq. (19).

For a general input state, write

$$H_S = \sum_\alpha \lambda_\alpha \Pi_\alpha, \quad |\psi\rangle = \sum_\alpha |\psi_\alpha\rangle, \quad |\psi_\alpha\rangle = \Pi_\alpha |\psi\rangle.$$

All stage Hamiltonians commute with H_S , so the evolution is block diagonal in the eigenspaces of H_S . Consequently,

$$\begin{aligned} \left\| \Pi_{\text{fail}} \mathcal{V}_A \left(|\psi\rangle_S |+\rangle_{\mathcal{A}}^{\otimes m} \right) \right\|^2 &= \sum_\alpha \| |\psi_\alpha\rangle \|^2 \| (\mathbb{I} - G^{\lambda_\alpha}) \psi_m^{\lambda_\alpha} \|^2 \\ &\leq \delta. \end{aligned} \quad (74)$$

This proves the correctness claim of Theorem 2.

It remains to compute the total time. By (68),

$$T_{\text{tot}} = m T_{\text{stage}} = O\left(\frac{m}{\nu} L [\log(e + L)]^q\right), \quad L = 1 + \log \frac{C_0 m}{\sqrt{\delta}}.$$

Since

$$m = \Theta(\log(1/\varepsilon)), \quad \nu = \Theta(\varepsilon), \quad L = O\left(1 + \log \frac{1}{\delta} + \log \log \frac{e}{\varepsilon}\right),$$

this is exactly the runtime bound stated in Theorem 2.

Application to local adiabatic Grover search

We briefly indicate why the same argument gives the high-confidence Grover statement in the main text. Let M marked items be known, set $a = M/N$, and assume $0 < a \leq 1/2$. The standard local adiabatic search Hamiltonian is

$$H_G(s) = (1 - s)(\mathbb{I} - |u\rangle\langle u|) + s(\mathbb{I} - \Pi_M), \quad (75)$$

where $|u\rangle$ is the uniform state and Π_M projects onto the marked subspace [13]. Starting from $|u\rangle$, the evolution is exactly confined to the two-dimensional subspace spanned by

$$|w\rangle = \frac{\Pi_M |u\rangle}{\sqrt{a}}, \quad |r\rangle = \frac{(\mathbb{I} - \Pi_M) |u\rangle}{\sqrt{1 - a}}. \quad (76)$$

In this subspace, after subtracting the scalar $\mathbb{I}/2$ and applying an s -independent rotation of Pauli axes, the traceless Hamiltonian is

$$\bar{H}_a(s) = -\frac{1}{2} \left[2\sqrt{1 - a} \left(s - \frac{1}{2} \right) X + \sqrt{a} Z \right], \quad (77)$$

with gap

$$g_a(s) = \sqrt{a + 4(1-a) \left(s - \frac{1}{2}\right)^2}. \quad (78)$$

Thus local adiabatic Grover search is another affine two-level avoided crossing.

Define the Roland–Cerf coordinate

$$\Theta_a(s) = \int_0^s \frac{du}{g_a(u)^2}. \quad (79)$$

A direct integration gives

$$\Theta_a(1) = \frac{\arctan \sqrt{(1-a)/a}}{\sqrt{a(1-a)}} = \Theta\left(\sqrt{\frac{N}{M}}\right). \quad (80)$$

Choose the DC-clocked local schedule by

$$\Theta_a(s(t)) = \Theta_a(1)F_q(t/T), \quad T = A\Theta_a(1), \quad (81)$$

up to an immaterial constant rescaling of A . To verify the hypotheses of Theorem 3, introduce the Bloch angle

$$\theta(s) = \arctan\left(\frac{2\sqrt{1-a}(s - \frac{1}{2})}{\sqrt{a}}\right), \quad \theta_0 = \arctan\sqrt{\frac{1-a}{a}}. \quad (82)$$

Then

$$d\Theta_a = \frac{d\theta}{2\sqrt{a(1-a)}}, \quad \theta(t) = -\theta_0 + 2\theta_0 F_q(t/T), \quad (83)$$

and the normalized traceless Hamiltonian is a fixed-gap two-level path

$$K_a(\theta) = -\sin \theta X - \cos \theta Z \quad (84)$$

with eigenvalues ± 1 . In the corresponding gap-time variable, d/dz has the form

$$\frac{d}{dz} = c_a(\theta)\partial_x + 2\theta_0 c_a(\theta)F'_q(x)\partial_\theta, \quad x = t/T, \quad (85)$$

where

$$c_a(\theta) = \frac{2\sqrt{1-a}\cos\theta}{\theta_0} \quad (86)$$

after absorbing a constant into A . Since $a \leq 1/2$, we have $\theta_0 \in [\pi/4, \pi/2]$, so c_a , its θ -derivatives, and the coefficient $2\theta_0 c_a$ are uniformly bounded. The same induction as in Lemma 4 therefore gives

$$\|K_a^{(n)}\|_\infty + \|K_a^{(n)}\|_1 \leq CR^n M_n, \quad n \geq 1, \quad (87)$$

with constants independent of N and M . The L^1 estimate again uses only the total angle variation

$$\int_0^1 |\theta'(x)| dx = 2\theta_0 \leq \pi, \quad (88)$$

and hence does not pay for the long Roland–Cerf coordinate length $\Theta_a(1) = \Theta(\sqrt{N/M})$. Endpoint flatness follows from the fact that F_q is constant near $x = 0, 1$.

Applying Theorem 3 gives leakage amplitude

$$\eta_G(A) \leq C \exp\left[-c \frac{A}{[\log(e+A)]^q}\right]. \quad (89)$$

At $s = 1$, the ground state in the invariant two-dimensional subspace is $|w\rangle$, which lies in the marked subspace. Hence the probability of measuring an unmarked item is bounded, after adjusting constants, by

$$p_{\text{fail}} \leq C \exp\left[-c \frac{A}{[\log(e+A)]^q}\right], \quad T = A\Theta_a(1) = \Theta\left(A\sqrt{\frac{N}{M}}\right). \quad (90)$$

Thus the DC-clocked Roland–Cerf schedule gives a single-run, high-confidence adiabatic Grover evolution with near-exponential failure suppression in the dimensionless adiabatic parameter A , assuming the marked fraction M/N is known.