Infinite dessin orders, projective resolutions and spectral determinants

Constructions, invariant theory and the arithmetic comparison problem

Research working draft

November 4, 2026

Abstract

We work throughout with infinite-dimensional locally gentle algebras and completed surface orders arising from explicit cellular clean dessins on compact Riemann surfaces. We construct their projective resolutions, Cartan determinants and dual dessins; develop determinant semi-invariants, Schur modules and cyclic-quiver character kernels; and calculate actual lattice almost-split sequences and an affine-type diamond quotient. A constrained-path Fock algebra retains the gentle relations and has a compact extension with incidence boundary map. Its degree spectral triple and an inserted arithmetic Fredholm family are explicit. Jacobi sums provide theta-dessin cellular Frobenius determinants with a positive local metric; inert primes obstruct the same real two-dimensional carrier. Recent spectral approaches are compared with their actual hypotheses. Selected power-series, module and matrix statements are verified in Lean4. A derived branch functor repairs projective determinant multiplicity; tame descent recovers a hereditary order but forces multiplicities for full-deck coefficient kernels. We also certify finite completed Dirichlet readouts and derive a cutoff-uniform original-family type-I remainder envelope, retaining its unresolved exponent loss. The arithmetic positive determinant comparison remains open; no proof of GRH is claimed.

Contents

1 Cellular dessins and their infinite orders **2**

2 Projective resolutions, dual dessins and determinants **4**
2.1 Auxiliary staircase resolutions over the order 5
2.2 A monotone quantum comparison with exact finite ranks 6
2.3 A correct DG model of the infinite node 7

3 Determinant semi-invariants and Schur constructions **7**
3.1 Young tableaux and a cyclic representation stratum 9

4 Lattice almost-split sequences and genuine diamond quotients **10**
4.1 Polynomial strings, completed lattices and the Fukaya category 11

5 Leavitt relations, the constrained Fock algebra and spectrum **12**
5.1 A graph presentation and an actual pullback 13
5.2 A separate diagonal-fiber C* completion retaining the pairing 13

6 Arithmetic attachment, descent and the remaining comparison **15**
6.1 A constructive arithmetic benchmark on the theta dessin 16
6.2 A genuine cellular carrier, and what still fails for the order 16
6.3 Arbitrary projective actions: an obstruction and a genuine determinant repair 17
6.4 Tame descent: a surviving order and a sharper obstruction 18
6.5 Whole components, derived endpoints and determinant relations 21

7 A genuine theta covering and its categorical limits **21**

8 Verification and research frontier **23**

9 Why an absolute coprime pair-square estimate cannot close the short rows **24**

10 A signed Möbius decomposition and its termwise obstruction **25**

11 A full-mask Hecke expansion for the type-I piece **26**

12 Actual renormalization retains the effective action **28**
12.1 Genuine dessin incidence models 28
12.2 Quantum thermal effective Hamiltonians 29

13 Linear relations and functorial branch filtrations **29**

14 A kernel-checked extension over the infinite dessin algebra **30**

15 Learned fields, width limits and quantum geometry **31**

16 Complementary physical datasets and checked dynamics **33**

17 Molecular structure, complex transport and learned chemistry **34**

18 Learning objectives with explicit analytic guarantees **34**

19 Learning nonlocal strategies and checking their certificates **34**

20 Hubs, spokes and the topology retained by a dessin **35**

21 A finite, validated approximation result **36**

22 What this does and does not certify **38**

23 An arithmetic-family moment with every original row retained **38**
23.1 The exact masked remainder and the remaining loss 39

24 Recombining the actual weighted cutoff **41**

25 A cutoff-uniform envelope for the original masked remainder **42**

1 Cellular dessins and their infinite orders

We start with the surface-order constructions of Schreiber [1, 2], retaining the original sources unchanged. The generic-resolution source [3] is background, not a substitute for exactness of a particular complex. All primary algebra examples below come from explicit clean Belyi maps. Finite-dimensional objects occur as modules or specified quotient strata of those infinite algebras, not as replacements for them. The initial arithmetic target is a primitive nonprincipal Dirichlet

L-function. Finite-order Hecke characters, Dedekind zeta functions and general Artin representations require separate comparison theorems; Artin holomorphy is not assumed.

A clean dessin on a compact oriented surface is specified by a nonempty finite dart set H, a fixed-point-free involution α, and a permutation σ such that $\langle \alpha, \sigma \rangle$ is transitive. Contract the white vertices of degree two. The resulting cellular graph has vertices the σ-orbits, edges the α-orbits, and faces the ϕ-orbits, where

$$\tau = \sigma^{-1}, \qquad \phi = \alpha\sigma^{-1}, \qquad 2 - 2g = \#\operatorname{cyc}(\sigma) - |H|/2 + \#\operatorname{cyc}(\phi).$$

Write $[h] = \{h, \alpha h\}$. Our quiver has vertices these edges and arrows $a_h : [h] \to [\tau h]$. Paths are composed in travel order: ab traverses a first. Set

$$A = kQ/(a_h a_{\phi h} : h \in H), \quad J = A_{>0}, \quad \Lambda = \varprojlim A/J^m.$$

Each vertex has two incoming and two outgoing arrows. Each arrow has precisely one permitted successor $a_{\tau h}$ and one forbidden successor $a_{\phi h}$; these are distinct because α has no fixed points. The same holds for predecessors. This is the *locally gentle* condition. Every permitted cycle repeats indefinitely, so A is infinite dimensional. We reserve "gentle" without qualification for the convention additionally imposing finite dimension.

Proposition 1.1 (The actual split order)**.** *For a permitted cycle c of length ℓ, let B_c be the hereditary cycle order in $M_\ell(k[[u_c]])$: diagonal and forward matrix entries are in $k[[u_c]]$, and entries crossing the chosen cycle cut are in $u_c k[[u_c]]$. Then Λ is the suborder of $B = \prod_c B_c$ whose two diagonal constant terms labelled $h, \alpha h$ agree. With $R = k[[u]]$ acting by $u \mapsto (u_c I)_c$, it is finite free of rank $\sum_c \ell(c)^2$; the conductor contains uB.*

Proof. In a cycle order a path of length less than ℓ is the corresponding matrix unit, multiplied by u_c exactly when it crosses the cut; another full turn multiplies it by u_c. Thus completed paths in each permitted branch give all entries of B_c. An arrow belongs to only that branch, whereas $e_{[h]}$ has diagonal entries at both occurrences $h, \alpha h$. Hence the only condition on branchwise formal coefficients is the displayed equality of diagonal constants. This proves both injectivity and surjectivity of the path-normal-form identification. Multiplication respects it: switching branches across a paired edge is precisely a forbidden pair. The quotient B/Λ is killed by u, so the submodule has the same generic rank as B. It is torsion free and finite over the DVR R, hence free of that rank. The path and u-adic topologies agree: $uA \subset J$, and if L is the longest permitted cycle, any path of length at least $L + 1$ is u times a shorter nonempty path. Thus $J^{L+1} \subset uA$. This also justifies the completion used in the identification. □

The path-degree variable q below is not u: one full turn has path degree $\ell(c)$ but u-degree one. Formal, Laurent and operator-norm completions will likewise remain distinct.

Here are five explicit global inputs. Permutations list darts starting at zero; $E = |H|/2$.

Surface and Belyi map β	σ	α	$(g, \operatorname{rank}_R \Lambda)$
$\mathbb{P}^1$, $4z(1-z)$	$(0)(1)$	(01)	$(0, 2)$
$\mathbb{P}^1$, $(z+1)^2/(4z)$	(01)	(01)	$(0, 4)$
$y^2 = x^3 - 1$, x^3	$(012)(345)$	$(03)(14)(25)$	$(1, 18)$
$\mathbb{P}^1$, $4z^3/(1+z^3)^2$	$(012)(345)$	$(03)(15)(24)$	$(0, 18)$
$y^2 = x^3 - x$, x^2	(0123)	$(02)(13)$	$(1, 16)$

The first is the nodal interval: $A = k[x, y]/(xy)$ and $\Lambda = k[[x, y]]/(xy)$. The second has $A = k\langle x, y \rangle/(x^2, y^2)$, still infinite dimensional. The two theta dessins have the same two permitted

three-cycles but different face cycles, respectively (051324) and (04)(13)(25). The elliptic square has one permitted and one forbidden four-cycle. For example x has pole order two on the cubic $y^2 = x^3 - 1$; x^3 has ramification $(3,3)$ above zero, $(2,2,2)$ above one since $x^3 - 1 = y^2$, and (6) above infinity. The rational theta map has passport $(3,3),(2,2,2),(2,2,2)$. These ramification computations, and the analogous direct zeros/poles in the other rows, verify cellular Belyi inputs rather than merely abstract quivers.

2 Projective resolutions, dual dessins and determinants

Use right modules $P_i = e_iA$, $S_i = P_i/P_iJ$. Internal shift $M\langle n\rangle$ places its old generator in degree n. Let $\mathcal{R}_n(i)$ be the *indexing words* of length n from i with every adjacent pair forbidden, including e_i for $n = 0$. Such a word is usually zero in A; it still labels a summand.

Theorem 2.1 (Exact resolutions)**.** *There is a minimal graded projective resolution, with $P_0(i) = e_iA$ and for $n \geq 1$,*

$$P_n(i) = \bigoplus_{p=a_1\cdots a_n \in \mathcal{R}_n(i)} e_{t(a_n)}A\langle n\rangle, \qquad d_n(x)_p = a_n x \text{ in the summand } a_1\cdots a_{n-1}.$$

The augmentation $P_0(i) \to S_i$ is the quotient. For the completed order the identical finite-rank projective terms and formal-series maps resolve the residue simple. Ext and Tor against residue simples count forbidden words with the prescribed endpoints. In the graded case Ext has matching internal shift n.

Proof. The maps are right linear and degree zero. Adjacent differentials multiply by a forbidden pair. The augmentation kernel is the direct sum of aA over initial arrows. For any arrow a, multiplication $e_{t(a)}A \to e_{s(a)}A$ kills exactly the permitted paths beginning with its forbidden successor $f(a)$. All other basis paths survive to distinct basis paths. Thus its kernel is $f(a)A$. For a fixed prefix, different last arrows have disjoint surviving supports; there is no cancellation between summands. This proves exactness inductively. For formal series the same assertion holds coefficient by coefficient and identifies the entire closed kernel, not just its dense polynomial subspace. Terms are idempotent projectives and all entries are arrows. Consequently applying Hom or tensor with a residue simple kills every differential, proving minimality and the Ext/Tor count. In A minimality is graded; it does not require that J be its ordinary Jacobson radical. □

The two arms never terminate in this closed class. Their labelled patterns follow ϕ-cycles. An ungraded positive syzygy tail can have a shorter period by permutation of summands: the nodal interval has labelled period two but unordered tail period one; the elliptic theta has labelled period six and unordered tail period three. A simple need not be isomorphic to a positive syzygy; for the node it is R-torsion, whereas the positive syzygies are lattices.

Let $C_{ij}(q) = \sum_{d\geq 0} \dim_k(e_iA_de_j)q^d$ and $E_{ij}(q) = \sum_{n\geq 0}(-1)^n \dim_k \operatorname{Ext}^n_A(S_i, S_j)q^n$. Degreewise Euler characteristic of the resolution gives $EC = I$: every coefficient is finite, because a summand in homological degree n starts in internal degree n. Constant terms are I, so the inverse is two-sided. Already signed $E(q)$ must not be evaluated at $-q$ again.

Proposition 2.2 (Dual dessin and cycle determinant)**.** *The complementary-relation algebra is another closed dessin algebra, with $\alpha^\# = \alpha$, $\sigma^\# = \phi^{-1}$, permitted successor ϕ, and forbidden successor τ. It exchanges graph vertices and faces and preserves genus. Its Cartan matrix satisfies*

$E(q) = C_{A^\#}(-q)$, *with the opposite convention if required for Yoneda composition. For independent commuting arrow weights* w_a,

$$\det C_A(\mathbf{w}) = \frac{\prod_{f\in \mathrm{cyc}(\phi)}(1-(-1)^{|f|}w(f))}{\prod_{c\in \mathrm{cyc}(\sigma)}(1-w(c))}.$$

Proof. $\alpha(\sigma^\#)^{-1} = \alpha\phi = \tau$, so the claimed relations and face data follow directly; the connectedness group is unchanged. To prove the determinant, put $S_{v,a} = w_a$ when $s(a) = v$, and $T_{a,v} = 1$ when $t(a) = v$, zero otherwise. On arrow states let $U_{a,b} = w_b$ for permitted adjacency and $V_{a,b} = w_b$ for forbidden adjacency. Then $TS = U + V$ and the path sum is $C = I + S(I - U)^{-1}T$. Sylvester's identity gives $\det C = \det(I + V)/\det(I - U)$. Each transition is a weighted permutation, whose cycle determinant is respectively $1 - (-1)^\ell w(c)$ and $1 - w(c)$. Everything holds in formal power series, hence as rational functions. This supplies a direct proof of the Bessenrodt–Holm formula in the closed dessin class [4]. □

The five univariate determinants are, in the order above,

$$\frac{1+q}{1-q},\quad \frac{1+q}{1-q},\quad \frac{1-q^6}{(1-q^3)^2},\quad \frac{(1-q^2)^3}{(1-q^3)^2},\quad 1.$$

The last cancellation does not imply finite global dimension: its forbidden resolution is infinite. These rational functions have cycle poles/zeros, not arithmetic prime Euler factors.

For clarity, a complex is a sequence with $d^2 = 0$, its homotopy category identifies maps differing by $dh + hd$, and $D(A)$ inverts maps inducing cohomology isomorphisms. A perfect complex is represented by a bounded complex of finite projectives. Our infinite simple resolutions compute derived Hom and tensor but are generally not perfect. The two-term presentation $[P_1 \to P_0]$ is perfect even if its kernel is nonzero. Quadratic linearity is a Koszul statement; it is not by itself an equivalence between all completed derived categories. Likewise a finite-dimensional gentle graded-curve theorem [6] cannot simply be transferred to these closed infinite orders. Here the surface walks explicitly compute the resolution endpoints: τ turns round a graph vertex and ϕ follows a face. For longer string objects all maps can be computed from the occurrence matrices; a universal geodesic-intersection classification for this completed category remains a separate theorem.

2.1 Auxiliary staircase resolutions over the order

The two filtration parameters below are not dessin arrows. Let $R = k[x, y]$ for a field k, and let $I = (g_0, \dots, g_{m-1})$, $m \geq 2$, where $g_i = x^{a_i}y^{b_i}$,

$$a_0 > \cdots > a_{m-1} = 0, \qquad 0 = b_0 < \cdots < b_{m-1}.$$

The positive outer exponents make R/I finite length. Write $R(-a, -b)$ for a free generator born in bidegree (a, b).

Proposition 2.3 (Adjacent syzygies)**.** *A minimal bigraded free resolution of* $M = R/I$ *is*

$$0 \longrightarrow \bigoplus_{i=0}^{m-2} R(-a_i, -b_{i+1}) \xrightarrow{d_2} \bigoplus_{i=0}^{m-1} R(-a_i, -b_i) \xrightarrow{d_1} R \longrightarrow M \longrightarrow 0,$$

where $d_1(e_i) = g_i$ *and* $d_2(f_i) = -y^{b_{i+1}-b_i}e_i + x^{a_i-a_{i+1}}e_{i+1}$. *Deleting row* i *from* d_2 *gives determinant* $(-1)^i g_i$. *The Hilbert series numerator is* $1 - \sum_i s^{a_i}t^{b_i} + \sum_{i=0}^{m-2} s^{a_i}t^{b_{i+1}}$.

Proof. Each column maps to zero. In a relation $\sum h_i g_i = 0$, all terms except the last contain $x^{a_{m-2}}$. Coprimality with $y^{b_{m-1}}$ implies $h_{m-1} = x^{a_{m-2}} q$. Subtract $q d_2(f_{m-2})$ from the coefficient vector; its last component vanishes. Factor $x^{a_{m-2}}$ from the remaining generators. They form a smaller staircase with the same strict inequalities, ending in a pure y-power. Induction expresses the remaining relation in the preceding adjacent columns. The base of one generator has no relations, since R is a domain. Hence $\ker d_1 = \operatorname{im} d_2$. If $d_2(q) = 0$, its first row gives $-y^{b_1} q_0 = 0$; successive rows then force all $q_i = 0$. This proves injectivity. Every differential entry has positive degree, so reduction modulo (x, y) kills the differentials and the resolution is minimal. In the row-deleted matrix the unique nonzero permutation term uses the negative y entries before row i and the positive x entries after it. Their exponents telescope to b_i, a_i, respectively, and the sign is $(-1)^i$. Finally take the degreewise Euler characteristic; the free summands have Hilbert denominator $(1 - s)(1 - t)$. □

For $I = (x^2, xy, y^2)$ the underlying sequence is also kernel checked in Lean 4 over the actual ring MvPolynomial(Fin 2, k). The verification proves the kernel/image equalities, the stated ideal image, surjective quotient augmentation, projective free terms, and vanishing of both induced Hom differentials to the constant-coefficient residue field. It uses regular multiplication on the free modules, distinct from the residue action on k. General staircases, graded minimality and the surface-order coefficient cone remain separately scoped prose arguments.

At grade $\nu \in \mathbb{N}^2$, M_ν has dimension one exactly when no generator degree is componentwise at most ν. Its comparable-grade map has rank one exactly when both endpoint grades satisfy this condition. A CW realization takes one loop and attaches a disk once along it at each generator degree. The active indices form an interval: the x inequality selects a final interval, the y inequality an initial interval. With c active disks, the cellular map $k^c \to k$ has every entry one. Thus $\dim H_1 = \mathbf{1}_{c=0}$ and $\dim H_2 = \max(c - 1, 0)$, the latter counting active adjacent syzygies. This auxiliary CW complex is not the dessin surface.

For the actual nodal order $\Lambda = k[[\xi, \eta]]/(\xi\eta)$ put $u = \xi + \eta$ and $E = \Lambda/u^2\Lambda$. The basis $(1, \xi, u, u\xi)$ of E satisfies $\xi^2 = u\xi$, $u^2 = 0$, $\eta = u - \xi$. Tensoring the resolution above with $0 \to \Lambda \xrightarrow{u^2} \Lambda \to E \to 0$ gives, over $\mathcal{B} = \Lambda[x, y]$, free ranks $(1, m + 1, 2m - 1, m - 1)$. Indeed the scalar resolution stays exact after tensoring over the field; its degree-zero module is free over Λ on the standard monomials. Multiplication by u^2 is injective there and on all free terms, so the long exact homology sequence of its mapping cone gives positive-degree homology zero and degree-zero module $E \otimes_k M$. In contrast, as an R-module this object has ranks $4(1, m, m - 1)$ in its minimal resolution. Neither finite-length E nor these parameter rings replace the infinite parent dessin algebra or its lattice category.

2.2 A monotone quantum comparison with exact finite ranks

Proposition 2.4 (One common channel)**.** *Let T be a positive trace-preserving linear map between finite complex matrix algebras. For density matrices ρ, σ, their trace distance $d(\rho, \sigma) = \frac{1}{2}\|\rho - \sigma\|_1$ satisfies $d(T\rho, T\sigma) \le d(\rho, \sigma)$. For finitely many labelled states, keep fixed vertex birth grades e_i. At (r, e) include vertices with $e_i \le e$ and edges with distance at most r, and take the flag complex. Then T gives an identity-on-labels inclusion from the input to the output bifiltration, natural in (r, e).*

Proof. Write the Hermitian difference as $A = A_+ - A_-$ with $A_\pm \ge 0$. Positivity, the trace norm triangle inequality and trace preservation give $\|TA\|_1 \le \|TA_+\|_1 + \|TA_-\|_1 = \operatorname{tr} A_+ + \operatorname{tr} A_- = \|A\|_1$. Thus every input edge is an output edge; the fixed births preserve vertices, and all cliques are preserved. Increasing either parameter preserves the simplices, and every map is the same label inclusion, so the squares commute. Complete positivity is sufficient but is not needed for this norm argument. □

For a finite inclusion $K \subset L$, work over a field and embed a basis Z_K of source one-cycles into the target edge space. If B_L is the matrix of target two-boundaries, the induced H_1 rank is

$$\rho_1(K, L) = \operatorname{rank}[B_L \; Z_K] - \operatorname{rank} B_L.$$

Its image is $(Z_K + \operatorname{im} B_L)/\operatorname{im} B_L$; source boundaries already map into target boundaries. This proves the formula, including empty complexes. Computing a flag two-skeleton suffices for H_0, H_1, but does not compute the full flag complex's H_2. Integer ceilings of scaled exact distances preserve the inclusion. For numerical distances, frozen integer labels define an exact combinatorial diagram only after their inequality is checked; this is not an interval certificate of the underlying quantum distances. None of these finite ranks identifies a completed arithmetic determinant or proves GRH.

2.3 A correct DG model of the infinite node

Use cohomological degrees and an uncurved differential of degree $+1$. An odd element ε with $d\varepsilon = 1$ contracts the underlying k-complex of every unital left DG module: $h(m) = \varepsilon m$ satisfies $dh + hd = 1$ by the graded Leibniz rule. Thus this example has zero cohomology, contrary to the original ground-field-resolution claim. An isomorphism on H^0 alone is also insufficient for a DG quasi-isomorphism; all degrees must be checked.

Proposition 2.5 (Nodal Koszul model)**.** *Let k be a field, $R = k[\xi, \eta]$ and $f = \xi\eta$ and $K = R \oplus R\varepsilon$, with $|\varepsilon| = -1$, $\varepsilon^2 = 0$, $dR = 0$ and $d\varepsilon = f$. The augmentation $K \to A = R/(\xi\eta)$, $a + b\varepsilon \mapsto [a]$, is a DG-algebra quasi-isomorphism onto the actual infinite nodal dessin algebra. Branch rescaling $\xi \mapsto q\xi$, $\eta \mapsto q^{-1}\eta$, $\varepsilon \mapsto \varepsilon$ for $q \in k^\times$ lifts to a DG automorphism.*

Proof. Multiplication is $(a + b\varepsilon)(c + e\varepsilon) = ac + (ae + bc)\varepsilon$. The differential kills degree zero and sends $b\varepsilon$ to fb. Its square is zero. For two odd inputs the Leibniz right side is $fbc\varepsilon - bfc\varepsilon = 0$; the other homogeneous cases follow by R-linearity. Multiplication by f is injective in the polynomial domain. Hence $H^{-1}(K) = 0$, $H^0(K) = R/(f)$, and all other groups vanish. The augmentation preserves products and kills boundaries. Rescaling fixes f, so its coefficientwise extension commutes with d and has inverse obtained by replacing q with q^{-1}. This is a one-relation Koszul construction [78, 79]. □

Replacing R by $k[[\xi, \eta]]$ gives the completed order: setting each variable separately to zero maps onto the branch pullback, and the kernel consists precisely of mixed monomials, hence equals $(\xi\eta)$. Rescaling is generally semilinear over the coefficient ring, rather than linear over the fixed diagonal DVR $k[[u]]$, $u = \xi + \eta$. It preserves that DVR as a subring exactly when $q^2 = 1$, and fixes it pointwise exactly when $q = 1$. No Hilbert norm or arithmetic spectrum follows.

The corresponding Lean file checks the actual exterior-line algebra, differential, sign-Leibniz, homotopy and polynomial ambient-resolution identities. Full graded/category/completion objects remain separately scoped. The two theta dessins likewise require endpoint-sensitive Ext: self Ext^1 is zero in both; for the elliptic theta self Ext^3 has dimension two, while for the spherical theta positive even self Ext has dimension two. Outdegree alone does not count self extensions. Minimal A_∞ models use additional graded data [77, §3.3]; m_3 is not automatically a physical curvature.

3 Determinant semi-invariants and Schur constructions

Assume k algebraically closed of characteristic zero for invariant-ring statements. A finite-dimensional representation of the *infinite* A has spaces N_i, arrow matrices $N(a)$ satisfying the monomial relations,

and base-change group $G = \prod_i \mathrm{GL}(N_i)$. With column vectors, $N(a) \mapsto g_{t(a)} N(a) g_{s(a)}^{-1}$. Let $P_0 = \bigoplus P_i^{a_i}$, $P_1 = \bigoplus P_i^{b_i}$, and $f : P_1 \to P_0$ be a finite path-matrix map. When $\sum a_i \dim N_i = \sum b_i \dim N_i$, define

$$c_f(N) = \det D_f(N), \qquad D_f(N) = \mathrm{Hom}_A(f, N).$$

It is intrinsically a determinant-line section; chosen bases give this scalar.

Proposition 3.1 (Weights and exact homological scope)**.** *$c_f(gN) = \prod_i \det(g_i)^{b_i - a_i} c_f(N)$. Restricted determinants of this form span all semi-invariants on a closed G-stable relation or nilpotence locus. Set $M = \operatorname{coker} f$. On the square locus,*

$$c_f(N) \neq 0 \quad \Longleftrightarrow \quad \mathrm{Hom}(M, N) = 0 \quad \Longleftrightarrow \quad \mathrm{Hom}(M, N) = \mathrm{Ext}^1(M, N) = 0.$$

If f is injective, $\operatorname{coker} D_f = \mathrm{Ext}^1(M, N)$ and higher Ext vanish. Without injectivity neither that cokernel equality nor higher vanishing follows.

Proof. Naturality gives $D_f(gN) = G_1 D_f(N) G_0^{-1}$, where G_j repeats the relevant g_i. Take determinants. The contragredient coordinate-ring action has the inverse character. Schofield–van den Bergh's spanning theorem allows oriented cycles [36, Theorem2.3]. Exactness of invariants for reductive G in characteristic zero lifts a fixed-weight function from any closed stable locus to the ambient representation space; restrict the spanning determinants. Injectivity of the path map need not survive that restriction. Finally $\ker D_f = \mathrm{Hom}(M, N)$. Writing $K = \operatorname{im} f$, the exact sequence with projective P_0 expresses $\mathrm{Ext}^1(M, N)$ as $\mathrm{Hom}(K, N)/\operatorname{im} \mathrm{Hom}(P_0, N)$; the epimorphism $P_1 \to K$ embeds this quotient in $\operatorname{coker} D_f$. A square finite matrix with zero kernel has zero cokernel. If f is injective it is the actual length-one resolution, giving the stronger assertion. □

For the node, $f = \times x$, $M = \Lambda/(x)$, $N = \Lambda/(x, y^2)$ give $\dim \operatorname{coker} D_f = 2$ but $\dim \mathrm{Ext}^1(M, N) = 1$. For the elliptic square name arrows so that $a, b : 0 \to 1$, $c, d : 1 \to 0$, permitted cycle *acbd* and forbidden cycle *adbc*. With $f = a$ and $N(a) = 1$, $N(b) = N(c) = N(d) = 0$, its Hom matrix is [1], yet the subsequent forbidden resolution gives $\mathrm{Ext}^{2,3}(M, N) = k$. Thus nonvanishing is not all-degree derived orthogonality.

Proposition 3.2 (What completion does to invariants)**.** *A continuous d-dimensional representation of Λ kills J^d. On the reduced nilpotent representation variety of fixed dimension vector every invariant regular function is constant. Nonzero-weight semi-invariants need not be constant.*

Proof. Continuity with discrete finite-dimensional target kills some J^m; its Loewy filtration has at most d nonzero strict steps. Choose vertexwise complements to this filtration. An arrow sends a layer p into layers $q \geq p + 1$. Scaling layer p by z^p conjugates each component by z^{q-p}. At $z = 0$ the representation tends to the zero-arrow representation. Every invariant therefore has the same value at all points; reducedness and the Nullstellensatz give the ring assertion. In the square example the scalar $N(a)$ varies on a nilpotent locus and has weight g_1/g_0, so semi-invariants survive. Formal entries in a projective map are evaluated on this locus by truncating modulo J^d, a legitimate finite polynomial operation. □

On the node, $u = x + y$ is regular and A is free of rank two over $k[u]$. The polynomial presentation $\times(u - \lambda)$ for $\lambda \neq 0$ is injective and its determinant on N is $\det(u_N - \lambda I)$, the characteristic polynomial of u_N up to the sign $(-1)^d$. But $u - \lambda$ is a unit in Λ, with its geometric-series inverse, so those modules do not extend continuously to the completion. An R-free lattice and its u-nilpotent quotients are different categories. On a lattice a separately specified endomorphism F has a meaningful relative characteristic polynomial over R; the scalar base action uI alone does not create Frobenius.

3.1 Young tableaux and a cyclic representation stratum

For a nonempty partition $\lambda \vdash m$, choose a numbering T of its boxes. On $V^{\otimes m}$ let $Y_T = R_T C_T$, the unnormalized row sum followed by the signed column sum. The classical identity $Y_T^2 = h_\lambda Y_T$, where h_λ is the hook product, makes Y_T/h_λ an idempotent in characteristic zero. Its image is $S_\lambda(V)$, one Schur copy, rather than the whole symmetric-group isotypic component [31, 32]. Its character is s_λ, computed by semistandard tableaux (weak rows, strict columns), and

$$\dim S_\lambda(k^d) = \prod_{(i,j)\in\lambda} \frac{d+j-i}{h_{ij}}.$$

Since $f^{\otimes m}$ commutes with the place permutations, it induces $S_\lambda(f)$ and respects composition and zero maps. Applying this at every vertex and arrow therefore gives a genuine representation of the monomial dessin algebra: $S_\lambda(N(b))S_\lambda(N(a)) = S_\lambda(0) = 0$ for a forbidden pair. The empty partition does not have this property: it sends zero arrow maps to the identity of k.

The Littlewood–Richardson rule gives natural intertwiners

$$S_\lambda(V) \otimes S_\mu(V) \cong \bigoplus_\nu S_\nu(V)^{\oplus c^\nu_{\lambda\mu}}.$$

Here $c^\nu_{\lambda\mu}$ counts semistandard tableaux of skew shape ν/λ and content μ whose row-reading word is Yamanouchi. Naturality makes the same decompositions intertwine all the arrows. The tensor operation is vertexwise with synchronous arrow maps, not $\otimes_A$ and not a Hopf-algebra tensor structure. There is no automatic tensor unit preserving these zero relations. These functors are not additive or exact; they cannot simply be applied degreewise to a projective resolution. They also need not preserve the fixed R-action on lattices.

In the elliptic theta set one of the two permitted three-cycle branches to zero. The other has arbitrary maps $X_1 : V_0 \to V_2$, $X_2 : V_2 \to V_1$, $X_3 : V_1 \to V_0$, with all spaces of dimension d. This is a closed representation stratum of our infinite dessin algebra. Put $M = X_3X_2X_1$.

Theorem 3.3 (Cyclic invariants and a local positive kernel)**.** *The invariant ring on this stratum is $k[c_1(M), \dots, c_d(M)]$, where c_j are characteristic coefficients. Its total-arrow-degree Hilbert series is $\prod_{j=1}^{d}(1-t^{3j})^{-1}$. Common multidegree (m,m,m) has basis $\chi_\lambda(M) = \operatorname{tr} S_\lambda(M)$, $\lambda \vdash m$, $\ell(\lambda) \le d$; other multidegrees have no invariants. Products obey the Littlewood–Richardson rule. Over $\mathbb{C}$, if $\|M_i\| \le R_0$ and $0 \le r$ and $rR_0^2 < 1$, the kernel*

$$K_r(M_i, M_j) = \det(I - rM_i \otimes \overline{M_j})^{-1} = \sum_\lambda r^{|\lambda|} \chi_\lambda(M_i)\overline{\chi_\lambda(M_j)}$$

is absolutely convergent and positive semidefinite.

Proof. Cycle-trace generation for quivers, including relation quotients, is the Le Bruyn–Procesi theorem [29, 30]. Every cycle here is a power of the full three-cycle, up to cyclic rotation. In characteristic zero Newton identities give the generators c_j. They are algebraically independent: take $X_1 = X_2 = I$ and arbitrary X_3. Scalar changes at the three vertices force equal arrow multidegrees. Restriction to that same section identifies degree (m,m,m) with degree-m symmetric polynomials in eigenvalues, whose Schur basis and product rule give the assertion. For the kernel use the ordinary Cauchy identity, first as a formal identity in eigenvalues. On Schur modules the operator norm of $S_\lambda(M)$ is at most $R_0^{|\lambda|}$, so its trace is bounded by $\dim S_\lambda(\mathbb{C}^d) R_0^{|\lambda|}$. The dimension Cauchy identity bounds the absolute series by $(1-rR_0^2)^{-d^2}$. Each summand is a rank-one Gram kernel, proving positivity by absolute summation. □

The term for the empty partition is a constant scalar feature, not an allowed module-valued tensor unit. In dimension two,

$$(\operatorname{tr} M)^2 = \operatorname{tr}\operatorname{Sym}^2 M + \det M, \qquad \operatorname{tr}\operatorname{Sym}^2 M = \tfrac{1}{2}((\operatorname{tr} M)^2 + \operatorname{tr} M^2).$$

More generally $\det(I - zM) = \sum_j (-z)^j \operatorname{tr} \bigwedge^j M$ and its reciprocal is $\sum_m z^m \operatorname{tr}\operatorname{Sym}^m M$. These are actual determinant/Schur connections, not identifications with an arithmetic L-function. On continuous finite-length Λ-modules the monodromy is nilpotent and all positive-degree characters vanish. The displayed Cauchy positivity is local representation-theoretic positivity, not the global Weil form.

4 Lattice almost-split sequences and genuine diamond quotients

For the interval dessin let $X = (x)$, $Y = (y)$ in $\Lambda = k[[x, y]]/(xy)$, $R = k[[u]]$, $u = x + y$. A lattice is a finitely generated Λ-module free over R.

Theorem 4.1 (The nodal lattice calculation)**.** *Every lattice is $\Lambda^r \oplus X^a \oplus Y^b$. The almost-split sequences in this lattice category are*

$$0 \to Y \hookrightarrow \Lambda \xrightarrow{\times x} X \to 0, \qquad 0 \to X \hookrightarrow \Lambda \xrightarrow{\times y} Y \to 0.$$

In particular $\tau X = Y$, $\tau Y = X$. Multiplication by u between two copies of either row, followed by reduction modulo u, gives an actual nonzero exact three-by-three diagram.

Proof. Normalize M *inside its total quotient module*: $\overline{M} = BM$, $B = k[[x]] \times k[[y]]$. The conductor gives $u\overline{M} \subset M \subset \overline{M}$. Both branch projections of $W = M/u\overline{M} \subset k^p \oplus k^q$ are onto. Intersect W with the two factors; its remaining quotient identifies their two residual quotients. Separate bases for these pure factors and the common quotient put W into pure first lines, pure second lines and diagonal pairs. Lift branch bases over the DVRs. These are respectively the summands X, Y, Λ. Their endomorphism rings are local.

Exactness of the first sequence is $\operatorname{ann} x = Y$. It does not split: a map $X \to \Lambda$ lands in $\operatorname{ann} y = X$, and its composite with $\times x$ is a nonunit in $\operatorname{End} X = k[[x]]$. Every map $\Lambda \to X$ factors through this epimorphism, $\operatorname{Hom}(Y, X) = 0$, and every nonunit endomorphism of X is divisible by x and factors through $X \hookrightarrow \Lambda \to X$. Classification handles all finite sums; a unit X component would be a retraction. This is right almost splitness. The dual argument for nonsections out of Y proves left almost splitness. Endomorphisms fixing either structure map have constant term one, hence are units, proving minimality. Swap x, y for the other sequence. Finally all three terms are R-free. Multiplication by u is injective and reduction preserves exactness, proving the diagram assertion. □

The lattice AR quiver has arrows $Y \to \Lambda \to X$ and $X \to \Lambda \to Y$; the mesh composites are zero. Its projective vertex Λ is not the endpoint of another nonsplit AR sequence. AR meshes do not mean that all rows and columns of any drawn grid are short exact.

For a second, explicit family fix $n \geq 3$ and the rational Belyi map

$$\beta_n(z) = \frac{1}{1 - T_n(z)^2} = \frac{1}{(1 - z^2)U_{n-1}(z)^2},$$

where T_n, U_{n-1} are Chebyshev polynomials. It has passport $(2n), (2^n), (2^{n-1}, 1, 1)$. Its contracted dual graph is a path, identifying its genus-zero dessin. The associated infinite algebra has vertices

$1, \dots, n$, endpoint loops l, r, opposite arrows $a_i : i \to i+1$, $b_i : i+1 \to i$, and precisely the relations $l^2, r^2, a_i b_i, b_i a_i$. Its single permitted cycle is $l a_1 \cdots a_{n-1} r b_{n-1} \cdots b_1$; the order rank is $(2n)^2$.

Retain one arrow at each edge in an orientation Ω, and kill its opposite. The quotient is exactly the string algebra $H_n(\Omega)$ of Huang–Lin–Su [46]; the killed-arrow ideal contains J^{n+2}, so it is also an actual quotient of Λ_n. Inflation identifies its module category with the full abelian quotient-annihilator subcategory $\mathcal{C}_\Omega$ of Λ_n-modules. This is generally not extension closed in the whole order category. The imported AR property is intrinsic to $\mathcal{C}_\Omega$, not automatically an ambient lattice or finite-length AR property. The extended edition gives nine explicit string modules, all twelve maps and the four exact meshes from the supplied diamond resource, now situated in this actual dessin quotient.

The other supplied resources also have a precise home. For the interval order the subcategory annihilated by J^2 is $\operatorname{Mod} B_0$, $B_0 = \Lambda/J^2$. This is a domestic string quotient, not itself a gentle algebra. Inflation is full and preserves purity, and $\operatorname{Zg} B_0$ is a closed subspace of $\operatorname{Zg} \Lambda$. Ringel's infinite-word/pure-injective program and its proof by Puninski–Prest apply there [43, 44]; the isolated band xy^{-1} gives Krull–Gabriel and relative Cantor–Bendixson dimension two by [45, Theorem10.19]. This is not an ambient rank calculation for all Λ-modules. The parameter in $x = J_2, y = \lambda J_2$ survives even though all cycle traces vanish; the pp condition $\exists v\ (z = xv\ \wedge\ (y - \lambda x)v = 0)$ detects it. These modules are u-torsion and are not nonzero order lattices. Infinite-string and locally finite AR theorems must retain their distinct finiteness hypotheses; a surviving permitted cycle has infinitely many paths at a vertex.

4.1 Polynomial strings, completed lattices and the Fukaya category

Crawley–Boevey's theorem [38, Theorems1.1–1.4] applies directly to the infinite polynomial dessin algebra: its finitely controlled modules decompose into strings and finite-dimensional bands; its pointwise artinian modules additionally allow primitive injective bands. For our finite quiver, these conditions mean finitely generated and artinian respectively. The words may be infinite, but their modules have algebraic direct-sum bases. This is not a classification of every module or of arbitrary formal completions.

In fact no nonzero completed-order lattice, restricted to A, meets either finiteness condition. The polynomial A is finite over $k[u]$: remove full turns from sufficiently long paths as in Proposition 1.1. If an R-free lattice were finitely generated over A, it would be finite torsion free, hence free, over the PID $k[u]$. Multiplication by $1 - u$ would then fail to be onto, whereas it is a unit over $R = k[[u]]$. Also $M \supsetneq uM \supsetneq u^2M \supsetneq \cdots$ excludes artinianity. For a concrete indecomposable outside the ordinary string/band list take $k[[x]]$ with $y = 0$: its endomorphisms are multiplication by formal series, so its endomorphism ring is local. Its x is injective and nonsurjective, while $1 - x$ is invertible; these properties exclude every pure-x ordinary string or band. The extended edition gives the full word argument.

The Fukaya connection requires equally precise categories. Lekili–Polishchuk explicitly allow infinite-dimensional locally gentle algebras, but their equivalence is for *perfect dg modules* of a homologically smooth algebra, with specified grading, line field and stops [66, Definition3.1.1, Lemma3.1.3, Theorem3.2.2]. Every nonempty closed dessin here has both permitted and forbidden cycles, so its polynomial algebra is neither proper nor smooth by their criterion. This does not exclude appropriate singular or completed variants; it prevents a direct use of that particular theorem. Infinite word, graded curve and geodesic are not interchangeable without a surface realization and boundary/grading data.

There is a genuine positive example inside our setting: the polynomial interval quotient $A/(y) = k[x]$ is still infinite dimensional, is smooth, and has no forbidden cycle. Its perfect category has the

partially wrapped annulus model with one stop total. In degree zero $\mathrm{Perf}(k[x]) = D^b(\mathrm{mod}_{\mathrm{fg}}\, k[x])$. Inflation of this quotient is fully faithful on ordinary modules, but not on derived categories: $\mathrm{Ext}^2_A(k[x], k[x]) = k$, whereas $\mathrm{Ext}^2_{k[x]}(k[x], k[x]) = 0$. The former follows from the alternating y, x resolution. Thus this quotient bridge does not identify the whole closed dessin's perfect category with the annulus category.

Finally, Leavitt path algebras *can* fail invariant basis number, but need not. The permitted-state Laurent algebras here have finite-dimensional matrix quotients, and therefore invariant basis number. The polynomial and completed dessin algebras have their vertex-simple quotients, giving the same conclusion. The complex Leavitt algebra of G_α below is a dense *-subalgebra of its graph C*-algebra, on that specified graph; it is not the original gentle algebra with ordinary CK1 imposed.

5 Leavitt relations, the constrained Fock algebra and spectrum

Work now over $\mathbb{C}$. Ordinary Leavitt and graph C* constructions [47, 48] cannot simply impose our original forbidden products: if $ab = 0$ and CK1 says $a^*a = p_{t(a)}$, then $0 = a^*ab = b$. Every arrow is a forbidden successor, hence every arrow and then every vertex is zero. CK2 is not needed. The ordinary graph-Toeplitz CK1 setting has the same collapse.

The valid permitted-arrow-state graph instead consists of the disjoint directed σ-cycles. Its Leavitt algebra is $\bigoplus_c M_{|c|}(k[z, z^{-1}])$, and its graph C* algebra is $\bigoplus_c M_{|c|}(C(\mathbb{T}))$. These forget the α-gluing; both theta dessins give the same answer. They are not the completed original order. Nor is the ordinary Toeplitz algebra of a cycle of length $\ell > 1$ equal to $M_\ell(\mathcal{T})$: their K_0 ranks are ℓ and one respectively [50].

Let $\mathcal{H} = \ell^2(\mathcal{P})$ for all permitted finite paths, including vacua e_v. Define $L_a\delta_p = \delta_{ap}$ when the prefix is permitted and composable, and zero otherwise. Let P_v project onto paths starting at v, $Q_a = L_aL_a^*$ and v_v onto the vacuum. Then

$$L_aL_{\phi(a)} = 0, \quad L_a^*L_a = P_{t(a)} - Q_{\phi(a)} = v_{t(a)} + Q_{\tau(a)}, \quad P_v - \sum_{s(a)=v} Q_a = v_v.$$

These follow on basis vectors. The different initial projection avoids CK1 collapse. Write $\mathcal{T}_A = C^*(P_v, L_a)$, a constrained-path Fock C* algebra in the monomial/subproduct setting [51, 52]. The polynomial A acts faithfully: applying a finite path sum to the sum of vacua recovers its coefficients. This representation need not extend to all formal Λ series: $\sum_{m\geq 0} c^m$ along a cycle would send its vacuum to a nonsquare-summable sum.

Theorem 5.1 (Compact extension and incidence index)**.** *Let $\mathcal{H}_v$ be the terminal-path sector. There is an exact sequence*

$$0 \longrightarrow \bigoplus_{v\in Q_0} \mathcal{K}(\mathcal{H}_v) \longrightarrow \mathcal{T}_A \longrightarrow \bigoplus_{c\in \mathrm{cyc}(\sigma)} M_{|c|}(C(\mathbb{T})) \longrightarrow 0.$$

With $\delta[u] = [1 - W^*W] - [1 - WW^*]$, its boundary map is $\delta(c) = -\sum_{h\in c} e_{[h]}$, minus the unsigned incidence map B_Γ of the cellular graph. Thus, for its E edges,

	$K_0(\mathcal{T}_A)$	$K_1(\mathcal{T}_A)$
Γ *bipartite*	$\mathbb{Z}^{E+1}$	$\mathbb{Z}$
Γ *nonbipartite*	$\mathbb{Z}^E \oplus \mathbb{Z}/2$	0.

Proof. All prefixes preserve the terminal sectors. For paths p, q ending at v, $L_pv_vL_q^* = |\delta_p\rangle\langle\delta_q|$. Therefore the vacuum ideal is exactly the displayed sum of compacts: it contains these matrix units

and that sum is itself an ideal. In the quotient, L_a has initial projection $Q_{\tau(a)}$ and range Q_a, a graph CK family for the permitted-state cycles. The Q_a have infinite rank and nonzero quotient images; the length gauge action descends. Gauge-invariant uniqueness [48, Theorem2.1] identifies the quotient as claimed.

For a cycle c based at arrow a, lift its corner unitary by $W = L_c Q_a + (1 - Q_a)$. Then $W^*W = 1$ and $1 - WW^* = Q_a - L_c Q_a L_c^*$ projects onto the first $|c|$ paths starting with a. Their terminal vertices count exactly its incident darts, including twice for a loop. This proves the index formula. The six-term exact sequence gives $K_1 = \ker B_\Gamma$ and $0 \to \operatorname{coker} B_\Gamma \to K_0 \to \mathbb{Z}^{\#\,\mathrm{cyc}\,\sigma} \to 0$. The free last term splits abstractly. Spanning-tree integer elimination gives Smith entries 1 repeated $V - 1$ times and one zero in the bipartite case; an odd cycle instead adds the entry two. This proves the groups. □

The nodal interval has $K_0 = \mathbb{Z}^2$, $K_1 = \mathbb{Z}$; the single-loop sphere has $\mathbb{Z} \oplus \mathbb{Z}/2$, zero. Both theta dessins have $\mathbb{Z}^4$, $\mathbb{Z}$. Coarse K-theory does not recover genus or the arithmetic factors.

5.1 A graph presentation and an actual pullback

The Fock algebra itself has a graph presentation, on a different graph from the original gentle quiver. Let G_α have vertices q_h for $h \in H$ and sinks b_e for $e \in Q_0$, and edges

$$s_h : q_h \to q_{\tau h}, \qquad t_h : q_h \to b_{[\tau h]}.$$

Proposition 5.2. *$C^*(G_\alpha) \cong \mathcal{T}_A$, by $q_h \mapsto Q_{a_h}$, $b_e \mapsto v_e$, $s_h \mapsto L_{a_h} Q_{a_{\tau h}}$ and $t_h \mapsto L_{a_h} v_{[\tau h]}$. The integer graph relation at h is $q_h - q_{\tau h} - b_{[\tau h]}$. Eliminating these relations recovers $K_1 = \ker B_\Gamma$ and $K_0 \cong \operatorname{coker} B_\Gamma \oplus \mathbb{Z}^{\#\,\mathrm{cyc}\,\sigma}$.*

Proof. The two edges at q_h have orthogonal initial pieces and orthogonal ranges summing to Q_{a_h}, by the Fock initial-projection identity. All vertex projections are nonzero. The length gauge acts with degree one on both edge families, so gauge-invariant uniqueness applies. They generate since $L_{a_h} = s_h + t_h$ and $P_e = v_e + \sum_{[h]=e} Q_{a_h}$. In the integer matrix a kernel coefficient is constant around each permitted cycle; its sink equations are exactly $B_\Gamma c = 0$. In the cokernel, eliminate all but one state per cycle; its remaining relation is the sum of its incident sink generators. The remaining cycle generators are free, proving the stated groups. □

For the nodal interval, G_α has two loop vertices, each with an exit to the common vacuum sink. Chirvasitu–Hajac–Tobolski's actual pullback theorem [65, Theorem3.4] gives

$$\{T \in \mathcal{T}_A : \pi(T) \in \mathbb{C}^2\} \cong C^*(E_1), \qquad \mathbb{C}^2 \hookrightarrow C(\mathbb{T})^2 \text{ by constants},$$

where E_1 has the same three vertices and countably many arrows from each loop vertex to the sink, with no loops. The arrow numbered $n \geq 0$ maps to loop^n followed by exit. This algebra is AF, with $K_0 = \mathbb{Z}^3$, $K_1 = 0$. The extended edition verifies every hypothesis. The theorem does not turn arbitrary cycle-vertex identifications into pullbacks: its first graph is acyclic and its path functor must cover all pointed paths. Its sink-extension Theorem4.3 has further injection and path-coverage requirements.

5.2 A separate diagonal-fiber C* completion retaining the pairing

The formal order's residue gluing cannot be copied by taking diagonals of arbitrary full matrices: $\operatorname{diag}(E_{12}E_{21}) \neq \operatorname{diag}(E_{12}) \operatorname{diag}(E_{21})$. For $\ell > 1$ there is no nonzero *-homomorphism from $M_\ell(C(\mathbb{T}))$

to a commutative C*-algebra, as restriction to its constant matrix units shows. Restrict the distinguished fiber first. Define

$$D_\ell = \{f \in C(\mathbb{T}, M_\ell(\mathbb{C})) : f(1) \text{ is diagonal}\}, \qquad \mathcal{D}_H = \Big(\prod_c D_{|c|}\Big) \underset{\mathbb{C}^H}{\times} \mathbb{C}^{Q_0},$$

where the first map evaluates diagonal entries at one and the second repeats λ_e at the two darts paired by α.

Theorem 5.3 (A faithful polynomial pullback model)**.** *There is a split exact sequence*

$$0 \to \bigoplus_c C_0(\mathbb{T} \setminus \{1\}, M_{|c|}) \to \mathcal{D}_H \to \mathbb{C}^{Q_0} \to 0,$$

so $K_0(\mathcal{D}_H) = \mathbb{Z}^{|Q_0|}$ and $K_1(\mathcal{D}_H) = \mathbb{Z}^{\#\,\mathrm{cyc}\,\sigma}$. The polynomial dessin algebra embeds faithfully and generates this C-algebra with its adjoints. This embedding does not extend to a unital algebra map from all of Λ.*

Proof. The section consists of the constant paired diagonal projections. The kernel is the displayed vanishing fiber, a sum of matrix suspensions, giving the groups by split exactness and suspension. For an explicit embedding set $g(z) = |1 - z|^2/4$, choose one cut arrow per permitted cycle, and send a_h to $g(z)z^{\epsilon_h}E_{h,\tau h}$ in its branch, where $\epsilon_h = 1$ at the cut and zero otherwise. Send e_e to the constant paired diagonal projection. Every forbidden product is zero. A full cycle at h maps to $w_c(z)E_{hh}$, $w_c = g^{|c|}z$. This scalar function is injective on the circle: away from its unique zero at one its argument recovers z. Continuous functional calculus therefore supplies every $f(z)E_{hh}$ with $f(1) = 0$. Multiplying by path images and approximating division away from one supplies every vanishing off-diagonal entry. Together with the constant paired diagonals these generate $\mathcal{D}_H$. Path normal forms in a fixed entry are a nonzero prefactor times a polynomial in w_c; its infinite range proves independence. Evaluation at one first separates the zero-length coefficients. Finally for $z_0 \neq 1$, $1 - w_c(z_0)^{-1}c_h$ is a unit in the formal order, but its proposed image is singular at z_0. A unital map preserves units, proving nonextension. □

This is the C* algebra generated by a faithful polynomial representation and its adjoints, distinct from the completed order and from $\mathcal{T}_A$. The norm closure of the polynomial image alone need not contain those adjoints. Their K groups already differ. The pairing belongs to the labelled presentation; an unmarked isomorphism class need not recover the ribbon ordering. No arithmetic descent, Frobenius action or completed-L positive determinant is inferred from this additional valid gluing model.

Proposition 5.4 (The degree spectral triple)**.** *Define $N\delta_p = |p|\delta_p$ on $\{\xi : \sum_p |p|^2|\xi_p|^2 < \infty\}$. With the polynomial creation/annihilation *-algebra this is an ungraded spectral triple, p-summable exactly for $p > 1$. With N_+^{-s} zero on the kernel,*

$$\operatorname{Tr} e^{-tN} = E + \frac{2E}{e^t - 1}, \quad \operatorname{Tr} N_+^{-s} = 2E\zeta(s), \quad \operatorname{Tr}\big(\chi(N)N_+^{-s}\big) = 2EL(s, \chi)$$

for $t > 0$ and $\operatorname{Re} s > 1$ in the last two formulas, where χ is an externally chosen Dirichlet character.

Proof. There are E vacua and $2E$ paths of every positive length: each starting arrow has a unique continuation. The real diagonal operator is self-adjoint on the stated maximal domain; finite supports are a core and its resolvent is compact. Prefixes preserve the domain and $[N, L_a] = L_a$, $[N, L_a^*] = -L_a^*$, so polynomial commutators are bounded. The trace formulas and summability reduce to the corresponding scalar series. The character multiplier is bounded; membership in $\mathcal{T}_A$ is not asserted. □

Length parity anticommutes with the creators but commutes with N, so it is not an even-spectral-triple grading. The ordinary odd Fredholm class of this positive N is trivial: $N(1+N^2)^{-1/2} - I$ is compact. The spectrum depends only on E; zeros of the scalar zeta trace are not eigenvalues of N.

6 Arithmetic attachment, descent and the remaining comparison

The compact ideal gives a precise home for an arithmetic determinant, with no claim that the arithmetic is derived from the dessin. Fix one terminal sector. Its two paths of each length $n \geq 1$ give orthonormal vectors $\xi_n = (\delta_{p_{n,1}} + \delta_{p_{n,2}})/\sqrt{2}$ and an isometry $U : \ell^2(\mathbb{N}_{\geq 1}) \to \mathcal{H}_v$.

For a nonprincipal character χ put $A_n = \sum_{j \leq n} \chi(j)$, $B_0 = \sup |A_n| \leq q_\chi$, and, on $2^k \leq n < 2^{k+1}$, set

$$w_n = \sqrt{n}(k+1), \quad u_n = A_n/w_n, \quad v_n(s) = w_n(n^{-s} - (n+1)^{-s}), \quad \varphi_s(z) = \sum_n v_n(s) z_n - z_1.$$

Here $u_1 = 1$. For $\operatorname{Re} s > 0$, $u, v(s) \in \ell^2$ and the pairing is complex linear, with no conjugation of s. Define

$$\widetilde{K_s} = U(u \otimes \varphi_s)U^*, \qquad \det_F(I + \widetilde{K_s}) = L(s, \chi).$$

Indeed dyadic blocks give $\|u\|^2 \leq 2B_0^2$, and on a compact set $\operatorname{Re} s \geq \sigma_0 > 0$, $|s| \leq M$, the squared v tail is bounded by $M^2 \sum_{k \geq J} (k+1)^2 2^{-2\sigma_0 k}$. Thus the rank-one family is trace-norm holomorphic. Its determinant is $1 + \varphi_s(u)$, equal by Abel summation to L. Replacing v by $P_N v$ and adding $A_N(N+1)^{-s} e_1^*$ to the functional gives exactly every raw partial sum S_N, with compact-uniform trace-norm convergence; $\|UXU^*\|_1 = \|X\|_1$. For primitive χ, completion replaces the functional by $g(s) \sum v_n(s) z_n - z_1$, with the standard nonzero gamma factor g, rather than multiplying the whole old functional. The extended edition supplies the quantitative estimates.

This inserts a scalar encoding into one actual dessin compact ideal; it is not known to be A-linear, a derived chain map or Frobenius compatible. A common fixed positive metric cannot make this same fixed-range family symmetric for all critical heights: symmetry forces its functionals to be proportional, whereas the coefficient ratio at heights $0, t \neq 0$ is asymptotic to $(\sigma + it)n^{-it}/\sigma$, with two opposite subsequential limits. Different positive realizations remain possible.

Proposition 6.1 (Complex dessin data do not determine arithmetic descent)**.** *The rational Belyi pairs $E_d : y^2 = x^3 - d$, $\beta_d = x^3/d$, for $d = 1, 2$, are isomorphic over $\mathbb{C}$ but have distinct Frobenius polynomials at the good prime seven: $1 - 4T + 7T^2$ and $1 - T + 7T^2$.*

Proof. Choose $a^3 = d$, $b^2 = d$. The projective isomorphism $(x, y) \mapsto (ax, by)$ from E_1 to E_d preserves the Belyi map. The degree-six clean passports are the elliptic theta passport already computed. Directly counting y for $x = 0, \ldots, 6$ gives respectively $(0, 1, 1, 0, 1, 0, 0)$ and $(0, 0, 0, 2, 0, 2, 2)$. Including infinity yields four and seven points, hence $a_7 = 8 - \#E_d(\mathbb{F}_7) = 4, 1$. Both discriminants $-432d^2$ are nonzero modulo seven. □

Any invariant of the complex dessin or its undecorated order/Fock algebra alone is the same for these two inputs, so cannot recover both factors without descent data. This does not obstruct a theory including a specified rational model and compatible Galois action. Elliptic H^1 has weight one and unshifted center $s = 1$, unlike Dirichlet center $1/2$; the example is not a counterexample to GRH.

6.1 A constructive arithmetic benchmark on the theta dessin

The preceding twists can be distinguished constructively once their finite-field descent data are retained. This closes the elementary local-factor computation, not the global positive-determinant problem. Let $\kappa = \mathbb{F}_q$, $\operatorname{char}\kappa \neq 2, 3$ and $d \in \kappa^*$. Put $E_d : y^2 = x^3 - d$ and $a_q(d) = q + 1 - \#E_d(\kappa)$. For $q \equiv 1 \pmod 3$, fix a nontrivial cubic character ρ and the quadratic character η, both extended by zero at zero. Define

$$J = \sum_{u \in \kappa} \rho(u)\eta(1-u), \qquad j = -\eta(-1)J, \qquad \lambda_d = \eta(d)\rho(d), \quad \pi_d = \lambda_d j.$$

Theorem 6.2 (Uniform local factor)**.** *For every such prime power and nonzero d,*

$$a_q(d) = \pi_d + \overline{\pi_d}, \quad |\pi_d|^2 = q, \quad P_d(T) := 1 - a_q(d)T + qT^2 = (1 - \pi_d T)(1 - \overline{\pi_d}T).$$

If $q \equiv 2 \pmod 3$, instead $a_q(d) = 0$. The split-case values lie in $\mathbb{Z}[\omega]$, $\omega^2 + \omega + 1 = 0$.

Proof. Cube fibers have size $1 + \rho(t) + \overline{\rho(t)}$, including at zero; the first term is the literal constant one, not the zero-extended trivial character. Counting square fibers gives $a_q(d) = -\sum_x \eta(x^3 - d)$. Grouping by $t = x^3$ removes the constant term since $\sum_t \eta(t - d) = 0$. Substitution $t = du$ gives $\sum_t \rho(t)\eta(t - d) = \rho(d)\eta(-d)J$; the other term is its conjugate. This proves the signed trace formula.

For a nontrivial additive character ψ, write $G(\chi) = \sum_x \chi(x)\psi(x)$. Character orthogonality gives $|G(\chi)|^2 = q$ for nontrivial χ: substitute $x = uy$ in its squared norm; the inner sum over $y \neq 0$ is $q - 1$ at $u = 1$ and -1 otherwise. Grouping $G(\rho)G(\eta)$ by $x + y$ gives $JG(\rho\eta)$, the zero-sum fiber vanishing since $\rho\eta$ is nontrivial. The three Gauss norms yield $|J|^2 = q$. Neither purity nor RH was assumed. The remaining signs and character values have modulus one. For $q \equiv 2 \pmod 3$ cubing is bijective, so the quadratic-character sum is zero. Finally all split-case summands are signs times cube roots of unity. These are the classical Gauss–Jacobi methods [68, pp.499–502], with our normalization proved explicitly. □

At $q = 7$, choose $\rho(3) = \omega$. Then $j = 3 + 2\omega$, $\lambda_2 = \omega^2$, and $\pi_2 = -1 - 3\omega$, giving the two factors already counted. At $q = 13$, $\rho(2) = \omega$ gives $J = 1 + 4\omega$ but $j = -1 - 4\omega$; this detects the essential sign hidden by the first example.

6.2 A genuine cellular carrier, and what still fails for the order

Assume $q \equiv 1 \pmod 3$ in this cellular construction. The elliptic theta dessin has cellular chains $C_2 = \mathbb{Q}$, $C_1 = \mathbb{Q}^3$, $C_0 = \mathbb{Q}^2$. Orient the three edges from v_+ to v_-; then $\partial_1(e_i) = v_- - v_+$ and $\partial_2 = 0$. Thus H_1 is the sum-zero plane, with basis $e_0 - e_1, e_1 - e_2$. Rotation σ acts there by

$$C = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}, \qquad G = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}.$$

G is the actual Euclidean edge Gram matrix in this basis. The deck involution α acts as $-I$. Consequently the six-element deck group $\langle \sigma, \alpha \rangle$ acts on this homology and permutes the original quiver and its entire projective resolution.

Write $\pi_d = A_d + B_d\omega$. Its multiplication action on this *actual cellular* plane is

$$M_d = A_d I + B_d C = \begin{pmatrix} A_d & -B_d \\ B_d & A_d - B_d \end{pmatrix}, \qquad \det(I - TM_d) = P_d(T), \qquad M_d^t G M_d = qG.$$

This constructs the local matrix from finite-field character sums and the dessin's deck action, rather than fitting its trace. Positivity follows from $2x^2 - 2xy + 2y^2 = (x-y)^2 + x^2 + y^2$. The normalized matrix is G-unitary, not asserted self-adjoint. The extended edition supplies a rational cellular chain lift, its exact determinant cancellation, and Kummer descent compatibility on homology. Its odd-over-even determinant ratio is the classical finite-field curve zeta function

$$Z(E_d/\kappa, T) = \frac{P_d(T)}{(1-T)(1-qT)}.$$

Identification with the geometric ℓ-adic Frobenius polynomial uses the standard elliptic trace and determinant theorem; a canonical integral comparison or an order-derived Frobenius functor has not been constructed here.

There is also a precise obstruction to extending this same real two-dimensional carrier to $q \equiv 2 \pmod 3$. Frobenius then conjugates the cubic deck action. The relation $MC = C^2M$ forces $M = \left(\begin{smallmatrix} a & b \\ a+b & -a \end{smallmatrix}\right)$, whose determinant is $-a^2 - ab - b^2 \le 0$, contradicting the required positive determinant q. This excludes that real carrier with its fixed deck action, not complex or enlarged Hilbert–Pólya constructions. The extended edition gives a compatible complex local replacement.

For the following insertion test use the characteristic-zero theta order $\Lambda_{\mathbb{Q}}$ over $R_{\mathbb{Q}} = \mathbb{Q}[[u]]$, with residue simple $S_e = \mathbb{Q}$. The limitation is concrete. The deck action has finite order and cannot alone supply the norm-$\sqrt{q}$ Jacobi coefficient. Inserting M_d into $S_e \otimes \mathbb{Q}^2$ gives an actual residue-module endomorphism with determinant P_d, but that simple is not perfect over the order and choosing it breaks full deck symmetry. Inserting the same matrix $\Lambda_{\mathbb{Q}}$-linearly into the regular projective instead gives the $R_{\mathbb{Q}}$-determinant P_d^{18}, or P_d^6 on $e\Lambda_{\mathbb{Q}} \otimes \mathbb{Q}^2$. Taking a root is not an established determinant-line comparison; a deck-semilinear lift is a different map. These surviving multiplicity/equivariance obligations prevent promoting the local computation to the required arithmetic order functor.

For the global comparison fix a nonzero rational d and use its good reductions. The finite-field curve zeta has center $1/2$ after $T = q^{-s}$. The *global elliptic* L-function over $\mathbb{Q}$ has unshifted center one, whereas Dirichlet L-functions have center $1/2$. Local weight-one purity does not determine either global zero set. The elementary local bound proves absolute convergence of the elliptic Euler product on $\operatorname{Re} s > 3/2$; the new local identities give no determinant-preserving continuation with the desired global positive spectrum.

6.3 Arbitrary projective actions: an obstruction and a genuine determinant repair

The preceding tensor test has a stronger replacement. Keep the characteristic-zero completed elliptic theta order Λ over $R = \mathbb{Q}[[u]]$, and write $K = \mathbb{Q}((u))$. All determinants in the next theorem are *ordinary underlying-R determinants*; they are not reduced norms.

Theorem 6.3 (Projective determinant multiplicity)**.** *Let P be any finitely generated projective right Λ-module and $f \in \operatorname{End}_\Lambda(P)$. Then $P \simeq \bigoplus_{i=0}^{2} (e_i\Lambda)^{n_i}$. Let $A_i \in M_{n_i}(\mathbb{Q})$ be the induced maps on its three top multiplicity spaces. Then*

$$\det_R(1 - Tf)\Big|_{u=0} = \left(\prod_{i=0}^{2} \det(1 - TA_i)\right)^6.$$

For any bounded complex of these projectives and any Λ-linear chain endomorphism, its alternating determinant ratio specializes to a sixth power in $\mathbb{Q}(T)$. Generically, the ordinary ratio is a cube in $K(T)$.

Proof. The arrow ideal is the Jacobson radical: its elements admit geometric-series inverses after subtracting from one, and the quotient is $\mathbb{Q}^3$. A projective's top lifts to a map from the displayed sum; Nakayama gives surjectivity, projectivity splits it, and another application of Nakayama kills the kernel.

Put $B = \Lambda/u\Lambda$. Each e_iB has path layers of dimensions $1, 2, 2, 1$: the identity, two length-one paths, two length-two paths, and one length-three path. The two full cycles based at i add to ue_i, so become negatives; all length-four paths vanish. A positive-length entry of a projective homomorphism strictly raises path length. Consequently the associated graded map has six copies of $A = \mathrm{diag}(A_0, A_1, A_2)$, giving the formula by triangular block determinants. Base change of determinants justifies specialization. Multiplicativity along filtrations is ordinary linear algebra, consistent with [69, Tags0FJ9,0GSX]; no purity assumption enters.

Each term's determinant has constant term one in T, so every denominator has nonzero specialization. Taking the finite alternating product therefore preserves the sixth power, even after cancellation. Finally $\Lambda_K = M_3(K) \times M_3(K)$; a projective's branch modules are sums of three-dimensional simples. A commuting endomorphism acts on each multiplicity space, repeating every factor three times. This proves the generic assertion. □

Thus no such ordinary perfect-complex ratio can equal the constant-in-u quadratic $P_d(T)$, or the full local curve zeta $P_d(T)/((1-T)(1-qT))$. For P_d, the valuation at infinity is -2, not a multiple of six. For the zeta ratio, $P_d(1) = q + 1 - a_q > 0$ by the local bound $|a_q| \le 2\sqrt{q} < q + 1$, so its pole at $T = 1$ is simple. This also covers square prime powers where the numerator may have a double root. It is an obstruction for the specified ordinary determinant and linear category, not all spectral or determinant theories.

There is nevertheless a *genuine integral functor* removing this multiplicity. Choose a permitted branch and its cycle cut. Its projection $\Lambda \to B_+ \subset M_3(R)$ makes the column lattice $W_+ = R^3$ a left Λ-module. Define

$$\Phi_+ : \mathrm{Perf}(\Lambda) \longrightarrow \mathrm{Perf}(R), \qquad P^\bullet \longmapsto P^\bullet \otimes_\Lambda W_+.$$

Bounded projectives compute this derived tensor functor; flatness of W_+ on all modules is unnecessary. Since $e_iW_+ \simeq R$, its determinant on $e_i\Lambda \otimes_{\mathbb{Q}} V$ with external action $1 \otimes M_d$ is exactly P_d, whereas on the regular module it is P_d^3. This is a defined change of determinant functor, not an arbitrary polynomial root. It does not construct the arithmetic action independently: the already computed M_d is still the input.

The price of this repair is precise. Branch swap α exchanges W_+ and W_-; pairing them yields P_d^2 on that basic-projective input. The chosen branch determinant cannot intertwine α on all endomorphisms: the corner unit $e_i + c_i^+$ acting on $e_i\Lambda$, with c_i^+ a plus full cycle, acts on its determinant by $1 + u$, whereas its α-image acts by one. Also no rank-one-per-vertex column lattice supported on one branch admits an R-linear lift of the rotation σ fixing u. Its three nonzero arrow scalars have nonnegative u-valuations summing to one; an invertible cyclic lift would make all three equal. This contradiction concerns that lattice, not every determinant-level rotation comparison. Ramification $u = v^3$ *and a new lattice* with all three arrow scalars v admits such a rotation, changing the stipulated base/lattice data. A descent-compatible arithmetic determinant retaining the required symmetry and a global positive realization remains open.

6.4 Tame descent: a surviving order and a sharper obstruction

We now test the proposed ramified repair rather than assume that its rotation descends. Extend coefficients explicitly to $k = \mathbb{Q}(\omega)$, $\omega^2 + \omega + 1 = 0$. Put $R = k[[u]]$, $S = k[[v]]$, $u = v^3$, and let

$g(v) = \omega v$, fixing k. On the balanced plus column $W = S^3$, let $C\epsilon_i = \epsilon_{i+1}$ and

$$\Theta = Dg, \qquad D = \mathrm{diag}(1, \omega^2, \omega).$$

The plus arrows act by $vE_{i,i-1}$, and minus arrows by zero. Direct calculation gives $\Theta^3 = 1$, compatibility of Θ with this Λ_R-action, and

$$\Theta C = \omega^2 C\Theta, \qquad L = W^\Theta = R\epsilon_0 \oplus Rv\epsilon_1 \oplus Rv^2\epsilon_2.$$

Thus invariants restore precisely the original cut lattice. The extension $R \to S$ is finite *free* of rank three and faithfully flat; the group quotient is not a torsor at $v = 0$. The comparison $S \otimes_R L \to W$ has matrix $\mathrm{diag}(1, v, v^2)$, determinant u and cokernel $S/(v) \oplus S/(v^2)$. Hence this semilinear group action alone is not effective descent of the column to R. This is the explicit local root-stack situation, consistent with the trivial-stabilizer criterion for vector-bundle descent [70, Theorem10.3]; no stack theorem is needed for the displayed calculation.

Two positive constructions survive. The determinant line $\bigwedge_S^3 W$ has trivial D and C actions and does descend, but the natural map

$$\det_R(L) \longrightarrow (\det_S W)^\Theta$$

has image uR inside R, not an isomorphism. Determinant and coarse invariants cannot be interchanged while dropping this nonunit comparison. Also the *carrier* endomorphism algebra $\mathrm{End}_S(W)^\Theta$, conjugated to the invariant basis, is exactly the hereditary order with entries R on and below the diagonal and uR above. Its rotation descends as $\mathrm{Ad}(A)$, where

$$A = \begin{pmatrix} 0 & 0 & u \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \qquad A^3 = uI.$$

The algebra automorphism has order three although A is not an integral unit. This is not $\mathrm{End}_\Lambda(W)$ or a descended arithmetic cohomology group.

The natural arithmetic attempt fails: $aI + bC$ does not commute with Θ when $b \neq 0$, as in both prime-seven Jacobi examples. The original cellular sum-zero plane is not Θ-stable either. Replacing C by vC does give the descended operator A on L, but

$$\det(1 - T(aI + bA)) = (1 - aT)^3 - ub^3T^3,$$

whose special fiber is $(1 - aT)^3$, not the required local quadratic. These carrier maps are not automatically Λ-linear arithmetic endomorphisms.

There is a stronger category-level conclusion.

Theorem 6.4 (Full-deck coefficient kernels)**.** *Let W_0 be a finite R-free left Λ_R-module on which the central u acts by the given scalar. Suppose it has invertible R-linear intertwiners lifting the order automorphisms σ and α. Then, for some $m \geq 0$, every e_iW_0 has rank $6m$. For any finite projective P and Λ_R-linear $f : P \to P$, let A_i be its top blocks. The genuine coefficient functor $\Phi_{W_0} = - \otimes^{\mathbf{L}}_\Lambda W_0$ satisfies*

$$\det_R(1 - T\Phi_{W_0}(f))\Big|_{u=0} = \left(\prod_i \det_k(1 - TA_i)\right)^{6m}.$$

Bounded perfect ratios retain this exponent. Thus this whole class of full-deck, fixed-center finite free coefficient kernels cannot supply either the single local quadratic or the full local curve zeta.

Proof. In the generic fiber intersect W_0 with its two central branch summands to obtain finite free integral lattices $W_\pm$; an integral direct-sum decomposition is not assumed. Rotation preserves them and branch swap exchanges them. On a branch let the common vertex rank be r. The three arrow matrices have cyclic product uI_r. Their determinant valuations are nonnegative, sum to r, and rotation intertwiners make them equal. Therefore $3 \mid r$. Branch swap makes the two ranks equal, so the full vertex rank is $6m$.

For the determinant formula, filter W_0/uW_0 by the arrow radical, whose fourth power vanishes. A positive-length entry of f raises the filtration; a top block A_i repeats across its i-th graded pieces a total of $\dim_k e_i(W_0/uW_0) = 6m$ times. Triangular determinants give the assertion. Finite alternating products preserve it. Quadratic degree and the uncancelled simple pole of the full local zeta exclude the targets as before. □

This theorem is sharp: $W_0 = \Lambda_R$ has vertex rank six. It requires neither left projectivity nor left flatness of a general kernel. It excludes this coefficient category, not arbitrary semilinear correspondences, twisted determinant lines or infinite spectral constructions. The extended edition identifies the full character-orbit repair with the hereditary normalization and explains why Schur commutator cancellation is insufficient. All determinant multiplicities here are relative to the stated k-valued base; restricting a rational quadratic factor to the former $\mathbb{Q}$-base squares it. No arithmetic descent or global positive determinant has been recovered by this ramification.

The recent NCG literature supplies concrete comparison targets, with strong hypotheses. Connes–Consani construct finite abelian covers of the adelic Riemann sector with prime-orbit length $\log p$ and arithmetic Frobenius monodromy [53, Theorem3.12]. The prolate/local-factor and moment constructions [54, 55] give particular finite-prime analytic models, not a uniform positive comparison for all completed L-functions. The zeta-spectral-triple determinant theorem [56, Theorem5.10] assumes a specified restricted Weil form, simple even bottom eigenfunction and boundary normalization. Its infinite comparison is still an additional step. The distributional real-zero theorem [57, Theorem6.1] concerns a particular interval quadratic form, essential self-adjointness on its trigonometric core and an even simple ground state; a Fock vacuum alone verifies none of those arithmetic identifications. Deninger's periodic-packet construction and real-cohomology obstruction [60, 62] further specify, rather than remove, these issues. Full theorem-level reading coverage and exact versions are recorded alongside this draft.

A complete approach must construct arithmetic descent and prime actions on actual order/derived or analytic objects, prove determinant-line and local-factor compatibility, control regularization and conductor/gamma normalization, and establish a positive realization for that *same* function. Finite-field cohomological trace formulas and Deligne purity supply the classical model [20, 21]; alternating duality alone is not positivity or purity. In a number-field construction these are obligations, not imported conclusions.

For a sharply testable final analytic target, put $F_\chi(z) = \Lambda(1/2 + iz, \chi)$. Suppose entire functions $h_m(z)\det(zI - H_m)$, with H_m finite self-adjoint and h_m entire nowhere zero, converge locally uniformly on $\mathbb{C}$ to this nonzero completed function. Hurwitz on every disk disjoint from $\mathbb{R}$ then proves all its zeros real, hence the Dirichlet critical-line assertion. Establishing this convergence with independently constructed H_m is exactly the missing load-bearing claim; a rank-one encoding or a reformulation does not establish it. Hecke, Dedekind and Artin extensions must state their analytic scope anew.

6.5 Whole components, derived endpoints and determinant relations

The full node lattice category has precisely the three indecomposables X, Λ, Y and arrows $X \to \Lambda \to Y$, $Y \to \Lambda \to X$. Its stable category has only X, Y, no irreducible arrows, and suspension exchanging them; its AR triangles have zero middle term. For the polynomial node at a smooth support $x = \lambda + t$, $y = 0$, $\lambda \neq 0$, the modules $T_r = k[t]/t^r$ form an ambient homogeneous tube:

$$0 \longrightarrow T_r \xrightarrow{(\pi,t)} T_{r-1} \oplus T_{r+1} \xrightarrow{(t,-\pi)} T_r \longrightarrow 0, \quad T_0 = 0, \quad \tau T_r = T_r.$$

The shift changes the derived placement, not the quasi-length. These nonzero-central-support objects do not extend to continuous finite-length modules of the arrow-adic completion. At its singular point even the candidate pure-branch tube fails ambient almost-splitness: $T_2^y \to k$ cannot factor through $T_2^x \to k$, since every intervening map lands in the x-socle, killed by the last map.

The nine HLS strings above sit in an entire intrinsic quotient component $T_{p,q}$, $(p, q) \in \mathbb{Z}^2$, with arrows increasing one coordinate and $\tau T_{p,q} = T_{p-1,q-1}$ [46]. Its central object $E = M(ab)$ has infinite projective dimension: restriction to the first corner gives the dual-number simple with its infinite loop resolution, while restricted projectives are free or zero. Its module AR sequence therefore yields a distinguished derived triangle but not a derived AR triangle ending at E. Ringel's four-quarter wind-wheel tile [43], with infinite-word diagonals and mixed center, is a different category's quilt; it is not silently identified with that HLS component or the Belyi surface.

There is no general equation "AR mesh equals polynomial semi-invariant syzygy." The intrinsic $k(1 \to 2)$ quotient of the infinite Chebyshev algebra has AR row $0 \to S_2 \to E \to S_1 \to 0$ and a free presentation determinant z on dimension $(1, 1)$, so its semi-invariant ring $k[z]$ has no generator relation. A genuine framed elliptic-square stratum has $Z = [x_1, x_2, y_1, y_2] \in \mathrm{Mat}_{2\times 4}$ with six minors and relation

$$p_{12}p_{34} - p_{13}p_{24} + p_{14}p_{23} = 0.$$

These minors have the same determinant character for the target GL_2 with framed source; they are not all semi-invariants for the full unframed vertex group. In contrast, the nodal lattice AR row gives a genuine relative determinant identity over R: multiplication by $a + bx$ on the basis $(y, 1)$ has matrix $\left(\begin{smallmatrix} a & -b \\ 0 & a+ub \end{smallmatrix}\right)$ and characteristic factor $(1 - Ta)(1 - T(a + ub))$. The endpoints have infinite order-projective dimension but are R-perfect.

Finally, the three-vertex theta algebra has no unital surjection onto a six-vertex radical-square-zero circular algebra: its semisimple image has at most three primitive idempotent images. Duplicating face vertex spaces instead gives an exact faithful unfolding which is not full; it does not inherit quotient or AR-transfer claims. Zwara Hom-control means that the Hom-dimension defect is bilinear on K_0, not merely a covering direct-sum formula [72, 73]. The book supplies explicit fixed-dimension counterexamples, corrected circular-complex incidence resolutions, complete drawn component windows and all category hypotheses. These additions do not supply the arithmetic determinant comparison.

7 A genuine theta covering and its categorical limits

Use column matrices, hence covariant quiver representations (left modules); opposite categories translate the right-module convention in [74]. Let A be the infinite elliptic theta dessin algebra: vertices $i \in \mathbb{Z}/3$, arrows $a_i, b_i : i \to i + 1$, and all mixed length-two paths zero. For the arrow ideal J and $L \geq 2$ put $B_L = A/J^L$. This is a finite string quotient of our actual infinite gentle algebra; for $L > 2$ its length-L relations prevent treating it as quadratic gentle.

Proposition 7.1 (A locally bounded integer cover)**.** *The category $\widetilde{B}_L$ with vertices $r \in \mathbb{Z}$, two arrows $a_r, b_r : r \to r+1$, and the same mixed and length-L zero relations, has orbit B_L under $r \mapsto r+3$. For finite-support representations, push-down F is exact and*

$$\operatorname{Hom}_{B_L}(FM, FN) \simeq \bigoplus_{g \in \mathbb{Z}} \operatorname{Hom}_{\widetilde{B}_L}(M, T_g N), \qquad (T_g N)_r = N_{r-3g}.$$

Only finitely many summands are nonzero. In general F is neither full, dense, nor Hom-controlled in Zwara's sense.

Proof. A nonzero path is constant-branch with length less than L. Thus the incoming and outgoing total Hom dimensions at a cover vertex are $2L-1$, and $\dim B_L = 3(2L-1)$. Paths and relations lift uniquely; the deck action is free and has the stated orbit. Push-down sums spaces of equal residue. A morphism decomposes uniquely into blocks with target lift index minus source index equal to $-3g$. The arrow commutation equations preserve that difference, proving the formula; finite supports bound the possible differences. Direct sums of vector spaces preserve exact sequences. The following explicit witnesses prove the three failures. □

For $L = 5$, let P be the cover interval on $0, 1, 2, 3, 4$ with all its a-maps one and b-maps zero, and let S be the sum of the five simples at the same vertices. They have the same K_0 class, but

	$\dim \operatorname{End}_{\widetilde{B}_5}$	$\dim \operatorname{End}_{B_5}(F(-))$	defect
S	5	9	4
P	1	2	1

Indeed the pushed-down simple multiplicities are $(2, 2, 1)$, whereas $\operatorname{End}_{B_5}(FP) = kI \oplus kX^3$ for its single nilpotent five-step chain X. The extra endomorphism X^3 proves non-fullness; the unequal defects at the same class exclude a bilinear K_0 defect. To use Zwara's finite-algebra definition literally, restrict the source to the finite convex interval $[0, 4]$, then extend by zero before push-down. This exact functor has the same endomorphism spaces and K_0 comparison.

Proposition 7.2 (An indecomposable module outside this cover image)**.** *Over a characteristic-zero field, B_3 has a nine-dimensional indecomposable representation which is not the push-down of any finite-support $\widetilde{B}_3$ representation.*

Proof. Label its basis by $0, \dots, 8$ with vertex residues $(0, 1, 0, 1, 2, 1, 2, 0, 2)$, and use matrix units E_{ij} acting on columns. Set

$$X = E_{10} + E_{32} + E_{43} + E_{65} + E_{76} + E_{08},$$
$$Y = E_{12} + E_{45} + E_{78}.$$

Their restrictions between adjacent vertex spaces give the a_i, b_i maps. They satisfy $XY = YX = X^3 = Y^3 = 0$, hence every length-three path is zero. The endomorphism ring preserving vertex spaces is $kI \oplus \mathfrak{r}$, where

$$\mathfrak{r} = \langle E_{02} + E_{13}, E_{15}, E_{35} + E_{46}, E_{48}, E_{68} + E_{70}, E_{72} \rangle, \qquad \mathfrak{r}^3 = 0.$$

Solving the arrow commutation equations gives exactly these seven basis elements. The ring is local: an element $cI + n$ is invertible exactly when $c \neq 0$, by the finite geometric series. It has no nontrivial idempotents, so the representation is indecomposable.

A finite-support lift would define an Euler operator D acting by r on the summand at lift vertex r. Then $[D, X] = X$ and $[D, Y] = Y$. Every displayed nonzero arrow is the unique nonzero entry in

its row and column in its own matrix. Its commutator entry is therefore $D_{ii} - D_{jj}$. Add the six X equations and subtract the three Y equations. Each diagonal term cancels, leaving $0 = 6 - 3 = 3$, a contradiction. This argument allows arbitrary D, not just a diagonal matrix in the displayed basis. □

Corollary 7.3 (Higher-genus and derived non-density). For the clean dessin with rotations $\sigma = (0 \cdots m-1)(m \cdots 2m-1)$ and $\alpha = \prod_{i=0}^{m-1}(i\, i+m)$, odd $m \geq 3$, and genus $(m-1)/2$, the analogous A_m/J^3 has a $3m$-dimensional representation outside its integer-cover image in characteristic zero. Its bounded-derived push-down is not dense either.

Proof. Use basis $v_0, \ldots, v_{3m-1}$, assign vertices $i, i+1, i$ to $v_{3i}, v_{3i+1}, v_{3i+2}$ modulo m, and set

$$Xv_{3i} = v_{3i+1}, \quad Xv_{3i+2} = v_{3i+3}, \quad Yv_{3i+2} = v_{3i+1},$$

with indices modulo $3m$ and all other values zero. Again mixed products and cubes vanish. An Euler operator would yield $2m - m = m = 0$ by the same telescoping calculation, impossible in characteristic zero. If a bounded finite-support complex pushed to this module in degree zero, exactness would make its zeroth cohomology a module lift, a contradiction. □

The source's derived result is a *precovering* in the stated bounded categories [74, Theorem 4.28]; density remains an additional condition. Neither this result nor the GLS equivariant classification [75] identifies our completed order with a GLS algebra. Graded surface/Fukaya models [76] likewise require their own dg category, stops and grading. These categorical tools do not construct the independent positive arithmetic determinant.

8 Verification and research frontier

The two nodal/invariant Lean files construct the node as an actual subring of $k[[X]] \times k[[X]]$ with equal constants, prove its annihilators, regular common parameter, augmentation, alternating projective resolution and a bijective pair of power-series coordinates. A second file proves CK1 collapse under the stated permutation/coverage hypotheses, determinant base change, actual 2×2 symmetric-square functoriality and relation preservation, a Schur character identity and the fixed-parameter lattice failure. These thirty printed theorem types have only standard Lean foundations. They do not formalize general dessins, abstract Ext/AR/Schur–Weyl, C* exactness, Fredholm determinants, arithmetic descent or GRH. Earlier formal layers remain preserved with their literal scopes.

Exact-rational diagnostics independently check the displayed dessins, path identities, Schur and Young examples and the nine-object grid. These finite checks supplement the proofs, rather than prove infinite spectral or arithmetic assertions. The next arithmetic target is an exact descent/Frobenius-compatible construction on the labelled order/Fock data, with a determinant-preserving positive comparison. The local Schur kernel and the simple degree spectrum alone do not supply it. All source clones and earlier research are retained; the earlier general focused PDFs are archived at the preceding commit.

Ten further declarations in `DessinPullbackComponents.lean` check actual matrix-unit identities, nonmultiplicativity of raw diagonal extraction, impossibility of a unital two-state matrix residue into a nonzero commutative ring, multiplicativity on diagonal fibers, and genuine equal-image pullback subring/projection components. The analytic suite then contained187printed declarations at unchanged pins. The circle-function C* model, functional calculus, generation, graph uniqueness

and K-theory remain prose/external-theorem arguments, not hidden assumptions of these component checks.

Sixteen additional declarations in `ThetaJacobi.lean` check genuine finite-field MulChar Jacobi conjugation, norm and rescaling with explicit nontriviality hypotheses; cubic–quadratic noncancellation; the cyclotomic multiplication determinant and positive Gram identity; and actual theta cellular boundary, homology-inclusion, restriction and rational chain-map matrices. The full analytic suite now contains203declarations. The point-count proof, deck/Kummer comparison, cohomological trace interpretation and global arithmetic comparison are not thereby fully formalized.

The final three `ThetaJacobi.lean` declarations verify the actual conjugation matrix identities and the real two-dimensional inert-prime determinant obstruction. They do not assume any unknown zero location or exclude complex/enlarged spectral realizations.

`ThetaProjectiveMultiplicity.lean` adds five declarations. Arbitrary upper-block matrices with six identical diagonal blocks have the sixth-power determinant and characteristic-factor formulas. A literal power-series constant-coefficient map specializes matrix determinants, for each scalar parameter; the coefficientwise polynomial/order interface remains prose. A nonzero quadratic cannot be a rational-function sixth power, proved by polynomial degrees. Three genuine power series with product X cannot have the cyclically equivalent residue-vanishing conditions forced by invertible rank-one arrow intertwiners. The derivation of the path blocks from the full theta order, projective decomposition, derived branch functor and deck intertwiners is not fully formalized. The complete analytic suite now has208printed declarations; no hidden arithmetic positivity assumption occurs.

`ThetaTameDescent.lean` adds twelve printed component declarations (and one auxiliary polynomial lemma). They use genuine formal-power-series rescaling and its coefficient operator, proving the coefficient action, invariant-coefficient vanishing and the tame operator's cube. Actual three-state matrices satisfy the clock/rotation commutator, comparison determinant and closed-fiber equations, weighted-cycle cube and full characteristic polynomial, noncommutation of a nonconstant rotation polynomial, trivial whole-column determinant actions, and the same-operator comparison for vC. The classification of all fixed series by residues, full order action and hereditary invariant algebra, stack interpretation, valuation/rank theorem, general coefficient-functor filtration and GRH are not thereby fully formalized. No unknown positivity premise is assumed.

9 Why an absolute coprime pair-square estimate cannot close the short rows

Keep the original sextic family

$$M_u(D) = D^{-1/2} \sum_n \mu(n)\nu(n) W(Nn/D)\chi_n(u), \qquad \chi_n(u) = (u/n)_6,$$

with squarefree primary column ideals outside the fixed excluded set, their full annular coefficients (fix W supported in $[1,2]$), and all nonzero Eisenstein element rows $Nu \le H$. The cubic symbol is χ_n^2, not χ_n. The following classical frame inequality [80] gives a new obstruction to one proposed estimate in this arithmetic program.

Proposition 9.1 (Coprime pair-square obstruction)**.** *Let A be an R-by-N complex matrix with $|A_{ui}| = 1$, and put $K_{ij} = \sum_u A_{ui}\overline{A_{uj}}$. Then*

$$\sum_{i \ne j} |K_{ij}|^2 \ge NR(N-R).$$

For the original character kernels, let $B_1(H,D)$ be the sum of $|K_H(n,m)|^2$ over all distinct coprime squarefree column ideals in $[D,2D]$. At $H=D^{3/4}$,

$$B_1(H,D) \gg \frac{D^{11/4}}{(\log D)^2}.$$

In particular the proposed sufficient input $B_1(H,D) \ll H^2 D^{\varepsilon}$ fails for each fixed $\varepsilon < 5/4$. This does not refute the signed Möbius mean square.

Proof. Set $F_{uv} = \sum_i A_{ui}\overline{A_{vi}}$. Reordering four finite sums proves $\sum_{i,j} |K_{ij}|^2 = \sum_{u,v} |F_{uv}|^2$. The row diagonal contributes RN^2; the column diagonal contributes NR^2. Removing the latter and retaining the former gives the inequality.

Restrict the column pairs in the nonnegative sum B_1 to distinct prime ideals p, q of norm in $[D,2D]$, outside the same fixed set. If $H < D$, no such prime divides a nonzero row element: divisibility would imply $Nu \geq Np$. Thus all original zero masks are exactly one on this ball, $|\chi_p(u)| = 1$, and the residual primitive finite character conductor is pq, of norm between D^2 and $4D^2$. All these pairs have common divisor1. The lattice count gives $R = (2\pi/\sqrt{3})H + O(\sqrt{H}+1)$. The classical unconditional fixed-field prime-ideal theorem gives $N \sim D/\log D$; this is an external analytic input, not a GRH hypothesis. At the stated endpoint $R = o(N)$, and the bound is therefore $(1-o(1))N^2R \gg D^{11/4}/(\log D)^2$. □

To see the exact failed step, write $c_n = \mu(n)\nu(n)W(Nn/D)$ and $C_D = \sum_n |c_n|^2 = O(D)$. Cauchy over coprime pairs bounds the absolute signed contribution by $D^{-1}C_D B_1^{1/2}$. The hoped-for pair-square upper bound would have closed this particular route, but is false. A large absolute pair-square mass is not a lower bound on the quadratic form with the fixed c_n. Nor is the prime part a positive summand of the full Möbius energy: prime–composite cross terms can cancel. The original low-gcd estimate and the positive arithmetic determinant bridge remain open.

10 A signed Möbius decomposition and its termwise obstruction

Keep the original squarefree good-ideal family, its fixed ν, its full $W(Nn/D)$ and every zero-extended sextic symbol. In ideal Dirichlet convolution, write $\zeta_0(n) = 1$ for every good ideal and let δ be the convolution unit. For $1 \leq U < D$ put $a_U = \mu \mathbf{1}_{Nn \leq U}$ and $b_U = \mu - a_U$. The classical Vaughan algebra gives

$$\mu = 2a_U - a_U * a_U * \zeta_0 + b_U * b_U * \zeta_0. \tag{1}$$

Indeed $\mu * \zeta_0 = \delta$, and expansion of the last product proves the identity without an analytic assumption. This is a classical decomposition, not a new cancellation theorem; compare [87, Lemma4.1].

On squarefree n all factors in $n = dc = abc$ are pairwise coprime. Define the actual, nonarbitrary coefficient functions

$$\beta_U(d) = \sum_{\substack{ab=d \\ Na, Nb \leq U}} \mu(a)\mu(b), \qquad g_U(s) = \sum_{\substack{b \mid s \\ Nb > U}} \mu(b).$$

For squarefree d, $\beta_U(d)$ is $\mu(d)$ times the number of the displayed ordered factorizations. For

$Ns > U$, $g_U(s) = -\sum_{b|s,\, Nb\le U} \mu(b)$. The two regrouped pieces are

$$I_u = D^{-1/2} \sum_{\substack{d,c \text{ squarefree} \\ (d,c)=1}} \beta_U(d)\nu(dc)\chi_d(u)\chi_c(u)W(NdNc/D),$$

$$II_u = D^{-1/2} \sum_{\substack{a,s \text{ squarefree} \\ Na,Ns>U,\,(a,s)=1}} \mu(a)g_U(s)\nu(as)\chi_a(u)\chi_s(u)W(NaNs/D).$$

Then $M_u = -I_u + II_u$. All ideals here remain outside the same fixed excluded set. The short $2a_U$ term vanishes because $U < D$ and W is supported in $[1,2]$. No phase or coprimality mask was removed.

Proposition 10.1 (An all-cutoff separate-piece bound is false)**.** *Take the allowed principal ν, $U = 1$, and a fixed nonnegative smooth W supported in $[1,2]$, bounded below by a positive constant on a closed subinterval of $(1,2)$ of positive length. For each of the two pieces,*

$$\sum_{0<Nu\le H} |I_u|^2 \gg_W \frac{D}{\log^2 D}, \qquad \sum_{0<Nu\le H} |II_u|^2 \gg_W \frac{D}{\log^2 D} \quad (H \ge 1).$$

Consequently a proposed bound HD^{ε} for each piece at $H = D^{3/4}$ fails for $\varepsilon < 1/4$. In the exact original notation $H(HD)^{\varepsilon}$, it fails for $\varepsilon < 1/7$. This refutes an all-cutoff termwise strengthening, not the original signed mean square or a specified moving-cutoff argument.

Proof. At $U = 1$, $a_U = \delta$, so $a_U * a_U * \zeta_0 = \zeta_0$ and $b_U * b_U * \zeta_0 = \mu + \zeta_0 - 2\delta$. On the annulus the two coefficients are 1 and $1 + \mu(n)$, respectively. On squarefree ideals the latter is zero or two. At the actual unit row $u = 1$, every original character equals one. A fixed-field prime-ideal theorem gives $\gg D/\log D$ good prime columns in the subinterval, so $I_1 \gg_W \sqrt{D}/\log D$. Fix a good prime ideal q_0. Products $q_0 p$ with varying distinct good primes in the rescaled subinterval have $\mu(q_0 p) = 1$. Their nonnegative contributions give $II_1 \gg_W \sqrt{D}/\log D$. Keeping that one row in each nonnegative energy proves the result. The prime-ideal theorem is a classical unconditional external input, with the field and q_0 fixed. □

The factorization does not shorten the primitive row conductor: for squarefree coprime a, s it is as, not a or s separately. For two resulting columns n, n' with common factor h, the oscillatory conductor is nn'/h^2 and a zero-preserving period is nn'/h. The signed low-gcd estimate and the independent positive arithmetic determinant comparison therefore remain open.

11 A full-mask Hecke expansion for the type-I piece

We now continue the signed decomposition through its actual squarefree inner sum. Fix $K = \mathbb{Q}(\sqrt{-3})$, the finite-order Hecke character ν, and the same finite excluded prime set S, containing primes above 6 and the conductor of ν. For $u \in \mathcal{O}_K \setminus \{0\}$, let ψ_u be the primitive Hecke character induced by $\nu(n)(u/n)_6$ away from $S \cup \mathrm{supp}(u)$. It is defined before, and independently of, the type-I divisor d. The abelian Kummer extension $K(u^{1/6})/K$ constructs the sextic factor; multiplying by ν gives the stated character. This uses classical abelian reciprocity, not general Artin holomorphy. Write $\psi_{u,2}$ for the primitive character inducing its square and $\mathfrak{f}_u, \mathfrak{f}_{u,2}$ for the two primitive conductors. For a squarefree good ideal d put

$$T = T(u,d) = S \cup \mathrm{supp}(u) \cup \mathrm{supp}(d), \qquad R_T = \prod_{p\in T} Np, \quad Q_u = |D_K| N\mathfrak{f}_u.$$

All primes in T are deleted even if absent from either primitive conductor.

Theorem 11.1 (Restoration and a uniform scalar contour bound)**.** *For $\Re s > 1$ the original type-I inner Dirichlet series is*

$$F_{u,d}(s) = \sum_{\substack{c \text{ squarefree} \\ (c,T)=1}} \frac{\psi_u(c)}{(Nc)^s} = \frac{L(s,\psi_u)}{L(2s,\psi_{u,2})}\, C_T(s), \tag{2}$$

$$C_T(s) = \frac{\displaystyle\prod_{p\in T,\ p\nmid \mathfrak{f}_u} (1-\psi_u(p)(Np)^{-s})}{\displaystyle\prod_{p\in T,\ p\nmid \mathfrak{f}_{u,2}} (1-\psi_{u,2}(p)(Np)^{-2s})}. \tag{3}$$

It continues meromorphically to $\Re s > 1/2$, with only a possible simple pole at $s = 1$ when $\psi_u = 1$. In that case its residue is

$$\rho_T = \frac{\operatorname{res}_{s=1}\zeta_K(s)}{\zeta_K(2)} \prod_{p\in T}(1+(Np)^{-1})^{-1}.$$

Fix $0 < \delta < 1/2$, $\varepsilon > 0$ and $W \in C_c^\infty((0,\infty))$ supported in $[1,2]$. Set

$$\mathcal{M}_4(W) = \sum_{j=0}^{4} \sup_x |(x\partial_x)^j W(x)|.$$

Uniformly for $Y \geq 1$, u and d,

$$\sum_{\substack{c \text{ squarefree} \\ (c,T)=1}} \psi_u(c) W(Nc/Y) = \mathbf{1}_{\psi_u=1}\rho_T Y \widehat{W}(1) + O_{\nu,S,\delta,\varepsilon}\Big(\mathcal{M}_4(W) Y^{1/2+\delta} Q_u^{1/4-\delta/2+\varepsilon} R_T^{\varepsilon}\Big). \tag{4}$$

Here $\widehat{W}(s) = \int_0^\infty W(x)x^{s-1}\,dx$. The continuation and functional equation of degree-one Hecke functions are classical external inputs; the contour argument is given below.

At a good prime with $v_p(u) = 3$, ψ_u is ramified while its square is unramified; the second product in (3) must restore a factor invisible to squaring the zero-extended primitive value. At $v_p(u) = 6$ both characters can forget the prime, but the original row still vanishes on columns divisible by it. In particular, for principal ν, $u = \pi^3$ and $u = \pi^6$ at $N\pi = 7$, the deleted local factors are

	primitive ratio at $s = 1$	restoration	full masked factor
$v_p(u) = 3$	48/49	49/48	1
$v_p(u) = 6$	8/7	7/8	1

These are exact local calculations, not numerical evidence about zeros.

For $1 \leq U$, $U^2 \leq D$, insert $Y = D/Nd$ into (4) and retain the actual $\beta_U(d)$ and outer $\nu(d)\chi_d(u)$. Classical tame Kummer conductors give $Q_u \ll_\nu Nu$. Thus the normalized I_u equals its displayed principal-residue sum plus an error E_u with

$$|E_u| \ll_{\nu,S,\delta,\varepsilon} \mathcal{M}_4(W) D^\delta U^{1-2\delta+\varepsilon} (Nu)^{1/4-\delta/2+\varepsilon}. \tag{5}$$

The resulting elementary energy envelope is only

$$\sum_{0<Nu\leq H} |E_u|^2 \ll \mathcal{M}_4(W)^2 D^{2\delta} U^{2-4\delta+\varepsilon} H^{3/2-\delta+\varepsilon}. \tag{6}$$

It does not prove the short-row target of order H up to small losses. This is an insufficient upper bound, not a lower bound on the actual error or a counterexample to coefficient-sensitive averaging. The original signed mean square and positive determinant bridge stay open.

12 Actual renormalization retains the effective action

A compression score does not specify an RG transformation. Following the real-space formulation of Wilson and Kogut [81], we specify the eliminated variables, effective action, normalization and scale map. Our exact finite decimation is distinct from an approximate tensor-network truncation such as Levin–Nave TRG [82].

Definition 12.1 (Finite classical RG step)**.** Let the fine configuration be $(s,t) \in S \times E$, with finite nonempty sets S, E, and dimensionless log Boltzmann weight $V(s,t) = -\beta H(s,t)$. Define

$$(\mathcal{R}V)(s) = \log \sum_{t \in E} e^{V(s,t)}, \qquad Z(V) = \sum_{s,t} e^{V(s,t)}.$$

The full action includes its additive constant. A truncation or a learned approximation is a further operation with its own error, not this identity.

Proposition 12.2 (Partition, observables and composition)**.** *One has $Z(\mathcal{R}V) = Z(V)$, and the normalized expectation of every observable depending only on s agrees in the two descriptions. Eliminating E_1 and then E_2 equals eliminating $E_1 \times E_2$ at once, with the same tensor labels and inverse temperature.*

Proof. Exponentiating the definition gives $e^{\mathcal{R}V(s)} = \sum_t e^{V(s,t)}$. Sum over s, optionally multiplying by an observable $O(s)$, and divide both sides by the same strictly positive Z. Reordering finite sums proves the composition statement. No projection onto pair interactions is used. □

For an even homogeneous periodic Ising chain, write the transfer matrix as $T_{ab} = \exp(C + Kab + h(a+b)/2)$, $a, b \in \{-1, 1\}$. Its partition function is $\operatorname{tr} T^N$. Summing alternate spins replaces T by T^2 and N by $N/2$; the physical spacing doubles. Writing $P = T^2$, its positive entries determine

$$K' = \tfrac{1}{4} \log \frac{P_{++}P_{--}}{P_{+-}^2}, \quad h' = \tfrac{1}{2} \log \frac{P_{++}}{P_{--}}, \quad C' = \tfrac{1}{4} \log(P_{++}P_{--}P_{+-}^2).$$

These three parameters reproduce every entry of P, so no normalization is lost. At $h = 0$, $\tanh K' = \tanh^2 K$ and the eigenvalue-ratio correlation length of the homogeneous extension in blocked units halves. Its product with the spacing stays fixed. The fixed points in $t = \tanh K$ are $t = 0, 1$; the latter is zero temperature, not a positive-temperature transition. Odd dessin examples below use the full-action definition, not this even-chain shortcut.

12.1 Genuine dessin incidence models

For odd m, the clean dessin $y^2 = x^m - 1$, $\beta = x^m$ has two black vertices, m white vertices and one face; its genus is $(m-1)/2$. Its incidence graph is $K_{2,m}$ with $2m$ clean edges. Its infinite locally gentle order quiver instead has m vertices and completed DVR-rank $2m^2$. We place physical spins $B_+, B_-, s_0, \ldots, s_{m-1}$ on the incidence vertices; this is a specified finite spin model, not a faithful representation of that infinite algebra or order. The abstract incidence graph alone does not recover the ribbon embedding or genus.

With $\eta_j = (-1)^j$ and fixed nonuniform fields h_j, choose

$$V = J_+ B_+ \sum_j s_j + J_- B_- \sum_j \eta_j s_j + bB_+ - \tfrac{3}{5} bB_- + \sum_j h_j s_j.$$

Summing the two black spins gives the *entire* effective action

$$V'(s) = \sum_j h_j s_j + \log\left[2\cosh\left(b + J_+ \sum_j s_j\right)\right] + \log\left[2\cosh\left(-\tfrac{3}{5}b + J_- \sum_j \eta_j s_j\right)\right].$$

Every one of the $2m$ physical edges is present, although its couplings belong to a tied three-parameter family. The Walsh expansion $V'(s) = \sum_{A\subseteq\{0,\dots,m-1\}} c_A \prod_{j\in A} s_j$, with $c_A = 2^{-m}\sum_s V'(s)\prod_{j\in A} s_j$, is unique by Boolean orthogonality. It retains the constant $c_\varnothing$ and genuinely generated higher-body interactions. Our genus-two and genus-four examples use $m = 5, 9$ and continue by exact nested marginalization. Dropping all terms of degree greater than two is displayed as a measured approximation.

12.2 Quantum thermal effective Hamiltonians

For a finite Hermitian Hamiltonian H on nonzero finite-dimensional $\mathcal{H}_S \otimes \mathcal{H}_E$ and $\beta > 0$, define

$$W' = \operatorname{Tr}_E e^{-\beta H}, \qquad H_{\text{eff}}(\beta) = -\beta^{-1}\log W'.$$

The matrix W' is strictly positive: for $v \neq 0$, sum the strictly positive quadratic forms of $v \otimes e_j$ over an orthonormal basis of $\mathcal{H}_E$. Its Hermitian spectral logarithm is therefore defined. The identity $e^{-\beta H_{\text{eff}}} = W'$ proves partition and retained *equal-time* observable preservation by the partial-trace property. Repeated traces compose. The full Pauli expansion, including its identity coefficient, is retained. It is generally nonlocal and depends on β; this statement does not preserve all fine real-time dynamics or supply a local zero-temperature/thermodynamic RG fixed point.

`FiniteRG.lean` checks nine actual finite scalar marginal, exponential-action, partition, observable, composition, error-sandwich and transfer-trace statements. The quantum spectral-log and complete thermodynamic/field-theory interfaces are not formalized by these lemmas. These physics constructions give no arithmetic determinant comparison or new GRH conclusion.

13 Linear relations and functorial branch filtrations

The original linear-relations chapter mixes graphs, partial relations and stable kernels. We keep these different objects separate. For a field k, a linear relation is a subspace $C \subseteq V \oplus V$; its inverse swaps coordinates and $DC = \{(x,z) : \exists y, (x,y) \in C, (y,z) \in D\}$. The full graph $\Gamma_T = \{(x, Tx) : x \in V\}$ always has dimension $\dim V$, including pairs with zero output. In particular $\Gamma_T^n(0) = 0$, whereas $(\Gamma_T^{-1})^n(0) = \ker T^n$. Thus the source's $\kappa(\Gamma_T) = \bigcup_n \Gamma_T^n(0)$ is zero, not the ordinary stable kernel.

For odd $m \geq 3$, use the actual clean dessin of $y^2 = x^m - 1$, $\beta = x^m$, on the compact smooth curve, in characteristic zero. Its genus is $(m-1)/2$; its infinite locally gentle algebra A_m has vertices $i \in \mathbb{Z}/m$, arrows $a_i, b_i : i \to i-1$, and mixed consecutive branch products zero. Below paths act in their direction of travel on left representations over any coefficient field k. Let $\mathcal{E}_m$ be the full category of finite-dimensional representations with every $b_i = 0$ and a_i invertible for $i \neq 0$.

Theorem 13.1 (A concrete relation category on an infinite dessin algebra)**.** *The category $\mathcal{E}_m$ is equivalent to finite-dimensional pairs (V, T) with intertwining maps. A normal form $N_m(T)$ puts V at each vertex, $a_0 = T$, and all other $a_i = 1$. For $n = \dim V$,*

$$V = K(T) \oplus I(T), \qquad K(T) = \ker T^n, \quad I(T) = \operatorname{im} T^n.$$

These summands and their kernel/image filtrations are functorial. The first restriction is nilpotent and the second invertible. The central full-turn parameter u acts as T. A finite $N_m(T)$ extends

continuously to the u-adically completed order precisely when T is nilpotent. Such modules are torsion, not nonzero order lattices.

Proof. Transport all vertex spaces to vertex zero along the invertible arrows; the remaining map is the full-turn monodromy T. Arrow-compatible maps become exactly $hT = T'h$. Finite kernel/image chains stabilize by step n. If $v = T^n w$ and $T^n v = 0$, then $w \in \ker T^{2n} = \ker T^n$, hence $v = 0$; rank-nullity proves the direct sum. Stability gives the two restrictions, and $hT^r = T'^r h$ proves functoriality. Continuity on a finite-dimensional discrete space means some u^r kills it. Conversely nilpotence makes every formal series act by a finite sum. This proves the completion assertion. □

Over an algebraically closed k, the indecomposable branch pairs are $J_r(\lambda)$. The value $\lambda = 1$ is permitted for a polynomial band; it is not a forbidden deformation parameter. The nilpotent blocks are strings, but are not all rigid: $N_m(J_2(0))$ is a nonsplit self-extension of $N_m(J_1(0))$. The Hom dimension between two such Jordan pairs is $\min(r, s)$ for equal eigenvalues and zero otherwise. These assertions concern this specified full subcategory, not all infinite modules.

If L transports once along the a-arrows on $\bigoplus_i V$, then

$$\det(1 - zL) = \det(1 - z^m T). \tag{7}$$

Eliminating the identity arrow blocks leaves the final monodromy block; the eliminated identity blocks have determinant one. For $J_r(\lambda)$ the answer is $(1 - \lambda z^m)^r$. Nilpotent strings therefore contribute one, and arbitrary bands have no forced unit-circle zero location. This determinant is an actual cyclic-quiver invariant; it is not the missing arithmetic positive determinant comparison.

14 A kernel-checked extension over the infinite dessin algebra

The preceding branch example can be realized on every element of the infinite algebra, rather than verified only on sampled arrow matrices. Use the complex genus-two clean dessin $y^2 = x^5 - 1$, $\beta = x^5$, and any coefficient field k. Put $H \subseteq M_5(k[t])$ equal to the matrices whose reduction modulo t is upper triangular. Let

$$B = H \times_{k^5} H,$$

where each map to k^5 takes the constant diagonal. Equivalently, B is generated inside $M_5(k[t])^2$ by

$$e_i = (E_{ii}, E_{ii}), \quad a_i = (w_i E_{i-1,i}, 0), \quad b_i = (0, w_i E_{i-1,i}), \qquad w_0 = t, \quad w_i = 1 \ (i \neq 0).$$

Indices are modulo five. Matrix multiplication acts on left column modules. The path-normal-form proof below identifies B with A_5; this is a polynomial algebra, not an arrow-truncated finite quotient.

Theorem 14.1 (An ambient module extension)**.** *Let $D = k[\epsilon]/(\epsilon^2)$, $M_1 = k^5$ and $M_2 = D^5$. For $(C(t), D'(t)) \in B$, act on M_1 by $C(0)$ and on M_2 by $C(\epsilon)$. These are unital B-modules, and*

$$0 \longrightarrow M_1 \xrightarrow{\ i\ } M_2 \xrightarrow{\ p\ } M_1 \longrightarrow 0, \qquad i(x) = \epsilon x, \quad p(x + \epsilon y) = x,$$

is exact and nonsplit. Moreover B is infinite dimensional over k. The assertions hold in every characteristic, including two.

Proof. Evaluation of polynomials gives both actions. Multiplying $C(\epsilon) = C_0 + \epsilon C_1$ proves B-linearity of i, p; their kernel and image follow coordinatewise. If s were a B-linear section, the constant coefficient of $(se_0)_0$ would be one. But a_0 kills M_1, while $(a_0 s e_0)_4 = \epsilon (se_0)_0$ has nonzero ϵ coefficient, a contradiction. Finally the permitted turn $c = a_1 a_2 a_3 a_4 a_0$ has first component tE_{00}. The unital algebra map $k[t] \to B$, $f \mapsto f(c)$, is injective: its $(0,0)$ first-component entry is $f(t)$. Thus B cannot be finite dimensional. □

In the preceding notation these modules are $N_5(J_1(0))$ and $N_5(J_2(0))$. They are finite torsion modules over an infinite algebra; they are not nonzero lattices over its completed surface order. The nonsplit sequence defines a nonzero Yoneda extension class in the ambient module category. No assertion that it is almost split in that entire category is needed or inferred. In particular it does not supply an arithmetic spectral determinant or a zero-location theorem.

Theorem 14.2 (Unbounded normal-image faithfulness)**.** *In $M_5(k[t])^2$, index the stationary images by i and the positive images by (b,i,j,q), where $b \in \{a,b\}$, $0 \le i,j < 5$, $q \ge 0$ and $i < j$ or $q > 0$. The positive image is $t^q E_{ij}$ in factor b, with zero in the other factor. Its path length is $5q+j-i > 0$. All these images, including the shared stationary images, are linearly independent over every field. Every finite normal expansion has uniquely recoverable coefficients, without a bound on degree or path length.*

Proof. Read a stationary coefficient from the first factor's constant diagonal; read a positive coefficient from the indicated factor, entry and degree. The index restriction ensures that no positive image contributes to a constant diagonal. These linear selectors are biorthogonal to the images, so applying each selector to a finite expansion recovers its coefficient. □

This theorem is now kernel checked for actual polynomial matrix pairs. The membership and spanning steps are supplied by the next theorem. Comparison with a separately constructed path quotient remains a distinct algebraic formalization task.

Theorem 14.3 (Normal coordinates in the generated algebra)**.** *Let $A \subseteq M_5(k[t])^2$ be the algebra generated by the displayed vertices and arrows, and let $\mathcal{K}$ be the normal index set of Theorem 14.2. Every normal image belongs to A. The actual A-valued finite expansion*

$$E : k^{(\mathcal{K})} \longrightarrow A, \qquad c \longmapsto \sum_{x \in \mathcal{K}} c_x v_x$$

is a linear isomorphism, whose inverse is $(E^{-1}a)_x = \lambda_x(a)$. Thus the normal images form a basis of the infinite generated algebra, not merely an independent family in its ambient matrix space.

Proof. Write $P_b(i,j)$ for the shortest positive branch walk from j to i, using a full turn when $i = j$. Its image is $t^d E_{ij}$ in branch b, where $d = 0$ if $i < j$ and $d = 1$ otherwise. The actual arrow product $P_b(i,i)^n P_b(i,j)$ has image $t^{n+d}E_{ij}$ for every $n \ge 0$. Admissibility gives $q \ge d$, so $n = q - d$ proves membership.

The normal span contains the generators and $1 = \sum e_i$ and is multiplicatively closed: stationary factors select endpoints; positive factors either multiply to zero or to $t^{q+r}E_{il}$ in the same branch. When $q + r = 0$, the two strict upper inequalities imply $i < l$, so this output slot is still permitted. Adjoin induction gives spanning of A. The previous coefficient selectors prove uniqueness and the inverse formula. □

All membership, spanning and finite-expansion assertions in this theorem are now Lean checked over arbitrary fields. The formal linear equivalence is not asserted to be an algebra map from an independently defined path quotient, nor a statement about completed orders, derived categories or GRH.

15 Learned fields, width limits and quantum geometry

Learned functions enter this program as candidate generators, physical surrogates and control laws. A network may suggest an arithmetic metric, operator or gluing datum; a successful fit does not

establish its positivity, analytic domain or identification with a completed L-function. Those are independent mathematical obligations. This section gives explicit interfaces and executed examples rather than treating "quantum", "tropical" or "infinite width" as a substitute for that comparison.

Proposition 15.1 (A specified quantum-feature width limit)**.** *Let $x \in \mathbb{R}^d$, $w \sim N(0, I_d)$ and $b \sim \mathrm{Unif}[0, 2\pi]$ be independent, and set $\phi_{w,b}(x) = \sqrt{2}\langle Z\rangle_{R_Y(w\cdot x+b)|0\rangle} = \sqrt{2}\cos(w \cdot x + b)$. For independent features,*

$$K_m(x, y) = m^{-1} \sum_{r=1}^{m} \phi_r(x)\phi_r(y) \longrightarrow K(x, y) = e^{-\|x-y\|^2/2}$$

almost surely on every specified finite set of pairs. Every K_m is a positive semidefinite Gram kernel. For a readout $f_a(x) = m^{-1/2} \sum_r a_r \phi_r(x)$ with only a trained, its tangent kernel is exactly K_m. These statements do not assert the same limit for a jointly trained finite quantum transformer.

Proof. Uniform-phase integration gives $\mathbb{E}_b[2\cos(u + b)\cos(v + b)] = \cos(u - v)$. The Gaussian characteristic function then gives K. Products are bounded by two, so the strong law applies to each pair and hence any finite union. For any finite coefficient vector c, the Gram quadratic form is $m^{-1} \sum_r |\sum_i c_i \phi_r(x_i)|^2 \geq 0$. Differentiating with respect to a_r gives the final assertion. □

The frozen-kernel squared-loss flow has residual $r(t) = \exp(-K_m t)r(0)$ when the loss normalization is absorbed into time. Null directions do not decay. A complex feature space instead uses the Hermitian kernel $\mathbb{E}[\phi(x)\overline{\phi(y)}]$. A finite coefficient field has neither this Gaussian law nor an ordered positive Hilbert metric; changing algebraic coefficients is a separate operation.

Proposition 15.2 (Tropical limit and a cancellation boundary)**.** *For real $s_1, \ldots, s_n$ and $\tau > 0$,*

$$0 \leq \tau \log \sum_j e^{s_j/\tau} - \max_j s_j \leq \tau \log n.$$

An attention maximum therefore has a controlled log-sum-exp approximation. It does not by itself define a tropicalization of a variety or an HMS functor. In a valued coefficient field, valuation of a sum can increase by cancellation.

Proof. After subtracting $M = \max s_j$, the exponential sum lies between one and n. For the second statement, both $1+T$ and $-1+T$ have valuation zero in $\mathbb{Q}((T))$, whereas their sum $2T$ has valuation one. In characteristic two that sum is zero. Thus replacing every addition by its tropical minimum without tracking leading coefficients loses information. □

Kontsevich–Soibelman constructions require valued analytic and integral-affine gluing data; appropriate gentle/Fukaya comparisons require specified categories, gradings, stops and smoothness hypotheses [92, 91, 93]. Learned coefficients can propose such data, but the compatibility equations and arithmetic comparison still need proofs.

Executed physical examples. On the actual clean genus-two dessin, put seven quantum spins on its $K_{2,5}$ incidence graph and compute a normalized Gibbs state. Exact white-spin partial trace gives a two-spin quantum RG marginal. Twelve RTX4090 runs train four-qubit-feature diffusion and flow-matching models for its smoothed Z-measurement law: two objectives, entangled/product circuit controls, and three seeds. They have 5,810 nominal parameters and 2,000 steps each. A separate four-qubit variational purification flow learns the full known density interpolation, with

48 controls. Its minimum fidelity on a 33-point grid is 0.999903; this is not a uniform theorem or universal noise inverse. The matched controls show no consistent entanglement advantage.

Actual cuStateVec/cuTensorNet execution also covers 7-, 11- and 23-qubit finite circuits on genus-two, genus-four and genus-ten dessin incidence graphs. The 23-qubit case has 8,388,608 amplitudes and 447 gates. These larger circuits are untrained; the small cases have independent NumPy comparisons, while the largest has normalization and inverse-circuit checks. The exact environment and all failed/preliminary records are preserved.

Geometric and hardware interfaces. For the contracted m-edge dessin, the graph fundamental group is F_{m-1}, and its universal graph cover is the m-regular tree. The theta case $m = 3$ has a three-regular cover, distinct from the four-regular Cayley graph of F_2. Neither is the coefficient field $\mathbb{F}_2$ or the universal cover of the filled compact surface. Explicit path modules over the infinite genus-two algebra give Hom, Ext and Tor computations; Frobenius pants tensors give exact TQFT gluing, with a heralded physical implementation. CHSH gives a finite game benchmark, distinct from the unrestricted correlation separation in MIP*=RE [99]. Three executable tropical hardware references and a source-pinned chip-design survey complement the circuits. They are arithmetic/control designs, not synthesized chips, validated CFT/string dualities or a proof of GRH.

16 Complementary physical datasets and checked dynamics

The learned program now has two additional, separately constructed corpora: 16,384 effective-material response records and22,951 molecular conformer/force records in their default configurations. Time samples are not counted as independent systems. Source-linked and dynamical configurations are additional. The active HolonForge/Aster and Mora-3 repositories were read only; no data, vocabulary, checkpoint, source enrollment or training process was changed.

The material corpus covers SSH, Kitaev, QWZ, Rice–Mele, Harper and actual clean-dessin classical RG models. The molecular corpus includes designed amides and peptides, with explicit-H MMFF94 calculations, geometric persistence, and method-labelled conformations. Forty-eight existing Aster MP20 structures, 32 existing Mora PubChem identities and eight public PDB structures supply separately labelled anchors. Their new computed responses are not new experimental observations, DFT labels, binding affinities or CISS measurements.

All conformer, gauge, intervention and task descendants share a partition. Specified physical/chemical similarity edges are closed transitively before component-level stratification. Large components are kept intact even when 80/10/10 proportions or rare-class coverage become impossible. This gives a finite, threshold-relative separation statement; it cannot guarantee every kind of statistical independence or prevent overfitting. Source anchors are train only. Molecular anchors are also checked against every default heldout using the same chemical and applicable geometric thresholds; conflicting parent groups are excluded. Global exposure to the models' previous corpora is a separate evaluation gate.

For a number-conserving one-particle Hamiltonian H, write $P_{ij} = \langle c_j^\dagger c_i \rangle$ and $U(t) = e^{-itH}$. The data use $P(t) = U(t)P(0)U(t)^*$, occupations $n_i = P_{ii}$ and outgoing currents $J_{ij} = -2\,\mathrm{Im}(H_{ij}P_{ji})$. Then $\dot{n}_i = -\sum_j J_{ij}$, directly from $i\dot{P} = [H, P]$. Paired BdG evolution includes its independently computed condensate source; it is not falsely treated as electron-number-conserving normal transport. An exact interacting Hubbard dimer supplements Gaussian exchange correlations. Molecular nuclear dynamics uses MMFF94, with a separately labelled, uncalibrated effective π-orbital model for charge modulation.

Parquet configurations retain condition, target and evaluation roles, source versions/hashes,

units, failures and audit evidence. The local dataset explorer shows actual structures, currents, time evolution, graph wavelets and checked circuit compilations. These computations support candidate discovery and model validation; the independent arithmetic determinant/zero-location bridge remains open. No new whole-chapter Lean formalization is asserted.

17 Molecular structure, complex transport and learned chemistry

The complementary-data laboratory now separates deposited protein structure, actual finite quantum transport and learned conformation generation. Covalent bonds use explicit molecular graphs or qualified PDB templates, with depth ordering and visible atom/bond scales. The ribbon view reads deposited `HELIX`/`SHEET` records; short synthetic peptide backbones remain unassigned rather than acquiring invented secondary structure.

Five additional finite propagation examples compare matched SSH/QWZ topological and trivial controls and an SSH domain wall. Complex phase, probability, current, coherence and selected eigenstates are distinct views. Analytic hydrogen volumes illustrate nodes, quantum beats and a phase vortex; they are not molecular electronic-structure calculations. Eight further CPU runs train classical and two-qubit hybrid score/flow models on periodic MMFF butane/pentane torsion spaces. All checkpoints, losses, controls and positive noise endpoints are explicit. These examples neither establish quantum advantage nor close the independent arithmetic bridge for GRH.

18 Learning objectives with explicit analytic guarantees

A useful learned contribution must distinguish approximation of known L-functions from proof of their zero locations. We propose holomorphic coefficient/readout learning, certified contour losses, arithmetic trace and functional-equation residuals, Schatten-controlled operator refinement, certificate search for the original signed moment, and exact-algebra representation constraints. Each has a different acceptance test.

For an analytic target on $|s-c| \le r$, let $|F| \le M_r$ there and choose $R < \rho < r$. A degree-N holomorphic readout trained on $m > N$ equally spaced points of the radius-ρ circle has the explicit uniform error budget

$$\|p_k - F\|_{|s-c|\le R} \le E_{\mathrm{opt}}(k) + E_{\mathrm{oracle}} + E_{\mathrm{alias}} + E_{\mathrm{Taylor}}.$$

The following proofs give all four terms, a primitive-Dirichlet arithmetic bound for M_r, and an infinite-feature kernel limit with its actual rates. Thus known analytic families can be approximated to arbitrary requested accuracy with independently certified evaluation error. This does not prove GRH. Uniform approximation by an already-real-zero determinant class is a different, potentially GRH-equivalent requirement; it must not be assumed through an alleged universal-approximation theorem.

19 Learning nonlocal strategies and checking their certificates

A two-player nonlocal game samples questions x, y and scores outputs a, b. The players may share a state, but communicate neither questions nor outcomes during play. Our finite tensor strategies have $p(a,b|x,y) = \mathrm{Tr}[\rho(P_a^x \otimes Q_b^y)]$. Offline joint optimization is allowed and is distinct from communication during the game. We train local measurement frames and a Schmidt angle using the exact Born objective, and replay actual checkpoints, probabilities, correlations, word moments and simulated shots in the interactive laboratory.

For $0 \le \alpha \le 2$ use the tilted score

$$S_\alpha = \alpha\langle A_0\rangle + \langle A_0B_0 + A_0B_1 + A_1B_0 - A_1B_1\rangle.$$

Its classical maximum is $2 + \alpha$; its quantum maximum is $\sqrt{8 + 2\alpha^2}$ [113]. At $\alpha = 0$ the uniform CHSH winning probability is $1/2 + S_0/8$. This conversion does not apply unchanged to the tilted objective. The three seeds for each of $\alpha = 0, 1, 3/2$ give nine trained runs. One $\alpha = 3/2$ run remains at the classical value 3.5; the other runs reach the stated quantum maxima numerically. This retained failure rules out a claim of universal convergence of the optimizer.

The magic-square task uses two Bell pairs, local four-dimensional joint measurements, row parity $+1$ and column parities $(+1, +1, -1)$. Its classical value is $8/9$ and its quantum value is 1, as proved below. Three further trained local-frame runs reach values within $3 \cdot 10^{-13}$ of one. All twelve runs are small CPU simulations with saved seeds and parameters, not hardware Bell experiments or evidence for quantum training advantage.

These examples belong to both the closure of finite tensor correlations C_{qa} and the commuting-operator set C_{qc}. The external theorem MIP* = RE separates these sets and implies the negative answer to Connes embedding [115]. We do not reproduce that theorem. Finite learned strategies, positive state Gram matrices and finite NPA constraints do not witness its separation.

20 Hubs, spokes and the topology retained by a dessin

The hubs-and-spokes construction in the HoTT book, Sections 6.6–6.8 [109], gives a useful language for attaching cells. For a loop $f : S^1 \to X$, form the homotopy pushout $X \cup_f CS^1$: add a fresh hub and paths from every $f(u)$ to that hub, continuously indexed by $u : S^1$. This attaches a disk, rather than merely a finite graph of radial edges. The book's Remark 6.7.1 explains why identifying the hub with an existing boundary point requires extra coherence; Remark 6.7.2 distinguishes propositional from judgmental computation.

For our dessins the useful distinction is geometric. A vertex star has a disk neighborhood with a specified cyclic ordering of its incident bands. Removing the disk center gives a punctured disk, which retains its peripheral loop. Removing only the vertex of a finite star instead leaves disconnected open rays. These operations cannot be interchanged. In a cellular dessin, filling a face imposes its attaching-word relation; leaving its center punctured retains a boundary component of the ribbon neighborhood.



disk
coherent circle of spokes

punctured disk
peripheral loop

The shaded regions are two-dimensional; the displayed finite spokes sample them. The annulus shown on the right is a deformation retract of the punctured disk, not a missing vertex in a one-dimensional graph.

Proposition 20.1 (Face filling changes the covering problem)**.** *Let a connected cellular dessin be a spine of an oriented genus-g ribbon surface with $n > 0$ boundary components. Its graph fundamental*

group is free of rank $2g + n - 1$. *Capping all boundary components gives the presentation*

$$\left\langle a_1, b_1, \ldots, a_g, b_g \;\middle|\; \prod_{i=1}^{g} [a_i, b_i] = 1 \right\rangle.$$

For the one-face clean dessins $K_{2,3}$ *and* $K_{2,5}$ *used here, these free ranks are respectively* 2 *and* 4, *before capping.*

Proof. A spanning-tree collapse of the connected spine leaves $E - V + 1$ circles. The ribbon surface retracts onto that spine and has Euler characteristic $2 - 2g - n = V - E$, proving the rank formula. Cutting a surface along a standard system of handle arcs gives the boundary word $\prod_i [a_i, b_i] \prod_j c_j$. Van Kampen therefore presents its group by these generators and this one relation. With a boundary present, eliminate c_n, leaving the stated free group. Capping attaches disks along the c_j, so van Kampen sets each $c_j = 1$, giving the closed-surface presentation. For our rotations on $K_{2,m}$, both black vertices have the same cyclic order on the m incident edges. Suppressing the bivalent white vertices gives dart rotations $(0 \cdots m-1)(m \cdots 2m-1)$ and involution $i \leftrightarrow i + m$. The face permutation is one $2m$-cycle for odd m. Thus $V = m + 2$, $E = 2m$, $F = 1$, and $g = (m-1)/2$, giving both examples. □

The once-punctured torus and closed torus both have $H_1 \cong \mathbb{Z}^2$, but their groups are F_2 and $\mathbb{Z}^2$; only the closed torus has $H_2 \cong \mathbb{Z}$. The analogous genus-two groups are F_4 and the four-generator surface group, with $H_1 \cong \mathbb{Z}^4$ in both cases. Indeed the standard closed-surface cellular complex has one vertex, $2g$ edges and one face. The face word is a product of commutators, so its abelianized boundary is zero. Removing the face leaves the wedge of circles. This computes the stated homology and also explains why H_1 alone misses the face relation. The torus group is abelian while F_2 is not; in higher genus the closed surface has nonzero H_2 while the graph does not.

Consequently the graph's universal cover is a tree, whereas the universal cover of the filled genus-two surface includes lifted faces and is a disk. Collapsing a spanning tree changes the combinatorial cover graph; the Cayley tree of a free basis is not literally the original biregular dessin cover. This distinction matters when lifting strings, periodic paths and boundary conditions in the infinite locally gentle algebra.

The gluing language organizes local patches and their compatibility maps. It does not reconstruct the ribbon cyclic order, complex structure, Belyi map, quadratic zero relations, coefficient ring or polynomial/formal/operator completion from the homotopy type alone. These data are retained explicitly in the existing dessin and order constructions. Descent of modules, derived categories, Hom-control, AR meshes or Fukaya categories needs a separately defined functor and hypotheses; a homotopy pushout of underlying spaces is not such a theorem. Nor does it imply an arithmetic determinant identity.

The accompanying tool computes the same one-face genus-one/two/four dart families and draws local disk/annulus patches and their surface polygon presentations, peripheral loops and face words. These are exact finite combinatorial checks plus the topological proof above. HoTT is referenced as a conceptual foundation; no univalence or higher inductive-type axiom is added to the pinned Lean4 development. Its complete cell-attachment and categorical formalization remains an explicit task.

21 A finite, validated approximation result

Let $\chi_3 = (0, 1, -1)$, $\chi_4 = (0, 1, 0, -1)$ and $\chi_5 = (0, 1, -1, -1, 1)$, extended periodically, with parities $a = 1, 1, 0$. We approximate the completed functions

$$\Lambda_q(s) = (q/\pi)^{(s+a)/2} \Gamma((s+a)/2) L(s, \chi_q).$$

The primitive theta/Mellin identity is the explicit classical analytic input; no hypothesis about zeros is used. These three examples do not assert a uniform result for all conductors, general Hecke functions or Artin holomorphy.

Theorem 21.1 (Scope of the supplied finite certificate)**.** *For each $q = 3, 4, 5$, the supplied exact-rational calculation constructs a complex polynomial P_q of degree 28 satisfying*

$$\sup_{|s-1/2|\le 7/10} |P_q(s) - \Lambda_q(s)| < 5 \cdot 10^{-15}.$$

The computation includes enclosures of all 64 true Fourier-circle nodes, completed-function sample error, analytic truncation, Fourier aliasing, the exact closed-form 180-step gradient-descent iterate and coefficient midpoint rounding. The Python implementation and informal analytic proof have independent checks; this is not a kernel-checked Lean theorem.

Proof. Write $F(c+z) = \sum_{n\ge 0} a_n z^n$, with $c = 1/2$, $R = 7/10$, $\rho = 6/5$, $r = 12/5$. The independently recomputed rational theta majorants are strictly below $M_q = 4, 10, 6$, respectively, on $|z| \le r$. Use true roots $\zeta_j = e^{2\pi i j/64}$ and rational labels y_j with proved error $|y_j - F(c + \rho\zeta_j)| \le \eta$. Use the real Euclidean gradient on complex coefficient coordinates for

$$\mathcal{L}(b) = \frac{1}{128} \sum_{j=0}^{63} \left| \sum_{n=0}^{28} b_n \zeta_j^n - y_j \right|^2.$$

Discrete Fourier orthogonality gives the least-squares optimum $b_n^* = 64^{-1} \sum_j y_j \bar{\zeta}_j^n$, $0 \le n \le 28$, and gradient $\nabla\mathcal{L}(b) = b - b^*$. For step $1/4$ and zero initialization, the exact 180-step iterate is $b_n^{180} = (1 - (3/4)^{180}) b_n^*$. The code evaluates this closed form; it does not claim an explicit180-update optimizer run.

Set $u = R/\rho = 7/12$, $v = R/r = 7/24$ and $w = (\rho/r)^{64} = 2^{-64}$. Let $\widehat{b}_n$ be an outward rectangle enclosing b_n^* and ϵ_n the sum of the coordinate halfwidths of the rectangle for b_n^{180}. Choose each actual polynomial coefficient as that rectangle's exact rational midpoint c_n. The normalized basis is explicit:

$$P_q(s) = \sum_{n=0}^{28} c_n \left(\frac{s - 1/2}{6/5} \right)^n.$$

Thus the exported coefficient with index n multiplies the normalized variable, rather than the raw power $(s - 1/2)^n$. Its error on the compact disk is bounded by

$$E_{\text{opt}} = (3/4)^{180} \sum_{n=0}^{28} |\widehat{b}_n|_1 u^n, \qquad E_{\text{arithmetic}} = \sum_{n=0}^{28} \epsilon_n u^n,$$

$$E_{\text{oracle}} = \eta \frac{1 - u^{29}}{1 - u}, \qquad E_{\text{alias}} = M_q \frac{w}{1 - w} \frac{1 - v^{29}}{1 - v},$$

$$E_{\text{Taylor}} = M_q \frac{v^{29}}{1 - v}.$$

Here $|\widehat{b}_n|_1$ is the sum of the maximum absolute real and imaginary coordinates, a safe upper bound for every enclosed complex modulus. Cauchy's estimate $|a_n| \le M_q r^{-n}$ and the Fourier identity $b_n^* = a_n \rho^n + \sum_{\ell\ge 1} a_{n+64\ell} \rho^{n+64\ell}$ for exact labels prove the alias and Taylor bounds. Label perturbations contribute at most η per Fourier coefficient. The remaining two terms follow directly from the exact iterate and coefficient rectangles. Each term is rounded upward to a 180-bit dyadic rational; the reported total is the exact sum of those reported terms. The certified totals below are strictly below the asserted threshold. Detailed sample enclosures are proved in the extended edition and encoded with exact integer endpoints. □

Conductor	Maximum sample radius	Uniform polynomial bound
3	3.026×10^{-17}	1.786×10^{-15}
4	7.206×10^{-17}	4.455×10^{-15}
5	1.404×10^{-16}	2.906×10^{-15}

The displayed values are rounded upward; exact rational records establish the strict inequalities.

22 What this does and does not certify

The final JSON records exact rational centers, interval endpoints, source hashes and all five error terms. Float values are display summaries only. A high-precision comparison is an independent diagnostic, not the reason an interval contains its mathematical target. Arithmetic uses outward integer operations, and invalid float/order inputs are rejected, including through typed caches. Earlier pilots and failed validation probes are retained.

This is a fixed finite tolerance/configuration. Increasing accuracy needs new truncation and precision choices; the fixed $N = 11, T = 128$ interface is not an arbitrary-accuracy oracle. The broader approximation existence theorem is separate. The certificate says nothing about the location of zeros: P_q is not required to be real-rooted or a self-adjoint spectral determinant. The independent arithmetic determinant identity, positivity and original signed short-row/covariance estimates remain open. No GRH proof or complete new Lean4 formalization is claimed.

23 An arithmetic-family moment with every original row retained

Let $K = \mathbb{Q}(\sqrt{-3})$, $\mathcal{O} = \mathbb{Z}[\omega]$, and write N for the ideal or element norm. Fix a finite-order Hecke character ν and a finite set S of prime ideals containing those above 6 and those dividing its conductor. For $0 \neq u \in \mathcal{O}$, let κ_u be the character of $K(u^{1/6})/K$ defined using arithmetic Frobenius, and put $\psi_u = \mathrm{prim}(\nu\kappa_u)$. Here prim means the underlying primitive Hecke character: it is formed before introducing any divisor variable or evaluating conductor-dependent zero extensions. The rows below are all actual Eisenstein elements, including units and prime powers supported in S.

The external inputs are abelian Kummer reciprocity and character uniqueness, the primitive Hecke functional equation and its usual vertical growth, Stirling's formula, scalar Dedekind-zeta convexity, and Blomer–Goldmakher–Louvel's large sieve for their specified sixth-order family. We use the pinned arXiv:1112.1650v1, Theorem 1.3, Lemma 2.1 and Appendix II; its Section 5 supplies the all-ideal decomposition whose fixed-twist proof is redone here. No statement about all primitive characters of comparable conductor, Artin holomorphy, Lindelöf bounds or zero locations is assumed.

Theorem 23.1 (The original-family numerator moment)**.** *For every $\varepsilon > 0$, uniformly for $H \geq 1$ and $t \in \mathbb{R}$,*

$$\sum_{0<Nu\leq H} |L(1/2 + it, \psi_u)|^2 \ll_{\nu,S,\varepsilon} H^{1+\varepsilon}(1 + |t|)^{1+\varepsilon}. \tag{8}$$

For every fixed $0 < \delta < 1/2$, the same estimate holds with $1/2 + it$ replaced by $1/2 + \delta + it$, with the constant also depending on δ. If the central bound has constant C_0 at an exponent $0 < \varepsilon_0 \leq 1/4$, an admissible off-central constant is

$$200C_0 \left(\frac{3/2 + \delta}{1/2 - \delta}\right)^2. \tag{9}$$

Proof. For the canonical BGL character $\vartheta_{\mathfrak{a}}$, Appendix II gives a radicand $\alpha_{\mathfrak{a}}$ with $(\alpha_{\mathfrak{a}})_S = \mathfrak{a}\mathfrak{g}^{-6}$. Since $\mathcal{O}_S$ is principal, comparison with any literal u whose outside-S ideal is $\mathfrak{a}$ gives $\kappa_u = \eta\vartheta_{\mathfrak{a}}$ for one

of finitely many fixed S-unit Kummer twists η. This is an equality of primitive characters, established outside a finite set and extended by uniqueness. For each outside-S ideal $\mathfrak{a}$ and S-supported integral ideal $\mathfrak{b}$, at most six elements generate $\mathfrak{ab}$. The resulting S-supported multiplicity sum converges at exponent $1+\varepsilon$.

For each fixed $\xi = \nu\eta$, split the canonical parameter as $\mathfrak{a} = \mathfrak{a}_0\mathfrak{a}_1\mathfrak{a}_2^6$, where $\mathfrak{a}_0$ is squarefree, $\mathfrak{a}_1$ is squarefull and sixth-power-free, and $(\mathfrak{a}_0, \mathfrak{a}_1) = 1$. The primitive conductor is at most a fixed multiple of $N(\mathfrak{a}_0\mathfrak{a}_1)$. The common-gamma approximate functional equation, with $X = H/(N\mathfrak{a}_2)^6$ and $T = 1 + |t|$, has common maximal length $B = X^{1/2}T(HT)^\eta$ for any sufficiently small fixed $\eta > 0$. Mellin inversion removes its row-dependent conductor weight. Split a column as $\mathfrak{b}_S\mathfrak{b}_1\mathfrak{b}_2$, with $\mathfrak{b}_S$ supported in S, $\mathfrak{b}_1$ squarefree and $\mathfrak{b}_2$ squarefull, the last two coprime and prime to S. The original primitive values at $\mathfrak{b}_S$ remain bounded row multipliers. The S-supported and squarefull column weights are summable at $1/2+\eta$. Multiplying squarefree column coefficients by the fixed ξ preserves their squared norm, whatever its finite order. BGL's large sieve therefore bounds the energy for fixed $\mathfrak{a}_1, \mathfrak{a}_2$ by

$$(HT)^{O(\eta)}\{M + B + (MB)^{2/3}\}, \qquad M = X/N\mathfrak{a}_1.$$

Squarefull counting and summation give $X + B\sqrt{X} + (XB)^{2/3} \ll (HT)^{O(\eta)}XT$. The $\mathfrak{a}_2$ sum converges at exponent 6. Principal rows are separated before the pole-free functional equation and bounded by scalar zeta convexity. This proves (8).

For the off-central statement divide each $L(s,\psi)$ by $(s+1)^{5/4}$ and, on principal rows, also multiply by $(s-1)/(s+1)$. These are bounded analytic functions on $\operatorname{Re} s > 1/2$, continuous on the boundary. Poisson representation and Jensen's inequality transfer the central squared moment. The convolution of $P_\delta(y) = \delta/[\pi(\delta^2+y^2)]$ with $(1+|t|)^{-3/2+\varepsilon_0}$ is at most eleven times that weight when $0 < \varepsilon_0 \le 1/4$. Restoring the denominator and the principal pole-removal factor gives (9). The extended edition gives the detailed derivation with all primitive local values. □

Separate fixed-unitary extension. The moment theorem also holds for any one fixed unitary Hecke character ν. For precision choose $\nu(\mathfrak{a}) = \nu_0(\mathfrak{a})(N\mathfrak{a})^{-i\tau}$, with angular parameter $m \in \mathbb{Z}$ and completed gamma factor

$$\gamma_\nu(s) = 2(2\pi)^{-(s+|m|/2+i\tau)}\Gamma(s + |m|/2 + i\tau).$$

All finite Kummer twists have this same infinity type. Use $T_\nu = 1 + |t+\tau| + |m|$, comparable to $1+|t|$ with ν-dependent constants. Pure norm rows have $L(s,\psi) = \zeta_K(s+i\tau)$ and are separated before the pole-free functional equation; their pole at $1-i\tau$ is removed by $(s-1+i\tau)/(s+1+i\tau)$. No uniformity in m, τ or the conductor of ν is asserted. This extension of the moment is separate from the following finite-order type-I residue formula.

23.1 The exact masked remainder and the remaining loss

Retain the original zero-extended symbol $\chi_d(u) = (u/d)_6$ on good squarefree ideals. Let μ be the ideal Möbius function and set

$$\beta_U(d) = \sum_{\substack{ab=d \\ Na, Nb \le U}} \mu(a)\mu(b), \quad T(u,d) = S \cup \operatorname{supp}(u) \cup \operatorname{supp}(d).$$

Let $\psi_{u,2} = \mathrm{prim}(\psi_u^2)$, formed independently of squaring any zero sentinel. With $\mathfrak{f}_u$ and $\mathfrak{f}_{u,2}$ the primitive conductors, the full-mask series is

$$F_{u,d}(s) = \frac{L(s,\psi_u)}{L(2s,\psi_{u,2})} C_{T(u,d)}(s), \tag{10}$$
$$C_T(s) = \frac{\prod_{p\in T,\ p\nmid \mathfrak{f}_u}(1-\psi_u(p)(Np)^{-s})}{\prod_{p\in T,\ p\nmid \mathfrak{f}_{u,2}}(1-\psi_{u,2}(p)(Np)^{-2s})}.$$

Initially it equals $\sum_{c\ \text{squarefree},\ (c,T)=1} \psi_u(c)(Nc)^{-s}$ for $\operatorname{Re} s > 1$. Every prime of T remains deleted, including those forgotten by either primitive conductor.

For finite-order ν, $D \geq 1$, $U \geq 1$, $U^2 \leq D$, and $W \in C_c^\infty((0,\infty))$ supported in $[1,2]$, put

$$\widehat{W}(s) = \int_0^\infty W(x) x^{s-1}\, dx, \qquad \mathcal{M}_4(W) = \sum_{j=0}^{4} \sup_x |(x\partial_x)^j W(x)|.$$

On $\sigma = 1/2 + \delta$, the exact residue-subtracted type-I remainder is

$$E_u = \frac{D^\delta}{2\pi} \int_{\mathbb{R}} \widehat{W}(\sigma+it) D^{it} \sum_{\substack{d\ \text{squarefree}\\ (d,S)=1}} \frac{\beta_U(d)\nu(d)\chi_d(u)}{(Nd)^{\sigma+it}} F_{u,d}(\sigma+it)\, dt. \tag{11}$$

This definition retains the actual β_U, all phases, d-masks, conductors and the independently primitive square.

Theorem 23.2 (An improved masked residual estimate)**.** *Under the preceding finite-order hypotheses, for every $\varepsilon > 0$,*

$$\sum_{0<Nu\leq H} |E_u|^2 \ll_{\nu,S,\delta,\varepsilon} \mathcal{M}_4(W)^2 D^{2\delta} U^{2-4\delta} H (HD)^\varepsilon. \tag{12}$$

Proof. The same deleted set in both factors of (10) gives $|F_{u,d}(\sigma+it)| \ll (Nu\,Nd)^\eta |L(\sigma+it,\psi_u)|$: the reciprocal incomplete Euler product at $2\sigma > 1$ is bounded by $\zeta_K(2\sigma)$, and its remaining finite numerator is bounded by a small power of its radical. Apply Minkowski to (11) in the exact row space and use Theorem 23.1. The actual coefficient sum is at most

$$\sum_d \frac{|\beta_U(d)|}{(Nd)^{\sigma-\eta}} \leq \left(\sum_{Na\leq U} (Na)^{-\sigma+\eta} \right)^2 \ll U^{1-2\delta+2\eta}.$$

Four Mellin integrations by parts bound the remaining height integral by a constant times $\mathcal{M}_4(W)$. Square the result and absorb the small losses using $U^2 \leq D$. □

At $H = D^{3/4}$ and $U = D^a$ with fixed $0 < a \leq 1/2$, the upper bound's excess over H is $2\delta + (2-4\delta)a$, up to small losses; as $\delta \downarrow 0$ it still leaves $2a$. This is a limitation of the upper bound, not an energy lower bound. The extracted principal residue, the original short term at smaller scales and the signed type-II minus type-I covariance are not controlled by (12). The next arithmetic task is to estimate the complete d-weighted moment before its absolute coefficient sum, or establish sufficient signed covariance. No positive arithmetic determinant, zero-location theorem or GRH proof follows. The analytic argument uses the external classical inputs above; finite diagnostics and selected existing Lean local identities do not formalize the Hecke functions, approximate functional equation, Poisson transfer or these moment bounds.

24 Recombining the actual weighted cutoff

Keep the finite-order Eisenstein family and original masks of Section 23. For $U \geq 1$ and a squarefree ideal m, put

$$A_U(m) = \sum_{d|m} \beta_U(d) = \sum_{\substack{ab|m \\ (a,b)=1,\ Na,Nb\leq U}} \mu(a)\mu(b).$$

Every summation variable here is an integral ideal. The last formula uses squarefreeness of m; an unconstrained square of a Möbius polynomial would also include forbidden overlapping supports.

Theorem 24.1 (Exact recombination and an absolute-half-plane rate)**.** *Write $T_0 = S \cup \mathrm{supp}(u)$ and $J_{u,U}(s) = \sum_{d \text{ squarefree},\ (d,S)=1} \beta_U(d)\nu(d)\chi_d(u)(Nd)^{-s}F_{u,d}(s)$. For $\mathrm{Re}\, s > 1$,*

$$J_{u,U}(s) = \sum_{m \text{ squarefree},\ (m,T_0)=1} A_U(m)\psi_u(m)(Nm)^{-s},$$
$$|A_U(m)| \leq 3^{\omega(m)}, \qquad A_U(m) = \mu(m) \quad (Nm \leq U).$$

For every $\kappa > 0$ and $0 < \epsilon < \kappa$, uniformly for $\mathrm{Re}\, s \geq 1 + \kappa$, all original rows and all $U \geq 1$,

$$\left| J_{u,U}(s) - \frac{1}{L^{T_0}(s,\psi_u)} \right| \leq U^{-\kappa+\epsilon}\{\zeta_K(1+\epsilon)^3 + \zeta_K(1+\epsilon)\}.$$

Here L^{T_0} removes every prime of T_0, including conductor-drop primes; the bound is in the absolute-convergence half-plane only.

Proof. On a surviving term $m = dc$, the two squarefree ideals c, d are coprime, and both avoid T_0. The original product $\nu(d)\chi_d(u)\psi_u(c)$ is then $\psi_u(m)$. Grouping by m gives A_U. Each prime of m has three possible positions in $(a, b, m/(ab))$, so the absolute sum is at most $3^{\omega(m)}$. When $Nm \leq U$ all divisors fit both cutoffs; the primewise sum is $1 - 1 - 1 = -1$, giving $\mu(m)$.

The same bound justifies absolute convergence and regrouping for $\mathrm{Re}\, s > 1$, because on squarefree ideals $3^{\omega(m)} = \tau_3(m)$ and $\sum_m \tau_3(m)(Nm)^{-v} = \zeta_K(v)^3$ for $v > 1$, where this latter sum runs over all nonzero integral ideals. Subtract the absolutely convergent Möbius series for $1/L^{T_0}$. Only $Nm > U$ remain. Rankin's inequality with exponent $\eta = \kappa - \epsilon$ bounds their absolute sum by

$$U^{-\eta} \sum_m \{\tau_3(m) + 1\}(Nm)^{-1-\epsilon},$$

which proves the stated constant independently of deleted primes, row multiplicities or character phases. □

For W supported in $[1, 2]$, the same exact regrouping gives the original normalized type-I polynomial

$$I_u = D^{-1/2} \sum_{m \text{ squarefree},\ (m,S)=1} A_U(m)\nu(m)\chi_m(u)W(Nm/D).$$

This formula retains all original zero sentinels. It permits a direct family large sieve before any absolute d sum; its quantitative moment is developed separately below. The displayed absolute-half-plane rate alone is not that critical-strip moment.

25 A cutoff-uniform envelope for the original masked remainder

Fix $K = \mathbb{Q}(\omega)$, $\mathcal{O} = \mathbb{Z}[\omega]$, a finite-order Hecke character ν, and a finite set S containing the primes above 6 and the conductor of ν. All unspecified ideal sums below are over nonzero integral ideals. Let $\chi_m(u) = (u/m)_6$ be the original symbol, zero when $(m,u) > 1$, and put

$$\beta_U(d) = \sum_{\substack{ab=d \\ Na, Nb \le U}} \mu(a)\mu(b), \qquad A_U(m) = \sum_{d \mid m} \beta_U(d).$$

The columns d, c, m are squarefree and prime to S where indicated. Retain the primitive characters $\psi_u = \mathrm{prim}(\nu\kappa_u)$, $\psi_{u,2} = \mathrm{prim}(\psi_u^2)$, and the full deleted set $T(u,d) = S \cup \mathrm{supp}(u) \cup \mathrm{supp}(d)$. For $H, D, U \ge 1$, $U^2 \le D$, and $W \in C_c^\infty((0,\infty))$ supported in $[1,2]$, the original normalized type-I piece is

$$I_u = D^{-1/2} \sum_{\substack{m \text{ squarefree} \\ (m,S)=1}} A_U(m)\nu(m)\chi_m(u) W(Nm/D). \tag{13}$$

Its exact principal residue is

$$P_u = \mathbf{1}_{\psi_u=1} \sqrt{D}\, \widehat{W}(1) \sum_{\substack{d \text{ squarefree} \\ (d,S)=1}} \frac{\beta_U(d)\nu(d)\chi_d(u)}{Nd} \rho_{T(u,d)},$$

$$\rho_T = \frac{\mathrm{res}_{s=1}\, \zeta_K(s)}{\zeta_K(2)} \prod_{p \in T} (1 + (Np)^{-1})^{-1}. \tag{14}$$

Define $E_u = I_u - P_u$; this is the same residue-subtracted remainder as the fully masked Mellin contour, with no cutoff limit or profile supremum.

Theorem 25.1 (Uniformity in the actual truncated cutoff)**.** *For every $\varepsilon > 0$,*

$$\sum_{0 < Nu \le H} |E_u|^2 \ll_{\nu,S,\varepsilon} \|W\|_\infty^2 (HD)^\varepsilon \{H + DH^{1/3} + (HD)^{2/3}\}. \tag{15}$$

The sum is over every actual nonzero Eisenstein element, including units, S-supported powers and sixth-power rows. The constant is independent of U under $U^2 \le D$. This is an application of the classical sextic large sieve with exact original coefficients, not a zero-location theorem.

Proof. On an absolutely convergent line, expand the restored squarefree series $F_{u,d}$. Its condition $(c, T(u,d)) = 1$ includes $(c,d) = 1$, and the original symbols multiply. Grouping $m = cd$ gives (13) by Mellin inversion. Each prime of m may belong to the first truncated Möbius divisor, to the second, or to neither. Consequently $|A_U(m)| \le 3^{\omega(m)} \ll_\eta (Nm)^\eta$, uniformly in U. The length-D coefficient sequence in (13) therefore has squared norm $\ll_\eta \|W\|_\infty^2 D^\eta$.

The all-ideal row norm in Section 3 of Blomer–Goldmakher–Louvel, arXiv:1112.1650v1 gives

$$B_2^1(X, D) \ll_\eta (XD)^\eta \{X + DX^{1/3} + (XD)^{2/3}\}.$$

For completeness, split $a = a_1 a_2^2 \cdots a_5^5 a_6^6$ dyadically; $a_1, \dots, a_5$ are squarefree and pairwise coprime. If their dyadic sizes are A_j, put $R = \prod_{j=1}^6 A_j$ and $M = \max(A_1, A_2)$. Pivot on the first or second variable, using the squarefree powered sieve proved simultaneously in that paper's Section 3.2. The other variables are bounded column multipliers. The block energy is at most

$$(XD)^\eta \{R + DR/M + D^{2/3}R/M^{1/3}\} \sum_m |\lambda_m|^2.$$

The inequalities $R \le X$, $R^3 \le XM^3$ and $R^3 \le X^2M$ follow from $\prod_j A_j^j \le X$ and the pivot choice. Summing the logarithmically many blocks proves the displayed bound.

Finite S-unit and ray-class partitions convert the original literal symbol into the canonical column value $\vartheta_m(a)$ with fixed modulus-one coefficient multipliers. On noncoprime pairs both values are zero: squarefree m has outside-S conductor exactly m. This orientation preserves sixth-power zeros. At most six elements generate each outside-S ideal times an S-supported ideal; the S-supported sums converge at powers 1, 1/3 and 2/3. Thus (15) holds first with I_u in place of E_u.

A principal ψ_u requires every outside-S valuation of u to be divisible by 6. There are $O_{\nu,S}(H^{1/6})$ such literal rows. Equation (14), $|\rho_T| \ll_K 1$, and the actual coefficient sum

$$\sum_d \frac{|\beta_U(d)|}{Nd} \le \left(\sum_{Na \le U} \frac{1}{Na} \right)^2 \ll \log^2(2U)$$

give $\sum |P_u|^2 \ll DH^{1/6}\|W\|_\infty^2 \log^4(2U)$. This is absorbed by the right side of (15), using $U^2 \le D$ and a smaller internal exponent. Finally $|E_u|^2 \le 2|I_u|^2 + 2|P_u|^2$. The extended proof specifies the canonical-family transfer and the powered-sieve input precisely. □

Combining this with the previously proved fixed-twist AFM transfer yields

$$\sum_{0<Nu\le H} |E_u|^2 \ll_{\nu,S,\delta,\varepsilon} \mathcal{M}_4(W)^2 (HD)^\varepsilon \min\{D^{2\delta}U^{2-4\delta}H,\ H + DH^{1/3} + (HD)^{2/3}\}. \tag{16}$$

At $H = D^{3/4}$ and $U = D^a$, $0 \le a \le 1/2$, the excess over H is at most $\min\{2\delta + (2-4\delta)a, 1/2\}$, up to arbitrarily small losses. The new envelope improves the old one for $a > 1/4$ as $\delta \downarrow 0$, but still leaves an exponent loss of 1/2. It does not settle the original signed short-row bound, type-II covariance, positive arithmetic determinant comparison or GRH. The principal-residue bound used here is sufficient for this envelope, not the stronger target needed to reassemble the full signed argument.

References

[1] A. Schreiber, *Surface Algebras I: Dessins D'enfants, Surface Algebras, and Dessin Orders*, arXiv:1810.06750v5 (2023).

[2] A. Schreiber, *Surface Algebras and Surface Orders II: Affine Bundles on Curves*, arXiv:1812.00621v1; record 2018, downloaded PDF internally dated November 27, 2024.

[3] A. Schreiber, *Multiparameter Persistent Homology—Generic Structures and Quantum Computing*, arXiv:2210.11433v1 (2022).

[4] C. Bessenrodt and T. Holm, *Weighted locally gentle quivers and Cartan matrices*, arXiv:math/0511610v1 (2005).

[5] S. Ford, A. Oswald and J.J. Zhang, *Homological conditions on locally gentle algebras*, arXiv:2409.08333v3 (5 September 2026).

[6] S. Opper, P.-G. Plamondon and S. Schroll, *A geometric model for the derived category of gentle algebras*, arXiv:1801.09659v7 (10 July 2025).

[7] W. Chang, *Geometric models for the algebraic hearts in the derived category of a gentle algebra*, arXiv:2305.03998v2 (4 September 2024).

[8] Y.-Z. Liu, C. Zhang and H. Zhang, *Constructing projective resolution and taking cohomology for gentle algebras in the geometric model*, arXiv:2308.07220v1 (2023).

[9] Z. Li, Y. Qiu and Y. Zhou, *A geometric realization of Koszul duality for graded gentle algebras*, arXiv:2403.15190v1 (2024).

[10] W. Chang, H. Jin and S. Schroll, *Recollements of partially wrapped Fukaya categories and surface cuts*, arXiv:2206.11196v2 (2023).

[11] B. Dequêne and M. Schoonheere, *Resolving subcategories for gentle algebras I: Monogeneous resolving subcategories for gentle trees*, arXiv:2502.20994v3 (2026).

[12] D.A. Buchsbaum and D. Eisenbud, *What makes a complex exact?*, J. Algebra 25 (1973), 259–268, primary paper.

[13] W. Bruns, *The existence of generic free resolutions and related objects*, Math. Scand. 55 (1984), 33–46.

[14] D. Eisenbud, *Homological algebra on a complete intersection, with an application to group representations*, Trans. AMS 260 (1980), 35–64.

[15] A. Beauville and Y. Laszlo, *Un lemme de descente*, C. R. Acad. Sci. Paris 320 (1995), 335–340, author text.

[16] P. Magyar, *Affine Schubert varieties and circular complexes*, arXiv:math/0210151v1 (2002).

[17] F.F. Knudsen and D. Mumford, *The projectivity of the moduli space of stable curves I: Preliminaries on 'det' and 'Div'*, Math. Scand. 39 (1976), 19–55, doi:10.7146/math.scand.a-11642.

[18] M. Breuning, *Determinant functors on triangulated categories*, arXiv:math/0610435v1 (2006).

[19] D. Eriksson and G. Freixas i Montplet, *Deligne–Riemann–Roch and intersection bundles*, J. Éc. polytech. Math. 11 (2024), 247–361, doi:10.5802/jep.254.

[20] P. Deligne, *La conjecture de Weil I*, Publ. Math. IHÉS 43 (1974), 273–307, primary text.

[21] P. Deligne, *La conjecture de Weil II*, Publ. Math. IHÉS 52 (1980), 137–252, primary text.

[22] P. Deligne, *Cohomologie étale (SGA $4\frac{1}{2}$)*, Lecture Notes in Mathematics 569, Springer-Verlag, 1977, doi:10.1007/BFb0091516.

[23] G. Tabuada, *Noncommutative Weil conjecture*, arXiv:1808.00950v7 (6 December 2019).

[24] C. Deninger, *Arithmetic geometry and analysis on foliated spaces*, arXiv:math/0505354v1 (2005).

[25] D.B. Ray and I.M. Singer, *R-Torsion and the Laplacian on Riemannian manifolds*, Advances in Math. 7 (1971), 145–210, doi:10.1016/0001-8708(71)90045-4.

[26] F. Bornemann, *On the numerical evaluation of Fredholm determinants*, arXiv:0804.2543v1; Math. Comp. 79 (2010), 871–915.

[27] B. Dequêne and M. Schoonheere, *Resolving subcategories for gentle algebras II: Resolving subcategories for gentle trees*, arXiv:2509.10120v2 (3 October 2025).

[28] C. Procesi, *The invariant theory of $n \times n$ matrices*, Advances in Math. 19 (1976), 306–381, doi:10.1016/0001-8708(76)90027-X.

[29] L. Le Bruyn and C. Procesi, *Semisimple representations of quivers*, Trans. Amer. Math. Soc. 317 (1990), 585–598, doi:10.1090/S0002-9947-1990-0958897-0.

[30] A. Craw and R. Yamagishi, *The Le Bruyn–Procesi theorem following Lusztig*, arXiv:2312.08527v4 (14 November 2025).

[31] P. Etingof, O. Golberg, S. Hensel, T. Liu, A. Schwendner, D. Vaintrob and E. Yudovina, *Introduction to Representation Theory*, with historical interludes by S. Gerovitch, Student Mathematical Library 59, American Mathematical Society, 2011, doi:10.1090/stml/059; author-hosted text.

[32] K. Akin, D.A. Buchsbaum and J. Weyman, *Schur functors and Schur complexes*, Advances in Math. 44 (1982), 207–278, doi:10.1016/0001-8708(82)90039-1.

[33] Q. Jiang, *Derived Grassmannians and derived Schur functors*, arXiv:2212.10488v3 (6 July 2023).

[34] A. Price, A. Stelzer and A. Yong, *Representations from matrix varieties, and filtered RSK*, arXiv:2403.09938v2 (6 October 2025).

[35] H. Derksen and J. Weyman, *Semi-invariants of quivers and saturation for Littlewood–Richardson coefficients*, J. Amer. Math. Soc. 13 (2000), 467–479, primary author text.

[36] A. Schofield and M. Van den Bergh, *Semi-invariants of quivers for arbitrary dimension vectors*, arXiv:math/9907174v1 (27 July 1999); Indag. Math. 12 (2001), 125–138.

[37] J.S. Milne, *Reductive Groups*, author's course notes, version 2.00 (10 March 2018), §5.18.

[38] W. Crawley-Boevey, *Classification of modules for infinite-dimensional string algebras*, Trans. Amer. Math. Soc. 370 (2018), 3289–3313, doi:10.1090/tran/7032; arXiv:1308.6410v2 (4 March 2016).

[39] M. Auslander and I. Reiten, *Almost split sequences for Cohen–Macaulay-modules*, Math. Ann. 277 (1987), 345–349, doi:10.1007/BF01457367.

[40] R. Bautista, S. Liu and C. Paquette, *Representation Theory of an Infinite Quiver*, arXiv:1109.3176v1 (14 September 2011); published as *Representation theory of strongly locally finite quivers*, Proc. London Math. Soc. 106 (2013), 97–162.

[41] C. Paquette, *Irreducible morphisms and locally finite dimensional representations*, arXiv:1508.04353v3 (4 January 2016).

[42] S. Liu, P. Ng and C. Paquette, *Almost split sequences and approximations*, arXiv:1203.5009v1 (19 March 2012).

[43] C.M. Ringel, *The Minimal Representation-Infinite Algebras which are Special Biserial*, author-hosted ICRA Tokyo 2010 proceedings text, 57 pages, retrieved 3 November 2026.

[44] G. Puninski and M. Prest, *Ringel's conjecture for domestic string algebras*, arXiv:1407.7470v2 (13 September 2015).

[45] R. Laking, M. Prest and G. Puninski, *Krull–Gabriel dimension of domestic string algebras*, arXiv:1506.07703v2 (29 June 2015).

[46] H.-L. Huang, Z. Lin and X. Su, *The Auslander–Reiten quivers of string algebras of affine type $\widetilde{C}$ and a conjecture by Geiss–Leclerc–Schröer*, arXiv:2110.02777v2 (16 November 2021).

[47] G. Abrams and G. Aranda Pino, *The Leavitt path algebra of a graph*, arXiv:math/0509494v1 (21 September 2005).

[48] T. Bates, D. Pask, I. Raeburn and W. Szymański, *The C^*-algebras of row-finite graphs*, New York J. Math. 6 (2000), 307–324, primary journal text.

[49] N.J. Fowler and I. Raeburn, *The Toeplitz algebra of a Hilbert bimodule*, arXiv:math/9806093v3 (23 June 1998).

[50] N. Brownlowe, M. Laca, D. Robertson and A. Sims, *Reconstructing directed graphs from generalised gauge actions on their Toeplitz algebras*, arXiv:1812.08903v1 (21 December 2018).

[51] E.T.A. Kakariadis and O.M. Shalit, *On operator algebras associated with monomial ideals in noncommuting variables*, arXiv:1501.06495v3 (21 November 2018).

[52] O.M. Shalit and B. Solel, *Subproduct systems*, arXiv:0901.1422v3 (26 May 2009).

[53] A. Connes and C. Consani, *Knots, primes and class field theory*, arXiv:2501.06560v1 (11 January 2025).

[54] A. Connes, C. Consani and H. Moscovici, *Zeta zeros and prolate wave operators*, arXiv:2310.18423v2 (4 May 2024).

[55] A. Connes, C. Consani and H. Moscovici, *On q-series and the moment problem associated to local factors*, arXiv:2403.01247v1 (2 March 2024).

[56] A. Connes, C. Consani and H. Moscovici, *Zeta Spectral Triples*, arXiv:2511.22755v1 (27 November 2025).

[57] A. Connes and W.D. van Suijlekom, *Quadratic Forms, Real Zeros and Echoes of the Spectral Action*, arXiv:2511.23257v1 (28 November 2025).

[58] A. Connes, *The Riemann Hypothesis: Past, Present and a Letter Through Time*, arXiv:2602.04022v1 (3 February 2026).

[59] A. Connes and C. Consani, *On the Jacobian of* $\overline{\mathrm{Spec}\,\mathbb{Z}}$, arXiv:2602.15941v1 (17 February 2026).

[60] C. Deninger, *Dynamical systems for arithmetic schemes*, arXiv:1807.06400v4 (7 February 2024); Indag. Math. 37 (2026), 25–136, doi:10.1016/j.indag.2024.05.007.

[61] C. Deninger, *Primes, knots and periodic orbits*, arXiv:2301.11643v1 (27 January 2023).

[62] C. Deninger, *There is no "Weil-"cohomology theory with real coefficients for arithmetic curves*, arXiv:2204.02714v2 (27 January 2023).

[63] R. Ponge and Y. Tian, *Nonclassical Weyl laws and Connes' Integration for weak Lorentz ideals, I*, arXiv:2605.25591v2 (28 September 2026).

[64] F.W.J. Olver et al. (eds.), *NIST Digital Library of Mathematical Functions*, finite summation by parts, Eq. 2.10.9 and Dirichlet L-functions, §25.15, consulted 3 November 2026.

[65] A. Chirvasitu, P.M. Hajac and M. Tobolski, *Non-surjective pullbacks of graph C*-algebras from non-injective pushouts of graphs*, arXiv:1907.10260v3 (31 December 2019), Theorems3.4 and4.3; doi:10.1112/blms.12389.

[66] Y. Lekili and A. Polishchuk, *Derived equivalences of gentle algebras via Fukaya categories*, arXiv:1801.06370v5 (27 August 2019), Definition3.1.1, Lemma3.1.3 and Theorem3.2.2.

[67] M. Tomforde, *Uniqueness theorems and ideal structure for Leavitt path algebras*, arXiv:math/0612628v6 (27 February 2007), Theorem7.3, pp.29–30.

[68] A. Weil, *Numbers of solutions of equations in finite fields*, Bull. Amer. Math. Soc.55 (1949),497–508, especially pp.499–502 and Eq.(4), doi:10.1090/S0002-9904-1949-09219-4; primary-paper scan.

[69] The Stacks Project, Tag0FJ9 and Tag0GSX (retrieved 3 November 2026).

[70] J. Alper, *Good moduli spaces for Artin stacks*, Ann. Inst. Fourier63 (2013),2349–2402, Theorem10.3, pp.2386–2387, doi:10.5802/aif.2833.

[71] The Stacks Project, Tag04V8: Taking roots of line bundles (retrieved 3 November 2026).

[72] G. Zwara, *Smooth morphisms of module schemes*, Proc. London Math. Soc. (3) 84 (2002), 539–558, primary publication.

[73] C. Riedtmann and G. Zwara, *Orbit closures and rank schemes*, Comment. Math. Helv. 88 (2013), 55–84, primary publication, Section6.

[74] Y. Liu, N. Li, B. Xing and P. Chen, *A lecture note on covering theory in representation theory of algebras*, arXiv:2508.01563v5, 28 May 2026.

[75] Q. Dong, Z. Lin, M. Lu and S. Ruan, *Representation-tame Geiß–Leclerc–Schröer algebras and a revised GLS conjecture on root systems*, arXiv:2609.26503v1, 22 September 2026.

[76] S. Schroll, *On geometric models in representation theory*, Proceedings of ICM 2026, vol.3, pp.250–271, doi:10.1137/25M1810647; author preprint arXiv:2601.14396v1.

[77] B. Keller, *Introduction to A_∞ algebras and modules*, arXiv:math/9910179v2 (2001), especially §§3.1,3.3,6.2.

[78] The Stacks Project Authors, *The Koszul complex*, Tag 0621, especially Definitions15.29.1–2 and Lemma15.29.5; snapshot November3,2026, hash recorded in the repository.

[79] The Stacks Project Authors, *Regular sequences are Koszul regular*, Tag 062F, Lemma15.31.2; snapshot November3,2026, hash recorded in the repository.

[80] S. Datta, S. Howard and D. Cochran, *Geometry of the Welch Bounds*, arXiv:0909.0206v2 (2012), Section2.1, Theorem2.2.

[81] K.G. Wilson and J. Kogut, *The renormalization group and the epsilon expansion*, Physics Reports12(1974),75–199, DOI10.1016/0370-1573(74)90023-4; pinned institutional scan and review scope in `planning/ACTUAL_RG_SOURCES.json`.

[82] M. Levin and C.P. Nave, *Tensor renormalization group approach to2Dclassical lattice models*, arXiv:cond-mat/0611687v2 (2008); Phys. Rev. Lett.99(2007),120601.

[83] J. Kim et al., *Pure Transformers are Powerful Graph Learners*, arXiv:2207.02505v2 (2022). Our finite measured-feature prototype does not reproduce its full benchmarks.

[84] S. Liu et al., *Unsupervised Discovery of Steerable Factors When Graph Deep Generative Models Are Entangled*, arXiv:2401.17123v1 (2024); scoped primary reading in the computational source manifest.

[85] S. Lahlou et al., *A theory of continuous generative flow networks*, arXiv:2301.12594v2 (2023); state measures and trajectory balance.

[86] M. Gross and B. Siebert, *Mirror Symmetry via Logarithmic Degeneration Data II*, arXiv:0709.2290v2; used as a geometric research context, not an implemented mirror comparison.

[87] B. Green and T. Tao, *Quadratic uniformity of the Möbius function*, arXiv:math/0606087v2, Section4, Lemma4.1, printed pages12–13. Focused identity reading only; no nilsequence estimate is asserted for the moving sextic family here.

[88] S.S. Kudla, *Tate's thesis*, expanded Jerusalem lecture notes (2001), Theorem4.3 and Corollary4.4, pinned primary notes.

[89] J.S. Milne, *Class Field Theory*, version4.03 (6August2020), Theorem0.8 and ChapterI,§1, author notes.

[90] A. Jacot, F. Gabriel and C. Hongler, *Neural Tangent Kernel: Convergence and Generalization in Neural Networks*, arXiv:1806.07572v4; specified classical scaling hypotheses.

[91] J. Liu et al., *Representation Learning via Quantum Neural Tangent Kernels*, arXiv:2111.04225v3; focused version/abstract review.

[92] M. Kontsevich and Y. Soibelman, *Affine structures and non-archimedean analytic spaces*, arXiv:math/0406564v1; valued and integral-affine data.

[93] Y. Lekili and A. Polishchuk, *Derived equivalences of gentle algebras via Fukaya categories*, arXiv:1801.06370v5; category and smoothness hypotheses retained.

[94] Z. Cui, P. Zhang and Y. Tang, *Quantum Flow Matching*, arXiv:2508.12413v1; abstract-level comparison, not an upstream reproduction.

[95] L. Abrams, *Two-dimensional topological quantum field theories and Frobenius algebras*, J. Knot Theory Ramifications 5 (1996), 569–587, author-hosted paper.

[96] G. Segal, *The definition of conformal field theory*, in *Topology, Geometry and Quantum Field Theory* (2004); primary publisher metadata and sewing-scope comparison.

[97] S. Ryu and T. Takayanagi, *Holographic Derivation of Entanglement Entropy from AdS/CFT*, arXiv:hep-th/0603001v2; external setting only.

[98] S. Mochizuki, *The Absolute Anabelian Geometry of Hyperbolic Curves*, author-hosted paper, introduction and Theorem 1.3.4; arithmetic extension hypotheses retained.

[99] Z. Ji, A. Natarajan, T. Vidick, J. Wright and H. Yuen, *MIP*=RE*, arXiv:2001.04383v4 (16 October 2026); external theorem and version record, not rerun locally.

[100] T. Fukui, Y. Hatsugai and H. Suzuki, *Chern Numbers in Discretized Brillouin Zone*, arXiv:cond-mat/0503172v2, 2005.

[101] RDKit contributors, *MMFF94 force fields and ETKDG conformer APIs*, `https://www.rdkit.org/docs/`; executed version2026.03.1.

[102] Worldwide Protein Data Bank, *Usage Policies*, CC0-1.0 coordinate archive, `https://www.wwpdb.org/about/usage-policies`.

[103] F. Evers et al., *Theory of Chirality Induced Spin Selectivity: Progress and Challenges*, arXiv:2108.09998v1, 2021.

[104] A. Privitera et al., *Direct detection of spin polarization in photoinduced charge transfer through a chiral bridge*, arXiv:2205.05353v1, 2022.

[105] wwPDB, Atomic Coordinate Entry Format3.3, secondary structure and coordinate sections, accessed2026-11-04. `https://www.wwpdb.org/documentation/file-format-content/format33/sect5.html`.

[106] Y. Song et al., Score-Based Generative Modeling through Stochastic Differential Equations, arXiv:2011.13456v1 (2020).

[107] Y. Lipman et al., Flow Matching for Generative Modeling, arXiv:2210.02747v1 (2022).

[108] R. T. Q. Chen and Y. Lipman, Flow Matching on General Geometries, arXiv:2302.03660v1 (2023).

[109] The Univalent Foundations Program, *Homotopy Type Theory: Univalent Foundations of Mathematics*, first-edition-257-g5561b73 (2013), Sections 6.6–6.8, especially Remarks 6.7.1–6.7.2, pinned official edition.

[110] A. Jacot, F. Gabriel and C. Hongler, *Neural Tangent Kernel: Convergence and Generalization in Neural Networks*, arXiv:1806.07572v4 (2020). This is background; our diagonal holomorphic readout limit is derived separately.

[111] NIST Digital Library of Mathematical Functions, *Dirichlet L-functions*, Section 25.15, equations 25.15.1–25.15.5, primary reference, accessed 4 November 2026.

[112] F. Bornemann, *On the Numerical Evaluation of Fredholm Determinants*, arXiv:0804.2543v2 (2008), Section 4, equation (4.1); Math. Comp. 79 (2010), 871–915.

[113] C. Bamps and S. Pironio, *Sum-of-squares decompositions for a family of CHSH-like inequalities and their application to self-testing*, arXiv:1504.06960v1 (2015).

[114] M. Navascués, S. Pironio and A. Acín, *A convergent hierarchy of semidefinite programs characterizing the set of quantum correlations*, arXiv:0803.4290v1 (2008).

[115] Z. Ji, A. Natarajan, T. Vidick, J. Wright and H. Yuen, *MIP*=RE*, arXiv:2001.04383v4 (16 October 2026). External theorem, not re-proved or newly formalized here.