Auditing Surface-Order Approaches to GRH: Detailed Proofs and Research Interfaces

GRH Research Project
Working manuscript; authorship to be confirmed

November 2, 2026

Research draft. GRH is not proved here.

Abstract

We audit proposed surface-order and Hilbert–Pólya arguments for GRH. We prove obstructions to the proposed prime/geodesic identification, integral cyclic induction, absolute prime-tail estimates and determinant preservation, and retain a conditional spectral closure criterion. We construct convergent arithmetic approximants with explicit errors and analyze coefficient-defined Jensen metrics. From the explicit scalar local factors of a separate exponent argument, we extend an Euler-correction region and its principal-residue interface, and isolate a shifted infinite-order angular Hecke factor. Under the stated classical analytic inputs, its zeros obstruct extending the unchanged correction's nonvanishing criterion below 7/12. We also reconstruct a frozen-capacity row count conditionally on the source moment and witness estimates, and extend the low interface under explicit reflected-energy and Gram hypotheses. The nonprincipal central contours also extend under their stated arithmetic and buffered-bound contracts. We verify the finite-field Gauss and zero-preserving correlation input, with prime-power and reciprocity dependencies distinguished. We also prove an annular periodic cancellation lemma and reconstruct the Gram summation under explicit arithmetic representation and period inputs. A classical conductor argument supplies an explicit common arithmetic period, distinguished from the smaller quadratic-phase modulus. We verify the local reflected coefficient separation and its common-profile normalization, retaining the global theta and hybrid-sieve dependencies. A pinned Lean 4 companion verifies selected arithmetic, complex-analytic, matrix and infinite-product components, with its remaining dependencies specified. These results do not establish GRH; the full arithmetic spectral bridge and the required mean-square and contour estimates remain unresolved.

Contents

1 Scope and conventions **4**

2 A corrected conditional closure theorem **4**

3 Arithmetic obstructions to the claimed completion **5**

4 Determinants and lost arithmetic information **5**

5 A necessary change of approximation class **7**

6 A convergent arithmetic replacement **7**

7 A certified test of the arithmetic pencil candidate **8**

8 Jensen approximants and a coefficient-defined metric **9**

9 Extending an Euler correction and locating an additional factor **9**

10 The compensated principal residue at perturbed parameters **11**

11 A frozen-capacity counting interface **11**

12 The perturbed low estimate **12**

13 The nonprincipal central-contour interface **13**

14 A finite arithmetic input: local correlations **14**

15 Periodic cancellation in the Gram estimate **15**

16 A concrete fixed arithmetic period **16**

17 Arithmetic extraction in a reflected block **18**

18 Consequences for a revised research article **19**

19 Why an absolute coprime pair-square estimate cannot close the short rows **19**

A Worked geodesic calculations and possible replacements **20**

B Character examples and induction bookkeeping **20**

C Local algebra and topology checks **21**

D Why weak convergence and boundary analogies are insufficient **21**

E Spectral approximation can conceal the original hypothesis **22**

F Hermitian dilation and the spectral parameter **22**

G Positive pencils, resolvent control, and a Herglotz test **23**

H Quantitative arithmetic approximation and prime regularization **24**

I Exact arithmetic behind the finite-level zero certificate **25**

J Jensen companions: exact normalization and the failed moment argument **26**
J.1 The coefficient construction 26
J.2 Why ordinary moment positivity fails 27

K Euler-correction estimates and the shifted angular character **28**
K.1 All valuation cases in the larger region 28
K.2 Principal factor, tails and nonvanishing 29
K.3 What the additional factor permits and prevents 29

L Principal multiplier, weighted slots and the extended residue comparison **30**
L.1 Local errors and the scalar normalizer 30
L.2 Contour powers and the perturbed geometry 31
L.3 A pointwise obstruction at 3/5 32

M Research tasks with mathematical acceptance criteria **32**

N Details of the frozen row-count extension **33**

O Support and offsets in the low-energy application **35**

P Central conductor, height and integral safeguards **36**

Q Prime-power and common-factor arithmetic details **38**

R Gram cancellation: masks, frequency volume and tails **39**

S From the phase table to the literal column period **41**

T Exact reflected phases, shared cubes and the common density **42**

U Computational examples and reproducibility **43**

V Weighted profiles, exact arithmetic examples and the surviving task **44**

1 Scope and conventions

For a primitive Dirichlet character χ of conductor q, write

$$L(s,\chi)=\sum_{n\geq 1}\frac{\chi(n)}{n^s},\qquad \Lambda(s,\chi)=\left(\frac{q}{\pi}\right)^{(s+a)/2}\Gamma\left(\frac{s+a}{2}\right)L(s,\chi),\quad a=\frac{1-\chi(-1)}{2}.$$

The series and Euler product initially apply for $\operatorname{Re} s>1$. Our target is Dirichlet GRH: every zero in $0<\operatorname{Re} s<1$ has real part $1/2$. For $q=1$, use the entire function $\xi(s)=\frac{1}{2}s(s-1)\Lambda(s,1)$ when an entire target is required. Finite-order Hecke characters and general Artin representations require separate constructions; a theorem for one family is not silently a theorem for another.

The supplied surface-order manuscripts [1, 2] propose arithmetic transfer operators and self-adjoint approximants. We isolate valid closure logic and prove obstructions to particular claimed constructions. These obstructions refute intermediate statements, not GRH. The separate `math` repository [3] includes claimed zero-free regions strictly weaker than GRH. Such a region cannot replace the missing critical-line argument.

2 A corrected conditional closure theorem

Theorem 2.1 (Spectral approximation criterion)**.** *Let $U\subset\mathbb{C}$ be connected and open, let f be holomorphic on U and not identically zero, and let*

$$F_N(s)=e^{a_N+b_N s}\det(sI-(\tfrac{1}{2}I+iH_N)),\qquad H_N=H_N^*\in M_{d_N}(\mathbb{C}).$$

If $F_N\to f$ locally uniformly on U, then every zero of f in U has real part $1/2$.

Proof. The eigenvalues $\lambda_{j,N}$ of H_N are real. Thus the zeros of F_N are $1/2+i\lambda_{j,N}$, with algebraic multiplicities; the exponential factor never vanishes. If $f(z)=0$ off that line, choose a closed disk $D\subset U$ about z disjoint from the line, with f nonzero on its boundary. Such a disk exists because the zeros of a nonzero holomorphic function on a connected domain are isolated. Let $m=\min_{\partial D}|f|>0$. Local uniform convergence gives $|F_N-f|<m$ on ∂D for sufficiently large N. Rouché's theorem forces F_N and f to have the same positive number of zeros inside D, contradicting the zero location of F_N. □

Remark 2.2. This is a conditional criterion, not a construction. Self-adjointness alone establishes only the zero location of the approximants. The difficult arithmetic statement is convergence to the particular completed L-function. Pointwise convergence, weak operator convergence, and agreement of finitely many zeros do not supply the displayed hypothesis.

Proposition 2.3 (A usable continuation interface)**.** *Suppose U is a connected domain, the entire functions F_N are locally uniformly bounded on U, and they converge pointwise on a nonempty open subset $V\subset U$ to a holomorphic function f defined on U. Then $F_N\to f$ locally uniformly on U.*

Proof. Montel's theorem makes the family normal. Every subsequence has a further subsequence converging locally uniformly to a holomorphic function g. On V, pointwise convergence gives $g=f$, so the identity theorem gives $g=f$ on U. If the original sequence failed to converge locally uniformly, a compact set, positive error, and violating subsequence would contradict this subsequence property. □

A proposed analytic continuation proof must actually provide the uniform bounds and a genuine overlap V. The disjoint domains $\operatorname{Re} s>1$ and $0<\operatorname{Re} s<1$ are not an overlap.

3 Arithmetic obstructions to the claimed completion

Theorem 3.1 (No integral prime-power geodesic norms)**.** *Let $\gamma \in \mathrm{SL}_2(\mathbb{Z})$ be hyperbolic. Its expanding eigenvalue in absolute value is $u > 1$, and its hyperbolic norm is $N(\gamma) = u^2$. Then $N(\gamma)$ is not an integer greater than one. In particular no closed geodesic arising from such a matrix has length $k \log p$ for a prime p and integer $k \geq 1$.*

Proof. Put $t = |\operatorname{tr} \gamma| \in \mathbb{Z}$, so $t = u + u^{-1}$. If $u^2 = n \in \mathbb{Z}$, $n > 1$, squaring gives

$$t^2 = n + 2 + \frac{1}{n}.$$

The left-hand side and $n + 2$ are integers, so $1/n$ is an integer, contradicting $0 < 1/n < 1$. Since length is $\log N(\gamma)$, length $k \log p$ would imply $N(\gamma) = p^k$, already excluded. □

The statement applies to every subgroup of $\mathrm{SL}_2(\mathbb{Z})$, hence to congruence surfaces. The prime geodesic theorem is an asymptotic counting analogy; it cannot provide the termwise identification claimed in the completion section of [1]. Hecke correspondences of determinant p must not be identified with hyperbolic deck transformations of determinant one.

Theorem 3.2 (Failure of integral cyclic induction)**.** *The trivial character of $G = C_2 \times C_2$ is not an integer linear combination of characters induced from one-dimensional characters of cyclic subgroups of G.*

Proof. Every cyclic subgroup has order one or two. The degree of any such induced character is its index, namely four or two. Evaluation at the identity sends every integer linear combination to an even integer, whereas the trivial character has degree one. □

This distinguishes rational Artin induction from integral Brauer induction. It invalidates the stated cyclic integral reduction, while leaving appropriate forms of these classical induction theorems available for a revised argument.

Theorem 3.3 (Prime-tail obstruction)**.** *For every real $\sigma \leq 1$, the positive series $\sum_p p^{-\sigma}$ diverges. Thus $\sum_{p>N} p^{-\sigma}$ is not a vanishing absolutely convergent tail in the critical strip.*

Proof. It suffices to show $\sum_p 1/p = \infty$, since $p^{-\sigma} \geq p^{-1}$. Suppose the reciprocal-prime series converged. For $0 \leq x \leq 1/2$, $-\log(1 - x) \leq 2x$, for example by integrating $(1 - t)^{-1} \leq 2$ from zero to x. Hence the finite Euler products $\prod_{p \leq X}(1 - 1/p)^{-1}$ would be bounded uniformly in X. Expanding each as a finite product of absolutely convergent geometric series gives the sum of $1/n$ over all positive integers with prime factors at most X. In particular it dominates $\sum_{n \leq \lfloor X \rfloor} 1/n$, which is unbounded. Contradiction. □

Any repair below $\operatorname{Re} s = 1$ must introduce proved cancellation or a regularization, with quantitative control; absolute prime-tail bounds cannot suffice.

4 Determinants and lost arithmetic information

For $T = \operatorname{diag}(\chi(p)p^{-s})_{p \leq N}$ one has exactly

$$\det(I - T) = \prod_{p \leq N} (1 - \chi(p)p^{-s}) = L_N(s, \chi)^{-1},$$

not L_N. On an absolute-convergence region,

$$\log\det(I-T) = -\sum_{k\geq 1}\frac{\operatorname{tr}(T^k)}{k}.$$

The reciprocal orientation matters before any analytic continuation. Moreover Hermitian-part extraction does not preserve this determinant:

$$T = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \frac{T+T^*}{2} = 0, \quad \det(I-T) = 2 \neq 1 = \det(I).$$

An identity for the transfer determinant therefore does not automatically identify the centered characteristic polynomial of a symmetrized operator.

Proposition 4.1 (A direct obstruction to the supplied diagonal diagnostic)**.** *For the $q = 1$ model evaluated at $s = 1/2$, the algorithm that sets $\lambda_p = -\arg(p^{-1/2})$ gives $H_N = 0$ and $F_N(s) = (s-1/2)^{d_N}$. These functions cannot converge locally uniformly to $\Lambda(s,1)$ or $\xi(s)$ on a neighborhood of $1/2$.*

Proof. Every $p^{-1/2}$ is positive, so its argument is zero and $F_N(1/2) = 0$. On the other hand the alternating Dirichlet series $\eta(1/2) = \sum_{n\geq 1}(-1)^{n-1}n^{-1/2}$ is positive: grouping into adjacent pairs gives positive terms and its sum is at least $1 - 1/\sqrt{2} > 0$. The continuation identity $\eta(s) = (1-2^{1-s})\zeta(s)$ gives $\zeta(1/2) < 0$. The gamma factor at $1/4$ is positive and finite, so $\Lambda(1/2,1) \neq 0$, and its pole-removal factor at $1/2$ is nonzero. Even pointwise convergence at that point is impossible. □

Proposition 4.2 (Holomorphic Hermitian rigidity)**.** *Let $U \subset \mathbb{C}$ be connected and open, with a fixed inner product on $\mathbb{C}^d$. If $H : U \to M_d(\mathbb{C})$ has holomorphic entries and $H(s)^* = H(s)$ for every complex $s \in U$, then H is constant.*

Proof. In fixed orthonormal coordinates, for each vector v, the holomorphic scalar $v^*H(s)v$ is real-valued, so it is constant by the open mapping theorem. Polarization recovers all matrix entries from these quadratic forms, hence every entry is constant. Equivalently, $H_{ij} + H_{ji}$ and $i(H_{ij} - H_{ji})$ are real-valued holomorphic functions whose constancy determines H_{ij}. □

Thus a nonconstant arithmetic transfer $T(s)$ cannot retain entrywise holomorphy after pointwise Hermitian-part extraction: a holomorphic $(T+T^*)/2$ would be constant, and the remaining holomorphic skew-Hermitian part would be constant as well. This excludes that particular shortcut, not fixed self-adjoint operators. The spectral criterion uses H_N independent of the spectral variable s. Requiring self-adjointness only on a real parameter line is also consistent with a nonconstant holomorphic extension, but needs its own zero-location theorem.

Proposition 4.3 (A positive generalized spectral pencil)**.** *Let $A = A^*$ and $B = B^* > 0$ be fixed $d \times d$ matrices. Every zero of $\det(A - zB)$ is real. Consequently every zero of $\det(A + i(s-1/2)B)$ lies on $\operatorname{Re} s = 1/2$.*

Proof. Singularity supplies $v \neq 0$ with $Av = zBv$. Hence $z = (v^*Av)/(v^*Bv)$, with real numerator and strictly positive real denominator. Thus z is real. Substitute $z = -i(s-1/2)$. □

This supplies a valid finite spectral mechanism, but not its arithmetic construction. To apply it one must construct A_N, B_N, certify $B_N > 0$, and prove the required completed-L determinant approximation independently. Hermitian coefficients alone are insufficient: for

$$A = \begin{pmatrix} 0 & \varepsilon \\ \varepsilon & 0 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \qquad \det(A - zB) = -(z^2 + \varepsilon^2),$$

$0 < \varepsilon < 1/2$ gives transported off-line zeros $s = 1/2 \pm \varepsilon$ inside the critical strip.

5 A necessary change of approximation class

Theorem 5.1 (Zero-free Euler truncations cannot approximate a zero)**.** *Let U be a connected open subset of $\{\operatorname{Re} s > 0\}$, let g_N be holomorphic and nowhere zero on U, and let P_N be arbitrary finite sets of rational primes. Put*

$$D_N(s) = \prod_{p \in P_N} (1 - \chi(p) e^{-s \log p}).$$

Each of $g_N D_N$ and g_N / D_N is holomorphic and nowhere zero on U. If either sequence converges locally uniformly to a holomorphic f which is not identically zero, then f has no zero in U.

Proof. For $\operatorname{Re} s > 0$, $|\chi(p) e^{-s \log p}| \le p^{-\operatorname{Re} s} < 1$, so every local factor is nonzero. Hence the finite product, its reciprocal, and their normalized versions have the asserted property. If f had a zero, choose a closed disk inside U with a positive number of zeros of f in its interior and none on its boundary. Local uniform convergence and Rouché's theorem would force the zero-free approximants to have that positive number of zeros, a contradiction. □

More quantitatively, on any such disk D, every zero-free holomorphic approximant G satisfies

$$\max_{\partial D} |G - f| \;\ge\; \min_{\partial D} |f| > 0.$$

Otherwise Rouché yields the same contradiction. Thus improved estimates cannot repair this approximation class near a nontrivial zero. The construction must change so that finite approximants can have zeros in the strip; a nonvanishing gamma or exponential normalization cannot do this.

6 A convergent arithmetic replacement

For a nonprincipal character modulo q, put $A(x) = \sum_{n \le x} \chi(n)$ and $B = \sup_x |A(x)| \le q$. Complete periods sum to zero, so B is finite. Partial summation gives, for $\sigma = \operatorname{Re} s > 0$,

$$L(s, \chi) - \sum_{n \le N} \chi(n) n^{-s} = -A(N) N^{-s} + s \int_N^\infty A(x) x^{-s-1} \, dx,$$

and therefore

$$\left| L(s, \chi) - \sum_{n \le N} \chi(n) n^{-s} \right| \le B \left(1 + \frac{|s|}{\sigma} \right) N^{-\sigma}.$$

The endpoint tends to zero and the integral is absolutely convergent. The limit agrees with the Euler definition on $\operatorname{Re} s > 1$, hence with its analytic continuation. Multiplication by the correct gamma/conductor factor gives locally uniform approximation on $\operatorname{Re} s > 0$ for either parity.

For zeta, reproduce the pole before removing it. Set

$$Z_N(s) = \sum_{n \le N} n^{-s} + \frac{N^{1-s}}{s-1}.$$

Euler summation gives, for $\sigma > 0$ and $s \neq 1$,

$$\zeta(s) - Z_N(s) = -s \int_N^\infty \{x\} x^{-s-1}\, dx, \qquad |\zeta(s) - Z_N(s)| \le \frac{|s|}{\sigma} N^{-\sigma}.$$

The integral continues holomorphically to $\sigma > 0$. Consequently

$$\xi_N(s) = \frac{s}{2}\pi^{-s/2}\Gamma(s/2)\left((s-1)\sum_{n \le N} n^{-s} + N^{1-s}\right)$$

is holomorphic on that half-plane, including $s = 1$, and converges locally uniformly to ξ; in fact $\xi_N(1) = 1/2$ for every N. These constructions use known arithmetic coefficients, not unknown zeros. They avoid Theorem 5.1 by allowing additive cancellation and unrestricted finite-level zeros. They establish convergence, but provide no self-adjoint realization or critical-line theorem for those zeros.

7 A certified test of the arithmetic pencil candidate

Let χ be the primitive nonprincipal character modulo 3. Its fifth Dirichlet partial sum is the entire function

$$f(s) = S_5(s, \chi) = 1 - 2^{-s} + 4^{-s} - 5^{-s}.$$

Theorem 7.1 (Certified zero of a finite partial sum)**.** *Put*

$$c = (337261 + 11027504i)/10^6, \qquad r = 10^{-4}.$$

The disk $|s - c| < r$ contains exactly one zero of f, counted with multiplicity, and lies inside $0 < \operatorname{Re} s < 1/2$.

Proof. The exact rational procedure detailed in the extended edition establishes

$$|f(c)| < 10^{-6}, \qquad |f'(c)| > 1, \qquad \sup_{|s-c| \le r} |f''(s)| \le 9.$$

It uses 60 terms of the logarithm's positive atanh series and terms 0 through 120 of the exponential Taylor series, with explicit remainders; no floating-point value is used in these inequalities. On the boundary, Taylor's formula gives

$$|f(s) - f'(c)(s-c)| < 10^{-6} + \tfrac{9}{2}r^2 < r < |f'(c)(s-c)|.$$

Rouché gives exactly one zero, which is therefore simple. The real part throughout the closed disk belongs to $[337161, 337361]/10^6 \subset (0, 1/2)$. □

The completed fifth partial sum has the same zero because its conductor/gamma factor is holomorphic and nonzero on this disk. It cannot equal a nowhere-zero holomorphic factor times the determinant of a fixed positive Hermitian pencil there. This rejects that exact finite-level identification. It does not reject convergence of the partial sums, modified approximants, later truncations, subsequences, or GRH itself: f is a finite partial sum, not the full L-function.

8 Jensen approximants and a coefficient-defined metric

This alternative currently concerns the Riemann target only. Define

$$\gamma_j = \frac{j!\,\xi^{(2j)}(1/2)}{(2j)!}, \qquad F(w) = \sum_{j\ge 0} \frac{\gamma_j}{j!} w^j, \qquad \Xi(t) = \xi(1/2 + it) = F(-t^2).$$

The classical positive Fourier kernel of Ξ gives $\gamma_j > 0$ [9]; evenness and entirety of ξ make F entire. Set, for $d \ge 1$,

$$J_d(x) = \sum_{j=0}^{d} \binom{d}{j} \gamma_j x^j, \qquad Q_d(w) = J_d(w/d), \qquad P_d(t) = Q_d(-t^2).$$

Proposition 8.1 (Explicit convergence without a zero-location premise)**.** *For $R, T \ge 0$,*

$$\sup_{|w|\le R} |Q_d(w) - F(w)| \le \frac{R^2 F''(R)}{2d}, \qquad \sup_{|t|\le T} |P_d(t) - \Xi(t)| \le \frac{T^4 F''(T^2)}{2d}.$$

Proof. The coefficient multiplier $w_{d,j} = j!\binom{d}{j}/d^j$ for $j \le d$, and 0 for $j > d$, satisfies $0 \le w_{d,j} \le 1$ and $1 - w_{d,j} \le j(j-1)/(2d)$. For $j \le d$ use $1 - \prod_{k<j}(1 - k/d) \le \sum_{k<j} k/d$; for $j > d$ use $j(j-1) \ge 2d$. Summing the absolutely convergent series with nonnegative coefficients proves the bounds. □

Let C_d be the multiplication-by-x companion of the monic J_d/γ_d, and $H_d = (m_{i+j})_{0\le i,j<d}$, where $m_k = \operatorname{tr}(C_d^k)$ and $m_0 = d$. Newton identities compute H_d from the coefficients, and $H_d C_d = C_d^T H_d$. The Hermite–Sylvester criterion gives $H_d \ge 0$ exactly when J_d is real-rooted [8]; then its positive coefficients force all roots negative. If $H_d > 0$, the extended edition constructs an explicit real symmetric K_d with

$$P_d(t) = \frac{(-1)^d \gamma_d}{d^d} \det(tI - K_d).$$

Strict positivity is needed for that construction; semidefinite metrics can be singular. The classical Jensen criterion makes $H_d \ge 0$ for every d equivalent to zeta RH [7]. This identifies the missing positivity; it does not prove it. Fixed-degree eventual hyperbolicity as the shift grows does not settle this unshifted growing-degree family.

Proposition 8.2 (Positive smooth kernels do not supply the metric)**.** *Positivity of an even smooth Fourier density, including Gaussian decay, does not imply positivity of its Jensen Newton-sum metric.*

Proof. Convolve $\frac{9}{10}\delta_0 + \frac{1}{20}(\delta_{-1} + \delta_1)$ with a centered Gaussian of variance $v > 0$. Its Fourier transform is $e^{-vt^2/2}(9/10 + \cos t/10)$ and its density is strictly positive, even and Schwartz. Writing its associated $F(w) = \sum \gamma_j w^j/j!$ gives $\gamma_0 = 1$, $\gamma_1 = (v + 1/10)/2$ and $\gamma_2 = (3v^2 + 3v/5 + 1/10)/12$. Thus $\gamma_1^2 - \gamma_0\gamma_2 = -7/1200 < 0$; the degree-two Newton-sum metric has negative determinant. These are model coefficients, not the arithmetic ξ coefficients. □

9 Extending an Euler correction and locating an additional factor

We now test one missing analytic input of the September argument in [3]. All deductions in this section use its displayed scalar local identity; its preceding Gauss/reciprocity derivation and main mean-square theorem are not certified here.

Proposition 9.1 (Extension of the explicit local product)**.** *The displayed correction product* $\mathcal{H}_{\eta,u}(x,w,z)$ *converges normally on a neighborhood of*

$$\operatorname{Re} x \geq 3/4, \qquad \operatorname{Re} w \geq 19/20, \qquad \operatorname{Re} z \geq 33/200,$$

and is $O_\epsilon(q_u^\epsilon)$ *there. On the displayed closed region, for* $Q = Np$*, its local defects are at most* $1024Q^{-149/100}$ *when* $p \nmid u$ *and* $32Q^{-7/10}$ *when* $p \mid u$*. The latter is an upper bound, not a nonvanishing statement.*

Proof. Substitution in the explicit local formulas gives the exponent and denominator bounds detailed in the extended edition. All five positive row valuations are included. The ideal count sums the first bound; the divisor-product estimate handles the finitely many primes of u. The bounds retain a positive margin on a larger open region. □

For the principal row put $H_\eta(s) = \mathcal{H}_{\eta,1}(s,1,1/6)$. At a prime set

$$v = Q^{-1}, \quad d = \eta(p)Q^{-s}, \quad \theta(p) = \alpha(p)^{-6}\eta(p)^6, \quad r = \theta(p)Q^{3-6s},$$

where $\alpha(p) = p/|p|$ for a generator. Sixth powers remove unit ambiguity. Exact simplification gives

$$H_p = \frac{(1-v^2)[1-vr-(1-v)d]}{(1-r)(1-d)}, \qquad A_p = (1-r)H_p.$$

For $\operatorname{Re} s > 1/2$, the numerator is nonzero because $|vr + (1-v)d| < v + (1-v) = 1$. Moreover

$$A_p - 1 = -v^2 + \frac{(1-v^2)v(d-r)}{1-d}.$$

Thus $A = \prod_{p \notin S} A_p$ is holomorphic and nonzero on $\operatorname{Re} s > 1/2$, while H_η itself converges normally and is nonzero on $\operatorname{Re} s > 2/3$. On the latter region the exact factorization is

$$H_\eta(s) = A(s) L_F^S(6s-3,\theta).$$

The character θ has nontrivial angular factor α^{-6}; it has infinite order and is outside the source's finite-order target family. The correction can be made uniformly close to 1 on each fixed $\operatorname{Re} s \geq \sigma_0 > 2/3$ by a sufficiently large finite deletion. This supplies the proposed tiny perturbation's local Euler input; the compensated contour errors and mean squares remain unproved.

Proposition 9.2 (Limitation of the unchanged correction)**.** *Assume the scalar identity and the classical Hecke completion properties listed in the extended edition, including order one and conjugation. The analytic continuation of this unchanged* H_η *has a zero with real part at least* $7/12$*. It therefore cannot be nonzero throughout* $\operatorname{Re} s > \sigma_0$ *for any* $\sigma_0 < 7/12$*.*

Proof. The entire order-one completion of $L(s,\theta)$ has infinitely many zeros: otherwise Hadamard factorization would contradict its $\sigma \log \sigma + O(\sigma)$ logarithmic growth on the positive axis. Euler nonvanishing, the functional equation and conjugation give a zero ρ with $1/2 \leq \operatorname{Re}\rho \leq 1$. Neither gamma nor deleted Euler factors vanish there. Hence the displayed continuation vanishes at $(\rho+3)/6$, where A is nonzero and $\operatorname{Re} s \geq 7/12$. □

The Hecke analytic inputs are classical external dependencies, not Lean theorems here [10, 11]. No GRH or Artin holomorphy premise is used. This limits the specified probe and nonvanishing criterion; it does not refute GRH, the claimed 7/8 result, or different constructions.

10 The compensated principal residue at perturbed parameters

We retain the source's scalar local identity, exact high representation, Mellin integrated/trace tail contracts, and polynomial reciprocal control on $\operatorname{Re} s = \beta_* + e$. The following verifies a principal interface, not the remaining energy or row-count estimates. On the enlarged Euler region, the four local multiplier errors have powers

$$Q^{-x_r}, \quad Q^{-6z_r}, \quad Q^{4-5x_r-6z_r}, \quad Q^{1-w_r-6z_r}.$$

Their common saving is at least $37/50$. The local lower bound for H_p therefore gives $\mathcal{B}_p = -1 + O(Q^{-37/50})$, uniformly in all heights. On fixed real boxes the full principal correction is $O(Z^{\ell \operatorname{Re} z})$, with height degree zero.

For a nonempty fixed slot system with positive lengths ℓ_i, set

$$S_i = \sum_{p \in \mathcal{P}_i(Z)} W_i(Q/P_i) Q^{-5/6}, \qquad A_T = (-1)^K Z^{-\ell/6} \prod_i S_i, \qquad \mu = \tfrac{7}{10} \min_i \ell_i > 0.$$

Fixed-ray prime counting gives $S_i > 0$ eventually and $A_T \sim c_T (\log Z)^{-K}$ with $c_T \neq 0$, of sign $(-1)^K$ [12]. No uniformity in a moving ray conductor is used. Nonnegative weights give the exact residue factorization

$$\mathfrak{H}_{\eta,1,Z}(s, 1, 1/6) = H_\eta(s) Z^{\ell/6} A_T(Z)(1 + \mathcal{R}_{\eta,Z}(s)), \qquad \mathcal{R}_{\eta,Z} = O(Z^{-\mu}).$$

Indeed the ith relative error is the negative weighted average of $\mathcal{B}_p + 1$, of size $O(P_i^{-37/50})$; a fixed finite product preserves that saving. Also $|A_T|^{-1} \ll_\epsilon Z^\epsilon$.

With the explicitly inherited contracts above, the source's principal contour comparison now extends to $3/4 \le \sigma_0 < \beta_*$. Its three remainders, relative to $C(\beta_*)$, have exponents

$$(1+h)e - l_y/20 + \epsilon, \qquad e - h/600 + \epsilon, \qquad e - \mu + \epsilon.$$

The extended edition derives these by crossing only $w = 1$ and $z = 1/6$. All three are negative for sufficiently small pretarget e, ϵ. The perturbed geometry at $t = 10^{-5}$ satisfies this interface, with $C(s) = s - 11/16$ and target $349999/400000$. This is not a new zero-free theorem: other low and nonprincipal contour inputs remain unverified; the following counting interface is conditional on its moment inputs.

There is a separate pointwise limitation. At the principal residue, putting $N = 1 - vr - (1-v)d$ gives

$$\mathcal{B}_p + 1 = \frac{(1-v)[r/d - vr - (1-v)d]}{N}.$$

For fixed $1/2 < \sigma = \operatorname{Re} s \le 3/5$ its norm is asymptotic to $Q^{3-5\sigma}$, uniformly in phases. In particular, for $\eta = 1$ and an inert rational prime, $\alpha^{-6} = 1$ and real $s = 3/5$ give $r = d$, so $\mathcal{B}_p = -1/Q$ exactly. The error around -1 tends to 1. Such primes can occur in the fixed identity ray. This excludes the pointwise power-saving argument there, without excluding cancellation after prime averaging or a different normalization.

11 A frozen-capacity counting interface

The next proposition concerns the September probe, not the spectral construction. Its arithmetic moment and witness inputs remain uncertified. The previously claimed $7/8$ bound is an explicit hypothesis here.

Proposition 11.1 (Conditional row count)**.** *Assume $\beta_* \le 7/8$ and the source's marked inverse moment, plain fourth moment, sixth-power amplification and simultaneous saturated witness statements, with their common profiles, masks and height bounds. For each $\epsilon > 0$, first choose capacity decrements and a slot-mesh threshold independent of the eventual fixed slot count. Then fix a finite slot system satisfying it and a retained nonfloor bin, $a > 51/100$, of sixth-power-free rows of norm $\asymp U = Z^d$, where $d \ge d_{\min} > 0$. Put $\delta = 2a - 1 \le 3/4$, $x = q/\delta \in [0, 1/2]$, where q is its length-weighted prime amplitude. Suppose the total available prime-slot length ℓ/d exceeds $7/37$ by a fixed margin. Then*

$$\#\mathcal{B} \ll U^{R_*+\epsilon}(1+T_1)^A, \quad R_* = 1 - \delta + \frac{(\alpha-\delta)\delta P}{2J}, \quad \alpha = 5/6,$$

where $D = 3 - 17x/9$, $P = (2 - 8x/9)(1-x)$ and $J = (\alpha-\delta)D + \delta P > 0$. The finite height order A is fixed before the external tail order. Without selected prime slots one instead has exponent $1 - 2\delta/3$, with no supply assumption.

Proof. The plain moment permits $\kappa = 3/4$ since $\beta_* \le (1+\kappa)/2$. Its exact zero-loss capacity is $z_P(m) = 2(1-2m)/9$; the inverse capacity is $z_M(r) = (1-r)/2$. Thus the source's $\Delta/4$ comparison loss is absent, rather than continued to a negative bootstrap parameter. At detector cutoff $\tau \in [1, 3/2]$, the saturated witnesses have $r + m \ge \tau - o(1)$, $|M_r|^2 \gg U^{\delta r - \epsilon}$ and $|S_m|^2 \gg U^{\delta m - \epsilon}$. Selecting whole positive slots of requested length z gives squared gain $U^{2qz - \epsilon_{\mathrm{round}}}$, with the fixed rounding loss $\epsilon_{\mathrm{round}}$ allocated within ϵ. Division of the moments by these spikes, using $|S_m|^4$ for the plain moment, gives the affine short counts

$$A_I(r) = 1 - \delta[x + (1-x)r], \quad S_\tau(r) = 1 - \delta[4x/9 + (2 - 8x/9)(\tau - r)].$$

They cross at $r_* = [(2 - 8x/9)\tau - 5x/9]/D$, with $23/37 \le r_* \le 1$ and $1/3 \le \tau - r_* \le 1/2$. Choose the plain count below r_* and the inverse count above it. Their common upper bound is $R_{\mathrm{short}}(\tau) = 1 - \delta + (\delta P/D)(3/2 - \tau)$. The requested inverse length is at most $7/37$, and the plain length at most $2/27 + o(1)$. A fixed capacity decrement preserves the strict marked-moment margins. Near zero capacity use the no-slot moments. For $r \ge 1$, amplification gives $L(\tau) = 1 - \delta + (\alpha - \delta)(\tau - 1)$. The maximum of these short and long bounds is minimized at $\tau = 1 + \delta P/(2J)$, giving R_*. With no selection, use $\tau = 1$; the short lines cross at $r = 2/3$, giving $1 - 2\delta/3$. All rounding and witness losses fit the prescribed ϵ. □

For $\ell_t = 1/6 + t$, $h_t = 13/16 + 3t/2$, the supply holds uniformly through $d \le h_t + \zeta$ whenever

$$0 \le \zeta < \frac{37\ell_t}{7} - h_t = \frac{23}{336} + \frac{53t}{14}.$$

This closes the frozen-capacity counting interface at $t = 10^{-5}$ under its explicit inputs. It does not certify those arithmetic moments, the earlier $7/8$ theorem, the low reflected-energy application or the remaining nonprincipal contours. Floor-bin rows need separate estimates.

12 The perturbed low estimate

Proposition 12.1 (Conditional low interface)**.** *Assume the source's general reflected-energy estimate, completed-row and sector identities, quantitative Gram bound and exact finite low separation, with their stated coefficient, support, smoothness and height uniformities. For $0 \le t \le 1/30$, put*

$$\ell_t = 1/6 + t, \quad M_t = 5/6 - t, \quad l_x = 17/48 - t/2, \quad l_y = 23/48 - t/2.$$

For every fixed finite system of disjoint physical prime slots with positive lengths of total ℓ_t and the original admissible smooth weights, the same finite compensated probe satisfies

$$|I_{\eta,\mathrm{modified},t}(Z)| \ll Z^{3/16-t/4+\epsilon} \qquad (\epsilon > 0).$$

The constant and threshold may depend on that slot system; the exponent has no mesh loss. No prior zero-free theorem is needed for this implication.

Proof. For a rescaled subset of length d, put $M' = M_t - 2d$ and $\ell' = \ell_t - d$. The retained dual-range calculation keeps $M' + \ell' - 1 = -3d$. The general energy formula gives

$$\sum_{0<q_m\ll Q} |B_m^J|^2 \ll Z^{M'+s_t(d)+\epsilon}, \qquad s_t(d) = \left(\frac{-1/6 + 5t + d}{4}\right)_+.$$

The extended edition verifies the support mapping and a uniform exponent bound including finite logarithmic offsets; the general arithmetic energy estimate remains a hypothesis. The tuple ratio $r_J = q_{p_J}/Z^d$ stays in a fixed compact subset of $(0,\infty)$, and

$$X' = Z^{l_x-d}/r_J, \quad Y' = Z^{l_y-d}/r_J, \quad Q = q_{b_*}X'Y', \quad P_a = Y'^2/Q = q_{b_*}^{-1}Z^{1/8}.$$

Uniformly over subsets, $l_x - d \ge 11/80$, $l_y - d \ge 21/80$, $M' \ge 2/5$ and $l_y - d - 11/48 \ge 1/30$. Thus $Q, Y', P_a \ge 1$ eventually and $P_a^2/Y' \ll P_a^{1/6}$. The assumed Gram bound and Cauchy–Schwarz give the unscaled tuple bound $(X')^{1/2}P_a^{1/12}Z^{s_t(d)/2+\epsilon}$, with a fixed polynomial height cost integrated against the external smooth Mellin transform. The number of tuples, their coefficient and $X'^{1/2}$ contribute $d - 3d/2 - d/2 = -d$. Since $t \le 1/30$ and $d \ge 0$,

$$s_t(d) \le d/4, \qquad -d + s_t(d)/2 \le -7d/8 \le 0.$$

Sum the finitely many subsets after allocating the small losses. The final exponent is $l_x/2+(1/8)/12 = 3/16 - t/4$. □

The potential row loss must be retained in this proof, rather than set to zero for every subset. The real-power absorption and reflected exponent inequalities are kernel-checked components; the arithmetic energy and Gram bounds are not. This closes the conditional low interface at $t = 10^{-5}$, leaving the nonprincipal contours and the deep source inputs unverified. It gives no unconditional new zero-free theorem.

13 The nonprincipal central-contour interface

The principal second Euler region does not govern the following move. Write $\mathcal{D}_1(\varepsilon_0)$ for the region $\operatorname{Re} s \ge 51/100$, $\operatorname{Re} z \ge 17/50$, $\operatorname{Re} w \ge -1/100$, $\operatorname{Re}(s+w) \ge 1+\varepsilon_0$.

Proposition 13.1 (Conditional central accounting)**.** *Retain the exact arithmetic high identity, full-correction holomorphy and polynomial height bounds on $\mathcal{D}_1$, the source's buffered detector bounds, and both integrated and trace-tail contracts. Fix $\beta_* > \sigma_0 \ge 51/100$ and a retained nonprincipal bin of physical rows $q_u \asymp U = Z^d$, with fixed $0 < d_{\min} \le d \le d_{\max} < \infty$ and $51/100 \le a \le \beta_* \le 1$. Exclude the detector's bounded exceptional rows. Take $T_1 = Z^\tau > 2$, with positive τ within $d_{\min}/100$ and the retained detector height ceiling. For $0 < e < 10^{-3}$ the original bin integral moves to*

$$\operatorname{Re} s = a + 16e, \qquad \operatorname{Re} w = 1 - a - 6e, \qquad \operatorname{Re} z = z_0 = 17/50$$

with error $O_{\eta,N}(Z^B T_1^{-N})$. Here B and internal height orders are fixed before N. On these retained contours suppose an admissible finite decomposition and pointwise row partitions satisfy

$$\sum_{u\in\mathcal{C}} |L^S(w,\chi(u))\mathfrak{H}_u^{(\lambda)}| \ll U^{R+\delta/2+\epsilon_c} Z^{\ell(z_0-1/2)+g+\epsilon_c}(1+T_1)^A,$$

uniformly, with $\epsilon_c \geq 0$, bounded R, g and polynomial logarithmic multiplicity. For every $\epsilon_d, \epsilon_p > 0$, the original bin contribution is bounded by

$$Z^{C(\sigma_0)+\sup E_{\sigma_0}(d;R,g)+(16-6l_y)e+(1+d)\epsilon_c+d\epsilon_d+\epsilon_p}(1+T_1)^A + O_{\eta,N}(Z^B T_1^{-N}),$$

where $\delta = 2a-1$, $h = 1-l_x+\ell$ and

$$E_\sigma = a-\sigma+h(z_0-1/6)-al_y-\ell/2+g+d(R+\delta/2-z_0).$$

Auxiliary retained coordinates and discarded integrations must obey the same single cumulative buffered-height allocation.

Proof. Move the full holomorphic tuple correction before introducing pieces. First move z to z_0 and s to β_*+20e with $w=3$. The reciprocal stays to the right of the actual supremum β_*. Next move w to $1-a-6e$; since $a \leq \beta_*$, $\mathrm{Re}(s+w) \geq 1+14e$. Discard the original height tails while the reciprocal is global, then move the retained s segment to $a+16e$. The rectangle has $\mathrm{Re}(s+w) \geq 1+10e$ and stays in $\mathcal{D}_1$. The buffered hypotheses exclude denominator zeros there. The numerator is entire and nonprincipal, and $6z_0 > 1$, so no pole is crossed. Horizontal joins use the trace contract; the buffered s-axis is never extended beyond its buffer.

Only after this move apply triangle and the aggregate bounds at each point. Dominate all occurring powers by one supremum and integrate the original analytic row sum; no individual pointwise partition is integrated. The outside power is $l_x(1/2-z_0)+a+z_0-1-al_y-dz_0+(16-6l_y)e$. Adding the assumed row power and subtracting $C(\sigma_0) = \sigma_0+l_x/2-1+h/6$ proves the formula. □

On the retained region only, local nonvanishing permits the main/error factorization with $g = q\ell$. Strict ramified errors spend the conductor deficit simultaneously: for distinct labels of valuation $j_p \geq 2$, $A_* = 1/2-w_r \geq 0$ gives $-w_r-(j_p-1)A_* \leq -1/2$. The extended edition records the allocation. For the perturbed geometry, at fixed algebraic δ, q, R, d,

$$E_t(d)-E_0(d) = t(13/50+\delta/4+q).$$

Thus the conditional central interface covers $t = 10^{-5}$ without a new $7/8$ restriction. It establishes neither negativity of every row exponent nor the underlying arithmetic inputs, outer rows or GRH.

14 A finite arithmetic input: local correlations

The following classical finite-field calculation is unconditional and is verified for actual finite fields and multiplicative characters. It does not establish a global character moment or GRH.

Theorem 14.1 (Zero-preserving local kernel)**.** *Let k be a finite field of size P, and let $A : k \to \mathbb{C}$ be a multiplicative character extended by zero at 0, including when principal. Put*

$$K_A(a,b;t) = \sum_{ax-by=t} A(x)\overline{A(y)}.$$

For $a, b \neq 0$,

$$K_A(a,b;t) = A(b/a)\begin{cases} P-1, & t=0, \\ P-2, & t\neq 0,\ A=1, \\ -1, & t\neq 0,\ A\neq 1. \end{cases}$$

If exactly one coefficient is zero, the value is $(P-1)1_{A=1}1_{t\neq 0}$. If both vanish, the value is $(P-1)^2$ when $A=1, t=0$, and zero otherwise.

Proof. The autocorrelation $C_A(h) = \sum_y A(y+h)\overline{A(y)}$ equals $P-1$ at $h=0$. For $h\neq 0$, scale $y = hz$ to reduce to $h=1$. Changing z to $-z$ gives $C_A(1) = A^{-1}(-1)J(A^{-1}, A)$. The Jacobi identities give -1 for nonprincipal A and $P-2$ for principal A; the pinned formal proofs are recorded in [13]. For $a, b \neq 0$, solve $x = (t+by)/a$ and factor $A(b/a)$ to obtain $A(b/a)C_A(t/b)$. If $b=0, a\neq 0$, the fixed value $x = t/a$ and the remaining character sum give the stated result. Swapping x, y and conjugating gives the other case. With both coefficients zero, the sum is $|\sum A|^2$ at $t=0$ and empty otherwise. □

For a good prime $\mathfrak{p}$ and common exponent $c \geq 1$, take $A = \chi_{\mathfrak{p}}^c$ and $(a,b) = (n_2, n_1)$ in the residue field. On the genuine coprime-residual locus at least one coefficient is a unit. Each field solution has exactly P^{c-1} lifts modulo $\mathfrak{p}^c$, so the common-prime local factor is $P^{c-1}K_A(a,b;t)$. Its norm is at most P^c. A residual modulus meeting the common prime introduces no extra multiplicity. CRT multiplies these local congruence sums.

The restriction matters. For the principal character on $\mathbb{F}_7$, $K_1(0,0;0) = 36 > 7$, an actual sum checked in Lean. Thus the genuine kernel bound cannot hold on all residual pairs. The source explicitly uses an artificial zero extension off the coprime-residual locus; this is not a refutation of its stated finite identity.

The Gauss normalization is also consistent. For nonprincipal A and nontrivial additive ψ, the checked finite identity is $|g(A,\psi)|^2 = P$. Choose the fixed primary generator π of $\mathfrak{p}$. Splitting a residue as $x = x_0 + \pi y$ gives the factor $P^{n/2-1}$ after additive orthogonality. Therefore

$$|G(\mathfrak{p}^n, t)| = P^{(n-1)/2}1_{v_{\mathfrak{p}}(t)=n-1} \quad (6 \nmid n),$$

For $6 \mid n$ the value, including its sign, is $G(\mathfrak{p}^n, t) = P^{n/2}1_{\mathfrak{p}^n|t} - P^{n/2-1}1_{\mathfrak{p}^{n-1}|t}$. Finite-field Gauss and kernel identities are formalized; quotient-ring lifting, CRT and the source's arithmetic phase identification remain prose. No full Gram, reflection or moment estimate follows from this finite input alone.

15 Periodic cancellation in the Gram estimate

Lemma 15.1 (Annular periodic Poisson)**.** *Let $\mathcal{O} = \mathbb{Z}[\omega] \subset \mathbb{C}$, $r \in \mathcal{O}\setminus\{0\}$, $q = |r|^2$, and let $a : \mathcal{O}^2 \to \mathbb{C}$ be periodic modulo r in both columns, with $|a| \leq M$ and complete mean zero. Let $W \in C_c^\infty(\mathbb{C}^2)$ be supported on fixed annuli $0 < c \leq |z_j| \leq C$, with sufficiently many fixed seminorms bounded by K. For $N > 0$ and $A > 2$,*

$$\left|\sum_{m\in\mathcal{O}^2} a(m)W(m/\sqrt{N})\right| \ll_{A,c,C} KMN^2 \min\{1, (q/N)^A\}.$$

Proof. Normalize the finite Fourier coefficient by q^{-2}, the reciprocal number of residue pairs. Its norm is at most M, and its trivial coefficient is zero. Fourier inversion followed by lattice

Poisson therefore gives N^2/v^2 times one sum over the dual of $(r\mathcal{O})^2$, where v is the covolume of $\mathcal{O}$. There is no additional q^2 factor. For $N \geq q$, remove the zero frequency and use Fourier decay of order $2A > 4$. The dual spacing is comparable to $q^{-1/2}$, uniformly in the angle of r, giving $\sum_{\xi \neq 0}(1+\sqrt{N}|\xi|)^{-2A} \ll (q/N)^A$. For $N < q$, fixed annular support makes the original sum empty below a fixed positive N, and otherwise allows $O(N^2)$ lattice pairs. □

The hypothesis at small N matters: a mean-zero periodic coefficient with $a(0) = 1$ and a smooth bump with $W(0) = 1$ can leave the single origin contribution as $N \to 0$. The source weights are annular; this example refutes dropping that condition, not the source estimate.

Proposition 15.2 (Conditional Gram reconstruction)**.** *Retain the September source's exact column/row Poisson representation, artificial common kernel, original zero masks and fixed reciprocity/ray period identification. Assume the stated joint smooth seminorm bounds. Retain its prescribed fixed polynomial ranges, in particular $Z \geq 2$ and $1 \leq Q, Y' \leq Z^{K_0}$ for fixed K_0. For its fixed ray coefficients, put $P = Y'^2/Q \geq 1$. Then the periodic cancellation and positive dyadic summation give*

$$\sum_{q_m \ll Q} |A_m|^2 \ll (1+|\nu|)^J \frac{Q}{Y'} \left(1 + P^{1/6} + \frac{P^2}{Y'}\right) Z^{\epsilon}.$$

Proof. Insert the full Möbius sum before separating residual columns. For a nonexceptional frequency k, CRT isolates a nonprincipal residue-field character; its sum is zero. The joint period satisfies $q_r \ll q_C q_k$. In the shell $q_C q_k \asymp RP$, with dyads $q_C \asymp C_0$, $q_d \asymp D_0$, use $N = Y'/(C_0 D_0)$ in Lemma 15.1. The label counts and outer normalization give

$$\frac{Q}{Y'^3}\left(C_0 D_0 \frac{RP}{C_0}\right) C_0 \left(\frac{Y'}{C_0 D_0}\right)^2 = \frac{Q}{Y'}\frac{RP}{C_0 D_0}.$$

For $\Lambda = Y'/(RP)$ and $A \geq 2$, the remaining dyadic sum is

$$S_A(\Lambda) = \sum_{n \geq 0} (n+1) 2^{-n} \min\{1, (2^n/\Lambda)^A\} \ll \frac{1 + \log(2+\Lambda)}{\Lambda}.$$

Indeed, if $2^L \leq \Lambda < 2^{L+1}$, the head is at most $[L2^{L+1}+1]/\Lambda^2$ and the tail equals $(L+3)2^{-L}$; if $\Lambda < 1$, the sum is 4. Summation in R with a sufficient smooth decay gives P^2/Y'. Exceptional k have sixth-power ideals apart from primes of C and a fixed modulus; their count is $\ll_\epsilon (RP/C_0)^{1/6} q_C^{\epsilon}$. The trivial column bound leaves summable factors $C_0^{-1/6+\epsilon} D_0^{-1}$, giving $P^{1/6}$. The exact zero-frequency diagonal gives 1. □

Nine Lean declarations check the finite normalized Fourier bound, zero mode, independent-coordinate cancellation, exact dyadic head, infinite total and tail, head inequality and scale identity. The lattice argument above is prose. Applying it to the source still requires the listed arithmetic representation and fixed period; neither the full upstream theorem nor a GRH scope is certified by this reduction.

16 A concrete fixed arithmetic period

The arithmetic period retained in Proposition 15.2 can be supplied in prose using classical reciprocity and class field theory. This does not certify its other analytic representation hypotheses.

Proposition 16.1 (A common defining modulus)**.** *Let S be the fixed source prime set, including primes above 6, and let $\mathfrak{m}_{\mathrm{pres}}$ contain the full defining moduli of its fixed ray/coefficient presentations, including redundant zero primes. Put*

$$\mathfrak{m}_0 = (6)^6 \prod_{\mathfrak{p}\in S} \mathfrak{p}^{25}, \qquad \mathfrak{l} = \operatorname{lcm}((12), \mathfrak{m}_{\mathrm{pres}}, \mathfrak{m}_0).$$

The source's artificial two-column Gram coefficient at fixed $C, d, k \neq 0$ is periodic in each column modulo

$$\mathfrak{r} = \operatorname{lcm}(\mathfrak{l}, \operatorname{rad}(C), \operatorname{rad}(k)), \qquad N\mathfrak{r} \leq N\mathfrak{l}\, q_C q_k.$$

If some prime $p \nmid C\mathfrak{l}$ has $6 \nmid v_p(k)$, its complete joint mean is zero. The defining modulus $\mathfrak{l}$ is independent of C, d, k and Z.

Proof. For $a = u\prod_{\mathfrak{p}\in S} \pi_{\mathfrak{p}}^{e_{\mathfrak{p}}}$, $0 \leq e_{\mathfrak{p}} \leq 5$, the extension $E = F(a^{1/6})/F$ is cyclic of degree dividing six. Its power-residue character has conductor dividing the relative discriminant by the classical conductor–discriminant formula [14, Theorem V.3.27]. The minimal polynomial g is a monic integral factor of $X^6 - a$. The field discriminant divides the power-basis discriminant, and $\operatorname{disc}(gh) = \operatorname{disc}(g)\operatorname{disc}(h)\operatorname{Res}(g,h)^2$. Thus its conductor divides $(6)^6(a)^5$, hence $\mathfrak{m}_0$. This also covers reducible $X^6 - a$. Arithmetic Frobenius identifies this character with the sextic residue symbol [14, Sections VIII.5.3–5.5]. Actual S-valuations reduce modulo six on S-units.

For odd $c = a + b\omega$, Gaussian Poisson summation gives

$$\Gamma(c) = \frac{1 + i^{-b} + i^a + i^{b-a}}{2}.$$

Its values on square classes $1, -1, \lambda, -\lambda$ modulo 4 are $1, 1, i, -i$. The ratio $\Gamma(ab)/(\Gamma(a)\Gamma(b))$ is the symmetric sign bicharacter $\mathcal{R}(a,b)$. Classical cubic reciprocity [16, Equation (1.4)] and CRT for quadratic Gauss sums identify it with $\chi_b(a)/\chi_a(b)$ on coprime primary pairs. For general pairs its separate finite definition is retained.

Write $k = \zeta k_S k_{\mathrm{good}}$. Reciprocity gives, with every zero retained,

$$\chi_m(k) = \kappa_{\zeta,\boldsymbol{e}}(m)\mathcal{R}(k_{\mathrm{good}}, m) \prod_{p\mid k_{\mathrm{good}}} \chi_p(m)^{e_p}.$$

The first factor has conductor dividing $\mathfrak{m}_0$, the second has period 4, and the product has period $\operatorname{rad}(k_{\mathrm{good}})$. Fixed coefficients use $\mathfrak{l}$ and the artificial kernel uses $\operatorname{rad}(C)$. The substitutions by C, d introduce no larger period. All nonunit/primary zero branches are taken before inverses. At a nonexceptional prime outside $C\mathfrak{l}$, one nonprincipal field coordinate remains independent of all other factors; its mean is zero. □

The larger bad-prime powers are essential. The primary primes $\pi_{31} = 1 + 6\omega$ and $\pi_{43} = 1 - 6\omega$ agree modulo 12, but

$$\chi_{\pi_{31}}(\lambda) = \zeta_6, \qquad \chi_{\pi_{43}}(\lambda) = \zeta_6^5, \qquad \zeta_6 = 1 + \omega.$$

Indeed $\omega = 5$ in $\mathbb{F}_{31}$ and $\omega = 36$ in $\mathbb{F}_{43}$, giving $11^5 = 6$ and $30^7 = 7$ in the respective fields. Thus modulus 12 for the quadratic phase cannot replace the full fixed numerator modulus. The source already allows those higher powers. Thirteen Lean components check the actual quotient-algebra phase and period; the Gaussian, cubic-reciprocity and conductor arguments remain classical prose dependencies, with no zero-location assumption.

17 Arithmetic extraction in a reflected block

Proposition 17.1 (Marked coefficient separation)**.** *Assume the canonical completed-theta reflection formula with its stated cusp support and coefficient bound. Retain the source block hypotheses: disjoint slot supports, original product-form slot coefficients independent of rows and dual variables, and fixed row restrictions independent of the active tuples and dual variables except the explicit (P, R) mask. Freeze every nonresidual base prime, including every $j = 4$ prime, and fix separate full M_{ref}^2 sectors of the residual row product R and active marked product P. Then the reflected block has the coefficient independences of the explicit quadratic–cubic norm in the source's equation (4.6a), with every original zero retained. Its per-tuple central exponent is*

$$-\frac{v}{2} - \ell_b - \frac{e_\lambda}{3} - \frac{S_0 + B_0 + z_a}{2}.$$

This is a conditional arithmetic reduction, not the hybrid mean-square bound.

Proof. At an active prime write $t = c/p$. It is a unit, and $\sigma_p = \lambda^2 t$, $\epsilon_p = -\lambda^{-5} t^{-2}$. The local scalar factors as $\xi_{p,j}\chi_p(c/p)^{2j+2}$, with $\xi_{p,j}$ depending on that prime and the fixed datum only. This follows directly from the two unit powers; at $j = 4$ the exponent is $-2 \equiv 2j + 2 \pmod 6$. For distinct active primes, cubic reciprocity gives

$$\chi_p(q)^{2j+2}\chi_q(p)^{2k+2} = \left(\frac{q}{p}\right)_3^{j+k+2}.$$

Thus residual $j = 1$ and marked $k = 0$ phases cancel. All other prime interactions and the angular conductor factor separate into row, tuple or fixed scalars. The full sectors fix the cusp data.

The moving columns, including their zeros, are

$$\chi_R(nb^3)^3 = \chi_R(nb)^3, \qquad \chi_P(nb^3)^{-2} = \chi_P(n)^{-2} 1_{(P,b)=1}.$$

A frozen Ramanujan divisibility term is partitioned into $p \mid n_0$ and $p \nmid n_0,\ p \mid b_0$. In the first case extract only its squarefree occurrence; in the second extract one cube occurrence. For $\mu = u\lambda^h N_F n (B_F b)^3$, the coefficient bound gives

$$\frac{d(\mu)\alpha(\mu)\vartheta(\lambda^4\mu)}{\sqrt{q_\mu}} = 27\delta(n,b) q_{\lambda^h}^{-1/3} q_{N_F n}^{-1/2} q_{B_F b}^{-1}, \quad |\delta| \le 1.$$

The extracted Ramanujan roots leave precisely $(S_0 + B_0)/2$; marked roots give $z_a/2$. The scalar $27/81$ is $1/3$. Shared cube factors remain in b and δ. Consequently the dual coefficient is independent of R, P, while the bounded tuple coefficient is independent of R, n, b. Allowed source primes of S in n, b are retained. □

The freezing condition is necessary for this separation. With primary $p = 1 + 3\omega$, $q = 4 + 3\omega$, the sextic phases are $a = \chi_p(q) = \zeta_6$, $b = \chi_q(p) = \zeta_6^4$. Then $a^4 b^2 = 1$, whereas $a^4 b^{10} = \omega \neq 1$ for a moving $j = 4$ prime. The source already freezes it. Likewise deleting $1_{(P,b)=1}$ changes the local value at $n = 1, b = 0$ from zero to one.

For completeness, the common-profile saving has a separate analytic proof. Let V and all its Euler derivatives decrease faster than every real power at zero and infinity. Put $\widehat{V}(t) = \int_0^\infty V(x) x^t \, dx/x$, $K = (2\pi)^4/27$, and

$$R(t) = \prod_{\pm} \frac{\Gamma(1 + t \pm 1/6)}{\Gamma(1 - t \pm 1/6)}, \qquad V^\sharp(x) = \frac{1}{2\pi i} \int_{(0)} \widehat{V}(-t) R(t) (Kx)^{-t} \, dt.$$

The finite seminorm used below is

$$\mathfrak{M}_{A,B}(V) = \sup_{-A\le\eta\le 1/4} \int_{\mathbb{R}} (1+|u|)^B |\widehat{V}(\eta+iu)|\,du.$$

The gamma quotient is holomorphic for $\Re t > -5/6$ and satisfies $|R(\sigma+iu)| \ll_A (1+|u|)^{4\sigma}$ on $-1/4 \le \sigma \le A$ by [17]. Rapid Mellin decay permits shifts to $-1/4, 0, A$, giving

$$|D_x^j V^\sharp(x)| \ll_{A,j} \min\{x^{1/4}, (1+x)^{-A}\}\mathfrak{M}_{A,\lceil 4A\rceil+j+2}(V).$$

On a fixed logarithmic box, $F_Y(\boldsymbol{y}) = W(\boldsymbol{y})V^\sharp(Y\prod y_i^{a_i})$ has one Fourier density common to every row and tuple. Euler derivatives introduce fixed a_i factors, with no extra Y power. Extract $m_{\mathrm{ker}} = Z^{-(T_d - v - 3\ell_b - e_\lambda)_+/4}$ from that linear vector before Minkowski; its square enters only afterward. Enlarge the positive row set after separation, with no kernel reevaluation on added rows. Nine new Lean components check the local characters, phase algebra and the integrable common-density norm step. The global theta/cusp identification and the hybrid analytic estimate remain separate, unverified inputs; no zero-free or GRH theorem follows.

18 Consequences for a revised research article

Theorems 3.1, 3.2, and 3.3 require replacements for the associated proof steps, not additional exposition. A credible next theorem would construct operators independently of unknown zeros and establish their exact arithmetic determinant, including ramified, archimedean, and conductor terms. It must then prove locally uniform convergence or the local boundedness in Proposition 2.3. Only then does Theorem 2.1 prove the desired zero location.

Weil/Li positivity, adelic trace methods, and canonical-product approximations are potential alternative interfaces. They move the research obligation to a specific positivity, trace, or approximation theorem; they do not discharge it by analogy. Lagarias [4] and Connes [5] provide relevant primary-source starting points. No novelty or complete proof is claimed for these classical conditional principles.

19 Why an absolute coprime pair-square estimate cannot close the short rows

Keep the original sextic family

$$M_u(D) = D^{-1/2}\sum_n \mu(n)\nu(n)W(Nn/D)\chi_n(u), \qquad \chi_n(u) = (u/n)_6,$$

with squarefree primary column ideals outside the fixed excluded set, their full annular coefficients (fix W supported in $[1,2]$), and all nonzero Eisenstein element rows $Nu \le H$. The cubic symbol is χ_n^2, not χ_n. The following classical frame inequality [18] gives a new obstruction to one proposed estimate in this arithmetic program.

Proposition 19.1 (Coprime pair-square obstruction)**.** *Let A be an R-by-N complex matrix with $|A_{ui}| = 1$, and put $K_{ij} = \sum_u A_{ui}\overline{A_{uj}}$. Then*

$$\sum_{i\ne j} |K_{ij}|^2 \ge NR(N-R).$$

For the original character kernels, let $B_1(H,D)$ be the sum of $|K_H(n,m)|^2$ over all distinct coprime squarefree column ideals in $[D,2D]$. At $H=D^{3/4}$,

$$B_1(H,D) \gg \frac{D^{11/4}}{(\log D)^2}.$$

In particular the proposed sufficient input $B_1(H,D) \ll H^2 D^{\varepsilon}$ fails for each fixed $\varepsilon < 5/4$. This does not refute the signed Möbius mean square.

Proof. Set $F_{uv} = \sum_i A_{ui}\overline{A_{vi}}$. Reordering four finite sums proves $\sum_{i,j} |K_{ij}|^2 = \sum_{u,v} |F_{uv}|^2$. The row diagonal contributes RN^2; the column diagonal contributes NR^2. Removing the latter and retaining the former gives the inequality.

Restrict the column pairs in the nonnegative sum B_1 to distinct prime ideals p, q of norm in $[D,2D]$, outside the same fixed set. If $H < D$, no such prime divides a nonzero row element: divisibility would imply $Nu \geq Np$. Thus all original zero masks are exactly one on this ball, $|\chi_p(u)| = 1$, and the residual primitive finite character conductor is pq, of norm between D^2 and $4D^2$. All these pairs have common divisor1. The lattice count gives $R = (2\pi/\sqrt{3})H + O(\sqrt{H}+1)$. The classical unconditional fixed-field prime-ideal theorem gives $N \sim D/\log D$; this is an external analytic input, not a GRH hypothesis. At the stated endpoint $R = o(N)$, and the bound is therefore $(1-o(1))N^2R \gg D^{11/4}/(\log D)^2$. □

To see the exact failed step, write $c_n = \mu(n)\nu(n)W(Nn/D)$ and $C_D = \sum_n |c_n|^2 = O(D)$. Cauchy over coprime pairs bounds the absolute signed contribution by $D^{-1}C_D B_1^{1/2}$. The hoped-for pair-square upper bound would have closed this particular route, but is false. A large absolute pair-square mass is not a lower bound on the quadratic form with the fixed c_n. Nor is the prime part a positive summand of the full Möbius energy: prime–composite cross terms can cancel. The original low-gcd estimate and the positive arithmetic determinant bridge remain open.

A Worked geodesic calculations and possible replacements

For trace $t = 3$, the expanding eigenvalue is $(3+\sqrt{5})/2$ and

$$N = \frac{7+3\sqrt{5}}{2}, \qquad \ell = 2\log\frac{3+\sqrt{5}}{2}.$$

This is a legitimate hyperbolic length, but not $\log p$. For $p = 2$, the proposed length $\log 2$ would require $t = 3/\sqrt{2}$, equivalently $2t^2 = 9$, which is impossible even modulo two. The Lean companion verifies this last exact integer obstruction. For arbitrary p^k, the integer argument in Theorem 3.1 is stronger than a numerical search over traces.

A Hecke operator attached to determinant-p double cosets is a different object from a determinant-one deck transformation. A replacement must specify the ambient space, coefficient system, trace class or regularized trace, and the precise arithmetic distribution. Reusing the same symbol for these objects does not construct an identification. An adelic framework may encode prime norms intrinsically; its trace and positivity obligations must then be proved in that framework.

B Character examples and induction bookkeeping

The nonprincipal character modulo three has values $\chi(1) = 1$, $\chi(2) = -1$, $\chi(0) = 0$ and is odd. For scalar weight-zero automorphy under $\Gamma_0(3)$,

$$f(\gamma z) = \chi(d) f(z),$$

the element $-I$ fixes z and has $\chi(-1) = -1$. Hence every such function would satisfy $f(z) = -f(z)$ and vanish. The proposed scalar bundle on the projective quotient is therefore unavailable as written. One can work on a different group or choose a compatible nonzero weight; either change requires rebuilding the operator and trace identity.

Modulo five, the quadratic character has $\chi(2) = -1$. Consequently the one-prime diagonal entry $\chi(2)2^{-1/2}$ is negative. Its 1×1 minor is already negative, so an unrestricted assertion of total nonnegativity of these transfer matrices is false. A positive state constructed as $TT^*/\operatorname{tr}(TT^*)$ discards phase information in this diagonal model: the nonzero character phases disappear from TT^* at real s, although the ramification zero mask remains. Thus positivity of that state cannot by itself recover character-sensitive Euler factors.

For $C_2 \times C_2$, the three order-two subgroups each admit two linear characters. All six induced characters have degree two; induction from the trivial subgroup gives degree four. Negative integer coefficients do not fix the parity obstruction. Rational coefficients can change parity, which explains why rational and integral induction cannot be interchanged. For a monomial representation $\operatorname{Ind}_H^G \theta$, the appropriate Artin formalism yields a Hecke L-function over the fixed field of H; it does not automatically reduce to a product of Dirichlet functions over $\mathbb{Q}$. Every extension must retain the base field and infinity type.

C Local algebra and topology checks

For the upper-Iwahori order

$$\mathcal{O} = \begin{pmatrix} R & R \\ \pi R & R \end{pmatrix}, \quad S = \begin{pmatrix} 0 & \pi \\ 1 & 0 \end{pmatrix},$$

$E_{12} \in \mathcal{O}$, whereas $SE_{12}S^{-1} = \pi^{-1}E_{21} \notin \mathcal{O}$. The displayed shift orientation therefore does not normalize this order. Changing an orientation may repair this isolated calculation, but all subsequent products and gluing maps must be rechecked with that convention.

For $R = \mathbb{C}[[t]]$ and $\lambda \in \mathbb{C}^\times$, there is no unital $\mathbb{C}$-algebra homomorphism sending t to λI in a nonzero matrix algebra: $t - \lambda$ is a unit in R and would be sent to zero. Thus substituting a unit complex phase for a formal uniformizer is not a representation of this allowed complete DVR. An algebra of Laurent polynomials can admit unitary phase evaluation, but it is a different algebra and needs its own construction.

A dessin on a sphere with two vertices and three edges has three faces, by Euler's formula $V - E + F = 2$, not two. Repeating a vertex along a face boundary does not create a new primitive vertex idempotent. In particular, a three-vertex path algebra with arrows between distinct vertices cannot have the claimed surjection to a six-vertex radical-square-zero cycle algebra: compose with the target's semisimple quotient k^6. Arrows map to zero, since the orthogonal source idempotents force their endpoint supports to be disjoint in the commutative target. The image is generated by at most three orthogonal idempotents, so it cannot be all of k^6. A valid replacement may require a cover or unfolded category rather than a quotient of the stated algebra.

D Why weak convergence and boundary analogies are insufficient

Let P_n be projection onto the nth basis vector of ℓ^2. For every $x \in \ell^2$, $\|P_n x\| = |x_n| \to 0$, so $P_n \to 0$ strongly, but $\|P_n\|_1 = 1$ for every n. Moreover $\det(I - P_n) = 0$ while $\det(I) = 1$. Thus even strong operator convergence does not imply determinant convergence. Trace-norm control is a natural sufficient condition, and must be established for the actual operator family, locally uniformly in s, before invoking Fredholm determinant continuity.

For a Phragmén–Lindelöf argument, one needs uniform control of approximants as the approximation level changes. Polynomial growth of each individual polynomial has constants and degrees depending on N; it does not supply a common bound. Dividing by $(1+|s|^2)^M$ produces a nonholomorphic expression, to which the maximum modulus principle does not apply. Dividing by $(1+s^2)^M$ instead introduces poles at $s=\pm i$; a denominator must be holomorphic and nonvanishing on the actual domain. Even after selecting such a denominator, the necessary uniform boundary and horizontal-side estimates are substantive mathematical work.

Likewise, a bound on a density of geodesics is not a bound on a weighted analytic remainder. The full test-function dependence, prime powers, scattering, elliptic and parabolic contributions must be retained. On a cusped finite-area surface the infimum of the injectivity radius is zero. It cannot provide the positive global mesh bound claimed in the source completion argument.

E Spectral approximation can conceal the original hypothesis

Proposition E.1 (Canonical-product converse for the Riemann target)**.** *If RH holds, the entire order-one function ξ admits locally uniform approximations of the form in Theorem 2.1 with diagonal self-adjoint H_N.*

Proof. The Hadamard factorization of ξ, which has no zero at zero, is

$$\xi(s) = e^{a+bs}\prod_{\rho}(1-s/\rho)e^{s/\rho},$$

with an enumeration for which the canonical product converges locally uniformly. Under RH every $\rho = 1/2 + i\gamma$ has real γ. Each finite product is an exponential of an affine function times a constant times $\prod_{j\le N}(s-\rho_j)$; absorb the nonzero constant into the exponential. Take $H_N = \operatorname{diag}(\gamma_1,\dots,\gamma_N)$. □

Together with Theorem 2.1, this shows why existence of the approximants, without an independent arithmetic construction, may merely restate RH. It is not progress to import the unknown zero locations into the input spectrum and then prove that the constructed spectrum is real. This proposition is a classical consequence, not a novel proof of RH.

F Hermitian dilation and the spectral parameter

A common attempt to preserve arithmetic input while obtaining self-adjointness is the dilation, for square $T(s) \in M_d(\mathbb{C})$,

$$K(s) = \begin{pmatrix} 0 & T(s) \\ T(s)^* & 0 \end{pmatrix}.$$

It is Hermitian pointwise. Singular-value decomposition shows that its spectrum is $\{\pm\sigma_j(T(s)) : 1 \le j \le d\}$, with multiplicities. Consequently

$$\det(\lambda I - K(s)) = \prod_{j=1}^{d}(\lambda^2 - \sigma_j(T(s))^2), \qquad \det(-K(s)) = (-1)^d|\det T(s)|^2.$$

These formulas retain singular values and erase determinant phase. For a nonconstant holomorphic scalar f, $|f|^2$ cannot be holomorphic: a real-valued holomorphic function is constant, and constant modulus then forces f constant. Thus the zero-matching scalar $|f|^2$ is not a holomorphic replacement

for f. Matrix nonholomorphy alone does not imply that every scalar expression formed from it is nonholomorphic; each proposed determinant identity must be examined separately. For example, $T(s) = \operatorname{diag}(e^s, e^{-s})$ is nonconstant but its dilation determinant is constantly one.

For the prime-diagonal model, the dilation eigenvalues are exactly $\pm|\chi(p)|p^{-\operatorname{Re} s}$. Both $\Im s$ and the nonzero character phases disappear. This cannot encode the proposed arithmetic zero ordinates. The source's parameter mismatch is visible where an s-dependent transfer is converted to a Hermitian/modular Hamiltonian and then used as though it had fixed eigenvalues in a characteristic polynomial [1], source lines 11465–11484 and 12200–12215; see also lines 13636–13643 for the explicit skew-Hermitian-part extraction. Its numerical code fixes a particular evaluation point instead, so the rigidity proposition does not itself refute that distinct operation. The earlier diagonal diagnostic obstruction addresses it directly.

A permitted architecture is a fixed self-adjoint H_N with spectral variable entering through $sI - (1/2 + iH_N)$. Alternatively one may use a holomorphic pencil that is Hermitian only on its real parameter axis, but this condition does not force real spectral roots. Already the scalar polynomial $h(t) = t^2 + 1$ is real for real t, whereas $z - h(z) = 0$ has the nonreal solutions $(1 \pm i\sqrt{3})/2$. An infinite-dimensional operator model would additionally need explicit domains, closure, adjoint and determinant regularization. No finite-matrix rigidity statement here is asserted to settle variable-domain unbounded operator constructions.

G Positive pencils, resolvent control, and a Herglotz test

The positive generalized pencil is precisely a fixed self-adjoint construction in different coordinates. Put $H = B^{-1/2}AB^{-1/2} = H^*$. Then

$$A - zB = B^{1/2}(H - zI)B^{1/2},$$

and therefore

$$\det(A + i(s - 1/2)B) = i^d \det(B) \det\bigl(sI - (\tfrac{1}{2}I + iH)\bigr).$$

The positive metric B must be supplied by the arithmetic construction; its existence is not inferred from a desire for real roots.

There is also a quantitative test. If $B \geq bI$ with $b > 0$ and z is nonreal, then for every v,

$$\|(A - zB)v\| \, \|v\| \geq |v^*(A - zB)v| \geq |\Im z| v^*Bv \geq b|\Im z| \, \|v\|^2.$$

For $v \neq 0$, divide by $\|v\|$. In finite dimension this proves invertibility and the operator-norm bound

$$\|(A - zB)^{-1}\| \leq \frac{1}{b|\Im z|} = \frac{1}{b|\operatorname{Re} s - 1/2|}.$$

Uniform approximation arguments must track dimension, normalization and b_N; a bound for each individual pencil does not by itself provide uniform determinant convergence.

A nonlinear extension uses a matrix-valued Herglotz condition: if $M(z)$ is holomorphic upstairs and

$$\Im M(z) := \frac{M(z) - M(z)^*}{2i} > 0,$$

then $M(z)v = 0$ would contradict $v^*\Im M(z)v > 0$. Thus $\det M(z) \neq 0$ there. A reflected lower-half-plane extension $M(\bar{z}) = M(z)^*$ is likewise invertible. A quantitative lower bound $\Im M(z) \geq \eta(z)I > 0$ gives $\|M(z)^{-1}\| \leq 1/\eta(z)$ by the same argument. For the positive linear pencil the Herglotz sign is $M(z) = zB - A$, not $A - zB$. Semidefinite positivity alone allows $M(z) = \operatorname{diag}(z, 0)$, which is always

singular. Without reflection, $M(z) = z + i$ is strictly Herglotz upstairs but has a lower-half-plane zero. Possible real-axis poles and the arithmetic determinant identity remain separate obligations.

For a candidate normalized identity written in the z-coordinate, $F_N(z)/g_N(z) = \det(A_N - zB_N)$ with a prescribed nowhere-zero g_N, a necessary diagnostic, for $d_N > 0$, is

$$-\partial_z \log(F_N/g_N) = \operatorname{tr}\big((A_N - zB_N)^{-1}B_N\big) = \sum_{j=1}^{d_N} \frac{1}{\lambda_{j,N} - z}.$$

Its imaginary part is strictly positive for $\Im z > 0$. A certified violation rejects that particular determinant identity; finitely many successful samples cannot prove it. One must not choose g_N using unknown zero locations or assume this Herglotz positivity as a substitute for establishing the arithmetic bridge.

H Quantitative arithmetic approximation and prime regularization

For the nonprincipal-character approximants, a compact set $K \subset \{\operatorname{Re} s > 0\}$ has $\delta = \min_K \operatorname{Re} s > 0$, $M = \max_K |s|$, and $G = \max_K |(q/\pi)^{(s+a)/2}\Gamma((s+a)/2)| < \infty$. The completed approximation error is at most $GB(1 + M/\delta)N^{-\delta}$ uniformly on K. This is a fixed-character compact bound, not a uniform-in-conductor GRH estimate. To verify the Abel formula's sign, first sum between integers $N < T$:

$$\sum_{N<n\le T} \chi(n)n^{-s} = A(T)T^{-s} - A(N)N^{-s} + s\int_N^T A(x)x^{-s-1}\,dx.$$

Then $A(T)T^{-s} \to 0$. Bounding $|A|$ by B gives exactly the main-text estimate. For zeta, $\lfloor x \rfloor = x - \{x\}$ in the same summation formula yields

$$\zeta(s) = \frac{s}{s-1} - s\int_1^\infty \{x\}x^{-s-1}\,dx.$$

The finite sum formula gives the stated remainder after subtracting Z_N. The integral and its derivatives converge uniformly on compact subsets of the positive half-plane, since logarithmic powers are integrable against $x^{-1-\delta}$. This justifies holomorphy and continuation. Multiplying the remainder by $s(s-1)\pi^{-s/2}\Gamma(s/2)/2$ gives the compact-uniform ξ_N error, including at $s = 1$. In contrast, multiplying a finite Euler product by $s(s-1)$ would force a zero at 1 for every truncation, although $\xi(1) = 1/2$.

A second exact interface isolates what a prime-based repair would need. For a nonprincipal primitive χ, define on $\operatorname{Re} s > 1$

$$P_\chi(s) = \sum_p \chi(p)p^{-s}, \qquad R_\chi(s) = \sum_p \sum_{k\ge 2} \frac{\chi(p)^k}{kp^{ks}}.$$

On each compact subset of $\operatorname{Re} s > 1/2$, the second series converges absolutely and uniformly. Indeed, for $\operatorname{Re} s \ge \delta > 1/2$,

$$\sum_{k\ge 2} \frac{|\chi(p)|^k}{kp^{k\delta}} \le \frac{p^{-2\delta}}{1 - 2^{-\delta}},$$

and the prime sum is dominated by the convergent integer series $\sum_{n\ge 2} n^{-2\delta}$. Thus R_χ is holomorphic there. On $\operatorname{Re} s > 1$, $L(s,\chi) = \exp(P_\chi(s) + R_\chi(s))$.

Proposition H.1 (The missing prime-linear term)**.** *For a nonprincipal primitive character, P_χ has a single-valued holomorphic continuation to $\operatorname{Re} s > 1/2$ if and only if $L(s,\chi)$ is zero-free there.*

Proof. If such a continuation exists, its exponential with R_χ agrees with L on $\operatorname{Re} s > 1$ and hence on the connected half-plane by the identity theorem. The exponential is nonzero. Conversely, if L is nonzero, the simply connected half-plane admits a holomorphic logarithm of L. Choose its additive constant to agree with the absolutely convergent Euler logarithm on $\operatorname{Re} s > 1$. Subtract R_χ to obtain the required continuation of P_χ. □

This proposition isolates a precise analytic obligation; it does not prove it. Regularizing away the terms $k \geq 2$ supplies no proof that the missing prime-linear function extends. The principal zeta case must first remove the pole, for example by using $(s-1)\zeta(s)$; an unqualified logarithm of ζ cannot be holomorphic through 1. The functional equation relates a fixed character's zeros across the critical line after complex conjugation, so a full right-half-plane nonvanishing theorem would give the desired Dirichlet GRH conclusion.

I Exact arithmetic behind the finite-level zero certificate

This section gives all parameters needed to reproduce Theorem 7.1. Write $C = |\operatorname{Re} c| + |\Im c| < 12$. Use the following rational logarithm approximations ℓ_n and error bounds δ_n:

n	ℓ_n	δ_n
2	$6931471805599453/10^{16}$	10^{-15}
4	$13862943611198906/10^{16}$	$2 \cdot 10^{-15}$
5	$16094379124341004/10^{16}$	10^{-15}

For $y = (n-1)/(n+1)$, set

$$L_n = 2\sum_{k=0}^{59} \frac{y^{2k+1}}{2k+1}, \qquad T_n = \frac{2y^{121}}{121(1-y^2)}.$$

The positive series for $\log n$ gives $L_n \leq \log n \leq L_n + T_n$. Exact rational comparisons give $\ell_n - \delta_n \leq L_n$ and $L_n + T_n \leq \ell_n + \delta_n$. They also give $0 < \log 2 < 1$ and $0 < \log 4, \log 5 < 2$.

Define Gaussian rationals

$$E_n = \sum_{k=0}^{120} \frac{(-c\ell_n)^k}{k!}, \qquad \tau = \frac{3^{24}24^{121}}{121!} < 10^{-18}, \qquad \eta_n = \tau + 2C\delta_n.$$

Since $|c\ell_n| \leq C\ell_n < 24$, the exponential series tail is at most $e^{24}24^{121}/121! < \tau$. Here $e < 3$ follows directly from its series. Also

$$|e^{-c\log n} - e^{-c\ell_n}| \leq e^{-\operatorname{Re}(c)\ell_n}|c|\delta_n e^{|c|\delta_n} \leq 2C\delta_n,$$

because $\ell_n, \operatorname{Re} c > 0$, $C\delta_n \leq 1/2$, and $e^u \leq (1-u)^{-1} \leq 2$ for $0 \leq u \leq 1/2$. Thus $|n^{-c} - E_n| \leq \eta_n$.

Put

$$F = 1 - E_2 + E_4 - E_5, \qquad D = \ell_2 E_2 - \ell_4 E_4 + \ell_5 E_5.$$

All entries are exact Gaussian rationals. With $\|(a,b)\|_1 = |a| + |b|$, the script checks the rational inequalities

$$\|F\|_1 + \eta_2 + \eta_4 + \eta_5 < 10^{-6}, \qquad |\operatorname{Re} D| - \sum_{n\in\{2,4,5\}} (\delta_n + \ell_n\eta_n) > 1.$$

The first bounds $|f(c)|$. For the second, note that $|n^{-c}| \le 1$ and

$$|(\log n)n^{-c} - \ell_n E_n| \le \delta_n + \ell_n \eta_n.$$

This proves $|f'(c)| > 1$. On the entire closed disk, $\operatorname{Re} s > 0$, so

$$|f''(s)| \le (\log 2)^2 + (\log 4)^2 + (\log 5)^2 < 1 + 4 + 4 = 9.$$

Integrating the second derivative along the line segment from c to s justifies the Taylor remainder bound $9|s-c|^2/2$. The exact strict boundary margin is

$$r - 10^{-6} - \tfrac{9}{2}r^2 = \frac{19791}{200000000} > 0.$$

Consequently $|f(s)|$ is greater than that margin on $|s-c| = r$. Any zero-free holomorphic positive-pencil model G on a neighborhood of this disk must have boundary error at least $\min_{|s-c|=r} |f(s)|$; otherwise another application of Rouché would contradict its zero count. This is a quantitative rejection of a prescribed finite-level model, not a proof about the zero locations of the limiting L-function.

The certificate program `scripts/check_arithmetic_zero.py` uses only Python integer and Fraction arithmetic. A separate independent exact-arithmetic reproduction passed. Lean verifies the disk's location and the last rational margin; the logarithm/exponential error estimates and the Rouché inference remain explicit prose proofs, not claims of a complete kernel-checked zero-existence theorem.

J Jensen companions: exact normalization and the failed moment argument

Our γ_j equals one eighth of the convention in equation 1 of [7], whose generating function is $8\xi(1/2+z)$. This common positive factor changes no polynomial zeros. Entirety of F follows directly from the even Taylor series: its absolute series at $|w| = R$ is the absolute series for $\xi(1/2+z)$ at $|z| = \sqrt{R}$. The finite-product inequality in the main-text convergence proof is also verified in Lean. The analytic summation and ξ specialization remain prose arguments, using classical properties of ξ. For an explicit coefficient interface, the classical representation in [9, equations 1.8–1.11] gives

$$\Phi(u) = 2\sum_{n\ge 1}(2\pi^2 n^4 e^{9u/2} - 3\pi n^2 e^{5u/2})e^{-\pi n^2 e^{2u}}, \qquad \gamma_j = \frac{j!}{(2j)!}\int_{\mathbb{R}} u^{2j}\Phi(u)\,du.$$

The kernel is even. Each summand is positive for $u \ge 0$ because $2\pi n^2 e^{2u} > 3$, so the moments are strictly positive. Its rapid decay permits differentiation of the Fourier integral. This arithmetic kernel representation is an external classical input, not a new Lean theorem in this project.

J.1 The coefficient construction

Write $p(x) = J_d(x)/\gamma_d = x^d + \sum_{i<d} a_i x^i$. The companion C has $C_{j+1,j} = 1$ for $0 \le j < d-1$, last column $C_{i,d-1} = -a_i$, and all other entries zero. For $1 \le k \le d$ Newton's recurrence is

$$m_k + \sum_{i=1}^{k-1} a_{d-i} m_{k-i} + k a_{d-k} = 0,$$

and for $k > d$ it is

$$m_k + \sum_{i=1}^{d} a_{d-i} m_{k-i} = 0.$$

Together with $m_0 = d$, this determines the entire required matrix $H = (m_{i+j})_{0 \le i,j < d}$. Multiplication in the quotient algebra $\mathbb{R}[x]/(p)$ gives its bilinear form

$$h(u, v) = \operatorname{tr}(M_{uv}), \qquad h(xu, v) = h(u, xv),$$

so $HC = C^T H$ without any zero-location assumption. Here M_{uv} means multiplication by uv in that finite quotient. When roots are real and distinct, the evaluation matrix V gives $H = V^T V > 0$. Repeated roots make V singular; the semidefinite Hermite criterion still applies, but inverse square roots do not. Nonnegative leading principal minors alone are not a PSD certificate; all principal minors, or a valid alternative test, are required.

Assume now $H > 0$. Then

$$S = H^{1/2}(-C)H^{-1/2} = S^T > 0, \qquad K = \begin{pmatrix} 0 & \sqrt{dS} \\ \sqrt{dS} & 0 \end{pmatrix} = K^T.$$

Indeed the similarity is symmetric by $HC = C^T H$, and its eigenvalues are negatives of the roots of p, all positive. The block determinant and companion characteristic polynomial give

$$\det(tI_{2d} - K) = \det(t^2 I_d - dS) = \det(t^2 I_d + dC),$$
$$J_d(-t^2/d) = \gamma_d(-1/d)^d \det(t^2 I_d + dC).$$

The identity holds also at $t = 0$, by polynomial equality. Using a companion for the scaled Q_d instead would remove the extra d inside the square root. With our stated convention it cannot be omitted. Thus the arithmetic data and approximation rate are specified; the unproved universal positivity is precisely where the RH difficulty lies. This construction supplies no general Dirichlet, Hecke, Dedekind or Artin conclusion. Their symmetry and coefficient interfaces must be established separately.

J.2 Why ordinary moment positivity fails

For $p = x^2 + bx + a$, the companion and Newton-sum metric are

$$C = \begin{pmatrix} 0 & -a \\ 1 & -b \end{pmatrix}, \qquad H = \begin{pmatrix} 2 & -b \\ -b & b^2 - 2a \end{pmatrix}.$$

Direct calculation gives $HC = (HC)^T$, $\det H = b^2 - 4a$, and

$$u^T H u = 2(u_0 - bu_1/2)^2 + \tfrac{1}{2}(b^2 - 4a)u_1^2.$$

These identities and positivity under $b^2 - 4a > 0$ are formalized in Lean. For $J_2 = \gamma_0 + 2\gamma_1 x + \gamma_2 x^2$, $\gamma_2 > 0$,

$$\det H = \frac{4(\gamma_1^2 - \gamma_0\gamma_2)}{\gamma_2^2}.$$

The unsmoothed atomic example has $\mu_0 = 1$, $\mu_2 = \mu_4 = 1/10$. Its ordinary even-moment Hankel matrix and Jensen metric are respectively

$$\begin{pmatrix} 1 & 1/10 \\ 1/10 & 1/10 \end{pmatrix} > 0, \qquad \begin{pmatrix} 2 & -12 \\ -12 & -96 \end{pmatrix}, \quad \det H = -336.$$

Gaussian smoothing gives the strictly positive densities in the main-text counterexample. The standard Gaussian moments yield $\mu_2 = v+1/10$ and $\mu_4 = 3v^2+3v/5+1/10$. With $\gamma_n = n!\mu_{2n}/(2n)!$, the negative gap is independent of v. Lean checks that exact algebraic cancellation; the measure integral and Fourier-transform identities remain prose. Thus even smooth positive kernel information cannot replace arithmetic-specific Jensen inequalities.

The exact program `scripts/check_jensen_metric.py` verifies the Newton recurrence, companion symmetry, positive LDL pivots for the $\cosh\sqrt{w}$ model at degrees 1 through 8, and the negative and singular witnesses. Its finite coefficient-bound checks supplement the universal proof; they do not replace it. No arithmetic ξ coefficient intervals or universal Jensen positivity are certified by this program. For example the sequence $(1,1,1)$ gives $J_2 = (1+x)^2$ and $H = \left(\begin{smallmatrix} 2 & -2 \\ -2 & 2 \end{smallmatrix}\right)$, which is PSD and singular. Adding an arbitrary positive diagonal to H need not preserve $HC = C^T H$.

K Euler-correction estimates and the shifted angular character

The immutable scalar input is the September manuscript's local identity, lines 3845–4206 at the pinned source in the project manifest. We verify its rational consequences, without claiming a new verification of the underlying Gauss identities. Write $Q = Np \geq 2$ and retain its notation $V = Q^{-6z}$, $R = a_p^2 Q^{4-6x-6z}$, $W = \chi_p(u)Q^{-w}$, $D = \eta(p)\overline{\chi_p(u)}Q^{-x}$. The unit phases have norm one. At $p \mid u$, $D = W = 0$ by zero extension, while the local unit phase in P_p^* is retained.

K.1 All valuation cases in the larger region

For $p \nmid u$, let $\mathcal{E} = P_p^* + D$. The $j = 0$ formula gives

$$\mathcal{E} = \frac{R((1-Q^{-1})/(1-V) + W - D) - DW(Q-1)V/(1-V)}{1-R}.$$

In the region of the main-text proposition, the exponents of $R, QDWV, DV, DW, DVW, VW$ are respectively at most

$$-\frac{149}{100},\ -\frac{169}{100},\ -\frac{174}{100},\ -\frac{17}{10},\ -\frac{269}{100},\ -\frac{194}{100}.$$

Each of $|R|, |V|, |D|, |W|$ is below $3/4$, so the inverses of the displayed $1-$factor denominators are at most 4. Consequently $|\mathcal{E}| \leq 40Q^{-149/100}$. The exact defect identity

$$H_p - 1 = \frac{D(V + W - VW) - VW + (1-V)(1-W)\mathcal{E}}{1-D}$$

then gives $|H_p - 1| \leq 656Q^{-149/100} < 1024Q^{-149/100}$.

For $p \mid u$, $1 \leq j = v_p(u) \leq 5$, cancellation gives

$$H_p - 1 = \frac{R(1-Q^{-1}) - \eta(p)(Q-1)Q^{-x-w}V^{1_{j\leq 1}} + (1-V)J_j}{1-R}.$$

The middle term has exponent at most $-7/10$. The J_j exponents are

j	1	2	3	4	5
exponents	$-17/10$	$-3/4$	$-6/5, -1$	$-29/20$	$-3/2.$

There are at most two such terms. Thus $|H_p - 1| \leq 24Q^{-7/10}$, and the stated constant 32 is safe. These calculations include every row valuation; primes in the excluded set S have no factors here. The larger open region

$$\operatorname{Re} x > 74/100, \quad \operatorname{Re} w > 94/100, \quad \operatorname{Re} z > 31/200$$

retains respective exponents $-137/100$ and $-68/100$ and the same denominator control. This proves convergence on neighborhoods, including the boundary of the stated closed region. For any $\epsilon > 0$, the finite row product is $O_\epsilon(q_u^\epsilon)$ since, outside a fixed finite set, $1 + 32Q^{-68/100} \le Q^\epsilon$ on that open neighborhood.

The source defines its full compensated correction as a finite sum over selected prime tuples. Each selected local replacement G_p has only $1 - V, 1 - R, 1 - D$ denominators, and its unselected factors are the H_p just bounded. That formula therefore also extends holomorphically to this neighborhood for each fixed Z. No quantitative principal-error estimate for the compensated expression follows merely from holomorphy.

K.2 Principal factor, tails and nonvanishing

At $u = 1, w = 1, z = 1/6$, the source's P_p^* becomes $(1 + v)(r - d)/(1 - r)$ with the main-text definitions. This proves the displayed principal formula. Its numerator cannot vanish on $\operatorname{Re} s > 1/2$: $0 < v < 1$, $|d| < 1$, $|r| < 1$ and the strict convexity estimate apply. To check convergence, direct subtraction gives

$$H_p - 1 = \frac{-v^2 + r(1 - v + v^3) + d(v + v^2 - v^3) - rd}{(1 - r)(1 - d)}.$$

For $\sigma = \operatorname{Re} s > 2/3$, $|r| < 1/2$ and $|d| < 2/3$. The numerator is at most $v^2 + 2|r| + 2v|d|$, while the denominator reciprocal is less than 6. This gives

$$|H_p - 1| \le 12(Q^{-2} + Q^{-\sigma-1} + Q^{3-6\sigma}).$$

For completeness, the Eisenstein norm satisfies $a^2 - ab + b^2 \ge (a^2 + b^2)/2$. The norm-Euclidean ring is a PID; counting generators in $|a|, |b| \le 2\sqrt{X}$ bounds the number of ideals of norm at most $X \ge 1$ by $25X$. Dyadic decomposition consequently gives

$$\sum_{Np > P_0} (Np)^{-1-\delta} \le \frac{50}{1 - 2^{-\delta}} P_0^{-\delta} \qquad (\delta > 0).$$

For a fixed $\sigma_0 > 2/3$, use $\delta = \min(1, \sigma_0, 6\sigma_0 - 4) > 0$. Choosing P_0 so that the local defect sum is at most $1/4$ gives

$$|H_\eta(s) - 1| \le e^{1/4} - 1 < 1/3 \quad (\operatorname{Re} s \ge \sigma_0).$$

This choice is uniform in phases for that fixed half-plane. Its constants degenerate as $\sigma_0 \downarrow 2/3$; no uniform bound on the whole open half-plane is being inferred.

K.3 What the additional factor permits and prevents

Removing $1 - r$ leaves

$$A_p - 1 = -v^2 + \frac{(1 - v^2)v(d - r)}{1 - d}.$$

On every fixed $\operatorname{Re} s \ge \sigma_0 > 1/2$, this is $O_{\sigma_0}(Q^{-2} + Q^{-\sigma_0-1} + Q^{2-6\sigma_0})$. The product A is thus normally convergent and nonzero there, by local nonvanishing and summable defects. The remaining Euler factors are exactly those of $L_F^S(6s - 3, \theta)$ with $\theta = \alpha^{-6}\eta^6$. Units have sixth power one, so the angular character is well-defined on ideals; a finite-order twist cannot cancel its nonzero angular type.

The stronger obstruction uses the following external classical inputs: the primitive completion Λ_θ is entire of order one, has gamma factor $\Gamma(s + 3)$ times a nonzero exponential, obeys $\Lambda_\theta(s) =$

$\epsilon_\theta \Lambda_{\bar\theta}(1-s)$, and $\Lambda_{\bar\theta}(\bar s) = \overline{\Lambda_\theta(s)}$. The nonvanishing Euler product holds on $\operatorname{Re} s > 1$. These are Hecke analytic theory, not GRH or Artin holomorphy assumptions; none is verified by the new Lean file. Tate's continuation/functional equation framework is [10, sections 4.4–4.5]. White's Theorem 2 [11] records the pure angular case, including the gamma shift; it is not a citation for arbitrary finite-conductor twists or a substitute for the stated order-one input.

Hadamard factorization with finitely many zeros would give $\Lambda_\theta(s) = e^{as+b}P(s)$. Along the positive real axis this has logarithmic modulus $\operatorname{Re}(a)\sigma + O(\log\sigma)$, whereas Stirling and $L(\sigma,\theta) \to 1$ give $\sigma\log\sigma + O(\sigma)$. Thus there are infinitely many zeros. Euler nonvanishing and the functional equation confine them to $0 \le \operatorname{Re}\rho \le 1$. Conjugation and the functional equation reflect a zero to $1-\bar\rho$ for the same character, so some zero has $\operatorname{Re}\rho \ge 1/2$. Gamma is finite and nonzero there, as are the deleted Euler factors; hence $L_F^S(\rho,\theta) = 0$. The unique continuation $A(s)L_F^S(6s-3,\theta)$ then vanishes at $(\rho+3)/6$. Enlarging finite S cannot remove that zero. This proves the stated 7/12 obstruction under the explicit classical data; a replacement normalization/probe would need new analytic estimates.

The exact program `scripts/check_euler_correction.py` checks all exponents and evaluates the identities at primary prime representatives $-2-3\omega$ and $1-3\omega$, of norms 7 and 13, using only $\mathbb{Q}(\sqrt{-3})$ arithmetic. Lean checks scalar cancellation, convex local nonvanishing, genuine complex infinite-product components under explicit summability hypotheses, and shifted-zero transport. The arithmetic convergence estimates, source Gauss identification, multivariable holomorphy and classical Hecke completion theory remain prose/external dependencies. This does not establish the missing compensated contour bounds, mean-square estimates or GRH.

L Principal multiplier, weighted slots and the extended residue comparison

The source passages are lines 6173–6328, 7004–7044, 9050–9097 and 15548–15669 of the pinned September paper. We retain its displayed scalar/high identities and Mellin/reciprocal contracts as stated inputs.

L.1 Local errors and the scalar normalizer

Its exact cancellation for $G_p + H_p$ consists of a bounded prefactor times

$$(1-Q^{-w})\{V/(1-V) - D\} + (\overline{\eta(p)}Q^x + 1 - Q^{-w})\mathcal{E}_p.$$

The first term costs $Q^{-6z_r} + Q^{-x_r}$. Multiplying the previous residual bound by Q^{x_r} costs $Q^{4-5x_r-6z_r} + Q^{1-w_r-6z_r}$; unamplified residual terms are smaller. Division by the uniformly nonzero H_p therefore gives

$$|\mathcal{B}_p + 1| \ll Q^{-\kappa}, \qquad \kappa = \min(x_r, 6z_r, 5x_r + 6z_r - 4, w_r + 6z_r - 1) \ge 37/50.$$

It follows that $G_p = O(1)$ on the enlarged principal region. On a fixed real box, an annular slot contributes at most $O(P_i^{z_r})$ by ideal counting, since its weight is Q^{z-1}. The unselected product has a bounded positive majorant. Thus the full tuple is $O(Z^{\ell z_r})$ uniformly in heights, as required. This is not a constant uniform over unbounded z_r.

For the prime normalizer, the fixed-extension consequence of [12, Theorem 1.1] gives $\pi_{1_T}(x) \sim \operatorname{Li}(x)/|T|$. Any exceptional zero is fixed below 1, so its secondary term is asymptotically smaller; it need not be ruled out. Stieltjes integration against the fixed annular weight gives

$$S_i(Z) \sim \frac{P_i^{1/6}}{|T|\log P_i}\int_0^\infty W_i(y)y^{-5/6}\,dy > 0.$$

A further fixed finite deletion eventually leaves the window entirely. The normalizer consequently has the signed asymptotic

$$A_T(Z) \sim \frac{c_T}{(\log Z)^K}, \qquad c_T = \frac{(-1)^K}{|T|^K} \prod_i \left(\ell_i^{-1} \int_0^\infty W_i(y) y^{-5/6}\, dy \right) \neq 0.$$

This proves the inverse subpower bound. For example, after setting $u = \log Z$, the exponential series shows that $e^{\epsilon u}$ eventually dominates the fixed power u^K. Thresholds may depend on the fixed ray and excluded set; the power-saving exponents do not.

At the residue write $a_p = W_i(Q/P_i)Q^{-5/6} \geq 0$ and $\varepsilon_p = \mathcal{B}_p + 1$. Then

$$\sum_p a_p \mathcal{B}_p = -S_i(1+\rho_i), \qquad \rho_i = -\frac{\sum_p a_p \varepsilon_p}{S_i}, \qquad |\rho_i| \ll P_i^{-37/50}.$$

The sign of ρ_i matters. Nonnegative individual weights permit division by their positive total without cancellation. A merely positive sum of signed weights would not give this inequality. For a fixed finite set of K errors bounded by $\delta \leq 1$,

$$\left| \prod_i (1+\rho_i) - 1 \right| \leq (2^K - 1)\delta.$$

Induct by writing $(1+\rho_i)P - 1 = (1+\rho_i)(P-1) + \rho_i$ and using $|1+\rho_i| \leq 2$. Thus the displayed residue identity holds with $\mu = (7/10)\ell_{\min}$, where $\ell_{\min} > 0$ is fixed before the target. The same signed A_T is used in every normalization of this probe.

L.2 Contour powers and the perturbed geometry

Use the inherited exact high representation for the same physical probe, the rapid integrated and trace tail contracts for the Mellin tests, and polynomial reciprocal control to the right of β_*. For $3/4 \leq \sigma_0 < \beta_*$, the principal paths can be moved to

$$\operatorname{Re} s = \beta_* + e, \quad \operatorname{Re} w = 1 + e, \quad \operatorname{Re} z = 1/6 + e$$

inside the enlarged domain. Move w left to $19/20$; in its residue move z left to $33/200$. Only the two scalar zeta poles are crossed. The reciprocal remains on the global line; at a principal pole its zero causes no additional residue. Horizontal joins use the trace estimates in the independent height coordinates $(\Im(s+z), \Im z, \Im w)$. After the w residue, use the corresponding two-dimensional kernel. Arithmetic height orders are fixed first; later external tail orders increase only the decay requested from the tests.

The raw exponent is

$$l_x/2 + \operatorname{Re} s - 1 + h \operatorname{Re} z + l_y(\operatorname{Re} w - 1), \qquad h = 1 - l_x + \ell.$$

On the unresidued w integral it is $C(\beta_*) + (1+h)e - l_y/20$; on the leftover z integral it is $C(\beta_*) + e - h/600$. At the double residue the normalization gives the main signal times $1 + \mathcal{R}_{\eta,Z}(s)$. Its error costs $Z^{C(\beta_*)+e-\mu+\epsilon}$ on the entire global line, using the Gaussian and reciprocal bound. Only the main signal moves right to $\operatorname{Re} s = 2$; no continuation of the relative error is used. The factor $1/6$ is already in c_S from the residue of $\zeta_F^S(6z)$. The normalizer's inverse is absorbed in the allocated ϵ. This proves the three remainder powers stated in the main text, conditional on the explicit inherited analytic contracts.

For $t = 10^{-5}$, the perturbed parameters are

$$\ell = 1/6 + t, \quad l_x = 17/48 - t/2, \quad l_y = 23/48 - t/2, \quad h = 13/16 + 3t/2.$$

They have $C(s) = s - 11/16$ and $\sigma_t = 349999/400000 > 3/4$. Put $m_w = l_y/20$, $m_z = h/600$, and choose

$$0 < e = \epsilon \le \min\left(\frac{1}{10000}, \frac{m_w}{8(1+h)}, \frac{m_z}{8}, \frac{\mu}{8}\right).$$

Each total error cost is at most one quarter of its corresponding margin, so all three remainders have a common positive saving fixed before the target. Nonprincipal contour, energy and frozen-capacity row-count extensions remain separate obligations.

L.3 A pointwise obstruction at $3/5$

At the principal residue put $N = 1 - vr - (1-v)d$. In addition to the main-text multiplier identity, exact algebra gives

$$\frac{\mathcal{B}_p + 1}{r/d} - 1 = \frac{-v(1-r) + (1-v)^2 d(1 - d/r)}{N}.$$

For fixed $1/2 < \sigma \le 3/5$ and $Q \ge 4$, $|N| \ge 1/4$, $|r| < 1$ and $|d| < 1/2$. Hence the relative error is at most

$$8Q^{-1} + 4Q^{-\sigma} + 4Q^{4\sigma-3} \longrightarrow 0.$$

Since $|r/d| = Q^{3-5\sigma}$, a pointwise negative-power bound around -1 fails in this range, uniformly in phases and heights.

For an exact arithmetic endpoint example, let $\eta = 1$ and let the rational prime $p \equiv 2 \pmod 3$ be inert. Its prime ideal has norm $Q = p^2$ and primary generator $-p$, so $\alpha^{-6} = 1$. At real $s = 3/5$, $r = Q^{3-18/5} = Q^{-3/5} = d$. Thus $P_p^* = 0$ and $\mathcal{B}_p = -Q^{-1}$. Given a fixed ray modulus $\mathfrak{m}$, choose a positive integer M with $M\mathcal{O} \subset \mathfrak{m}$. Primes $p \equiv -1 \pmod{3M}$ have primary generator $-p \equiv 1 \pmod{\mathfrak{m}}$ and occur infinitely often by the classical fixed-progression prime theorem. They obey the identity-ray restriction. Nevertheless their ideals have count $O(\sqrt{P})$ up to norm P, a zero-density subset of the prime ideals. This supplies no exclusion of cancellation in the full slot average.

The exact checker `scripts/check_principal_residue.py` verifies the multiplier and slot identities, rational gaps and illustrative principal loss allocations. Its slot counts do not certify row-count admissibility. The six Lean statements verify scalar identities, actual complex weighted normalization and finite-product error transfer. Prime counting, the arithmetic endpoint identification and all contour integrations remain prose/external inputs. No new zero-free half-plane follows from this principal interface alone.

M Research tasks with mathematical acceptance criteria

Track A: repair the determinant interface. Start with the exact diagonal Euler transfer on $\operatorname{Re} s > 1$. Determine which regularized operator determinant could continue the reciprocal Euler product without poles or zeros being silently exchanged. Prove a relation to a self-adjoint operator before replacing the transfer by its Hermitian part. Acceptance requires an identity of functions, not similarity of zero plots. The displayed 2×2 example rejects unrestricted determinant preservation immediately.

Track B: trace norms and normal families. Choose a connected domain containing an Euler-product overlap and the critical strip, after removing the known poles in the target. Supply estimates uniform in N on every compact set. Either construct a trace-class holomorphic family with uniform trace-norm convergence or prove the hypotheses of Proposition 2.3. The divergence in Theorem 3.3 prevents an absolute prime-sum shortcut. Any cancellation estimate must be independently proved without a zero-free region equivalent to the desired conclusion.

Track C: positivity and canonical products. Compare the exact spectral quadratic form with Weil positivity and the Li coefficients for the chosen L-function family. First state the test-function space and regularization, then prove equality of the forms. Positivity of a finite matrix or of finitely many Li coefficients is insufficient; a uniform infinite-family statement is required. Canonical products provide a convergence benchmark, but the preceding converse explains the circularity risk of using zeros as input.

Track D: make formalization match mathematics. The standalone Lean companion checks exact integer and finite-matrix obstructions. A separate mathlib layer now verifies complex Euler-factor nonvanishing, a local boundary-convergence theorem, and holomorphic Hermitian rigidity. It does not yet formalize the Dirichlet partial-sum convergence or an arithmetic self-adjoint operator. Further formalization must use the actual standard L-functions and explicit analytic domains. An abstract theorem of the form "if an assumed bridge implies GRH, then GRH" is logically valid but would not verify the bridge. Print every theorem's axioms and review its full type and dependencies, not merely its name.

N Details of the frozen row-count extension

We give the selection and uniformity details of Proposition 11.1. The immutable source locations and full dependency list are recorded in `research/frozen-row-count.md`. This proof uses the stated source moments and witnesses as hypotheses; it does not establish them. Given the requested exponent loss, first choose capacity decrements and a moment/rounding mesh threshold independent of the eventual slot count; then fix the finite slot system and form the bins. Only afterward fix its finite internal profile and height orders. An arbitrary loss is not asserted for a fixed coarse mesh.

Spikes and selection. After subdivision by a fixed finite character presentation and a dyadic pair, the witnesses have common row character, smooth profiles and lengths r, m. The number of subdivisions is $O((\log U)^2)$. At base U, slot i has length $w_i = \ell_i/d$ and amplitude $g_i \in [0, \delta/2]$, with $q = (\sum_i w_i g_i)/(\sum_i w_i)$. Only positive amplitudes provide lower bounds for the corresponding physical slot. Fill requested length z with the largest positive amplitudes first. The fractional selection has gain at least qz: its initial weighted average is at least the total mean; after the positive slots are exhausted their full gain is $q\ell/d \geq qz$. Dropping the one possibly fractional slot gives

$$\left| \prod_{i \text{ selected}} Q_i \right|^2 \geq U^{2qz - \delta \max_i w_i}.$$

A fixed capacity decrement ν_0 adds a loss at most $2q\nu_0$. Since $d \geq d_{\min}$, a sufficiently fine fixed physical mesh makes both losses as small as required. Error slots with $g_i = 0$ do not acquire fictitious lower bounds.

The inverse moment, divided by the two spikes, bounds the row count by $U^{1-\delta r-2qz+\epsilon}(1+T_1)^A$. For the plain moment use two copies of S_m, so division is by $|S_m|^4$, giving exponent $1-2\delta m-2qz+\epsilon$.

A formal component verifies this finite-moment step for actual complex values: if $A_0 \le |w_u|^2$, $B_0 \le |p_u|^2$, $B_0 \ge 0$ and $\sum_{u\in\mathcal{B}} |w_u p_u|^2 \le M_0$, then $\#\mathcal{B}\, A_0 B_0 \le M_0$. It supplies no arithmetic moment bound.

Arithmetic presentations. With negative common orientation write the row as $\nu(n)\overline{\chi_n(u)}$, $\nu \in \Theta$. The slot coefficient is $\overline{\nu(p)}1_{p\in 1_T}$, a fixed finite expansion in Θ, independent of the moving row in the current sum. For the plain moment conjugate the entire product, both witness profiles and all selected slots. Its positive orientation has coefficients $\nu(p)1_{p\in 1_T}$, still in the allowed class. Conjugation preserves zero extensions and underlying disjoint supports. The same admissible fixed-within-the-sum mask is retained in both plain copies and every selected slot. A moving mask is never absorbed into the fixed arithmetic datum. Large physical sixth-power-free rows induce outside Θ by ramification outside S; exceptional physical rows supported on S remain a separate finite family. These family properties are indispensable inherited hypotheses.

Crossing and strict margins. The frozen capacity satisfies

$$2m + 6(3/4)\frac{1-2m}{9/2} = 1.$$

The plain exponent decreases with m, with derivative $-2\delta + 8q/9 < 0$, so $m \ge \tau - r - o(1)$ gives $S_\tau(r)$. Both affine counts have opposite slopes. Their crossing satisfies

$$r_*(3/2) = 1, \quad r_*(1) \ge 23/37, \quad \tau - r_*(\tau) = \frac{(1-x)\tau + 5x/9}{D} \in [1/3, 1/2].$$

Thus the chosen inverse capacity is at most $7/37$ and the plain capacity at most $2/27 + O(\epsilon)$. On the inverse side, $z = (1-r)/2 - \nu_0 > 0$ gives

$$1 - r - 2z = 2\nu_0, \qquad 3 - 2r - 8z = 4(1 - r - 2z) + (2r - 1) \ge 9/37.$$

These are fixed positive widths for the marked moment. If either capacity is at most ν_0, no-slot moments give at most $1 - \delta + O(\nu_0)$, below the short envelope. For $r \ge 1$, the amplified inverse exponent is $1 - \alpha + (\alpha - \delta)r \le L(\tau) + O(\epsilon)$; for $r \le 1$ the no-slot exponent is $1 - \delta r$. Their uniform agreement at $r = 1$ avoids an invalid marked-moment application with a vanishing width. If $m \ge 1/2$, use the no-slot plain moment. The floor $a = 51/100$ is excluded because its rows need not have an actual zero and hence need not have witnesses.

Balancing and the height ceiling. One has $D \ge 37/18$, $P \ge 7/9$ and $\alpha - \delta \ge 1/12$. Consequently $J > \delta P \ge 0$ and $1 \le \tau_* < 3/2$. The exact identity

$$D\{R_{\mathrm{short}}(\tau) - L(\tau)\} = \delta P/2 - J(\tau - 1)$$

proves the displayed balance and its minimax property. The algebra includes $\delta = 0$; physical nonfloor bins satisfy $\delta > 1/50$. Neither the algebraic endpoint nor finitely many rational test cases imply the source's arithmetic estimates.

Fix the internal moment and detector smoothness orders before the external Fourier-tail order. If A_{ht} dominates their finitely many height degrees, choose a positive analysis exponent ω_{ht} within the inherited ceilings and at most $d_{\min}\epsilon_{\mathrm{ht}}/[20(A_{\mathrm{ht}} + 1)]$. For $T_1 = Z^{\omega_{\mathrm{ht}}}$, eventually

$$(1 + T_1)^{A_{\mathrm{ht}}} \le 2^{A_{\mathrm{ht}}} Z^{d_{\min}\epsilon_{\mathrm{ht}}/20} \le U^{\epsilon_{\mathrm{ht}}/10}.$$

The later external tail order changes only external input seminorms, not any previously fixed internal height degree. This preserves the source's order of choices. It is not a height-uniform estimate without a ceiling, and it leaves the low-energy and nonprincipal-contour obligations separate.

O Support and offsets in the low-energy application

We record the analytic interface and the full real exponent behind Proposition 12.1. The pinned source passages and the remaining dependencies are listed in `research/low-energy.md`. The general reflected-energy and Gram estimates remain hypotheses.

A common Gaussian profile. For the surviving marked slots let $\varrho_i = q_{p_i}/P_i$. The physical completed series contains $W_{\mathrm{G}}(q_A/(Zq_{p_{J^c}}))$. On a retained annulus it becomes one joint profile

$$w_{k,J}(x,\boldsymbol{\varrho}) = \chi(\log x) W_{\mathrm{G}}\left(\frac{e^k x}{\prod_{i\notin J}\varrho_i}\right)\prod_{i\notin J} W_i(\varrho_i).$$

The logarithmic ratios lie in fixed compact sets. Every fixed weighted Fourier/seminorm sum is finite because each Gaussian derivative is a polynomial times a Gaussian in $k + O(1)$. Whole annuli with $|k| > \kappa_{\mathrm{G}} \log Z + O(1)$ have arbitrarily small polynomial total norm after fixing the orders. For the retained annuli, $N_* = 1 + \ell' + \theta$ with $|\theta| \le \kappa_{\mathrm{G}} + O(1/\log Z)$. Separate one density common to all rows and surviving tuples, and use Minkowski before squaring. No tuplewise or rowwise choice of Fourier density is allowed. The original normalization $q_c^{-1/2} q_n^{-1}$ stays unchanged when the annulus is recentered.

The sector identity covers all nonzero element rows, including those meeting S, with their original zero extensions. Here $f = 1$ and the puncture is 1; the rescaled tuple introduces no new moving puncture in B^J. Its row ball Q depends only on the fixed tuple p_J and not on surviving active slots or dual variables. Keep the actual residual-row annulus while extracting the kernel saving; only afterward may the positive row range be enlarged. Do not evaluate that saving on the enlarged rows.

Explicit real exponent. Write $z = z_a$, $S = S_0$, $N = N_0$, $B = B_0$, $e = e_\lambda$ and $u = \min\{v, z, (v+z)/3\}$. The support offsets can be bounded by $\varepsilon_Z = O_{\mathrm{fixed}}(1/\log Z) \ge 0$, with

$$O \ge -\varepsilon_Z, \quad H \le M' - O + \varepsilon_Z, \quad A_0 \ge 0, \quad 2A_0 \le O + \varepsilon_Z,$$
$$0 \le N \le A_0, \quad S, B \ge 0, \quad 0 \le z \le \ell', \quad v, \ell_b \ge 0, \quad e \ge -\varepsilon_Z.$$

The nonnegative v, ℓ_b are the actual ideal centers specified in the energy statement, not merely a weaker lower bound by $-\varepsilon_Z$. Source dual ideals may share primes and retain permitted primes of S; no additional coprimality is inserted. The count of frozen row parts is already charged by $O/2$. For a retained dyad put

$$T = 2H + 2A_0 + 2z - 1 - \ell' - \theta - N - 3B, \quad y = v + 3\ell_b + e \le T + \tau, \quad \tau \ge 0.$$

The exact source exponent is

$$E = O/2 + \max(H, v + \ell_b) - S - B + z - u - \ell_b - 2e/3 - (T-y)_+/2.$$

Since $M' + \ell' \le 1$, the support bounds imply $T \le H + 2\varepsilon_Z + |\theta|$, hence $\max(H, v+\ell_b) \le H + 3\varepsilon_Z + |\theta| + \tau$. Either $u = z$, in which case the positive z term cancels, or $u \ge v/2$. In the latter branch,

$$T/4 \le y/4 + (T-y)_+/2, \qquad u + \ell_b + 2e/3 + (T-y)_+/2 \ge T/4 - 5\varepsilon_Z/12.$$

The last allowance cannot simply be omitted: equality holds at $v = \ell_b = 0$, $e = T = -\varepsilon_Z$. Substitution of T, using $2A_0 - N + B + 4S \ge 0$, gives in both branches

$$E \le M' + \left(\frac{1 + 3\ell' - 2M'}{4}\right)_+ + 6\varepsilon_Z + 2|\theta| + \tau.$$

This full real inequality is proved in Lean with its explicit support/retention hypotheses. It does not assert an arithmetic mean-square estimate. For example choose $\kappa_{\mathrm{G}} \le \epsilon/32$, $\tau \le \epsilon/16$ and then Z large enough that the combined support/annulus offset $\varepsilon_Z \le \epsilon/128$. Since $|\theta| \le \kappa_{\mathrm{G}} + \varepsilon_Z$, the displayed offset is at most $3\epsilon/16$. Reserve the remaining allowance for the source output, common separation and dyad-count losses, each collectively at most $\epsilon/4$. Finite local choices and logarithmically many actual dyads fit the same budget. The general energy hypothesis now yields the displayed B^J norm with $s_t(d)$.

The loss and the external height order. The identity $1 + 3\ell' - 2M' = -1/6 + 5t + d$ follows before substituting any numerical target. Its positive part can be nonzero; for example its formal $s_t(d)$ envelope is $3t/2$ at $d = \ell_t$. That does not assert a positive actual norm loss in the empty-mark special case. The proof retains the envelope and absorbs it through $-d + s_t(d)/2 \le -7d/8$. For $Z \ge 1$, Lean also checks the actual real-power bound $Z^{-d+s_t(d)/2} \le 1$.

All internal reflection and Gram smoothness/height orders are fixed first. Their finite polynomial costs are integrated against weighted common Fourier densities and the transform of the fixed external smooth W_0. Choosing that external decay order later does not increase an already fixed internal height degree. Constants may depend on the fixed finite slot system, but no mesh power or Z-dependent height constant is introduced. This verifies the conditional low application; it does not certify the general reflection, Gram/correlation proof, exact arithmetic identities or the remaining nonprincipal contours.

P Central conductor, height and integral safeguards

These details support Proposition 13.1; pinned passages and its unverified arithmetic dependencies are in `research/central-contour.md`.

The correction is continued before it is divided. Contour movement uses the full finite tuple sum with selected G_p and unselected H_p, including at zeros of individual raw factors. On the retained dynamic lines only, the local estimates give $H_p - 1 = O(Q^{-1-10e})$ for $p \nmid u$ and $O(Q^{-10e})$ for $p \mid u$. A fixed sufficiently large $P_0(e)$ makes these H_p nonzero there. Only now define G_p/H_p and split each slot into main and error parts. The resulting factorization is not used for continuation or global tails. The four good-prime error exponents are

$$-a - 4e, \quad -51/25 + 12e, \quad 49/25 - 5a - 68e, \quad a - 51/25 + 12e,$$

all at most $-51/100$ on the stated range. The fixed identity ray does not impose $\eta(p) = 1$.

The strict ramified term requires one simultaneous conductor inequality. For its distinct labels J_0, retain the source arithmetic assertion

$$q_{\mathfrak{f}_u} \ll_S q_u \prod_{p \in J_0} Q_p^{-(j_p - 1)}.$$

The primitive functional equation and buffered reflected numerator bound cost $q_{\mathfrak{f}_u}^{A_*}$, up to small row powers and a fixed polynomial height, where $A_* = 1/2 - w_r = a - 1/2 + 6e \ge 0$. For every label,

$$-w_r - (j_p - 1)A_* = -1/2 - (j_p - 2)A_* \le -1/2.$$

Spend the deficit once for the whole set, before summing labels. Disjoint slot supports prevent the same prime from spending it twice. Divisor-many ramified choices cost a small row power.

Good-prime errors have the stronger 51/100 saving. Together these give the joint numerator/error estimate, with zero amplitude assigned to every error slot. Lean checks the simultaneous real-power inequality, not the arithmetic conductor identification or the buffered L-bound. Main slots contribute $\sum_i \ell_i g_i = q\ell$, a base-$Z$ power; it must not be replaced by $dq\ell$.

A cumulative height budget. For each buffered argument list once all its coordinates: $\gamma_j + \sum_{k=1}^{m} a_{jk} v_k$. Put

$$c_k = \frac{1}{2m(1 + \max_j |a_{jk}|)}, \qquad |v_k| \le c_k T_1.$$

Then $|\sum_k a_{jk} v_k| \le T_1/2$. If $|\gamma_j| \le (3i+1)T_1$, the resulting height is at most $(3i + 3/2)T_1$, within the $(3i+2)T_1$ buffer. A fixed bounded shift fits after increasing the threshold. The allowance is not renewed at successive prime, witness or row estimates.

An internal coefficient measure known only through a fixed weighted L^1 bound stays integrated inside the arithmetic factor. It does not become a pointwise external trace density. Extra integrals used for a central bound must be included before asserting its aggregate estimate; discarded parts stay on global or absolute lines. In particular integrated tails alone do not justify horizontal joins. The external trace estimate does not differentiate the arithmetic factor, so increasing its decay order cannot raise an internal moment-profile or height order.

The detector excludes bounded rows for which any inducing presentation in its finite family is principal. Numerator nonprincipality alone does not encode this full retained-family condition. Floor-bin rows remain permitted for central accounting, but have direct cardinality bounds; a saturated witness theorem is not asserted for them.

Only the original analytic sum is integrated. At a retained point the amplitude/witness partition may depend on that point. Apply triangle, bound the number of labels by a fixed logarithmic power, and dominate every exponent by a uniform supremum. The original analytic sum, times its kernel, is the function integrated. The pointwise pieces need not be integrated or assigned measurable labels. For an actual integrable complex function f and integrable real kernel k, the Lean component proves

$$\|f(x)\| \le Ck(x) \text{ a.e.} \quad \Longrightarrow \quad \left\| \int f \, d\mu \right\| \le C \int k \, d\mu.$$

The original integrability and pointwise envelope are explicit inputs; this theorem does not assert an arithmetic majorant from no hypotheses.

Boundary reference and the late height choice. The exact accounting also gives $E_\sigma - (\beta_* - \sigma) = E_{\beta_*}$. For frozen algebraic δ, q, R, the fixed-d perturbation is $t(13/50 + \delta/4 + q)$. At the moving endpoint $h_t = h_0 + 3t/2$,

$$E_t(h_t) - E_0(h_0) = t(-1/4 + \delta + q + 3R/2).$$

These identities do not equate the actual pointwise row partitions of two different probes or prove their exponents negative. If a positive common saving m is independently established and the internal height order A is fixed, choose a positive height exponent τ within its detector ceiling and $m/[4(A+1)]$. Only afterward choose N so $B - N\tau < C(\beta_*) - m/2$. Both choices can depend on the target, while m remains common. Lean checks the real and natural-number choices. It does not establish the hypothesized saving or a complete zero-free theorem.

Q Prime-power and common-factor arithmetic details

The finite-field results in Theorem 14.1 use actual `MulChar` and `AddChar` objects over arbitrary finite fields. The full ledger, pinned source passages and exact diagnostics are in `research/finite-correlation.md`.

Principal powers keep their zeros. A character of order six has principal cth character power exactly when $6 \mid c$. That power is the unit indicator, not the constant function 1 on the whole residue ring. The formal companion proves $(A^n)(0) = 0$ even for the zeroth character power; this is a character-group power, not the numerical convention 0^0. For every A the diagonal sum $\sum_x A(x)\overline{A(x)}$ is therefore $P - 1$. At a nonzero shift, the Jacobi reduction proves the exact autocorrelation before it is used to solve the affine field equation. No assumed correlation or zero-location property is inserted into that proof.

Lifts and normalized Gauss sums. At each successive level from $\mathfrak{p}$ to $\mathfrak{p}^c$, one coefficient in the genuine common congruence is a unit. One variable is free and the other determined, giving exactly P lifts per previous solution. The residue character values are unchanged, so the total factor is P^{c-1}. This remains true when the common prime divides one residual modulus. If both residuals meet it, the unit argument fails; the source no longer asserts this congruence-sum identity there.

Choose the fixed primary π with $(\pi) = \mathfrak{p}$. For $G(\mathfrak{p}^n, t)$, splitting $x = x_0 + \pi y$ makes the inner additive sum vanish unless $\mathfrak{p}^{n-1} \mid t$. When $t = \pi^{n-1}t_1$, it contributes P^{n-1}, and the remaining normalized factor is

$$P^{n/2-1} \sum_{x_0 \bmod \mathfrak{p}} \chi_{\mathfrak{p}}(x_0)^n e(t_1 x_0/\pi).$$

The good-prime trace pairing is primitive, since the excluded set contains the different; identifying this particular pairing is a source arithmetic input beyond the finite-field theorem. For nonprincipal multiplicative and nontrivial additive characters, conjugating the Gauss sum gives $g(A^{-1}, \psi^{-1})$. The formal Gauss product then gives $|g(A, \psi)|^2 = P$. For a principal multiplicative character with nontrivial additive phase, the sum is -1, also checked formally; at trivial phase it is $P - 1$. These are exactly the two prime-power formulas in the core text. The field statements are kernel-checked; the quotient-ring splitting and its relation to the source pairing are prose.

The full correlation and its phase. Expansion of the source's normalized Fourier formula gives the prefactor $1/(N(u)N(v))$. Additive orthogonality over $\mathcal{O}/(uv)$ contributes $N(uv) = N(u)N(v)$, even for shared prime powers. Thus no gcd factor is missing. At zero frequency a nonzero summand has unit variables; reducing modulo u and v forces mutual divisibility. For the chosen primary generators this is $u = v$, and the answer is $\varphi(u)$.

For $C = (u, v)$, $u = Cn_1$, $v = Cn_2$, the sum vanishes unless $C \mid j$. At $j = Ct$, reduction modulo the coprime residuals extracts

$$\chi_{n_1}(t)\overline{\chi_{n_2}(-t)}\frac{\chi_{n_2}(n_1)}{\chi_{n_1}(n_2)}.$$

All inverted symbols are units. CRT leaves exactly the common kernel L_C, even when C meets one residual. The source identifies the unit ratio with its fixed reciprocity factor $\mathcal{R}(n_1, n_2)$. This audit checks its orientation, but does not re-prove that earlier reciprocity theorem or its fixed ray-table implementation.

For complete common support $u = Da$, $v = Eb$, $(a,b) = 1$, $(ab, DE) = 1$, the changes $x' = bx \bmod D$, $y' = ay \bmod E$ leave the phase

$$\frac{\chi_E(a)}{\chi_a(E)}\frac{\chi_b(D)}{\chi_D(b)}\frac{\chi_b(a)}{\chi_a(b)} = \mathcal{R}(a,E)\overline{\mathcal{R}(b,D)}\mathcal{R}(a,b).$$

Every denominator is again a symbol of a unit. Off this locus the stipulated extension and complete Möbius divisor sum must be retained; an individual divisor term does not turn a newly common prime into genuine D, E support.

Genuine and artificial kernels are different. With both coefficients zero and principal A, the genuine field kernel at zero frequency is $(P-1)^2$. The actual $\mathbb{F}_7$ example is $36 > 7$. By contrast, the source's artificial extension assigns zero. The latter keeps the desired periodicity and bound, but is not the original congruence sum at an added pair. Subsequent positive enlargements must preserve the exact Möbius identity before estimating the columns separately. A failure to retain this distinction would require repair; no such failure is established by the present finite audit.

The deterministic checker covers every coefficient pair, right side and character power $1, \dots, 6$ in actual Eisenstein fields of sizes $7, 13, 25$. The inert size-25 field is $\mathbb{F}_5[\omega]/(\omega^2 + \omega + 1)$, not $\mathbb{F}_5$. Additional exact Hensel and CRT tests cover repeated common primes and vanishing symbols. These samples complement the universal Lean field proofs; they do not establish the quantitative Gram estimate, the reflected-energy identity or any global zero-free theorem.

R Gram cancellation: masks, frequency volume and tails

The exact audit target is lines 8345–8560 of the pinned September source [3], especially its periodic-Poisson equation. The following expands Proposition 15.2 and separates the surviving arithmetic inputs from the general analytic lemma. The prescribed fixed polynomial ranges in Z are retained when logarithmic and arithmetic small-power losses are absorbed in Z^ϵ.

Complete extension before separation. Write $s_i = Cn_i$ with $(n_1, n_2) = 1$. The complete correlation expansion cancels the row Poisson denominator $q_{s_1}q_{s_2}$ and leaves Q/Y'^3. The source's artificial L_C is bounded by q_C, periodic modulo $\mathrm{rad}(C)$ and zero when a prime of C meets both residuals. Insert the entire Möbius divisor sum for coprimality before taking absolute values. In a term $n_i = dm_i$, the original zero masks at primes of k, and the artificial zero at primes of C, force $(d, Ck) = 1$. Principal character powers retain these zeros. No new coprimality condition is imposed on m_1, m_2.

Assume that the exact earlier reciprocity/ray theorem supplies a fixed modulus $\mathfrak{l}$ independent of C, d, k. The joint period divides

$$\mathfrak{r} = \mathrm{lcm}(\mathfrak{l}, \mathrm{rad}(C), \mathrm{rad}(k)), \qquad N\mathfrak{r} \ll q_C q_k.$$

If a prime $\mathfrak{p} \nmid C\mathfrak{l}$ has $e = v_\mathfrak{p}(k) \not\equiv 0 \pmod 6$, CRT leaves one nonprincipal factor $\chi_\mathfrak{p}(m_1)^e$; every other factor is independent of that coordinate. Multiplication by d is invertible there. Arbitrary coupling among the remaining residues and the second column cannot change the vanishing complete mean. The finite inference is formalized; the source-specific fixed modulus is still an input.

One dual sum, not two period volumes. For a coefficient periodic modulo a generator r of $\mathfrak{r}$, let $q = N\mathfrak{r}$ and

$$\widehat{a}(\psi) = q^{-2} \sum_{x \in (\mathcal{O}/(r))^2} a(x)\overline{\psi(x)}.$$

There are q^2 pairs, so $|\widehat{a}| \le q_C$ in this application. Poisson applied after finite Fourier inversion yields

$$\frac{N^2}{v^2} \sum_{\xi \in (r\mathcal{O})^{*2}} \widehat{a}(\psi_{-\xi})\widehat{W}(\sqrt{N}\xi).$$

The residue character and its ordinary dual-lattice translate already index this single lattice. Multiplying by an additional q^2 would count the same frequencies twice. The coefficient at zero vanishes; so do all frequencies restricting to the trivial character. Since principal ideals of $\mathbb{Z}[\omega]$ are similar to one fixed lattice, there is no changing lattice-shape constant. Fourier decay of order $2A > 4$ controls the nonzero sum uniformly up to $N \asymp q$.

For $N < q$, source support in both column annuli excludes zero coordinates and forces N above a fixed positive threshold whenever the sum is nonempty. The direct count is then $O((1+N)^2) = O(N^2)$. For comparison, a coefficient $(-1)^{u_1}$, where u_1 is the first integral coordinate of the first Eisenstein column, has mean zero modulo 2 and value 1 at the origin. A smooth bump supported sufficiently near the origin leaves that term alone as $N \to 0$. Thus an unrestricted Schwartz formulation allowing $a(0)W(0) \neq 0$ cannot have the same N^2 bound. Appropriate vanishing and decay could replace annularity, but no such replacement is needed here.

The exact dyadic tail. For $2^L \le \Lambda < 2^{L+1}$ and $A \ge 2$,

$$\sum_{n=0}^{L} (n+1)2^{-n}(2^n/\Lambda)^A \le \Lambda^{-2} \sum_{n=0}^{L} (n+1)2^n = \frac{L2^{L+1}+1}{\Lambda^2},$$

$$\sum_{n=L+1}^{\infty} (n+1)2^{-n} = (L+3)2^{-L}.$$

The first expression is at most $2(L+1)/\Lambda$ and the second less than $2(L+3)/\Lambda$. Consequently $S_A(\Lambda) \le 4(L+2)/\Lambda$. At the exact cutoff,

$$S_2(2^L) = 3(L+1)2^{-L} + 4^{-L}.$$

For example $S_2(1) = 4$, $S_2(2) = 13/4$ and $S_2(4) = 37/16$. These include the closed infinite tail, rather than silently replacing an infinite sum by a truncated experiment. The new Lean module proves the head, total, tail and head bound; the diagonal regrouping and full comparison remain prose.

Exceptional frequencies and remaining scope. With $K_R = RP/C_0$, exceptional ideals have $(k) = \mathfrak{a}_1^6\mathfrak{e}$, where $\mathfrak{e}$ has exponents $0, \dots, 5$ supported on $C\mathfrak{l}$. There are $O_\epsilon(q_C^\epsilon)$ choices and $O(K_R^{1/6})$ sixth-power parts in a nonempty shell. Their total before the outer Q/Y'^3 is

$$Y'^2(RP)^{1/6}C_0^{-1/6+\epsilon}D_0^{-1}.$$

Choose $\epsilon < 1/6$ before summing the two dyads. A sufficiently large fixed smooth decay order then sums the R shells. This accounts for every scale in the claimed Gram bound and exposes no extra period loss. Exact diagnostics check the finite Fourier normalization, character cancellation, scale

identity and dyadic series. They do not prove the source reciprocity theorem, its trace pairing, the reflected identity, moments, outer-row estimates or final closure. These and the arithmetic spectral bridge remain separate research tasks.

S From the phase table to the literal column period

A nonminimal modulus with explicit exponents. The modulus $\mathfrak{m}_0$ of Proposition 16.1 has exponent $6v_{\mathfrak{p}}(6)+25$ at every $\mathfrak{p}\in S$. For S consisting just of the primes above $2,3$, this is 31 at (2) and 37 at (λ). These are upper defining exponents, not exact conductors. Existing complete ray presentations remain in $\mathfrak{m}_{\mathrm{pres}}$. In particular a primitive conductor never erases a redundant zero prime.

To justify the discriminant step even in smaller degree, let $\xi^6=a$ and $g=\mathrm{minpoly}_F(\xi)$. The roots of g are integral, so its coefficients lie in the normal ring $\mathcal{O}$; division gives a monic integral h with $X^6-a=gh$. The root-difference discriminant identity and resultant are integral, and the polynomial is separable. Locally at every prime, the trace matrix for the power basis changes to an integral basis with determinant an integral index, so the field discriminant divides the order one [15, Proposition 2.11, Lemma 2.23 and Proposition 2.34]. Conductor–discriminant includes every character of the cyclic group; the faithful μ_6 character is among them, also for degrees $1,2,3$. This proves all ideal divisibilities used in the core proof without assuming irreducibility of X^6-a or a power integral basis.

The Gaussian normalization and square classes. For the self-dual source measure $d\mu=(2/\sqrt{3})dx\,dy$, apply Poisson to $e(z^2/c)e^{-\pi\varepsilon|z|^2}$. The quadratic determinant is $\varepsilon^2+16/(3|c|^2)$. Writing $r_\varepsilon=1+3|c|^2\varepsilon^2/16$, its transform is

$$\frac{|c|}{2\sqrt{r_\varepsilon}}e^{-\pi\varepsilon|c|^2|y|^2/(4r_\varepsilon)}e\left(-\frac{cy^2}{4r_\varepsilon}\right).$$

Divide by $I_\varepsilon=2/(\sqrt{3}\varepsilon)$. Original residue means modulo c give $\Gamma(c)/|c|$. On the dual side replacing r_ε by 1 costs $O_c(1)=o(I_\varepsilon)$ in the phase, and smaller normalized amplitude/damping errors. Each of the four classes modulo 2 has damped normalized mean $1/|c|^2$. The limit is the four-term formula. This argument fixes c before letting $\varepsilon\downarrow 0$; it asserts no uniform moving-modulus limit.

The twelve units modulo 4 have square subgroup $\{1,\omega,\omega^2\}$. If square classes are written $(-1)^e\lambda^f$, then

$$\mathcal{R}((-1)^e\lambda^f,(-1)^g\lambda^h)=(-1)^{eh+fg+fh}.$$

Its diagonal is multiplicative. The element residue class of -1 is used here; the ideal (-1) is the identity ideal and cannot be substituted for that residue class. For coprime primary pairs cubic reciprocity makes the sextic quotient a sign; quadratic CRT computes its cube as the displayed sign. This establishes the orientation. No GRH-conditional analytic theorem of [16] is invoked.

Every literal factor. After the complete Möbius insertion, the joint coefficient is, up to fixed scalar conventions,

$$\begin{aligned}a_{C,d,k}(m_1,m_2)=\mu(d)c_\sigma(Cdm_1)\overline{c_\sigma(Cdm_2)}\,\chi_{dm_1}(k)\overline{\chi_{dm_2}(-k)}\\ \cdot\mathcal{R}(dm_1,dm_2)L_C(dm_1,dm_2;k).\end{aligned}$$

Here $c_\sigma(s)=1_{s\in\sigma}\chi_s(b_*)/(\tau\xi(s))$ is extended by zero off primary elements prime to S before division. The symbol with numerator b_* has conductor dividing $\mathfrak{m}_0$; ξ and the full σ presentation are fixed. On

the nonzero branch, multiplication by primary good C, d only changes an input residue. Congruent unit elements modulo a defining ray modulus generate ideals with the same character value, since their quotient has local representative 1 there. On nonunits the common zero branch is used first.

The factor $\mathcal{R}(dm_1, dm_2)$ depends only on residues modulo 4. The artificial L_C has period $\mathrm{rad}(C)$, including when C meets a residual. If d meets C, its artificial zero kills the term; if d meets k, the original symbol zero kills it. Otherwise factor $\chi_{dm}(\pm k) = \chi_d(\pm k)\chi_m(\pm k)$. For $k = \zeta k_S k_{\mathrm{good}}$, the fixed numerator reduces the S-valuations modulo six and uses $\mathfrak{m}_0$. The good reciprocity phase uses 4 and the remaining local powers use only their good residue fields. The sign change uses $-\zeta$ in the same fixed family. No moving prime enters $\mathfrak{l}$.

At a nonexceptional prime outside $C\mathfrak{l}$, the first column contains just the nonprincipal $\chi_p(m_1)^{e_p}$ in that CRT coordinate; arbitrary dependence in the other coordinates is harmless. Since $(d, k) = 1$, multiplication by d is invertible at p. This supplies the complete zero mode needed by the Gram proof.

The formal companion uses QuadraticAlgebra$(\mathbb{Z}/4\mathbb{Z}, -1, -1)$, with actual Eisenstein multiplication, and its integer counterpart. Thirteen declarations check the ring relation, unit inverse/count/closure, square-class homomorphism/kernel, four-term table/refinement, sign bicharacter/diagonal and integer period. They do not formalize class field theory, the Gaussian integral, cubic reciprocity or the full analytic moments. Exact split and inert field diagnostics include repeated powers, shared zeros, actual signed row conversions and the norms $31, 43$ witness. Finite samples do not replace the classical universal conductor proof or establish GRH.

T Exact reflected phases, shared cubes and the common density

Proposition 17.1 audits September lines 7747–8005 and their canonical scalar inputs. It assumes the stated theta/cusp reflection identity, rather than replacing that identity by finite-field experiments. The branch denominator is $c = c_F F_{\mathrm{act}} RP$. Separate full sectors of R, P fix the class of their product and hence d, ϑ, κ_F without imposing a pair-dependent restriction. An inactive marked prime is absent from the conductor and carries its q_p^{-1} weight. Its absolute mass is removed by Hilbert-space triangle before applying the active-block norm.

The phase computation and its obstruction. For a unit z and unit w, the exact integer-exponent identity is

$$z^{-2}(wz^{-2})^{-(j+2)} = w^{-(j+2)} z^{2j+2}.$$

It explains the local dependence on c/p without dividing a zero symbol. For $a^2 = b^2$ one has $a^{2j+2} b^{2k+2} = (a^2)^{j+k+2}$. If $a^6 = 1$, the row–mark case is one; the moving $j = 4$ case equals a^2. The actual norms $7, 13$ example in the core text has $a^2 = \omega$. Thus freezing those nonresidual primes is an arithmetic condition, not an optional simplification. The changed marked sign remains a one-prime scalar. Interactions with frozen primes depend on only one moving product. The angular factor $\bar{\alpha}(c)^2$ separates multiplicatively.

A shared cube prime is not extracted twice. Let a frozen Ramanujan prime have $v_p(n_0) \in \{0, 1\}$ and $v_p(b_0) = b_p \geq 0$. If $v_p(n_0) = 1$, the N_F assignment removes that one occurrence from n_0 and leaves all b_p occurrences in b_0. The dual valuation remains $1 + 3b_p$. If $v_p(n_0) = 0$ and $b_p \geq 1$, the B_F assignment extracts one occurrence from b_0 and leaves $b_p - 1$; the dual valuation is $3 + 3(b_p - 1) = 3b_p$. The divisibility coefficient contributes $q_p^{1/2}$ only once. It cancels $q_p^{-1/2}$ in the

first assignment and leaves $q_p^{-1/2}$ in the second. These calculations remain valid when other frozen primes occur in both dual ideals.

The bound $|d(\mu)| \le 27\,3^{h/6}|B_F b|$ and $q_\mu = 3^h q_{N_F n} q_{B_F b}^3$ give the exact extraction in the core. The ramified power is $3^{h/6-h/2} = 3^{-h/3}$, including $h \ge -4$. Together with the canonical scalar $1/81$, this leaves $1/3$. The normalized dual coefficient is bounded by one and is independent of the row and active tuple after the full sectors are fixed. It still includes the original source coefficient and all frozen local restrictions. A numerator such as -2 is allowed in the source dual support; evaluating a good moving character there does not permit adding an S-mask to the dual sum.

The one joint profile. Write $y_R = q_R/Z^H$, $y_n = q_n/Z^v$, $y_b = q_b/Z^{\ell_b}$ and $y_i = q_{p_i}/Z^{z_i}$. Then, on the actual cutoffs,

$$\frac{q_\mu X}{q_c^2} = Y y_n y_b^3 y_R^{-2} \prod_i y_i^{-2}, \qquad Y = q_{c_F}^{-2} Z^{v+3\ell_b+e_\lambda-T_d}.$$

This uses actual products, not a coprimality approximation. One may retain $q_P^{-1/2}$ explicitly and put the remaining inverse roots, norm windows and kernel in a single profile. All norm labels are coordinates of that profile; they do not index independently selected measures.

Let $D_x = x\partial_x$ and $\mathfrak{m}_A(Y) = \min\{Y^{1/4}, (1+Y)^{-A}\}$. For the kernel in the core proof, the first numerator gamma pole is $-5/6$ and reciprocal denominator gamma is entire. Vertical Stirling bounds cancel the exponential factors and leave the power $4\Re t$. The Mellin transform of a rapidly decreasing smooth test is entire and rapidly decreasing on fixed strips, by integration by parts. Consequently the shifts and Euler differentiation under the integral are justified. Three line bounds give the displayed small-, bounded- and large-argument estimate, with constants depending on the fixed order.

If $h(\boldsymbol{y}) = \prod y_i^{a_i}$ and the logarithmic support is fixed and compact, then $D_i V^\sharp(Yh) = a_i D_x V^\sharp(Yh)$. Leibniz and logarithmic Fourier integration by parts give

$$\|F_Y\|_{J,\mathrm{sep}} \ll \mathfrak{m}_A(Y) p_{J+d+2}(W) \mathfrak{M}_{A,\lceil 4A\rceil+J+d+4}(V).$$

The Fourier weight is integrable; each fixed annular row/tuple/dual block is finite, so interchange with the one density is legitimate. The extracted scalar therefore enters the linear row vector once. The new actual Bochner theorem, for an explicitly integrable density in a complete complex normed space, proves

$$\left\| \int (c\phi(t)) B(t)\, dt \right\| \le |c| \int |\phi(t)|\, \|B(t)\|\, dt.$$

It does not assert that the arithmetic representation or gamma estimates are formally proved. Squaring this inequality supplies $|c|^2$; positive row enlargement occurs only after separation. Whole discarded dyads are bounded on their original kernel arguments before taking Fourier absolute values.

The nine Lean declarations and exact diagnostics verify these local and norm components, including zero inputs and repeated cube primes. They do not prove the canonical theta transformation, identify its three cusp coefficient families, establish the hybrid quadratic–cubic large sieve or complete the final common-signal argument. Those remain distinct mathematical obligations.

U Computational examples and reproducibility

The Python checks use integers and `fractions.Fraction`, with no random choices, model weights, scientific datasets or floating-point zero finding. They reproduce the Iwahori witness at $\pi = 2$, the

determinant change $2 \to 1$, the parity obstruction, and the vanishing diagonal model. The Lean statements verify selected finite calculations and the parity argument in the kernel of a pinned Lean 4 toolchain. Neither executable claims to verify the full complex-analytic arguments in this article.

Both editions share the same core source. The longer edition adds these calculations and research interfaces; extra exposition does not strengthen an unproved hypothesis into a theorem. The source manifest and audit coverage record distinguish full text review, focused review, and unverified dependencies. Publication readiness also requires independent mathematical review, confirmed authorship, and journal-specific preparation.

V Weighted profiles, exact arithmetic examples and the surviving task

The finite matrix proof also retains an arbitrary row profile. Suppose $|A_{ui}|^2 = w_u$ for every column, set $S = \sum_u w_u$ and $Q = \sum_u w_u^2$. The row diagonals are Nw_u and column diagonals are S, so the same identity proves

$$\sum_{i \neq j} |K_{ij}|^2 \geq N^2 Q - NS^2.$$

This is a finite statement. Applying it to an infinite Schwartz tail would require its convergence and every newly encountered prime collision; these are not discarded. If a weighted pair-square attempt includes the annular W, one can instead choose a closed subannulus on which a nonzero $|W|$ is bounded below. Its prime count still has order $D/\log D$. The fixed ν has unit modulus outside its conductor. These facts do not replace the original coefficients in a signed estimate.

Here are exact sextic computations with all nonzero elements of the Eisenstein norm ball, and both prime ideals above each split rational prime in the column annulus. These kernel diagnostics use $\nu = 1$ and exclude only primes above2,3; they do not compute a signed Möbius energy. Values are integer norms in $\mathbb{Z}[\omega]$.

D	H	R	N	$\sum_{i \neq j} \lvert K_{ij} \rvert^2$	$NR(N-R)$
128	4	18	22	6840	1584
1024	9	36	138	651672	506736
4096	81	300	456	60080328	21340800
256	64	240	40	363672	−1920000

The final row has $H = D^{3/4}$ but $N < R$; its negative lower bound is correctly vacuous. The asymptotic theorem follows from the proof, not an extrapolation of these examples. For $p \equiv 1 \pmod 3$ and $w^2 + w + 1 = 0$ in $\mathbf{F}_p$, evaluate $a + b\omega$ as $a + bw$ and take its $(p-1)/6$ power, lifting the six roots via $\zeta_6 = 1 + \omega$. Every zero is tested before a unit value is assigned. For instance $1 + 3\omega$ has norm7 and vanishes at $\omega = 2$ modulo7, so the unimodularity hypothesis cannot be used beyond its stated domain.

`ShortRowPairBarrier.lean` verifies the finite frame identity, row-diagonal bound, weighted inequality and unimodular specialization, with actual complex sums and explicit hypotheses. It does not formalize the ideal-character comparison or prime-ideal theorem. The deterministic integer checker also verifies weighted profiles and256 scalar pair sums independently of its matrix computation. No numerical model or learned coefficient is used.

The next signed route must retain μ, ν, W before Cauchy. An exact starting identity in ideal Dirichlet convolution is

$$\mu = 2a - a * a * \mathbf{1} + b * b * \mathbf{1}, \qquad a = \mu\, \mathbf{1}_{Nn \leq U}, \quad b = \mu - a.$$

Indeed $\mu * \mathbf{1} = \delta$ implies $(\mu - a) * (\mu - a) * \mathbf{1} = \mu - 2a + a * a * \mathbf{1}$. On squarefree n, each divisor factorization retains pairwise coprimality; the original row zeros still multiply along it. Estimating these type-I/type-II kernels with their true conductors is a new open arithmetic task. The identity alone supplies no cancellation or GRH proof.

References

[1] A. Schreiber, *Emergent Cyclic Time from Surface Orders and a Hilbert–Pólya Program for the Generalized Riemann Hypothesis*, repository source, commit `212ff730`, `https://github.com/amelie-iska/Surface-Orders-Hilbert-Polya`.

[2] A. Schreiber, *Surface-Order Determinants, Monomial Artin L-Functions, and the General Hilbert–Pólya Extension Problem*, repository PDF, commit `0c77122d`, `https://github.com/amelie-iska/GRH`.

[3] *Mathematical manuscripts and supporting proof artifacts*, supplied repository, `https://github.com/amelie-iska/math`. See project source manifest for the exact local pin and review scope.

[4] J. C. Lagarias, Li coefficients for automorphic L-functions, *Ann. Inst. Fourier* **57** (2007), 1689–1740, `https://arxiv.org/abs/math/0404394v4`.

[5] A. Connes, Trace formula in noncommutative geometry and the zeros of the Riemann zeta function, *Selecta Math.* **5** (1999), 29–106, `https://arxiv.org/abs/math/9811068`.

[6] The mathlib community, *Riemann zeta function documentation*, `https://leanprover-community.github.io/mathlib4_docs/Mathlib/NumberTheory/LSeries/RiemannZeta.html`.

[7] M. Griffin, K. Ono, L. Rolen and D. Zagier, Jensen polynomials for the Riemann zeta function and other sequences, *Proc. Natl. Acad. Sci. USA* **116** (2019), 11103–11110, `https://doi.org/10.1073/pnas.1902572116`.

[8] M. B. Nathanson, The Hermite–Sylvester criterion for real-rooted polynomials, version 2 (2019), `https://arxiv.org/abs/1911.01745v2`.

[9] D. Romik, Orthogonal polynomial expansions for the Riemann xi function, version 3 (2019), equations 1.8–1.11, `https://arxiv.org/abs/1902.06330v3`.

[10] J. Tate, *Fourier Analysis in Number Fields and Hecke's Zeta-Functions*, Princeton dissertation (1950), sections 4.4–4.5, `https://sites.math.rutgers.edu/~alexk/2023S572/Tate1950.pdf`.

[11] D. White, Extreme values of Hecke L-functions to angular characters, version 1 (2022), Theorem 2, `https://arxiv.org/abs/2208.02309v1`.

[12] J. Thorner and A. Zaman, A unified and improved Chebotarev density theorem, *Algebra Number Theory* **13** (2019), 1039–1068, Theorem 1.1, `https://doi.org/10.2140/ant.2019.13.1039`.

[13] M. Stoll and mathlib contributors, Jacobi sums, Gauss sums and multiplicative characters, Lean source at commit `f897ebcf72cd16f89ab4577d0c826cd14afaafc7`, `https://github.com/leanprover-community/mathlib4/blob/f897ebcf72cd16f89ab4577d0c826cd14afaafc7/Mathlib/NumberTheory/JacobiSum/Basic.lean`.

[14] J. S. Milne, *Class Field Theory*, version 4.03 (6 August 2020), Theorem V.3.27, Remark V.3.28 and Sections VIII.5.3–5.5, https://www.jmilne.org/math/CourseNotes/CFT.pdf.

[15] J. S. Milne, *Algebraic Number Theory*, version 3.08 (19 July 2020), Proposition 2.11, Lemma 2.23 and Proposition 2.34, https://www.jmilne.org/math/CourseNotes/ANT.pdf.

[16] A. Dunn and M. Radziwiłł, Bias in cubic Gauss sums: Patterson's conjecture, version 3 (14 May 2024), equation (1.4), https://arxiv.org/abs/2109.07463v3.

[17] NIST Digital Library of Mathematical Functions, Sections 5.2 and 5.11, especially equation (5.11.9), https://dlmf.nist.gov/5.2, https://dlmf.nist.gov/5.11#E9.

[18] S. Datta, S. Howard and D. Cochran, *Geometry of the Welch Bounds*, arXiv:0909.0206v2 (2012), Section2.1, Theorem2.2.