Infinite dessin orders, projective resolutions and spectral determinants

Extended proofs, Schur functors and Auslander–Reiten calculations

Research working draft

November 3, 2026

Abstract

We work throughout with infinite-dimensional locally gentle algebras and completed surface orders arising from explicit cellular clean dessins on compact Riemann surfaces. We construct their projective resolutions, Cartan determinants and dual dessins; develop determinant semi-invariants, Schur modules and cyclic-quiver character kernels; and calculate actual lattice almost-split sequences and an affine-type diamond quotient. A constrained-path Fock algebra retains the gentle relations and has a compact extension with incidence boundary map. Its degree spectral triple and an inserted arithmetic Fredholm family are explicit. Jacobi sums provide theta-dessin cellular Frobenius determinants with a positive local metric; inert primes obstruct the same real two-dimensional carrier. Recent spectral approaches are compared with their actual hypotheses. Selected power-series, module and matrix statements are verified in Lean4. A derived branch functor repairs projective determinant multiplicity; tame descent recovers a hereditary order but forces multiplicities for full-deck coefficient kernels. The arithmetic positive determinant comparison remains open; no proof of GRH is claimed.

Contents

1 Cellular dessins and their infinite orders **4**

2 Projective resolutions, dual dessins and determinants **5**
2.1 Auxiliary staircase resolutions over the order 7
2.2 A monotone quantum comparison with exact finite ranks 8
2.3 A correct DG model of the infinite node 8

3 Determinant semi-invariants and Schur constructions **9**
3.1 Young tableaux and a cyclic representation stratum 10

4 Lattice almost-split sequences and genuine diamond quotients **11**
4.1 Polynomial strings, completed lattices and the Fukaya category 12

5 Leavitt relations, the constrained Fock algebra and spectrum **13**
5.1 A graph presentation and an actual pullback 14
5.2 A separate diagonal-fiber C* completion retaining the pairing 15

6 Arithmetic attachment, descent and the remaining comparison **16**
6.1 A constructive arithmetic benchmark on the theta dessin 17
6.2 A genuine cellular carrier, and what still fails for the order 18
6.3 Arbitrary projective actions: an obstruction and a genuine determinant repair 19
6.4 Tame descent: a surviving order and a sharper obstruction 20
6.5 Whole components, derived endpoints and determinant relations 22

7 A genuine theta covering and its categorical limits **23**

8 Verification and research frontier **24**

9 Why an absolute coprime pair-square estimate cannot close the short rows **26**

10 A signed Möbius decomposition and its termwise obstruction **27**

11 A full-mask Hecke expansion for the type-I piece **28**

12 Actual renormalization retains the effective action **29**
12.1 Genuine dessin incidence models 30
12.2 Quantum thermal effective Hamiltonians 30

13 Linear relations and functorial branch filtrations **31**

14 A kernel-checked extension over the infinite dessin algebra **31**

15 Learned fields, width limits and quantum geometry **33**

A Expanded dessin, resolution and derived calculations **34**
A.1 The cycle order in coordinates 34
A.2 The two-arm complex and its calculations 35

B Semi-invariants, tableaux and explicit intertwiners **36**
B.1 Derived information retained by a presentation 36
B.2 One Young operator and one LR decomposition in coordinates 37
B.3 Two precise functorial limitations 38

C Full lattice and diamond diagrams **38**
C.1 The nodal exact diagram 38
C.2 Chebyshev geometry and the exact quotient 39
C.3 Nine actual strings and all maps 39
C.4 The pure-injective quotient and what its dimension means 40

D The operator extension in detail **41**
D.1 A corrected cycle lift and the integer boundary map 41
D.2 Spectral domain and arithmetic trace 41

E Abel determinant estimates and the arithmetic program **42**
E.1 Precisely how classical and recent sources enter 43
E.2 A concrete sequence of remaining constructions 44

F Graph pullbacks and completions with specified fibers **44**
F.1 The complete Fock graph presentation 44
F.2 A genuine application of the supplied pullback theorem 45
F.3 Diagonal residue obstruction and the corrected pullback 46
F.4 Generation, faithfulness and the exact failure of extension 46
F.5 Invariant basis number and the Leavitt comparison 47

G String finiteness and the smooth quotient surface **47**

H Arithmetic character sums and the cellular determinant **48**
H.1 The uniform calculation and its exceptional cases . . . 48
H.2 Deck descent and the actual cellular basis . . . 49
H.3 An explicit chain lift and determinant cancellation . . . 50
H.4 A decisive restriction at inert primes . . . 50
H.5 The remaining order and global spectral obligations . . . 51

I The theta order interface beyond tensor insertion **52**
I.1 Normal forms and the special fiber . . . 52
I.2 All projectives and all linear endomorphisms . . . 52
I.3 An actual integral determinant functor of multiplicity one . . . 53
I.4 The exact rotation obstruction and its allowed escape . . . 54

J Explicit tame descent and the full-deck coefficient test **55**
J.1 The action on genuine formal series . . . 55
J.2 A projective symmetry survives on the endomorphism order . . . 56
J.3 Determinants, Schur weights and the boundary correction . . . 56
J.4 Why the local Jacobi plane does not descend by this repair . . . 57
J.5 The full orbit is the normalization, and arbitrary kernels still fail . . . 57

K Literal formal and computational coverage **59**

L Explicit covering calculations and derived transport **60**

M Weighted profiles, exact arithmetic examples and the surviving task **61**

N Exact factorization, retained zeros and the failed Cauchy route **62**
N.1 Proof of the regrouping with original coefficients . . . 62
N.2 The two conductor scales are unchanged . . . 62
N.3 Why a first generic Cauchy step does not supply the saving . . . 63
N.4 Exact finite checks and formal boundaries . . . 64

O Primitive conductors, deleted factors and the scalar contour **64**
O.1 Constructing the column character without losing its zeros . . . 64
O.2 Why the half-plane above one half is available . . . 65
O.3 Mellin inversion, residues and profile uniformity . . . 65
O.4 Verification boundaries . . . 66

P Controlled errors and a trained full-action RG approximation **67**
P.1 Executed finite calculations . . . 67
P.2 A quantum-feature model trained on actual RG . . . 67
P.3 Quantum TokenGT and the geometric error-correction boundary . . . 68
P.4 An actual finite gentle cut for the categorical interface . . . 68

Q Relation calculations, splitting functors and determinant limits **69**
Q.1 Correct graphs and stable spaces . . . 69
Q.2 Proof of the dessin-category equivalence . . . 69
Q.3 Functoriality and the actual filtration . . . 70
Q.4 Strings, bands, Hom and one non-rigid string . . . 71
Q.5 A determinant comparison with its exact limits . . . 71

R Normal forms and the full module action **72**
R.1 Why the polynomial matrix algebra is the path algebra . . . 72
R.2 Coefficient selectors without a truncation . . . 72
R.3 Constructing every normal image from actual arrows . . . 73
R.4 Spanning, multiplication and the inverse coordinate map . . . 74
R.5 Exactness for arbitrary algebra elements . . . 75
R.6 Formal and computational coverage . . . 75

S Physical generative models and their verification **76**
S.1 From a dessin to an actual quantum RG target . . . 76
S.2 Diffusion and exact versus approximate reverse controls . . . 76
S.3 Flow matching with a measurable target velocity . . . 77
S.4 Architecture, matched controls and actual GPU results . . . 77
S.5 A coherent learned purification flow . . . 79
S.6 Width, fields, RG and arithmetic obligations . . . 79

T Covers, paths, pants and quantum games **80**
T.1 Which free group and which universal cover? . . . 80
T.2 Actual infinite-algebra path modules and unbounded resolutions . . . 81
T.3 Pants tensors, physical dilation and sewing boundaries . . . 83
T.4 Quantum games and the Connes-embedding distinction . . . 83
T.5 Tropical accelerator reference designs and reproducibility . . . 84

1 Cellular dessins and their infinite orders

We start with the surface-order constructions of Schreiber [1, 2], retaining the original sources unchanged. The generic-resolution source [3] is background, not a substitute for exactness of a particular complex. All primary algebra examples below come from explicit clean Belyi maps. Finite-dimensional objects occur as modules or specified quotient strata of those infinite algebras, not as replacements for them. The initial arithmetic target is a primitive nonprincipal Dirichlet L-function. Finite-order Hecke characters, Dedekind zeta functions and general Artin representations require separate comparison theorems; Artin holomorphy is not assumed.

A clean dessin on a compact oriented surface is specified by a nonempty finite dart set H, a fixed-point-free involution α, and a permutation σ such that $\langle \alpha, \sigma \rangle$ is transitive. Contract the white vertices of degree two. The resulting cellular graph has vertices the σ-orbits, edges the α-orbits, and faces the ϕ-orbits, where

$$\tau = \sigma^{-1}, \qquad \phi = \alpha\sigma^{-1}, \qquad 2 - 2g = \#\operatorname{cyc}(\sigma) - |H|/2 + \#\operatorname{cyc}(\phi).$$

Write $[h] = \{h, \alpha h\}$. Our quiver has vertices these edges and arrows $a_h : [h] \to [\tau h]$. Paths are composed in travel order: ab traverses a first. Set

$$A = kQ/(a_h a_{\phi h} : h \in H), \quad J = A_{>0}, \quad \Lambda = \varprojlim A/J^m.$$

Each vertex has two incoming and two outgoing arrows. Each arrow has precisely one permitted successor $a_{\tau h}$ and one forbidden successor $a_{\phi h}$; these are distinct because α has no fixed points. The same holds for predecessors. This is the *locally gentle* condition. Every permitted cycle repeats indefinitely, so A is infinite dimensional. We reserve "gentle" without qualification for the convention additionally imposing finite dimension.

Proposition 1.1 (The actual split order)**.** *For a permitted cycle c of length ℓ, let B_c be the hereditary cycle order in $M_\ell(k[[u_c]])$: diagonal and forward matrix entries are in $k[[u_c]]$, and entries crossing the chosen cycle cut are in $u_c k[[u_c]]$. Then Λ is the suborder of $B = \prod_c B_c$ whose two diagonal constant terms labelled $h, \alpha h$ agree. With $R = k[[u]]$ acting by $u \mapsto (u_c I)_c$, it is finite free of rank $\sum_c \ell(c)^2$; the conductor contains uB.*

Proof. In a cycle order a path of length less than ℓ is the corresponding matrix unit, multiplied by u_c exactly when it crosses the cut; another full turn multiplies it by u_c. Thus completed paths in each permitted branch give all entries of B_c. An arrow belongs to only that branch, whereas $e_{[h]}$ has diagonal entries at both occurrences $h, \alpha h$. Hence the only condition on branchwise formal coefficients is the displayed equality of diagonal constants. This proves both injectivity and surjectivity of the path-normal-form identification. Multiplication respects it: switching branches across a paired edge is precisely a forbidden pair. The quotient B/Λ is killed by u, so the submodule has the same generic rank as B. It is torsion free and finite over the DVR R, hence free of that rank. The path and u-adic topologies agree: $uA \subset J$, and if L is the longest permitted cycle, any path of length at least $L+1$ is u times a shorter nonempty path. Thus $J^{L+1} \subset uA$. This also justifies the completion used in the identification. □

The path-degree variable q below is not u: one full turn has path degree $\ell(c)$ but u-degree one. Formal, Laurent and operator-norm completions will likewise remain distinct.

Here are five explicit global inputs. Permutations list darts starting at zero; $E = |H|/2$.

Surface and Belyi map β	σ	α	$(g, \operatorname{rank}_R \Lambda)$
$\mathbb{P}^1$, $4z(1-z)$	(0)(1)	(01)	$(0, 2)$
$\mathbb{P}^1$, $(z+1)^2/(4z)$	(01)	(01)	$(0, 4)$
$y^2 = x^3 - 1$, x^3	(012)(345)	(03)(14)(25)	$(1, 18)$
$\mathbb{P}^1$, $4z^3/(1+z^3)^2$	(012)(345)	(03)(15)(24)	$(0, 18)$
$y^2 = x^3 - x$, x^2	(0123)	(02)(13)	$(1, 16)$

The first is the nodal interval: $A = k[x, y]/(xy)$ and $\Lambda = k[[x, y]]/(xy)$. The second has $A = k\langle x, y\rangle/(x^2, y^2)$, still infinite dimensional. The two theta dessins have the same two permitted three-cycles but different face cycles, respectively (051324) and (04)(13)(25). The elliptic square has one permitted and one forbidden four-cycle. For example x has pole order two on the cubic $y^2 = x^3 - 1$; x^3 has ramification $(3, 3)$ above zero, $(2, 2, 2)$ above one since $x^3 - 1 = y^2$, and (6) above infinity. The rational theta map has passport $(3, 3), (2, 2, 2), (2, 2, 2)$. These ramification computations, and the analogous direct zeros/poles in the other rows, verify cellular Belyi inputs rather than merely abstract quivers.

2 Projective resolutions, dual dessins and determinants

Use right modules $P_i = e_i A$, $S_i = P_i/P_i J$. Internal shift $M\langle n\rangle$ places its old generator in degree n. Let $\mathcal{R}_n(i)$ be the *indexing words* of length n from i with every adjacent pair forbidden, including e_i for $n = 0$. Such a word is usually zero in A; it still labels a summand.

Theorem 2.1 (Exact resolutions)**.** *There is a minimal graded projective resolution, with $P_0(i) = e_i A$ and for $n \geq 1$,*

$$P_n(i) = \bigoplus_{p = a_1 \cdots a_n \in \mathcal{R}_n(i)} e_{t(a_n)} A\langle n\rangle, \qquad d_n(x)_p = a_n x \text{ in the summand } a_1 \cdots a_{n-1}.$$

The augmentation $P_0(i) \to S_i$ is the quotient. For the completed order the identical finite-rank projective terms and formal-series maps resolve the residue simple. Ext and Tor against residue simples count forbidden words with the prescribed endpoints. In the graded case Ext has matching internal shift n.

Proof. The maps are right linear and degree zero. Adjacent differentials multiply by a forbidden pair. The augmentation kernel is the direct sum of aA over initial arrows. For any arrow a, multiplication $e_{t(a)}A \to e_{s(a)}A$ kills exactly the permitted paths beginning with its forbidden successor $f(a)$. All other basis paths survive to distinct basis paths. Thus its kernel is $f(a)A$. For a fixed prefix, different last arrows have disjoint surviving supports; there is no cancellation between summands. This proves exactness inductively. For formal series the same assertion holds coefficient by coefficient and identifies the entire closed kernel, not just its dense polynomial subspace. Terms are idempotent projectives and all entries are arrows. Consequently applying Hom or tensor with a residue simple kills every differential, proving minimality and the Ext/Tor count. In A minimality is graded; it does not require that J be its ordinary Jacobson radical. □

The two arms never terminate in this closed class. Their labelled patterns follow ϕ-cycles. An ungraded positive syzygy tail can have a shorter period by permutation of summands: the nodal interval has labelled period two but unordered tail period one; the elliptic theta has labelled period six and unordered tail period three. A simple need not be isomorphic to a positive syzygy; for the node it is R-torsion, whereas the positive syzygies are lattices.

Let $C_{ij}(q) = \sum_{d\geq 0} \dim_k(e_i A_d e_j) q^d$ and $E_{ij}(q) = \sum_{n\geq 0} (-1)^n \dim_k \operatorname{Ext}^n_A(S_i, S_j) q^n$. Degreewise Euler characteristic of the resolution gives $EC = I$: every coefficient is finite, because a summand in homological degree n starts in internal degree n. Constant terms are I, so the inverse is two-sided. Already signed $E(q)$ must not be evaluated at $-q$ again.

Proposition 2.2 (Dual dessin and cycle determinant)**.** *The complementary-relation algebra is another closed dessin algebra, with $\alpha^{\#} = \alpha$, $\sigma^{\#} = \phi^{-1}$, permitted successor ϕ, and forbidden successor τ. It exchanges graph vertices and faces and preserves genus. Its Cartan matrix satisfies $E(q) = C_{A^{\#}}(-q)$, with the opposite convention if required for Yoneda composition. For independent commuting arrow weights w_a,*

$$\det C_A(\mathbf{w}) = \frac{\prod_{f\in \mathrm{cyc}(\phi)} (1 - (-1)^{|f|} w(f))}{\prod_{c\in \mathrm{cyc}(\sigma)} (1 - w(c))}.$$

Proof. $\alpha(\sigma^{\#})^{-1} = \alpha\phi = \tau$, so the claimed relations and face data follow directly; the connectedness group is unchanged. To prove the determinant, put $S_{v,a} = w_a$ when $s(a) = v$, and $T_{a,v} = 1$ when $t(a) = v$, zero otherwise. On arrow states let $U_{a,b} = w_b$ for permitted adjacency and $V_{a,b} = w_b$ for forbidden adjacency. Then $TS = U + V$ and the path sum is $C = I + S(I - U)^{-1}T$. Sylvester's identity gives $\det C = \det(I + V)/\det(I - U)$. Each transition is a weighted permutation, whose cycle determinant is respectively $1 - (-1)^{\ell} w(c)$ and $1 - w(c)$. Everything holds in formal power series, hence as rational functions. This supplies a direct proof of the Bessenrodt–Holm formula in the closed dessin class [4]. □

The five univariate determinants are, in the order above,

$$\frac{1+q}{1-q}, \quad \frac{1+q}{1-q}, \quad \frac{1-q^6}{(1-q^3)^2}, \quad \frac{(1-q^2)^3}{(1-q^3)^2}, \quad 1.$$

The last cancellation does not imply finite global dimension: its forbidden resolution is infinite. These rational functions have cycle poles/zeros, not arithmetic prime Euler factors.

For clarity, a complex is a sequence with $d^2 = 0$, its homotopy category identifies maps differing by $dh + hd$, and $D(A)$ inverts maps inducing cohomology isomorphisms. A perfect complex is represented by a bounded complex of finite projectives. Our infinite simple resolutions compute derived Hom and tensor but are generally not perfect. The two-term presentation $[P_1 \to P_0]$ is perfect even if its kernel is nonzero. Quadratic linearity is a Koszul statement; it is not by itself an equivalence between all completed derived categories. Likewise a finite-dimensional gentle graded-curve theorem [6] cannot simply be transferred to these closed infinite orders. Here the surface walks explicitly compute the resolution endpoints: τ turns round a graph vertex and ϕ follows a face. For longer string objects all maps can be computed from the occurrence matrices; a universal geodesic-intersection classification for this completed category remains a separate theorem.

2.1 Auxiliary staircase resolutions over the order

The two filtration parameters below are not dessin arrows. Let $R = k[x, y]$ for a field k, and let $I = (g_0, \dots, g_{m-1})$, $m \geq 2$, where $g_i = x^{a_i} y^{b_i}$,

$$a_0 > \cdots > a_{m-1} = 0, \qquad 0 = b_0 < \cdots < b_{m-1}.$$

The positive outer exponents make R/I finite length. Write $R(-a, -b)$ for a free generator born in bidegree (a, b).

Proposition 2.3 (Adjacent syzygies)**.** *A minimal bigraded free resolution of $M = R/I$ is*

$$0 \longrightarrow \bigoplus_{i=0}^{m-2} R(-a_i, -b_{i+1}) \xrightarrow{d_2} \bigoplus_{i=0}^{m-1} R(-a_i, -b_i) \xrightarrow{d_1} R \longrightarrow M \longrightarrow 0,$$

where $d_1(e_i) = g_i$ and $d_2(f_i) = -y^{b_{i+1}-b_i} e_i + x^{a_i - a_{i+1}} e_{i+1}$. Deleting row i from d_2 gives determinant $(-1)^i g_i$. The Hilbert series numerator is $1 - \sum_i s^{a_i} t^{b_i} + \sum_{i=0}^{m-2} s^{a_i} t^{b_{i+1}}$.

Proof. Each column maps to zero. In a relation $\sum h_i g_i = 0$, all terms except the last contain $x^{a_{m-2}}$. Coprimality with $y^{b_{m-1}}$ implies $h_{m-1} = x^{a_{m-2}} q$. Subtract $q d_2(f_{m-2})$ from the coefficient vector; its last component vanishes. Factor $x^{a_{m-2}}$ from the remaining generators. They form a smaller staircase with the same strict inequalities, ending in a pure y-power. Induction expresses the remaining relation in the preceding adjacent columns. The base of one generator has no relations, since R is a domain. Hence $\ker d_1 = \operatorname{im} d_2$. If $d_2(q) = 0$, its first row gives $-y^{b_1} q_0 = 0$; successive rows then force all $q_i = 0$. This proves injectivity. Every differential entry has positive degree, so reduction modulo (x, y) kills the differentials and the resolution is minimal. In the row-deleted matrix the unique nonzero permutation term uses the negative y entries before row i and the positive x entries after it. Their exponents telescope to b_i, a_i, respectively, and the sign is $(-1)^i$. Finally take the degreewise Euler characteristic; the free summands have Hilbert denominator $(1 - s)(1 - t)$. □

For $I = (x^2, xy, y^2)$ the underlying sequence is also kernel checked in Lean 4 over the actual ring MvPolynomial(Fin 2, k). The verification proves the kernel/image equalities, the stated ideal image, surjective quotient augmentation, projective free terms, and vanishing of both induced Hom differentials to the constant-coefficient residue field. It uses regular multiplication on the free modules, distinct from the residue action on k. General staircases, graded minimality and the surface-order coefficient cone remain separately scoped prose arguments.

At grade $\nu \in \mathbb{N}^2$, M_ν has dimension one exactly when no generator degree is componentwise at most ν. Its comparable-grade map has rank one exactly when both endpoint grades satisfy this condition. A CW realization takes one loop and attaches a disk once along it at each generator degree.

The active indices form an interval: the x inequality selects a final interval, the y inequality an initial interval. With c active disks, the cellular map $k^c \to k$ has every entry one. Thus $\dim H_1 = \mathbf{1}_{c=0}$ and $\dim H_2 = \max(c-1, 0)$, the latter counting active adjacent syzygies. This auxiliary CW complex is not the dessin surface.

For the actual nodal order $\Lambda = k[[\xi, \eta]]/(\xi\eta)$ put $u = \xi + \eta$ and $E = \Lambda/u^2\Lambda$. The basis $(1, \xi, u, u\xi)$ of E satisfies $\xi^2 = u\xi$, $u^2 = 0$, $\eta = u - \xi$. Tensoring the resolution above with $0 \to \Lambda \xrightarrow{u^2} \Lambda \to E \to 0$ gives, over $\mathcal{B} = \Lambda[x, y]$, free ranks $(1, m+1, 2m-1, m-1)$. Indeed the scalar resolution stays exact after tensoring over the field; its degree-zero module is free over Λ on the standard monomials. Multiplication by u^2 is injective there and on all free terms, so the long exact homology sequence of its mapping cone gives positive-degree homology zero and degree-zero module $E \otimes_k M$. In contrast, as an R-module this object has ranks $4(1, m, m-1)$ in its minimal resolution. Neither finite-length E nor these parameter rings replace the infinite parent dessin algebra or its lattice category.

2.2 A monotone quantum comparison with exact finite ranks

Proposition 2.4 (One common channel)**.** *Let T be a positive trace-preserving linear map between finite complex matrix algebras. For density matrices ρ, σ, their trace distance $d(\rho, \sigma) = \frac{1}{2}\|\rho - \sigma\|_1$ satisfies $d(T\rho, T\sigma) \le d(\rho, \sigma)$. For finitely many labelled states, keep fixed vertex birth grades e_i. At (r, e) include vertices with $e_i \le e$ and edges with distance at most r, and take the flag complex. Then T gives an identity-on-labels inclusion from the input to the output bifiltration, natural in (r, e).*

Proof. Write the Hermitian difference as $A = A_+ - A_-$ with $A_\pm \ge 0$. Positivity, the trace norm triangle inequality and trace preservation give $\|TA\|_1 \le \|TA_+\|_1 + \|TA_-\|_1 = \operatorname{tr} A_+ + \operatorname{tr} A_- = \|A\|_1$. Thus every input edge is an output edge; the fixed births preserve vertices, and all cliques are preserved. Increasing either parameter preserves the simplices, and every map is the same label inclusion, so the squares commute. Complete positivity is sufficient but is not needed for this norm argument. □

For a finite inclusion $K \subset L$, work over a field and embed a basis Z_K of source one-cycles into the target edge space. If B_L is the matrix of target two-boundaries, the induced H_1 rank is

$$\rho_1(K, L) = \operatorname{rank}[B_L \; Z_K] - \operatorname{rank} B_L.$$

Its image is $(Z_K + \operatorname{im} B_L)/\operatorname{im} B_L$; source boundaries already map into target boundaries. This proves the formula, including empty complexes. Computing a flag two-skeleton suffices for H_0, H_1, but does not compute the full flag complex's H_2. Integer ceilings of scaled exact distances preserve the inclusion. For numerical distances, frozen integer labels define an exact combinatorial diagram only after their inequality is checked; this is not an interval certificate of the underlying quantum distances. None of these finite ranks identifies a completed arithmetic determinant or proves GRH.

2.3 A correct DG model of the infinite node

Use cohomological degrees and an uncurved differential of degree $+1$. An odd element ε with $d\varepsilon = 1$ contracts the underlying k-complex of every unital left DG module: $h(m) = \varepsilon m$ satisfies $dh + hd = 1$ by the graded Leibniz rule. Thus this example has zero cohomology, contrary to the original ground-field-resolution claim. An isomorphism on H^0 alone is also insufficient for a DG quasi-isomorphism; all degrees must be checked.

Proposition 2.5 (Nodal Koszul model)**.** *Let k be a field, $R = k[\xi, \eta]$ and $f = \xi\eta$ and $K = R \oplus R\varepsilon$, with $|\varepsilon| = -1$, $\varepsilon^2 = 0$, $dR = 0$ and $d\varepsilon = f$. The augmentation $K \to A = R/(\xi\eta)$, $a + b\varepsilon \mapsto [a]$, is*

a DG-algebra quasi-isomorphism onto the actual infinite nodal dessin algebra. Branch rescaling $\xi \mapsto q\xi$, $\eta \mapsto q^{-1}\eta$, $\varepsilon \mapsto \varepsilon$ for $q \in k^\times$ lifts to a DG automorphism.

Proof. Multiplication is $(a + b\varepsilon)(c + e\varepsilon) = ac + (ae + bc)\varepsilon$. The differential kills degree zero and sends $b\varepsilon$ to fb. Its square is zero. For two odd inputs the Leibniz right side is $fbc\varepsilon - bfc\varepsilon = 0$; the other homogeneous cases follow by R-linearity. Multiplication by f is injective in the polynomial domain. Hence $H^{-1}(K) = 0$, $H^0(K) = R/(f)$, and all other groups vanish. The augmentation preserves products and kills boundaries. Rescaling fixes f, so its coefficientwise extension commutes with d and has inverse obtained by replacing q with q^{-1}. This is a one-relation Koszul construction [78, 79]. □

Replacing R by $k[[\xi, \eta]]$ gives the completed order: setting each variable separately to zero maps onto the branch pullback, and the kernel consists precisely of mixed monomials, hence equals $(\xi\eta)$. Rescaling is generally semilinear over the coefficient ring, rather than linear over the fixed diagonal DVR $k[[u]]$, $u = \xi + \eta$. It preserves that DVR as a subring exactly when $q^2 = 1$, and fixes it pointwise exactly when $q = 1$. No Hilbert norm or arithmetic spectrum follows.

The corresponding Lean file checks the actual exterior-line algebra, differential, sign-Leibniz, homotopy and polynomial ambient-resolution identities. Full graded/category/completion objects remain separately scoped. The two theta dessins likewise require endpoint-sensitive Ext: self Ext^1 is zero in both; for the elliptic theta self Ext^3 has dimension two, while for the spherical theta positive even self Ext has dimension two. Outdegree alone does not count self extensions. Minimal A_∞ models use additional graded data [77, §3.3]; m_3 is not automatically a physical curvature.

3 Determinant semi-invariants and Schur constructions

Assume k algebraically closed of characteristic zero for invariant-ring statements. A finite-dimensional representation of the *infinite* A has spaces N_i, arrow matrices $N(a)$ satisfying the monomial relations, and base-change group $G = \prod_i \mathrm{GL}(N_i)$. With column vectors, $N(a) \mapsto g_{t(a)} N(a) g_{s(a)}^{-1}$. Let $P_0 = \bigoplus P_i^{a_i}$, $P_1 = \bigoplus P_i^{b_i}$, and $f : P_1 \to P_0$ be a finite path-matrix map. When $\sum a_i \dim N_i = \sum b_i \dim N_i$, define

$$c_f(N) = \det D_f(N), \qquad D_f(N) = \mathrm{Hom}_A(f, N).$$

It is intrinsically a determinant-line section; chosen bases give this scalar.

Proposition 3.1 (Weights and exact homological scope)**.** $c_f(gN) = \prod_i \det(g_i)^{b_i - a_i} c_f(N)$. *Restricted determinants of this form span all semi-invariants on a closed G-stable relation or nilpotence locus. Set $M = \mathrm{coker}\, f$. On the square locus,*

$$c_f(N) \neq 0 \quad \Longleftrightarrow \quad \mathrm{Hom}(M, N) = 0 \quad \Longleftrightarrow \quad \mathrm{Hom}(M, N) = \mathrm{Ext}^1(M, N) = 0.$$

If f is injective, $\mathrm{coker}\, D_f = \mathrm{Ext}^1(M, N)$ *and higher Ext vanish. Without injectivity neither that cokernel equality nor higher vanishing follows.*

Proof. Naturality gives $D_f(gN) = G_1 D_f(N) G_0^{-1}$, where G_j repeats the relevant g_i. Take determinants. The contragredient coordinate-ring action has the inverse character. Schofield–van den Bergh's spanning theorem allows oriented cycles [36, Theorem2.3]. Exactness of invariants for reductive G in characteristic zero lifts a fixed-weight function from any closed stable locus to the ambient representation space; restrict the spanning determinants. Injectivity of the path map need not survive that restriction. Finally $\ker D_f = \mathrm{Hom}(M, N)$. Writing $K = \mathrm{im}\, f$, the exact sequence with projective P_0 expresses $\mathrm{Ext}^1(M, N)$ as $\mathrm{Hom}(K, N)/\,\mathrm{im}\,\mathrm{Hom}(P_0, N)$; the epimorphism $P_1 \to K$

embeds this quotient in $\operatorname{coker} D_f$. A square finite matrix with zero kernel has zero cokernel. If f is injective it is the actual length-one resolution, giving the stronger assertion. □

For the node, $f = \times x$, $M = \Lambda/(x)$, $N = \Lambda/(x, y^2)$ give $\dim \operatorname{coker} D_f = 2$ but $\dim \operatorname{Ext}^1(M, N) = 1$. For the elliptic square name arrows so that $a, b : 0 \to 1$, $c, d : 1 \to 0$, permitted cycle *acbd* and forbidden cycle *adbc*. With $f = a$ and $N(a) = 1$, $N(b) = N(c) = N(d) = 0$, its Hom matrix is [1], yet the subsequent forbidden resolution gives $\operatorname{Ext}^{2,3}(M, N) = k$. Thus nonvanishing is not all-degree derived orthogonality.

Proposition 3.2 (What completion does to invariants)**.** *A continuous d-dimensional representation of Λ kills J^d. On the reduced nilpotent representation variety of fixed dimension vector every invariant regular function is constant. Nonzero-weight semi-invariants need not be constant.*

Proof. Continuity with discrete finite-dimensional target kills some J^m; its Loewy filtration has at most d nonzero strict steps. Choose vertexwise complements to this filtration. An arrow sends a layer p into layers $q \geq p+1$. Scaling layer p by z^p conjugates each component by z^{q-p}. At $z = 0$ the representation tends to the zero-arrow representation. Every invariant therefore has the same value at all points; reducedness and the Nullstellensatz give the ring assertion. In the square example the scalar $N(a)$ varies on a nilpotent locus and has weight g_1/g_0, so semi-invariants survive. Formal entries in a projective map are evaluated on this locus by truncating modulo J^d, a legitimate finite polynomial operation. □

On the node, $u = x+y$ is regular and A is free of rank two over $k[u]$. The polynomial presentation $\times(u-\lambda)$ for $\lambda \neq 0$ is injective and its determinant on N is $\det(u_N - \lambda I)$, the characteristic polynomial of u_N up to the sign $(-1)^d$. But $u-\lambda$ is a unit in Λ, with its geometric-series inverse, so those modules do not extend continuously to the completion. An R-free lattice and its u-nilpotent quotients are different categories. On a lattice a separately specified endomorphism F has a meaningful relative characteristic polynomial over R; the scalar base action uI alone does not create Frobenius.

3.1 Young tableaux and a cyclic representation stratum

For a nonempty partition $\lambda \vdash m$, choose a numbering T of its boxes. On $V^{\otimes m}$ let $Y_T = R_T C_T$, the unnormalized row sum followed by the signed column sum. The classical identity $Y_T^2 = h_\lambda Y_T$, where h_λ is the hook product, makes Y_T/h_λ an idempotent in characteristic zero. Its image is $S_\lambda(V)$, one Schur copy, rather than the whole symmetric-group isotypic component [31, 32]. Its character is s_λ, computed by semistandard tableaux (weak rows, strict columns), and

$$\dim S_\lambda(k^d) = \prod_{(i,j)\in\lambda} \frac{d+j-i}{h_{ij}}.$$

Since $f^{\otimes m}$ commutes with the place permutations, it induces $S_\lambda(f)$ and respects composition and zero maps. Applying this at every vertex and arrow therefore gives a genuine representation of the monomial dessin algebra: $S_\lambda(N(b))S_\lambda(N(a)) = S_\lambda(0) = 0$ for a forbidden pair. The empty partition does not have this property: it sends zero arrow maps to the identity of k.

The Littlewood–Richardson rule gives natural intertwiners

$$S_\lambda(V) \otimes S_\mu(V) \cong \bigoplus_\nu S_\nu(V)^{\oplus c^\nu_{\lambda\mu}}.$$

Here $c^\nu_{\lambda\mu}$ counts semistandard tableaux of skew shape ν/λ and content μ whose row-reading word is Yamanouchi. Naturality makes the same decompositions intertwine all the arrows. The tensor

operation is vertexwise with synchronous arrow maps, not $\otimes_A$ and not a Hopf-algebra tensor structure. There is no automatic tensor unit preserving these zero relations. These functors are not additive or exact; they cannot simply be applied degreewise to a projective resolution. They also need not preserve the fixed R-action on lattices.

In the elliptic theta set one of the two permitted three-cycle branches to zero. The other has arbitrary maps $X_1 : V_0 \to V_2$, $X_2 : V_2 \to V_1$, $X_3 : V_1 \to V_0$, with all spaces of dimension d. This is a closed representation stratum of our infinite dessin algebra. Put $M = X_3X_2X_1$.

Theorem 3.3 (Cyclic invariants and a local positive kernel)**.** *The invariant ring on this stratum is $k[c_1(M), \dots, c_d(M)]$, where c_j are characteristic coefficients. Its total-arrow-degree Hilbert series is $\prod_{j=1}^{d}(1 - t^{3j})^{-1}$. Common multidegree (m, m, m) has basis $\chi_\lambda(M) = \operatorname{tr} S_\lambda(M)$, $\lambda \vdash m$, $\ell(\lambda) \le d$; other multidegrees have no invariants. Products obey the Littlewood–Richardson rule. Over $\mathbb{C}$, if $\|M_i\| \le R_0$ and $0 \le r$ and $rR_0^2 < 1$, the kernel*

$$K_r(M_i, M_j) = \det(I - rM_i \otimes \overline{M_j})^{-1} = \sum_\lambda r^{|\lambda|} \chi_\lambda(M_i)\overline{\chi_\lambda(M_j)}$$

is absolutely convergent and positive semidefinite.

Proof. Cycle-trace generation for quivers, including relation quotients, is the Le Bruyn–Procesi theorem [29, 30]. Every cycle here is a power of the full three-cycle, up to cyclic rotation. In characteristic zero Newton identities give the generators c_j. They are algebraically independent: take $X_1 = X_2 = I$ and arbitrary X_3. Scalar changes at the three vertices force equal arrow multidegrees. Restriction to that same section identifies degree (m, m, m) with degree-m symmetric polynomials in eigenvalues, whose Schur basis and product rule give the assertion. For the kernel use the ordinary Cauchy identity, first as a formal identity in eigenvalues. On Schur modules the operator norm of $S_\lambda(M)$ is at most $R_0^{|\lambda|}$, so its trace is bounded by $\dim S_\lambda(\mathbb{C}^d) R_0^{|\lambda|}$. The dimension Cauchy identity bounds the absolute series by $(1 - rR_0^2)^{-d^2}$. Each summand is a rank-one Gram kernel, proving positivity by absolute summation. □

The term for the empty partition is a constant scalar feature, not an allowed module-valued tensor unit. In dimension two,

$$(\operatorname{tr} M)^2 = \operatorname{tr} \operatorname{Sym}^2 M + \det M, \qquad \operatorname{tr} \operatorname{Sym}^2 M = \tfrac{1}{2}((\operatorname{tr} M)^2 + \operatorname{tr} M^2).$$

More generally $\det(I - zM) = \sum_j (-z)^j \operatorname{tr} \bigwedge^j M$ and its reciprocal is $\sum_m z^m \operatorname{tr} \operatorname{Sym}^m M$. These are actual determinant/Schur connections, not identifications with an arithmetic L-function. On continuous finite-length Λ-modules the monodromy is nilpotent and all positive-degree characters vanish. The displayed Cauchy positivity is local representation-theoretic positivity, not the global Weil form.

4 Lattice almost-split sequences and genuine diamond quotients

For the interval dessin let $X = (x)$, $Y = (y)$ in $\Lambda = k[[x, y]]/(xy)$, $R = k[[u]]$, $u = x + y$. A lattice is a finitely generated Λ-module free over R.

Theorem 4.1 (The nodal lattice calculation)**.** *Every lattice is $\Lambda^r \oplus X^a \oplus Y^b$. The almost-split sequences in this lattice category are*

$$0 \to Y \hookrightarrow \Lambda \xrightarrow{\times x} X \to 0, \qquad 0 \to X \hookrightarrow \Lambda \xrightarrow{\times y} Y \to 0.$$

In particular $\tau X = Y$, $\tau Y = X$. Multiplication by u between two copies of either row, followed by reduction modulo u, gives an actual nonzero exact three-by-three diagram.

Proof. Normalize M *inside its total quotient module*: $\overline{M} = BM$, $B = k[[x]] \times k[[y]]$. The conductor gives $u\overline{M} \subset M \subset \overline{M}$. Both branch projections of $W = M/u\overline{M} \subset k^p \oplus k^q$ are onto. Intersect W with the two factors; its remaining quotient identifies their two residual quotients. Separate bases for these pure factors and the common quotient put W into pure first lines, pure second lines and diagonal pairs. Lift branch bases over the DVRs. These are respectively the summands X, Y, Λ. Their endomorphism rings are local.

Exactness of the first sequence is $\operatorname{ann} x = Y$. It does not split: a map $X \to \Lambda$ lands in $\operatorname{ann} y = X$, and its composite with $\times x$ is a nonunit in $\operatorname{End} X = k[[x]]$. Every map $\Lambda \to X$ factors through this epimorphism, $\operatorname{Hom}(Y, X) = 0$, and every nonunit endomorphism of X is divisible by x and factors through $X \hookrightarrow \Lambda \to X$. Classification handles all finite sums; a unit X component would be a retraction. This is right almost splitness. The dual argument for nonsections out of Y proves left almost splitness. Endomorphisms fixing either structure map have constant term one, hence are units, proving minimality. Swap x, y for the other sequence. Finally all three terms are R-free. Multiplication by u is injective and reduction preserves exactness, proving the diagram assertion. □

The lattice AR quiver has arrows $Y \to \Lambda \to X$ and $X \to \Lambda \to Y$; the mesh composites are zero. Its projective vertex Λ is not the endpoint of another nonsplit AR sequence. AR meshes do not mean that all rows and columns of any drawn grid are short exact.

For a second, explicit family fix $n \geq 3$ and the rational Belyi map

$$\beta_n(z) = \frac{1}{1 - T_n(z)^2} = \frac{1}{(1 - z^2)U_{n-1}(z)^2},$$

where T_n, U_{n-1} are Chebyshev polynomials. It has passport $(2n), (2^n), (2^{n-1}, 1, 1)$. Its contracted dual graph is a path, identifying its genus-zero dessin. The associated infinite algebra has vertices $1, \ldots, n$, endpoint loops l, r, opposite arrows $a_i : i \to i + 1$, $b_i : i + 1 \to i$, and precisely the relations $l^2, r^2, a_i b_i, b_i a_i$. Its single permitted cycle is $l a_1 \cdots a_{n-1} r b_{n-1} \cdots b_1$; the order rank is $(2n)^2$.

Retain one arrow at each edge in an orientation Ω, and kill its opposite. The quotient is exactly the string algebra $H_n(\Omega)$ of Huang–Lin–Su [46]; the killed-arrow ideal contains J^{n+2}, so it is also an actual quotient of Λ_n. Inflation identifies its module category with the full abelian quotient-annihilator subcategory $\mathcal{C}_\Omega$ of Λ_n-modules. This is generally not extension closed in the whole order category. The imported AR property is intrinsic to $\mathcal{C}_\Omega$, not automatically an ambient lattice or finite-length AR property. The extended edition gives nine explicit string modules, all twelve maps and the four exact meshes from the supplied diamond resource, now situated in this actual dessin quotient.

The other supplied resources also have a precise home. For the interval order the subcategory annihilated by J^2 is $\operatorname{Mod} B_0$, $B_0 = \Lambda/J^2$. This is a domestic string quotient, not itself a gentle algebra. Inflation is full and preserves purity, and $\operatorname{Zg} B_0$ is a closed subspace of $\operatorname{Zg} \Lambda$. Ringel's infinite-word/pure-injective program and its proof by Puninski–Prest apply there [43, 44]; the isolated band xy^{-1} gives Krull–Gabriel and relative Cantor–Bendixson dimension two by [45, Theorem10.19]. This is not an ambient rank calculation for all Λ-modules. The parameter in $x = J_2, y = \lambda J_2$ survives even though all cycle traces vanish; the pp condition $\exists v\ (z = xv \ \wedge \ (y - \lambda x)v = 0)$ detects it. These modules are u-torsion and are not nonzero order lattices. Infinite-string and locally finite AR theorems must retain their distinct finiteness hypotheses; a surviving permitted cycle has infinitely many paths at a vertex.

4.1 Polynomial strings, completed lattices and the Fukaya category

Crawley–Boevey's theorem [38, Theorems1.1–1.4] applies directly to the infinite polynomial dessin algebra: its finitely controlled modules decompose into strings and finite-dimensional bands; its

pointwise artinian modules additionally allow primitive injective bands. For our finite quiver, these conditions mean finitely generated and artinian respectively. The words may be infinite, but their modules have algebraic direct-sum bases. This is not a classification of every module or of arbitrary formal completions.

In fact no nonzero completed-order lattice, restricted to A, meets either finiteness condition. The polynomial A is finite over $k[u]$: remove full turns from sufficiently long paths as in Proposition 1.1. If an R-free lattice were finitely generated over A, it would be finite torsion free, hence free, over the PID $k[u]$. Multiplication by $1-u$ would then fail to be onto, whereas it is a unit over $R = k[[u]]$. Also $M \supsetneq uM \supsetneq u^2M \supsetneq \cdots$ excludes artinianity. For a concrete indecomposable outside the ordinary string/band list take $k[[x]]$ with $y = 0$: its endomorphisms are multiplication by formal series, so its endomorphism ring is local. Its x is injective and nonsurjective, while $1-x$ is invertible; these properties exclude every pure-x ordinary string or band. The extended edition gives the full word argument.

The Fukaya connection requires equally precise categories. Lekili–Polishchuk explicitly allow infinite-dimensional locally gentle algebras, but their equivalence is for *perfect dg modules* of a homologically smooth algebra, with specified grading, line field and stops [66, Definition3.1.1, Lemma3.1.3, Theorem3.2.2]. Every nonempty closed dessin here has both permitted and forbidden cycles, so its polynomial algebra is neither proper nor smooth by their criterion. This does not exclude appropriate singular or completed variants; it prevents a direct use of that particular theorem. Infinite word, graded curve and geodesic are not interchangeable without a surface realization and boundary/grading data.

There is a genuine positive example inside our setting: the polynomial interval quotient $A/(y) = k[x]$ is still infinite dimensional, is smooth, and has no forbidden cycle. Its perfect category has the partially wrapped annulus model with one stop total. In degree zero $\mathrm{Perf}(k[x]) = D^b(\mathrm{mod}_{\mathrm{fg}}\, k[x])$. Inflation of this quotient is fully faithful on ordinary modules, but not on derived categories: $\mathrm{Ext}^2_A(k[x], k[x]) = k$, whereas $\mathrm{Ext}^2_{k[x]}(k[x], k[x]) = 0$. The former follows from the alternating y, x resolution. Thus this quotient bridge does not identify the whole closed dessin's perfect category with the annulus category.

Finally, Leavitt path algebras *can* fail invariant basis number, but need not. The permitted-state Laurent algebras here have finite-dimensional matrix quotients, and therefore invariant basis number. The polynomial and completed dessin algebras have their vertex-simple quotients, giving the same conclusion. The complex Leavitt algebra of G_α below is a dense *-subalgebra of its graph C*-algebra, on that specified graph; it is not the original gentle algebra with ordinary CK1 imposed.

5 Leavitt relations, the constrained Fock algebra and spectrum

Work now over $\mathbb{C}$. Ordinary Leavitt and graph C* constructions [47, 48] cannot simply impose our original forbidden products: if $ab = 0$ and CK1 says $a^*a = p_{t(a)}$, then $0 = a^*ab = b$. Every arrow is a forbidden successor, hence every arrow and then every vertex is zero. CK2 is not needed. The ordinary graph-Toeplitz CK1 setting has the same collapse.

The valid permitted-arrow-state graph instead consists of the disjoint directed σ-cycles. Its Leavitt algebra is $\bigoplus_c M_{|c|}(k[z, z^{-1}])$, and its graph C* algebra is $\bigoplus_c M_{|c|}(C(\mathbb{T}))$. These forget the α-gluing; both theta dessins give the same answer. They are not the completed original order. Nor is the ordinary Toeplitz algebra of a cycle of length $\ell > 1$ equal to $M_\ell(\mathcal{T})$: their K_0 ranks are ℓ and one respectively [50].

Let $\mathcal{H} = \ell^2(\mathcal{P})$ for all permitted finite paths, including vacua e_v. Define $L_a\delta_p = \delta_{ap}$ when the prefix is permitted and composable, and zero otherwise. Let P_v project onto paths starting at v,

$Q_a = L_a L_a^*$ and v_v onto the vacuum. Then

$$L_a L_{\phi(a)} = 0, \quad L_a^* L_a = P_{t(a)} - Q_{\phi(a)} = v_{t(a)} + Q_{\tau(a)}, \quad P_v - \sum_{s(a)=v} Q_a = v_v.$$

These follow on basis vectors. The different initial projection avoids CK1 collapse. Write $\mathcal{T}_A = C^*(P_v, L_a)$, a constrained-path Fock C* algebra in the monomial/subproduct setting [51, 52]. The polynomial A acts faithfully: applying a finite path sum to the sum of vacua recovers its coefficients. This representation need not extend to all formal Λ series: $\sum_{m\geq 0} c^m$ along a cycle would send its vacuum to a nonsquare-summable sum.

Theorem 5.1 (Compact extension and incidence index)**.** *Let $\mathcal{H}_v$ be the terminal-path sector. There is an exact sequence*

$$0 \longrightarrow \bigoplus_{v\in Q_0} \mathcal{K}(\mathcal{H}_v) \longrightarrow \mathcal{T}_A \longrightarrow \bigoplus_{c\in \mathrm{cyc}(\sigma)} M_{|c|}(C(\mathbb{T})) \longrightarrow 0.$$

With $\delta[u] = [1 - W^*W] - [1 - WW^*]$, its boundary map is $\delta(c) = -\sum_{h\in c} e_{[h]}$, minus the unsigned incidence map B_Γ of the cellular graph. Thus, for its E edges,

	$K_0(\mathcal{T}_A)$	$K_1(\mathcal{T}_A)$
Γ *bipartite*	$\mathbb{Z}^{E+1}$	$\mathbb{Z}$
Γ *nonbipartite*	$\mathbb{Z}^E \oplus \mathbb{Z}/2$	0.

Proof. All prefixes preserve the terminal sectors. For paths p, q ending at v, $L_p v_v L_q^* = |\delta_p\rangle\langle\delta_q|$. Therefore the vacuum ideal is exactly the displayed sum of compacts: it contains these matrix units and that sum is itself an ideal. In the quotient, L_a has initial projection $Q_{\tau(a)}$ and range Q_a, a graph CK family for the permitted-state cycles. The Q_a have infinite rank and nonzero quotient images; the length gauge action descends. Gauge-invariant uniqueness [48, Theorem2.1] identifies the quotient as claimed.

For a cycle c based at arrow a, lift its corner unitary by $W = L_c Q_a + (1 - Q_a)$. Then $W^*W = 1$ and $1 - WW^* = Q_a - L_c Q_a L_c^*$ projects onto the first $|c|$ paths starting with a. Their terminal vertices count exactly its incident darts, including twice for a loop. This proves the index formula. The six-term exact sequence gives $K_1 = \ker B_\Gamma$ and $0 \to \operatorname{coker} B_\Gamma \to K_0 \to \mathbb{Z}^{\#\,\mathrm{cyc}\,\sigma} \to 0$. The free last term splits abstractly. Spanning-tree integer elimination gives Smith entries 1 repeated $V - 1$ times and one zero in the bipartite case; an odd cycle instead adds the entry two. This proves the groups. □

The nodal interval has $K_0 = \mathbb{Z}^2$, $K_1 = \mathbb{Z}$; the single-loop sphere has $\mathbb{Z} \oplus \mathbb{Z}/2$, zero. Both theta dessins have $\mathbb{Z}^4$, $\mathbb{Z}$. Coarse K-theory does not recover genus or the arithmetic factors.

5.1 A graph presentation and an actual pullback

The Fock algebra itself has a graph presentation, on a different graph from the original gentle quiver. Let G_α have vertices q_h for $h \in H$ and sinks b_e for $e \in Q_0$, and edges

$$s_h : q_h \to q_{\tau h}, \qquad t_h : q_h \to b_{[\tau h]}.$$

Proposition 5.2. *$C^*(G_\alpha) \cong \mathcal{T}_A$, by $q_h \mapsto Q_{a_h}$, $b_e \mapsto v_e$, $s_h \mapsto L_{a_h} Q_{a_{\tau h}}$ and $t_h \mapsto L_{a_h} v_{[\tau h]}$. The integer graph relation at h is $q_h - q_{\tau h} - b_{[\tau h]}$. Eliminating these relations recovers $K_1 = \ker B_\Gamma$ and $K_0 \cong \operatorname{coker} B_\Gamma \oplus \mathbb{Z}^{\#\,\mathrm{cyc}\,\sigma}$.*

Proof. The two edges at q_h have orthogonal initial pieces and orthogonal ranges summing to Q_{a_h}, by the Fock initial-projection identity. All vertex projections are nonzero. The length gauge acts with degree one on both edge families, so gauge-invariant uniqueness applies. They generate since $L_{a_h} = s_h + t_h$ and $P_e = v_e + \sum_{[h]=e} Q_{a_h}$. In the integer matrix a kernel coefficient is constant around each permitted cycle; its sink equations are exactly $B_\Gamma c = 0$. In the cokernel, eliminate all but one state per cycle; its remaining relation is the sum of its incident sink generators. The remaining cycle generators are free, proving the stated groups. □

For the nodal interval, G_α has two loop vertices, each with an exit to the common vacuum sink. Chirvasitu–Hajac–Tobolski's actual pullback theorem [65, Theorem3.4] gives

$$\{T \in \mathcal{T}_A : \pi(T) \in \mathbb{C}^2\} \cong C^*(E_1), \qquad \mathbb{C}^2 \hookrightarrow C(\mathbb{T})^2 \text{ by constants},$$

where E_1 has the same three vertices and countably many arrows from each loop vertex to the sink, with no loops. The arrow numbered $n \geq 0$ maps to loop^n followed by exit. This algebra is AF, with $K_0 = \mathbb{Z}^3$, $K_1 = 0$. The extended edition verifies every hypothesis. The theorem does not turn arbitrary cycle-vertex identifications into pullbacks: its first graph is acyclic and its path functor must cover all pointed paths. Its sink-extension Theorem4.3 has further injection and path-coverage requirements.

5.2 A separate diagonal-fiber C* completion retaining the pairing

The formal order's residue gluing cannot be copied by taking diagonals of arbitrary full matrices: $\operatorname{diag}(E_{12}E_{21}) \neq \operatorname{diag}(E_{12})\operatorname{diag}(E_{21})$. For $\ell > 1$ there is no nonzero *-homomorphism from $M_\ell(C(\mathbb{T}))$ to a commutative C*-algebra, as restriction to its constant matrix units shows. Restrict the distinguished fiber first. Define

$$D_\ell = \{f \in C(\mathbb{T}, M_\ell(\mathbb{C})) : f(1) \text{ is diagonal}\}, \qquad \mathcal{D}_H = \Big(\prod_c D_{|c|}\Big) \underset{\mathbb{C}^H}{\times} \mathbb{C}^{Q_0},$$

where the first map evaluates diagonal entries at one and the second repeats λ_e at the two darts paired by α.

Theorem 5.3 (A faithful polynomial pullback model)**.** *There is a split exact sequence*

$$0 \to \bigoplus_c C_0(\mathbb{T} \setminus \{1\}, M_{|c|}) \to \mathcal{D}_H \to \mathbb{C}^{Q_0} \to 0,$$

so $K_0(\mathcal{D}_H) = \mathbb{Z}^{|Q_0|}$ *and* $K_1(\mathcal{D}_H) = \mathbb{Z}^{\#\,\mathrm{cyc}\,\sigma}$*. The polynomial dessin algebra embeds faithfully and generates this C*-algebra with its adjoints. This embedding does not extend to a unital algebra map from all of* Λ*.*

Proof. The section consists of the constant paired diagonal projections. The kernel is the displayed vanishing fiber, a sum of matrix suspensions, giving the groups by split exactness and suspension. For an explicit embedding set $g(z) = |1 - z|^2/4$, choose one cut arrow per permitted cycle, and send a_h to $g(z)z^{\epsilon_h}E_{h,\tau h}$ in its branch, where $\epsilon_h = 1$ at the cut and zero otherwise. Send e_e to the constant paired diagonal projection. Every forbidden product is zero. A full cycle at h maps to $w_c(z)E_{hh}$, $w_c = g^{|c|}z$. This scalar function is injective on the circle: away from its unique zero at one its argument recovers z. Continuous functional calculus therefore supplies every $f(z)E_{hh}$ with $f(1) = 0$. Multiplying by path images and approximating division away from one supplies every vanishing off-diagonal entry. Together with the constant paired diagonals these generate $\mathcal{D}_H$.

Path normal forms in a fixed entry are a nonzero prefactor times a polynomial in w_c; its infinite range proves independence. Evaluation at one first separates the zero-length coefficients. Finally for $z_0 \neq 1$, $1 - w_c(z_0)^{-1}c_h$ is a unit in the formal order, but its proposed image is singular at z_0. A unital map preserves units, proving nonextension. □

This is the C* algebra generated by a faithful polynomial representation and its adjoints, distinct from the completed order and from $\mathcal{T}_A$. The norm closure of the polynomial image alone need not contain those adjoints. Their K groups already differ. The pairing belongs to the labelled presentation; an unmarked isomorphism class need not recover the ribbon ordering. No arithmetic descent, Frobenius action or completed-L positive determinant is inferred from this additional valid gluing model.

Proposition 5.4 (The degree spectral triple)**.** *Define $N\delta_p = |p|\delta_p$ on $\{\xi : \sum_p |p|^2|\xi_p|^2 < \infty\}$. With the polynomial creation/annihilation *-algebra this is an ungraded spectral triple, p-summable exactly for $p > 1$. With N_+^{-s} zero on the kernel,*

$$\operatorname{Tr} e^{-tN} = E + \frac{2E}{e^t - 1}, \quad \operatorname{Tr} N_+^{-s} = 2E\zeta(s), \quad \operatorname{Tr}\big(\chi(N)N_+^{-s}\big) = 2EL(s,\chi)$$

for $t > 0$ and $\operatorname{Re} s > 1$ in the last two formulas, where χ is an externally chosen Dirichlet character.

Proof. There are E vacua and $2E$ paths of every positive length: each starting arrow has a unique continuation. The real diagonal operator is self-adjoint on the stated maximal domain; finite supports are a core and its resolvent is compact. Prefixes preserve the domain and $[N, L_a] = L_a$, $[N, L_a^*] = -L_a^*$, so polynomial commutators are bounded. The trace formulas and summability reduce to the corresponding scalar series. The character multiplier is bounded; membership in $\mathcal{T}_A$ is not asserted. □

Length parity anticommutes with the creators but commutes with N, so it is not an even-spectral-triple grading. The ordinary odd Fredholm class of this positive N is trivial: $N(1+N^2)^{-1/2} - I$ is compact. The spectrum depends only on E; zeros of the scalar zeta trace are not eigenvalues of N.

6 Arithmetic attachment, descent and the remaining comparison

The compact ideal gives a precise home for an arithmetic determinant, with no claim that the arithmetic is derived from the dessin. Fix one terminal sector. Its two paths of each length $n \geq 1$ give orthonormal vectors $\xi_n = (\delta_{p_{n,1}} + \delta_{p_{n,2}})/\sqrt{2}$ and an isometry $U : \ell^2(\mathbb{N}_{\geq 1}) \to \mathcal{H}_v$.

For a nonprincipal character χ put $A_n = \sum_{j \leq n} \chi(j)$, $B_0 = \sup |A_n| \leq q_\chi$, and, on $2^k \leq n < 2^{k+1}$, set

$$w_n = \sqrt{n}(k+1), \quad u_n = A_n/w_n, \quad v_n(s) = w_n(n^{-s} - (n+1)^{-s}), \quad \varphi_s(z) = \sum_n v_n(s)z_n - z_1.$$

Here $u_1 = 1$. For $\operatorname{Re} s > 0$, $u, v(s) \in \ell^2$ and the pairing is complex linear, with no conjugation of s. Define

$$\widetilde{K_s} = U(u \otimes \varphi_s)U^*, \qquad \det_F(I + \widetilde{K_s}) = L(s,\chi).$$

Indeed dyadic blocks give $\|u\|^2 \leq 2B_0^2$, and on a compact set $\operatorname{Re} s \geq \sigma_0 > 0$, $|s| \leq M$, the squared v tail is bounded by $M^2 \sum_{k \geq J}(k+1)^2 2^{-2\sigma_0 k}$. Thus the rank-one family is trace-norm holomorphic. Its determinant is $1 + \varphi_s(u)$, equal by Abel summation to L. Replacing v by $P_N v$ and adding $A_N(N+1)^{-s}e_1^*$ to the functional gives exactly every raw partial sum S_N, with compact-uniform

trace-norm convergence; $\|UXU^*\|_1 = \|X\|_1$. For primitive χ, completion replaces the functional by $g(s)\sum v_n(s)z_n - z_1$, with the standard nonzero gamma factor g, rather than multiplying the whole old functional. The extended edition supplies the quantitative estimates.

This inserts a scalar encoding into one actual dessin compact ideal; it is not known to be A-linear, a derived chain map or Frobenius compatible. A common fixed positive metric cannot make this same fixed-range family symmetric for all critical heights: symmetry forces its functionals to be proportional, whereas the coefficient ratio at heights $0, t \neq 0$ is asymptotic to $(\sigma + it)n^{-it}/\sigma$, with two opposite subsequential limits. Different positive realizations remain possible.

Proposition 6.1 (Complex dessin data do not determine arithmetic descent)**.** *The rational Belyi pairs $E_d : y^2 = x^3 - d$, $\beta_d = x^3/d$, for $d = 1, 2$, are isomorphic over $\mathbb{C}$ but have distinct Frobenius polynomials at the good prime seven: $1 - 4T + 7T^2$ and $1 - T + 7T^2$.*

Proof. Choose $a^3 = d$, $b^2 = d$. The projective isomorphism $(x, y) \mapsto (ax, by)$ from E_1 to E_d preserves the Belyi map. The degree-six clean passports are the elliptic theta passport already computed. Directly counting y for $x = 0, \ldots, 6$ gives respectively $(0, 1, 1, 0, 1, 0, 0)$ and $(0, 0, 0, 2, 0, 2, 2)$. Including infinity yields four and seven points, hence $a_7 = 8 - \#E_d(\mathbb{F}_7) = 4, 1$. Both discriminants $-432d^2$ are nonzero modulo seven. □

Any invariant of the complex dessin or its undecorated order/Fock algebra alone is the same for these two inputs, so cannot recover both factors without descent data. This does not obstruct a theory including a specified rational model and compatible Galois action. Elliptic H^1 has weight one and unshifted center $s = 1$, unlike Dirichlet center $1/2$; the example is not a counterexample to GRH.

6.1 A constructive arithmetic benchmark on the theta dessin

The preceding twists can be distinguished constructively once their finite-field descent data are retained. This closes the elementary local-factor computation, not the global positive-determinant problem. Let $\kappa = \mathbb{F}_q$, $\operatorname{char}\kappa \neq 2, 3$ and $d \in \kappa^*$. Put $E_d : y^2 = x^3 - d$ and $a_q(d) = q + 1 - \#E_d(\kappa)$. For $q \equiv 1 \pmod 3$, fix a nontrivial cubic character ρ and the quadratic character η, both extended by zero at zero. Define

$$J = \sum_{u \in \kappa} \rho(u)\eta(1 - u), \qquad j = -\eta(-1)J, \qquad \lambda_d = \eta(d)\rho(d), \quad \pi_d = \lambda_d j.$$

Theorem 6.2 (Uniform local factor)**.** *For every such prime power and nonzero d,*

$$a_q(d) = \pi_d + \overline{\pi_d}, \quad |\pi_d|^2 = q, \quad P_d(T) := 1 - a_q(d)T + qT^2 = (1 - \pi_d T)(1 - \overline{\pi_d}T).$$

If $q \equiv 2 \pmod 3$, instead $a_q(d) = 0$. The split-case values lie in $\mathbb{Z}[\omega]$, $\omega^2 + \omega + 1 = 0$.

Proof. Cube fibers have size $1 + \rho(t) + \overline{\rho(t)}$, including at zero; the first term is the literal constant one, not the zero-extended trivial character. Counting square fibers gives $a_q(d) = -\sum_x \eta(x^3 - d)$. Grouping by $t = x^3$ removes the constant term since $\sum_t \eta(t - d) = 0$. Substitution $t = du$ gives $\sum_t \rho(t)\eta(t - d) = \rho(d)\eta(-d)J$; the other term is its conjugate. This proves the signed trace formula.

For a nontrivial additive character ψ, write $G(\chi) = \sum_x \chi(x)\psi(x)$. Character orthogonality gives $|G(\chi)|^2 = q$ for nontrivial χ: substitute $x = uy$ in its squared norm; the inner sum over $y \neq 0$ is $q - 1$ at $u = 1$ and -1 otherwise. Grouping $G(\rho)G(\eta)$ by $x + y$ gives $JG(\rho\eta)$, the zero-sum fiber vanishing since $\rho\eta$ is nontrivial. The three Gauss norms yield $|J|^2 = q$. Neither purity nor RH was assumed. The remaining signs and character values have modulus one. For $q \equiv 2 \pmod 3$

cubing is bijective, so the quadratic-character sum is zero. Finally all split-case summands are signs times cube roots of unity. These are the classical Gauss–Jacobi methods [68, pp.499–502], with our normalization proved explicitly. □

At $q = 7$, choose $\rho(3) = \omega$. Then $j = 3 + 2\omega$, $\lambda_2 = \omega^2$, and $\pi_2 = -1 - 3\omega$, giving the two factors already counted. At $q = 13$, $\rho(2) = \omega$ gives $J = 1 + 4\omega$ but $j = -1 - 4\omega$; this detects the essential sign hidden by the first example.

6.2 A genuine cellular carrier, and what still fails for the order

Assume $q \equiv 1 \pmod 3$ in this cellular construction. The elliptic theta dessin has cellular chains $C_2 = \mathbb{Q}$, $C_1 = \mathbb{Q}^3$, $C_0 = \mathbb{Q}^2$. Orient the three edges from v_+ to v_-; then $\partial_1(e_i) = v_- - v_+$ and $\partial_2 = 0$. Thus H_1 is the sum-zero plane, with basis $e_0 - e_1, e_1 - e_2$. Rotation σ acts there by

$$C = \begin{pmatrix} 0 & -1 \\ 1 & -1 \end{pmatrix}, \qquad G = \begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}.$$

G is the actual Euclidean edge Gram matrix in this basis. The deck involution α acts as $-I$. Consequently the six-element deck group $\langle \sigma, \alpha \rangle$ acts on this homology and permutes the original quiver and its entire projective resolution.

Write $\pi_d = A_d + B_d\omega$. Its multiplication action on this *actual cellular* plane is

$$M_d = A_d I + B_d C = \begin{pmatrix} A_d & -B_d \\ B_d & A_d - B_d \end{pmatrix}, \qquad \det(I - TM_d) = P_d(T), \qquad M_d^t G M_d = qG.$$

This constructs the local matrix from finite-field character sums and the dessin's deck action, rather than fitting its trace. Positivity follows from $2x^2 - 2xy + 2y^2 = (x - y)^2 + x^2 + y^2$. The normalized matrix is G-unitary, not asserted self-adjoint. The extended edition supplies a rational cellular chain lift, its exact determinant cancellation, and Kummer descent compatibility on homology. Its odd-over-even determinant ratio is the classical finite-field curve zeta function

$$Z(E_d/\kappa, T) = \frac{P_d(T)}{(1 - T)(1 - qT)}.$$

Identification with the geometric ℓ-adic Frobenius polynomial uses the standard elliptic trace and determinant theorem; a canonical integral comparison or an order-derived Frobenius functor has not been constructed here.

There is also a precise obstruction to extending this same real two-dimensional carrier to $q \equiv 2 \pmod 3$. Frobenius then conjugates the cubic deck action. The relation $MC = C^2 M$ forces $M = \left(\begin{smallmatrix} a & b \\ a+b & -a \end{smallmatrix}\right)$, whose determinant is $-a^2 - ab - b^2 \le 0$, contradicting the required positive determinant q. This excludes that real carrier with its fixed deck action, not complex or enlarged Hilbert–Pólya constructions. The extended edition gives a compatible complex local replacement.

For the following insertion test use the characteristic-zero theta order $\Lambda_{\mathbb{Q}}$ over $R_{\mathbb{Q}} = \mathbb{Q}[[u]]$, with residue simple $S_e = \mathbb{Q}$. The limitation is concrete. The deck action has finite order and cannot alone supply the norm-$\sqrt{q}$ Jacobi coefficient. Inserting M_d into $S_e \otimes \mathbb{Q}^2$ gives an actual residue-module endomorphism with determinant P_d, but that simple is not perfect over the order and choosing it breaks full deck symmetry. Inserting the same matrix $\Lambda_{\mathbb{Q}}$-linearly into the regular projective instead gives the $R_{\mathbb{Q}}$-determinant P_d^{18}, or P_d^6 on $e\Lambda_{\mathbb{Q}} \otimes \mathbb{Q}^2$. Taking a root is not an established determinant-line comparison; a deck-semilinear lift is a different map. These surviving

multiplicity/equivariance obligations prevent promoting the local computation to the required arithmetic order functor.

For the global comparison fix a nonzero rational d and use its good reductions. The finite-field curve zeta has center $1/2$ after $T = q^{-s}$. The *global elliptic* L-function over $\mathbb{Q}$ has unshifted center one, whereas Dirichlet L-functions have center $1/2$. Local weight-one purity does not determine either global zero set. The elementary local bound proves absolute convergence of the elliptic Euler product on $\operatorname{Re} s > 3/2$; the new local identities give no determinant-preserving continuation with the desired global positive spectrum.

6.3 Arbitrary projective actions: an obstruction and a genuine determinant repair

The preceding tensor test has a stronger replacement. Keep the characteristic-zero completed elliptic theta order Λ over $R = \mathbb{Q}[[u]]$, and write $K = \mathbb{Q}((u))$. All determinants in the next theorem are *ordinary underlying-R determinants*; they are not reduced norms.

Theorem 6.3 (Projective determinant multiplicity)**.** *Let P be any finitely generated projective right Λ-module and $f \in \operatorname{End}_\Lambda(P)$. Then $P \simeq \bigoplus_{i=0}^{2} (e_i\Lambda)^{n_i}$. Let $A_i \in M_{n_i}(\mathbb{Q})$ be the induced maps on its three top multiplicity spaces. Then*

$$\det_R(1 - Tf)\Big|_{u=0} = \left(\prod_{i=0}^{2} \det(1 - TA_i)\right)^6.$$

For any bounded complex of these projectives and any Λ-linear chain endomorphism, its alternating determinant ratio specializes to a sixth power in $\mathbb{Q}(T)$. Generically, the ordinary ratio is a cube in $K(T)$.

Proof. The arrow ideal is the Jacobson radical: its elements admit geometric-series inverses after subtracting from one, and the quotient is $\mathbb{Q}^3$. A projective's top lifts to a map from the displayed sum; Nakayama gives surjectivity, projectivity splits it, and another application of Nakayama kills the kernel.

Put $B = \Lambda/u\Lambda$. Each e_iB has path layers of dimensions $1, 2, 2, 1$: the identity, two length-one paths, two length-two paths, and one length-three path. The two full cycles based at i add to ue_i, so become negatives; all length-four paths vanish. A positive-length entry of a projective homomorphism strictly raises path length. Consequently the associated graded map has six copies of $A = \operatorname{diag}(A_0, A_1, A_2)$, giving the formula by triangular block determinants. Base change of determinants justifies specialization. Multiplicativity along filtrations is ordinary linear algebra, consistent with [69, Tags0FJ9,0GSX]; no purity assumption enters.

Each term's determinant has constant term one in T, so every denominator has nonzero specialization. Taking the finite alternating product therefore preserves the sixth power, even after cancellation. Finally $\Lambda_K = M_3(K) \times M_3(K)$; a projective's branch modules are sums of three-dimensional simples. A commuting endomorphism acts on each multiplicity space, repeating every factor three times. This proves the generic assertion. □

Thus no such ordinary perfect-complex ratio can equal the constant-in-u quadratic $P_d(T)$, or the full local curve zeta $P_d(T)/((1 - T)(1 - qT))$. For P_d, the valuation at infinity is -2, not a multiple of six. For the zeta ratio, $P_d(1) = q + 1 - a_q > 0$ by the local bound $|a_q| \le 2\sqrt{q} < q + 1$, so its pole at $T = 1$ is simple. This also covers square prime powers where the numerator may have a double root. It is an obstruction for the specified ordinary determinant and linear category, not all spectral or determinant theories.

There is nevertheless a *genuine integral functor* removing this multiplicity. Choose a permitted branch and its cycle cut. Its projection $\Lambda \to B_+ \subset M_3(R)$ makes the column lattice $W_+ = R^3$ a left Λ-module. Define

$$\Phi_+ : \mathrm{Perf}(\Lambda) \longrightarrow \mathrm{Perf}(R), \qquad P^\bullet \longmapsto P^\bullet \otimes_\Lambda W_+.$$

Bounded projectives compute this derived tensor functor; flatness of W_+ on all modules is unnecessary. Since $e_i W_+ \simeq R$, its determinant on $e_i\Lambda \otimes_{\mathbb{Q}} V$ with external action $1 \otimes M_d$ is exactly P_d, whereas on the regular module it is P_d^3. This is a defined change of determinant functor, not an arbitrary polynomial root. It does not construct the arithmetic action independently: the already computed M_d is still the input.

The price of this repair is precise. Branch swap α exchanges W_+ and W_-; pairing them yields P_d^2 on that basic-projective input. The chosen branch determinant cannot intertwine α on all endomorphisms: the corner unit $e_i + c_i^+$ acting on $e_i\Lambda$, with c_i^+ a plus full cycle, acts on its determinant by $1 + u$, whereas its α-image acts by one. Also no rank-one-per-vertex column lattice supported on one branch admits an R-linear lift of the rotation σ fixing u. Its three nonzero arrow scalars have nonnegative u-valuations summing to one; an invertible cyclic lift would make all three equal. This contradiction concerns that lattice, not every determinant-level rotation comparison. Ramification $u = v^3$ *and a new lattice* with all three arrow scalars v admits such a rotation, changing the stipulated base/lattice data. A descent-compatible arithmetic determinant retaining the required symmetry and a global positive realization remains open.

6.4 Tame descent: a surviving order and a sharper obstruction

We now test the proposed ramified repair rather than assume that its rotation descends. Extend coefficients explicitly to $k = \mathbb{Q}(\omega)$, $\omega^2 + \omega + 1 = 0$. Put $R = k[[u]]$, $S = k[[v]]$, $u = v^3$, and let $g(v) = \omega v$, fixing k. On the balanced plus column $W = S^3$, let $C\epsilon_i = \epsilon_{i+1}$ and

$$\Theta = Dg, \qquad D = \mathrm{diag}(1, \omega^2, \omega).$$

The plus arrows act by $vE_{i,i-1}$, and minus arrows by zero. Direct calculation gives $\Theta^3 = 1$, compatibility of Θ with this Λ_R-action, and

$$\Theta C = \omega^2 C\Theta, \qquad L = W^\Theta = R\epsilon_0 \oplus Rv\epsilon_1 \oplus Rv^2\epsilon_2.$$

Thus invariants restore precisely the original cut lattice. The extension $R \to S$ is finite *free* of rank three and faithfully flat; the group quotient is not a torsor at $v = 0$. The comparison $S \otimes_R L \to W$ has matrix $\mathrm{diag}(1, v, v^2)$, determinant u and cokernel $S/(v) \oplus S/(v^2)$. Hence this semilinear group action alone is not effective descent of the column to R. This is the explicit local root-stack situation, consistent with the trivial-stabilizer criterion for vector-bundle descent [70, Theorem10.3]; no stack theorem is needed for the displayed calculation.

Two positive constructions survive. The determinant line $\bigwedge_S^3 W$ has trivial D and C actions and does descend, but the natural map

$$\det_R(L) \longrightarrow (\det_S W)^\Theta$$

has image uR inside R, not an isomorphism. Determinant and coarse invariants cannot be interchanged while dropping this nonunit comparison. Also the *carrier* endomorphism algebra $\mathrm{End}_S(W)^\Theta$, conjugated to the invariant basis, is exactly the hereditary order with entries R on and below the diagonal and uR above. Its rotation descends as $\mathrm{Ad}(A)$, where

$$A = \begin{pmatrix} 0 & 0 & u \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \qquad A^3 = uI.$$

The algebra automorphism has order three although A is not an integral unit. This is not $\mathrm{End}_\Lambda(W)$ or a descended arithmetic cohomology group.

The natural arithmetic attempt fails: $aI + bC$ does not commute with Θ when $b \neq 0$, as in both prime-seven Jacobi examples. The original cellular sum-zero plane is not Θ-stable either. Replacing C by vC does give the descended operator A on L, but

$$\det(1 - T(aI + bA)) = (1 - aT)^3 - ub^3T^3,$$

whose special fiber is $(1 - aT)^3$, not the required local quadratic. These carrier maps are not automatically Λ-linear arithmetic endomorphisms.

There is a stronger category-level conclusion.

Theorem 6.4 (Full-deck coefficient kernels)**.** *Let W_0 be a finite R-free left Λ_R-module on which the central u acts by the given scalar. Suppose it has invertible R-linear intertwiners lifting the order automorphisms σ and α. Then, for some $m \geq 0$, every e_iW_0 has rank $6m$. For any finite projective P and Λ_R-linear $f : P \to P$, let A_i be its top blocks. The genuine coefficient functor $\Phi_{W_0} = - \otimes^{\mathbf{L}}_{\Lambda} W_0$ satisfies*

$$\det_R(1 - T\Phi_{W_0}(f))\Big|_{u=0} = \left(\prod_i \det_k(1 - TA_i)\right)^{6m}.$$

Bounded perfect ratios retain this exponent. Thus this whole class of full-deck, fixed-center finite free coefficient kernels cannot supply either the single local quadratic or the full local curve zeta.

Proof. In the generic fiber intersect W_0 with its two central branch summands to obtain finite free integral lattices $W_\pm$; an integral direct-sum decomposition is not assumed. Rotation preserves them and branch swap exchanges them. On a branch let the common vertex rank be r. The three arrow matrices have cyclic product uI_r. Their determinant valuations are nonnegative, sum to r, and rotation intertwiners make them equal. Therefore $3 \mid r$. Branch swap makes the two ranks equal, so the full vertex rank is $6m$.

For the determinant formula, filter W_0/uW_0 by the arrow radical, whose fourth power vanishes. A positive-length entry of f raises the filtration; a top block A_i repeats across its i-th graded pieces a total of $\dim_k e_i(W_0/uW_0) = 6m$ times. Triangular determinants give the assertion. Finite alternating products preserve it. Quadratic degree and the uncancelled simple pole of the full local zeta exclude the targets as before. □

This theorem is sharp: $W_0 = \Lambda_R$ has vertex rank six. It requires neither left projectivity nor left flatness of a general kernel. It excludes this coefficient category, not arbitrary semilinear correspondences, twisted determinant lines or infinite spectral constructions. The extended edition identifies the full character-orbit repair with the hereditary normalization and explains why Schur commutator cancellation is insufficient. All determinant multiplicities here are relative to the stated k-valued base; restricting a rational quadratic factor to the former $\mathbb{Q}$-base squares it. No arithmetic descent or global positive determinant has been recovered by this ramification.

The recent NCG literature supplies concrete comparison targets, with strong hypotheses. Connes–Consani construct finite abelian covers of the adelic Riemann sector with prime-orbit length $\log p$ and arithmetic Frobenius monodromy [53, Theorem3.12]. The prolate/local-factor and moment constructions [54, 55] give particular finite-prime analytic models, not a uniform positive comparison for all completed L-functions. The zeta-spectral-triple determinant theorem [56, Theorem5.10] assumes a specified restricted Weil form, simple even bottom eigenfunction and boundary normalization. Its infinite comparison is still an additional step. The distributional real-zero theorem [57, Theorem6.1]

concerns a particular interval quadratic form, essential self-adjointness on its trigonometric core and an even simple ground state; a Fock vacuum alone verifies none of those arithmetic identifications. Deninger's periodic-packet construction and real-cohomology obstruction [60, 62] further specify, rather than remove, these issues. Full theorem-level reading coverage and exact versions are recorded alongside this draft.

A complete approach must construct arithmetic descent and prime actions on actual order/derived or analytic objects, prove determinant-line and local-factor compatibility, control regularization and conductor/gamma normalization, and establish a positive realization for that *same* function. Finite-field cohomological trace formulas and Deligne purity supply the classical model [20, 21]; alternating duality alone is not positivity or purity. In a number-field construction these are obligations, not imported conclusions.

For a sharply testable final analytic target, put $F_\chi(z) = \Lambda(1/2 + iz, \chi)$. Suppose entire functions $h_m(z)\det(zI - H_m)$, with H_m finite self-adjoint and h_m entire nowhere zero, converge locally uniformly on $\mathbb{C}$ to this nonzero completed function. Hurwitz on every disk disjoint from $\mathbb{R}$ then proves all its zeros real, hence the Dirichlet critical-line assertion. Establishing this convergence with independently constructed H_m is exactly the missing load-bearing claim; a rank-one encoding or a reformulation does not establish it. Hecke, Dedekind and Artin extensions must state their analytic scope anew.

6.5 Whole components, derived endpoints and determinant relations

The full node lattice category has precisely the three indecomposables X, Λ, Y and arrows $X \to \Lambda \to Y$, $Y \to \Lambda \to X$. Its stable category has only X, Y, no irreducible arrows, and suspension exchanging them; its AR triangles have zero middle term. For the polynomial node at a smooth support $x = \lambda + t$, $y = 0$, $\lambda \neq 0$, the modules $T_r = k[t]/t^r$ form an ambient homogeneous tube:

$$0 \longrightarrow T_r \xrightarrow{(\pi,t)} T_{r-1} \oplus T_{r+1} \xrightarrow{(t,-\pi)} T_r \longrightarrow 0, \quad T_0 = 0, \quad \tau T_r = T_r.$$

The shift changes the derived placement, not the quasi-length. These nonzero-central-support objects do not extend to continuous finite-length modules of the arrow-adic completion. At its singular point even the candidate pure-branch tube fails ambient almost-splitness: $T_2^y \to k$ cannot factor through $T_2^x \to k$, since every intervening map lands in the x-socle, killed by the last map.

The nine HLS strings above sit in an entire intrinsic quotient component $T_{p,q}$, $(p, q) \in \mathbb{Z}^2$, with arrows increasing one coordinate and $\tau T_{p,q} = T_{p-1,q-1}$ [46]. Its central object $E = M(ab)$ has infinite projective dimension: restriction to the first corner gives the dual-number simple with its infinite loop resolution, while restricted projectives are free or zero. Its module AR sequence therefore yields a distinguished derived triangle but not a derived AR triangle ending at E. Ringel's four-quarter wind-wheel tile [43], with infinite-word diagonals and mixed center, is a different category's quilt; it is not silently identified with that HLS component or the Belyi surface.

There is no general equation "AR mesh equals polynomial semi-invariant syzygy." The intrinsic $k(1 \to 2)$ quotient of the infinite Chebyshev algebra has AR row $0 \to S_2 \to E \to S_1 \to 0$ and a free presentation determinant z on dimension $(1, 1)$, so its semi-invariant ring $k[z]$ has no generator relation. A genuine framed elliptic-square stratum has $Z = [x_1, x_2, y_1, y_2] \in \mathrm{Mat}_{2\times 4}$ with six minors and relation

$$p_{12}p_{34} - p_{13}p_{24} + p_{14}p_{23} = 0.$$

These minors have the same determinant character for the target GL_2 with framed source; they are not all semi-invariants for the full unframed vertex group. In contrast, the nodal lattice AR row gives a genuine relative determinant identity over R: multiplication by $a + bx$ on the basis $(y, 1)$

has matrix $\left(\begin{smallmatrix} a & -b \\ 0 & a+ub \end{smallmatrix}\right)$ and characteristic factor $(1-Ta)(1-T(a+ub))$. The endpoints have infinite order-projective dimension but are R-perfect.

Finally, the three-vertex theta algebra has no unital surjection onto a six-vertex radical-square-zero circular algebra: its semisimple image has at most three primitive idempotent images. Duplicating face vertex spaces instead gives an exact faithful unfolding which is not full; it does not inherit quotient or AR-transfer claims. Zwara Hom-control means that the Hom-dimension defect is bilinear on K_0, not merely a covering direct-sum formula [72, 73]. The book supplies explicit fixed-dimension counterexamples, corrected circular-complex incidence resolutions, complete drawn component windows and all category hypotheses. These additions do not supply the arithmetic determinant comparison.

7 A genuine theta covering and its categorical limits

Use column matrices, hence covariant quiver representations (left modules); opposite categories translate the right-module convention in [74]. Let A be the infinite elliptic theta dessin algebra: vertices $i \in \mathbb{Z}/3$, arrows $a_i, b_i : i \to i+1$, and all mixed length-two paths zero. For the arrow ideal J and $L \geq 2$ put $B_L = A/J^L$. This is a finite string quotient of our actual infinite gentle algebra; for $L > 2$ its length-L relations prevent treating it as quadratic gentle.

Proposition 7.1 (A locally bounded integer cover)**.** *The category $\widetilde{B}_L$ with vertices $r \in \mathbb{Z}$, two arrows $a_r, b_r : r \to r+1$, and the same mixed and length-L zero relations, has orbit B_L under $r \mapsto r+3$. For finite-support representations, push-down F is exact and*

$$\operatorname{Hom}_{B_L}(FM, FN) \simeq \bigoplus_{g \in \mathbb{Z}} \operatorname{Hom}_{\widetilde{B}_L}(M, T_g N), \qquad (T_g N)_r = N_{r-3g}.$$

Only finitely many summands are nonzero. In general F is neither full, dense, nor Hom-controlled in Zwara's sense.

Proof. A nonzero path is constant-branch with length less than L. Thus the incoming and outgoing total Hom dimensions at a cover vertex are $2L-1$, and $\dim B_L = 3(2L-1)$. Paths and relations lift uniquely; the deck action is free and has the stated orbit. Push-down sums spaces of equal residue. A morphism decomposes uniquely into blocks with target lift index minus source index equal to $-3g$. The arrow commutation equations preserve that difference, proving the formula; finite supports bound the possible differences. Direct sums of vector spaces preserve exact sequences. The following explicit witnesses prove the three failures. □

For $L = 5$, let P be the cover interval on $0, 1, 2, 3, 4$ with all its a-maps one and b-maps zero, and let S be the sum of the five simples at the same vertices. They have the same K_0 class, but

	$\dim \operatorname{End}_{\widetilde{B}_5}$	$\dim \operatorname{End}_{B_5}(F(-))$	defect
S	5	9	4
P	1	2	1

Indeed the pushed-down simple multiplicities are $(2, 2, 1)$, whereas $\operatorname{End}_{B_5}(FP) = kI \oplus kX^3$ for its single nilpotent five-step chain X. The extra endomorphism X^3 proves non-fullness; the unequal defects at the same class exclude a bilinear K_0 defect. To use Zwara's finite-algebra definition literally, restrict the source to the finite convex interval $[0, 4]$, then extend by zero before push-down. This exact functor has the same endomorphism spaces and K_0 comparison.

Proposition 7.2 (An indecomposable module outside this cover image)**.** *Over a characteristic-zero field, B_3 has a nine-dimensional indecomposable representation which is not the push-down of any finite-support $\widetilde{B}_3$ representation.*

Proof. Label its basis by $0, \dots, 8$ with vertex residues $(0, 1, 0, 1, 2, 1, 2, 0, 2)$, and use matrix units E_{ij} acting on columns. Set

$$X = E_{10} + E_{32} + E_{43} + E_{65} + E_{76} + E_{08},$$
$$Y = E_{12} + E_{45} + E_{78}.$$

Their restrictions between adjacent vertex spaces give the a_i, b_i maps. They satisfy $XY = YX = X^3 = Y^3 = 0$, hence every length-three path is zero. The endomorphism ring preserving vertex spaces is $kI \oplus \mathfrak{r}$, where

$$\mathfrak{r} = \langle E_{02} + E_{13}, E_{15}, E_{35} + E_{46}, E_{48}, E_{68} + E_{70}, E_{72} \rangle, \qquad \mathfrak{r}^3 = 0.$$

Solving the arrow commutation equations gives exactly these seven basis elements. The ring is local: an element $cI + n$ is invertible exactly when $c \neq 0$, by the finite geometric series. It has no nontrivial idempotents, so the representation is indecomposable.

A finite-support lift would define an Euler operator D acting by r on the summand at lift vertex r. Then $[D, X] = X$ and $[D, Y] = Y$. Every displayed nonzero arrow is the unique nonzero entry in its row and column in its own matrix. Its commutator entry is therefore $D_{ii} - D_{jj}$. Add the six X equations and subtract the three Y equations. Each diagonal term cancels, leaving $0 = 6 - 3 = 3$, a contradiction. This argument allows arbitrary D, not just a diagonal matrix in the displayed basis. □

Corollary 7.3 (Higher-genus and derived non-density)*.* For the clean dessin with rotations $\sigma = (0 \cdots m-1)(m \cdots 2m-1)$ and $\alpha = \prod_{i=0}^{m-1}(i\, i+m)$, odd $m \geq 3$, and genus $(m-1)/2$, the analogous A_m/J^3 has a $3m$-dimensional representation outside its integer-cover image in characteristic zero. Its bounded-derived push-down is not dense either.

Proof. Use basis $v_0, \dots, v_{3m-1}$, assign vertices $i, i+1, i$ to $v_{3i}, v_{3i+1}, v_{3i+2}$ modulo m, and set

$$Xv_{3i} = v_{3i+1}, \quad Xv_{3i+2} = v_{3i+3}, \quad Yv_{3i+2} = v_{3i+1},$$

with indices modulo $3m$ and all other values zero. Again mixed products and cubes vanish. An Euler operator would yield $2m - m = m = 0$ by the same telescoping calculation, impossible in characteristic zero. If a bounded finite-support complex pushed to this module in degree zero, exactness would make its zeroth cohomology a module lift, a contradiction. □

The source's derived result is a *precovering* in the stated bounded categories [74, Theorem 4.28]; density remains an additional condition. Neither this result nor the GLS equivariant classification [75] identifies our completed order with a GLS algebra. Graded surface/Fukaya models [76] likewise require their own dg category, stops and grading. These categorical tools do not construct the independent positive arithmetic determinant.

8 Verification and research frontier

The two nodal/invariant Lean files construct the node as an actual subring of $k[[X]] \times k[[X]]$ with equal constants, prove its annihilators, regular common parameter, augmentation, alternating

projective resolution and a bijective pair of power-series coordinates. A second file proves CK1 collapse under the stated permutation/coverage hypotheses, determinant base change, actual 2×2 symmetric-square functoriality and relation preservation, a Schur character identity and the fixed-parameter lattice failure. These thirty printed theorem types have only standard Lean foundations. They do not formalize general dessins, abstract Ext/AR/Schur–Weyl, C* exactness, Fredholm determinants, arithmetic descent or GRH. Earlier formal layers remain preserved with their literal scopes.

Exact-rational diagnostics independently check the displayed dessins, path identities, Schur and Young examples and the nine-object grid. These finite checks supplement the proofs, rather than prove infinite spectral or arithmetic assertions. The next arithmetic target is an exact descent/Frobenius-compatible construction on the labelled order/Fock data, with a determinant-preserving positive comparison. The local Schur kernel and the simple degree spectrum alone do not supply it. All source clones and earlier research are retained; the earlier general focused PDFs are archived at the preceding commit.

Ten further declarations in `DessinPullbackComponents.lean` check actual matrix-unit identities, nonmultiplicativity of raw diagonal extraction, impossibility of a unital two-state matrix residue into a nonzero commutative ring, multiplicativity on diagonal fibers, and genuine equal-image pullback subring/projection components. The analytic suite then contained187printed declarations at unchanged pins. The circle-function C* model, functional calculus, generation, graph uniqueness and K-theory remain prose/external-theorem arguments, not hidden assumptions of these component checks.

Sixteen additional declarations in `ThetaJacobi.lean` check genuine finite-field MulChar Jacobi conjugation, norm and rescaling with explicit nontriviality hypotheses; cubic–quadratic noncancellation; the cyclotomic multiplication determinant and positive Gram identity; and actual theta cellular boundary, homology-inclusion, restriction and rational chain-map matrices. The full analytic suite now contains203declarations. The point-count proof, deck/Kummer comparison, cohomological trace interpretation and global arithmetic comparison are not thereby fully formalized.

The final three `ThetaJacobi.lean` declarations verify the actual conjugation matrix identities and the real two-dimensional inert-prime determinant obstruction. They do not assume any unknown zero location or exclude complex/enlarged spectral realizations.

`ThetaProjectiveMultiplicity.lean` adds five declarations. Arbitrary upper-block matrices with six identical diagonal blocks have the sixth-power determinant and characteristic-factor formulas. A literal power-series constant-coefficient map specializes matrix determinants, for each scalar parameter; the coefficientwise polynomial/order interface remains prose. A nonzero quadratic cannot be a rational-function sixth power, proved by polynomial degrees. Three genuine power series with product X cannot have the cyclically equivalent residue-vanishing conditions forced by invertible rank-one arrow intertwiners. The derivation of the path blocks from the full theta order, projective decomposition, derived branch functor and deck intertwiners is not fully formalized. The complete analytic suite now has208printed declarations; no hidden arithmetic positivity assumption occurs.

`ThetaTameDescent.lean` adds twelve printed component declarations (and one auxiliary polynomial lemma). They use genuine formal-power-series rescaling and its coefficient operator, proving the coefficient action, invariant-coefficient vanishing and the tame operator's cube. Actual three-state matrices satisfy the clock/rotation commutator, comparison determinant and closed-fiber equations, weighted-cycle cube and full characteristic polynomial, noncommutation of a nonconstant rotation polynomial, trivial whole-column determinant actions, and the same-operator comparison for vC. The classification of all fixed series by residues, full order action and hereditary invariant algebra, stack interpretation, valuation/rank theorem, general coefficient-functor filtration and GRH are not

thereby fully formalized. No unknown positivity premise is assumed.

9 Why an absolute coprime pair-square estimate cannot close the short rows

Keep the original sextic family

$$M_u(D) = D^{-1/2} \sum_n \mu(n)\nu(n)W(Nn/D)\chi_n(u), \qquad \chi_n(u) = (u/n)_6,$$

with squarefree primary column ideals outside the fixed excluded set, their full annular coefficients (fix W supported in $[1,2]$), and all nonzero Eisenstein element rows $Nu \le H$. The cubic symbol is χ_n^2, not χ_n. The following classical frame inequality [80] gives a new obstruction to one proposed estimate in this arithmetic program.

Proposition 9.1 (Coprime pair-square obstruction)**.** *Let A be an R-by-N complex matrix with $|A_{ui}| = 1$, and put $K_{ij} = \sum_u A_{ui}\overline{A_{uj}}$. Then*

$$\sum_{i \ne j} |K_{ij}|^2 \ge NR(N-R).$$

For the original character kernels, let $B_1(H,D)$ be the sum of $|K_H(n,m)|^2$ over all distinct coprime squarefree column ideals in $[D,2D]$. At $H = D^{3/4}$,

$$B_1(H,D) \gg \frac{D^{11/4}}{(\log D)^2}.$$

In particular the proposed sufficient input $B_1(H,D) \ll H^2 D^{\varepsilon}$ fails for each fixed $\varepsilon < 5/4$. This does not refute the signed Möbius mean square.

Proof. Set $F_{uv} = \sum_i A_{ui}\overline{A_{vi}}$. Reordering four finite sums proves $\sum_{i,j} |K_{ij}|^2 = \sum_{u,v} |F_{uv}|^2$. The row diagonal contributes RN^2; the column diagonal contributes NR^2. Removing the latter and retaining the former gives the inequality.

Restrict the column pairs in the nonnegative sum B_1 to distinct prime ideals p, q of norm in $[D,2D]$, outside the same fixed set. If $H < D$, no such prime divides a nonzero row element: divisibility would imply $Nu \ge Np$. Thus all original zero masks are exactly one on this ball, $|\chi_p(u)| = 1$, and the residual primitive finite character conductor is pq, of norm between D^2 and $4D^2$. All these pairs have common divisor1. The lattice count gives $R = (2\pi/\sqrt{3})H + O(\sqrt{H}+1)$. The classical unconditional fixed-field prime-ideal theorem gives $N \sim D/\log D$; this is an external analytic input, not a GRH hypothesis. At the stated endpoint $R = o(N)$, and the bound is therefore $(1-o(1))N^2R \gg D^{11/4}/(\log D)^2$. □

To see the exact failed step, write $c_n = \mu(n)\nu(n)W(Nn/D)$ and $C_D = \sum_n |c_n|^2 = O(D)$. Cauchy over coprime pairs bounds the absolute signed contribution by $D^{-1}C_D B_1^{1/2}$. The hoped-for pair-square upper bound would have closed this particular route, but is false. A large absolute pair-square mass is not a lower bound on the quadratic form with the fixed c_n. Nor is the prime part a positive summand of the full Möbius energy: prime–composite cross terms can cancel. The original low-gcd estimate and the positive arithmetic determinant bridge remain open.

10 A signed Möbius decomposition and its termwise obstruction

Keep the original squarefree good-ideal family, its fixed ν, its full $W(Nn/D)$ and every zero-extended sextic symbol. In ideal Dirichlet convolution, write $\zeta_0(n) = 1$ for every good ideal and let δ be the convolution unit. For $1 \le U < D$ put $a_U = \mu \mathbf{1}_{Nn \le U}$ and $b_U = \mu - a_U$. The classical Vaughan algebra gives

$$\mu = 2a_U - a_U * a_U * \zeta_0 + b_U * b_U * \zeta_0. \tag{1}$$

Indeed $\mu * \zeta_0 = \delta$, and expansion of the last product proves the identity without an analytic assumption. This is a classical decomposition, not a new cancellation theorem; compare [87, Lemma4.1].

On squarefree n all factors in $n = dc = abc$ are pairwise coprime. Define the actual, nonarbitrary coefficient functions

$$\beta_U(d) = \sum_{\substack{ab=d \\ Na, Nb \le U}} \mu(a)\mu(b), \qquad g_U(s) = \sum_{\substack{b \mid s \\ Nb > U}} \mu(b).$$

For squarefree d, $\beta_U(d)$ is $\mu(d)$ times the number of the displayed ordered factorizations. For $Ns > U$, $g_U(s) = -\sum_{b \mid s,\, Nb \le U} \mu(b)$. The two regrouped pieces are

$$I_u = D^{-1/2} \sum_{\substack{d,c \text{ squarefree} \\ (d,c)=1}} \beta_U(d)\nu(dc)\chi_d(u)\chi_c(u)W(NdNc/D),$$

$$II_u = D^{-1/2} \sum_{\substack{a,s \text{ squarefree} \\ Na, Ns > U,\, (a,s)=1}} \mu(a)g_U(s)\nu(as)\chi_a(u)\chi_s(u)W(NaNs/D).$$

Then $M_u = -I_u + II_u$. All ideals here remain outside the same fixed excluded set. The short $2a_U$ term vanishes because $U < D$ and W is supported in $[1, 2]$. No phase or coprimality mask was removed.

Proposition 10.1 (An all-cutoff separate-piece bound is false)**.** *Take the allowed principal ν, $U = 1$, and a fixed nonnegative smooth W supported in $[1, 2]$, bounded below by a positive constant on a closed subinterval of $(1, 2)$ of positive length. For each of the two pieces,*

$$\sum_{0 < Nu \le H} |I_u|^2 \gg_W \frac{D}{\log^2 D}, \qquad \sum_{0 < Nu \le H} |II_u|^2 \gg_W \frac{D}{\log^2 D} \quad (H \ge 1).$$

Consequently a proposed bound HD^{ε} for each piece at $H = D^{3/4}$ fails for $\varepsilon < 1/4$. In the exact original notation $H(HD)^{\varepsilon}$, it fails for $\varepsilon < 1/7$. This refutes an all-cutoff termwise strengthening, not the original signed mean square or a specified moving-cutoff argument.

Proof. At $U = 1$, $a_U = \delta$, so $a_U * a_U * \zeta_0 = \zeta_0$ and $b_U * b_U * \zeta_0 = \mu + \zeta_0 - 2\delta$. On the annulus the two coefficients are 1 and $1 + \mu(n)$, respectively. On squarefree ideals the latter is zero or two. At the actual unit row $u = 1$, every original character equals one. A fixed-field prime-ideal theorem gives $\gg D/\log D$ good prime columns in the subinterval, so $I_1 \gg_W \sqrt{D}/\log D$. Fix a good prime ideal q_0. Products $q_0 p$ with varying distinct good primes in the rescaled subinterval have $\mu(q_0 p) = 1$. Their nonnegative contributions give $II_1 \gg_W \sqrt{D}/\log D$. Keeping that one row in each nonnegative energy proves the result. The prime-ideal theorem is a classical unconditional external input, with the field and q_0 fixed. □

The factorization does not shorten the primitive row conductor: for squarefree coprime a, s it is as, not a or s separately. For two resulting columns n, n' with common factor h, the oscillatory conductor is nn'/h^2 and a zero-preserving period is nn'/h. The signed low-gcd estimate and the independent positive arithmetic determinant comparison therefore remain open.

11 A full-mask Hecke expansion for the type-I piece

We now continue the signed decomposition through its actual squarefree inner sum. Fix $K = \mathbb{Q}(\sqrt{-3})$, the finite-order Hecke character ν, and the same finite excluded prime set S, containing primes above 6 and the conductor of ν. For $u \in \mathcal{O}_K \setminus \{0\}$, let ψ_u be the primitive Hecke character induced by $\nu(n)(u/n)_6$ away from $S \cup \operatorname{supp}(u)$. It is defined before, and independently of, the type-I divisor d. The abelian Kummer extension $K(u^{1/6})/K$ constructs the sextic factor; multiplying by ν gives the stated character. This uses classical abelian reciprocity, not general Artin holomorphy. Write $\psi_{u,2}$ for the primitive character inducing its square and $\mathfrak{f}_u, \mathfrak{f}_{u,2}$ for the two primitive conductors. For a squarefree good ideal d put

$$T = T(u,d) = S \cup \operatorname{supp}(u) \cup \operatorname{supp}(d), \qquad R_T = \prod_{p \in T} Np, \quad Q_u = |D_K| N\mathfrak{f}_u.$$

All primes in T are deleted even if absent from either primitive conductor.

Theorem 11.1 (Restoration and a uniform scalar contour bound)**.** *For $\Re s > 1$ the original type-I inner Dirichlet series is*

$$F_{u,d}(s) = \sum_{\substack{c \text{ squarefree} \\ (c,T)=1}} \frac{\psi_u(c)}{(Nc)^s} = \frac{L(s,\psi_u)}{L(2s,\psi_{u,2})}\, C_T(s), \tag{2}$$

$$C_T(s) = \frac{\displaystyle\prod_{p \in T,\ p \nmid \mathfrak{f}_u} (1 - \psi_u(p)(Np)^{-s})}{\displaystyle\prod_{p \in T,\ p \nmid \mathfrak{f}_{u,2}} (1 - \psi_{u,2}(p)(Np)^{-2s})}. \tag{3}$$

It continues meromorphically to $\Re s > 1/2$, with only a possible simple pole at $s = 1$ when $\psi_u = 1$. In that case its residue is

$$\rho_T = \frac{\operatorname{res}_{s=1} \zeta_K(s)}{\zeta_K(2)} \prod_{p \in T} (1 + (Np)^{-1})^{-1}.$$

Fix $0 < \delta < 1/2$, $\varepsilon > 0$ and $W \in C_c^\infty((0,\infty))$ supported in $[1,2]$. Set

$$\mathcal{M}_4(W) = \sum_{j=0}^{4} \sup_x |(x\partial_x)^j W(x)|.$$

Uniformly for $Y \geq 1$, u and d,

$$\sum_{\substack{c \text{ squarefree} \\ (c,T)=1}} \psi_u(c) W(Nc/Y) = \mathbf{1}_{\psi_u = 1} \rho_T Y \widehat{W}(1) + O_{\nu,S,\delta,\varepsilon}\Big(\mathcal{M}_4(W) Y^{1/2+\delta} Q_u^{1/4-\delta/2+\varepsilon} R_T^{\varepsilon}\Big). \tag{4}$$

Here $\widehat{W}(s) = \int_0^\infty W(x) x^{s-1}\, dx$. The continuation and functional equation of degree-one Hecke functions are classical external inputs; the contour argument is given below.

At a good prime with $v_p(u) = 3$, ψ_u is ramified while its square is unramified; the second product in (3) must restore a factor invisible to squaring the zero-extended primitive value. At $v_p(u) = 6$

both characters can forget the prime, but the original row still vanishes on columns divisible by it. In particular, for principal ν, $u=\pi^3$ and $u=\pi^6$ at $N\pi=7$, the deleted local factors are

	primitive ratio at $s=1$	restoration	full masked factor
$v_p(u)=3$	48/49	49/48	1
$v_p(u)=6$	8/7	7/8	1

These are exact local calculations, not numerical evidence about zeros.

For $1\le U$, $U^2\le D$, insert $Y=D/Nd$ into (4) and retain the actual $\beta_U(d)$ and outer $\nu(d)\chi_d(u)$. Classical tame Kummer conductors give $Q_u\ll_\nu Nu$. Thus the normalized I_u equals its displayed principal-residue sum plus an error E_u with

$$|E_u|\ll_{\nu,S,\delta,\varepsilon}\mathcal{M}_4(W)D^{\delta}U^{1-2\delta+\varepsilon}(Nu)^{1/4-\delta/2+\varepsilon}. \tag{5}$$

The resulting elementary energy envelope is only

$$\sum_{0<Nu\le H}|E_u|^2\ll\mathcal{M}_4(W)^2D^{2\delta}U^{2-4\delta+\varepsilon}H^{3/2-\delta+\varepsilon}. \tag{6}$$

It does not prove the short-row target of order H up to small losses. This is an insufficient upper bound, not a lower bound on the actual error or a counterexample to coefficient-sensitive averaging. The original signed mean square and positive determinant bridge stay open.

12 Actual renormalization retains the effective action

A compression score does not specify an RG transformation. Following the real-space formulation of Wilson and Kogut [81], we specify the eliminated variables, effective action, normalization and scale map. Our exact finite decimation is distinct from an approximate tensor-network truncation such as Levin–Nave TRG [82].

Definition 12.1 (Finite classical RG step)**.** Let the fine configuration be $(s,t)\in S\times E$, with finite nonempty sets S,E, and dimensionless log Boltzmann weight $V(s,t)=-\beta H(s,t)$. Define

$$(\mathcal{R}V)(s)=\log\sum_{t\in E}e^{V(s,t)},\qquad Z(V)=\sum_{s,t}e^{V(s,t)}.$$

The full action includes its additive constant. A truncation or a learned approximation is a further operation with its own error, not this identity.

Proposition 12.2 (Partition, observables and composition)**.** *One has $Z(\mathcal{R}V)=Z(V)$, and the normalized expectation of every observable depending only on s agrees in the two descriptions. Eliminating E_1 and then E_2 equals eliminating $E_1\times E_2$ at once, with the same tensor labels and inverse temperature.*

Proof. Exponentiating the definition gives $e^{\mathcal{R}V(s)}=\sum_t e^{V(s,t)}$. Sum over s, optionally multiplying by an observable $O(s)$, and divide both sides by the same strictly positive Z. Reordering finite sums proves the composition statement. No projection onto pair interactions is used. □

For an even homogeneous periodic Ising chain, write the transfer matrix as $T_{ab}=\exp(C+Kab+h(a+b)/2)$, $a,b\in\{-1,1\}$. Its partition function is $\operatorname{tr}T^N$. Summing alternate spins replaces T by T^2 and N by $N/2$; the physical spacing doubles. Writing $P=T^2$, its positive entries determine

$$K'=\tfrac14\log\frac{P_{++}P_{--}}{P_{+-}^2},\quad h'=\tfrac12\log\frac{P_{++}}{P_{--}},\quad C'=\tfrac14\log(P_{++}P_{--}P_{+-}^2).$$

These three parameters reproduce every entry of P, so no normalization is lost. At $h = 0$, $\tanh K' = \tanh^2 K$ and the eigenvalue-ratio correlation length of the homogeneous extension in blocked units halves. Its product with the spacing stays fixed. The fixed points in $t = \tanh K$ are $t = 0, 1$; the latter is zero temperature, not a positive-temperature transition. Odd dessin examples below use the full-action definition, not this even-chain shortcut.

12.1 Genuine dessin incidence models

For odd m, the clean dessin $y^2 = x^m - 1$, $\beta = x^m$ has two black vertices, m white vertices and one face; its genus is $(m-1)/2$. Its incidence graph is $K_{2,m}$ with $2m$ clean edges. Its infinite locally gentle order quiver instead has m vertices and completed DVR-rank $2m^2$. We place physical spins $B_+, B_-, s_0, \ldots, s_{m-1}$ on the incidence vertices; this is a specified finite spin model, not a faithful representation of that infinite algebra or order. The abstract incidence graph alone does not recover the ribbon embedding or genus.

With $\eta_j = (-1)^j$ and fixed nonuniform fields h_j, choose

$$V = J_+ B_+ \sum_j s_j + J_- B_- \sum_j \eta_j s_j + bB_+ - \tfrac{3}{5} bB_- + \sum_j h_j s_j.$$

Summing the two black spins gives the *entire* effective action

$$V'(s) = \sum_j h_j s_j + \log\left[2\cosh\left(b + J_+ \sum_j s_j\right)\right] + \log\left[2\cosh\left(-\tfrac{3}{5} b + J_- \sum_j \eta_j s_j\right)\right].$$

Every one of the $2m$ physical edges is present, although its couplings belong to a tied three-parameter family. The Walsh expansion $V'(s) = \sum_{A \subseteq \{0,\ldots,m-1\}} c_A \prod_{j \in A} s_j$, with $c_A = 2^{-m} \sum_s V'(s) \prod_{j \in A} s_j$, is unique by Boolean orthogonality. It retains the constant $c_\varnothing$ and genuinely generated higher-body interactions. Our genus-two and genus-four examples use $m = 5, 9$ and continue by exact nested marginalization. Dropping all terms of degree greater than two is displayed as a measured approximation.

12.2 Quantum thermal effective Hamiltonians

For a finite Hermitian Hamiltonian H on nonzero finite-dimensional $\mathcal{H}_S \otimes \mathcal{H}_E$ and $\beta > 0$, define

$$W' = \operatorname{Tr}_E e^{-\beta H}, \qquad H_{\text{eff}}(\beta) = -\beta^{-1} \log W'.$$

The matrix W' is strictly positive: for $v \neq 0$, sum the strictly positive quadratic forms of $v \otimes e_j$ over an orthonormal basis of $\mathcal{H}_E$. Its Hermitian spectral logarithm is therefore defined. The identity $e^{-\beta H_{\text{eff}}} = W'$ proves partition and retained *equal-time* observable preservation by the partial-trace property. Repeated traces compose. The full Pauli expansion, including its identity coefficient, is retained. It is generally nonlocal and depends on β; this statement does not preserve all fine real-time dynamics or supply a local zero-temperature/thermodynamic RG fixed point.

`FiniteRG.lean` checks nine actual finite scalar marginal, exponential-action, partition, observable, composition, error-sandwich and transfer-trace statements. The quantum spectral-log and complete thermodynamic/field-theory interfaces are not formalized by these lemmas. These physics constructions give no arithmetic determinant comparison or new GRH conclusion.

13 Linear relations and functorial branch filtrations

The original linear-relations chapter mixes graphs, partial relations and stable kernels. We keep these different objects separate. For a field k, a linear relation is a subspace $C \subseteq V \oplus V$; its inverse swaps coordinates and $DC = \{(x,z) : \exists y, (x,y) \in C, (y,z) \in D\}$. The full graph $\Gamma_T = \{(x, Tx) : x \in V\}$ always has dimension $\dim V$, including pairs with zero output. In particular $\Gamma_T^n(0) = 0$, whereas $(\Gamma_T^{-1})^n(0) = \ker T^n$. Thus the source's $\kappa(\Gamma_T) = \bigcup_n \Gamma_T^n(0)$ is zero, not the ordinary stable kernel.

For odd $m \geq 3$, use the actual clean dessin of $y^2 = x^m - 1$, $\beta = x^m$, on the compact smooth curve, in characteristic zero. Its genus is $(m-1)/2$; its infinite locally gentle algebra A_m has vertices $i \in \mathbb{Z}/m$, arrows $a_i, b_i : i \to i-1$, and mixed consecutive branch products zero. Below paths act in their direction of travel on left representations over any coefficient field k. Let $\mathcal{E}_m$ be the full category of finite-dimensional representations with every $b_i = 0$ and a_i invertible for $i \neq 0$.

Theorem 13.1 (A concrete relation category on an infinite dessin algebra)**.** *The category $\mathcal{E}_m$ is equivalent to finite-dimensional pairs (V,T) with intertwining maps. A normal form $N_m(T)$ puts V at each vertex, $a_0 = T$, and all other $a_i = 1$. For $n = \dim V$,*

$$V = K(T) \oplus I(T), \qquad K(T) = \ker T^n, \quad I(T) = \operatorname{im} T^n.$$

These summands and their kernel/image filtrations are functorial. The first restriction is nilpotent and the second invertible. The central full-turn parameter u acts as T. A finite $N_m(T)$ extends continuously to the u-adically completed order precisely when T is nilpotent. Such modules are torsion, not nonzero order lattices.

Proof. Transport all vertex spaces to vertex zero along the invertible arrows; the remaining map is the full-turn monodromy T. Arrow-compatible maps become exactly $hT = T'h$. Finite kernel/image chains stabilize by step n. If $v = T^n w$ and $T^n v = 0$, then $w \in \ker T^{2n} = \ker T^n$, hence $v = 0$; rank-nullity proves the direct sum. Stability gives the two restrictions, and $hT^r = T'^r h$ proves functoriality. Continuity on a finite-dimensional discrete space means some u^r kills it. Conversely nilpotence makes every formal series act by a finite sum. This proves the completion assertion. □

Over an algebraically closed k, the indecomposable branch pairs are $J_r(\lambda)$. The value $\lambda = 1$ is permitted for a polynomial band; it is not a forbidden deformation parameter. The nilpotent blocks are strings, but are not all rigid: $N_m(J_2(0))$ is a nonsplit self-extension of $N_m(J_1(0))$. The Hom dimension between two such Jordan pairs is $\min(r,s)$ for equal eigenvalues and zero otherwise. These assertions concern this specified full subcategory, not all infinite modules.

If L transports once along the a-arrows on $\bigoplus_i V$, then

$$\det(1 - zL) = \det(1 - z^m T). \tag{7}$$

Eliminating the identity arrow blocks leaves the final monodromy block; the eliminated identity blocks have determinant one. For $J_r(\lambda)$ the answer is $(1 - \lambda z^m)^r$. Nilpotent strings therefore contribute one, and arbitrary bands have no forced unit-circle zero location. This determinant is an actual cyclic-quiver invariant; it is not the missing arithmetic positive determinant comparison.

14 A kernel-checked extension over the infinite dessin algebra

The preceding branch example can be realized on every element of the infinite algebra, rather than verified only on sampled arrow matrices. Use the complex genus-two clean dessin $y^2 = x^5 - 1$,

$\beta = x^5$, and any coefficient field k. Put $H \subseteq M_5(k[t])$ equal to the matrices whose reduction modulo t is upper triangular. Let

$$B = H \times_{k^5} H,$$

where each map to k^5 takes the constant diagonal. Equivalently, B is generated inside $M_5(k[t])^2$ by

$$e_i = (E_{ii}, E_{ii}), \quad a_i = (w_i E_{i-1,i}, 0), \quad b_i = (0, w_i E_{i-1,i}), \qquad w_0 = t, \quad w_i = 1 \ (i \neq 0).$$

Indices are modulo five. Matrix multiplication acts on left column modules. The path-normal-form proof below identifies B with A_5; this is a polynomial algebra, not an arrow-truncated finite quotient.

Theorem 14.1 (An ambient module extension)**.** *Let $D = k[\epsilon]/(\epsilon^2)$, $M_1 = k^5$ and $M_2 = D^5$. For $(C(t), D'(t)) \in B$, act on M_1 by $C(0)$ and on M_2 by $C(\epsilon)$. These are unital B-modules, and*

$$0 \longrightarrow M_1 \xrightarrow{i} M_2 \xrightarrow{p} M_1 \longrightarrow 0, \qquad i(x) = \epsilon x, \quad p(x + \epsilon y) = x,$$

is exact and nonsplit. Moreover B is infinite dimensional over k. The assertions hold in every characteristic, including two.

Proof. Evaluation of polynomials gives both actions. Multiplying $C(\epsilon) = C_0 + \epsilon C_1$ proves B-linearity of i, p; their kernel and image follow coordinatewise. If s were a B-linear section, the constant coefficient of $(se_0)_0$ would be one. But a_0 kills M_1, while $(a_0 s e_0)_4 = \epsilon (se_0)_0$ has nonzero ϵ coefficient, a contradiction. Finally the permitted turn $c = a_1 a_2 a_3 a_4 a_0$ has first component tE_{00}. The unital algebra map $k[t] \to B$, $f \mapsto f(c)$, is injective: its $(0,0)$ first-component entry is $f(t)$. Thus B cannot be finite dimensional. □

In the preceding notation these modules are $N_5(J_1(0))$ and $N_5(J_2(0))$. They are finite torsion modules over an infinite algebra; they are not nonzero lattices over its completed surface order. The nonsplit sequence defines a nonzero Yoneda extension class in the ambient module category. No assertion that it is almost split in that entire category is needed or inferred. In particular it does not supply an arithmetic spectral determinant or a zero-location theorem.

Theorem 14.2 (Unbounded normal-image faithfulness)**.** *In $M_5(k[t])^2$, index the stationary images by i and the positive images by (b, i, j, q), where $b \in \{a, b\}$, $0 \leq i, j < 5$, $q \geq 0$ and $i < j$ or $q > 0$. The positive image is $t^q E_{ij}$ in factor b, with zero in the other factor. Its path length is $5q + j - i > 0$. All these images, including the shared stationary images, are linearly independent over every field. Every finite normal expansion has uniquely recoverable coefficients, without a bound on degree or path length.*

Proof. Read a stationary coefficient from the first factor's constant diagonal; read a positive coefficient from the indicated factor, entry and degree. The index restriction ensures that no positive image contributes to a constant diagonal. These linear selectors are biorthogonal to the images, so applying each selector to a finite expansion recovers its coefficient. □

This theorem is now kernel checked for actual polynomial matrix pairs. The membership and spanning steps are supplied by the next theorem. Comparison with a separately constructed path quotient remains a distinct algebraic formalization task.

Theorem 14.3 (Normal coordinates in the generated algebra)**.** *Let $A \subseteq M_5(k[t])^2$ be the algebra generated by the displayed vertices and arrows, and let $\mathcal{K}$ be the normal index set of Theorem 14.2. Every normal image belongs to A. The actual A-valued finite expansion*

$$E : k^{(\mathcal{K})} \longrightarrow A, \qquad c \longmapsto \sum_{x \in \mathcal{K}} c_x v_x$$

is a linear isomorphism, whose inverse is $(E^{-1}a)_x = \lambda_x(a)$*. Thus the normal images form a basis of the infinite generated algebra, not merely an independent family in its ambient matrix space.*

Proof. Write $P_b(i,j)$ for the shortest positive branch walk from j to i, using a full turn when $i = j$. Its image is $t^d E_{ij}$ in branch b, where $d = 0$ if $i < j$ and $d = 1$ otherwise. The actual arrow product $P_b(i,i)^n P_b(i,j)$ has image $t^{n+d}E_{ij}$ for every $n \geq 0$. Admissibility gives $q \geq d$, so $n = q - d$ proves membership.

The normal span contains the generators and $1 = \sum e_i$ and is multiplicatively closed: stationary factors select endpoints; positive factors either multiply to zero or to $t^{q+r}E_{il}$ in the same branch. When $q + r = 0$, the two strict upper inequalities imply $i < l$, so this output slot is still permitted. Adjoin induction gives spanning of A. The previous coefficient selectors prove uniqueness and the inverse formula. □

All membership, spanning and finite-expansion assertions in this theorem are now Lean checked over arbitrary fields. The formal linear equivalence is not asserted to be an algebra map from an independently defined path quotient, nor a statement about completed orders, derived categories or GRH.

15 Learned fields, width limits and quantum geometry

Learned functions enter this program as candidate generators, physical surrogates and control laws. A network may suggest an arithmetic metric, operator or gluing datum; a successful fit does not establish its positivity, analytic domain or identification with a completed L-function. Those are independent mathematical obligations. This section gives explicit interfaces and executed examples rather than treating "quantum", "tropical" or "infinite width" as a substitute for that comparison.

Proposition 15.1 (A specified quantum-feature width limit)**.** *Let* $x \in \mathbb{R}^d$*,* $w \sim N(0, I_d)$ *and* $b \sim \mathrm{Unif}[0, 2\pi]$ *be independent, and set* $\phi_{w,b}(x) = \sqrt{2}\langle Z\rangle_{R_Y(w\cdot x+b)|0\rangle} = \sqrt{2}\cos(w \cdot x + b)$*. For independent features,*

$$K_m(x,y) = m^{-1}\sum_{r=1}^{m} \phi_r(x)\phi_r(y) \longrightarrow K(x,y) = e^{-\|x-y\|^2/2}$$

almost surely on every specified finite set of pairs. Every K_m *is a positive semidefinite Gram kernel. For a readout* $f_a(x) = m^{-1/2}\sum_r a_r\phi_r(x)$ *with only a trained, its tangent kernel is exactly* K_m*. These statements do not assert the same limit for a jointly trained finite quantum transformer.*

Proof. Uniform-phase integration gives $\mathbb{E}_b[2\cos(u + b)\cos(v + b)] = \cos(u - v)$. The Gaussian characteristic function then gives K. Products are bounded by two, so the strong law applies to each pair and hence any finite union. For any finite coefficient vector c, the Gram quadratic form is $m^{-1}\sum_r |\sum_i c_i\phi_r(x_i)|^2 \geq 0$. Differentiating with respect to a_r gives the final assertion. □

The frozen-kernel squared-loss flow has residual $r(t) = \exp(-K_m t)r(0)$ when the loss normalization is absorbed into time. Null directions do not decay. A complex feature space instead uses the Hermitian kernel $\mathbb{E}[\phi(x)\overline{\phi(y)}]$. A finite coefficient field has neither this Gaussian law nor an ordered positive Hilbert metric; changing algebraic coefficients is a separate operation.

Proposition 15.2 (Tropical limit and a cancellation boundary)**.** *For real* $s_1, \ldots, s_n$ *and* $\tau > 0$*,*

$$0 \leq \tau \log \sum_j e^{s_j/\tau} - \max_j s_j \leq \tau \log n.$$

An attention maximum therefore has a controlled log-sum-exp approximation. It does not by itself define a tropicalization of a variety or an HMS functor. In a valued coefficient field, valuation of a sum can increase by cancellation.

Proof. After subtracting $M = \max s_j$, the exponential sum lies between one and n. For the second statement, both $1+T$ and $-1+T$ have valuation zero in $\mathbb{Q}((T))$, whereas their sum $2T$ has valuation one. In characteristic two that sum is zero. Thus replacing every addition by its tropical minimum without tracking leading coefficients loses information. □

Kontsevich–Soibelman constructions require valued analytic and integral-affine gluing data; appropriate gentle/Fukaya comparisons require specified categories, gradings, stops and smoothness hypotheses [92, 91, 93]. Learned coefficients can propose such data, but the compatibility equations and arithmetic comparison still need proofs.

Executed physical examples. On the actual clean genus-two dessin, put seven quantum spins on its $K_{2,5}$ incidence graph and compute a normalized Gibbs state. Exact white-spin partial trace gives a two-spin quantum RG marginal. Twelve RTX4090 runs train four-qubit-feature diffusion and flow-matching models for its smoothed Z-measurement law: two objectives, entangled/product circuit controls, and three seeds. They have 5,810 nominal parameters and 2,000 steps each. A separate four-qubit variational purification flow learns the full known density interpolation, with 48 controls. Its minimum fidelity on a 33-point grid is 0.999903; this is not a uniform theorem or universal noise inverse. The matched controls show no consistent entanglement advantage.

Actual cuStateVec/cuTensorNet execution also covers 7-, 11- and 23-qubit finite circuits on genus-two, genus-four and genus-ten dessin incidence graphs. The 23-qubit case has 8,388,608 amplitudes and 447 gates. These larger circuits are untrained; the small cases have independent NumPy comparisons, while the largest has normalization and inverse-circuit checks. The exact environment and all failed/preliminary records are preserved.

Geometric and hardware interfaces. For the contracted m-edge dessin, the graph fundamental group is F_{m-1}, and its universal graph cover is the m-regular tree. The theta case $m = 3$ has a three-regular cover, distinct from the four-regular Cayley graph of F_2. Neither is the coefficient field $\mathbb{F}_2$ or the universal cover of the filled compact surface. Explicit path modules over the infinite genus-two algebra give Hom, Ext and Tor computations; Frobenius pants tensors give exact TQFT gluing, with a heralded physical implementation. CHSH gives a finite game benchmark, distinct from the unrestricted correlation separation in MIP*=RE [99]. Three executable tropical hardware references and a source-pinned chip-design survey complement the circuits. They are arithmetic/control designs, not synthesized chips, validated CFT/string dualities or a proof of GRH.

A Expanded dessin, resolution and derived calculations

A.1 The cycle order in coordinates

Number a permitted cycle in travel order $0, \dots, \ell-1$. Map an arrow $i \to i+1$ to $E_{i,i+1}$ for $i < \ell-1$, and the final arrow to $u_c E_{\ell-1,0}$. A path from i to j then gives $u_c^r E_{ij}$, where r counts cut crossings. These are linearly independent path normal forms, and every positive-length permitted path has exactly one such expression on exactly one branch. The full-cycle sum in a branch is $u_c I$. At a paired edge its two diagonal occurrences come from a single original idempotent; no positive-length coefficients are identified. This proves the order description including self-pairings within one cycle.

Although the suborder is generically a product of full matrix algebras, it is not their unglued product at the special fiber.

For the interval this becomes the literal fiber product

$$\Lambda = \{(f,g) \in k[[X]]^2 : f(0) = g(0)\}, \qquad x = (X,0),\ y = (0,X),\ u = (X,X).$$

Every element has a unique expression $a(u) + b(u)x$: take $a = g$ and $b = (f-g)/X$. Thus it is R-free on $1, x$ with $x^2 = ux$ and $y = u - x$. The annihilator of x consists of pairs $(0,g)$ with $g(0) = 0$, exactly $y\Lambda$; the other branch is symmetric. Multiplication by u is injective, coefficientwise. These are actual formal power-series statements, not conclusions from finite truncations.

A.2 The two-arm complex and its calculations

For an arrow a write $f(a)$ for its unique forbidden successor. The summand in $P_n(i)$ indexed by the initial arrow a has endpoint $t(f^{n-1}(a))$. Its map multiplies by that last arrow. Thus the resolution can be computed by iterating a finite permutation, while the projectives themselves remain infinite-dimensional. The Hom complex against a module N is the direct sum of the endpoint spaces with differentials given by the same arrow matrices. Tensor against a left module instead uses the same path entries acting on the left. No rank or exactness claim for an arbitrary N is inferred by counting endpoints alone.

For the nodal residue field the maps are particularly explicit:

$$\cdots \xrightarrow{\mathrm{diag}(x,y)} \Lambda^2 \xrightarrow{\mathrm{diag}(y,x)} \Lambda^2 \xrightarrow{(x \quad y)} \Lambda \longrightarrow k \longrightarrow 0.$$

If $xz_1 + yz_2 = 0$, its first branch says the first branch of z_1 is zero, and its second branch says the second branch of z_2 is zero. Hence $z_1 = yw_1$, $z_2 = xw_2$. The same annihilator calculation proves the alternating kernels in all degrees. Projectivity is genuine: the terms are regular modules or their finite direct sums. It is not a finite free resolution, despite the period of its positive tail.

For the elliptic theta the quiver vertices are $v_0 = \{0,3\}$, $v_1 = \{1,4\}$, $v_2 = \{2,5\}$. The arrows a_0, a_3 go $v_0 \to v_2$, a_2, a_5 go $v_2 \to v_1$, and a_1, a_4 go $v_1 \to v_0$. The two forbidden arms from v_0 are

$$a_0, a_5, a_1, a_3, a_2, a_4, \ldots, \qquad a_3, a_2, a_4, a_0, a_5, a_1, \ldots.$$

Their top pairs are $(v_2,v_2), (v_1,v_1), (v_0,v_0)$, repeated. The three steps exchange the arms. Their distinct top pairs exclude shorter unordered positive periods. The spherical theta instead has forbidden two-cycles and successive top pairs $(v_2,v_1), (v_0,v_0)$, giving period two. Thus even identical hereditary overorders need not give identical resolutions.

Quadratic Koszul duality here has a directly checkable local meaning: relations complement the two possible successors. The dual dessin replaces the graph rotation by ϕ^{-1}. In the Ext algebra against A/J, concatenating the forbidden-word basis gives the product when the join is forbidden; a permitted join gives zero. One can see this by lifting a length-r basis class to the resolution: discard the prescribed initial r letters and identify the remaining summand with its suffix resolution, taking zero on other prefixes. These prefix-discard comparison maps commute with the suffix differentials, with the standard shift signs when viewed as shifted complexes. The concatenated indexing word exists exactly when its join is forbidden; evaluation of the two basis classes on that word gives one. If the join is permitted there is no such indexing word and the product is zero. Composition therefore concatenates indexing words, with the chosen right-module/Yoneda opposite convention. This is a graded algebra statement; it neither makes the unbounded resolution perfect nor constructs an equivalence of all formal-completion derived categories.

For an arbitrary bounded projective complex $P^\bullet$, its determinant line over a commutative coefficient ring is $\bigotimes_n (\det P^n)^{(-1)^n}$, with determinant duals for negative exponents. A chain automorphism acts on this line; a quasi-isomorphism gives the corresponding determinant-functor isomorphism. For a general endomorphism F, the expression for $1 - zF$ uses the locus where it is invertible and gives a rational determinant-line action/trivialization, not an everywhere invertible line map at singular parameters. Additivity is compatible with exact triangles under the usual determinant-functor conventions [17, 18]. In our order context an R-linear determinant uses the finite free underlying R-modules; it must specify a separate endomorphism. Determinants of a differential presentation, alternating determinants of a chain action, regularized analytic determinants and the Cartan path matrix are four distinct constructions. Their equality requires a proof, not their shared name.

The geometric model literature [6, 8, 9] is useful for identifying which crossings and gradings would have to be represented by a functor. To apply it here one must specify the category of polynomial strings, completed lattices or finite-length quotients; construct its surface/curve objects; and prove that degree-r intersections compute the Hom group into shift r, with local systems and boundary exceptions included. We have supplied explicit walks and maps in the examples, not claimed this general completed-order theorem. On a finite word, a map can already be computed rigorously without such a theorem: assign a coefficient matrix between equal-vertex occurrences and solve $F_{t(a)} N(a) = N'(a) F_{s(a)}$ for every arrow. The diamond calculations below implement exactly this system.

B Semi-invariants, tableaux and explicit intertwiners

B.1 Derived information retained by a presentation

Let $K = \operatorname{im} f$. The injection $\operatorname{Hom}(K, N) \to \operatorname{Hom}(P_1, N)$ descends to the Ext injection used in Theorem 3.1; its kernel modulo $\operatorname{im}\operatorname{Hom}(P_0, N)$ is zero because the map $P_1 \to K$ is onto. The missing part of the cokernel detects maps on P_1 not descending through K. This explains exactly why an injective resolution map is needed for its full Euler interpretation.

In the node example the resolution of $M = \Lambda/(x)$ has maps $x, y, x, y, \ldots$. On $N = k[y]/(y^2)$ the Hom cochain maps are $0, y, 0, y, \ldots$. Thus $\operatorname{Hom}(M, N) = N$ and $\operatorname{Ext}^1(M, N) = \ker y$ have dimensions two and one; the cokernel of the first zero map is two-dimensional. In the elliptic-square example the first four Hom maps are $1, 0, 0, 0$, periodically. Hence degrees zero and one vanish, degrees two and three are k, and degree four vanishes. This also makes clear why a nonzero scalar semi-invariant does not give a derived orthogonality theorem.

With coordinate action $(gF)(N) = F(g^{-1}N)$ the Schofield weight is $a - b$, the negative of the point-evaluation weight. For a single arrow it is the same elementary distinction as the action on $\operatorname{Hom}(V, W)$ versus on its coordinate functions $V \otimes W^*$. The Cauchy decomposition for coordinate functions is

$$\operatorname{Sym}^m(V \otimes W^*) \cong \bigoplus_{\lambda \vdash m} S_\lambda(V) \otimes S_\lambda(W)^*.$$

A relation quotient can kill these components; its ambient multiplicities do not descend unchanged. Acyclic-quiver semi-invariant/LR results such as [35] retain their acyclicity assumptions. The explicit two-arrow slice below is acyclic, whereas the whole theta quiver is not.

B.2 One Young operator and one LR decomposition in coordinates

For $\lambda = (2,1)$ number the first row $1,2$ and second row 3. Then

$$Y = (I + P_{12})(I - P_{13}), \quad Y^2 = 3Y, \quad \dim \operatorname{im}(Y \mid (k^2)^{\otimes 3}) = 2.$$

The hook product is three; its Specht dimension is $3!/3 = 2$, also two here for a different reason. The normalized projector extracts one $S_{(2,1)}$ copy. The trace $\operatorname{tr}((Y/3)M^{\otimes 3})$ is therefore the Schur character, not twice that character. This fixes the common row/column normalization ambiguity.

Write $t = \operatorname{tr} M$, $\Delta = \det M$ for a two-dimensional endomorphism. Its complete characters obey $h_0 = 1$, $h_1 = t$, $h_m = t h_{m-1} - \Delta h_{m-2}$, and

$$\chi_{(a,b)}(M) = \Delta^b h_{a-b}(M), \quad a \geq b \geq 0.$$

In particular $h_2 = t^2 - \Delta$, $h_3 = t^3 - 2t\Delta$, and $\chi_{(2,1)} = t\Delta$. Thus $t h_2 = h_3 + \chi_{(2,1)}$ is the LR identity $V \otimes \operatorname{Sym}^2 V \cong \operatorname{Sym}^3 V \oplus S_{(2,1)}V$, with $S_{(2,1)}V \cong \det V \otimes V$.

An explicit natural splitting avoids relying just on characters. In characteristic zero multiplication $m : V \otimes \operatorname{Sym}^2 V \to \operatorname{Sym}^3 V$ has natural section

$$s(v_1 v_2 v_3) = \tfrac{1}{3} \sum_{i=1}^{3} v_i \otimes \prod_{j \neq i} v_j.$$

The map $j : \bigwedge^2 V \otimes V \to V \otimes \operatorname{Sym}^2 V$ sends $(v \wedge w) \otimes z$ to $v \otimes wz - w \otimes vz$. In dimension two it is injective: for a basis e_1, e_2 its images at $z = e_1, e_2$ have distinct independent coordinates. They lie in $\ker m$, whose dimension is $6 - 4 = 2$. Hence (s, j) is the required isomorphism. Tensorial definitions make both maps commute with every linear endomorphism, including singular arrow maps. This is a genuine LR intertwiner on our dessin representation stratum.

For a concrete theta representation take

$$X_1 = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad X_2 = \begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}, \quad X_3 = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}.$$

Then $M = X_3 X_2 X_1 = \left(\begin{smallmatrix} 2 & 2 \\ 2 & 3 \end{smallmatrix}\right)$, $t = 5$, $\Delta = 2$, $h_2 = 23$, $h_3 = 105$ and $\chi_{(2,1)} = 10$. The identity is $115 = 105 + 10$. Since the other permitted branch is zero, every forbidden product is zero; this is an actual representation of the infinite elliptic dessin algebra.

In the invertible-arrow open stratum one can gauge X_1, X_2 to the identity, leaving conjugation of M. The invariant-ring proof in the core works also at singular arrows by cycle-trace generation, not by assuming that this gauge choice extends there. The degree-one monodromy coefficients have total arrow degree three; this is why the Hilbert-series denominator uses $3j$, not j.

For the Cauchy kernel choose the Schur subspace inside the tensor power with its inherited Hilbert norm. $M^{\otimes m}$ preserves it and has norm at most $\|M\|^m$. This proves the trace bound for nonnormal matrices as well as diagonal ones. Formal Cauchy identities extend by polynomial continuity in matrix entries; the stated absolute bound then justifies analytic summation. Gram positivity does not assert entrywise positivity or positivity of an unrelated arithmetic form. For three exact test matrices M, $\left(\begin{smallmatrix} 1 & 1 \\ 0 & 2 \end{smallmatrix}\right)$, $\left(\begin{smallmatrix} 0 & 1 \\ -1 & 1 \end{smallmatrix}\right)$, the choice $r = 1/100$ satisfies $r R_0^2 \leq 21/100 < 1$ using Frobenius norms, and all seven principal minors of the rational kernel matrix are nonnegative.

B.3 Two precise functorial limitations

At any actual vertex simple S of A, the split exact sequence $0 \to S \to S \oplus S \to S \to 0$ uses first inclusion and second projection. Termwise Sym^2 has dimensions $1, 3, 1$ at that vertex; its middle kernel contains the mixed monomial as well as the first square, but its first image contains only the first square. Middle homology is S. Thus even split exactness is not preserved. Genuine derived Schur constructions instead use mixed tensor degrees, signed total differentials and suitable flat/projective replacements [32, 33]; their membership in a particular order category must still be checked.

The R-lattice $M = Re_x \oplus Re_y$ with $x = \mathrm{diag}(u, 0)$, $y = \mathrm{diag}(0, u)$ has $x + y = uI$. On $\mathrm{Sym}^2_R M$ the images are $\mathrm{diag}(u^2, 0, 0)$ and $\mathrm{diag}(0, 0, u^2)$. Their sum kills the mixed vector, hence equals neither uI nor $u^2 I$. This is a specific failure to retain the original central action, not a failure of the polynomial monomial representation functor. Even a common ramified base substitution does not repair this mixed vector.

Semi-invariants retain useful moduli on nilpotent modules. In the theta quiver keep only $a_0, a_3 : v_0 \to v_2$, put their scalar values $(x, y) = (1, \lambda)$ and all other arrows zero. The v_1 simple splits off. Every length-two path acts by zero, so these are continuous completed modules. The two scalars have the same weight g_{v_2}/g_{v_0}, and y/x is a rational invariant on $x \neq 0$, giving the projective pencil $[x : y]$. It is not a regular invariant on the whole affine locus, whose orbit closures meet zero.

For the interval quotient take $x = J_2$, $y = \lambda J_2$. With its top and socle specified, the separated Kronecker coordinates are again $(1, \lambda)$; the relation to the same-weight ratio is exact. The pp formula in the core is stronger than merely noticing a ratio: on M_λ its solution set is $\mathrm{im}\, x$, whereas on M_μ, $\mu \neq \lambda$, the equation $(\mu - \lambda)xv = 0$ forces $z = xv = 0$. Different parameters are therefore distinguishable by actual positive-primitive conditions even though every cycle trace is zero.

C Full lattice and diamond diagrams

C.1 The nodal exact diagram

The normalization proof of Theorem 4.1 uses BM inside the generic module. Replacing it without justification by $B \otimes_\Lambda M$ could introduce torsion and would not prove the same statement. Once the subdirect residue subspace is decomposed, the branch basis changes lift over complete DVRs; taking inverse images modulo the conductor gives the direct decomposition, not just a decomposition of residues.

Here is the promised diagram, with the terminal short exact column for every position:

$$\begin{array}{ccccccccc}
0 & \longrightarrow & Y & \hookrightarrow & \Lambda & \xrightarrow{x} & X & \longrightarrow & 0 \\
 & & \downarrow{\scriptstyle u} & & \downarrow{\scriptstyle u} & & \downarrow{\scriptstyle u} & & \\
0 & \longrightarrow & Y & \hookrightarrow & \Lambda & \xrightarrow{x} & X & \longrightarrow & 0 \\
 & & \downarrow & & \downarrow & & \downarrow & & \\
0 & \longrightarrow & Y/uY & \hookrightarrow & \Lambda/u\Lambda & \longrightarrow & X/uX & \longrightarrow & 0
\end{array}$$

Each vertical column is completed by zero above and below. In the bottom row identify $Y/uY = k$ via generator y, $X/uX = k$ via x, and $\Lambda/u\Lambda = k[\epsilon]/(\epsilon^2)$ with $x = \epsilon$, $y = -\epsilon$. Its first map is $1 \mapsto -\epsilon$, its second the augmentation. It is nonsplit as a dual-number sequence. This explicit order reduction, not an unrelated finite algebra, explains its appearance.

For a further Ext calculation, resolve $X \cong \Lambda/(y)$ with successive maps $y, x, \ldots$. On Y the first is multiplication by y and the second zero, so $\mathrm{Ext}^1(X,Y) = Y/yY = k$; on X the first is zero and the second injective, so $\mathrm{Ext}^1(X,X) = 0$. Interchanging branches gives the other two cases. The two nonsplit lattice rows realize the nonzero cross classes. Their Hom spaces are finite over R, generally infinite over k.

C.2 Chebyshev geometry and the exact quotient

The identity $1 - T_n(z)^2 = (1-z^2)U_{n-1}(z)^2$ follows from $z = \cos\theta$ and the sine/cosine multiple-angle formulas, hence as a polynomial identity. $T_n' = nU_{n-1}$. Therefore β_n has its only zero at infinity, of order $2n$; its n points over one are double since $\beta_n - 1 = T_n^2/(1-T_n^2)$; and its poles are simple at ± 1 and double at the $n-1$ interior roots of U_{n-1}. The derivative has no additional critical values. The total ramification is $(2n-1) + n + (n-1) = 4n-2$, as required for a degree-$2n$ rational Belyi map. The dual graph has $n+1$ vertices of degrees $1, 1, 2, \ldots, 2$, and n edges; it is a connected path. Its planar ribbon embedding is unique, so this identifies the particular nested-loop dessin, not only its passport.

In the quotient retaining orientation Ω, any surviving path traverses each chain edge at most once and either endpoint loop at most once. Its length is at most $n+1$; thus the killed-arrow ideal contains J^{n+2}. The quotient of A_n and of its completion is the same $H_n(\Omega)$. For $n = 3$ with forward chain arrows, its path counts in lengths zero through four are $3, 4, 3, 2, 1$, total thirteen. These counts describe a quotient representation category of the infinite order, not a change of primary object.

Inflation is full because the two algebras have exactly the same actions on these annihilated objects. It preserves internal exact sequences. It does not preserve arbitrary ambient extension closure: the two adjacent simples are annihilated by the killed reverse arrow, but the two-dimensional extension having only that arrow nonzero is not. Likewise endpoint-local freeness over $k[l]/l^2$ or $k[r]/r^2$ in [46] is not R-freeness. To transfer an almost-split property to a larger category one must prove the requisite approximation/factorization statements; locally finite quiver criteria [40, 41, 42] do not apply just because each vertex has two arrows. The permitted cycles violate interval/Hom finiteness.

C.3 Nine actual strings and all maps

Take $n = 3$ with $1 \xrightarrow{a} 2 \xrightarrow{b} 3$, loops l, r and $l^2 = r^2 = 0$ in the quotient. Here b denotes the retained forward a_2, not a reverse arrow of the original doubled chain. The following words are in travel order. A superscript $-$ means a formal inverse in a word, not an inverse algebra element.

Object	word	start	dimension vector
A	$l^- abrb^- a^- l^-$	1	$(4,2,2)$
B	$l^- ab$	1	$(2,1,1)$
C	$l^- abr^-$	1	$(2,1,2)$
D	$abrb^- a^- l^-$	1	$(3,2,2)$
E	ab	1	$(1,1,1)$
F	abr^-	1	$(1,1,2)$
G	$r^- b^- a^- labrb^- a^- l^-$	3	$(4,3,4)$
H	$r^- b^- a^- lab$	3	$(2,2,3)$
I	$r^- b^- a^- labr^-$	3	$(2,2,4)$

A basis vector is assigned to each visited position $0, \ldots, \ell(w)$ at its vertex. A direct letter sends its preceding vector to its following vector; an inverse letter sends the following vector to the preceding

vector. Sum these elementary maps for each arrow and put all unmentioned entries zero. This defines every matrix, including repeated loops, and verifies the relations. Set the killed reverse arrows to zero to obtain actual Λ_3-modules.

Their signed grid is

$$\begin{array}{ccccc} A & \xrightarrow{p} & B & \xrightarrow{i} & C \\ {\scriptstyle -p}\downarrow & & \downarrow{\scriptstyle p} & & \downarrow{\scriptstyle -p} \\ D & \xrightarrow{p} & E & \xrightarrow{i} & F \\ {\scriptstyle -i}\downarrow & & \downarrow{\scriptstyle i} & & \downarrow{\scriptstyle -i} \\ G & \xrightarrow{p} & H & \xrightarrow{i} & I. \end{array}$$

Horizontal projections keep the initial word positions and kill the remaining positions. Horizontal inclusions append the final r^-, keeping old positions. Upper vertical projections delete the initial l^-: old position $j \geq 1$ maps to position $j-1$, and old position zero maps to zero. Lower vertical inclusions prepend $r^-b^-a^-l$: position j maps to $j+4$. Multiply these maps by the indicated signs. For every arrow these formulas commute with the string action, including the two positions adjacent to a cut or join; those are exactly where the loop inverse and chosen orientation matter.

Every small square has two length-two composites whose sum is zero. The exact sequences are

$$0 \to A \to B \oplus D \to E \to 0, \qquad 0 \to B \to C \oplus E \to F \to 0,$$
$$0 \to D \to E \oplus G \to H \to 0, \qquad 0 \to E \to F \oplus H \to I \to 0.$$

For example the maps out of A retain its initial positions $0,1,2,3$ in B and delete its first position in D, so their joint kernel is zero. The maps to E are onto, and the dimension vectors give $\dim A + \dim E = \dim B + \dim D$ vertexwise. The zero composite plus this equality proves exactness. The other three use precisely the same keep/delete or append/prepend rules; their vertexwise dimension equalities and jointly injective/surjective maps prove exactness. The shipped exact checker constructs all matrices and verifies these assertions independently over $\mathbb{Q}$.

The stronger almost-split property is imported by identifying $E = w = ab$ as the minimal string of type $(2,2)$ and these exact hook/cohook words with [46, Proposition3.12], using that paper's Propositions2.6/2.9 and Lemma3.10. Opposite-module conventions are reconciled by reversing travel/composition conventions; the closed doubled quiver is isomorphic to its opposite by exchanging the two arrows at each chain edge. Thus intrinsically $\tau E = A$, $\tau F = B$, $\tau H = D$, $\tau I = E$. The rows and columns of this grid are *not* generally short exact, unlike the nodal diagram. Two small cautions in that source are not needed here: a tube-rank label in Example3.9(3) is inconsistent with its listed mouth, and a general homogeneous-tube statement must not be read as a converse classifying every such object as a band. Neither is used to import these four meshes.

C.4 The pure-injective quotient and what its dimension means

For $B_0 = \Lambda/J^2 = k\langle x,y\rangle/(x,y)^2$, words alternate direct and inverse letters because every direct/direct and inverse/inverse adjacent pair is forbidden. Up to rotation and inverse its only primitive band is xy^{-1}; the bridge quiver has two oriented band vertices with no connecting nontrivial bridge. The longest path has length zero. The domestic string results [44, 45] therefore give relative dimension $0+2=2$, not an automatic dimension three for every one-domestic algebra. Finite-dimensional points have relative CB rank zero; the appropriate Prüfer/adic and one-sided periodic string points have rank one; the unique generic point has rank two. These statements concern the quotient category with its inherited pp structure.

Indeed a pp formula is a finite system of linear equations with existentially quantified unknowns. On a module annihilated by J^2 every coefficient reduces to B_0, so purity and pure-injectivity are preserved in both directions under inflation. The condition $MJ^2 = 0$ is a definable closed condition, giving the closed Ziegler subspace. Its relative ranks need not be ranks in the entire Λ spectrum. This subcategory is not extension closed: $\Lambda/(x^3, y)$ gives an extension of objects killed by J^2 that is not itself killed by J^2. Since u^2 annihilates every object of the subcategory, it contains no nonzero R-free lattice. Ringel's representation-theoretic node construction is also not a Morita equivalence identifying the geometric nodal order with the Kronecker algebra. Infinite-dimensional polynomial string classifications [38] retain finitely-controlled or pointwise-artinian assumptions; an arbitrary complete lattice does not automatically satisfy them as an algebraic A-module.

D The operator extension in detail

D.1 A corrected cycle lift and the integer boundary map

In the permitted-state graph each length-ℓ cycle has a full matrix-corner Laurent/circle coordinate. Its chosen K_1 generator is the full cycle at one state plus the complementary identity. In Fock space $L_c^* L_c = Q_a + v_{s(a)}$. Thus the tempting lift $L_c + (1 - Q_a)$ has overlapping initial pieces and is not an isometry. The corrected lift in Theorem 5.1 first restricts to Q_a:

$$W = L_c Q_a + (1 - Q_a), \quad W^* W = I, \quad I - WW^* = Q_a - L_c Q_a L_c^*.$$

A path starting with a and length greater than ℓ has uniquely the full prefix c followed by another path starting with a. The missing range consists exactly of lengths $1, \dots, \ell$. Its terminal multiset is the original graph's incident-edge multiset at that cycle vertex. These finite defect vectors all lie in terminal compact ideals; their classes give the claimed unsigned incidence entries. The sign convention affects only the overall sign of the boundary map.

Here is the Smith calculation without spectral assumptions. Choose a spanning tree of the connected underlying graph and one root. By changing signs on one bipartition of the tree, its $E \times V$ incidence submatrix becomes the ordinary oriented tree incidence, with $V - 1$ unit pivots. After eliminating these pivots, a remaining column represents the common root coordinate, and each nontree edge evaluates on it as zero when its fundamental cycle is even, or ± 2 when odd. A loop is the latter case. If all are even the graph is bipartite and this column is zero; otherwise its entries have gcd two, giving the final pivot two. Remaining rows are free. Consequently the kernel/cokernel, then the six-term sequence, give the K groups stated in the core. No theorem reconstructing a cyclic embedding from those groups is asserted.

The compact ideal is block diagonal, not $\mathcal{K}(\mathcal{H})$: prefixes never move terminal vertices, so cross-terminal rank-one maps do not belong to the displayed ideal. This is why the arithmetic insertion must choose one terminal sector. A graph/Cuntz boundary quotient kills the vacua and hence this ideal, so that insertion need not survive passage to the quotient.

D.2 Spectral domain and arithmetic trace

A vector belongs to the adjoint domain of the diagonal N precisely when its coefficientwise product by the real sequence $|p|$ is square summable; this proves self-adjointness on the given domain. The truncations by length prove finite supports are a core. Resolvent coefficients $(|p| - i)^{-1}$ tend to zero and each length space is finite dimensional, giving compactness. Prefix and deletion shift the length by at most one, so preserve the domain and prove the commutator formulas there. Leibniz then handles the dense polynomial *-algebra.

Every terminal sector has one vacuum and two paths at each length $n \geq 1$, even when some graph endpoints coincide: distinct formal paths remain distinct basis vectors. Thus restriction to a sector still has a simple vacuum, but it need not be a faithful representation of the whole concrete algebra. The trace of e^{-tN^2} is $E + 2E\sum_{n\geq 1} e^{-tn^2}$, distinct from the semigroup trace of N in the core. The trace identity with $\chi(N)$ is only on $\operatorname{Re} s > 1$, where the product is trace class. Analytic continuation of the scalar L-function does not make that same operator product trace class on $\operatorname{Re} s > 0$.

E Abel determinant estimates and the arithmetic program

For completeness the inserted family has a full elementary convergence proof. Since complete nonprincipal character periods sum to zero, $|A_n| \leq B_0 \leq q_\chi$. On $2^k \leq n < 2^{k+1}$,

$$\sum |u_n|^2 \leq \frac{B_0^2}{(k+1)^2}, \qquad \sum |v_n(s)|^2 \leq M^2(k+1)^2 r^k, \quad r = 2^{-2\sigma_0} < 1.$$

The second follows from $n^{-s} - (n+1)^{-s} = s\int_n^{n+1} x^{-s-1}\,dx$. Define

$$Q_J(r) = r^J\left(\frac{(J+1)^2}{1-r} + \frac{2(J+1)r}{(1-r)^2} + \frac{r(1+r)}{(1-r)^3}\right).$$

Expanding $(J+j+1)^2$ in the three geometric moment sums proves $Q_J = \sum_{k\geq J}(k+1)^2 r^k \to 0$. Locally uniform ℓ^2 convergence of the holomorphic finite-coordinate vectors proves Banach holomorphy. The linear functional $z \mapsto \sum v_n z_n$ has norm $\|v\|$; this is the complex-linear coefficient pairing, not a conjugate-linear dependence on s.

Write $\mathcal{I}(v)(z) = \sum_n v_n z_n$ for the coefficient functional, distinct from the arrow ideal J. For $N_J = 2^J - 1$, $J \geq 1$, let $K_N = u \otimes (\mathcal{I}(P_N v) - e_1^*)$. Then

$$\|K - K_{N_J}\|_1 \leq \sqrt{2} B_0 M \sqrt{Q_J(r)}.$$

Finite summation by parts gives

$$T_N = \det_F(I + K_N) = \sum_{n\leq N} A_n(n^{-s} - (n+1)^{-s}) = S_N - A_N(N+1)^{-s}.$$

Adding the last endpoint times e_1^* to the functional recovers S_N exactly because $u_1 = 1$. Its trace-norm error is at most

$$\sqrt{2} B_0(M\sqrt{Q_J(r)} + B_0(N_J+1)^{-\sigma_0}).$$

For arbitrary cutoffs use the last whole dyadic block for an enclosing tail. Galerkin compressions also converge: $\|u - P_{N_J}u\| \leq B_0/\sqrt{J}$. With $C_K = 1 + M\sqrt{(1+r)/(1-r)^3}$,

$$\|K - P_{N_J} K P_{N_J}\|_1 \leq \frac{B_0 C_K}{\sqrt{J}} + \sqrt{2} B_0 M \sqrt{Q_J(r)}.$$

The raw endpoint adds the same endpoint bound as before. Isometric insertion by U preserves nonzero singular values, every norm estimate, and the Fredholm determinant. Everything is in one compact ideal of the actual infinite dessin Fock algebra.

For primitive parity a use $g(s) = (q_\chi/\pi)^{(s+a)/2}\Gamma((s+a)/2)$, holomorphic nonzero on $\operatorname{Re} s > 0$. The completed functional is $g(s)\mathcal{I}(v) - e_1^*$, and the completed raw functional is $g(s)(\mathcal{I}(P_N v) + A_N(N+$

$1)^{-s}e_1^*) - e_1^*$. Their determinants are exactly gL and gS_N. Scaling the entire old perturbation would instead give $1 + g(L-1)$, the wrong normalization. Trace-class background is standard [26], while these special rank-one identities follow directly from $\det_F(I + u \otimes \varphi) = 1 + \varphi(u)$.

The metric obstruction is likewise elementary. With inner product conjugate-linear in its first variable, symmetry of $T = u \otimes \varphi$ implies

$$\varphi(z)\langle u, u\rangle = \overline{\varphi(u)}\langle u, z\rangle.$$

A nonzero φ cannot have $\varphi(u) = 0$; thus all symmetric maps with that range have proportional functionals. For $n \geq 2$ the coefficient ratios at $\sigma + it$ and σ satisfy

$$\frac{v_n(\sigma + it)}{v_n(\sigma)} = \frac{\sigma + it}{\sigma}\, n^{-it}(1 + O_{\sigma,t}(1/n)).$$

Integers nearest $\exp(2\pi k/|t|)$ and $\exp((2\pi k + \pi)/|t|)$ give opposite nonzero subsequential limits. Hence no common positive metric symmetrizes this family for all critical heights. Restricting any proposed metric on the inserted family to $U\ell^2$ gives the same contradiction. An independently constructed positive comparison is still a possible different route.

E.1 Precisely how classical and recent sources enter

Over a finite field the Grothendieck trace-formula architecture writes a zeta function as alternating determinants on finite-dimensional cohomology; Deligne's purity supplies the weights of eigenvalues [22, 20, 21]. If a weight-w Frobenius is $q^{w/2}$ times a positive-metric unitary, each eigenvalue has modulus $q^{w/2}$. The norm conclusion is immediate; constructing that form or proving purity is the deep step. Symplectic/alternating duality $F^t J F = q^w J$ alone only pairs reciprocal eigenvalues and does not give their individual moduli. An order resolution or Schur-positive character expansion alone cannot replace this step in number-field cohomology.

The recent primary comparisons used here are specific:

- [53, Theorem3.12, pp.15–17] gives finite abelian covers with local orbit length $\log p$ and genuine arithmetic monodromy. It suggests what a descent-equipped order functor should match; it does not identify our u or a face cycle with a prime without such a functor.

- [54, Propositions4.1–4.3, Theorem4.6] builds finite-prime local models with archimedean Fourier factors. Its Hilbertian isomorphisms are not an asserted uniformly controlled common isometry for our arithmetic approximants. [55, Proposition3.1] gives determinacy of a particular local moment problem; positive finite Hankel matrices are not the missing global Weil inequality.

- [56, Theorem3.6] gives discreteness/lower boundedness for the stated restricted Weil form. Theorem5.10, pp.23–24, additionally assumes a finite truncation, $\lambda > 1$, a simple even bottom eigenfunction ξ and $\delta_N \xi = 1$. Its quotient metric is positive and its regularized determinant is $-i\lambda^{-iz}\widehat{\xi}(z)$. Passing to the infinite arithmetic comparison, including the ground-state and convergence conditions in its §8, is not automatic.

- [57, Theorem6.1, p.13] assumes a specified real distribution on an interval, the corresponding trigonometric-core form, essential self-adjointness, and an isolated simple lowest eigenvalue with reflection-even eigenfunction. Its Fourier transform then has only real zeros. Our degree operator's vacuum and length parity are not those data.

- [60, Theorem6.1] gives prime-indexed periodic packets in its dynamical construction, not one proved orbit per prime with the full desired cohomological trace formula. [62] isolates an obstruction to a particular real-coefficient cohomology with full functorial Artin multiplicities; it is not a theorem excluding every Hilbert–Pólya operator. The current survey [58] distinguishes remaining steps. The recent Picard-monoid construction [59] is not already a polarized abelian variety with a number-field purity theorem. [63] uses RH as an input in the cited spectral identification, so cannot be used circularly as its proof.

These are targeted theorem/proof readings, not claims that every page of every cited work has been audited. The preserved review records specify actual page coverage and hashes, including inaccessible original PDFs where the modern primary proof was used instead.

E.2 A concrete sequence of remaining constructions

First choose a number-field model of a Belyi pair, its finite covers and conductor/ramification data. Construct actions or exact correspondences on the labelled order/module or analytic categories that recover the local prime Frobenius polynomials. The elliptic twists show why omitting this datum loses information. Test at more than one arithmetic model of the same complex dessin, not just its topological passport.

Second specify a finite projective or otherwise determinant-admissible complex and its *independent* arithmetic endomorphism. Prove exact-sequence/determinant-line compatibility. Use semi-invariants to encode presentation degeneracies, and Schur/exterior characters to calculate local factors, retaining every central-action and nilpotence distinction above. General derived Schur replacements require their own construction; termwise Schur on the resolution fails the exhibited split sequence test.

Third prove the same local factors, gamma terms and conductor give a globally convergent or correctly regularized determinant. Trace-norm convergence must be established for the chosen analytic perturbations; strong convergence is insufficient in general. The inserted Abel family supplies this topology for an elementary scalar encoding, but has neither the independent arithmetic spectrum nor the positive metric needed for zero location.

Finally produce an independently proved positive form or self-adjoint determinant-preserving realization. One acceptance criterion is the Hurwitz target in the core; another is equality with the precise arithmetic quadratic form in the recent spectral theorems, with all their domain, ground-state, normalization and limit hypotheses verified. The local Cauchy kernel, K-group computation and graph spectral dimension alone meet none of these final comparison hypotheses. No remaining condition is silently asserted true here. Dirichlet, Hecke, Dedekind and Artin scopes must remain individually recorded.

F Graph pullbacks and completions with specified fibers

F.1 The complete Fock graph presentation

For graph C* conventions the edge operator has initial projection at its range vertex and final projection under its source vertex. Put

$$S_h = L_{a_h} Q_{a_{\tau h}}, \qquad T_h = L_{a_h} v_{[\tau h]}.$$

The two right-hand projections are orthogonal subprojections of $L_{a_h}^* L_{a_h} = Q_{a_{\tau h}} + v_{[\tau h]}$. Thus $S_h^* S_h = Q_{a_{\tau h}}$, $T_h^* T_h = v_{[\tau h]}$, and $S_h^* T_h = 0$. For distinct h the prefix ranges are orthogonal, so every other cross-edge CK1 product is zero. Their final projections sum to Q_{a_h}. The vacua have

no outgoing edges. The resulting graph has $|H| + |Q_0|$ vertices and $2|H|$ edges, with cycles only among its state vertices and exits to the paired sinks. Gauge covariance and nonzero vertices prove faithfulness as well as surjectivity. This explains precisely how a graph C* algebra is obtained without imposing the incompatible original-quiver CK1 relations.

The graph integer map has one column for each regular vertex h and no sink columns:

$$d : \mathbb{Z}^H \longrightarrow \mathbb{Z}^H \oplus \mathbb{Z}^{Q_0}, \qquad d(e_h) = e_h - e_{\tau h} - b_{[\tau h]}.$$

The state part of $d(x) = 0$ says $x_h = x_{\tau^{-1}h}$; the sink part then says the constant cycle coefficients are in $\ker B_\Gamma$. For the cokernel choose a base state on each cycle and successively use $q_h = q_{\tau h} + b_{[\tau h]}$ to eliminate all other states. Closing the cycle leaves $\sum_{h \in c} b_{[h]} = 0$, while its base state remains free. Hence

$$\operatorname{coker} d \cong \mathbb{Z}^{\#\,\mathrm{cyc}\,\sigma} \oplus \operatorname{coker} B_\Gamma.$$

This gives an independent integral check of the previous compact-extension computation. The full labelled graph, diagonal projections and edge families carry more information than those coarse groups.

F.2 A genuine application of the supplied pullback theorem

For the interval dessin let $E_2 = G_\alpha$ have loop vertices q_x, q_y, common sink b, loops s_x, s_y and exits t_x, t_y. Remove b to obtain F_2, two disjoint loop graphs. Its quotient is exactly $C(\mathbb{T})^2$. Let E_1 have the same three vertices and edges $e_{x,n} : q_x \to b$, $e_{y,n} : q_y \to b$, $n \geq 0$, and no other edges. Remove b to obtain the edgeless F_1, whose algebra is $\mathbb{C}^2$.

The inclusions $F_i \subset E_i$ are admissible in the precise sense of [65, Definition3.1]: the removed sink set is hereditary/saturated; all edges ending at retained vertices are retained; no vertex emits infinitely many edges into the removed set and a positive finite number outside it. For E_1 the latter outside count is zero, which is allowed. Define a path functor identity on objects by

$$e_{x,n} \longmapsto s_x^n t_x, \qquad e_{y,n} \longmapsto s_y^n t_y.$$

Images of distinct positive edges are prefix incomparable: at the first different length one has an exit and the other a loop. Source vertices and identities cause no exception. The image consists of all positive pointed paths in E_2, whose last edge is not a short loop; vertices are included separately. The first graph is acyclic and the removed sink has no short loop in the second. All hypotheses of Theorem3.4 are now verified, giving

$$C^*(E_1) \cong \mathcal{T}_A \underset{C(\mathbb{T})^2}{\times} \mathbb{C}^2.$$

The quotient of $C^*(E_1)$ by its sink ideal is $\mathbb{C}^2$, and its sink ideal is compact operators on the paths ending at b. The graph is acyclic, hence its algebra is AF. There are no regular vertices in E_1: the two sources are infinite emitters and b is a sink. The graph K matrix has no columns and three rows, yielding $K_0 = \mathbb{Z}^3$ and $K_1 = 0$.

This is a positive instance of the new source inside our infinite dessin framework. It is a constants-symbol subalgebra of the Fock model, not the entire Fock algebra or the formal order. For a permitted cycle of length greater than one, its full closed path is itself pointed in this paper's convention, so an object-preserving acyclic path functor cannot cover all such paths. That simple obstruction prevents copying this example to every cycle by analogy. The sink-extension Theorem4.3 also requires a specified original pullback, injective gluing maps, annihilation of glued vertex projections and coverage of all paths ending at the gluing sinks. A two-to-one identification of sink occurrences within one branch does not satisfy these assumptions without another construction.

F.3 Diagonal residue obstruction and the corrected pullback

Suppose one tries to glue full matrix-circle blocks by their diagonal values. In M_2, $E_{12}E_{21} = E_{11}$, although both factors have zero diagonal. Thus the proposed boundary map is not a homomorphism. For $\ell > 1$, more generally any *-homomorphism from the full matrix-circle algebra to a commutative C*-algebra has equal images of $E_{ii} = E_{ij}E_{ji}$ and $E_{jj} = E_{ji}E_{ij}$. These images are both orthogonal and equal, hence zero; summing the diagonal units kills the image of one and then the whole homomorphism. This works for nonunital *-maps as well: a nonzero image of one would give a nonzero unital corner map. The failure concerns full matrix-circle blocks, not the already-restricted fiber algebra D_ℓ.

Evaluation $D_\ell \to \mathbb{C}^\ell$ is a genuine surjective *-homomorphism since at one both factors are diagonal. Its kernel is $C_0(\mathbb{T} \setminus \{1\}, M_\ell)$. Pulling it back along the repeated-coordinate embedding $\Delta_\alpha : \mathbb{C}^{Q_0} \to \mathbb{C}^H$ gives exactly the fiber conditions of $\mathcal{D}_H$. The paired constant diagonal matrices give a unital *-section. Hence this is a split C*-extension, not a claim deduced from a merely commutative diagram of nonmultiplicative maps. Since $\mathbb{T} \setminus \{1\} \cong \mathbb{R}$, each kernel block is a matrix suspension of $\mathbb{C}$, with $K_0 = 0$, $K_1 = \mathbb{Z}$. Split exactness now proves Theorem 5.3.

The pairing is retained in this specified presentation, not recovered from the unmarked isomorphism class. In particular arbitrary constant permutation matrices can relabel a branch's diagonal positions, so the spherical and elliptic theta examples have isomorphic unmarked $\mathcal{D}_H$ despite different ribbon orderings. Keep the labelled arrows, cycle cuts and distinguished fiber as additional data when using it as a dessin model. Also the central branch unitary zI_ℓ represents ℓ times the primitive corner generator in K_1, not a new independent generator per matrix entry.

F.4 Generation, faithfulness and the exact failure of extension

Let ρ denote the polynomial representation in Theorem 5.3. Its vertex images are orthogonal projections summing to one. Every arrow image lies under the correct source and range vertex projections. A forbidden product vanishes because the intermediate state indices are τh and $\alpha\tau h$, which are distinct. Thus ρ is an actual algebra map, not a fitted characteristic polynomial.

For a full turn starting at h, $\rho(c_h) = w_c E_{hh}$ with $w_c(z) = g(z)^{|c|}z$. It is a normal element of $\mathcal{D}_H$. The function w_c vanishes exactly at one; away from that point $w_c/|w_c| = z$. Therefore it is injective on all of $\mathbb{T}$, and a homeomorphism to its image. Nonunital continuous functional calculus at $\rho(c_h)$, using functions zero at zero, gives every fE_{hh} with $f \in C(\mathbb{T})$, $f(1) = 0$. The zero condition makes the result vanish in all other matrix entries and branches.

To obtain an off-diagonal entry choose the unique shortest permitted path from state h to j within its cycle, of length $r > 0$ and cut-crossing count $\kappa \in \{0, 1\}$. Its image is $g^r z^\kappa E_{hj}$. Given $f(1) = 0$, multiply it on the left by the already available diagonal element

$$\frac{f(z)z^{-\kappa}}{g(z)^r + \varepsilon} E_{hh}, \qquad \varepsilon > 0.$$

The product is $fg^r/(g^r + \varepsilon)E_{hj}$ and converges uniformly to fE_{hj}. Indeed near one $|f|$ is small, and on the complement of that neighborhood g^r has a positive minimum. These two bounds prove uniform convergence without assuming division by a function vanishing at the fiber. Thus the generated algebra contains the whole kernel ideal. Its quotient contains the entire paired diagonal algebra, so it is $\mathcal{D}_H$.

For injectivity let a finite linear combination of path images vanish. At one, all positive paths vanish, forcing the coefficients of the original vertex idempotents to vanish. Fix a state-matrix entry

in a branch. Its remaining paths have lengths $r + m\ell$ and images

$$g^r z^\kappa \sum_m a_m (g^\ell z)^m E_{hj},$$

with a possible initial zero coefficient if $r = 0$. On $\mathbb{T} \setminus \{1\}$ the prefactor is nonzero, and w_c has infinite range, so the polynomial has infinitely many distinct zeros. All coefficients are zero. Repeating by entries proves faithfulness.

No boundedness improvement repairs full formal extension. Pick any $z_0 \neq 1$, let $\lambda = w_c(z_0)^{-1}$, and consider $1 - \lambda c_h$. Its inverse $\sum_{m \geq 0} \lambda^m c_h^m$ exists in the arrow-adic order. The matrix image at z_0 has a zero eigenvalue on state h, so cannot be invertible. A unital algebra homomorphism would preserve the inverse identity, an immediate contradiction. This proves nonextension without any continuity assumption. It does not rely on the false claim that a norm greater than one alone forces a geometric-series denominator to be singular. Scaling g to have norm at most one still does not allow arbitrary formal series.

The two valid operator completions are structurally different. In the bipartite case $K_0(\mathcal{T}_A) = \mathbb{Z}^{E+1}$ differs from $K_0(\mathcal{D}_H) = \mathbb{Z}^E$; in the nonbipartite case the former has the extra $\mathbb{Z}/2$ torsion. Neither is automatically the norm completion of the arrow-adic topology. No invariant constructed solely from these complex objects distinguishes the rational twists already exhibited; arithmetic descent remains a separate input.

F.5 Invariant basis number and the Leavitt comparison

If a unital k-algebra R_1 has a unital representation on a nonzero finite-dimensional vector space V, an isomorphism $R_1^m \cong R_1^n$ induces an isomorphism $V^m \cong V^n$ by tensoring. Comparing dimensions gives $m = n$. This proves invariant basis number without requiring the representation to be faithful. Every original dessin algebra and completed order has a vertex-simple representation, and each permitted-state Laurent algebra has evaluation onto a nonzero matrix block. The Leavitt algebra of G_α has the same finite-dimensional representation after killing the vacuum sinks and evaluating one boundary cycle. Thus all of these examples have invariant basis number. Leavitt's free-rank phenomena are important elsewhere but cannot be assumed in this class simply from the name LPA.

For the specified finite graph G_α, the usual complex Leavitt path algebra embeds in $C^*(G_\alpha)$ with norm-dense image, using its canonical generators and involution [67, Theorem7.3]. Those generators are the split S_h, T_h, not the unsplit original gentle arrows equipped with ordinary CK1. This is the exact algebraic/analytic twin relationship applicable here.

G String finiteness and the smooth quotient surface

For a finite quiver, finitely controlled implies finitely generated: choose one finitely generated submodule containing each vertex space and sum the finitely many choices. Conversely the module itself works. Stabilization at each of finitely many vertices likewise gives ordinary artinianity. Crawley–Boevey's polynomial theorem therefore applies directly to our finitely generated and artinian polynomial modules, with its primitive injective bands in the latter case. It does not assert a classification of every possible algebraic module.

The formal branch $M = k[[x]]$, $y = 0$, gives a precise counterexample to that stronger assertion. An A-endomorphism F preserves each $x^n M$. For the polynomial truncation f_n of f, one has $F(f) - f_n F(1) \in x^n M$. Separatedness and all n imply $F(f) = fF(1)$. Hence $\mathrm{End}_A(M) = k[[x]]$ is local, so M is indecomposable. If an ordinary word module has $y = 0$, its word cannot contain y or y^{-1}, since such a letter would create a nonzero basis map. A reduced word using only $x^{\pm 1}$

cannot change direction. Its nonperiodic choices are finite strings, the free ray $k[x]$, and the opposite locally nilpotent ray. The first and last fail injectivity of x, and the free ray fails invertibility of $1-x$. A two-sided pure-x word is periodic, and in every associated band x acts invertibly. On M, x is injective but not onto. None of these possibilities equals M. This includes arbitrary nonzero indecomposable Laurent coefficient modules, not just finite bands.

For a general nonzero completed lattice, finite generation over A would imply finite generation over $k[u]$. Every nonzero polynomial in u is a power of u times a unit of $k[[u]]$, so it acts injectively. The lattice would be a nonzero finite free $k[u]$-module. Multiplication by $1-u$ on such a free module has a nonzero cokernel, but the same map is invertible over $k[[u]]$. This proves the contradiction in the core. The strictly descending u-adic chain separately excludes artinianity. These are actual failed hypotheses, not merely unverified applicability.

Lekili–Polishchuk's precise convention takes $D(A)$ to mean perfect dg modules. Their locally gentle definition allows infinite dimension, and their smoothness criterion says no forbidden cycle; properness says no permitted cycle. Both fail for every nonempty closed dessin here. Their Theorem3.2.2 therefore does not apply to the full polynomial dessin algebra, even though it applies to suitable infinite-dimensional quotients. The OPS finite-dimensional geometric-intersection theorem has its own grading, local-system, boundary and exceptional coincident-curve conditions. An unrestricted statement that every gentle/string derived object is an ungraded geodesic crossing is not justified by either source.

For the interval quotient $B = A/(y) = k[x]$, global dimension one follows from its PID structure: every submodule of a finite free module is free, giving finite projective resolutions of all finitely generated modules. Thus $\mathrm{Perf}(B) = D^b(\mathrm{mod}_{\mathrm{fg}}\, B)$ in grading zero. Its permitted loop gives one unmarked boundary component and its length-one forbidden thread x gives one stop on the other boundary, the annulus of the smooth partially wrapped model. This construction uses the infinite polynomial quotient, not the completed $k[[x]]$ category.

Finally the quotient does not give a derived fully faithful embedding by ordinary inflation. Resolve B over A by the alternating maps $y, x, y, x, \ldots$. Hom into B gives $0, x, 0, x, \ldots$, so $\mathrm{Ext}^{2r}_A(B,B) = B/xB = k$ for every $r \geq 1$, whereas all positive self-Ext over B vanish. Consequently its inflation is not derived fully faithful, and the perfect B-module B becomes an infinite-projective-dimension A-module. A surface functor transferring these categories must account for that difference explicitly.

H Arithmetic character sums and the cellular determinant

H.1 The uniform calculation and its exceptional cases

In the finite-field calculation write $k = \mathbb{F}_q$; the final order-insertion paragraphs specify a separate characteristic-zero coefficient field. The curve is nonsingular because its discriminant is $-432d^2 \neq 0$, and has one point at infinity. Thus counting $1 + \eta(x^3 - d)$ in each affine fiber proves the trace formula in Theorem 6.2 directly. The multiplicative group of k is cyclic. When $3 \mid q-1$, a nonzero cube has three roots and a noncube none, detected by $1 + \rho(t) + \bar{\rho}(t)$. At zero this expression is one, using the *constant* one. This convention differs from mathlib's zero-extended trivial multiplicative character, whose value there is zero. The proof and formal components never replace the constant by that character.

The substitution is valid including $u = 0, 1$ because the characters are zero extended:

$$\sum_t \rho(t)\eta(t-d) = \sum_u \rho(du)\eta(du-d) = \rho(d)\eta(-d)\sum_u \rho(u)\eta(1-u).$$

Multiplying by the initial minus sign gives exactly $\pi_d + \bar{\pi}_d$. Changing ρ to ρ^{-1} conjugates both

j and λ_d, so does not change the unordered roots or the integer trace. Characteristics two and three and $d = 0$ are excluded throughout. For a residue characteristic $p \equiv 2 \pmod 3$, odd extension degrees have trace zero by cube bijectivity, whereas even degrees lie in the split case; no claim of zero trace for all extensions of that characteristic is made.

For completeness, choose a nonzero $\mathbb{F}_p$-linear functional on k and compose it with $t \mapsto e^{2\pi i t/p}$ to obtain ψ. For a nontrivial multiplicative character χ,

$$\begin{aligned} |G(\chi)|^2 &= \sum_{u \neq 0} \chi(u) \sum_{y \neq 0} \psi((u-1)y) \\ &= (q-1) - \sum_{u \neq 0,1} \chi(u) = q. \end{aligned}$$

Both orthogonality identities follow by translating a finite character sum by an element on which its character is not one. The $z = 0$ terms in $G(\chi)G(\theta)$ sum to $\theta(-1)\sum_x (\chi\theta)(x)$, zero if the product is nontrivial. For $z \neq 0$ set $x = zu$, $y = z(1-u)$ to obtain $J(\chi,\theta)G(\chi\theta)$. Coprime orders three and two ensure our product character is nontrivial. Taking norms proves $|J|^2 = q$, rather than presuming a desired positivity. This is the classical calculation reviewed directly in [68]; no claim of novelty is made.

At seven the five nonzero summands are $-\omega^2, -\omega, \omega, -\omega^2, 1$, so $J = 3 + 2\omega$. Multiplication by $\lambda_2 = \omega^2$ gives $-1 - 3\omega$. Their matrices are

$$\begin{pmatrix} 3 & -2 \\ 2 & 1 \end{pmatrix}, \qquad \begin{pmatrix} -1 & 3 \\ -3 & 2 \end{pmatrix}.$$

Both have determinant seven and the same positive Gram form; their traces are four and one. The arithmetic distinction between the rational models now comes from explicit finite-field characters, while their complex dessin is unchanged. Exact prime-field diagnostics test every nonzero twist for the listed nineteen primes; the theorem itself is uniform in prime powers and rests on the proof, not that finite list.

H.2 Deck descent and the actual cellular basis

The CM cellular construction in this and the next subsection assumes $q \equiv 1 \pmod 3$. The cover $\beta = x^3/d$ has deck transformations $(x,y) \mapsto (\zeta_3 x, y)$ and $(x,y) \mapsto (x,-y)$, generating $C_3 \times C_2 = C_6$. Label their actions on darts by $\sigma = (012)(345)$ and $\alpha = (03)(14)(25)$; they commute. Their product is $(0\,4\,2\,3\,1\,5)$. They preserve the edge pairing and both successor permutations. Consequently they define algebra automorphisms, preserve J and its completion and fix the common u. The projective-resolution maps are compatible since $\theta_g(a_h x) = a_{gh}\theta_g(x)$. Twisting module actions gives an exact functor and the corresponding relabelled resolution, without an assumption of arithmetic Frobenius.

Over an algebraic closure choose $a^3 = d$, $b^2 = d$ and $\varphi(x,y) = (ax, by)$ from E_1 to E_d. For the q-power Frobenius morphisms,

$$\varphi^{-1} \circ F_d \circ \varphi = h_d \circ F_1, \qquad h_d(x,y) = (a^{q-1}x, b^{q-1}y).$$

When $q \equiv 1 \pmod 6$, the first coordinate is the cubic residue $d^{(q-1)/3}$ and the second $\eta(d)$. Choose the embedding of cube roots defining ρ consistently. On the deck eigenspace labelled by dx/y, the *deck* character is their ratio, $\rho(d)/\eta(d) = \lambda_d$. This labels the twist, not a nonzero Frobenius action on differentials in characteristic p: that pullback is zero. An inverse convention for arithmetic Galois

Frobenius changes the action convention; our displayed formula uses the geometric curve morphism explicitly. The character-sum polynomial fixes its trace without an unstated convention.

Orient the three geometric edges as in the core. Each occurs twice with opposite orientation in the single face, so $\partial_2 = 0$; ∂_1 has columns $(-1, 1)^t$. The homology basis $b_0 = e_0 - e_1$, $b_1 = e_1 - e_2$ has Gram matrix G inherited from the orthonormal cellular edges. Rotation sends $b_0 \mapsto b_1$, $b_1 \mapsto -b_0 - b_1$, exactly C. The involution swaps vertices and reverses each edge, hence acts as $-I$ on this plane. Thus multiplication by $A + B\omega$ is literally $AI + BC$, not merely an abstract matrix with the correct trace. Expanding its determinant gives the norm $A^2 - AB + B^2$, and multiplication of this norm proves $M^t G M = qG$. The displayed positive quadratic decomposition proves the metric independently of unknown zeros.

H.3 An explicit chain lift and determinant cancellation

Let P be the three-cycle permutation on C_1. Write $\pi_d = A + B\omega$ and $\gamma = A + B$. Define

$$F_1 = AI + BP, \qquad F_0 = \frac{1}{2}\begin{pmatrix} 1+\gamma & 1-\gamma \\ 1-\gamma & 1+\gamma \end{pmatrix}, \qquad F_2 = q.$$

These are rational cellular maps. The sum-zero plane of F_1 has restriction M_d, while the edge-sum line has eigenvalue γ. On C_0 the vertex-sum line has eigenvalue one and the boundary line eigenvalue γ. It follows directly that $F_0\partial_1 = \partial_1 F_1$; the other chain condition is zero. The induced maps on H_0, H_1, H_2 have characteristic factors $1 - T$, $P_d(T)$, $1 - qT$. In particular

$$\det(I - TF_0) = (1 - T)(1 - \gamma T), \qquad \det(I - TF_1) = (1 - \gamma T)P_d(T).$$

The contractible boundary factor cancels, giving

$$\frac{\det(I - TF_1)}{\det(I - TF_0)\det(I - TF_2)} = \frac{P_d(T)}{(1 - T)(1 - qT)}.$$

This equality holds as a formal rational identity and by cancellation at apparent intermediate singularities. The alternating determinant with exponent $(-1)^i$ is its inverse; signs are not changed silently. The standard elliptic trace and determinant theorem identifies the right side with the finite-field curve zeta function [22, 20]. The elementary proof above supplies its actual trace and norm here; it does not independently reprove the general cohomological trace formalism.

This lift also has the required descent character on H_1. Write the base Jacobi value as $j = A_1 + B_1\omega$. Then h_d acts there by $\eta(d)C^r$ when $\rho(d) = \omega^r$, so twisting $A_1 I + B_1 C$ multiplies its eigenvalue j by λ_d, giving $M_d = A_d I + B_d C$. Reduction modulo $C^2 + C + I = 0$ gives the coefficients of π_d. If the corresponding full three-edge polynomials differ by $c(I + P + P^2)$, the coefficients are $c = 0, \eta(d)B_1, \eta(d)A_1$ for $r = 0, 1, 2$ respectively; they have the same homology action and differ only on the contractible boundary pair. An explicit homotopy sends $v_- - v_+$ to $c(e_0 + e_1 + e_2)$ and the vertex sum to zero. Its $\partial H + H\partial$ is the edge difference and the boundary scalar difference $3c$. Thus this normalization is compatible up to rational chain homotopy. It is not asserted to arise from a continuous Frobenius map of the complex Riemann surface, or to give a canonical integral Tate-lattice comparison.

H.4 A decisive restriction at inert primes

For $q \equiv 2 \pmod 3$, Frobenius sends ζ_3 to ζ_3^{-1}. Thus its cohomological relation with the cubic deck action requires $MC = C^2 M$ (the pullback convention yields the equivalent reversed relation). On a

real two-dimensional carrier with our fixed C, comparing entries forces

$$M = \begin{pmatrix} a & b \\ a+b & -a \end{pmatrix}, \qquad \det M = -a^2 - ab - b^2 = -\tfrac{1}{2}\big(a^2 + b^2 + (a+b)^2\big) \le 0.$$

The genuine elliptic Frobenius determinant is $q > 0$, so such a real carrier is impossible. This is an elementary instance of the real-coefficient cohomology issue discussed by Deninger [62]. It does not rule out a complex realization, a larger real space, a different coefficient theory or an infinite-dimensional height operator. In particular it is not a refutation of an unrestricted Hilbert–Pólya approach.

There is a local complex replacement, with its limitations explicit. The conjugation matrix

$$K = \begin{pmatrix} 1 & -1 \\ 0 & -1 \end{pmatrix} \quad \text{satisfies} \quad K^2 = I,\ KC = C^2K,\ K^tGK = G.$$

Set $F = i\sqrt{q}\,K$ on the complexification. Then $\operatorname{tr} F = 0$, $\det F = q$, $F^2 = -qI$ and $F^*GF = qG$, giving the correct trace-zero local factor. This is a two-dimensional complex, hence four-dimensional real, local model. This is a one-q model; independently choosing $i\sqrt{q}K$ for each extension field does not give Frobenius-power compatibility. Such a family would need to start from one F_p and prove the comparison for its powers. The inserted $i\sqrt{q}$ is not a canonical rational or integral Tate comparison or a construction of a global positive determinant. The fixed real two-dimensional model has genuinely failed; its surviving split-prime construction and this complex alternative must not be conflated.

H.5 The remaining order and global spectral obligations

The deck automorphism has finite order, so over characteristic zero its eigenvalues are roots of unity. Its permutation action alone cannot supply a weight-one norm $\sqrt{q} > 1$. The independently computed Jacobi coefficient is essential. The above construction closes the specified local cellular benchmark, not a reconstruction of arithmetic from bare complex dessin data.

For these order-insertion tests take $\Lambda = \Lambda_{\mathbb{Q}}$ over $R = \mathbb{Q}[[u]]$, with rational residue simples and $V_d = \mathbb{Q}^2$. This is distinct from reducing the matrix in characteristic p, where its determinant p becomes zero. There are two concrete failures of a naive order lift. First, a residue simple S_e is annihilated by all arrows; $S_e \otimes V_d$ with independent endomorphism $I \otimes M_d$ has the desired underlying finite-$\mathbb{Q}$ determinant. But its projective dimension is infinite by the nonterminating forbidden resolution, and σ permutes all three vertex labels, so choosing S_e does not preserve full deck symmetry. This is not a perfect-order determinant realization.

Second, $\Lambda \otimes V_d$ is free of rank two over Λ, but the underlying R-rank of Λ is eighteen. In an R-basis, the Λ-*linear* $I \otimes M_d$ is eighteen copies of M_d, hence its determinant is P_d^{18}. On $e\Lambda$ the rank is six: each of its two hereditary branches contributes a row of length three, with finite conductor gluing changing no rank. This gives P_d^6. Restoring all three residue vertices similarly gives P_d^3 for this untwisted tensor action. A genuine deck-semilinear map permutes factors as well and need not have these powers; it requires its own comparison. No justified determinant-line root or multiplicity-removing functor is supplied here.

Finally fix a nonzero rational d when forming its global Euler product from good reductions. Local purity cannot be multiplied into a global zero-location proof. Since each reciprocal root has modulus $\sqrt{p}$, the product of good elliptic factors is absolutely convergent on $\operatorname{Re} s > 3/2$. The local zeros at $\operatorname{Re} s = 1/2$ are outside that initial product argument. The finite-field curve zeta's center is $1/2$; the global elliptic L-function's unshifted center is one, or $1/2$ only after a half-unit shift.

Neither is an identification with the primitive Dirichlet function targeted earlier. Ramified factors, archimedean normalization, adelic descent and the global determinant-preserving positive spectral limit remain separate obligations.

I The theta order interface beyond tensor insertion

We prove the multiplicity theorem and construct its actual functorial escape, retaining the completed infinite order as the primary algebra. The finite algebra used below is its specified special fiber, not a replacement primary example.

I.1 Normal forms and the special fiber

Index the three quiver vertices by $i \in \mathbb{Z}/3$. Let $a_i^{\pm} : i \to i-1$ be the two permitted branches. Only paths of constant sign survive. Write $p_{i,r}^{\pm}$ for their length-r paths and $c_i^{\pm} = p_{i,3}^{\pm}$. The split order model gives

$$ue_i = c_i^+ + c_i^-, \qquad p_{i,r+3}^{\pm} = up_{i,r}^{\pm} \quad (r \geq 1).$$

The second identity follows because multiplying by the opposite full cycle switches branch and vanishes. An R-basis of $e_i\Lambda$ is

$$e_i,\ a_i^+, a_i^-,\ p_{i,2}^+, p_{i,2}^-,\ c_i^+.$$

These span by the two identities and completion. They are independent in the two hereditary branch orders: off-diagonal entries distinguish the four shorter paths; the remaining combination $xe_i + yc_i^+$ has paired diagonal entries $(x + uy, x)$, forcing $x = y = 0$. This is an actual power-series basis, not just a generic rank calculation.

In $B = \Lambda/u\Lambda$, the same eighteen basis vectors remain. A product of nonempty paths is zero if branches switch or the total length exceeds three; the length-three minus path is the negative of the plus path. A length-four path is u times a length-one path, hence zero. The radical $\mathfrak{r}$ is the positive-length ideal, with $\mathfrak{r}^4 = 0$ and $B/\mathfrak{r} = \mathbb{Q}^3$. Each e_iB has radical-layer dimensions $(1, 2, 2, 1)$ and has two composition factors at each vertex, so the Cartan matrix is

$$C_B = \begin{pmatrix} 2 & 2 & 2 \\ 2 & 2 & 2 \\ 2 & 2 & 2 \end{pmatrix}.$$

The Cartan matrix alone does not establish the endomorphism formula; the path filtration supplies that step.

I.2 All projectives and all linear endomorphisms

Let J be the completed arrow ideal. Every $1 - x$, $x \in J$, has its convergent formal inverse; the same applies to $1 - yx$ for arbitrary $y \in \Lambda$. Thus J is contained in the Jacobson radical, and $\Lambda/J = \mathbb{Q}^3$ implies equality. If P is finitely generated projective, choose a basis of each multiplicity space in P/PJ and lift its vectors into Pe_i. This gives

$$Q = \bigoplus_i (e_i\Lambda)^{n_i} \longrightarrow P$$

inducing an isomorphism on tops. Nakayama makes this surjective. Its split kernel N is finitely generated and satisfies $N/NJ = 0$, hence vanishes by Nakayama. This proves the decomposition for every such projective without an unproved freeness assumption over Λ.

An entry $e_i\Lambda \to e_j\Lambda$ is left multiplication by an element of $e_j\Lambda e_i$. Its degree-zero part is possible only when $i = j$ and is a scalar multiple of e_i. For f on P, write its top blocks A_i and set $A = \operatorname{diag}_i A_i$. The remainder has positive path length. On P/uP it strictly raises the path filtration. At each of the two length-one and two length-two slots, and the length-zero and length-three slots, the degree-preserving map is exactly A. In an increasing-degree basis the matrix is lower block triangular. For the formal upper-block convention order the six slots instead as

$$3+,\ 2+,\ 2-,\ 1+,\ 1-,\ 0.$$

There are six diagonal copies of A, allowing arbitrary positive-length off-diagonal entries. Hence

$$\det_{\mathbb{Q}}(1 - T\overline{f}) = \det_{\mathbb{Q}}(1 - TA)^6.$$

Specializing the finite R-matrix determinant coefficientwise gives Theorem 6.3. No simultaneous diagonalizability, tensor form of f, arithmetic purity or deck equivariance is required.

For a bounded complex $P^\bullet$ with a linear chain map $f^\bullet$, fix either parity convention and write

$$D_R(P^\bullet, f^\bullet; T) = \prod_j \det_R(1 - Tf^j)^{(-1)^j}.$$

The inverse convention gives the reciprocal and the same obstruction. Each factor has constant coefficient one, so its reduction is nonzero. Specialization of this displayed rational expression is defined and equals

$$\left[\prod_{j,i} \det_{\mathbb{Q}}(1 - TA_i^j)^{(-1)^j}\right]^6.$$

Cancellations cannot change the divisibility of divisor multiplicities. In particular adding contractible projective pairs cannot repair it. Here a perfect endomorphism means a bounded-projective chain representative; no statement about an arbitrary infinite resolution or nonperfect lattice is inferred.

Generically put $N = \sum_i n_i$. As a right module over $M_3(K) \times M_3(K)$, P_K is the sum of N copies of each branch's three-dimensional row simple. Any commuting map has matrices $F_+, F_- \in M_N(K)$ on their multiplicity spaces, so

$$\det_K(1 - Tf_K) = \big(\det(1 - TF_+)\det(1 - TF_-)\big)^3.$$

For example, left multiplication by c_i^+ on $e_i\Lambda$ has ordinary characteristic determinant $(1 - uT)^3$, so a sixth power is forced only at the special fiber. This also shows why generic reduced determinants and ordinary determinants must not be conflated. We do not need an integral choice of either generic matrix to derive the cube obstruction.

For the local numerator, suppose a sixth power $A(T)^6/B(T)^6$ equals a nonzero quadratic $P(T)$ with $B \neq 0$. Polynomial degrees would give $6\deg A = 2 + 6\deg B$, impossible. This includes repeated roots. For the complete curve zeta, $q > 1$ and $|a_q| \leq 2\sqrt{q}$ imply $P(1) > 0$ and $P(1/q) = P(1)/q > 0$. Its two denominator roots are distinct uncancelled simple poles, again impossible for a sixth power. These arguments do not assume global RH or Artin holomorphy.

I.3 An actual integral determinant functor of multiplicity one

Project to one hereditary cycle order $B_+ \subset M_3(R)$. Its standard column $W_+ = R^3$ is a left Λ-module with $e_iW_+ = R\epsilon_i$. More explicitly a_i^+ acts by $u^{w_i}E_{i,i-1}$, with $(w_0, w_1, w_2) = (1, 0, 0)$

for one cut, while every a_i^- acts by zero. These matrices respect travel-order multiplication as written: $E_{ij}E_{jk} = E_{ik}$. Each plus full cycle is uE_{ii}, so the common central u acts by uI. Nonempty paths tend to zero u-adically, giving the action of the *full completed order*, not just its polynomial subalgebra.

This (Λ, R)-bimodule defines $- \otimes^{\mathbf{L}}_{\Lambda} W_+$. On a bounded complex of finite projectives it is computed termwise, and its terms are finite free over R. An acyclic bounded projective complex is split contractible, and tensor preserves that contraction. Thus it descends to a genuine triangulated functor $\Phi_+ : \mathrm{Perf}(\Lambda) \to \mathrm{Perf}(R)$. No assertion of exactness on all un-derived modules or flatness of W_+ is needed. In fact W_+ has infinite projective dimension as a left Λ-module: Λ is noetherian, being finite over the central noetherian ring R, and W_+ is finitely generated. A finite projective resolution could thus be chosen with finite terms. Every left projective term has R-rank divisible by six by the same top argument, whereas W_+ has rank three, contradicting the alternating rank identity. This does not prevent tensoring bounded right-projective complexes.

The isomorphism $e_i\Lambda \otimes_\Lambda W_+ \to e_iW_+$ sends $e_i\lambda \otimes w \mapsto e_i\lambda w$. Its inverse is $e_iw \mapsto e_i \otimes e_iw$, proving rank one directly. Consequently $e_i\Lambda \otimes_{\mathbb{Q}} V$, with action $1 \otimes M$, maps to $R \otimes_{\mathbb{Q}} V$ with action M. Its characteristic determinant is $\det(1 - TM)$ *once*. On the regular input, the three vertex lines survive and the factor appears three times. A full symmetric vertex orbit therefore is not silently replaced by a chosen vertex.

Different cycle cuts give different integral column lattices. Their generic modules are isomorphic, and branch endomorphism characteristic determinants agree by generic conjugation; equality over K implies equality over R. Nevertheless an equivariant lattice is stronger data. Branch selection also changes the functor: for the unit $f = e_i + c_i^+$ on $e_i\Lambda$, the plus determinant is $1 + u$ and its characteristic factor is $1 - (1 + u)T$. Under α, c_i^+ becomes c_i^-, whose plus action is zero; the determinant becomes one and the characteristic factor $1 - T$. Since these scalar automorphisms differ, no natural isomorphism of determinant lines can make this chosen branch functor α-invariant for all endomorphisms. Pairing both branch functors makes this branch-swap data available but doubles the local factor on the basic-projective tensor input.

I.4 The exact rotation obstruction and its allowed escape

Consider any left Λ-lattice supported on the plus branch, free of rank one at every e_i, and with central u acting as multiplication by u. Write its plus arrow scalars as $a_0, a_1, a_2 \in R$. Their product is u, so they are nonzero and their nonnegative valuations sum to one. A strict R-linear lift of σ that intertwines the order rotation permutes the three vertex lines by units. Its arrow-intertwining identities make the three arrow valuations equal. Thus $3m = 1$ for an integer $m \geq 0$, impossible. The argument already excludes an invertible intertwiner; imposing that its cube be one cannot help.

There is an elementary coefficient proof of the same obstruction. Intertwining by units makes vanishing of the three arrow constant coefficients equivalent. Their product has constant coefficient zero, so all three vanish. Factoring u from two of them would make $a_0a_1a_2$ divisible by u^2, contradicting that it equals u. This is the literal power-series component checked in Lean.

After the explicit ramified change $R' = \mathbb{Q}[[v]]$, $u = v^3$, *choose a new column lattice* with all three plus arrow maps $vE_{i,i-1}$. Cyclic permutation of the basis is an order-three intertwiner fixing v. The full-cycle relation is now $v^3 = u$ as required. Merely base-changing the old cut lattice gives arrow valuations $(3, 0, 0)$ and does not produce that linearization. This construction changes the base and lattice and does not establish an unramified descent comparison. It also still needs both branches to address α.

The rotation obstruction is about actual central-fixed rank-one branch lattices, not every possible determinant-level natural transformation. The branch characteristic determinants may be

rotation-invariant through their generic reduced interpretation even without a lattice lift. Neither this theorem nor the ordinary-determinant obstruction excludes semilinear correspondences, other determinant categories or infinite analytic models.

The next arithmetic task is therefore narrower and more concrete: choose a branch functor, a ramified lattice or a different equivariant replacement only after proving the required descent comparison, and construct its prime action from arithmetic data rather than feed in the already known M_d. Bad primes, integral/Tate comparison, local-to-global conductor and gamma factors, and a positive spectral limit for the same completed function remain unresolved.

J Explicit tame descent and the full-deck coefficient test

The preceding ramified column is a precisely specified object of the original infinite theta-order framework. We compute its descent instead of treating a semilinear group action as automatically effective integral descent. All constructions below are over the new *mathematical* coefficient field $k = \mathbb{Q}(\omega)$; they do not replace the original rational model or its source history.

J.1 The action on genuine formal series

Write $R = k[[u]]$, $S = k[[v]]$ with $u = v^3$. Every $f(v) \in S$ has a unique expansion $f_0(u) + vf_1(u) + v^2 f_2(u)$. Hence S is finite free of rank three over R. Let $g(f)(v) = f(\omega v)$. On $W = S^3$ with basis $\epsilon_0, \epsilon_1, \epsilon_2$, put

$$\Theta(f_i\epsilon_i) = \omega^{-i} f_i(\omega v)\epsilon_i, \qquad C\epsilon_i = \epsilon_{i+1} \quad (i \bmod 3).$$

The plus arrow a_i^+ acts by $vE_{i,i-1}$, the minus arrow by zero. The cyclic product is uE_{ii} and arbitrarily long permitted paths tend to zero v-adically, so this is a module over the full completed order. For $i = 0$, the coefficient in the target under Θa_i^+ is ωv and in the source under $a_i^+\Theta$ it is $v\omega^{-2}$; these agree. The same calculation for $i = 1, 2$ proves compatibility with every arrow and therefore the completed algebra. Also $\Theta^3 = 1$.

For all coefficients, not just a truncated diagnostic,

$$[v^n](\Theta w)_i = \omega^{n-i}[v^n]w_i.$$

Thus invariance is equivalent to support in $n \equiv i \pmod 3$, and

$$L = W^\Theta = \bigoplus_{i=0}^{2} Rb_i, \qquad b_i = v^i\epsilon_i.$$

The arrows on this basis have scalar valuations $(1, 0, 0)$, since $a_0^+ b_2 = ub_0$, $a_1^+ b_0 = b_1$ and $a_2^+ b_1 = b_2$. Tame invariants return the old cut column rather than a new strict unramified rotation carrier.

The extension $R \subset S$ is faithfully flat. Its group quotient has a fixed point at $v = 0$, however, and is not a μ_3-torsor there: $S \otimes_R S$ is not the product of the three group graphs at that point. Semilinear μ_3 data are therefore not by themselves the full effective fpqc descent data for a vector bundle on $\operatorname{Spec} R$. The adjunction comparison is explicitly

$$S \otimes_R L \longrightarrow W, \qquad \operatorname{diag}(1, v, v^2),$$

with determinant $v^3 = u$, cokernel $S/(v) \oplus S/(v^2)$ and closed-fiber rank one. This proves its failure to be an isomorphism directly. The quotient stack $[\operatorname{Spec} S/\mu_3]$ retains the stabilizer at zero; its coarse space is $\operatorname{Spec} R$. This local root construction is described in [71]. Alper's vector-bundle criterion [70, Theorem10.3, pp.2386–2387] distinguishes precisely this nontrivial closed-fiber action from effective coarse descent. We use these sources for interpretation; all displayed module comparisons have already been calculated explicitly.

J.2 A projective symmetry survives on the endomorphism order

Directly on the three basis vectors,

$$\Theta C = \omega^{-1} C\Theta = \omega^2 C\Theta.$$

Thus the rotation does not preserve L. The corrected vC does preserve it, since the factor ω from $g(v)$ cancels the commutator. In the invariant basis it has matrix A from the core, with $A^3 = uI$ and $\det A = u$. This is generically invertible and gives an order-three projective rotation after passing to the fraction field; it is not an invertible integral lattice intertwiner.

There is nevertheless a true algebra-level escape. Let $F = (f_{ij})$ be an arbitrary S-linear endomorphism of the *carrier* W, not assumed Λ-linear. Its tame conjugate has entries $\omega^{j-i} g(f_{ij})$. Invariance therefore means

$$f_{ij} \in v^{[i-j]_3} R,$$

where $[r]_3 \in \{0, 1, 2\}$ is the least residue. After replacing ϵ_i by $b_i = v^i \epsilon_i$, the entry becomes

$$v^{[i-j]_3 + j - i} R = \begin{cases} R & i \geq j, \\ uR & i < j. \end{cases}$$

Consequently $\mathrm{End}_S(W)^\Theta$ is exactly the hereditary three-cycle order B_+ in the previously fixed cut convention. Multiplication is preserved by conjugation in the generic matrix algebra; no entrywise projection is being misused as a ring map.

The scalar commutator cancels under conjugation. Hence $\mathrm{Ad}(C)$ commutes with tame conjugation, descends to B_+, and is represented there by $\mathrm{Ad}(A)$ because v is central. Since $A^3 = uI$, this algebra automorphism has order three and preserves the hereditary order despite A^{-1} not lying in its unit group. This explains the surviving rotation of the branch order; it does not construct a rotated integral column or a Frobenius endomorphism. The other branch is identical and the paired-diagonal residue gluing still defines the original Λ_R inside $B_+ \times B_-$. Its label automorphisms remain the already constructed dessin symmetries.

J.3 Determinants, Schur weights and the boundary correction

On $\det_S W = \bigwedge_S^3 W$, both $\det D = 1$ and $\det C = 1$. Thus this line has the scalar action g and genuinely descends to the trivial R-line with trivial rotation. This is a real positive determinant-level result. On the other hand,

$$b_0 \wedge b_1 \wedge b_2 = v^3 (\epsilon_0 \wedge \epsilon_1 \wedge \epsilon_2),$$

so the natural comparison

$$\det_R(W^\Theta) \longrightarrow (\det_S W)^\Theta$$

is multiplication by u, with cokernel $R/(u)$. Both lines are abstractly free of rank one, but their given comparison is not an isomorphism. It records the divisor of the ramified lattice change and cannot be discarded when claiming a determinant-preserving integral comparison. For the *same* operator vC, the scalar on both determinant lines is u; naturality is respected. Comparing $\det C = 1$ with $\det A = u$ without the extra factor v would instead compare different operators.

For an induced nonzero homogeneous tensor/Schur carrier of degree d, the same functorial calculation gives commutator ω^{-d}. The two induced invertible operators commute exactly when $3 \mid d$. This is a group-action statement, not sufficient for coarse vector-bundle descent: $\mathrm{Sym}^3(S^3)$ has closed-fiber inertia multiplicities $(4, 3, 3)$ for the three characters, so its inertia is still nontrivial

even though the commutator vanishes. A mixed tensor $W^{\otimes a} \otimes (W^*)^{\otimes b}$ has net exponent $a - b$; $\mathrm{End}_S(W)$ is the concrete net-zero escape computed above.

These carrier functors are not automatically functors on unital Λ-modules. Applying global $\bigwedge^3$ to the rank-one vertex idempotents kills all three, although their sum is the identity before applying this nonadditive functor. Vertexwise $\bigwedge^3$ of the rank-one column is zero. Vertexwise Sym^d for $d > 0$ does retain the monomial zero relations but changes the full-cycle scalar from u to u^d. It leaves the fixed central-action problem unless that change of base is included. None of the Schur cancellations supplies the missing two-dimensional arithmetic carrier.

J.4 Why the local Jacobi plane does not descend by this repair

On constant vectors Θ acts by D. It sends the sum-zero cycle $\epsilon_0 - \epsilon_1$ to $\epsilon_0 - \omega^2\epsilon_1$, whose coordinate sum is nonzero. It also does not preserve the edge-sum line. Thus this order column is not already a tame descent of the earlier cellular chain complex or its H_1.

For the split local coefficient $\pi = a + b\omega$, the natural rotation polynomial on the balanced carrier is $F = aI + bC$. Since g fixes $a, b \in k$, its commutator is

$$\Theta F - F\Theta = b(\omega^2 - 1)C\Theta.$$

It is nonzero when $b \neq 0$. In particular both prime-seven twists $(a, b) = (3, 2), (-1, -3)$ fail this test. Prime powers with $b = 0$ are not excluded by this particular calculation; that exceptional commutation does not prove a global comparison.

Replacing C by vC gives an invariant carrier map, whose matrix on L is $aI + bA$. Direct determinant expansion gives

$$\det(1 - T(aI + bA)) = (1 - aT)^3 - ub^3T^3.$$

At $u = 0$ this is $(1 - aT)^3$, not $P_d(T)$, which has degree two. The weighted replacement therefore does not recover the established local factor. Neither C nor vC is generally Λ-linear: they intertwine its rotation. Their sums with the identity have not thereby become endomorphisms in the ordinary order-module category. They are legitimate carrier tests, with exactly the failures just exhibited.

J.5 The full orbit is the normalization, and arbitrary kernels still fail

Twist the tame action by the three characters: $\Theta_r = \omega^r\Theta$, $r \in \mathbb{Z}/3$. Its fixed lattice is

$$L_r = \bigoplus_i Rv^{[i-r]_3}\epsilon_i,$$

with the cut at arrow r. Rotation gives an actual R-linear isomorphism $C : L_r \to L_{r+1}$, taking each displayed basis vector to the corresponding next one. Thus the direct sum of all three cuts has a strict rotation, at the cost of vertex rank three. Before invariants the orbit is S-rank nine and underlying R-rank twenty-seven; its invariant lattice is R-rank nine. The original orbit S-lattice still has nontrivial inertia at zero and need not be the base change of these invariants.

There is an explicit integral identification of the invariant orbit. In the standard hereditary order B_+, the j-th column has entries $u^{t_{ij}}R$ with $t_{ij} = 1$ for $i < j$ and zero otherwise. In its natural column basis the arrow coefficient has exponent

$$\delta_{i0} + t_{i-1,j} - t_{ij} = \delta_{ij}.$$

Its cut is therefore exactly j, identifying B_+e_j with L_j. Consequently $\bigoplus_r L_r \simeq B_+$ as left Λ_R-modules. Adding the exchanged minus-branch orbit gives the hereditary normalization $\widetilde{\Lambda} = B_+ \times B_-$,

with full deck action and vertex rank six. On a basic projective with external arithmetic matrix this yields P_d^6.

In fact this normalization replacement restores the *ordinary determinant for every linear endomorphism*. For a finite projective P, the inclusion $P \to P \otimes_\Lambda \widetilde{\Lambda}$ becomes an isomorphism over $K = k((u))$, since $\widetilde{\Lambda}_K = \Lambda_K$. Its extended endomorphism and the original f therefore have the same generic characteristic determinant. Both polynomials have coefficients in R, which injects into K, so they agree already over R. The bounded-complex ratios also agree. This symmetry repair returns the previously refuted ordinary determinant, not merely its external tensor exponent.

For completeness we now prove Theorem 6.4 for arbitrary finite free kernels, without assuming they split integrally into branches or are projective over the order. Let

$$p_\pm = u^{-1} \sum_i c_i^\pm \in \Lambda_K.$$

These are central generic branch idempotents. Put $W_\pm = W_0 \cap p_\pm(W_0 \otimes_R K)$. They are finite free R-lattices spanning those generic summands, stable under Λ_R. Rotation fixes them, branch swap exchanges them. Their vertex ranks are constant on each branch: the arrow maps become invertible over K because a full cycle acts by u. Write these ranks r_+, r_-. No equality $W_0 = W_+ \oplus W_-$ over R has been assumed.

On the plus branch choose integral vertex bases. The three square arrow matrices X_0, X_1, X_2 have product uI_{r_+} around every cycle. Thus

$$\sum_i \mathrm{val}_u(\det X_i) = r_+.$$

The rotation lift is an invertible matrix between each pair of vertex lattices, hence its determinant is an R-unit. Its intertwining identities make all three arrow determinant valuations equal. Therefore $r_+ = 3m_+$; similarly $r_- = 3m_-$. The branch-swap lift gives $r_+ = r_-$, so generic rank additivity gives $\mathrm{rank}_R\, e_i W_0 = 6m$. Only invertible intertwiners were used; imposing the full deck group relations is stronger than necessary.

For the general determinant statement reduce modulo u and filter each vertex by $e_i J^a(W_0/uW_0)$. Since $J^4 \subset u\Lambda$, this is a finite filtration. In $\Phi_{W_0}(P) = \bigoplus_i (e_i W_0)^{n_i}$, positive-length entries of f raise the filtration, while the degree-zero matrix A_i repeats on its i-th graded pieces. Their total dimension is $6m$. Multiplying the graded characteristic determinants proves the asserted power for *every* f. Flatness of the kernel as a left Λ-module was not used; bounded right projectives compute the derived functor and give finite free R-outputs. Denominator constant terms one permit specialization of every bounded alternating ratio. The regular left kernel Λ_R attains $m = 1$, making the rank bound sharp.

The remaining arithmetic problem therefore requires a different hypothesis: a genuinely twisted determinant target, a specified non-linear/semilinear correspondence with its own comparison, or an infinite analytic realization. One cannot continue searching inside this closed finite-free, fixed-center, full-deck kernel class for a missing small estimate. The local purity and descent computations supply no global positive spectral limit. Moreover $k/\mathbb{Q}$ has degree two: ordinary restriction of scalars sends a rational local factor P_d to P_d^2. A relative k-determinant must not silently be identified with the former rational determinant. Finite-field curve zeta, global elliptic L-functions and Dirichlet GRH keep their distinct scopes and centers.

K Literal formal and computational coverage

`DessinNodeOrder.lean` defines the fiber-product subring of two genuine power-series rings. Its twenty printed declarations establish diagonal injectivity, nonzero branches, branch zero products, factorizations and annihilator identities, regular u, augmentation surjectivity and exactness, bijective power-series coordinates, square-zero alternating maps, three kernel-equals-range statements and projectivity of the actual regular/product terms. The coordinate statement is a bijection; the file does not yet package it as an R-linear equivalence or declare the general order freeness theorem.

`DessinInvariantComponents.lean` has ten printed declarations: three CK1-collapse components, determinant change of bases with an actual inverse, symmetric-square zero, identity, composition and relation preservation, the degree-two Schur character identity, and the nodal central-parameter obstruction. Its symmetric-square matrix is

$$\operatorname{Sym}^2\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} a^2 & ab & b^2 \\ 2ac & ad+bc & 2bd \\ c^2 & cd & d^2 \end{pmatrix}.$$

This is the ordinary monomial basis convention; composition is proved universally over any commutative ring. The full project suite prints types and foundational dependencies; no theorem about general LR decompositions, abstract derived equivalences, C* K-theory or GRH is inferred from these component proofs.

The exact Python checks use fixed rational matrices and exhaustive finite sets, with no random seeds. They construct the displayed permutations and paths, evaluate rational Cartan identities, check formal Euler products through a stated degree, verify unrestricted-output Fock basis identities on bounded input sets, compute incidence Smith data, construct all nine string representations and signed maps, and test Young/LR/Cauchy examples and the seven-element-field counts. Those diagnostics are reproducible finite corroboration. The universal infinite claims rest on the proofs or explicitly cited theorems, and the arithmetic comparison remains open.

The ten additional `DessinPullbackComponents.lean` statements use literal two-state matrix units and a subring of a product of arbitrary rings, with equal images under two ring homomorphisms. The section hypothesis of its projection-surjectivity theorem is explicit. It does not construct a topological C*-pullback or assert that a commutative diagram is automatically a pullback. The no-residue theorem is universal algebra, with a nontrivial commutative target; it is not an assumed arithmetic positivity premise.

`ThetaJacobi.lean` adds sixteen declarations, with actual finite fields and MulChar rather than a hypothetical arithmetic kernel. Its norm theorem invokes mathlib's proved Jacobi inverse-product identity and derives characteristic inequality from finiteness; all three character nontrivialities are explicit. Other declarations verify the rescaling substitution, CM multiplication/polynomial/Gram positivity components, exact prime-seven matrix polynomials, and actual 2-by-3 boundary/3-by-2 cellular inclusion, homology restriction, three-edge determinant and rational vertex chain-map equations. It does not formalize elliptic point counting or an order-to-Frobenius functor.

The final three `ThetaJacobi.lean` declarations verify the actual conjugation matrix identities and the real two-dimensional inert-prime determinant obstruction. They do not assume any unknown zero location or exclude complex/enlarged spectral realizations.

`ThetaProjectiveMultiplicity.lean` adds five declarations. Arbitrary upper-block matrices with six identical diagonal blocks have the sixth-power determinant and characteristic-factor formulas. A literal power-series constant-coefficient map specializes matrix determinants, for each scalar parameter; the coefficientwise polynomial/order interface remains prose. A nonzero quadratic cannot be a rational-function sixth power, proved by polynomial degrees. Three genuine power

series with product X cannot have the cyclically equivalent residue-vanishing conditions forced by invertible rank-one arrow intertwiners. The derivation of the path blocks from the full theta order, projective decomposition, derived branch functor and deck intertwiners is not fully formalized. The complete analytic suite now has208printed declarations; no hidden arithmetic positivity assumption occurs.

`ThetaTameDescent.lean` adds twelve printed component declarations (and one auxiliary polynomial lemma). They use genuine formal-power-series rescaling and its coefficient operator, proving the coefficient action, invariant-coefficient vanishing and the tame operator's cube. Actual three-state matrices satisfy the clock/rotation commutator, comparison determinant and closed-fiber equations, weighted-cycle cube and full characteristic polynomial, noncommutation of a nonconstant rotation polynomial, trivial whole-column determinant actions, and the same-operator comparison for vC. The classification of all fixed series by residues, full order action and hereditary invariant algebra, stack interpretation, valuation/rank theorem, general coefficient-functor filtration and GRH are not thereby fully formalized. No unknown positivity premise is assumed.

L Explicit covering calculations and derived transport

Here are the linear details behind Proposition 7.2. Write $x_i = F_{ii}$ for diagonal entries of an endomorphism F preserving the three vertex spaces. Solving $FX = XF$, $FY = YF$ leaves a common diagonal c and six free entries. More explicitly its only off-diagonal entries are

$$F_{02} = F_{13} = t_0, \quad F_{15} = t_1, \quad F_{35} = F_{46} = t_2, \quad F_{48} = t_3, \quad F_{68} = F_{70} = t_4, \quad F_{72} = t_5.$$

All others vanish. This can be checked column by column using the displayed partial permutation matrices X, Y: their nonzero columns specify all constraints, and the zero columns force the remaining entries to zero. If $r_0, \dots, r_5$ are the six radical generators in the displayed order, the only nonzero products are

$$r_0 r_2 = r_1, \qquad r_2 r_4 = r_3, \qquad r_4 r_0 = r_5.$$

Every product of three radical generators vanishes. Thus the listed space is a local algebra, not merely a seven-dimensional vector-space count.

For clarity, the Euler equations used in the obstruction are

$$\begin{aligned} &d_1 - d_0 = 1, && d_3 - d_2 = 1, && d_4 - d_3 = 1, \\ &d_6 - d_5 = 1, && d_7 - d_6 = 1, && d_0 - d_8 = 1, \\ &d_1 - d_2 = 1, && d_4 - d_5 = 1, && d_7 - d_8 = 1, \end{aligned}$$

where $d_i = D_{ii}$. Subtract the last row from the first two rows after summing: the left side is zero and the right side is three. This is an identity in every characteristic-zero coefficient field. In characteristic three this particular obstruction disappears; no conclusion there is asserted. The closed walk

$$a_0 b_0^{-1} a_0 \, a_1 b_1^{-1} a_1 \, a_2 b_2^{-1} a_2$$

explains the computation: six forward steps and three backward steps have net lift displacement three. Arbitrary changes of basis cannot remove the Euler contradiction. This is an algebraic lift obstruction, not an assertion about the topological fundamental group of the compact torus.

The Hom formula also has a constructive perfect-complex version. Each cover representable is finite support and pushes to its corresponding vertex projective. For a bounded complex $P^\bullet$ of finite sums of such representables and a bounded finite-support complex $N^\bullet$, decompose the entire

Hom complex by deck displacement. The differential preserves that displacement, and only finitely many displacements occur across the finitely many degrees. Taking cohomology gives

$$\operatorname{Hom}_{D(B_L)}(FP^\bullet, FN^\bullet[q]) \simeq \bigoplus_g \operatorname{Hom}_{D(\widetilde{B}_L)}(P^\bullet, T_g N^\bullet[q]).$$

Bounded projective complexes compute these derived Hom spaces, which justifies the equality rather than assuming every module perfect. This elementary proof does not assert an equivalence of entire bounded derived categories or cover objects of infinite projective dimension. The broader precovering statement has its exact hypotheses in [74, Theorem 4.28].

Finally the branch involution $\sigma(a_i) = b_i$, $\sigma(b_i) = a_i$ preserves B_L. In characteristic different from two one can study $B_L \# k[\mathbb{Z}/2]$ through equivariant representations. For the nine-dimensional module above, $\operatorname{rank} X = 6$ and $\operatorname{rank} Y = 3$, so $M \not\simeq \sigma M$. The sum $M \oplus \sigma M$ has canonical swap equivariance. This is an explicit orbit construction; it neither chooses a nonexistent fixed-object linearization nor identifies this skew group algebra with the particular GLS algebras in [75].

`CoveringBand.lean` verifies the actual matrices' mixed products, cubic nilpotence and absence of an arbitrary Euler matrix. The general covering, module endomorphism classification, derived functors and the surface/Fukaya equivalence are not fully formalized by these three matrix statements. Exact rational code additionally checks thirty-six Hom pairs, the full radical multiplication and the two Hom-control defects.

M Weighted profiles, exact arithmetic examples and the surviving task

The finite matrix proof also retains an arbitrary row profile. Suppose $|A_{ui}|^2 = w_u$ for every column, set $S = \sum_u w_u$ and $Q = \sum_u w_u^2$. The row diagonals are $N w_u$ and column diagonals are S, so the same identity proves

$$\sum_{i \neq j} |K_{ij}|^2 \geq N^2 Q - N S^2.$$

This is a finite statement. Applying it to an infinite Schwartz tail would require its convergence and every newly encountered prime collision; these are not discarded. If a weighted pair-square attempt includes the annular W, one can instead choose a closed subannulus on which a nonzero $|W|$ is bounded below. Its prime count still has order $D/\log D$. The fixed ν has unit modulus outside its conductor. These facts do not replace the original coefficients in a signed estimate.

Here are exact sextic computations with all nonzero elements of the Eisenstein norm ball, and both prime ideals above each split rational prime in the column annulus. These kernel diagnostics use $\nu = 1$ and exclude only primes above2,3; they do not compute a signed Möbius energy. Values are integer norms in $\mathbb{Z}[\omega]$.

D	H	R	N	$\sum_{i \neq j} \lvert K_{ij}\rvert^2$	$NR(N-R)$
128	4	18	22	6840	1584
1024	9	36	138	651672	506736
4096	81	300	456	60080328	21340800
256	64	240	40	363672	−1920000

The final row has $H = D^{3/4}$ but $N < R$; its negative lower bound is correctly vacuous. The asymptotic theorem follows from the proof, not an extrapolation of these examples. For $p \equiv 1 \pmod 3$ and $w^2 + w + 1 = 0$ in $\mathbf{F}_p$, evaluate $a + b\omega$ as $a + bw$ and take its $(p-1)/6$ power, lifting the six roots via $\zeta_6 = 1 + \omega$. Every zero is tested before a unit value is assigned. For instance $1 + 3\omega$

has norm7 and vanishes at $\omega = 2$ modulo7, so the unimodularity hypothesis cannot be used beyond its stated domain.

`ShortRowPairBarrier.lean` verifies the finite frame identity, row-diagonal bound, weighted inequality and unimodular specialization, with actual complex sums and explicit hypotheses. It does not formalize the ideal-character comparison or prime-ideal theorem. The deterministic integer checker also verifies weighted profiles and256 scalar pair sums independently of its matrix computation. No numerical model or learned coefficient is used.

The next signed route must retain μ, ν, W before Cauchy. An exact starting identity in ideal Dirichlet convolution is

$$\mu = 2a - a * a * \mathbf{1} + b * b * \mathbf{1}, \qquad a = \mu\, \mathbf{1}_{Nn \le U}, \quad b = \mu - a.$$

Indeed $\mu * \mathbf{1} = \delta$ implies $(\mu - a) * (\mu - a) * \mathbf{1} = \mu - 2a + a * a * \mathbf{1}$. On squarefree n, each divisor factorization retains pairwise coprimality; the original row zeros still multiply along it. Estimating these type-I/type-II kernels with their true conductors is a new open arithmetic task. The identity alone supplies no cancellation or GRH proof.

N Exact factorization, retained zeros and the failed Cauchy route

N.1 Proof of the regrouping with original coefficients

The monoid consists of nonzero integral ideals prime to the fixed excluded set S. Every divisor of a good ideal is good, so the identity $\mu * \zeta_0 = \delta$ holds on this same monoid. Expanding convolution in (1) gives finite divisor sums. Multiply the identity by $\mu(n)^2 \nu(n) \chi_n(u) W(Nn/D)$ and sum. The extra squarefree indicator does not change $\mu(n)$, since $\mu(n)^3 = \mu(n)$. On its support $n = abc$ forces each factor to be squarefree and any two to be coprime. Sextic multiplicativity, including zeros, gives $\chi_n(u) = \chi_a(u)\chi_b(u)\chi_c(u)$; a collision with any factor makes both sides zero. The fixed character also satisfies $\nu(n) = \nu(a)\nu(b)\nu(c)$ on these good ideals.

In the small-factor piece group $d = ab$. The remaining variable c is squarefree and coprime to d; $Nd \le U^2$ is necessary, but not sufficient for a nonzero $\beta_U(d)$: its specified truncated factorizations must still exist. For each such factorization on squarefree d, $\mu(a)\mu(b) = \mu(d)$, proving the coefficient formula. In the large-factor piece group $s = bc$. The condition $Nb > U$ implies $Ns > U$, and summing that divisor gives exactly $g_U(s)$. Since $Ns > U \ge 1$, s is not the unit ideal; its complete Möbius divisor sum vanishes, proving the alternative formula for g_U. This regrouping never replaces the two-variable weight by $W(Ns/D)$, never deletes $(a, s) = 1$, and never replaces the resulting coefficients by a favorable arbitrary family.

One can instead keep the unrestricted convolution identity on nonsquarefree columns, but then both individual pieces and their local conductor description change. For $U = 1$ and a good prime p, the column p^2 contributes one to each of $a_U * a_U * \zeta_0$ and $b_U * b_U * \zeta_0$, even though $\mu(p^2) = 0$; these cancel only after the pieces are recombined. Its squared sextic symbol has a cubic local component. Thus dropping total squarefreeness is not a harmless piecewise simplification, although the full unrestricted algebraic identity still holds.

N.2 The two conductor scales are unchanged

For a squarefree $n = as$ with $(a, s) = 1$, every prime of n has one nontrivial sextic local component in the row variable. Hence its primitive finite conductor is the ideal as; $NaNs \asymp D$ on the column support. On multiplying a column by the conjugate of another, the local exponent is zero at common

primes, 1 at a prime of n/h, and -1 at a prime of n'/h. The common prime still imposes its original zero on rows. Writing $n = hr$, $n' = ht$ therefore gives exactly

$$\chi_n(u)\overline{\chi_{n'}(u)} = \mathbf{1}_{(u,h)=1}\chi_r(u)\overline{\chi_t(u)}, \qquad q = rt, \quad \mathfrak{l} = hrt.$$

Here q is the primitive oscillatory conductor and $\mathfrak{l}$ a period of the full zero extension. These are ideals, not necessarily squarefree rational norms. In particular, conjugate primes over one split rational prime are distinct coprime ideal factors. Full defining moduli of ν and primary conventions remain fixed in S; this row-conductor calculation does not assert a new uniform reciprocity theorem in the column variable. In the coprime bulk $Nq \asymp D^2$ remains large.

N.3 Why a first generic Cauchy step does not supply the saving

On one type-II dyad let $Na \asymp A$, $Ns \asymp B$, $AB \asymp D$. Keep the full profile and coprimality restrictions in

$$X_a(u) = \nu(a)\chi_a(u) \sum_{\substack{s \text{ squarefree} \\ Ns \asymp B,\ (a,s)=1}} g_U(s)\nu(s)\chi_s(u)W(NaNs/D).$$

The exact elementary Cauchy inequality is

$$\sum_{0<Nu\le H} \left| D^{-1/2} \sum_{Na\asymp A} \mu(a)X_a(u) \right|^2 \le \frac{\sum_{Na\asymp A}|\mu(a)|^2}{D} \sum_{Na\asymp A} \sum_{0<Nu\le H} |X_a(u)|^2.$$

The a-mask has been kept inside X_a. Even if one granted an optimistic uniform slice bound $(H+B)\sum_{Ns\asymp B}|g_U(s)|^2 D^{\varepsilon}$, the ordinary divisor estimate $|g_U(s)| \le \tau(s)$ and the fixed-field ideal count would only yield the envelope

$$A(H+B)D^{\varepsilon} \asymp (AH+D)D^{\varepsilon}.$$

That hypothetical slice estimate is not established here. This is an insufficiency of the generic upper-bound calculation, not a lower bound on the signed energy. At balanced $A, B \asymp D^{1/2}$ and $H = D^{3/4}$, the AH term loses $D^{1/2}$ relative to the target; the D term also exceeds H. A sharper use of the actual g_U, cross-factor constraints or covariance between the original pieces is required. This calculation does not refute all bilinear, moving-cutoff or dispersion methods.

The fixed-$U = 1$ lower bounds in Proposition 10.1 are different: they are genuine energy lower bounds for the separate pieces. If $W \ge w_0 > 0$ on $[c_1, c_2] \subset (1,2)$, count primes in $[c_1 D, c_2 D]$ for I_1 and in $[c_1 D/Nq_0, c_2 D/Nq_0]$ for II_1. Removing S and q_0 removes finitely many primes. The prime-ideal theorem gives positive constants times $D/\log D$ in both cases; the latter contribution has coefficient two. Because every coefficient of II_1 is nonnegative, keeping these products is valid. This is unlike isolating the prime part of the full signed Möbius energy, where uncontrolled cross terms remain. For the exact requested bound $H(HD)^{\varepsilon}$ at $H = D^{3/4}$, the competing power is $D^{3/4+7\varepsilon/4}$, giving the threshold $\varepsilon < 1/7$. Replacing $(HD)^{\varepsilon}$ by D^{ε} reparametrizes the loss and gives the equivalent $1/4$ formulation. The positivity argument is specific to $U = 1$. For a concrete control, the primary element $7 - 21\omega$ has norm $637 = 7 \cdot 7 \cdot 13$ and three distinct ideal prime factors. At $U = 7$ its type-I coefficient is -1, its type-II coefficient is -2, and its Möbius value is -1. Thus the separate pieces need not have nonnegative coefficients at another cutoff; no moving-cutoff obstruction follows from the preceding positivity proof.

N.4 Exact finite checks and formal boundaries

The deterministic checker represents every good ideal of norm at most 1024 once by its primary generator, including inert prime ideals and squarefree products of three distinct prime ideals. It checks 1535 coefficient/regrouping cases at five cutoffs, retains 19680 encountered original zeros in actual sextic sums, and verifies 900 pair-conductor exponent ledgers. It evaluates principal ν and the fixed Kummer character $\nu(n) = \chi_n(2)$ on the same good ideals. The finite diagnostic weight is the rational profile $((Nn - D)(2D - Nn))^2/D^4$ on $[D, 2D]$, zero elsewhere; it is not claimed to be the smooth analytic weight of the theorem. Every identity is algebraic for arbitrary finite weights; these samples do not prove seminorm or scale-supremum estimates.

For example, at $D = 128$, $H = 16$, principal ν and $U = 1$, the unit-row normalized sums have the following exact common scaling:

$$\begin{array}{r|r} D^{9/2}M_1 & -122721831 \\ D^{9/2}I_1 & 314726661 \\ D^{9/2}II_1 & 192004830 \end{array}$$

Their exact difference is $II_1 - I_1 = M_1$. The lower-bound theorem uses the prime-ideal proof, not extrapolation from these small numbers. An independent no-production-import oracle also checks convolution, squarefree regrouping and weighted sextic sums.

`MobiusSplit.lean` proves eight actual statements: the inverse-ring identity, natural-number arithmetic truncation/tail, the Möbius Dirichlet identity, the unit cutoff, its exact tail convolution and coefficient, and nonnegativity. The natural-number Dirichlet ring is an actual arithmetic object; it is not the ideal convolution ring. Ideal arithmetic, sextic/Hecke character construction, analytic prime counting and the unproved signed mean square remain prose or named classical inputs. No new zero-free region, full GRH proof or positive determinant bridge has been obtained.

O Primitive conductors, deleted factors and the scalar contour

O.1 Constructing the column character without losing its zeros

Let $\alpha^6 = u$. Since K contains μ_6, the extension $K(\alpha)/K$ is abelian, with character $g \mapsto g(\alpha)/\alpha$. At $p \nmid 6u$, arithmetic Frobenius gives the residue $u^{(Np-1)/6}$, exactly the source's sextic symbol. Abelian reciprocity makes it a finite-order Hecke character; multiply by ν and take its primitive inducing character. The classical reciprocity input is [89, Theorem 0.8 and Chapter I,§1]. The finite-order archimedean character is trivial, since $\mathbb{C}^\times$ is connected.

At $p \nmid 6$, write $u = \pi^e v$ with v a local unit. A sixth root of v lies in an unramified extension: its residue root is separable and lifts by Hensel's lemma. Inertia acts on a sixth root of π through μ_6, so on α the inertia character has order $6/\gcd(6, e)$. It is trivial exactly when $6 \mid e$; its square is trivial exactly when $3 \mid e$. A nontrivial tame one-dimensional character has conductor exponent one. Since ν is unramified outside S, it cannot cancel this inertia there. At primes above 6, the groups $K_p^\times/(K_p^\times)^6$ are finite; their finitely many characters have a uniform maximum conductor. Consequently

$$\mathfrak{f}_u \mid \mathfrak{c}_\nu \prod_{p \mid u,\ p \nmid 6} p, \qquad Q_u \ll_\nu Nu.$$

The fixed ideal $\mathfrak{c}_\nu$ includes the wild and fixed ν factors. No assertion identifies $\mathfrak{f}_u$ with the larger full zero modulus.

For $p \notin T$ put $z = \psi_u(p)(Np)^{-s}$. Its squarefree factor $1 + z = (1 - z^2)/(1 - z)$ proves the ratio using the *same* deleted set in numerator and denominator. Restoring the primitive Euler factors

yields (3). At a prime where the primitive square becomes unramified its value cannot be obtained by squaring the sentinel zero $\psi_u(p) = 0$. For principal ν, $u = \pi^3$ has a quadratic Kummer character whose primitive square is globally principal. Its local value in that square is one, not zero. For $u = \pi^6$ both primitive characters are principal, although the full family still deletes p. These two cases prove the local table directly.

O.2 Why the half-plane above one half is available

For a character η write $L^T(s, \eta)$ for its Euler product with all primes of T removed. For $\sigma > 1/2$ the product $1/L^T(2s, \psi_{u,2})$ converges absolutely and is bounded by $\zeta_K(2\sigma)$, uniformly in the character, T and $t = \Im s$. Every deleted finite factor is nonzero there. Therefore

$$F_{u,d}(s) = L(s, \psi_u) \prod_{p \in T, p \nmid \mathfrak{f}_u} (1 - \psi_u(p)(Np)^{-s})/L^T(2s, \psi_{u,2})$$

has just the stated possible principal pole. This uses absolute convergence at $2\sigma > 1$, not a conjectural zero-free assertion on the critical line. For any fixed $\sigma > 0$ and $\eta > 0$,

$$\prod_{p \in T} (1 + (Np)^{-\sigma}) \ll_{\sigma,\eta} R_T^{\eta}.$$

Indeed $1 + q^{-\sigma} \le q^{\eta}$ for all sufficiently large q; the remaining finitely many prime ideals give a uniform fixed constant. This controls every moving deleted prime; the denominator has already been bounded by the full convergent zeta product.

We use the classical completed Hecke functional equation in the form

$$\Lambda(s, \psi_u) = Q_u^{s/2}(2\pi)^{-s}\Gamma(s)L(s, \psi_u), \qquad \Lambda(s, \psi_u) = w_u \Lambda(1 - s, \overline{\psi_u}), \quad |w_u| = 1.$$

This is a named external theorem, not a newly formalized analytic result; compare [88, Theorem 4.3, Corollary 4.4]. The cited notes sketch some of Tate's calculations; their incomplete displayed epsilon formulas are not used as independently checked constants here. At $\Re s = 1 + \eta$, absolute convergence bounds L by $\zeta_K(1 + \eta)$. The functional equation and Stirling's formula give on $\Re s = -\eta$ the bound $Q_u^{1/2+\eta}(1 + |t|)^{1+2\eta}$ up to a fixed constant. Phragmén–Lindelöf on this strip therefore gives, for fixed $0 < \sigma < 1$,

$$|L(\sigma + it, \psi_u)| \ll_{\sigma,\eta} Q_u^{(1-\sigma)/2+\eta}(1 + |t|)^{1-\sigma+2\eta}.$$

For the principal character apply the strip argument to $(s - 1)L(s, 1)/(s + 1)$, then divide back on the fixed line $\sigma < 1$. The constants need not be uniform as $\sigma \to 1$. Polynomial vertical strip growth and finite order, as supplied by classical Hecke theory, justify this application. No Lindelöf or subconvexity bound is inserted.

O.3 Mellin inversion, residues and profile uniformity

Initially, absolute convergence and compact smooth support give

$$S_{u,d}(Y) = \frac{1}{2\pi i} \int_{(1+\eta)} \widehat{W}(s) Y^s F_{u,d}(s)\, ds.$$

Four integrations by parts using $x\partial_x$ give $|\widehat{W}(\sigma + it)| \ll_{\sigma} \mathcal{M}_4(W)(1 + |t|)^{-4}$. Choose $0 < \eta < \min(\varepsilon/8, 1/8)$, decreasing it further when distributing the small losses, and shift to $\sigma = 1/2 + \delta$. The convergent reciprocal Euler product, the finite factor bound and convexity above control the

complete integrand. Their polynomial t growth is integrable against this Mellin decay and makes the horizontal edges tend to zero. The only crossed pole is $s = 1$ when $\psi_u = 1$; evaluating its finite factors gives exactly ρ_T. Taking absolute values on the new vertical line proves (4), after renaming the small losses. This also proves uniformity over any family of profiles with bounded $\mathcal{M}_4$; it does not interchange a profile supremum with an arithmetic mean square.

In the original I_u, the inner profile is exactly $W(Nc/Y)$ with $Y = D/Nd$; no d-dependent derivative enlargement occurs. On the support of β_U, $Nd \le U^2 \le D$, hence $Y \ge 1$. Also $R_T \le C_S N u\, N d$. The principal term is explicitly

$$P_u = \mathbf{1}_{\psi_u=1}\sqrt{D}\,\widehat{W}(1) \sum_{\substack{d \text{ squarefree} \\ (d,S)=1}} \frac{\beta_U(d)\nu(d)\chi_d(u)}{Nd}\rho_{T(u,d)}.$$

This sum retains all original phases and zeros; no positivity of it is assumed. For the error use the actual truncated factorizations, not a free coefficient bound:

$$\sum_{d \text{ squarefree}} \frac{|\beta_U(d)|}{(Nd)^{1/2+\delta-\eta}} \le \left(\sum_{Na\le U} (Na)^{-1/2-\delta+\eta}\right)^2 \ll_{\delta,\eta} U^{1-2\delta+2\eta}.$$

The fixed-field ideal count is $O_K(X)$; dyadic summation proves the last inequality for sufficiently small $\eta < \delta + 1/2$. Inserting this and the conductor/radical bounds proves (5). The elementary lattice count $\#\{u : 0 < Nu \le H\} = O_K(H)$ then gives (6). Rows are not divided by units.

At $H = D^{3/4}$ and $U = D^a$ with $0 \le a \le 1/2$, the power of the resulting error envelope *relative to* H is

$$\frac{3}{8} + \frac{5}{4}\delta + (2 - 4\delta)a$$

up to the arbitrarily small losses. It remains positive for every $0 < \delta < 1/2$. Thus this scalar convexity argument alone supplies no short-row saving. This is a limitation of the estimate just derived, not an obstruction to averaging the numerators over Kummer rows, retaining signed coefficients, or exploiting covariance with type II.

The seminorm uniformity is not a proof of the original scale supremum. For each independently chosen scale D' the displayed outer argument requires $U^2 \le D'$. If a fixed cutoff is retained while D' shrinks, then eventually even $D' \le U$: the short $2a_U$ term in the Vaughan identity can meet the support and must be restored. Neither that term nor the covariance with type II may be omitted in a supremum or reassembly.

O.4 Verification boundaries

`HeckeTypeI.lean` checks seven actual complex local/finite-product statements, including independent primitive coefficient arrays and their full deleted-factor restoration. It does not identify an ideal Hecke series with an abstract finite product or formalize a Mellin contour. The exact checker includes split and inert prime ideals, actual sextic phases, conductor drops, sixth-power rows and original mask zeros. The analytic theorem above uses classical abelian reciprocity, Hecke continuation/functional equation and standard complex-analytic tools as external inputs. Full ideal-character/formal analytic objects, the original signed low-gcd mean square and every GRH completion obligation remain open.

P Controlled errors and a trained full-action RG approximation

Proposition P.1 (Classical action-error propagation)**.** *If* $|\widehat{V}(s,t) - V(s,t)| \le \varepsilon$ *at every finite configuration, then* $|\mathcal{R}\widehat{V}(s) - \mathcal{R}V(s)| \le \varepsilon$*. The partition ratio lies in* $[e^{-\varepsilon}, e^{\varepsilon}]$*. Every further* exact *marginalization preserves the same action bound.*

Proof. The pointwise inequalities $e^{-\varepsilon}e^{V} \le e^{\widehat{V}} \le e^{\varepsilon}e^{V}$ survive finite summation. Taking logarithms proves the first assertion; summing also over the retained variables proves the second. Iterate the same argument. For normalized measures the likelihood ratio lies between $e^{-2\varepsilon}$ and $e^{2\varepsilon}$, so a bounded observable error is at most $\|O\|_\infty \min(2, e^{2\varepsilon} - 1)$. Per-step learned errors would add; a single initial error does not grow under subsequent exact elimination. This scalar proof is not an assertion that the matrix exponential is operator-monotone. □

P.1 Executed finite calculations

The implementation checks full enumeration against the closed black-spin formula, Walsh reconstruction, raw partition equality, retained-spin magnetizations and two-stage versus direct elimination. At the quantum $\Gamma = 0$, $\beta = 1$ limit, every Pauli coefficient agrees with the classical effective action, with the required minus sign between energy and logweight. At nonzero transverse field the model is

$$H = -J_+ Z_+ \sum_j Z_j - J_- Z_- \sum_j \eta_j Z_j - bZ_+ + \tfrac{3}{5} bZ_- - \sum_j h_j Z_j - \Gamma \sum_i X_i.$$

At general β, the classical zero-transverse-field limit uses βV, including the fixed white fields, rather than scaling only the three varied controls. For $m = 5$, its full dimension is 128 and the first retained dimension is 32. Every one of the 1024 five-spin Pauli coefficients is computed; later retained dimensions are 8,4,2. Spectral-log/exponential and full Pauli reconstruction residuals, exact-trace composition and retained-observable comparisons are recorded separately. Floating-point residuals are numerical checks of the proved finite identities, not interval certificates.

The local laboratory exposes every scale, coupling/field controls for the classical model, selected generated interaction supports on actual dessin radial lifts, the even-chain K, h, C trajectory and physical correlation length, and three fixed quantum matrix calculations. Live quantum parameter sliders are not offered for uncomputed Hamiltonians. Pair truncation carries the identity term and reports both partition and probability/density errors. A late-scale floating-point loss of the small eigenvalue is avoided by transporting the exact log-eigenratio coordinate $\log r \mapsto 2 \log r$; this is explicitly the transfer RG identity, not a newly diagonalized value.

P.2 A quantum-feature model trained on actual RG

The learner targets all 32 unnormalized logweights of the $7 \to 5$ spin $m = 5$ decimation. Its three inputs are (J_+, J_-, b), with ranges $[0.15, 1.1]^2 \times [-0.6, 0.6]$. Fixed fields are $h_j = 0.045 \cos(1.7j) + 0.02j/5$. Seed 20261103 fixes 160 input samples, a 120/40 train/test split, and 40 OOD samples with $J_\pm \in [1.15, 1.5]$, $b \in [-0.8, 0.8]$. The three feature qubits encode the couplings by $R_y(1.5J_+) \otimes R_y(1.5J_-) \otimes R_y(2b)|000\rangle$. Two local R_y, R_z layers, with a CZ triangle after each, contain 12 nominal optimizer angles. The final three R_z phases commute with the measured Z products and are structurally inactive for this loss; they still change the displayed state phases. We count them as nominal parameters, not additional effective expressive directions. Seven nonempty Pauli-Z products feed an affine 8×32 readout. There are 268 parameters. Feature registers are not the seven physical spin sites or five quiver vertices.

The training objective is full-action MSE, including the constant; KL alone would ignore that constant. A training-only ridge fit initializes the head, then 120 Adam updates optimize both circuit and head. Actual PennyLane analytic expectation calls implement the feature circuit. Parameter-shift derivatives of each feature are combined with the classical head gradient; full-loss finite differences agree to 6.46×10^{-11}. All data, parameters, intermediate checkpoints and complete loss curves are preserved.

The held-out full-action MSE is approximately 0.00985, mean $D_{\mathrm{KL}}(p_{\mathrm{exact}} \| p_{\mathrm{learned}}) = 0.00300$, mean TV 0.0290, and mean absolute log-partition error 0.0610. A classical quadratic-feature baseline with the same 256 readout parameters obtains 0.00613,0.00240,0.0233 and 0.0138, respectively. OOD action MSE increases to 0.267 for the quantum hybrid, versus 0.0555 for the baseline. The quantum-feature constant MAE is 0.0444 and higher-body coefficient RMSE is 0.00900, versus 0.0124 and 0.0130 for the classical baseline; different metrics favor different aspects. The aggregate results show no quantum advantage. They expose genuine full-action approximation errors rather than substituting state-compression success for RG.

Each saved example computes its maximum error over all 32 coarse states; the proposition then applies to its subsequent exact marginalizations. These per-example finite bounds are not uniform over the input parameter box. The synchronized architecture inspector shows literal gate outputs, Pauli features, head, logweights, generated interactions and partition value, linked to the exact genus-two dessin/order context. No full order representation, continuum universality, material-specific accuracy or arithmetic zero-location bridge is inferred.

P.3 Quantum TokenGT and the geometric error-correction boundary

A further executed node/edge-token hybrid uses the same exact RG teacher, with frozen three-qubit features, trained Q/K/V and a full-action head. Softmax, componentwise max-plus and hybrid variants are compared with a matched classical feature model; the latter has lower held-out action error. An explicit intervention-code/change variational MI objective learns concept directions; a 61-candidate latent forest and a held-out analogy transfer are scored on exact physical observables. Rips H_1 is actually reduced over $\mathbf{F}_2$, and a continuous Gaussian GFlowNet trains a trajectory-balance objective with specified root/Lebesgue measures. Empirical residuals do not prove exact terminal sampling, semantic reasoning or categorical equivalence.

The physical bit-flip repetition channel is separately executed and has $p_L = 3p^2 - 2p^3$ under ideal recovery. A genuine cellular CSS calculation on the genus-two one-face dessin gives $[[10, 4, 1]]$: vertex-incidence rank 6, zero face boundary, X-distance 1 and Z-distance 4. Thus this cellulation provides no useful protection merely from its genus. These finite layers, full formulas, proofs, parameters and scope are developed in the research book and interactive laboratories; the physical models are not faithful representations of the infinite completed order and supply no GRH bridge.

P.4 An actual finite gentle cut for the categorical interface

On the genuine $m = 5$ dessin quiver write $a_i, b_i : i \to i - 1$ with indices modulo 5; cross-branch consecutive pairs are forbidden. Remove both arrows with source 0. The surviving algebra B has vertices 0 through 4, arrows a_i, b_i for $i = 1, \ldots, 4$, and relations $a_i b_{i-1} = b_i a_{i-1} = 0$ for $i = 2, \ldots, 4$. It is an explicit finite gentle quotient, not the completed order itself.

Every surviving positive path stays on one branch and has length at most 4. For length $\ell = 1, \ldots, 4$ there are $2(5 - \ell)$ paths, while length 0 contributes five idempotents. Therefore $\dim_k B = 5 + 2(4 + 3 + 2 + 1) = 25$. The forbidden-chain resolution already proved for monomial gentle algebras has two alternating chains of length n from vertex i to $i - n$ when $1 \le n \le i$,

and none for $n > i$. Applying Hom to simples kills its arrow differentials. Hence $\operatorname{pd}_B S_i = i$ and $\operatorname{Ext}^4_B(S_4, S_0) \cong k^2$. The split finite-dimensional simple-module criterion gives $\operatorname{gldim} B = 4$.

Every full branch cycle crosses a removed arrow, so the common completed order parameter u maps to zero. Thus B is a finite u-torsion quotient, not a DVR-free surface order. The difference is visible in derived Hom: S_0 is projective over B, whereas in the original completed order the two forbidden arms of length 5 return to vertex 0 and give $\operatorname{Ext}^5_\Lambda(S_0, S_0) \cong k^2$. The exact restriction along $\Lambda \twoheadrightarrow B$ is consequently not derived fully faithful. This supplies a concrete cut on which to investigate graded gentle/Fukaya models, while proving why its equivalence cannot simply be transported to the original completed order. Boundary/stops/line-field data and an actual module-valued neural decoder would still have to be supplied. Exact path and endpoint enumeration accompanies the prose proof; this cut/category comparison is not claimed wholly Lean formalized.

Q Relation calculations, splitting functors and determinant limits

Q.1 Correct graphs and stable spaces

In the original basis let $fe_1 = e_2$, $fe_2 = e_3$, $fe_3 = 0$ and $ge_1 = 0$, $ge_2 = e_1$, $ge_3 = e_2$. Then

$$\begin{aligned}\Gamma_f &= \langle (e_1, e_2), (e_2, e_3), (e_3, 0)\rangle,\\ \Gamma_{f^2} &= \langle (e_1, e_3), (e_2, 0), (e_3, 0)\rangle,\\ \Gamma_{gf^2} &= \langle (e_1, e_2), (e_2, 0), (e_3, 0)\rangle.\end{aligned}$$

Each graph has dimension three. Omitting zero-output generators changes the relation into a partial graph and changes its powers and domain. The following are subspaces of V, not lists of pairs in $V \oplus V$:

T	$\bigcup_r \ker T^r$	$\bigcap_r \operatorname{im} T^r$
f, g	V	0
$gf = \operatorname{diag}(1,1,0)$	ke_3	$ke_1 \oplus ke_2$
$fg = \operatorname{diag}(0,1,1)$	ke_1	$ke_2 \oplus ke_3$

For all four full graphs $\bigcup_r \Gamma^r_T(0) = 0$. This corrects both the source's graph convention and its interchanged gf, fg kernel entries. For an arbitrary multivalued relation there need not even be a linear map on its domain; its regular-part theorem requires the appropriate relation subquotients and finiteness conditions. We do not replace those hypotheses by the graph calculation. For comparison, in [38, Section 4, Lemmas 4.2,4.5–4.6] write $C' = \bigcup_n C^n(0)$ and let C'' consist of vectors with a compatible infinite backward chain. In finite dimension $C'' = \bigcap_n C^n V$; this equality is not automatic for arbitrary infinite-dimensional spaces. The regular quotient is $C^\sharp/C^\flat$, where

$$C^\sharp = C'' \cap (C^{-1})'', \qquad C^\flat = (C'' \cap (C^{-1})') + (C' \cap (C^{-1})'').$$

The automorphism statement uses this quotient for an arbitrary relation. The splitting statement additionally requires its finite-dimensionality, rather than the source's unqualified ι/κ expression.

Q.2 Proof of the dessin-category equivalence

Write $a_i : V_i \to V_{i-1}$ for $i = 1, \dots, m-1$ and $p_i = a_1 \cdots a_i : V_i \to V_0$, with $p_0 = 1$; the rightmost map acts first. Every p_i is an isomorphism. Conjugating the arrow maps by these identifications makes $a_i = 1$ for $i \neq 0$, and makes $T = p_{m-1}a_0$ at the remaining arrow. All b-relations vanish. An intertwiner is transported to maps h_i with $h_i = h_{i-1}$ for $i \neq 0$ and $h_{m-1}T = T'h_0$. Hence every $h_i = h$ and $hT = T'h$. This proves full faithfulness and essential surjectivity, with inverse given

by $(V,T) \mapsto N_m(T)$. It is also an exact equivalence: kernels and cokernels of intertwiners are the usual invariant kernels/cokernels; the identity arrow maps remain identities. Finite dimension is a condition on modules, while A_m itself is infinite dimensional through its arbitrarily long permitted branch paths.

The sum of the full permitted turns on both branches is central in this closed dessin algebra. On the chosen modules the b-turns vanish and each a-turn equals T, so u acts as T at every vertex. If $T^n = 0$, the action factors through $A_m/(u^n)$ and extends to the completion. If T has a nonzero invertible Fitting summand, no power of u kills that summand, precluding continuity. For $T = J_r(\lambda)$ with $\lambda \neq 0$, the obstruction is stronger: $u - \lambda$ is a unit of $k[[u]]$ but its rth power annihilates the proposed module, impossible for a nonzero unital module. This rules out even an unqualified algebraic extension of that band, not just a chosen topology.

Q.3 Functoriality and the actual filtration

Kernel dimensions are nondecreasing and at most n. Once two successive kernels agree, applying T^{-1} shows that all later kernels agree. Thus they stabilize by n; rank-nullity gives image stabilization as well. On $K(T)$, $T^n = 0$. On $I(T)$, $T(I(T)) = I(T)$; finite dimension makes this a bijection. If $hT = T'h$, induction gives $hT^r = T'^r h$, hence

$$h(\ker T^r) \subseteq \ker T'^r, \qquad h(\operatorname{im} T^r) \subseteq \operatorname{im} T'^r.$$

This proves functoriality even when $\dim V \neq \dim V'$: choose a common stabilization exponent for the two finite objects. The decomposition projection onto $K(T)$ along $I(T)$ is canonical and commutes with all intertwiners; the existence of a Jordan basis is not needed. Finite dimension is essential. On $V = k[t]$, multiplication by t has stable kernel zero and stable image $\bigcap_r t^r k[t] = 0$, although $V \neq 0$. The same construction $N_m(V,t)$ is an actual infinite-dimensional module of A_m. Thus the finite Fitting split cannot be extended to every module of our infinite algebra merely by repeating its notation.

Let $k_r = \dim \ker T^r$, with $k_0 = 0$, and form $G_r = \ker T^r / \ker T^{r-1}$. For a nilpotent Jordan block of size s, $k_r = \min(r,s)$, so $\dim G_r$ counts the blocks of size at least r, and $2k_r - k_{r-1} - k_{r+1}$ counts blocks of size exactly r. For $J_3(0) \oplus J_1(0) \oplus J_2(1)$ the stable dimensions are 4 and 2; the kernel dimensions begin $0, 2, 3, 4, 4$. The G_r are functorial subquotients. Their dimensions do not specify a natural choice of basis or split each filtration step.

Proposition Q.1 (What the source's splitting-functor assumptions imply)**.** *Let $\mathcal{A}, \mathcal{B}_i$ be additive categories with finite biproducts. Suppose additive functors $S_i : \mathcal{B}_i \to \mathcal{A}$ and $F_i : \mathcal{A} \to \mathcal{B}_i$ satisfy $F_i S_i \cong 1$, $F_i S_j \cong 0$ for $i \neq j$; only finitely many $F_i M$ are nonzero for each M; the F_i jointly reflect isomorphisms; and maps $\gamma_{i,M} : S_i F_i M \to M$ have invertible F_i-images. Then $M \cong \bigoplus_i S_i F_i M$. Every nonzero indecomposable is $S_i B$ for a unique index and an indecomposable B up to isomorphism.*

Proof. Sum the finitely many nonzero $\gamma_{i,M}$ to a map γ_M. After applying F_j, all off-diagonal summands vanish, leaving the assumed isomorphism $F_j \gamma_{j,M}$, or $0 \to 0$. Joint reflection therefore makes γ_M invertible. For an indecomposable M, exactly one of the nonzero summands can occur. Conversely, if $S_i B = X \oplus Y$ and B is indecomposable, applying F_i makes one summand vanish, say $F_i X = 0$; every other $F_j X$ also vanishes. Joint reflection applied to $0 \to X$ gives $X = 0$ up to isomorphism. Finally apply F_i to an isomorphism $S_i B \cong S_j B'$ to recover both index and object. □

The direction of γ matters: it is into M, contrary to one source diagram. Local finiteness is objectwise in i, not a finiteness assertion over all objects of a category. Abelian categories require kernels, cokernels and the usual exactness axioms; addition is on Hom groups, not on an alleged

group of quiver vertices. The abstract assumptions above do not assert that the family F_i is faithful on arbitrary morphisms. Our Fitting example is concrete: $\mathcal{B}_0$ consists of nilpotent pairs, $\mathcal{B}_\times$ of invertible pairs, S are inclusions and $F_0, F_\times$ the two stable summands.

Q.4 Strings, bands, Hom and one non-rigid string

In this branch category, Jordan normal form classifies indecomposables over an algebraically closed field. Over a general field replace Jordan eigenvalues by powers of monic irreducible polynomials. A Jordan block $J_r(0)$ is one directed string with mr basis visits and $mr-1$ nonzero successor arrows; $J_r(\lambda)$ for $\lambda \neq 0$ is a permitted cycle band. Formal inverse letters are reverse traversals, not inverse elements of A_m. Strings and their reverses describe the same string module under the usual isomorphism convention. Arbitrary infinite modules are not all classified by this finite calculation; the actual finiteness hypotheses in [38] remain essential.

Identify a Jordan pair with $k[t]/(t-\lambda)^r$. An intertwiner from size r to size s is determined by the image of 1, annihilated by $(t-\lambda)^r$. For equal eigenvalues this subspace has dimension $\min(r,s)$. For different eigenvalues, Bezout's identity makes $t-\lambda$ invertible on the target, so only zero remains. Full faithfulness above transfers this Hom computation to A_m. The sequence $0 \to J_1(0) \to J_2(0) \to J_1(0) \to 0$ is not split: with $J_2(0)e_1 = e_2$, a lift of the quotient generator has nonzero first coordinate and is not killed by $J_2(0)$. Its image under N_m is a nonsplit ambient-A_m self-extension. Thus even the string $N_m(J_1(0))$ is not rigid. No assertion that this exact full subcategory is derived fully faithful is needed for the nonsplitting argument.

For this particular closed A_m, every oriented or inverse letter has exactly two admissible successors: continue on its branch in the same orientation, or switch branch and reverse orientation. Cancellation and forbidden pairs, including inverse forbidden pairs, remove the other choices. A fixed ordering of these choices identifies each first-letter cylinder with $\{0,1\}^{\mathbb{N}}$. The whole infinite-word space is a finite disjoint union of Cantor spaces, hence itself Cantor. There are no terminating words inside that space to remove. Arbitrary gentle quivers can have isolated or eventually forced paths, so this conclusion is not universal. Algebraic covers retain the lifted zero-relations and deck equivariance data; arbitrary periodic cover representations are not automatically bands or a unitary time evolution.

Q.5 A determinant comparison with its exact limits

On $\bigoplus_i V_i$, $L^m = \mathrm{diag}(T, \dots, T)$. For a direct determinant proof of (7), successively eliminate the blocks 1 at vertices $1, \dots, m-1$ in $1 - zL$. Block row additions preserve determinant and leave $1 - z^m T$ at vertex zero. This works over $k[z]$ without inverting z or T; the zero-size case gives one on both sides. The Fitting splitting separates a nilpotent factor 1 and the determinant on the invertible summand. For instance $m = 5$, $T = J_2(2)$ gives $(1 - 2z^5)^2$, whose complex roots have modulus $2^{-1/5}$, despite being a genuine genus-two dessin module.

Under conjugation of T this is an invariant polynomial. With independent source/target bases a presentation determinant is instead a semi-invariant with its determinant-character weight; those actions must not be conflated. The polynomial equality does not construct a positive Hermitian metric, an arithmetic Euler factor, a regularized determinant or a zero-preserving map to completed L-functions. Functorial kernel filtrations similarly do not construct an equivalence of derived/Fukaya categories. A fixed graded surface category and a determinant-compatible arithmetic realization remain separate obligations.

The exact rational checker evaluates actual genus-two/four quivers, Hom equations, full-turn matrices, both filtration directions and the nonsplit extension. `lean-analytic/LinearRelations.`

`lean` proves full graph identities, semiconjugacy and filtration preservation, actual finite-space Fitting splitting/stabilization and unit-power annihilation. It does not package the entire dessin category, Jordan classification, general splitting-functor proposition or determinant proof in Lean.

R Normal forms and the full module action

R.1 Why the polynomial matrix algebra is the path algebra

With our convention a path from j to i acts as E_{ij}; the rightmost arrow acts first. Every nonzero positive path uses only the a-branch or only the b-branch, because mixed consecutive arrows are the defining quadratic zero relations. Along either branch the path of length ℓ starting at j is unique. Write

$$\ell = j - i + 5q, \qquad 0 \le i, j < 5.$$

Its image is $t^q E_{ij}$ in the chosen factor and zero in the other. For $i < j$, $q \ge 0$; for $i > j$, $q \ge 1$; for $i = j$ and positive length, $q \ge 1$. The stationary path is the shared (E_{ii}, E_{ii}).

These images are linearly independent: off-diagonal matrix units, positive diagonal monomials, branch factors and degrees are independent; the shared constant diagonal units are independent of all positive paths. They span $H \times_{k^5} H$, since upper entries permit all polynomials, lower entries are divisible by t, and only the constant diagonals of the two factors must agree. Products of paths agree with matrix products. Thus the arrow assignment induces a bijective algebra map from $kQ/(a_i b_{i+1}, b_i a_{i+1})$ to B. This argument includes arbitrarily long paths; it imposes no nilpotence relation on a full turn.

The central parameter u, the sum of all full turns on both branches, has image (tI_5, tI_5). Completing in this parameter is a separate operation. The polynomial algebra and its completion are not identified by the normal-form calculation. The finite modules below are annihilated by u^2, so their continuous completed actions require only a finite Taylor polynomial.

R.2 Coefficient selectors without a truncation

Here is an explicit proof of Theorem 14.2 and the coordinate convention used by the implementation. Let

$$\mathcal{S} = \{(i, j, q) : 0 \le i, j < 5,\ q \in \mathbb{N},\ i < j \text{ or } q > 0\}, \qquad \mathcal{K} = \{0, \dots, 4\} \sqcup (\{a, b\} \times \mathcal{S}).$$

For a stationary index set $v_i = (E_{ii}, E_{ii})$; for a positive index set $v_{a,i,j,q} = (t^q E_{ij}, 0)$ and $v_{b,i,j,q} = (0, t^q E_{ij})$. These are elements of the ambient polynomial matrix-pair space, with no degree cutoff. For $M = (M_a, M_b)$ define

$$\lambda_i(M) = [t^0](M_a)_{ii}, \qquad \lambda_{b,i,j,q}(M) = [t^q](M_b)_{ij}.$$

Here the subscript b in the second formula denotes either branch. Every λ_x is k-linear, since extraction of a fixed matrix entry and polynomial coefficient is linear.

We verify $\lambda_x(v_y) = \delta_{xy}$, including the cases at degree zero. For two stationary indices this is the ordinary diagonal matrix-unit identity. A stationary selector vanishes on a positive image in the other factor. In its own factor it could see a constant diagonal only if $i = j$ and $q = 0$, which the definition of $\mathcal{S}$ excludes. Conversely, a positive selector applied to a stationary image could be nonzero only at precisely the same forbidden slot. For two positive indices, different branches have disjoint supports. In one branch the matrix entry forces both endpoints to agree and the monomial coefficient forces the degrees to agree. These conditions give equality of the full indices. The witness

of the condition defining $\mathcal{S}$ carries no extra algebraic data. This proves biorthogonality in every characteristic, using no division.

For any finitely supported $c : \mathcal{K} \to k$, linearity now gives

$$\lambda_x \left(\sum_{y \in \mathcal{K}} c_y v_y \right) = c_x.$$

If the sum vanishes, every c_x vanishes; if two sums agree, their coefficients agree. This proves linear independence and injectivity of normal expansion on the actual k-module of finitely supported functions. It does not replace finite support by a convergent or formal infinite sum.

The length formula also contains useful information beyond independence. If $i < j$ then $5q+j > i$ even at $q = 0$; otherwise $q > 0$ and $i \leq 4$ give the same strict inequality. Hence $\ell = 5q + j - i$ is a positive integer, and $\ell + i = 5q + j$. For fixed starting vertex j and length ℓ, two such equations imply $i - i' = 5(q - q')$. Since $i, i' \in \{0, \ldots, 4\}$, this forces $i = i'$ and then $q = q'$. In Lean the subtraction is natural subtraction; positivity is proved before using the rearranged equation.

For example the a-path of length 12 from vertex 4 ends at vertex 2 and has image $(t^2 E_{24}, 0)$. The length-4 a-path from 3 to 4 has image $(tE_{43}, 0)$. Their product, with the rightmost path first, is $(t^3 E_{23}, 0)$, of length 16. Replacing the second path by its b-branch version makes the product zero. Degree-zero upper entries such as $(E_{24}, 0)$ are permitted, but $(E_{22}, 0)$ is not a stationary image: the stationary element is the shared pair (E_{22}, E_{22}). This last distinction prevents a spurious extra idempotent in the proposed basis.

The selector theorem alone gives the faithfulness part of the intended quotient comparison. The next construction proves membership and spanning inside the generated algebra without assuming that comparison. A separately defined path algebra, mixed-zero quotient and its universal algebra map still require formal packaging. Completion, module-category and almost-split universal properties require their own subsequent constructions. None of these finite-support identities supplies an arithmetic self-adjoint determinant or controls zeta zeros.

R.3 Constructing every normal image from actual arrows

We now prove Theorem 14.3. This closes the membership and spanning gap left by ambient coefficient recovery. Let $p(j) = j - 1$ modulo five and, inside the actual generated algebra, define

$$W_b(0, j) = e_j, \qquad W_b(n + 1, j) = W_b(n, p(j))\, b_j.$$

In this formula b_j denotes the arrow in the chosen branch; it may be an a-arrow. The rightmost arrow acts first. This recursive definition is an element of A for every n, because it uses only its generators and multiplication. Set

$$d_{ij} = \begin{cases} 0 & i < j, \\ 1 & i \geq j, \end{cases} \quad L_{ij} = j - i + 5d_{ij}, \quad P_b(i, j) = W_b(L_{ij}, j), \quad C_b(i) = P_b(i, i).$$

The integer L_{ij} lies between one and five. In particular $P_b(i, i)$ is a positive full turn, whereas $W_b(0, i)$ is the shared stationary idempotent. These two elements are not interchangeable.

Directly following the arrows, the path $P_b(i, j)$ crosses the weighted arrow b_0 exactly d_{ij} times. Its image is therefore $t^{d_{ij}} E_{ij}$ in factor b, with zero in the other factor. The Lean proof checks all five-by-five endpoint pairs for each of the two branches; this is a finite verification of a finite combinatorial case split, not a bound on permitted path lengths. For all $n \in \mathbb{N}$, induction gives

$$\mathrm{view}\big(C_b(i)^n P_b(i, j)\big) = t^{n + d_{ij}} E_{ij} \quad \text{in factor } b.$$

For $n = 0$ the power is the unit of the whole two-factor algebra, so multiplication by $P_b(i, j)$ still selects the correct branch. The induction step uses $C_b(i)$ with image tE_{ii} and $E_{ii}E_{ij} = E_{ij}$. No assertion that a branch-only degree-zero diagonal belongs to the algebra is needed. Since a permitted slot satisfies $q \geq d_{ij}$, the choice $n = q - d_{ij}$ constructs its preimage in A. This proves membership at every polynomial degree.

For clarity, if $i = 2$, $j = 4$, $q = 2$, then $P_a(2, 4) = a_3a_4$ and $C_a(2)^2P_a(2, 4)$ realizes $(t^2E_{24}, 0)$ by a length-12 arrow word. For $i = 4$, $j = 0$, $q = 1$, we have $P_a(4, 0) = a_0$ and the exponent $q - d_{40}$ is zero. The corresponding exponent must never be replaced by a negative power or by an inverse arrow. The construction works over $\mathbb{F}_2$ as well as characteristic zero.

R.4 Spanning, multiplication and the inverse coordinate map

Let S be the k-linear span of all normal images in the ambient space. Membership just proved gives $S \subseteq A$. We prove the reverse inclusion by showing that S is closed under the operations used to generate A. The product rules are

$$e_ie_j = \delta_{ij}e_i, \qquad e_rv_{b,i,j,q} = \delta_{ri}v_{b,i,j,q}, \qquad v_{b,i,j,q}e_r = \delta_{jr}v_{b,i,j,q},$$

and for two positive images

$$v_{b,i,j,q}v_{c,r,l,s} = \begin{cases} v_{b,i,l,q+s} & b = c \text{ and } j = r, \\ 0 & \text{otherwise.} \end{cases}$$

The displayed nonzero output is always an admissible positive image. If $q + s > 0$ this follows from its degree. If $q + s = 0$, nonnegativity gives $q = s = 0$, so the input conditions imply $i < j = r < l$. Thus $i < l$ and the output is admissible even at degree zero. This explains why closure cannot be justified merely by ignoring the lower-triangular constraints.

By bilinearity the product of two finite linear combinations lies in S. Every vertex is one of its normal images, and each arrow is a permitted image with degree zero except the wrap arrow, which has degree one. The sum of the shared vertices is the unit. Scalar multiples of this unit belong to S. Induction on the algebra generated by the arrows and vertices therefore gives $A \subseteq S$. In Lean this is actual `Algebra.adjoin_induction`, not a predicate that assumes the desired normal form for every element. Consequently $A = S$ as submodules of the polynomial matrix-pair space.

Now lift each v_x to an element $\widetilde{v}_x \in A$ using its proved membership. Their images under the injective inclusion are the already independent ambient images, so the lifts are independent. The equality $A = S$ gives surjectivity of finite expansion; independence gives injectivity. This constructs the linear equivalence E on the genuine generated algebra. For any $a \in A$, apply the ambient coefficient recovery identity to $E^{-1}a$ to obtain

$$(E^{-1}a)_x = \lambda_x(\mathrm{view}(a)).$$


The formula holds for every element of A, not only on a sampled list of paths or generators. Its finite support follows from the equivalence with finitely supported functions. No convergence of an infinite expansion is asserted. Although the product table has been used to prove span closure, we have not yet supplied the formal universal map from a separately constructed bound-quiver path quotient. The distinction between a linear coordinate equivalence and that algebraic quotient interface is retained.

R.5 Exactness for arbitrary algebra elements

In coordinates $M_2 = k^5 \oplus \epsilon k^5$ let $C(t) = \sum_{r \geq 0} C_r t^r$ be the first component of an arbitrary B-element. Its action is

$$(x, y) \longmapsto (C_0 x, C_1 x + C_0 y), \qquad \rho_2(C) = \begin{pmatrix} C_0 & 0 \\ C_1 & C_0 \end{pmatrix}.$$

Evaluation is multiplicative: the constant coefficient of a product is $C_0 C_0'$ and the linear coefficient is $C_0 C_1' + C_1 C_0'$. Consequently the formula is a representation of the entire algebra. The second branch acts by zero but has not been removed from the algebra. Both maps in the sequence commute with this general block action, not merely with a finite test list of generators.



Every vector in $\ker p$ is $(0, y) = i(y)$, i is injective, and p is surjective by $x = p(x, 0)$. The section obstruction uses only a_0: it is zero on M_1, while on M_2 it sends the coordinate-zero constant coefficient to the coordinate-four infinitesimal part. A section commuting with this arrow would force $1 = 0$. No algebraic closure, eigenvalue classification, division by two or positivity premise occurs.

R.6 Formal and computational coverage

The file `lean-analytic/BranchExtension.lean` defines B as the actual generated subalgebra of polynomial matrix pairs, proves its infinite dimensionality through injective polynomial evaluation, and constructs i, p as B-linear maps. The original 21 printed theorem types and axiom reports cover the generators, infinite embedding, actions, exactness and absence of a module section. Eight additional reports in `NormalPath` cover slot extensionality, positive length, the length equation, recovery from start and length, biorthogonality, coefficient recovery, linear independence and injective normal expansion. Ten further printed reports now establish actual generated-algebra membership, equality with the normal span, lifted independence, bijective finite expansion and its explicit inverse coefficients. The normal-form isomorphism with the separately constructed path quotient above is proved in prose; it is not packaged as a Lean categorical equivalence. Nor is a Yoneda-Ext object or the universal factorization property of an almost-split sequence formalized by these maps alone.

The exact checker `examples/relations/check_extension.py` independently uses unbounded normal paths and polynomial matrix pairs. It checks 405 distinct paths through length 40, 7,225 products, 30 distinct loop powers and 80 whole-polynomial module actions. The full generator system for a hypothetical section has coefficient rank 49 and augmented rank 50 over $\mathbb{Q}, \mathbb{F}_2, \mathbb{F}_3, \mathbb{F}_5$. These finite diagnostics complement the symbolic proof; their finite cutoffs do not establish infinite dimensionality by themselves.

The separate exact checker `examples/relations/check_normal_paths.py` checks 475 normal images at degrees up to 1,000,003 (length 5,000,019), all 225,625 selector pairs, 1,030 direct arrow steps, 15,625 composable-or-zero product comparisons and 200 finite expansions over $\mathbb{Q}, \mathbb{F}_2, \mathbb{F}_3, \mathbb{F}_5$. Invalid constant slots are rejected. These are finite diagnostics of the code and conventions; the unbounded assertion is established by the Lean theorem, not by the size of the tests.

A further deterministic checker, `examples/relations/check_normal_algebra.py`, constructs all 50 shortest arrow words and 375 actual preimages, including degree 1,000,003, using sparse binary powers of the full turns. It checks 5,625 product round trips and 200 exact expansions over the four stated fields. Three ambient matrices with forbidden constant shapes fail reconstruction,

as required. These diagnostics test the implementation independently of finite degree bounds; the general membership/span/equivalence result is the Lean theorem, not an extrapolation from these samples.

S Physical generative models and their verification

S.1 From a dessin to an actual quantum RG target

Take $y^2 = x^5 - 1$, $\beta = x^5$. Its clean incidence graph has two black vertices and five white vertices, hence seven physical spin sites and ten incidence edges. These differ from the five quiver vertices and the infinite-dimensional polynomial algebra. The physical Hilbert space below is finite; no faithful action of that whole algebra on it is assumed.

Let X_q, Z_q be Pauli matrices on physical site q, with the black sites numbered zero and one and the white sites $2 + j$. Our executed Hamiltonian is

$$\begin{aligned} H = & -.50 \sum_{j=0}^{4} Z_0 Z_{2+j} - .35 \sum_{j=0}^{4} (-1)^j Z_1 Z_{2+j} \\ & - .55 \sum_{q=0}^{6} X_q - .12 Z_0 + .072 Z_1 - \sum_{j=0}^{4} h_j Z_{2+j}. \end{aligned}$$

Here $h_j = .045 \cos(1.7j) + .02j/5$. All coefficients are dimensionless and inverse temperature is one. Diagonalizing the actual 128×128 Hermitian matrix gives $\rho = e^{-H} / \operatorname{tr}(e^{-H})$. We retain the two black spins:

$$\rho_R = \operatorname{Tr}_{\text{white}} \rho, \qquad (\rho_R)_{ab} = \sum_{j=0}^{31} \rho_{aj,bj}.$$

This is an actual coarse quantum state, with trace one and positive spectrum. An independent Pauli–Kronecker NumPy reconstruction agrees with the saved matrix to $2.04 \cdot 10^{-10}$. Its minimum eigenvalue is approximately 0.0806414, and its off-diagonal Frobenius norm is 0.300551. There is therefore quantum coherence that a probability histogram alone cannot describe. No chemical or experimental material dataset is involved: this is an exactly specified finite quantum-matter benchmark.

Let $p_s = (\rho_R)_{ss}$ for $s \in \{\pm 1\}^2$. The hybrid samplers learn

$$p_\sigma(x) = \sum_s p_s \mathcal{N}(x; s, \sigma^2 I_2), \qquad \sigma = .2.$$

This is a continuous dequantization of the measured spin law. A sign measurement of these samples differs from p by at most $2\Phi(-1/\sigma)$ in total variation, by a coordinatewise sign-flip coupling and the union bound. This is below $6 \cdot 10^{-7}$ here. It is small compared with finite-sampling error, but it is not zero. The samplers do not recover phases or the entire ρ_R. For example, ρ_R and $(Z \otimes I)\rho_R(Z \otimes I)$ have the same diagonal but different density matrices.

S.2 Diffusion and exact versus approximate reverse controls

For $T = 64$ use the cosine cumulative noise schedule, with endpoint offset .008, clipped step variances between .0001 and .999. Write $\alpha_k = 1 - \beta_k$ and $\bar{\alpha}_k = \prod_{j \le k} \alpha_j$. The forward noising law is

$$x_k = \sqrt{\bar{\alpha}_k}\, x_0 + \sqrt{1 - \bar{\alpha}_k}\, \epsilon, \qquad \epsilon \sim N(0, I_2).$$

A network $\epsilon_\theta(x_k, k/T)$ minimizes the ordinary mean squared noise-prediction error. It is a classical stochastic sampler containing quantum-circuit features, not an implemented coherent quantum channel. The executed reverse update uses

$$\mu_\theta = \frac{x_k - \beta_k \epsilon_\theta / \sqrt{1-\bar{\alpha}_k}}{\sqrt{\alpha_k}}, \qquad \widetilde{\beta}_k = \beta_k \frac{1-\bar{\alpha}_{k-1}}{1-\bar{\alpha}_k}, \qquad x_{k-1} = \mu_\theta + \sqrt{\widetilde{\beta}_k}\, z.$$

The last step is deterministic. Even an exact conditional noise predictor does not make this Gaussian reverse kernel the exact reverse mixture at finite T: posterior uncertainty of the mixture component remains. The oracle-noise control therefore retains a reverse-kernel approximation, finite schedule and Monte Carlo error, not just a numerical integration error.

There is also an exact, known-teacher reverse-mixture control. Conditional on center s, put $v_k = \bar{\alpha}_k \sigma^2 + 1 - \bar{\alpha}_k$ and $m_{k,s} = \sqrt{\bar{\alpha}_k} s$. At a supplied x_k, draw the center with posterior weights proportional to $p_s \exp(-\|x_k - m_{k,s}\|^2/(2v_k))$. Then draw

$$x_{k-1} \mid x_k, s \sim N\left(m_{k-1,s} + \frac{\sqrt{\alpha_k} v_{k-1}}{v_k}(x_k - \sqrt{\alpha_k} m_{k-1,s}), \frac{v_{k-1}\beta_k}{v_k} I_2 \right).$$

These formulas follow by conditioning a jointly Gaussian pair with covariance $\sqrt{\alpha_k} v_{k-1} I_2$. Starting from the exact forward terminal mixture and composing these reverse conditionals recovers p_σ exactly in distribution, up to numerical and sampling error. It uses the full teacher, so it is a diagnostic rather than a learned solution.

S.3 Flow matching with a measurable target velocity

Independently sample $z \sim N(0, I_2)$ and $x_1 \sim p_\sigma$, set $x_t = (1-t)z + t x_1$, and train $v_\theta(x_t, t)$ toward $x_1 - z$. This is the conditional flow-matching regression objective. Conditional on center s, its time-t mean and variance are

$$ts, \qquad v_t = (1-t)^2 + t^2\sigma^2.$$

Let $\pi_s(x,t)$ be the corresponding mixture posterior. The exact conditional velocity is

$$v_*(x,t) = \sum_s \pi_s(x,t) \left[s + \frac{t\sigma^2 - (1-t)}{v_t}(x - ts) \right].$$

Indeed $\mathrm{Cov}(x_1 - z, x_t \mid s) = (t\sigma^2 - (1-t))I_2$; Gaussian conditioning gives the displayed formula. The positive variance v_t includes both endpoints. The code verifies this formula and the diffusion-score formula by independent finite differences of a separately written mixture log density. Learned sampling uses 128 Heun steps. Oracle-velocity sampling isolates finite integration and sampling error for this known smooth density; it is not an upstream QFM reproduction [94].

S.4 Architecture, matched controls and actual GPU results

The neural input is $[x_0, x_1, t, \sin 2\pi t, \cos 2\pi t, \sin 4\pi t, \cos 4\pi t]$. An affine encoder maps its seven channels to four angles through $\pi \tanh$. A two-layer four-qubit circuit uses angle-reuploading R_Y, R_Z gates and, for the entangled architecture, a CNOT ring. Fifteen commuting Z-string expectations are concatenated with the seven input channels. A $22 \to 64 \to 64 \to 2$ SiLU head predicts noise or velocity. The product control uses the same nominal parameters and removes the entanglers. Final diagonal phase rotations are invisible to the all-Z readout; equality of nominal parameter counts does not establish equality of effective Fisher capacity.

Every run has 5,810 nominal parameters, batch size 512 and 2,000 Adam steps. There are 16,384 training draws and 4,096 disjoint held-out draws, with seeds 20261104, 20261105 and 20261106. The four architectures/objectives were rerun at every seed. The following figures are mean and sample standard deviation across those three runs, not confidence intervals or asymptotic theorems.

Objective	circuit	sign-law TV	sliced W_1
Diffusion	product	$.0380 \pm .0113$	$.0768 \pm .0306$
Diffusion	entangled	$.0451 \pm .0133$	$.0939 \pm .0183$
Flow	product	$.0217 \pm .0079$	$.0460 \pm .0077$
Flow	entangled	$.0293 \pm .0130$	$.0522 \pm .0159$

The second metric uses sixteen fixed projection directions and an independent teacher draw. The experiment demonstrates learning but no consistent quantum or entanglement advantage; the product control has lower mean error on both reported metrics. No physical quantum processor, shot noise, industrial chemical prediction, OOD material transfer or complexity speedup was tested. All checkpoints, seeds, histories, runtimes and teacher controls are retained.




Figure 1: Actual final samples from seed20261104, with exact mixture centers and two-standard-deviation circles. Titles report mean sign-law TV over three seeds. The product control is competitive; no entanglement advantage is inferred.

S.5 A coherent learned purification flow

We separately implement quantum evolution, without identifying it with the classical samplers. Exact depolarizing diffusion is

$$D_t(\rho_R) = e^{-t}\rho_R + (1 - e^{-t})I_4/4.$$

It preserves trace and positivity and satisfies $D_s D_t = D_{s+t}$ by direct substitution. Its linear inverse on all matrices is not positive: $D_t^{-1}(P) = e^t P - (e^t - 1)I_4/4$ has a negative eigenvalue on the orthogonal complement of any rank-one projection P when $t > 0$. Thus a learned inverse cannot be a universal deterministic physical channel.

For a constructive state-preparation benchmark, take the family $R_t = (1 - t)I_4/4 + t\rho_R$, $0 \le t \le 1$, and its canonical purification $|\Psi_t\rangle = \operatorname{vec}\sqrt{R_t}$ in $\mathbb{C}^4 \otimes \mathbb{C}^4$. It is normalized because $\|\sqrt{R_t}\|_F^2 = \operatorname{tr} R_t = 1$. The initial purification is two Bell pairs. We train a six-block four-qubit circuit: each block conjugates local R_Y rotations by a fixed CNOT network and its inverse; angles are $t\theta + t^2\omega$. All blocks are identity at the control time zero. The 48 learned controls are trained against full known purification states, using the fidelity loss. This ideal supervision is stronger than access to measured samples and is accounted for explicitly.

An RTX4090 run of 2,000 steps, batch 64, gives minimum fidelity 0.999903 on a 33-point grid and maximum reduced-density Frobenius error 0.009642. The reduced state is obtained by an actual ancilla partial trace and passes trace/positivity checks. These are grid diagnostics, not a uniform or out-of-family guarantee. The construction is a learned coherent control path for known targets, not a universally learned inverse to depolarization. Its full 76-gate replay and intermediate complex states are exported to the same numerical architecture inspector as the other models.

S.6 Width, fields, RG and arithmetic obligations

Proposition 15.1 fixes which width is being increased: the number of independent one-qubit features, with only their readout trained. At initialization let $a_r \sim N(0, 1)$ be independent standard centered readout coefficients, independent of the frozen features. Conditional finite-dimensional outputs are then exactly centered Gaussian with covariance K_m. The almost-sure kernel convergence implies convergence of these finite-dimensional characteristic functions to the Gaussian law with covariance K. This proves the specified finite-dimensional GP limit. It does not prove tightness in a function-space norm, parameter-feature learning, a many-qubit limit or a continuum QFT. For frozen K_m, squared-loss gradient flow gives $\dot{r} = -K_m r$, whose eigencomponents decay at their kernel eigenvalues. A zero eigenvalue leaves a nonzero residual unchanged.

The executable width experiment fits the existing exact 32-configuration *classical* black-spin-decimated action on genus two, including its constant. This is a second true RG teacher, distinct from the quantum white-spin partial trace above. With 96 parameter triples for training, 32 for testing, eight feature seeds and ridge 10^{-5}, widths 16 and 4,096 give mean relative kernel errors 0.2990 and 0.01726 and mean full-action test MSE 0.02005 and 0.000743. The limiting RBF readout is also evaluated. Finite numerical convergence supports the implementation; the proposition supplies the limit proof. Quantum NTK literature requires its own parameterization, ensemble and lazy/large-width hypotheses [91, 90].

For complex embeddings the Hermitian Gram form, realification $A + iB \mapsto \left(\begin{smallmatrix} A & -B \\ B & A \end{smallmatrix}\right)$, and conjugation must be respected. An arbitrary real network on twice as many channels need not be complex linear. In characteristic two, $1+1 = 0$; an embedding into real vectors cannot be a unital field homomorphism preserving that addition. Finite-field path algebras and cellular/CSS computations are still meaningful, but Gaussian integration, ordered Gibbs weights and real backpropagation do




Figure 2: Left: eight-seed frozen-feature kernel errors, with sample standard deviations. Right: the separate learned coherent purification flow, evaluated on33time points. These are distinct width and quantum-control experiments.

not transfer by renaming the field. An explicit discrete learner over $\mathbb{F}_p$, $p = 2, 3, 5$, uses features $1, x, y, xy$ and coefficients searched in the finite field to learn $f(x, y) = xy$. All p^2 inputs are verified; an affine-only learner has nonzero minimum error. The trained map compiles to the permutation unitary $U_f|x, y, z\rangle = |x, y, z + f(x, y) \bmod p\rangle$ on a complex Hilbert space of dimension p^3. Retaining x, y makes its inverse subtraction explicit. This is reversible encoding of a field-valued polynomial learner, not a field embedding into complex amplitudes or backpropagation in $\mathbb{F}_p$. Its training is exact finite coefficient search. A tropical semiring is not a field; coefficient cancellation, valuation signs and the underlying valued algebra must be specified.

For GRH, each architecture offers an experimentally testable role: quantum features can parameterize candidate metrics or probe observables; graph tokens can encode actual local incidence, masks and monodromy; tropical attention can search extremal valuations or active constraints; flow/diffusion can propose continuous operator/control parameters; RG can organize a genuine hierarchy of coarse actions. None of these roles proves that its Gram matrix is the arithmetic positive metric. One must still construct the arithmetic object with correct conductor, gamma factors, zero divisor and domain, prove compact-uniform control and zero location, and keep Dirichlet, Hecke, Dedekind and Artin hypotheses separate. A learned candidate is accepted only after its exact defining equations and analytic estimates are supplied.

T Covers, paths, pants and quantum games

T.1 Which free group and which universal cover?

Write $F_2 = \langle x, y\rangle$, $\mathbb{F}_2$ and $\widehat{F}_2$ for the free group, coefficient field and profinite completion, respectively. For a theta graph with vertices A, B and edges a, b, c, an explicit cover has vertices $(g, A), (g, B)$, $g \in F_2$, and edges

$$a : (g, A) - (g, B), \qquad b : (g, A) - (gx, B), \qquad c : (g, A) - (gy, B).$$

Left multiplication is the deck action. Contracting all a-edges produces the usual four-regular Cayley tree of F_2; before contraction the graph is three-regular. Equivalently, vertices are reduced nonbacktracking edge paths from A, with an immediately reversed edge cancelled. This gives a

tree and unique path lifting in every length. Its radius-r patch has $1 + 3\sum_{k=0}^{r-1} 2^k$ vertices, hence 766 at radius eight. With a planar ribbon structure the abstract graph is a spine of a topological pair of pants. The actual $m = 3$ dessin instead has the ribbon structure of a once-punctured torus. Forgetting the cyclic orders identifies the abstract graphs, not these thickenings. Graph distance is not automatically hyperbolic geodesic length.

For the contracted clean m-edge dessin there are two vertices and m parallel edges, so π_1 is F_{m-1} and the analogous universal tree is m-regular. The new examples include $m = 5, 9$, of genus two and four. Reinserting clean white bivalent vertices subdivides its edges. Attaching the face to recover the compact surface introduces a relation; the closed genus-two surface has its surface group and a disk universal cover, not one of these graph trees. Separately, $\mathbb{P}^1 \setminus \{0, 1, \infty\}$ has fundamental group F_2. The unbranched cover induced by a clean dessin removes all preimages of those three values. For our degree-$2m$ map there are $m + 3$ punctures; its Euler characteristic is $-2m$, not that of the dessin's one-skeleton. Thus the occurrences of F_2 in monodromy, graph covers and arithmetic profinite theory must not be silently identified.




Figure 3: Finite windows of universal covers of contracted actual clean dessins. These are schematic graph trees, not uniformizing embeddings of the compact surfaces. Clean white bivalent vertices are suppressed.

For finite-supported representations of a suitably graded bound quiver, pushdown maps split into deck-homogeneous blocks. The arrow equations preserve the degree, and monomial zero relations are homogeneous. The resulting Hom direct sum controls all deck pieces, not just degree one. For an explicit graph-only diagnostic two lifted edge intervals have upstairs endomorphism dimension two, but downstairs endomorphism dimension four: the deck pieces have dimensions $2 + 1 + 1$. This graph-quiver calculation is *not* substituted for a dessin-algebra module example. The actual infinite polynomial module examples below supply that part. A Hom-controlled functor needs its specified exceptional/source/target subcategory and control statement; exactness, a deck-sum formula or equal dimensions alone is not an equivalence or an AR universality theorem. A derived comparison must control all $\mathrm{Hom}(X, Y[n])$ and composition, together with an actual derived/enhanced functor.

T.2 Actual infinite-algebra path modules and unbounded resolutions

To fix conventions, let A_{col} be the earlier rightmost-arrow-first polynomial matrix algebra acting on left columns. Here we use travel-order path multiplication and write $A_{\mathrm{trav}} = A_{\mathrm{col}}^{\mathrm{op}}$ for the prose-identified bound-quiver algebra. Right A_{trav} representations therefore encode the earlier

covariant left-column representations. There are vertices $i \in \mathbb{Z}/5$ and arrows $a_i, b_i : i \to i-1$, with

$$a_i b_{i-1} = b_i a_{i-1} = 0.$$

Both pure branch paths survive at every length. No arrow-truncated finite algebra is being used. We work with ordinary polynomial modules, not with all completed lattices or all infinite strings.

Put $P_i = e_i A_{\mathrm{trav}}$ and let S_i be its residue simple. The augmentation kernel is $a_i A \oplus b_i A$. Multiplication by an arrow kills exactly the opposite-branch positive paths at its target, while each surviving path has a distinct normal image. Its kernel is therefore the right ideal generated by its unique forbidden successor, without cancellation. Iteration yields a projective resolution with

$$P_0 = P_i, \qquad P_n = P_{i-n} \oplus P_{i-n} \quad (n \geq 1).$$

The two arms are indexed by alternating-branch relation words. The *differential entry is the last arrow*, not the product of the zero relation word. This kernel description proves exactness in every degree. Applying Hom to a simple kills every arrow entry, hence

$$\dim \mathrm{Hom}(S_i, S_j) = \mathbf{1}_{i=j}, \qquad \dim \mathrm{Ext}^n(S_i, S_j) = 2\mathbf{1}_{j=i-n} \quad (n \geq 1).$$

Tensoring with the left residue simple gives the same Tor dimensions. A graded version needs its internal degree shifts. In particular the residue simples have infinite projective dimension in this ambient algebra.

Let B be the one-arrow string supported at vertices zero and four, with $a_0 = 1$ and every other arrow zero. Its presentation is

$$P_4 \oplus P_3 \xrightarrow{(b_0,\ a_0 a_4)} P_0 \longrightarrow B \longrightarrow 0.$$

The two images have disjoint normal-path support and span the kernel, because B retains exactly e_0, a_0. The next entries are a_4 on the first arm and b_3 on the second; continue by their unique forbidden successors. This gives an unbounded exact resolution by the same normal-path argument. Full arrow-commuting equations give

$$\dim \mathrm{Hom}(B, B) = 1, \quad \dim \mathrm{Hom}(S_0, B) = 0, \quad \dim \mathrm{Hom}(B, S_0) = 1, \quad \dim \mathrm{Hom}(S_4, B) = 1.$$

In the Hom complex for $\mathrm{Ext}^1(B, B)$ both adjacent differentials vanish, so $\mathrm{Ext}^1(B, B) \simeq \mathbb{Q}$. A concrete nonsplit self-extension has $\mathbb{Q}^2$ at zero and four, $a_0 = I_2$, $b_0 = \left(\begin{smallmatrix} 0 & 1 \\ 0 & 0 \end{smallmatrix}\right)$, all other arrows zero. The first-coordinate submodule and quotient are B. A section at vertex zero would lift the quotient basis to a vector with second coordinate one; b_0 sends it to a nonzero first-coordinate vector, contradicting commutation with the zero b_0 action on B. Thus B is not rigid in this actual ambient category. No almost-split universality is inferred from this nonsplitting.

For finite-dimensional N let $DN = \mathrm{Hom}_{\mathbb{Q}}(N, \mathbb{Q})$. On each finite idempotent projective the evaluation pairing identifies $P \otimes_A DN$ with $D \mathrm{Hom}_A(P, N)$. Dualization is exact on these finite-dimensional terms, so the resulting complexes give $\mathrm{Tor}^A_n(M, DN) \simeq D \mathrm{Ext}^n_A(M, N)$ for the stated finitely generated resolutions. In particular $\mathrm{Tor}^A_1(B, DB) \simeq \mathbb{Q}$. The exact program checks six source/target pairs through degree twelve, all five simple-target formulas and 250 normal-kernel samples. The unbounded resolution assertions rest on the normal-form proof, not on the finite cutoff. These new homological proofs are prose/exact calculations; full corresponding resolution objects are not yet Lean packaged.

A geometric path may label a string after specifying the ribbon structure, relations and category. Turning it into a geodesic needs a metric and appropriate boundary/stop data. Intersections compute morphisms only under the hypotheses of an actual graded/Fukaya comparison. The homologically smooth graded-gentle perfect-category theorems do not identify our entire ordinary or completed module category with the Fukaya category by inspection [93]. In particular the residue resolutions warn against silently imposing smoothness or finite projective dimension.

T.3 Pants tensors, physical dilation and sewing boundaries

For $C = \mathbb{Q}[\epsilon]/\epsilon^2$, take $\varepsilon(1) = 0$, $\varepsilon(\epsilon) = 1$. In the ordered bases $(1, \epsilon)$ and $(1 \otimes 1, 1 \otimes \epsilon, \epsilon \otimes 1, \epsilon \otimes \epsilon)$, multiplication and comultiplication are

$$\mu = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \end{pmatrix}, \qquad \Delta = \begin{pmatrix} 0 & 0 \\ 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

The pairing is $\left(\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right)$, with determinant -1. Associativity follows from $\epsilon^2 = 0$; coassociativity sends one to the three placements of a single one among two ϵ's and sends ϵ to $\epsilon^{\otimes 3}$. On the four tensor basis vectors, all three maps in

$$\Delta\mu = (\mu \otimes I)(I \otimes \Delta) = (I \otimes \mu)(\Delta \otimes I)$$

give, respectively, $1 \otimes \epsilon + \epsilon \otimes 1$, $\epsilon \otimes \epsilon$, $\epsilon \otimes \epsilon$, and zero. This proves the Frobenius identity. The unit and counit identities follow directly on the same basis. These are exact oriented 2D TQFT gluing data [95], not a presumed positive Hilbert structure.

The handle map $H = \mu\Delta$ sends $1 \mapsto 2\epsilon$ and $\epsilon \mapsto 0$. Closed genus-g contractions $\varepsilon H^g \eta$ are $0, 2, 0, 0, \ldots$. The genus-two value really vanishes in this example. In characteristic two the torus value also vanishes, illustrating a meaningful coefficient change rather than an embedding-space relabelling. After extending to $\mathbb{C}$ with the Euclidean metric, $\Delta^*\Delta = \mathrm{diag}(2, 1)$ and $\Delta \neq \mu^*$; $\mu : \mathbb{C}^4 \to \mathbb{C}^2$ is not an isometry. The exact code constructs $K = \mu/\sqrt{2}$ and a 4×4 failure block L with $K^*K + L^*L = I_4$. Stacking them gives an isometry into $\mathbb{C}^2 \oplus \mathbb{C}^4$, extendable to a unitary with ancillas. Its success probabilities on the four basis states are $1/2, 1/2, 1/2, 0$. Postselection implements K with state-dependent success; normalizing the selected outcome is not the original linear map μ.

For a closed genus-$g \geq 2$ surface a pants decomposition has $2g - 2$ pieces and $3g - 3$ sewn curves. Conformal sewing adds moduli-dependent propagators and analytic consistency [96]; string amplitudes need worldsheet, ghost and integration data beyond this finite tensor. A pants network or a hyperbolic graph is likewise not an AdS/CFT duality or a Ryu–Takayanagi derivation [97]. Such interpretations remain specified research targets until their extra structure is constructed. Arithmetic/anabelian reconstruction additionally requires arithmetic profinite fundamental-group data and its Galois structure under the proper field/curve hypotheses. Neither a discrete F_2 graph nor a coefficient $\mathbb{F}_2$ module supplies that arithmetic extension [98].

T.4 Quantum games and the Connes-embedding distinction

For uniform input bits x, y, CHSH accepts outputs u, v when $u \oplus v = xy$. With ± 1 observables its success probability is $1/2 + \langle T \rangle/8$, where

$$T = A_0 \otimes (B_0 + B_1) + A_1 \otimes (B_0 - B_1).$$

For a deterministic classical assignment one bracket vanishes and the other is ± 2, giving success at most $3/4$; all sixteen assignments are checked. On $|\Phi^+\rangle$, take $A_0 = Z, A_1 = X$, $B_0 = (Z + X)/\sqrt{2}, B_1 = (Z - X)/\sqrt{2}$. Then $T = \sqrt{2}(Z \otimes Z + X \otimes X)$ gives success $(2 + \sqrt{2})/4$. For arbitrary self-adjoint involutions, $T^2 = 4I - [A_0, A_1] \otimes [B_0, B_1]$ and each commutator has norm at most two. Hence $\|T\| \leq 2\sqrt{2}$, proving optimality. The exact matrix checker verifies the entire spectrum $\{-2\sqrt{2}, 0, 0, 2\sqrt{2}\}$, not just a numerical sample.

MIP*=RE, in its pinned v4, gives an external unrestricted correlation separation and a negative answer to Connes embedding [99]. We do not rerun its external formalization or claim a new proof of it. CHSH has its optimum on two qubits and the same commuting optimum by the preceding bound; it does not exhibit the general separation. A learned game strategy is useful for finite witness search and checked bounds, but unrestricted computability and dimension quantifiers cannot be settled by a finite game simulation.

T.5 Tropical accelerator reference designs and reproducibility

The chip-design campaign pins twenty public repositories, twenty-four focused document ranges and thirteen primary papers. It covers AlphaChip/Circuit Training, independent macro-placement benchmarks, OpenROAD-family flows, quantum layout and low-level control. The source ledger records what was actually read, not a full audit or reproduction of every model. An exact public identity for the phrase "bare-metal Qiskit" was not established; no repository was invented.

Three original executable references implement a masked signed max-plus tile, an ordered mixed MAC/max-plus scheduler and a fail-closed synthetic IQ steering controller. Wide addition precedes winner selection and final saturation; earliest-index tie handling is explicit. Empty masks remain invalid, not an extra candidate of value zero. The controller requires a current calibration epoch, enough samples and a valid margin/deadline before allowlisting a steering action. Checks include all 65,536 signed eight-bit additions, 262,144 masked two-lane cases, reordered completions and twelve negative controls. These are digital reference models, not synthesized RTL, routed chips, measured RF feedback, power/area estimates or universal hardware speedups. A tropical context block can avoid multiplications inside that primitive while its projections, scores and other layers still require their own arithmetic.

The Linux snapshot preserves the Mac project, source clones, histories, checkpoints and environment archives. Linux GPU execution uses the existing Torch CUDA environment and a separately approved isolated cuQuantum prefix; macOS binaries are not treated as a Linux installation. The conda lock, setup instructions, gate/head replay records and all failures are retained. The on-page book contains one canonical laboratory for this chapter and links all five new checkpoint architectures into the existing tensor inspector. New physics, width and homological results have their explicit prose/numerical/external status; selected earlier Lean components do not constitute a full formalization of this extension or a proof of GRH.

References

[1] A. Schreiber, *Surface Algebras I: Dessins D'enfants, Surface Algebras, and Dessin Orders*, arXiv:1810.06750v5 (2023).

[2] A. Schreiber, *Surface Algebras and Surface Orders II: Affine Bundles on Curves*, arXiv:1812.00621v1; record 2018, downloaded PDF internally dated November 27, 2024.

[3] A. Schreiber, *Multiparameter Persistent Homology—Generic Structures and Quantum Computing*, arXiv:2210.11433v1 (2022).

[4] C. Bessenrodt and T. Holm, *Weighted locally gentle quivers and Cartan matrices*, arXiv:math/0511610v1 (2005).

[5] S. Ford, A. Oswald and J.J. Zhang, *Homological conditions on locally gentle algebras*, arXiv:2409.08333v3 (5 September 2026).

[6] S. Opper, P.-G. Plamondon and S. Schroll, *A geometric model for the derived category of gentle algebras*, arXiv:1801.09659v7 (10 July 2025).

[7] W. Chang, *Geometric models for the algebraic hearts in the derived category of a gentle algebra*, arXiv:2305.03998v2 (4 September 2024).

[8] Y.-Z. Liu, C. Zhang and H. Zhang, *Constructing projective resolution and taking cohomology for gentle algebras in the geometric model*, arXiv:2308.07220v1 (2023).

[9] Z. Li, Y. Qiu and Y. Zhou, *A geometric realization of Koszul duality for graded gentle algebras*, arXiv:2403.15190v1 (2024).

[10] W. Chang, H. Jin and S. Schroll, *Recollements of partially wrapped Fukaya categories and surface cuts*, arXiv:2206.11196v2 (2023).

[11] B. Dequêne and M. Schoonheere, *Resolving subcategories for gentle algebras I: Monogeneous resolving subcategories for gentle trees*, arXiv:2502.20994v3 (2026).

[12] D.A. Buchsbaum and D. Eisenbud, *What makes a complex exact?*, J. Algebra 25 (1973), 259–268, primary paper.

[13] W. Bruns, *The existence of generic free resolutions and related objects*, Math. Scand. 55 (1984), 33–46.

[14] D. Eisenbud, *Homological algebra on a complete intersection, with an application to group representations*, Trans. AMS 260 (1980), 35–64.

[15] A. Beauville and Y. Laszlo, *Un lemme de descente*, C. R. Acad. Sci. Paris 320 (1995), 335–340, author text.

[16] P. Magyar, *Affine Schubert varieties and circular complexes*, arXiv:math/0210151v1 (2002).

[17] F.F. Knudsen and D. Mumford, *The projectivity of the moduli space of stable curves I: Preliminaries on 'det' and 'Div'*, Math. Scand. 39 (1976), 19–55, doi:10.7146/math.scand.a-11642.

[18] M. Breuning, *Determinant functors on triangulated categories*, arXiv:math/0610435v1 (2006).

[19] D. Eriksson and G. Freixas i Montplet, *Deligne–Riemann–Roch and intersection bundles*, J. Éc. polytech. Math. 11 (2024), 247–361, doi:10.5802/jep.254.

[20] P. Deligne, *La conjecture de Weil I*, Publ. Math. IHÉS 43 (1974), 273–307, primary text.

[21] P. Deligne, *La conjecture de Weil II*, Publ. Math. IHÉS 52 (1980), 137–252, primary text.

[22] P. Deligne, *Cohomologie étale (SGA $4\frac{1}{2}$)*, Lecture Notes in Mathematics 569, Springer-Verlag, 1977, doi:10.1007/BFb0091516.

[23] G. Tabuada, *Noncommutative Weil conjecture*, arXiv:1808.00950v7 (6 December 2019).

[24] C. Deninger, *Arithmetic geometry and analysis on foliated spaces*, arXiv:math/0505354v1 (2005).

[25] D.B. Ray and I.M. Singer, *R-Torsion and the Laplacian on Riemannian manifolds*, Advances in Math. 7 (1971), 145–210, doi:10.1016/0001-8708(71)90045-4.

[26] F. Bornemann, *On the numerical evaluation of Fredholm determinants*, arXiv:0804.2543v1; Math. Comp. 79 (2010), 871–915.

[27] B. Dequêne and M. Schoonheere, *Resolving subcategories for gentle algebras II: Resolving subcategories for gentle trees*, arXiv:2509.10120v2 (3 October 2025).

[28] C. Procesi, *The invariant theory of $n \times n$ matrices*, Advances in Math. 19 (1976), 306–381, doi:10.1016/0001-8708(76)90027-X.

[29] L. Le Bruyn and C. Procesi, *Semisimple representations of quivers*, Trans. Amer. Math. Soc. 317 (1990), 585–598, doi:10.1090/S0002-9947-1990-0958897-0.

[30] A. Craw and R. Yamagishi, *The Le Bruyn–Procesi theorem following Lusztig*, arXiv:2312.08527v4 (14 November 2025).

[31] P. Etingof, O. Golberg, S. Hensel, T. Liu, A. Schwendner, D. Vaintrob and E. Yudovina, *Introduction to Representation Theory*, with historical interludes by S. Gerovitch, Student Mathematical Library 59, American Mathematical Society, 2011, doi:10.1090/stml/059; author-hosted text.

[32] K. Akin, D.A. Buchsbaum and J. Weyman, *Schur functors and Schur complexes*, Advances in Math. 44 (1982), 207–278, doi:10.1016/0001-8708(82)90039-1.

[33] Q. Jiang, *Derived Grassmannians and derived Schur functors*, arXiv:2212.10488v3 (6 July 2023).

[34] A. Price, A. Stelzer and A. Yong, *Representations from matrix varieties, and filtered RSK*, arXiv:2403.09938v2 (6 October 2025).

[35] H. Derksen and J. Weyman, *Semi-invariants of quivers and saturation for Littlewood–Richardson coefficients*, J. Amer. Math. Soc. 13 (2000), 467–479, primary author text.

[36] A. Schofield and M. Van den Bergh, *Semi-invariants of quivers for arbitrary dimension vectors*, arXiv:math/9907174v1 (27 July 1999); Indag. Math. 12 (2001), 125–138.

[37] J.S. Milne, *Reductive Groups*, author's course notes, version 2.00 (10 March 2018), §5.18.

[38] W. Crawley-Boevey, *Classification of modules for infinite-dimensional string algebras*, Trans. Amer. Math. Soc. 370 (2018), 3289–3313, doi:10.1090/tran/7032; arXiv:1308.6410v2 (4 March 2016).

[39] M. Auslander and I. Reiten, *Almost split sequences for Cohen–Macaulay-modules*, Math. Ann. 277 (1987), 345–349, doi:10.1007/BF01457367.

[40] R. Bautista, S. Liu and C. Paquette, *Representation Theory of an Infinite Quiver*, arXiv:1109.3176v1 (14 September 2011); published as *Representation theory of strongly locally finite quivers*, Proc. London Math. Soc. 106 (2013), 97–162.

[41] C. Paquette, *Irreducible morphisms and locally finite dimensional representations*, arXiv:1508.04353v3 (4 January 2016).

[42] S. Liu, P. Ng and C. Paquette, *Almost split sequences and approximations*, arXiv:1203.5009v1 (19 March 2012).

[43] C.M. Ringel, *The Minimal Representation-Infinite Algebras which are Special Biserial*, author-hosted ICRA Tokyo 2010 proceedings text, 57 pages, retrieved 3 November 2026.

[44] G. Puninski and M. Prest, *Ringel's conjecture for domestic string algebras*, arXiv:1407.7470v2 (13 September 2015).

[45] R. Laking, M. Prest and G. Puninski, *Krull–Gabriel dimension of domestic string algebras*, arXiv:1506.07703v2 (29 June 2015).

[46] H.-L. Huang, Z. Lin and X. Su, *The Auslander–Reiten quivers of string algebras of affine type $\widetilde{C}$ and a conjecture by Geiss–Leclerc–Schröer*, arXiv:2110.02777v2 (16 November 2021).

[47] G. Abrams and G. Aranda Pino, *The Leavitt path algebra of a graph*, arXiv:math/0509494v1 (21 September 2005).

[48] T. Bates, D. Pask, I. Raeburn and W. Szymański, *The C^*-algebras of row-finite graphs*, New York J. Math. 6 (2000), 307–324, primary journal text.

[49] N.J. Fowler and I. Raeburn, *The Toeplitz algebra of a Hilbert bimodule*, arXiv:math/9806093v3 (23 June 1998).

[50] N. Brownlowe, M. Laca, D. Robertson and A. Sims, *Reconstructing directed graphs from generalised gauge actions on their Toeplitz algebras*, arXiv:1812.08903v1 (21 December 2018).

[51] E.T.A. Kakariadis and O.M. Shalit, *On operator algebras associated with monomial ideals in noncommuting variables*, arXiv:1501.06495v3 (21 November 2018).

[52] O.M. Shalit and B. Solel, *Subproduct systems*, arXiv:0901.1422v3 (26 May 2009).

[53] A. Connes and C. Consani, *Knots, primes and class field theory*, arXiv:2501.06560v1 (11 January 2025).

[54] A. Connes, C. Consani and H. Moscovici, *Zeta zeros and prolate wave operators*, arXiv:2310.18423v2 (4 May 2024).

[55] A. Connes, C. Consani and H. Moscovici, *On q-series and the moment problem associated to local factors*, arXiv:2403.01247v1 (2 March 2024).

[56] A. Connes, C. Consani and H. Moscovici, *Zeta Spectral Triples*, arXiv:2511.22755v1 (27 November 2025).

[57] A. Connes and W.D. van Suijlekom, *Quadratic Forms, Real Zeros and Echoes of the Spectral Action*, arXiv:2511.23257v1 (28 November 2025).

[58] A. Connes, *The Riemann Hypothesis: Past, Present and a Letter Through Time*, arXiv:2602.04022v1 (3 February 2026).

[59] A. Connes and C. Consani, *On the Jacobian of* $\overline{\mathrm{Spec}\,\mathbb{Z}}$, arXiv:2602.15941v1 (17 February 2026).

[60] C. Deninger, *Dynamical systems for arithmetic schemes*, arXiv:1807.06400v4 (7 February 2024); Indag. Math. 37 (2026), 25–136, doi:10.1016/j.indag.2024.05.007.

[61] C. Deninger, *Primes, knots and periodic orbits*, arXiv:2301.11643v1 (27 January 2023).

[62] C. Deninger, *There is no "Weil-"cohomology theory with real coefficients for arithmetic curves*, arXiv:2204.02714v2 (27 January 2023).

[63] R. Ponge and Y. Tian, *Nonclassical Weyl laws and Connes' Integration for weak Lorentz ideals, I*, arXiv:2605.25591v2 (28 September 2026).

[64] F.W.J. Olver et al. (eds.), *NIST Digital Library of Mathematical Functions*, finite summation by parts, Eq. 2.10.9 and Dirichlet L-functions, §25.15, consulted 3 November 2026.

[65] A. Chirvasitu, P.M. Hajac and M. Tobolski, *Non-surjective pullbacks of graph C*-algebras from non-injective pushouts of graphs*, arXiv:1907.10260v3 (31 December 2019), Theorems3.4 and4.3; doi:10.1112/blms.12389.

[66] Y. Lekili and A. Polishchuk, *Derived equivalences of gentle algebras via Fukaya categories*, arXiv:1801.06370v5 (27 August 2019), Definition3.1.1, Lemma3.1.3 and Theorem3.2.2.

[67] M. Tomforde, *Uniqueness theorems and ideal structure for Leavitt path algebras*, arXiv:math/0612628v6 (27 February 2007), Theorem7.3, pp.29–30.

[68] A. Weil, *Numbers of solutions of equations in finite fields*, Bull. Amer. Math. Soc.55 (1949),497–508, especially pp.499–502 and Eq.(4), doi:10.1090/S0002-9904-1949-09219-4; primary-paper scan.

[69] The Stacks Project, Tag0FJ9 and Tag0GSX (retrieved 3 November 2026).

[70] J. Alper, *Good moduli spaces for Artin stacks*, Ann. Inst. Fourier63 (2013),2349–2402, Theorem10.3, pp.2386–2387, doi:10.5802/aif.2833.

[71] The Stacks Project, Tag04V8: Taking roots of line bundles (retrieved 3 November 2026).

[72] G. Zwara, *Smooth morphisms of module schemes*, Proc. London Math. Soc. (3) 84 (2002), 539–558, primary publication.

[73] C. Riedtmann and G. Zwara, *Orbit closures and rank schemes*, Comment. Math. Helv. 88 (2013), 55–84, primary publication, Section6.

[74] Y. Liu, N. Li, B. Xing and P. Chen, *A lecture note on covering theory in representation theory of algebras*, arXiv:2508.01563v5, 28 May 2026.

[75] Q. Dong, Z. Lin, M. Lu and S. Ruan, *Representation-tame Geiß–Leclerc–Schröer algebras and a revised GLS conjecture on root systems*, arXiv:2609.26503v1, 22 September 2026.

[76] S. Schroll, *On geometric models in representation theory*, Proceedings of ICM 2026, vol.3, pp.250–271, doi:10.1137/25M1810647; author preprint arXiv:2601.14396v1.

[77] B. Keller, *Introduction to A_∞ algebras and modules*, arXiv:math/9910179v2 (2001), especially §§3.1,3.3,6.2.

[78] The Stacks Project Authors, *The Koszul complex*, Tag 0621, especially Definitions15.29.1–2 and Lemma15.29.5; snapshot November3,2026, hash recorded in the repository.

[79] The Stacks Project Authors, *Regular sequences are Koszul regular*, Tag 062F, Lemma15.31.2; snapshot November3,2026, hash recorded in the repository.

[80] S. Datta, S. Howard and D. Cochran, *Geometry of the Welch Bounds*, arXiv:0909.0206v2 (2012), Section2.1, Theorem2.2.

[81] K.G. Wilson and J. Kogut, *The renormalization group and the epsilon expansion*, Physics Reports12(1974),75–199, DOI10.1016/0370-1573(74)90023-4; pinned institutional scan and review scope in `planning/ACTUAL_RG_SOURCES.json`.

[82] M. Levin and C.P. Nave, *Tensor renormalization group approach to2Dclassical lattice models*, arXiv:cond-mat/0611687v2 (2008); Phys. Rev. Lett.99(2007),120601.

[83] J. Kim et al., *Pure Transformers are Powerful Graph Learners*, arXiv:2207.02505v2 (2022). Our finite measured-feature prototype does not reproduce its full benchmarks.

[84] S. Liu et al., *Unsupervised Discovery of Steerable Factors When Graph Deep Generative Models Are Entangled*, arXiv:2401.17123v1 (2024); scoped primary reading in the computational source manifest.

[85] S. Lahlou et al., *A theory of continuous generative flow networks*, arXiv:2301.12594v2 (2023); state measures and trajectory balance.

[86] M. Gross and B. Siebert, *Mirror Symmetry via Logarithmic Degeneration Data II*, arXiv:0709.2290v2; used as a geometric research context, not an implemented mirror comparison.

[87] B. Green and T. Tao, *Quadratic uniformity of the Möbius function*, arXiv:math/0606087v2, Section4, Lemma4.1, printed pages12–13. Focused identity reading only; no nilsequence estimate is asserted for the moving sextic family here.

[88] S.S. Kudla, *Tate's thesis*, expanded Jerusalem lecture notes (2001), Theorem4.3 and Corollary4.4, pinned primary notes.

[89] J.S. Milne, *Class Field Theory*, version4.03 (6August2020), Theorem0.8 and ChapterI,§1, author notes.

[90] A. Jacot, F. Gabriel and C. Hongler, *Neural Tangent Kernel: Convergence and Generalization in Neural Networks*, arXiv:1806.07572v4; specified classical scaling hypotheses.

[91] J. Liu et al., *Representation Learning via Quantum Neural Tangent Kernels*, arXiv:2111.04225v3; focused version/abstract review.

[92] M. Kontsevich and Y. Soibelman, *Affine structures and non-archimedean analytic spaces*, arXiv:math/0406564v1; valued and integral-affine data.

[93] Y. Lekili and A. Polishchuk, *Derived equivalences of gentle algebras via Fukaya categories*, arXiv:1801.06370v5; category and smoothness hypotheses retained.

[94] Z. Cui, P. Zhang and Y. Tang, *Quantum Flow Matching*, arXiv:2508.12413v1; abstract-level comparison, not an upstream reproduction.

[95] L. Abrams, *Two-dimensional topological quantum field theories and Frobenius algebras*, J. Knot Theory Ramifications 5 (1996), 569–587, author-hosted paper.

[96] G. Segal, *The definition of conformal field theory*, in *Topology, Geometry and Quantum Field Theory* (2004); primary publisher metadata and sewing-scope comparison.

[97] S. Ryu and T. Takayanagi, *Holographic Derivation of Entanglement Entropy from AdS/CFT*, arXiv:hep-th/0603001v2; external setting only.

[98] S. Mochizuki, *The Absolute Anabelian Geometry of Hyperbolic Curves*, author-hosted paper, introduction and Theorem 1.3.4; arithmetic extension hypotheses retained.

[99] Z. Ji, A. Natarajan, T. Vidick, J. Wright and H. Yuen, *MIP*=RE*, arXiv:2001.04383v4 (16 October 2026); external theorem and version record, not rerun locally.