
Compute grids of radiative transfer models of dense cores and filaments
Compute the attenuating column Σ_A of every model shell with the dust temperature T, Eq. (4)
Fit the model data with $T(\Sigma_A) = T_{min} + (T_0 - T_{min})(1 + \Sigma_A/\Sigma_\star)^{-\gamma}$, Eq. (5)
Define a model line of sight of true Σ trough a uniform sphere and a uniform layer
Attenuating column $\Sigma_A(u)$ at every depth, averaged over directions and shapes, Eq. (4)
Temperature $T(\Sigma_A(u))$ at every depth, from the fitted relation, Eq. (5)
Emission of all the depths added, giving the intensities $I_\lambda(\Sigma)$, Eq. (6)
One modified blackbody fitted as the image was, giving Σ_F, Eq. (7)
Tabulate $f(\Sigma_F) = \Sigma/\Sigma_F$, what an image holds, Eq. (8)
Below 4×10^{21} cm^{-2}, average the sphere with a radiative transfer cloud, blend to 10^{22}, Sect. 4.3
Multiply by the field factor $g(\Sigma_F, G_0)$ for the field of the cloud and hold at the maximum, Eq. (10)
Evaluate f_{G_0} on the original image $\mathcal{D}$, create the correction factors image $\mathcal{F}$
Multiply the original image by the correction factors: $\mathcal{F} \cdot \mathcal{D}$, Eq. (11)